

2148-13-13



This is an electronic reprint of the original article. This reprint *may differ* from the original in pagination and typographic detail.

Author(s):	Piiroinen, Saija; Lindström, Leena; Lyytinen, Anne; Mappes, Johanna; Chen, Yolanda H; Izzo, Victor; Grapputo, Alessandro
Title:	Pre-invasion history and demography shape the genetic variation in the insecticide resistance-related acetylcholinesterase 2 gene in the invasive Colorado potato beetle
Year:	2013
Version:	
Please cite	the original version:
	Piiroinen, S., Lindström, L., Lyytinen, A., Mappes, J., Chen, Y. H., Izzo, V., & Grapputo, A. (2013). Pre-invasion history and demography shape the genetic variation in the insecticide resistance-related acetylcholinesterase 2 gene in the invasive Colorado potato beetle. BMC Evolutionary Biology, 13(13). https://doi.org/10.1186/1471-

All material supplied via JYX is protected by copyright and other intellectual property rights, and duplication or sale of all or part of any of the repository collections is not permitted, except that material may be duplicated by you for your research use or educational purposes in electronic or print form. You must obtain permission for any other use. Electronic or print copies may not be offered, whether for sale or otherwise to anyone who is not an authorised user.

Online Supplementary material: Piiroinen et al. "*Pre-invasion history and demography shape the genetic variation in the insecticide resistance-related acetylcholinesterase 2 gene in the invasive Colorado potato beetle*".

						Diversity indices						
Region	N	Н	S	Tr.	Pr.	k	h	h (SD	π	π (SD)	$\theta_{\mathbf{W}}$	θ_{W} (SD)
	hE2 ((1037	'bp)									
Mexico	58	18	59	17	41	12.540	0.906	0.021	0.0121	0.0013	0.0123	0.0036
US	38	25	45	13	16	9.764	0.967	0.015	0.0094	0.0086	0.0103	0.0033
Europe	100	9	22	5	1	7.009	0.836	0.018	0.0068	0.0023	0.0041	0.0013
Total	196	49	87	25	87	14.048	0.944	0.007	0.0136	0.0005	0.0143	0.0034
DP1 (931 bp, including a short 58 bp intron site)												
Mexico	58	33	41	10	22	9.154	0.971	0.009	0.0098	0.0007	0.0095	0.0029
US	36	21	41	10	16	7.270	0.946	0.023	0.0078	0.0009	0.0106	0.0035
Europe	88	5	23	10	3	5.576	0.719	0.018	0.0060	0.0005	0.0049	0.0015
Total	182	56	67	18	67	10.837	0.916	0.012	0.0116	0.0004	0.0125	0.0031
<i>JHE-b</i> (583 bp)												
Mexico*	36	21	15	6	8	4.497	0.957	0.019	0.0077	0.0007	0.0062	0.0024
US (Kansas) 6	2	19	7	5	10.133	0.824	0.098	0.0104	0.0016	0.0081	0.0036
Europe [†]	36	15	16	7	4	4.514	0.798	0.059	0.0077	0.0010	0.0066	0.0025
Total	78	38	32	13	32	8.732	0.946	0.016	0.0150	0.0002	0.0111	0.0034

Additional file 1 Summary statistics for the Colorado potato beetle populations pooled
according to geographical regions

Abbreviations: N, number of gene copies; H, number of haplotypes; S, number of polymorphic sites; Tr., number of transversions; Pr., number of private polymorphic sites; k, the average number of nucleotide differences; h, haplotype diversity; π , nucleotide diversity; θ_w , Watterson's theta estimate. *Morelos and Oaxaca populations, †Russian, Finnish and Italian populations.