



## This is an electronic reprint of the original article. This reprint *may differ* from the original in pagination and typographic detail.

- Author(s): Sun, Ren X; Chong, Lauren C; Simmons, Trent T; Houlahan, Kathleen E; Prokopec, Stephenie D; Watson, John D; Moffat, Ivy D; Lensu, Sanna; Lindén, Jere; P'ng, Christine; Okey, Allan B; Pohjanvirta, Raimo; Boutros, Paul C
- Title:Cross-species transcriptomic analysis elucidates constitutive aryl hydrocarbon<br/>receptor activity
- Year: 2014

Version:

## Please cite the original version:

Sun, R., Chong, L., Simmons, T., Houlahan, K., Prokopec, S., Watson, J., Moffat, I.D, Lensu, S. Lindén, J., P'ng, C., Okey, A.B, Pohjavirta, R. & Boutros, P.C. (2014). Cross-species transcriptomic analysis elucidates constitutive aryl hydrocarbon receptor activity. BMC Genomics, 15 (December), 1053. doi:10.1186/1471-2164-15-1053

All material supplied via JYX is protected by copyright and other intellectual property rights, and duplication or sale of all or part of any of the repository collections is not permitted, except that material may be duplicated by you for your research use or educational purposes in electronic or print form. You must obtain permission for any other use. Electronic or print copies may not be offered, whether for sale or otherwise to anyone who is not an authorised user.







