

**This is an electronic reprint of the original article.
This reprint *may differ* from the original in pagination and typographic detail.**

Author(s): Auranen, Kalle; Uusitalo, Juha; Juutinen, Sakari; Jakobsson, Ulrika; Grahn, Tuomas; Greenlees, Paul; Hauschild, Karl; Herzan, Andrej; Julin, Rauno; Konki, Joonas; Leino, Matti; Pakarinen, Janne; Partanen, Jari; Peura, Pauli; Rahkila, Panu; Ruotsalainen, Panu; Sandzelius, Mikael; Sarén, Jan; Scholey, Catherine; Sorri, Juha; Stolze, Sanna

Title: Spectroscopy of At 201 including the observation of a shears band and the 29/2 + isomeric state

Year: 2015

Version:

Please cite the original version:

Auranen, K., Uusitalo, J., Juutinen, S., Jakobsson, U., Grahn, T., Greenlees, P., . . .
Stolze, S. (2015). Spectroscopy of At 201 including the observation of a shears band
and the 29/2 + isomeric state. *Physical Review C*, 91 (2), 024324 .
doi:10.1103/PhysRevC.91.024324

All material supplied via JYX is protected by copyright and other intellectual property rights, and duplication or sale of all or part of any of the repository collections is not permitted, except that material may be duplicated by you for your research use or educational purposes in electronic or print form. You must obtain permission for any other use. Electronic or print copies may not be offered, whether for sale or otherwise to anyone who is not an authorised user.

Erratum: Spectroscopy of ^{201}At including the observation of a shears band and the $29/2^+$ isomeric state [Phys. Rev. C **91**, 024324 (2015)]

K. Auranen, J. Uusitalo, S. Juutinen, U. Jakobsson, T. Grahn, P. T. Greenlees, K. Hauschild, A. Herzáň, R. Julin, J. Konki, M. Leino, J. Pakarinen, J. Partanen, P. Peura, P. Rahkila, P. Ruotsalainen, M. Sandzelius, J. Sarén, C. Scholey, J. Sorri, and S. Stolze

(Received 7 May 2015; published 9 September 2015)

DOI: [10.1103/PhysRevC.92.039901](https://doi.org/10.1103/PhysRevC.92.039901) PACS number(s): 23.20.Lv, 23.20.Nx, 23.35.+g, 27.80.+w, 99.10.Cd

The $B(E2)$ value of the 269-keV $E2$ transition and the $B(E3)$ value of the 339-keV $E3$ transition, each depopulating the 2319-keV isomeric state, contain errors owing to a mistake in the calculation of the values. The correct values are $B(E2) = 1.26(4) \times 10^{-3}$ and $B(E3) = 21(3)$ W.u. [replacing the incorrect values of $1.33(4) \times 10^{-3}$ and $11(2)$ W.u., respectively].

In addition, Table II of the original article contains four typographical errors: The spin values were interchanged between rows 2 and 4. The correct table is presented here. These changes do not affect the conclusions reached in the article.

TABLE II. γ -ray transitions associated with the shears band. I_γ normalized such that $I_\gamma(635 \text{ keV}) = 100$. Intensities and energies are deduced from the sum of gates (746- and 427-keV transitions) γ - γ data.

E_γ (keV)	I_γ	A_2	I_i^π (\hbar)	I_f^π (\hbar)	$B(M1)/B(E2)$ (μ_N^2/e^2b^2)
145.0(4)	3.7(3)	-0.5(2)	(25/2 ⁻)	(23/2 ⁻)	
197.9(4)	1.6(2)	-0.80(12)	(35/2 ⁻)	(33/2 ⁻)	>8
244.4(4)	8.3(5)	-0.59(9)	(27/2 ⁻)	(25/2 ⁻)	>30
272.3(4)	3.5(3)	-0.45(11)	(33/2 ⁻)	(31/2 ⁻)	>35
286.9(4)	6.6(4)	-0.47(3)	(29/2 ⁻)	(27/2 ⁻)	>25
317.3(4)	6.7(4)	-0.81(9)	(31/2 ⁻)	(29/2 ⁻)	>30
335.0(4)	1.5(2)	-0.66(4)	(37/2 ⁻)	(35/2 ⁻)	>2
1068.9(4)	4.5(3)	-0.47(5)	(23/2 ⁻)	21/2 ⁺	