

**This is an electronic reprint of the original article.
This reprint *may differ* from the original in pagination and typographic detail.**

Author(s): Kangas, Reeta; Laakkonen, Eija; Kovanen, Vuokko

Title: Aging and systemic hormonal status affects the circulating miR-21, miR-146a and FasL levels

Year: 2015

Version:

Please cite the original version:

Kangas, R., Laakkonen, E., & Kovanen, V. (2015). Aging and systemic hormonal status affects the circulating miR-21, miR-146a and FasL levels. *RNA and Disease*, 2(2), Article e552. <https://doi.org/10.14800/rd.552>

All material supplied via JYX is protected by copyright and other intellectual property rights, and duplication or sale of all or part of any of the repository collections is not permitted, except that material may be duplicated by you for your research use or educational purposes in electronic or print form. You must obtain permission for any other use. Electronic or print copies may not be offered, whether for sale or otherwise to anyone who is not an authorised user.

RESEARCH HIGHLIGHT

Aging and systemic hormonal status affects the circulating miR-21, miR-146a and FasL levels

R. Kangas, E. Pöllänen, V. Kovanen

Gerontology Research Center, Department of Health Sciences, University of Jyväskylä, Finland

Correspondence: Reeta Kangas

E-mail: reeta.kangas@jyu.fi

Received: January 28, 2015

Published online: February 22, 2015

MicroRNAs are small molecules, found in all cell types and body fluids, which most commonly affect negatively to gene expressions by translational repression. Their role in various physiological conditions and diseases has been emphasized during the last twenty years. In our recent studies with postmenopausal monozygotic twin sisters (n=11), we have investigated how different systemic hormonal status affects the levels of specific circulating microRNAs and other molecules related to inflammation and apoptosis, both processes associated with aging. Our results have shown that the systemic levels of miR-21, miR-146a and Fas ligand are lower within the postmenopausal women who are using estrogen-based hormonal replacement (HRT), compared to their non-using co-twins (p=0.018, p=0.039, p=0.021, respectively). To get further knowledge about the aging effect, we also measured the same markers from the premenopausal control women (n=8), with natural hormonal status, and found out that the inflammatory profile was healthier among the young women and that the serum miR-21 profile was more similar with the HRT users than non-users, and miR-146 and FasL profile more similar to non-users. These findings demonstrate that HRT has effects on the circulating inflammation related regulatory molecules. Whether we can state that the effects are clearly positive or negative, needs further investigations and understanding of the regulatory system.

To cite this article: R.Kangas, et al. Aging and systemic hormonal status affects the circulating miR-21, miR-146a and FasL levels. RNA Dis 2015; 2: e552. doi: 10.14800/rd.552.

Aging is a process influenced by the combination of genetics and environmental factors. During the last decades, epigenetics, including microRNA (miR) regulation, has revolutionized our understanding of how our genes are being switched on and off. It is known that a single miR can regulate several biological processes as well as a group of miRs are affecting the expression of a single gene. To make it more complex, it is not only about how miRs regulate different processes: In addition, also miRs themselves are regulated by various molecules such as sex steroid hormones. Interest towards the interplay between hormonal regulation and different stages of miR maturation and targeting has emerged among many research groups in the recent years^[1]. The studies have shown that the regulatory system among these molecules is bi-directional, meaning both hormonal

signaling and miR biogenesis and function are affecting each other. It has also been stated that miRs deliver their messages similarly to hormones themselves. In our recent study we demonstrated that circulating miR-21 (onco-miR) and miR-146a (inflamma-miR) levels differ between postmenopausal monozygotic twin sisters whose hormonal status differs from each other, especially what comes to systemic estrogen concentrations^[2]. We have also shown with the same data, that HRT enhances the IGF-1 signaling by lowering the levels of miR-223 and miR-182 in the skeletal muscle^[3]. In addition, we have reported that the HRT users possess better muscle quality properties than the non-user twin sisters^[4]. Nothnick et al^[5] have also studied the interplay of hormones and miR regulation within the uterus of ovariectomized mice by demonstrating that steroid

hormones not only affect the miR expression at the transcription level but also the expression of the molecules, Exportin-5 and Dicer1, involved in miR maturation process. Several studies are focusing on the associations with miR levels and cancers associated with tissues sensitive to estrogens and androgens^[6-8].

One of the most intriguing questions in miR research is whether we should consider a change in a single miR level significant or should we examine the miR profile from a specific tissue as a whole. Many studies have shown that changes in specific miR levels are characteristics to certain diseases such as various cancers, autoimmune disorders and neurological diseases^[9]. Another set of studies has shown changes in miR profiles in different conditions^[10]. As we showed in our recent twin paper, the levels of specific circulating miRs are partly genetically regulated (miR-21: $r=0.66$, miR-146a: $r=0.38$)^[2]. Due to that, setting common reference values for normal and abnormal miR levels for the clinical use, might be challenging.

In our ongoing studies we are focusing on the serum levels of the specific miRs and FasL in another longitudinal data set consisting of aged male sprint athletes^[11]. It seems like an aging effect can be detected in this unique study population what comes to the studied molecules. In addition, our miR work continues with the already discussed twin data, in order to get a better picture of the entire miR profile of the serum samples.

Overall, it is not so obvious, whether we express certain inherited properties or not, it is all about the interplay between the genes and the environment and the bunch of molecules working in between. MicroRNAs seem to be one of the sensitive mechanisms affecting the fine-tuning of the gene expression. Due to their already detected variation in different conditions and diseases and high stability in the biological samples, miRs are promising tools to be used in diagnostics and therapeutics. However, in order to generalize and take the miR analysis into official clinical use, the methods need to be standardized.

Acknowledgements

We thank warmly our international collaborators at the Department of Clinical and Molecular Sciences and at the Department of Clinical Pathology and Innovative Therapy at the Università Polytechnica delle Marche, Ancona, Italy and at the Department of Experimental, Diagnostic and Specialty Medicine and Interdepartmental Center Galvani "CIG" at the University of Bologna, Bologna, Italy, and our national collaborators at the Institute for Molecular Medicine and National Institute for Health and Welfare at the University of

Helsinki, Department of Microbiology and Immunology at the University of Tampere and Gerontology Research Center at the University of Tampere and Jyväskylä, whose contribution to the original article was essential (see ref 2). This research has been supported by the Finnish Cultural Foundation (R.K), Department of Health Sciences (University of Jyväskylä), Sohlberg Foundation (E.P), Academy of Finland (grants # 265240 and 114310 to Jaakko Kaprio and V.K, respectively) and EFCP7 Collaborative project MYOAGE (#GA-223576, Sarianna Sipilä).

References

1. Fletcher CE, Dart DA, Bevan CL. Interplay between steroid signalling and microRNAs: Implications for hormone-dependent cancers. *Endocr Relat Cancer* 2014; 21:R409-29.
2. Kangas R, Pollanen E, Rippo MR, Lanzarini C, Prattichizzo F, Niskala P, et al. Circulating miR-21, miR-146a and fas ligand respond to postmenopausal estrogen-based hormone replacement therapy - A study with monozygotic twin pairs. *Mech Ageing Dev* 2014; 143-144:1-8.
3. Olivieri F, Ahtiainen M, Lazzarini R, Pollanen E, Capri M, Lorenzi M, et al. Hormone replacement therapy enhances IGF-1 signaling in skeletal muscle by diminishing miR-182 and miR-223 expressions: A study on postmenopausal monozygotic twin pairs. *Aging Cell* 2014; 13:850-61.
4. Ronkainen PH, Kovanen V, Alen M, Pollanen E, Palonen EM, Ankarberg-Lindgren C, et al. Postmenopausal hormone replacement therapy modifies skeletal muscle composition and function: A study with monozygotic twin pairs. *J Appl Physiol* (1985) 2009; 107:25-33.
5. Nothnick WB, Healy C, Hong X. Steroidal regulation of uterine miRNAs is associated with modulation of the miRNA biogenesis components exportin-5 and Dicer1. *Endocrine* 2010; 37:265-73.
6. Kumar S, Keerthana R, Pazhanimuthu A, Perumal P. Overexpression of circulating miRNA-21 and miRNA-146a in plasma samples of breast cancer patients. *Indian J Biochem Biophys* 2013; 50:210-4.
7. Waltering KK, Porkka KP, Jalava SE, Urbanucci A, Kohonen PJ, Latonen LM, et al. Androgen regulation of micro-RNAs in prostate cancer. *Prostate* 2011; 71:604-1.
8. Liu J, Sun H, Wang X, Yu Q, Li S, Yu X, et al. Increased exosomal microRNA-21 and microRNA-146a levels in the cervicovaginal lavage specimens of patients with cervical cancer. *Int J Mol Sci* 2014; 15:758-73.
9. Wu K, Li L, Li S. Circulating microRNA-21 as a biomarker for the detection of various carcinomas: An updated meta-analysis based on 36 studies. *Tumour Biol* 2014;doi:10.1007/s13277-014-2803-2
10. Di Leva G, Croce CM. miRNA profiling of cancer. *Curr Opin Genet Dev* 2013; 23:3-11.
11. Korhonen MT. Effects of aging and training on sprint performance, muscle structure and contractile function in athletes [dissertation]. *Studies in Sport, Physical Education and Health*, 137. University of Jyväskylä; 2009.