

**This is an electronic reprint of the original article.
This reprint *may differ* from the original in pagination and typographic detail.**

Author(s): Eklund, Daniela; Schumann, Moritz; Kraemer, William J.; Izquierdo, Mikel; Taipale, Ritva; Häkkinen, Keijo

Title: Acute Endocrine and Force Responses and Long-Term Adaptations to Same-Session Combined Strength and Endurance Training in Women

Year: 2016

Version:

Please cite the original version:

Eklund, D., Schumann, M., Kraemer, W. J., Izquierdo, M., Taipale, R., & Häkkinen, K. (2016). Acute Endocrine and Force Responses and Long-Term Adaptations to Same-Session Combined Strength and Endurance Training in Women. *Journal of Strength and Conditioning Research*, 30(1), 164-175.
<https://doi.org/10.1519/JSC.0000000000001022>

All material supplied via JYX is protected by copyright and other intellectual property rights, and duplication or sale of all or part of any of the repository collections is not permitted, except that material may be duplicated by you for your research use or educational purposes in electronic or print form. You must obtain permission for any other use. Electronic or print copies may not be offered, whether for sale or otherwise to anyone who is not an authorised user.

Title: Acute endocrine and force responses and long-term adaptations to same-session combined strength and endurance training in women

Journal of Strength and Conditioning Research

Running head: Order effect in women: loading and training

Eklund, Daniela¹, Schumann, Moritz¹, Kraemer, William J.², Izquierdo, Mikel³, Taipale, Ritva S¹, Häkkinen, Keijo¹. 2014.

¹Department of Biology of Physical Activity, University of Jyväskylä, Finland; ²Department of Human Sciences, The Ohio State University, OH, USA; ³Department of Health Sciences, Public University of Navarra, Spain

Corresponding author:

Eklund, Daniela

Department of Biology of Physical Activity, University of Jyväskylä

P.O. Box 35, 40014 University of Jyväskylä, Finland

Tel.: +358408054591, Fax: +358142602071

daniela eklund@jyu.fi

Co-authors:

Schumann, Moritz.

Department of Biology of Physical Activity, University of Jyväskylä

P.O. Box 35, 40014 University of Jyväskylä, Finland

Tel.: +358408053757, Fax: +358142602071

moritz.schumann@jyu.fi

Kraemer, William J.

Department of Human Sciences, The Ohio State University

A054 PAES Building, 305 W 17th Avenue, OH 43210, USA

Tel.: +1 614-688-2354, Fax.: +1 614-688-3432

kraemer.44@osu.edu

Izquierdo, Mikel

Department of Health Sciences, Public University of Navarra

Campus of Tudela, 31500-Tudela Spain

tel.: +34948417876

mikel.izquierdo@gmail.com

Taipale, Ritva S

Department of Biology of Physical Activity

P.O. Box 35, 40014 University of Jyväskylä, Finland

Tel.: +358408053731, Fax: +358142602071

ritva.taipale@jyu.fi

Häkkinen, Keijo

Department of Biology of Physical Activity

P.O. Box 35, 40014 University of Jyväskylä, Finland

Tel.: +358504285289, Fax: +358142602071

keijo.hakkinen@jyu.fi

1

2 **ABSTRACT**

3 This study examined acute hormone and force responses as well as strength and endurance
4 performance and muscle hypertrophy before and after 24 weeks of same-session combined
5 strength and endurance training in previously untrained women. Subjects were assigned one
6 of two training orders: endurance preceding strength (E+S, n=15) or vice versa (S+E, n=14).
7 Acute force and hormone responses to a combined loading (continuous cycling and a leg
8 press protocol in the assigned order) were measured. Additionally, leg press one-repetition
9 maximum (1RM), maximal workload during cycling (W_{max}) and muscle cross-sectional-area
10 (CSA) were assessed. Loading-induced decreases in force were significant ($p<0.01-0.001$)
11 before (E+S $20\pm 11\%$, S+E $18\pm 5\%$) and after (E+S $24\pm 6\%$, S+E $22\pm 8\%$) training. Recovery
12 was completed within 24h in both groups. The acute growth hormone response was
13 significantly ($p<0.001$) higher after S+E than E+S at both Week 0 and Week 24. Testosterone
14 was significantly ($p<0.001$) elevated only after the S+E loading at Week 24, but was not
15 significantly different from E+S. Both groups significantly ($p<0.001$) improved 1RM (E+S
16 $13\pm 12\%$, S+E $16\pm 10\%$), W_{max} (E+S $21\pm 10\%$, S+E $16\pm 12\%$) and CSA (E+S $15\pm 10\%$, S+E
17 $11\pm 8\%$). This study showed that the acute growth hormone response to combined endurance
18 and strength loadings was significantly larger in S+E compared to E+S both before and after
19 24 weeks of same-session combined training. Strength and endurance performance and CSA
20 increased to similar extents in both groups during 24 weeks despite differences in the kinetics
21 of growth hormone. Previously untrained women can improve performance and increase
22 muscle CSA utilizing either exercise order.

23

24 **Keywords:** concurrent training, testosterone, growth hormone, performance adaptations,
25 order effect

1

2 INTRODUCTION

3 It has been well established in male populations, that metabolically demanding resistance
4 exercise elicits large acute elevations of serum testosterone, growth hormone and cortisol (18,
5 25, 35). These acute anabolic responses in men have in some studies been linked to long-term
6 physiological adaptations such as gains in muscle strength and hypertrophy (22, 31, 41),
7 while in other studies this phenomenon has not been found (47). Even though the magnitude
8 of exercise-induced elevations in hormonal concentrations may not be correlated to long-term
9 adaptations per se, the hormonal responses are known to create the metabolic environment
10 involved in tissue remodelling (e.g. 19, 45).

11 The hormonal responses to resistance exercise in women are similar to those of men, albeit
12 smaller in magnitude. Typically, only minor or no acute elevations in testosterone
13 concentrations are reported in women following strenuous resistance exercise protocols (e.g.
14 (9, 19, 29). These limited magnitudes of testosterone responses are likely related to the
15 intensity of exercise and amount of activated muscle mass (10, 26, 29, 32), but may possibly
16 be counterbalanced by acute growth hormone release to meet the anabolic needs of resistance
17 exercise sessions (25).

18 When combining strength (S) and endurance (E) into the same training session, the question
19 arises regarding which exercise order (i.e. E+S or S+E) should be preferred. The acute effect
20 of the exercise order on circulating hormones is of relevance considering the possible
21 implications for long-term adaptations. As data from female populations is scarce, current
22 knowledge of the hormonal responses to combined loadings relies mainly on findings from
23 men. Based on earlier reports, a bout of endurance exercise seems to blunt the growth
24 hormone response to subsequent resistance exercise, thus resulting in lower post-exercise
25 concentrations than in the opposite order (16, 39). The findings regarding cortisol and
26 testosterone (4, 37, 39) are less conclusive and may be related to the intensity or volume of

1 the utilized exercise protocols or the training status of the subjects (4, 16, 39). Since most of
2 these studies have incorporated a cross-sectional design, possible changes in the exercise-
3 induced hormonal responses are not well understood. A previous study by our group noted
4 that the S+E order could initially result in faster recovery of testosterone in men in
5 comparison to the opposite order, possibly indicating different recovery needs (39). However,
6 this difference was found to diminish with prolonged training, and did not influence the long-
7 term strength gains. Furthermore, even though endurance exercise acutely impairs subsequent
8 force production (12, 28) and has been suggested to attenuate strength development following
9 prolonged E+S training (3), recent reports from both men and women show similar strength
10 gains following long-term training (11, 14).

11 Despite a growing interest towards research regarding concurrent training in female
12 populations (e.g. 11, 40), there is currently paucity in the knowledge regarding hormonal
13 responses to combined exercise sessions in women. Although strength and endurance
14 performance as well as lean mass are likely to increase to a similar extent following training
15 in either order (11), the effects of prolonged training on exercise induced hormonal responses
16 and the relevance for training adaptations has not been elucidated. Thus, the main purpose of
17 the present study was to investigate the influence of the exercise order of combined strength
18 and endurance loadings on acute hormone and force responses both before and after 24 weeks
19 of combined training. A secondary purpose was to investigate whether the acute exercise-
20 induced changes in hormone concentrations are associated with long-term training
21 adaptations in strength and endurance performance or muscle cross-sectional area.

22

23 **METHODS**

24 **Experimental approach to the problem**

1 In order to examine the effect of prolonged training on acute exercise-induced force and
2 hormone responses to combined E+S or S+E loadings and the chronic adaptations in strength
3 and endurance performance and muscle cross-sectional area, a 24-week training intervention
4 was conducted. As the focus of this study was to compare training-induced adaptations in
5 acute loading responses, a cross-over design was not used and the subjects performed the
6 experimental loading in their assigned loading order only. The acute loading responses and
7 long-term adaptations in strength and endurance performance were determined before (Week
8 0) and after (Week 24) the intervention (Figure 1).

9 **Subjects**

10 Twenty-nine women participated in the present study. Recruitment was conducted by several
11 public postings. Subjects were 1) recreationally physically active but without systematic
12 strength or endurance training for at least 1 year prior to participation, 2) below a body mass
13 index of 30 m²/kg, 3) non-smokers 4) free from chronic illnesses and injuries and 5) not
14 pregnant or lactating. A resting ECG screening was approved by a cardiologist. The subjects
15 were informed about the study design, measurements and procedures. The subjects were
16 matched by physical fitness at baseline into two training groups: endurance preceding
17 strength (E+S, n=15, 29.1 ± 5.6 years, 168 ± 7 cm, 67 ± 10 kg and BMI 23.7 ± 3.3 kg/m²) and
18 strength preceding endurance (S+E, n=14, 28.9 ± 4.4 years, 164 ± 5 cm, 62.4 ± 8 kg and BMI
19 23.2 ± 3.4 kg/m²). Due to organizational constraints, acute loading responses were assessed
20 from 23 subjects (E+S n=12, S+E n=11), while changes in strength and endurance
21 performance as well as muscle cross-sectional area were assessed for all subjects. The study
22 received ethical approval from the Ethics Committee of the University of Jyväskylä, Finland,
23 and was conducted in accordance with the Declaration of Helsinki. After written and verbal
24 information about the study and its procedures had been provided, written informed consent
25 was obtained from all subjects.

26 **Procedures**

1 Prior to the start of the measurements and training, subjects reported to the laboratory for a
2 familiarization session during which the strength measurements were practiced and the
3 equipment was adjusted to the specifics of the individual. Subjects wore the same shoes for
4 all measurements and loading sessions. Blood sampling and all physical tests were conducted
5 at the same time of day \pm 1 h throughout the study. The measurements of maximal strength
6 and endurance performance were separated from each other and the loading measurements by
7 at least two days. The last training session of the 24-week training intervention was separated
8 from the following basal measurements by 2-4 days of rest. Nutritional information according
9 to the national guidelines was provided before the start of the study and the subjects were
10 asked to keep their energy intake constant throughout the intervention. Ingestion of caffeine
11 and alcohol was not allowed 12 h and 24 h (respectively) prior to the measurements and
12 subjects were required to keep the nutritional intake prior to the measurements similar at
13 Weeks 0 and 24.

14 As recent reports have shown minimal influence of the menstrual cycle phase on anabolic
15 hormone responses to strength (43) and endurance exercise (33), the measurements were
16 conducted across several phases of the menstrual cycle. Four subjects from the E+S and three
17 subjects from the S+E groups reported oral contraceptive use.

18

19 *** Figure 1 near here ***

20

21 **Basal measurements**

22 *Strength.* Maximal bilateral dynamic leg press one-repetition maximum (1RM) was measured
23 using a David 210 weight stack horizontal leg press device (David Health Solutions Ltd.,
24 Helsinki, Finland). Three warm-up sets (5 x 70-75%, 3 x 80-85% and 2 x 90-95% of
25 estimated 1RM) with 1 min of rest between sets were performed before the 1 RM trials. Upon

1 verbal instruction, subjects performed a full leg extension (knee angle 180°) from a starting
2 knee angle of below 60° (58°±2°). After each successful completion the load was increased.
3 Subjects were allowed a maximum of five trials. The trial with the highest completed load
4 was accepted as the 1 RM.

5 *Endurance.* The maximal endurance test was conducted on a cycle ergometer (Ergometrics
6 800, Ergoline, Bitz, Germany) using a graded exercise protocol. The test was initiated at 50
7 watts (W) for all subjects with 25W increments applied every 2 min until volitional
8 exhaustion. Maximal workload (W_{\max}) was calculated as $W_{\max} = W_{\text{com}} + (t/120) * 25$ (39), where
9 W_{com} represents the load of the last completed and t the time of the last incomplete stage.
10 Aerobic and anaerobic thresholds were determined for each subject based on the points of
11 deflection in the curves of ventilation, oxygen consumption, production of carbon dioxide and
12 blood lactate (2).

13 *Muscle cross-sectional area.* Cross-sectional area (CSA) of the vastus lateralis muscle of the
14 right limb was measured using a B-mode axial-plane ultrasound device (SSD-a10, Aloka,
15 Tokyo, Japan) and a panoramic imaging technique (1). Images were taken at 50% and 70% of
16 the femur length, i.e. the distance between the greater trochanter and the joint space on the
17 lateral side of the knee. The 10 MHz linear-array probe was moved across the thigh from the
18 medial to the lateral side with the subject lying in a supine position. Leg position was fixed
19 using a Styrofoam support. Lines perpendicular to the measurement table were drawn across
20 the thigh to ensure that the probe was moved in a straight line. Three images were taken at
21 both 50% and 70% of the muscle length. Images were analyzed using ImageJ -software
22 version 1.44 (National Institute of Health, USA) by manually marking the outlines of the
23 muscles onto the image. The mean of the two closest values of 50% and 70%, respectively,
24 were averaged and used in the statistical analyses to assess total CSA. The reproducibility of
25 the measurement has been reported earlier by our research group (38).

26 **Experimental loading protocol**

1 The experimental loading was intended to reflect the content of the 24-week training
2 program, which was designed to reflect the exercise recommendations for physically active
3 individuals (42). The loading consisted of both endurance cycling and a leg press protocol.
4 Loadings were conducted for each subject at the same time of day (± 1 h) at Week 0 and 24
5 and were performed in the order specific to the training. Measurements of force and hormone
6 responses during the loading were conducted before the initiation of the loading (“Pre”), after
7 the first part (“Mid”: E or S, respective to order) and after the complete loading (“Post”).
8 Recovery was monitored 24 ± 1 h and 48 ± 1 h after the cessation of exercise (“24h” and “48h”,
9 respectively).

10 Subjects were verbally encouraged throughout the loadings. Proper hydration on the day
11 preceding the loading was encouraged. Consumption of 0.2 l of water was allowed between
12 the two loading modes, after the “Mid” blood sample was taken.

13 *Strength loading.* A David 210 weight-stack horizontal leg press (David Health Solutions
14 Ltd., Helsinki, Finland) was used to conduct the strength loading. A detachable handle was
15 available for assisting if necessary. The loading consisted of three protocols typically used in
16 training for explosive strength (3x10 repetitions at 40% 1RM with 3 min rest between sets),
17 maximal strength (4x3 repetitions at 75-90% 1RM with 3 min rest between sets) and muscle
18 hypertrophy (4x10 repetitions at 75-80% 1RM with 2 min rest between sets). Loads were
19 calculated from the 1 RM obtained during the basal measurements. Additional resistance was
20 added to at least one maximal and one hypertrophic set in order to complete a true repetition
21 maximum and standardize the loading conditions. In the explosive sets, subjects were
22 instructed to perform the concentric phase as fast as possible and the eccentric phase in a
23 controlled manner, without pausing between repetitions. For the hypertrophy and maximal
24 sets, subjects were instructed to fully extend their legs without locking their knees and to keep
25 an even pace throughout the movement.

1 *Endurance loading.* The endurance loading consisted of 30 minutes of continuous cycling at
2 an intensity of 65% W_{\max} (39) on a Monark cycling ergometer (Ergomedic 839E, Monark
3 Exercise AB, Vansbro, Sweden) equipped with electric resistance. The intensity was
4 calculated based on W_{\max} from the basal measurement. Subjects were instructed to keep the
5 pedalling pace at 70 revolutions per minute (rpm). The rpm was visible to the subjects
6 throughout the loading and was additionally monitored by a member of staff. In case of the
7 rpm dropping below 65 with the subject unable to increase it, the workload was lowered by
8 15W. If the subject was unable to keep up the pace for a full minute after the reduction, the
9 workload was further reduced by 15W. If necessary, the procedure was repeated until the
10 subject was able to keep up the required pace and complete the loading.

11 **Measurements during the experimental loading**

12 *Isometric force production.* Maximal isometric force (MVC) was measured on a leg press
13 device (Department of Biology of Physical Activity, University of Jyväskylä, Jyväskylä,
14 Finland) with a knee joint angle of 107° (180° representing full extension) (20). The greater
15 trochanter of the femur and lateral malleolus of the ankle of the right limb were used as
16 anatomical reference points.

17 Subjects were instructed to perform an isometric bilateral leg press action as rapidly as
18 possible with the aim of reaching the maximum force at the beginning of the trial and
19 maintaining it for a duration of approximately 3s. At Pre, 24h and 48h subjects were allowed
20 to perform three trials with 1 min rest between. At Mid and Post subjects immediately
21 proceeded to the measurement and performed two trials with only 10s rest between for the
22 purpose of recording exercise-induced fatigue. The trial with the highest force was selected
23 for analysis. Force signals were recorded with Signal 2.16 software (CED, Cambridge, UK),
24 sampled at 2000 Hz and processed with a low-pass filter of 20 Hz. Trials were analyzed for
25 MVC and average force produced between 0 and 500 ms (MVC_{500}).

1 *Blood samples.* To determine blood lactate concentrations, capillary blood samples were
2 taken from the fingertip at Pre, Mid and Post into a reaction tube containing an anti-coagulant
3 and hemolyzing agent. The samples were analyzed using a Biosen lactate analyzer (S-line
4 Lab+ EKF, Magdeburg, Germany). In addition, venous blood samples were drawn at Pre,
5 Mid, Post, 24h and 48h for determination of total testosterone (T), cortisol (C) and growth
6 hormone (GH, 22 kDa) concentrations. Resting concentrations of the same hormones as well
7 as sex-hormone binding globulin (SHBG) and T/C and T/SHBG -ratios were determined on
8 the morning of the loading (7:00-9:00 am) in a fasted state. Samples were drawn by a
9 laboratory technician from the antecubital vein into a serum tube (Venosafe, Terum Medical
10 Co, Leuven, Belgium). Samples were centrifuged for 10 min at 3500 rpm, after which serum
11 was removed and frozen until analysed. The hormones were analysed with a chemical
12 luminescence technique (Immulite 1000, Siemens, New York, USA) using hormone-specific
13 immune-assay kits (Siemens, New York, USA). Creatine kinase (CK) was analysed using
14 chemical analysis (KoneLab 20 XTi, Thermo Fisher Scientific Oy, Vantaa, Finland).
15 Sensitivities for T, C, GH, SHBG and CK were 0.5 nmol l^{-1} , 5.5 nmol l^{-1} , 0.03 mIU l^{-1} and
16 0.02 nmol l^{-1} and 0.7 mIU l^{-1} , respectively. Intra-assay coefficients of variation for T, C, GH
17 and SHBG were 9.8 ± 3.9 , 7.1 ± 1.1 , 6.0 ± 0.5 , $3.1 \pm 1.3\%$, and $1.5 \pm 0.7\%$, respectively.
18 Inter-assay coefficients of variation for T, C, GH, SHBG and CK were 12.0 ± 6.3 , 7.9 ± 1.2 ,
19 5.8 ± 0.3 , $5.0 \pm 1.0\%$ and $3.6 \pm 0.8\%$, respectively. Serum hormone concentrations were not
20 corrected for changes in plasma volume. To monitor haemoconcentration (13), haemoglobin
21 (HGB) and haematocrit (HCR) were analysed with Sysmex KX 21 N (Sysmex America Inc.,
22 Mundelein, IL, USA) automated haematology analyser with a cyanide-free and cumulative
23 pulse height detection method, respectively.

24 **Training**

25 The training program has been described in detail previously (14). Briefly, the training was
26 aimed to reflect recommendations for physically active individuals (e.g. 42) and was targeted
27 at improving both maximal strength and endurance performance. During the first 12 weeks,

1 the subjects completed two weekly sessions of [1E+1S] or [1S+1E] (respective to the
2 assigned training order) and five sessions per two weeks (5x [1E+1S] or [1S+1E]) during
3 weeks 13-24. Time between training modes was 5-10 min and recovery time between training
4 sessions 48-72 h. Training sessions were supervised by research staff. Maintenance of normal
5 daily activity was encouraged.

6 Strength training mainly targeted knee extensors and flexors as well as hip extensors.
7 Exercises consisted of horizontal leg press, seated hamstring curls and seated knee extensions.
8 The program was initiated with the exercises performed in a circuit (2-4 sets of 15-20
9 repetitions with up to 60% of 1RM) and continued through hypertrophy-inducing training (2-
10 5 x 8-12 at 80-85% of 1RM, 1-2 min rest) towards maximal strength training (2-5 x 3-5 at 85-
11 95% of 1RM, 3-4 min rest). A similar pattern of periodization was used for the upper body.
12 Dumbbells and cable pulley machines were used for the upper body exercises and both
13 machines and body weight for exercises of the trunk. The periodization was repeated during
14 weeks 13-24 with increased training intensity and volume. The duration of each strength
15 session was 50-60 min.

16 Endurance training sessions were performed on a cycle ergometer. Training intensities were
17 controlled by heart rate zones corresponding to the threshold values of aerobic and anaerobic
18 thresholds. Training consisted of 30-50 min continuous cycling near the AT (weeks 1-7 and
19 13-16), including interval training at and above the anaerobic threshold from weeks 8 and 17
20 onwards. The interval sessions were initiated and ended with 10-15 minute bouts below the
21 aerobic threshold with 5-minute altering bouts on the anaerobic threshold and below the
22 aerobic threshold in between.

23

24 **Statistical analysis.** Data are presented as means±SD. Statistical analysis for changes during
25 the experimental loadings at Week 0 and 24 was performed using a five-level ANCOVA (i.e.
26 Pre, Mid, Post, 24h and 48h) with absolute values for within-group changes and values
27 relative to Pre for between-group differences, with Pre-values used as covariates. As GH

1 during the experimental loading and basal SHBG were non-normally distributed even after a
2 log transformation, non-parametric statistics were used both for the within-group changes
3 (Wilcoxon signed-rank test) and between-group comparisons (Mann-Whitney U-test). For the
4 non-parametric tests a Bonferroni adjustment was applied by multiplying the pairwise p-
5 values with the number of comparisons. To compare the experimental loading-induced
6 within-group changes (i.e. Mid, Post, 24h and 48h) across 24 weeks, paired-samples t-tests
7 were applied for each measurement point.

8 Training induced changes in basal hormones and basal measurements of 1 RM, W_{\max} and
9 CSA were analyzed with a two-way ANCOVA with baseline-values used as covariates, and
10 between-group differences with an independent-samples t-test. The individual ratios of
11 changes in 1 RM and W_{\max} were calculated as the percentage change in 1 RM divided by the
12 percentage change in W_{\max} .

13 Reported effect sizes (ES) are Cohen's d except for non-normally distributed data, where ES
14 was defined as $Z\text{-score}/\sqrt{n}$. Associations between the exercise-induced changes in serum
15 hormone concentrations and training-induced adaptations were examined using bivariate
16 Pearson correlation coefficient for normally and Spearman's rank correlation coefficient for
17 non-normally distributed data. A trend was accepted for p-values <0.06 .

18

19 **RESULTS**

20 Training adherence was 99% in both E+S and S+E. All subjects completed at least 90% of the
21 training sessions.

22

23 **Acute loading responses at week 0**

24 No significant changes in body weight ($-0.3\pm 0.3\%$), HGB ($+3.6\pm 3.0\%$) or HCR ($+2.6\pm 2.9\%$)
25 were observed during the loading.

1 MVC decreased significantly during the loading in both groups by Mid (E+S $-18\pm 13\%$ from
 2 1740 ± 235 N, $P<0.01$, $ES=-1.305$; S+E $-17\pm 7\%$ from 1810 ± 633 N, $P<0.01$, $ES=-0.515$) and
 3 by Post (E+S $-20\pm 11\%$, $P<0.001$, $ES=-1.587$; S+E $-18\pm 5\%$ $P<0.001$, $ES=-0.532$) (Figure 2).
 4 MVC_{500} decreased significantly in E+S by mid (E+S $-18\pm 12\%$, $P<0.01$, $ES=-1.24$) and post ($-$
 5 $20\pm 14\%$, $P<0.01$, $ES=-1.15$) and for S+E by post ($-16\pm 7\%$, $P<0.05$, $ES=-0.611$). No
 6 significant differences of MVC or MVC_{500} to Pre were observed for either group at 24h or
 7 48h.

8

9

*** Figure 2 near here ***

10

11 A significant increase in T was observed in E+S at Mid (from 0.5 ± 0.4 to 8.9 ± 1.1 $\text{nmol}\cdot\text{l}^{-1}$,
 12 $P<0.05$, $ES=0.513$) (Figure 3). C remained statistically unaltered throughout the loading for
 13 both groups (Table 1). A trend ($P=0.051$, $ES=0.256$) was observed in C for E+S at Mid. At
 14 24h and 48h, C was significantly lowered from Pre for S+E (24h: $-29\pm 23\%$, $P<0.01$, $ES=-$
 15 1.53 and 48h: $-29\pm 14\%$ $P<0.01$, $ES=-1.70$). A 5.6-fold increase from pre in GH was observed
 16 at Mid for E+S ($P<0.05$, $ES=0.888$) and a 5.2-fold increase at Post for S+E ($P<0.05$,
 17 $ES=0.830$) (Figure 4). A trend was found in S+E from Mid to Post ($P=0.055$, $ES=0.402$). The
 18 change from Mid to Post was significantly different between groups ($P<0.001$, $ES=0.886$).

19 Blood lactate increased significantly in both groups by Mid (E+S by 4.2-fold $P<0.001$,
 20 $ES=4.80$; S+E by 4.0-fold, $P<0.001$, $ES=2.8$) and Post (E+S by 5.0-fold $P<0.001$, $ES=2.84$;
 21 S+E by 4.0-fold, $P<0.001$, $ES=2.05$) (Table 1). CK was significantly elevated in comparison
 22 to Pre in E+S at Mid ($13\pm 9\%$ from 93 ± 21 $\text{mIU}\cdot\text{l}^{-1}$, $P<0.01$, $ES=0.544$), Post ($16\pm 9\%$, $P<0.01$,
 23 $ES=0.681$) and 24h ($57\pm 32\%$, $P<0.05$, $ES=1.73$) and for S+E at Post ($26\pm 17\%$ from 95 ± 31
 24 $\text{mIU}\cdot\text{l}^{-1}$, $P<0.01$, $ES=0.679$) but not at 24h ($96\pm 89\%$, $ES=1.29$) (Table 1). CK further
 25 increased in both groups between Post and 24h (E+S $P<0.05$, $ES=1.337$ and S+E $P<0.05$,
 26 $ES=0.935$). No between-group differences were observed during loading or recovery.

1

2

*** Figure 3 and 4 near here ***

3

4 **Acute loading responses at week 24**

5 No significant changes in body weight ($-0.5\pm 0.2\%$), HGB ($+4.2\pm 3.8\%$) or HCR ($+4.4\pm 2.9\%$)
6 were observed during the loading.

7 MVC decreased significantly for both groups by Mid (E+S $-16\pm 9\%$ from 1833 ± 322 N,
8 $P<0.001$, ES= -0.890 ; S+E $-21\pm 8\%$ from 1966 ± 690 N, $P<0.001$, ES= -0.623) and by Post
9 (E+S $-24\pm 6\%$, $P<0.001$, ES= -1.66 ; S+E $-22\pm 8\%$, $P<0.01$, ES= -0.731) (Figure 2). MVC₅₀₀
10 decreased significantly for both groups by Mid (E+S $-18\pm 17\%$ $P<0.05$, ES= -0.683 and S+E -
11 $20\pm 12\%$, $P<0.05$, ES= -0.678) and post (E+S $-24\pm 6\%$, $P<0.001$, ES= -1.43 and S+E $-25\pm 23\%$
12 $P<0.05$, ES= -0.637). Both groups recovered significantly ($P<0.05-0.001$) from Post to 24h
13 and 48h. No between-group differences were observed during loading or recovery.

14 A significant increase in T was found in E+S at Mid (from 0.8 ± 0.3 to 1.2 ± 0.4 nmol \cdot l $^{-1}$,
15 $P<0.05$, ES= 1.247) and S+E at Post (from 0.9 ± 0.9 to 1.5 ± 1.2 nmol \cdot l $^{-1}$, $P<0.001$, ES= 0.536)
16 (Figure 3). For S+E, T significantly increased from Mid to Post ($P<0.01$, ES= 0.371). C was
17 decreased for S+E from Pre at 48h (-28% , $P<0.001$, ES= -0.974) and near-significantly
18 decreased at 24h (-26% , $P=0.051$, ES= -0.8254) in comparison to Pre (Table 1). A significant
19 increase in GH was noted for both groups at Mid (E+S 7.0-fold from, $P<0.01$, ES= 0.885 and
20 S+E 2.5-fold, $P<0.05$, ES= 0.790) and at Post for S+E (3.6-fold, $P<0.05$, ES= 0.886) (Figure
21 4). For E+S a decrease took place from Mid to Post (from 68.3 ± 31.7 to 14.7 ± 11.3 mIU \cdot l $^{-1}$,
22 $P<0.01$, ES= -0.885). A between-group difference was observed at Mid ($P<0.01$), and the
23 change from Mid to Post was significantly different between groups ($P<0.001$).

24 Blood lactate increased 5-fold for E+S by Mid ($P<0.001$, ES= 3.03) and 6-fold for S+E
25 ($P<0.001$, ES= 3.24) (Table 1). Lactate at Post was increased 5-fold for E+S ($P<0.001$,

1 ES=4.81) and 5.5-fold for S+E ($P<0.001$, ES=3.23) (Table 1). CK was significantly ($P<0.05$)
2 elevated in comparison to Pre for S+E at Mid ($14\pm 9\%$, from 108 ± 72 $\text{mIU}\cdot\text{l}^{-1}$, $P<0.001$,
3 ES=0.169) and Post ($25\pm 17\%$, $P<0.01$, ES=0.299), but not at 24h or 48h (Table 1). No
4 between-group differences were observed during loading or recovery.

5

6
7
8 *** Table 1 near here ***

9

10 **Differences in the acute loading responses before and after training**

11 In E+S, the GH response was significantly different at Pre-Mid ($P<0.01$), Pre-Post and Mid-
12 Post ($P<0.05$) in comparison to the corresponding changes at Week 0 (Figure 4). The relative
13 change in blood lactate was significantly larger after 24 weeks of training than the
14 corresponding change at Week 0 for S+E during Pre-Mid ($P<0.01$), Pre-Post ($P<0.05$) and
15 Mid-Post ($P<0.05$) (Table 1).

16

17 **Basal measurements**

18 Both groups increased 1RM (E+S by $13\pm 12\%$ from 102 ± 21 kg, $P<0.001$, ES=0.569 and S+E
19 by $17\pm 10\%$ from 99 ± 18 kg, $P<0.001$, ES=0.884) (Figure 5), W_{max} (E+S by $21\pm 10\%$ from
20 170 ± 26 W, $P<0.001$, ES=1.36 and S+E by $16\pm 12\%$ from 182 ± 27 W, $P<0.001$ ES=1.05) and
21 CSA (E+S by $15\pm 10\%$ from $17\pm 2\text{cm}^2$, $P<0.001$, ES=1.32 and S+E by $11\pm 8\%$ from $19\pm 3\text{cm}^2$,
22 $P<0.001$, ES=0.680). Basal hormone concentrations are presented in Table 2.

23

24 **Correlations**

1 No significant correlations were observed between the acute changes in testosterone, cortisol
2 or growth hormone and long-term 1RM, W_{\max} or CSA development in either E+S or S+E.
3 Basal levels of T, T/SHBG and T/C were not correlated with changes in 1 RM or W_{\max} or
4 CSA.

5

6 *** Figure 5 and Table 2 near here ***

7

8 **DISCUSSION**

9 The main findings of the present study were that following the experimental loading at Week
10 0, significantly elevated serum GH was observed only in S+E, while serum T remained
11 unchanged in both groups. At Week 24, both T and GH were significantly elevated in S+E
12 but not in E+S at Post. The exercise order did not affect the magnitude of loading-induced
13 fatigue measured as maximal voluntary isometric force and rapid force production either at
14 Week 0 or 24. Additionally, muscle force production was recovered by 24 h following both
15 exercise orders both at Week 0 and 24. The present 24-week combined strength and
16 endurance training period resulted in significant increases in 1RM strength, W_{\max} and muscle
17 cross-sectional of similar magnitudes in both groups. These chronic adaptations were not
18 associated with the acute exercise-induced changes of serum hormones in either order.

19 In accordance with our previous study with men performing the same experimental loading
20 with the same relative intensity (39), we observed no acutely elevated concentrations of T in
21 the present study at Post before the prolonged training period following either order. This
22 outcome was expected, considering the combination of explosive, maximal and hypertrophic
23 sets in the present strength loading. Thus, the protocol was likely not strenuous enough to
24 elicit acute anabolic responses (39), as large elevations in T would be expected in women
25 mainly following hypertrophic type protocols with a large stress on the metabolic system (26,

1 29). However, this design was purposefully chosen to reflect the content of the 24-week
2 training program which was created based on common exercise recommendations (42).

3 Interestingly, as elevated concentrations of serum T were observed during loading for E+S at
4 Mid (Week 0 and 24), and for S+E at Post (Week 24), our results suggest that the observed
5 elevations may primarily have been a result of the present endurance exercise. Considering
6 the likely absence of haemoconcentration in the present study, this supports previous findings
7 of endurance exercise inducing elevations in T in female populations (15, 24). The lack of
8 significantly increased T at Post for the S+E group at Week 0 is in line with earlier
9 investigations in men (4, 37, 39), with unchanged concentrations of T following a combined
10 loading in the S+E order. However, the significantly elevated concentration of serum T in
11 S+E at Week 24 could be related to training-induced increased sensitivity to
12 adrenocorticotrophic hormone (30), which stimulates the adrenal cortex and releases androgens
13 as a byproduct of cortisol secretion (23, 34). This together with the relatively higher rise in
14 lactate after training could be related to why elevated T was observed in the S+E group at
15 Post at Week 24, but not at Week 0. However, no such observation was made in the E+S
16 group. Furthermore, as we only measured total testosterone and did also not detect any
17 significant elevations in cortisol during the loadings in either order, this hypothesis remains
18 speculative.

19 It also needs to be acknowledged that the underlying causes of exercise-induced elevations in
20 T in women are not fully comprehended, and not all plausible mechanisms were monitored in
21 the present study. Possible mechanisms include e.g. the time course of androgen receptor
22 regulation (44) and reduced hepatic clearance as observed in men (5). Haemoconcentration as
23 an indirect cause for elevated T can likely be ruled out in the present study due to unchanged
24 hemoglobin and hematocrit during loading. It is also possible that oral contraceptive use
25 affects the secretion of T as well as the metabolites of dehydroepiandrosterone (15) and,
26 consequently, the biosynthesis pathway of T during exercise. However, in the present study, a

1 similar number of subjects in both groups reported oral contraceptive use and, on a group
2 level, the pattern of T response to endurance exercise was comparable in both orders.

3 Similarly to the exercise-related variables affecting the acute responses of T, the intensity of
4 exercise is a major contributor to the magnitude of responses of GH in women (29, 46). As
5 expected based on previous findings (16), the GH concentrations in the E+S order both before
6 and after training were significantly elevated after endurance exercise but diminished
7 following the strength loading. This pattern in the kinetics of exercise-induced growth
8 hormone release was significantly different between groups both before and after training as
9 S+E demonstrated elevated GH throughout the loading in contrast to E+S. These differing
10 GH responses may have been caused by endurance exercise-induced lipolysis (36). The
11 release of free fatty acids (FFA) is likely a major influence for suppressed GH release,
12 possibly through affecting anterior pituitary function (6). It needs to be noted, that even
13 though oral contraceptive use could amplify lipolysis during continuous cycling exercise, the
14 FFA concentration is likely to remain unaffected (7).

15 Due to a critical relative threshold for GH secretion, the intensity of training would need to be
16 continuously progressive in order for significant GH responses to occur within a loading
17 session after prolonged training (8). In the present study, the loading was conducted with
18 values relative to 1RM and W_{\max} in order to keep the relative intensity the same at both weeks
19 0 and 24 and to be matched for the current training status and improved performance level of
20 the subjects. This was reflected in the GH responses in both groups at Week 24 as higher
21 absolute concentrations at both Mid and Post in comparison to corresponding time points at
22 Week 0. In the S+E order, despite the fact that the magnitude of the loading-induced changes
23 (Pre-Mid and Pre-Post) were not statistically larger than the corresponding magnitudes at
24 Week 0, the GH responses during loading at Week 24 were statistically significant. This
25 serves as an indication of adaptation to training, as the same relative exercise intensity was
26 potent in significantly elevating serum GH concentrations. Similar indications for training
27 adaptations were found in the E+S group, as the magnitudes of the Pre-Mid and Pre-Post

1 changes were significantly larger at Week 24 than at Week 0. Interestingly, while the
2 adaptations in GH release seem to indicate that the loadings were still strenuous at Week 24,
3 changes in the behaviour of CK may suggest better tolerance of the experimental loading.
4 CK was slightly elevated both during loading and recovery in both groups at Week 0, but
5 during Week 24 only elevated in S+E during loading and similar to resting levels during
6 recovery. As CK can be considered to be an indirect indicator of muscle damage, the lack of
7 its presence during recovery after training may indicate an increased tolerance for combined
8 strength and endurance loadings, similarly to what was recently observed in men (39).

9 Interestingly, although the GH responses clearly differed between the present exercise orders
10 during loading, no between-group differences were observed in training-induced increases in
11 muscle CSA. The implications of the present findings thus require further clarification e.g.
12 through examining additional forms of GH than solely the present 22 kDa variant. While an
13 acute bout of exercise may stimulate variants of GH that are incapable of generating increases
14 in biological activity, chronic resistance exercise may increase the circulating concentrations
15 of biologically active growth hormone (27). This may in part explain why the GH responses
16 in the present study were not related to the changes in muscle cross-sectional area in either
17 group and warrants further investigation of the mechanisms of several GH variants both
18 during combined strength and endurance loadings as well as prolonged combined training.

19 In addition to a lack of a relationship between changes in muscle cross-sectional area and the
20 magnitudes of acute GH release, we also observed no associations between acute responses of
21 T and long-term performance adaptations. While it has been suggested that tissue exposure to
22 acute elevations in anabolic hormones would not be associated with hypertrophy or strength
23 performance (47), such correlations have been demonstrated in male populations following
24 pure strength training (e.g. 22, 31). Furthermore, previously reported correlations of basal
25 levels of T and T/SHBG-ratios and strength development and changes in CSA following
26 strength training in women (e.g. 17, 21) were not found in present study despite significantly
27 increased basal levels of T. Thus, it seems reasonable to suggest that the detection of possible

1 linkages of resistance exercise-induced anabolic responses and gains in strength and
2 hypertrophy may be interfered when strength training is simultaneously accompanied by
3 endurance training both in men (39) and women. However, further studies with different
4 training protocols are needed for more definite conclusions.

5 It is noteworthy that both exercise orders resulted in similar training-induced long-term gains
6 in 1RM, thus challenging earlier suggestions of the order of S+E being superior to E+S in
7 terms of adaptations in strength performance (3, 28). The experimental loading showed that
8 the acute fatigue in terms of exercise-induced decreases in MVC and MVC₅₀₀ were of similar
9 magnitudes following both loading conditions both before and after training. Furthermore, as
10 neither MVC nor MVC₅₀₀ were no longer significantly depressed from Pre-loading values by
11 24h, the experimental loadings indicate that the recovery of maximal and rapid strength
12 performance was completed within 24 hours of cessation of exercise. Consequently, it can be
13 assumed that the recovery between individual training sessions was sufficient, as the sessions
14 were consistently separated by at least 48-72h. Even though we only monitored recovery
15 before and after 24 weeks of training, the loads utilized in the experimental loading were
16 similar to those used during training. This may, in part, explain similar gains in 1RM in both
17 groups.

18 However, it also needs to be noted that the findings regarding the effect of the mode of
19 endurance exercise (i.e. running or cycling) on changes in strength performance are to date
20 equivocal (40, 48). Thus, comparisons of the present training program, consisting of cycling
21 endurance training, to other protocols should be done with caution. Interestingly, while no
22 between-group differences were observed in long-term training-induced changes, the
23 magnitude of gains in strength in relation to endurance performance was highly individual in
24 both exercise orders (Figure 5). This warrants further investigation regarding the mechanisms
25 of the underlying adaptations to same-session combined strength and endurance training in
26 women.

1 To conclude, this study demonstrated that the acute hormone and force responses to a
2 combined strength and endurance loading were by large similar between exercise orders in
3 previously untrained women both before and after training, with the exception of differences
4 in the kinetics of serum concentrations of GH during exercise. Furthermore, our results
5 showed that strength and endurance performance as well as muscle cross-sectional area
6 following 24 weeks of same-session combined strength and endurance training were similar
7 in both exercise orders. Therefore, our data indicates that despite some differences in the
8 acute anabolic responses to exercise, the present 24-week combined strength and endurance
9 training program resulted in similar long-term performance and morphological adaptations in
10 both groups.

11 **PRACTICAL APPLICATIONS**

12 As the present study did not show order-specific responses of recovery of force, the findings
13 indicate that the exercise order does not seem to be of great importance for previously
14 untrained women when combining strength and endurance into the same training session.
15 Even though the growth hormone responses to exercise were significantly larger in S+E
16 compared to E+S both before and after the training, this was not reflected in or associated
17 with the long-term adaptations. Consequently, the gains in strength and endurance
18 performance as well as muscle size were of similar magnitudes in the two training groups
19 following 24 weeks of combined strength and endurance training. Thus, previously untrained
20 women can achieve performance improvements and increases in muscle size by combining
21 strength and endurance into the same training session with either exercise order, when
22 sufficient recovery is allowed.

1

2 **Captions and legends for figures**

3

4 **Figure 1.** Overview of the experimental design. E+S = Endurance preceding strength, S+E =
5 Strength preceding endurance.

6 **Figure 2.** MVC for E+S and S+E during loading and recovery at Week 0 and at Week 24.
7 Within-group differences: *=Significant from Pre, §=Significant from Post. ** $P<0.01$,
8 *** $P<0.001$, § $P<0.05$, §§ $P<0.01$ and §§§ $P<0.001$.

9 **Figure 3.** Responses in total testosterone for E+S and S+E during loading and recovery at
10 Week 0 and at Week 24. Within-group differences: *=Significant from Pre, +=Significant
11 from Mid, §=Significant from Post. * $P<0.05$, *** $P<0.001$, + $P<0.05$ and §§§ $P<0.001$.

12 **Figure 4.** Growth hormone responses for E+S and S+E during loading at Week 0 and at
13 Week 24. Within-group differences: *=Significant from Pre, +=Significant from Mid,
14 §=Significant from Post, •=significant from Week 0. #=between-group difference at given
15 time point. * $P<0.05$. ** $P<0.01$, + $P<0.05$, + $P<0.01$, • $P<0.05$, •• $P<0.01$, # $P<0.05$, ## $P<0.01$
16 and ### $P<0.001$.

17 **Figure 5.** Changes in 1RM (left), Maximal workload (middle) during the cycling endurance
18 test and the individual ratios of the magnitude of gains in 1RM and workload (right).
19 *=significant from Week 0. *** $P<0.001$.

1

2 **Captions and legends for tables**

3 **Table 1.** Exercise-induced changes in serum cortisol (C), creatine kinase (CK) and blood
 4 lactate (La) during loading and recovery for E+S and S+E at Week 0 and 24. Within-group
 5 differences: *=significant from Pre, +=significant from Mid, §=significant from Post, α =
 6 significant from 24 h, •=significant from corresponding value at Week 0. * $P<0.05$. ** $P<0.01$,
 7 *** $P<0.001$, + $P<0.05$, + $P<0.01$, • $P<0.05$, •• $P<0.01$, α $P=0.05$, § $P<0.05$. §§ $P<0.01$, §§§
 8 $P<0.001$.

9 **Table 2.** Basal serum concentrations of total testosterone (T), cortisol (C), growth hormone
 10 (GH), sex-hormone binding globulin (SHBG), and ratios of testosterone and cortisol (T/C)
 11 and testosterone and sex-hormone binding globulin (T/SHBG) before and after the training
 12 intervention.

13 * within-group difference to Week 0; * $P<0.05$ and *** $P<0.001$

1

2 **REFERENCES**

3 1. Ahtiainen, JP, Hoffren, M, Hulmi, JJ, Pietikainen, M, Mero, AA, Avela, J, and Häkkinen,
4 K. Panoramic ultrasonography is a valid method to measure changes in skeletal muscle cross-
5 sectional area. *Eur. J. Appl. Physiol.* 108: 273-279, 2010.

6 2. Aunola, S, and Rusko, H. Aerobic and anaerobic thresholds determined from venous
7 lactate or from ventilation and gas exchange in relation to muscle fiber composition. *Int. J.*
8 *Sports Med.* 7: 161-166, 1986.

9 3. Cadore, EL, Izquierdo, M, Alberton, CL, Pinto, RS, Conceicao, M, Cunha, G, Radaelli, R,
10 Bottaro, M, Trindade, GT, and Kruel, LF. Strength prior to endurance intra-session exercise
11 sequence optimizes neuromuscular and cardiovascular gains in elderly men. *Exp. Gerontol.*
12 47: 164-169, 2012.

13 4. Cadore, EL, Izquierdo, M, dos Santos, MG, Martins, JB, Rodrigues Lhullier, FL, Pinto,
14 RS, Silva, RF, and Kruel, LF. Hormonal responses to concurrent strength and endurance
15 training with different exercise orders. *J. Strength Cond Res.* 26: 3281-3288, 2012.

16 5. Cadoux-Hudson, TA, Few, JD, and Imms, FJ. The effect of exercise on the production and
17 clearance of testosterone in well trained young men. *Eur. J. Appl. Physiol. Occup. Physiol.*
18 54: 321-325, 1985.

19 6. Casanueva, FF, Villanueva, L, Dieguez, C, Diaz, Y, Cabranes, JA, Szoke, B, Scanlon, MF,
20 Schally, AV, and Fernandez-Cruz, A. Free fatty acids block growth hormone (GH) releasing
21 hormone-stimulated GH secretion in man directly at the pituitary. *J. Clin. Endocrinol. Metab.*
22 65: 634-642, 1987.

- 1 7. Casazza, GA, Jacobs, KA, Suh, SH, Miller, BF, Horning, MA, and Brooks, GA. Menstrual
2 cycle phase and oral contraceptive effects on triglyceride mobilization during exercise. *J.*
3 *Appl. Physiol.* (1985) 97: 302-309, 2004.
- 4 8. Chang, FE, Dodds, WG, Sullivan, M, Kim, MH, and Malarkey, WB. The acute effects of
5 exercise on prolactin and growth hormone secretion: comparison between sedentary women
6 and women runners with normal and abnormal menstrual cycles. *J. Clin. Endocrinol. Metab.*
7 62: 551-556, 1986.
- 8 9. Copeland, JL, Consitt, LA, and Tremblay, MS. Hormonal responses to endurance and
9 resistance exercise in females aged 19-69 years. *J. Gerontol. A Biol. Sci. Med. Sci.* 57: B158-
10 65, 2002.
- 11 10. Cumming, DC, Wall, SR, Galbraith, MA, and Belcastro, AN. Reproductive hormone
12 responses to resistance exercise. *Med. Sci. Sports Exerc.* 19: 234-238, 1987.
- 13 11. Davitt, PM, Pellegrino, JK, Schanzer, JR, Tjionas, H, and Arent, SM. The effects of a
14 combined resistance training and endurance exercise program in inactive college female
15 subjects: does order matter? *J. Strength Cond Res.* 28: 1937-1945, 2014.
- 16 12. de Souza, EO, Tricoli, V, Franchini, E, Paulo, AC, Regazzini, M, and Ugrinowitsch, C.
17 Acute effect of two aerobic exercise modes on maximum strength and strength endurance. *J.*
18 *Strength Cond Res.* 21: 1286-1290, 2007.
- 19 13. Dill, DB, and Costill, DL. Calculation of percentage changes in volumes of blood,
20 plasma, and red cells in dehydration. *J. Appl. Physiol.* 37: 247-248, 1974.
- 21 14. Eklund, D, Pulverenti, T, Bankers, S, Avela, J, Newton, R, Schumann, M, and Häkkinen,
22 K. Neuromuscular adaptations to different modes of combined strength and endurance
23 training. *Int. J. Sports Med.* 36: 120-129, 2015.

- 1 15. Enea, C, Boisseau, N, Ottavy, M, Mulliez, J, Millet, C, Ingrand, I, Diaz, V, and Dugue, B.
2 Effects of menstrual cycle, oral contraception, and training on exercise-induced changes in
3 circulating DHEA-sulphate and testosterone in young women. *Eur. J. Appl. Physiol.* 106:
4 365-373, 2009.
- 5 16. Goto, K, Higashiyama, M, Ishii, N, and Takamatsu, K. Prior endurance exercise
6 attenuates growth hormone response to subsequent resistance exercise. *Eur. J. Appl. Physiol.*
7 94: 333-338, 2005.
- 8 17. Häkkinen, K, Pakarinen, A, and Kallinen, M. Neuromuscular adaptations and serum
9 hormones in women during short-term intensive strength training. *Eur. J. Appl. Physiol.*
10 *Occup. Physiol.* 64: 106-111, 1992.
- 11 18. Häkkinen, K, and Pakarinen, A. Acute hormonal responses to two different fatiguing
12 heavy-resistance protocols in male athletes. *J. Appl. Physiol.* (1985) 74: 882-887, 1993.
- 13 19. Häkkinen, K, and Pakarinen, A. Acute hormonal responses to heavy resistance exercise in
14 men and women at different ages. *Int. J. Sports Med.* 16: 507-513, 1995.
- 15 20. Häkkinen, K, Kallinen, M, Izquierdo, M, Jokelainen, K, Lassila, H, Mälkiä, E, Kraemer,
16 WJ, Newton, RU, and Alen, M. Changes in agonist-antagonist EMG, muscle CSA, and force
17 during strength training in middle-aged and older people. *J. Appl. Physiol.* (1985) 84: 1341-
18 1349, 1998.
- 19 21. Häkkinen, K, Pakarinen, A, Kraemer, WJ, Newton, RU, and Alen, M. Basal
20 concentrations and acute responses of serum hormones and strength development during
21 heavy resistance training in middle-aged and elderly men and women. *J. Gerontol. A Biol.*
22 *Sci. Med. Sci.* 55: B95-105, 2000.

- 1 22. Hansen, S, Kvorning, T, Kjaer, M, and Sjøgaard, G. The effect of short-term strength
2 training on human skeletal muscle: the importance of physiologically elevated hormone
3 levels. *Scand. J. Med. Sci. Sports* 11: 347-354, 2001.
- 4 23. Johnson, LG, Kraemer, RR, Haltom, R, Kraemer, GR, Gaines, HE, and Castracane, VD.
5 Effects of estrogen replacement therapy on dehydroepiandrosterone, dehydroepiandrosterone
6 sulfate, and cortisol responses to exercise in postmenopausal women. *Fertil. Steril.* 68: 836-
7 843, 1997.
- 8 24. Keizer, HA, Kuipers, H, de Haan, J, Janssen, GM, Beckers, E, Habets, L, van
9 Kranenburg, G, and Geurten, P. Effect of a 3-month endurance training program on metabolic
10 and multiple hormonal responses to exercise. *Int. J. Sports Med.* 8 Suppl 3: 154-160, 1987.
- 11 25. Kraemer, WJ, Gordon, SE, Fleck, SJ, Marchitelli, LJ, Mello, R, Dziados, JE, Friedl, K,
12 Harman, E, Maresh, C, and Fry, AC. Endogenous anabolic hormonal and growth factor
13 responses to heavy resistance exercise in males and females. *Int. J. Sports Med.* 12: 228-235,
14 1991.
- 15 26. Kraemer, WJ, Fleck, SJ, Dziados, JE, Harman, EA, Marchitelli, LJ, Gordon, SE, Mello,
16 R, Frykman, PN, Koziris, LP, and Triplett, NT. Changes in hormonal concentrations after
17 different heavy-resistance exercise protocols in women. *J. Appl. Physiol.* (1985) 75: 594-604,
18 1993.
- 19 27. Kraemer, WJ, Nindl, BC, Marx, JO, Gotshalk, LA, Bush, JA, Welsch, JR, Volek, JS,
20 Spiering, BA, Maresh, CM, Mastro, AM, and Hymer, WC. Chronic resistance training in
21 women potentiates growth hormone in vivo bioactivity: characterization of molecular mass
22 variants. *Am. J. Physiol. Endocrinol. Metab.* 291: E1177-87, 2006.
- 23 28. Leveritt, M, and Abernethy, PJ. Acute effects of high-intensity endurance exercise on
24 subsequent resistance activity. *Sports Med.* 13: 47-51, 1999.

- 1 29. Linnamo, V, Pakarinen, A, Komi, PV, Kraemer, WJ, and Häkkinen, K. Acute hormonal
2 responses to submaximal and maximal heavy resistance and explosive exercises in men and
3 women. *J. Strength Cond Res.* 19: 566-571, 2005.
- 4 30. Loucks, AB, Mortola, JF, Girton, L, and Yen, SS. Alterations in the hypothalamic-
5 pituitary-ovarian and the hypothalamic-pituitary-adrenal axes in athletic women. *J. Clin.*
6 *Endocrinol. Metab.* 68: 402-411, 1989.
- 7 31. McCall, GE, Byrnes, WC, Fleck, SJ, Dickinson, A, and Kraemer, WJ. Acute and chronic
8 hormonal responses to resistance training designed to promote muscle hypertrophy. *Can. J.*
9 *Appl. Physiol.* 24: 96-107, 1999.
- 10 32. Nindl, BC, Kraemer, WJ, Gotshalk, LA, Marx, JO, Volek, JS, Bush, FA, Häkkinen, K,
11 Newton, RU, and Fleck, SJ. Testosterone responses after resistance exercise in women:
12 influence of regional fat distribution. *Int. J. Sport Nutr. Exerc. Metab.* 11: 451-465, 2001.
- 13 33. O'Leary, CB, Lehman, C, Koltun, K, Smith-Ryan, A, and Hackney, AC. Response of
14 testosterone to prolonged aerobic exercise during different phases of the menstrual cycle. *Eur.*
15 *J. Appl. Physiol.* 113: 2419-2424, 2013.
- 16 34. Petrides, JS, Mueller, GP, Kalogeras, KT, Chrousos, GP, Gold, PW, and Deuster, PA.
17 Exercise-induced activation of the hypothalamic-pituitary-adrenal axis: marked differences in
18 the sensitivity to glucocorticoid suppression. *J. Clin. Endocrinol. Metab.* 79: 377-383, 1994.
- 19 35. Raastad, T, Bjoro, T, and Hallen, J. Hormonal responses to high- and moderate-intensity
20 strength exercise. *Eur. J. Appl. Physiol.* 82: 121-128, 2000.
- 21 36. Romijn, JA, Coyle, EF, Sidossis, LS, Gastaldelli, A, Horowitz, JF, Endert, E, and Wolfe,
22 RR. Regulation of endogenous fat and carbohydrate metabolism in relation to exercise
23 intensity and duration. *Am. J. Physiol.* 265: E380-91, 1993.

- 1 37. Rosa, C, Vilaca-Alves, J, Fernandes, HM, Saavedra, FJ, Pinto, RS, and Machado Dos
2 Reis, V. Order effects of combined strength and endurance training on testosterone, cortisol,
3 growth hormone and IGFBP-3 in concurrent-trained men. *J. Strength Cond Res.* , 2014.
- 4 38. Schumann, M, Kiiusmaa, M, Newton, RU, Sirparanta, AI, Syvaoja, H, Hakkinen, A, and
5 Hakkinen, K. Fitness and lean mass increases during combined training independent of
6 loading order. *Med. Sci. Sports Exerc.* , 2014.
- 7 39. Schumann, M, Walker, S, Izquierdo, M, Newton, RU, Kraemer, WJ, and Hakkinen, K.
8 The order effect of combined endurance and strength loadings on force and hormone
9 responses: effects of prolonged training. *Eur. J. Appl. Physiol.* 114: 867-880, 2014.
- 10 40. Silva, RF, Cadore, EL, Kothe, G, Guedes, M, Alberton, CL, Pinto, SS, Pinto, RS,
11 Trindade, G, and Krueel, LF. Concurrent training with different aerobic exercises. *Int. J.*
12 *Sports Med.* 33: 627-634, 2012.
- 13 41. Spiering, BA, Kraemer, WJ, Anderson, JM, Armstrong, LE, Nindl, BC, Volek, JS, and
14 Maresh, CM. Resistance exercise biology: manipulation of resistance exercise programme
15 variables determines the responses of cellular and molecular signalling pathways. *Sports Med.*
16 38: 527-540, 2008.
- 17 42. Thompson, W, Gordon, N, and Pescatello, L. ACSM's Guidelines for Exercise Testing
18 and Prescription. Philadelphia; Lippincott Williams & Wilkins.
- 19 43. Timon, R, Corvillo, M, Brazo, J, Robles, MC, and Maynar, M. Strength training effects
20 on urinary steroid profile across the menstrual cycle in healthy women. *Eur. J. Appl. Physiol.*
21 113: 1469-1475, 2013.
- 22 44. Vingren, JL, Kraemer, WJ, Hatfield, DL, Volek, JS, Ratamess, NA, Anderson, JM,
23 Hakkinen, K, Ahtiainen, J, Fragala, MS, Thomas, GA, Ho, JY, and Maresh, CM. Effect of

1 resistance exercise on muscle steroid receptor protein content in strength-trained men and
2 women. *Steroids* 74: 1033-1039, 2009.

3 45. Walker, S, Santolamazza, F, Kraemer, W, and Häkkinen, K. Effects of Prolonged
4 Hypertrophic Resistance Training on Acute Endocrine Responses in Young and Older Men.
5 *J. Aging Phys. Act.* , 2014.

6 46. Weltman, A, Weltman, JY, Schurrer, R, Evans, WS, Veldhuis, JD, and Rogol, AD.
7 Endurance training amplifies the pulsatile release of growth hormone: effects of training
8 intensity. *J. Appl. Physiol. (1985)* 72: 2188-2196, 1992.

9 47. West, DW, Burd, NA, Tang, JE, Moore, DR, Staples, AW, Holwerda, AM, Baker, SK,
10 and Phillips, SM. Elevations in ostensibly anabolic hormones with resistance exercise
11 enhance neither training-induced muscle hypertrophy nor strength of the elbow flexors. *J.*
12 *Appl. Physiol. (1985)* 108: 60-67, 2010.

13 48. Wilson, JM, Marin, PJ, Rhea, MR, Wilson, SM, Loenneke, JP, and Anderson, JC.
14 Concurrent training: a meta-analysis examining interference of aerobic and resistance
15 exercises. *J. Strength Cond Res.* 26: 2293-2307, 2012.

16

Table 1. Exercise-induced changes in serum cortisol (C), creatine kinase (CK) and blood lactate (La) during loading and recovery for E+S and S+E at Week 0 and 24. Within-group differences: *=significant from Pre, +=significant from Mid, §=significant from Post, ♂ = significant from 24 h, •=significant from corresponding value at Week 0. **P*<0.05. ***P*<0.01, ****P*<0.001, +*P*0.05, +*P*<0.01, •*P*<0.05, ••*P*<0.01, ♂ *P*=0.05, § *P*<0.05. §§ *P*<0.01, §§§ *P*<0.001.

	WEEK 0		WEEK 24	
	E+S	S+E	E+S	S+E
<i>C</i> (nmol·l ⁻¹)	(<i>n</i> =10)	(<i>n</i> =10)	(<i>n</i> =8)	(<i>n</i> =10)
Pre	397±128	479±105	413±154	616±178
Mid	437±176	367±139	471±96	567±199
Post	426±207	435±211	423±108	778±252
24 h	312±111 [*] <i>p</i> =0.051	332±87 **	284±134	459±200 [*] <i>p</i> =0.051 §§
48 h	347±141	333±60 **	322±116	445±170*** §§§
<i>CK</i> (mlU·l ⁻¹)	(<i>n</i> =9)	(<i>n</i> =9)	(<i>n</i> =9)	(<i>n</i> =9)
Pre	93±21	95±31	103±44	109±72
Mid	103±19 **	106±32	121±49	121±71 ***
Post	105±17 **	118±35 ** +	123±55	131±75 **
24 h	143±36 *	181±89 §	145±80	148±81
48 h	128±38	132±55 ♂	141±116	123±70
<i>La</i> (mmol·l ⁻¹)	(<i>n</i> =12)	(<i>n</i> =11)	(<i>n</i> =12)	(<i>n</i> =11)
Pre	1.1±0.3	1.3±0.3	1.2±0.4	1.1±0.4
Mid	4.7±1.0 ***	5.0±1.9 ***	6.3±2.3 ***	6.8±2.4 *** ••
Post	5.6±2.2 ***	5.2±2.3 ***	6.5±1.5 ***	6.0±2.1 *** •

Table 2. Basal serum concentrations of total testosterone (T), cortisol (C), growth hormone (GH), sex-hormone binding globulin (SHBG), and ratios of testosterone and cortisol (T/C) and testosterone and sex-hormone binding globulin (T/SHBG) before and after the training intervention.

* within-group difference to Week 0; * $P < 0.05$ and *** $P < 0.001$

	WEEK 0		WEEK 24	
	E+S	S+E	E+S	S+E
T (nmol·l ⁻¹)	0.6±0.3	0.6±0.4	1.1±0.3 ***	1.3±0.9 ***
C (nmol·l ⁻¹)	546±186	564±185	614±225	673±136 *
GH (mIU·l ⁻¹)	7.9±12.1	11.0±10.0	6.1±11.1	14.0±20.6
SHBG (nmol·l ⁻¹)	52.3±17.5	65.3±42.0	68.6±12.7 *	81.6±43.4 *
T/C *10 ³	1.4±1.1	1.3±1.0	1.9±1.1 *	2.1±1.8 *
T/SHBG *10 ³	15.3±8.8	13.1±10.7	17.3±6.7	21.5±11.4









