

**This is an electronic reprint of the original article.  
This reprint *may differ* from the original in pagination and typographic detail.**

**Author(s):** Kenig, Carlos; Salo, Mikko

**Title:** The Calderón problem with partial data on manifolds and applications

**Year:** 2013

**Version:**

**Please cite the original version:**

Kenig, C., & Salo, M. (2013). The Calderón problem with partial data on manifolds and applications. *Analysis & PDE*, 6(8), 2003-2048.  
<https://doi.org/10.2140/apde.2013.6.2003>

All material supplied via JYX is protected by copyright and other intellectual property rights, and duplication or sale of all or part of any of the repository collections is not permitted, except that material may be duplicated by you for your research use or educational purposes in electronic or print form. You must obtain permission for any other use. Electronic or print copies may not be offered, whether for sale or otherwise to anyone who is not an authorised user.

# ANALYSIS & PDE

Volume 6

No. 8

2013

CARLOS KENIG AND MIKKO SALO

**THE CALDERÓN PROBLEM WITH PARTIAL DATA ON  
MANIFOLDS  
AND APPLICATIONS**



# THE CALDERÓN PROBLEM WITH PARTIAL DATA ON MANIFOLDS AND APPLICATIONS

CARLOS KENIG AND MIKKO SALO

We consider Calderón’s inverse problem with partial data in dimensions  $n \geq 3$ . If the inaccessible part of the boundary satisfies a (conformal) flatness condition in one direction, we show that this problem reduces to the invertibility of a broken geodesic ray transform. In Euclidean space, sets satisfying the flatness condition include parts of cylindrical sets, conical sets, and surfaces of revolution. We prove local uniqueness in the Calderón problem with partial data in admissible geometries, and global uniqueness under an additional concavity assumption. This work unifies two earlier approaches to this problem — one by Kenig, Sjöstrand, and Uhlmann, the other by Isakov — and extends both. The proofs are based on improved Carleman estimates with boundary terms, complex geometrical optics solutions involving reflected Gaussian beam quasimodes, and invertibility of (broken) geodesic ray transforms. This last topic raises questions of independent interest in integral geometry.

1. Introduction	2003
2. Statement of results	2006
3. The Euclidean case	2010
4. Carleman estimate	2015
5. Reflection approach	2021
6. Local uniqueness on simple manifolds	2025
7. Quasimodes concentrating near broken rays	2030
8. Recovering the broken ray transform	2043
Acknowledgements	2045
References	2045

## 1. Introduction

This article is concerned with inverse problems where measurements are made only on part of the boundary. A typical example is the inverse problem of Calderón, where the objective is to determine the electrical conductivity of a medium from voltage and current measurements on its boundary. The mathematical formulation of this problem is as follows. Let  $\Omega \subset \mathbb{R}^n$ ,  $n \geq 2$ , be a bounded domain with smooth boundary. Given a positive function  $\gamma \in L^\infty(\Omega)$  (the electrical conductivity of the medium) and two open subsets  $\Gamma_D, \Gamma_N$  of  $\partial\Omega$ , consider the partial Cauchy data set

$$C_\gamma^{\Gamma_D, \Gamma_N} = \{(u|_{\Gamma_D}, \gamma \partial_\nu u|_{\Gamma_N}) : \operatorname{div}(\gamma \nabla u) = 0 \text{ in } \Omega, u \in H^1(\Omega), \operatorname{supp}(u|_{\partial\Omega}) \subset \Gamma_D\}.$$

*MSC2010:* primary 35R30; secondary 35J10, 58J32.

*Keywords:* Calderón problem, partial data, inverse problem.

The Calderón problem with partial data is to determine the conductivity  $\gamma$  from the knowledge of  $C_\gamma^{\Gamma_D, \Gamma_N}$  for possibly very small sets  $\Gamma_D, \Gamma_N$ . Here  $\partial_\nu$  is the normal derivative, and the conormal derivative  $\gamma \partial_\nu u|_{\partial\Omega}$  is interpreted in the weak sense as an element of  $H^{-1/2}(\partial\Omega)$ .

A closely related problem is to determine a potential  $q \in L^\infty(\Omega)$  from partial boundary measurements for the Schrödinger equation, given by the partial Cauchy data set

$$C_q^{\Gamma_D, \Gamma_N} = \{(u|_{\Gamma_D}, \partial_\nu u|_{\Gamma_N}) : (-\Delta + q)u = 0 \text{ in } \Omega, u \in H_\Delta(\Omega), \text{supp}(u|_{\partial\Omega}) \subset \Gamma_D\}.$$

Here we use the space

$$H_\Delta(\Omega) = \{u \in L^2(\Omega) : \Delta u \in L^2(\Omega)\},$$

and the trace  $u|_{\partial\Omega}$  and normal derivative  $\partial_\nu u|_{\partial\Omega}$  are in  $H^{-1/2}(\partial\Omega)$  and  $H^{-3/2}(\partial\Omega)$ ; see [Bukhgeim and Uhlmann 2002]. Above, one thinks of  $u|_{\partial\Omega}$  as Dirichlet data prescribed only on  $\Gamma_D$ , and one measures the Neumann data of the corresponding solution on  $\Gamma_N$ . If  $\Lambda_\gamma : H^{1/2}(\Omega) \rightarrow H^{-1/2}(\partial\Omega)$  is the Dirichlet-to-Neumann map (DN map) given by

$$\Lambda_\gamma : u|_{\partial\Omega} \mapsto \gamma \partial_\nu u|_{\partial\Omega}, \quad \text{where } u \in H^1(\Omega) \text{ solves } \text{div}(\gamma \nabla u) = 0 \text{ in } \Omega,$$

then the partial Cauchy data set is a restriction of the graph of  $\Lambda_\gamma$ ,

$$C_\gamma^{\Gamma_D, \Gamma_N} = \{(f|_{\Gamma_D}, \Lambda_\gamma f|_{\Gamma_N}) : f \in H^{1/2}(\partial\Omega), \text{supp}(f) \subset \Gamma_D\}.$$

A similar interpretation is valid for  $C_q^{\Gamma_D, \Gamma_N}$  provided that 0 is not a Dirichlet eigenvalue of  $-\Delta + q$  in  $\Omega$ .

The problems above are well studied questions in the theory of inverse problems. The case of full data ( $\Gamma_D = \Gamma_N = \partial\Omega$ ) has received the most attention. Major results include [Sylvester and Uhlmann 1987; Haberman and Tataru 2013] in dimensions  $n \geq 3$  and [Nachman 1996; Astala and Päivrinta 2006; Bukhgeim 2008] in the case  $n = 2$ . In particular, it is known that the set  $C_\gamma^{\partial\Omega, \partial\Omega}$  determines uniquely a conductivity  $\gamma \in C^1(\bar{\Omega})$  if  $n \geq 3$  and a conductivity  $\gamma \in L^\infty(\Omega)$  if  $n = 2$ . These results are based on the method of complex geometrical optics solutions developed in [Sylvester and Uhlmann 1987] for  $n \geq 3$  and in [Nachman 1996; Bukhgeim 2008] in the case  $n = 2$ .

The partial data question where the sets  $\Gamma_D$  or  $\Gamma_N$  may not be the whole boundary has also attracted considerable attention. We mention here four approaches, each of which gives a slightly different partial data result. Formulated in terms of the Schrödinger problem, it is known that  $C_q^{\Gamma_D, \Gamma_N}$  determines  $q$  in  $\Omega$  in the following cases:

- (1)  $n \geq 3$ , the set  $\Gamma_D$  is possibly very small, and  $\Gamma_N$  is slightly larger than  $\partial\Omega \setminus \bar{\Gamma}_D$ ; proved by Kenig, Sjöstrand, and Uhlmann [Kenig et al. 2007].
- (2)  $n \geq 3$  and  $\Gamma_D = \Gamma_N = \Gamma$ , and  $\partial\Omega \setminus \Gamma$  is either part of a hyperplane or part of a sphere; proved in [Isakov 2007].
- (3)  $n = 2$  and  $\Gamma_D = \Gamma_N = \Gamma$ , where  $\Gamma$  can be an arbitrary open subset of  $\partial\Omega$ ; proved by Imanuvilov, Uhlmann, and Yamamoto [Imanuvilov et al. 2010].
- (4)  $n \geq 2$ , linearized partial data problem,  $\Gamma_D = \Gamma_N = \Gamma$ , where  $\Gamma$  can be an arbitrary open subset of  $\partial\Omega$ ; proved by dos Santos Ferreira, Kenig, Sjöstrand, and Uhlmann [Ferreira 2009b].

Approaches (1)–(3) also give a partial data result of determining  $\gamma$  from  $C_{\gamma}^{\Gamma_D, \Gamma_N}$  with the same assumptions on the dimension and the sets  $\Gamma_D, \Gamma_N$ . In (4), the linearized partial data problem is to show injectivity of the Fréchet derivative of  $\Lambda_q$  at  $q = 0$  instead of injectivity of the full map  $q \mapsto \Lambda_q$ , when restricted to the sets  $\Gamma_D$  and  $\Gamma_N$ .

It is interesting that, although each of the four approaches is based on a version of complex geometrical optics solutions, the approaches are distinct in the sense that none of the above results is contained in any of the others. The result in [Kenig et al. 2007] uses Carleman estimates with boundary terms, given for special limiting weights, that allow one to control the solutions on parts of the boundary, whereas [Isakov 2007] is based on the full data arguments of [Sylvester and Uhlmann 1987] and a reflection argument. The result in [Imanuvilov et al. 2010] is a strong one that only requires Dirichlet and Neumann data on any small set, but the method involves complex analysis and Carleman weights with critical points and does not obviously extend to higher dimensions. Finally, [Ferreira 2009b] is based on analytic microlocal analysis but is so far restricted to the linearized problem.

Nevertheless, given that there exist several approaches to the same problem, one expects that a combination of ideas from different approaches might lead to improved partial data results. In this paper we unify the Carleman estimate approach of [Kenig et al. 2007] and the reflection approach of [Isakov 2007], and, in fact, we obtain the main results of both these papers as special cases.

The method also allows us to improve both approaches. Concerning [Isakov 2007], we are able to relax the hypothesis on the inaccessible part  $\Gamma_i = \partial\Omega \setminus \Gamma$  of the boundary: instead of requiring  $\Gamma_i$  to be completely flat (or spherical), we can deal with  $\Gamma_i$  that satisfy a flatness condition only in one direction. Compared with [Kenig et al. 2007], we remove the need for measurements on certain parts of the boundary that are flat in one direction; and, in certain cases where  $\partial\Omega$  may not have any symmetries, we eliminate the overlap of  $\Gamma_D$  and  $\Gamma_N$  needed in [Kenig et al. 2007]. The method eventually boils down to inverting geodesic ray transforms (possibly for broken geodesics). In some cases the invertibility of the ray transform is known, but in other cases it is not, and in these cases we obtain a reduction from the Calderón problem with partial data to integral geometry problems of independent interest.

The survey [Kenig and Salo 2013] describes earlier results on the Calderón problem with partial data and also the results in the present paper. However, we also list here some further references for partial data results, first for the case  $n \geq 3$ . The Carleman estimate approach was initiated in [Bukhgeim and Uhlmann 2002; Kenig et al. 2007]. Based on this approach, there are low regularity results [Knudsen 2006; Zhang 2012], results for other scalar equations [Ferreira 2007; Knudsen and Salo 2007; Chung 2012] and systems [Salo and Tzou 2010; Chung et al. 2013], stability results [Heck and Wang 2006], and reconstruction results [Nachman and Street 2010]. The reflection approach was introduced in [Isakov 2007], and has been employed for the Maxwell system [Caro et al. 2009]. Partial data results for slab geometries are given in [Li and Uhlmann 2010; Krupchyk et al. 2012]. Also, at the same time as this preprint was first submitted, a preprint of Imanuvilov and Yamamoto [2013a] appeared that independently proves a result similar to that in Section 3A in this paper.

In two dimensions, the main partial data result is that of [Imanuvilov et al. 2010], which has been extended in [Imanuvilov et al. 2011a; 2011b; Imanuvilov and Yamamoto 2012a; 2012b] to, respectively,

more general equations, combinations of measurements on disjoint sets, less regular coefficients, and certain systems. An earlier result is [Astala et al. 2005]. In the case of Riemann surfaces with boundary, corresponding partial data results are given in [Guillarmou and Tzou 2011a; 2011b; Albin et al. 2013]. See also the surveys [Guillarmou and Tzou 2013; Imanuvilov and Yamamoto 2013b].

In the case when the conductivity is known near the boundary, the partial data problem can be reduced to the full data problem [Ammari and Uhlmann 2004; Alessandrini and Kim 2012; Hyvönen et al. 2012]. Also, we remark that in the corresponding problem for the wave equation, it has been known for a long time (see [Katchalov et al. 2001]) that measuring the Dirichlet and Neumann data of waves on an arbitrary open subset of the boundary is sufficient to determine the coefficients uniquely up to natural gauge transforms. Partial results for the case where Dirichlet and Neumann data are measured on disjoint sets are in [Lassas and Oksanen 2010; 2012].

The structure of this paper is as follows. Section 2 states our main partial data results in the setting of Riemannian manifolds, and Section 3 considers some consequences of the Calderón problem with partial data in Euclidean space. Section 4 gives a Carleman estimate that is used to control solutions on parts of the boundary, and Section 5 discusses a reflection approach that can be used as an alternative to Carleman estimates in some cases. In Section 6 we give the proofs of the local uniqueness results for simple transversal manifolds, based on complex geometrical optics solutions involving WKB type quasimodes. In Section 7 we discuss a more sophisticated quasimode construction based on reflected Gaussian beams, and in Section 8 we show how complex geometrical optics solutions involving reflected Gaussian beam quasimodes can be used to recover the broken ray transform of a potential from partial Cauchy data.

## 2. Statement of results

Our method is based on ideas developed for the anisotropic Calderón problem in [Ferreira 2009a], and even though much of the motivation comes from the Calderón problem with partial data in Euclidean domains, it is convenient to formulate our main results in the setting of manifolds. The Riemannian geometry notation we use is mostly that of [Ferreira 2009a].

**Definition.** Let  $(M, g)$  be a compact oriented Riemannian manifold with  $C^\infty$  boundary, and let  $n = \dim(M) \geq 3$ .

1. We say that  $(M, g)$  is *conformally transversally anisotropic* (or CTA) if

$$(M, g) \in (\mathbb{R} \times \widehat{M}_0, g), \quad g = c(e \oplus g_0),$$

where  $(\widehat{M}_0, g_0)$  is some compact  $(n - 1)$ -dimensional manifold with boundary,  $e$  is the Euclidean metric on the real line, and  $c$  is a smooth positive function in the cylinder  $\mathbb{R} \times \widehat{M}_0$ .

2. We say that  $(M, g)$  is *admissible* if it is CTA and additionally the transversal manifold  $(\widehat{M}_0, g_0)$  is *simple*, meaning that the boundary  $\partial\widehat{M}_0$  is strictly convex (the second fundamental form is positive definite) and for each  $p \in \widehat{M}_0$ , the exponential map  $\exp_p$  is a diffeomorphism from its maximal domain of definition in  $T_p\widehat{M}_0$  onto  $\widehat{M}_0$ .

The uniqueness results in [Ferreira 2009a] were given for admissible manifolds. In this paper we will give results both for admissible and CTA manifolds. In the main results, we will also assume that there is a compact  $(n - 1)$ -dimensional manifold  $(M_0, g_0)$  with smooth boundary such that

$$(M, g) \subset (\mathbb{R} \times M_0, g) \Subset (\mathbb{R} \times \widehat{M}_0, g), \quad g = c(e \oplus g_0), \tag{2-1}$$

and the following intersection is nonempty:

$$\partial M \cap (\mathbb{R} \times \partial M_0) \neq \emptyset.$$

Under some conditions, it will be possible to ignore boundary measurements in the set  $\partial M \cap (\mathbb{R} \times \partial M_0)$ . In the results below, we will implicitly assume that the various manifolds satisfy (2-1), and if  $(M, g)$  is admissible, it is also assumed that  $(\widehat{M}_0, g_0)$  is simple (but  $(M_0, g_0)$  need not be simple, since its boundary may not be strictly convex).

Write  $x = (x_1, x')$  for points in  $\mathbb{R} \times \widehat{M}_0$ , where  $x_1$  is the Euclidean coordinate. The approaches of [Kenig et al. 2007; Ferreira 2009a] are based on complex geometrical optics solutions of the form  $u = e^{\tau\varphi}(m + r)$ , where  $\varphi$  is a special *limiting Carleman weight*. We refer to the latter paper for the definition and properties of limiting Carleman weights on manifolds. For present purposes, we only mention that the functions  $\varphi(x) = \pm x_1$  are natural limiting Carleman weights in the cylinder  $(\mathbb{R} \times \widehat{M}_0, g)$ .

The weight  $\varphi(x) = x_1$  allows us to decompose the boundary  $\partial M$  as the disjoint union

$$\partial M = \partial M_+ \cup \partial M_- \cup \partial M_{\text{tan}},$$

where

$$\partial M_{\pm} = \{x \in \partial M : \pm \partial_\nu \varphi(x) > 0\} \quad \text{and} \quad \partial M_{\text{tan}} = \{x \in \partial M : \partial_\nu \varphi(x) = 0\}.$$

Here the normal derivative is understood with respect to the metric  $g$ . Note that  $\partial_\nu \varphi = 0$  on  $\mathbb{R} \times \partial M_0$  whenever  $(M_0, g_0) \Subset (\widehat{M}_0, g_0)$ . We think of  $\partial M_{\text{tan}}$  as being flat in one direction (the direction of the gradient of  $\varphi$ ). For the sake of definiteness, the sets  $\partial M_{\pm} = \partial M_{\pm}(\varphi)$  will refer to the weight  $\varphi(x) = x_1$  in this section, but all results remain true when  $\partial M_+$  and  $\partial M_-$  are interchanged (this amounts to replacing the weight  $x_1$  by  $-x_1$ ).

Next we give the local results for the Calderón problem with partial data on manifolds. In these results we say that a unit speed geodesic  $\gamma : [0, L] \rightarrow M_0$  is *nontangential* if its endpoints are on  $\partial M_0$ , the vectors  $\dot{\gamma}(0), \dot{\gamma}(L)$  are nontangential, and  $\gamma(t) \in M_0^{\text{int}}$  for  $0 < t < L$ . We also define the partial Cauchy data set as

$$C_{g,q}^{\Gamma_D, \Gamma_N} = \{(u|_{\Gamma_D}, \partial_\nu u|_{\Gamma_N}) : (-\Delta_g + q) = 0 \text{ in } M, u \in H_{\Delta_g}(M), \text{supp}(u|_{\partial M}) \subset \Gamma_D\},$$

where  $H_{\Delta_g}(M) = \{u \in L^2(M) : \Delta_g u \in L^2(M)\}$  and  $u|_{\partial M} \in H^{-1/2}(\partial M), \partial_\nu u|_{\partial M} \in H^{-3/2}(\partial M)$  by the same arguments as in [Bukhgeim and Uhlmann 2002].

To explain the results, it is convenient to think in terms of the following special case.

**Example.** Let  $M = M_{\text{left}} \cup M_{\text{mid}} \cup M_{\text{right}}$  be a compact manifold with boundary consisting of three parts:  $M_{\text{mid}} = [a, b] \times M_0$  for some compact manifold  $(M_0, g_0)$  with boundary,  $M_{\text{left}} \subset \{x_1 < a\} \times M_0$ , and

$M_{\text{right}} \subset \{x_1 > b\} \times M_0$ . We also assume that  $\partial M_- = M_{\text{left}} \cap \partial M$  and  $\partial M_+ = M_{\text{right}} \cap \partial M$ . In this case  $\partial M_{\text{tan}} = [a, b] \times \partial M_0$ .

The methods developed in this paper suggest that it should suffice to measure Neumann data on  $\partial M_+$  for Dirichlet data supported in  $\partial M_-$ , with no measurements required on  $\partial M_{\text{tan}}$ . However, in the results below we need a part  $\Gamma_a \subset \partial M_{\text{tan}}$  that is accessible to measurements, and  $\Gamma_i = \partial M_{\text{tan}} \setminus \Gamma_a$  is the inaccessible part. Suppose for simplicity that

$$\Gamma_a = [a, b] \times E, \quad \Gamma_i = [a, b] \times (\partial M_0 \setminus E)$$

for some nonempty open subset  $E$  of  $\partial M_0$ .

In this setting, [Theorem 2.1](#) implies that from Neumann data measured near  $\partial M_+ \cup \Gamma_a$  with Dirichlet data input near  $\partial M_- \cup \Gamma_a$ , one can determine certain integrals of the potential  $q$  in the set

$$\mathbb{R} \times \bigcup_{\gamma} \gamma([0, L]),$$

where the union is over all nontangential geodesics in  $M_0$  with endpoints on  $E$ . Moreover, if the local ray transform is injective in this set in a suitable sense, one can determine the potential in this set by [Theorem 2.2](#). [Theorem 2.4](#) shows that one can go beyond this set and extract information about integrals of  $q$  over all nontangential broken rays with endpoints on  $E$ , and [Theorem 2.3](#) gives a global uniqueness result in the case where  $\partial M_{\text{tan}}$  has zero measure.

**Theorem 2.1.** *Let  $(M, g)$  be an admissible manifold as in (2-1), and let  $q_1, q_2 \in C(M)$ . Let  $\Gamma_i$  be a closed subset of  $\partial M_{\text{tan}}$ , and suppose that, for some nonempty open subset  $E$  of  $\partial M_0$ , one has*

$$\Gamma_i \subset \mathbb{R} \times (\partial M_0 \setminus E).$$

Let  $\Gamma_a = \partial M_{\text{tan}} \setminus \Gamma_i$ , and assume that

$$C_{g, q_1}^{\Gamma_D, \Gamma_N} = C_{g, q_2}^{\Gamma_D, \Gamma_N},$$

where  $\Gamma_D$  and  $\Gamma_N$  are any open sets in  $\partial M$  such that  $\Gamma_D \supset \partial M_- \cup \Gamma_a$  and  $\Gamma_N \supset \partial M_+ \cup \Gamma_a$ .

Given any nontangential geodesic  $\gamma : [0, L] \rightarrow M_0$  with endpoints on  $E$ , and given any real number  $\lambda$ , one has

$$\int_0^L e^{-2\lambda t} (c(q_1 - q_2))^{\wedge}(2\lambda, \gamma(t)) dt = 0.$$

Here  $q_1 - q_2$  is extended by zero outside  $M$ , and  $(\cdot)^{\wedge}$  denotes the Fourier transform in the  $x_1$  variable.

The previous theorem allows us to conclude uniqueness of potentials in sets where the local ray transform is injective in the following sense.

**Definition.** Let  $(M_0, g_0)$  be a compact oriented manifold with smooth boundary, and let  $O$  be an open subset of  $M_0$ . We say that the local ray transform is injective on  $O$  if any function  $f \in C(M_0)$  with

$$\int_{\gamma} f dt = 0 \quad \text{for all nontangential geodesics } \gamma \text{ contained in } O$$

must satisfy  $f|_O = 0$ .



**Theorem 2.2.** *Assume the conditions in Theorem 2.1. Then  $q_1 = q_2$  in  $M \cap (\mathbb{R} \times O)$  for any open subset  $O$  of  $M_0$  such that the local ray transform is injective on  $O$  and  $O \cap \partial M_0 \subset E$ .*

The local ray transform is known to be injective in the next three cases (the second case will be used in Section 3):

1.  $(M_0, g_0) = (\bar{\Omega}_0, e)$ , where  $\Omega_0 \subset \mathbb{R}^{n-1}$  is a bounded domain with  $C^\infty$  boundary,  $e$  is the Euclidean metric,  $E$  is an open subset of  $\partial\Omega_0$ , and  $O$  is the intersection of  $\bar{\Omega}_0$  with the union of all hyperplanes in  $\mathbb{R}^{n-1}$  that have  $\partial\Omega_0 \setminus E$  on one side. The complement of this union is the intersection of half-spaces and is thus convex. If the integral of  $f \in C(\bar{\Omega}_0)$ , extended by zero to  $\mathbb{R}^{n-1}$ , vanishes over all line segments in  $O$ , the integral over all hyperplanes that do not meet  $\partial\Omega_0 \setminus E$  also vanishes, and it follows from the Helgason support theorem [1999] that the local ray transform is injective on  $O$ .
2.  $(M_0, g_0) \in (\tilde{M}_0, g_0)$  are simple manifolds with real-analytic metric, and  $\tilde{\mathcal{F}}$  is an open set of nontangential geodesics in  $(\tilde{M}_0, g_0)$  such that any curve in  $\tilde{\mathcal{F}}$  can be deformed to a point on  $\partial\tilde{M}_0$  through curves in  $\tilde{\mathcal{F}}$ . In such a case, by a result of Krishnan [2009] the local ray transform is injective on the set  $O$  of all points in  $M_0$  that lie on some geodesic in  $\tilde{\mathcal{F}}$ .
3. If  $\dim(M_0) \geq 3$  and if  $\partial M_0$  is strictly convex at a point  $p \in \partial M_0$ , then  $p$  has a neighborhood  $O$  in  $M_0$  on which the local ray transform is injective. This is a result from [Uhlmann and Vasy 2012].

In Theorem 2.2, if the nontangential geodesics with endpoints on  $E$  cover a dense subset  $O$  of  $M_0$  and if the local ray transform is injective in  $O$ , we obtain a global uniqueness result stating that  $q_1 = q_2$  in  $M$ . An example of such a result under a concavity assumption is given in Section 3F.

The method for proving Theorems 2.1 and 2.2 also allows us to reduce the overlap for  $\Gamma_D$  and  $\Gamma_N$  needed in [Kenig et al. 2007]. An example of such a result is the following (a similar result was also proved in [Imanuvilov and Yamamoto 2013b]).

**Theorem 2.3.** *Let  $(M, g)$  be an admissible manifold and assume that  $q_1, q_2 \in C(M)$ . If  $\partial M_{\tan}$  has zero measure in  $\partial M$ , then*

$$C_{g, q_1}^{\partial M_-, \partial M_+} = C_{g, q_2}^{\partial M_-, \partial M_+} \implies q_1 = q_2.$$

Next we wish to gather information on the potentials beyond the set that can be reached by transversal geodesics with endpoints on  $E$ . To do this, we will use broken geodesics in the transversal manifold that go inside  $M_0$ , reflect finitely many times, and eventually return to  $E$ .

**Definition.** Let  $(M_0, g_0)$  be a compact manifold with boundary.

- (a) We call a continuous curve  $\gamma : [a, b] \rightarrow M_0$  a *broken ray* if  $\gamma$  is obtained by following unit speed geodesics that are reflected according to geometrical optics (angle of incidence equals angle of reflection) whenever they hit a point of  $\partial M_0$ .
- (b) A broken ray  $\gamma : [0, L] \rightarrow M_0$  is called *nontangential* if  $\dot{\gamma}(t)$  is nontangential whenever  $\gamma(t) \in \partial M_0$ , and additionally all points of reflection are distinct.

The next theorem is a generalization of [Theorem 2.1](#) in the sense that it allows arbitrary transversal manifolds and recovers integrals over all nontangential broken rays (instead of just nontangential geodesics) with endpoints on  $E$ . However, it is stated with a weaker partial data condition.

**Theorem 2.4.** *Let  $(M, g)$  be a CTA manifold as in (2-1), and let  $q_1, q_2 \in C(M)$ . Let  $\Gamma_i$  be a closed subset of  $\partial M_{\text{tan}}$ , and suppose that, for some nonempty open subset  $E$  of  $\partial M_0$ , one has*

$$\Gamma_i \subset \mathbb{R} \times (\partial M_0 \setminus E).$$

Let  $\Gamma_a = \partial M_{\text{tan}} \setminus \Gamma_i$ , and assume that

$$C_{g, q_1}^{\Gamma_D, \Gamma_N} = C_{g, q_2}^{\Gamma_D, \Gamma_N},$$

where  $\Gamma_D = \Gamma_N = \Gamma$  for some neighborhood  $\Gamma$  of the set  $\overline{\partial M_+ \cup \partial M_- \cup \Gamma_a}$  in  $\partial M$ .

Given any nontangential broken ray  $\gamma : [0, L] \rightarrow M_0$  with endpoints on  $E$ , and given any real number  $\lambda$ , one has

$$\int_0^L e^{-2\lambda t} (c(q_1 - q_2))^\wedge(2\lambda, \gamma(t)) dt = 0.$$

Here  $q_1 - q_2$  is extended by zero outside  $M$ , and  $(\cdot)^\wedge$  denotes the Fourier transform in the  $x_1$  variable.

It is natural to ask whether a function in  $M_0$  is determined by its integrals over broken rays with endpoints in some subset  $E$  of  $\partial M_0$  (that is, whether the broken ray transform is injective). Combined with [Theorem 2.4](#) and with the proof of [Theorem 2.2](#), such a result would imply unique recovery of the potential in the whole manifold  $M$ . However, it seems that there are very few results in this direction, except for the case where  $E$  is the whole boundary and the question reduces to the injectivity of the usual ray transform; see [[Sharafutdinov 1994](#)].

[Eskin \[2004\]](#) has proved injectivity in the case of Euclidean broken rays reflecting off several convex obstacles, with  $E$  being the boundary of a smooth domain enclosing all the obstacles, if the obstacles satisfy additional restrictions (in particular, the obstacles must have corner points and they cannot be smooth). [Hubenthal \[2013a; 2013b\]](#) and [Ilmavirta \[2013a; 2013b; 2013c\]](#) have given partial results for the broken ray transform in special geometries. See also [[Florescu et al. 2011](#); [Lozev 2013](#)] for related results. However, the following question seems to be open even in convex Euclidean domains except when  $E = \partial M_0$ .

**Question.** *Let  $(M_0, g_0)$  be a simple manifold, let  $E$  be a nonempty open subset of  $\partial M_0$ , and assume that  $f \in C(M_0)$  satisfies*

$$\int_0^L f(\gamma(t)) dt = 0$$

for all nontangential broken rays  $\gamma : [0, L] \rightarrow M_0$  with endpoints on  $E$ . Does this imply that  $f = 0$ ?

### 3. The Euclidean case

In this section, we indicate some consequences of the previous results to the Calderón problem with partial data in Euclidean space. We assume that  $\Omega \subset \mathbb{R}^3$  is a bounded domain with smooth boundary

equipped with the Euclidean metric  $g = e$ , and  $q_1, q_2 \in C(\bar{\Omega})$ . We also assume that

$$C_{q_1}^{\Gamma, \Gamma} = C_{q_2}^{\Gamma, \Gamma},$$

where  $\Gamma$  is some strict open subset of  $\partial\Omega$ . Write

$$\Gamma_i = \partial\Omega \setminus \Gamma$$

for the inaccessible part of the boundary. The results in this section show that in cases where  $\Gamma_i$  satisfies certain geometric restrictions, it is possible to conclude that

$$q_1 = q_2 \quad \text{in } \bar{\Omega} \cap (\mathbb{R} \times O),$$

where the sets  $O \subset \mathbb{R}^2$  will be described below.

**Remark.** We also obtain results for the conductivity equation by making a standard reduction to the Schrödinger equation. More precisely, if  $\gamma_1, \gamma_2 \in C^2(\bar{\Omega})$  are positive functions such that  $C_{\gamma_1}^{\Gamma, \Gamma} = C_{\gamma_2}^{\Gamma, \Gamma}$ , the corresponding DN maps satisfy

$$\Lambda_{\gamma_1} f|_{\Gamma} = \Lambda_{\gamma_2} f|_{\Gamma} \quad \text{for } f \in H^{1/2}(\partial\Omega) \text{ with } \text{supp}(f) \subset \Gamma.$$

Boundary determination [Kohn and Vogelius 1984; Sylvester and Uhlmann 1988] implies that

$$\gamma_1|_{\Gamma} = \gamma_2|_{\Gamma}, \quad \partial_\nu \gamma_1|_{\Gamma} = \partial_\nu \gamma_2|_{\Gamma}.$$

Writing  $q_j = \Delta \gamma_j^{1/2} / \gamma_j^{1/2}$ , the relation

$$\Lambda_{q_j} f = \gamma_j^{-1/2} \Lambda_{\gamma_j} (\gamma_j^{-1/2} f) + \frac{1}{2} \gamma_j^{-1} (\partial_\nu \gamma_j) f|_{\partial\Omega}$$

and the above conditions imply that the DN maps  $\Lambda_{q_j}$  for the Schrödinger equations satisfy

$$\Lambda_{q_1} f|_{\Gamma} = \Lambda_{q_2} f|_{\Gamma} \quad \text{for } f \in H^{1/2}(\partial\Omega) \text{ with } \text{supp}(f) \subset \Gamma.$$

Thus  $C_{q_1}^{\Gamma, \Gamma} = C_{q_2}^{\Gamma, \Gamma}$ , and we obtain that

$$q_1 = q_2 \quad \text{in } \bar{\Omega} \cap (\mathbb{R} \times O).$$

Write  $q = q_1 = q_2$  in  $\bar{\Omega} \cap (\mathbb{R} \times O)$ . Then  $\gamma_1^{1/2}$  and  $\gamma_2^{1/2}$  are both solutions of  $(-\Delta + q)u = 0$  in  $\bar{\Omega} \cap (\mathbb{R} \times O)$  having identical Cauchy data on  $\Gamma$ . It follows that  $\gamma_1 = \gamma_2$  in any connected component of  $\bar{\Omega} \cap (\mathbb{R} \times O)$  whose intersection with  $\Gamma$  contains a nonempty open subset of  $\partial\Omega$ .

In the following we will use some general facts on limiting Carleman weights from [Ferreira 2009a], where it was proved that any limiting Carleman weight in  $\mathbb{R}^3$  has, up to translation, rotation and scaling, one of the following six forms:

$$x_1, \quad \log|x|, \quad \arg(x_1 + ix_2), \quad \frac{x_1}{|x|^2}, \quad \log \frac{|x + e_1|^2}{|x - e_1|^2}, \quad \arg(e^{i\theta}(x + ie_1)^2).$$

Here  $\theta \in [0, 2\pi)$ , and the argument function is defined by

$$\arg(z) = 2 \arctan \frac{\text{Im}(z)}{|z| + \text{Re}(z)}, \quad z \in \mathbb{C} \setminus \{t \in \mathbb{R} : t \leq 0\}.$$

It was also proved in Section 2 of [Ferreira 2009a] that if  $\varphi$  is a limiting Carleman weight near  $(\bar{\Omega}, e)$ , then  $\nabla_{\tilde{g}}\varphi$  is a unit parallel vector field near  $(\bar{\Omega}, \tilde{g})$  where

$$\tilde{g} = c^{-1}e, \quad c = |\nabla_e\varphi|_e^{-2}.$$

Furthermore, by the proof of Lemma A.5 of the same reference, if  $(y_1, y')$  are coordinates so that  $\nabla_{\tilde{g}}\varphi = \partial_{y_1}$  and if the coordinates  $y'$  parametrize a 2-dimensional manifold  $S$  such that  $\nabla_{\tilde{g}}\varphi$  is orthogonal to  $S$  with respect to the  $\tilde{g}$  metric, then the metric has the form

$$\tilde{g}(y_1, y') = \begin{pmatrix} 1 & 0 \\ 0 & \tilde{g}_0(y') \end{pmatrix},$$

where  $\tilde{g}_0$  is the metric on  $S$  induced by  $\tilde{g}$ .

**3A. Cylindrical sets.** This case corresponds to the limiting Carleman weight  $\varphi(x) = x_1$ . Suppose that  $\Omega \subset \mathbb{R} \times \Omega_0$ , where  $\Omega_0$  is a bounded domain with smooth boundary in  $\mathbb{R}^2$ . Let  $E$  be an open subset of  $\partial\Omega_0$ , and assume that

$$\Gamma_i \subset \mathbb{R} \times (\partial\Omega_0 \setminus E).$$

If  $\Omega_0$  has strictly convex boundary, Theorem 2.2 and the result of [Krishnan 2009] imply that

$$q_1 = q_2 \quad \text{in } \bar{\Omega} \cap (\mathbb{R} \times O),$$

where  $O$  is the intersection of  $\bar{\Omega}_0$  with the union of all lines in  $\mathbb{R}^2$  that have  $\partial\Omega_0 \setminus E$  on one side.

The above conclusion holds true also when  $\Omega_0$  does not have strictly convex boundary. To see this, let  $\Omega_0 \Subset B \Subset \tilde{B}$ , where  $B$  and  $\tilde{B}$  are balls. The extensions of the line segments in  $O$  to  $\tilde{B}$  form a class  $\tilde{\mathcal{F}}$  such that any curve in  $\tilde{\mathcal{F}}$  can be deformed to a point through curves in  $\tilde{\mathcal{F}}$ . It is then enough to extend  $q_1 - q_2$  by zero to  $\mathbb{R} \times \tilde{B}$ , and to use the proof of Theorem 2.2 with  $M_0$  replaced by  $\tilde{B}$ , together with [Krishnan 2009].

**3B. Conical sets.** Consider the limiting Carleman weight  $\varphi(x) = \log|x|$ . Suppose that  $\bar{\Omega} \subset \{x_3 > 0\}$ , let  $(S^2, g_0)$  be the sphere with its standard metric, let  $S^2_+ = \{\omega \in S^2 : \omega_3 > 0\}$ , and let  $(M_0, g_0)$  be a compact submanifold of  $(S^2_+, g_0)$  with smooth boundary. Let  $E$  be an open subset of  $\partial M_0$ , and assume that

$$\Gamma_i \subset \{r\omega : r > 0, \omega \in \partial M_0 \setminus E\}.$$

We have  $c = |\nabla\varphi|^{-2} = |x|^2$  and  $\tilde{g} = |x|^{-2}e$ ,  $\nabla_{\tilde{g}}\varphi = x$ . Choose coordinates so that

$$y_1 = \log|x|, \quad y' = x/|x|.$$

The coordinates  $y'$  parametrize the manifold  $S^2$  and the metric  $\tilde{g}_0$  on  $S^2$  induced by  $\tilde{g}$  is just the standard metric  $g_0$ . The discussion in the beginning of this section shows that

$$\tilde{g}(y_1, y') = \begin{pmatrix} 1 & 0 \\ 0 & g_0(y') \end{pmatrix}.$$

Now  $(M_0, g_0)$  is contained in some simple submanifold  $(\widehat{M}_0, g_0)$  of the hemisphere  $(S_+^2, g_0)$  (just remove a neighborhood of the equator). Since geodesics in  $S_+^2$  are restrictions of great circles, [Theorem 2.2](#) and the local injectivity result [[Krishnan 2009](#)] imply, as in [Section 3A](#), that

$$q_1 = q_2 \quad \text{in } \bar{\Omega} \cap \{r\omega : r > 0, \omega \in O\},$$

where  $O$  is the union of all great circle segments in  $S_+^2$  such that  $\partial M_0 \setminus E$  is on one side of the hyperplane containing the great circle segment.

**3C. Surfaces of revolution.** Let  $\Omega \subset \mathbb{R}^3 \setminus \{x : x_1 \leq 0\}$ , and consider the limiting Carleman weight

$$\varphi(x) = \arg(x_1 + ix_2).$$

Then

$$\nabla\varphi = \left( \frac{-x_2}{x_1^2 + x_2^2}, \frac{x_1}{x_1^2 + x_2^2}, 0 \right)$$

and

$$c = x_1^2 + x_2^2, \quad \tilde{g} = \frac{1}{x_1^2 + x_2^2}e, \quad \nabla_{\tilde{g}}\varphi = (-x_2, x_1, 0).$$

We make the change of coordinates valid near  $\bar{\Omega}$ ,

$$y_1 = \arg(x_1 + ix_2), \quad y_2 = \sqrt{x_1^2 + x_2^2}, \quad y_3 = x_3.$$

The coordinates  $y'$  parametrize the manifold  $S = \{(x_1, 0, x_3) : x_1 > 0\}$  and  $\nabla_{\tilde{g}}\varphi$  is orthogonal to  $S$ . Furthermore, we may also think of  $S$  as the set  $\{(0, y_2, y_3) : y_2 > 0\}$ , and the metric on  $S$  induced by  $\tilde{g}$  is the hyperbolic metric  $\tilde{g}_0 = (1/y_2^2)e$ . The discussion in the beginning of this section shows that

$$\tilde{g}(y_1, y') = \begin{pmatrix} 1 & 0 \\ 0 & \tilde{g}_0(y') \end{pmatrix}.$$

Let  $(M_0, g_0)$  be a compact submanifold of  $S$  with smooth boundary, let  $E$  be an open subset of  $\partial M_0$ . We think of  $M_0$  as lying in  $\{(x_1, 0, x_3) : x_1 > 0\}$ . Now, assume that

$$\Gamma_i \subset \{R_\theta(\partial M_0 \setminus E) : \theta \in (-\pi, \pi)\},$$

where  $R_\theta x = (\tilde{R}_\theta(x_1, x_2)^t, x_3)^t$  and  $\tilde{R}_\theta$  rotates vectors in  $\mathbb{R}^2$  by angle  $\theta$  counterclockwise. That is, we assume that the inaccessible part  $\Gamma_i$  is contained in a surface of revolution obtained by rotating the boundary curve  $\partial M_0 \setminus E$ .

Now, the geodesics in  $S$  (and, after restriction, also in  $M_0$ ) have either the form

$$(y_2(t), y_3(t)) = (R \sin t, R \cos t + \alpha),$$

where  $t \in (0, \pi)$ ,  $R > 0$ , and  $\alpha \in \mathbb{R}$ , or the form  $(y_2(t), y_3(t)) = (t, \alpha)$ , where  $t > 0$  and  $\alpha \in \mathbb{R}$  (these are not unit speed parametrizations). In the  $x$  coordinates, these are either the half circles in the  $\{x_2 = 0\}$  plane given by

$$(x_1(t), x_2(t), x_3(t)) = (R \sin t, 0, R \cos t + \alpha), \quad t \in (0, \pi),$$

or the lines

$$(x_1(t), x_2(t), x_3(t)) = (t, 0, \alpha), \quad t > 0.$$

Enclosing  $M_0$  in some ball  $B$  in  $S$ , the manifold  $(\bar{B}, g_0)$  is simple and [Theorem 2.2](#) and [\[Krishnan 2009\]](#) imply, as in [Section 3A](#), that

$$q_1 = q_2 \quad \text{in } \bar{\Omega} \cap \{R_\theta(O) : \theta \in (-\pi, \pi)\},$$

where  $O$  is the union of all geodesics in  $S$  that have  $\partial M_0 \setminus E$  on one side.

**3D. Other limiting Carleman weights.** So far we have considered three of the six possible forms of limiting Carleman weights in  $\mathbb{R}^3$ . The fourth one,  $\varphi(x) = x_1/|x|^2$ , is the Kelvin transform of the linear weight, and corresponds to inaccessible parts of the boundary that are Kelvin transforms of cylindrical domains. In particular, if part of the cylindrical domain is on the hyperplane  $\{x_3 = 1\}$ , its Kelvin transform lies on the sphere centered at  $(0, 0, 1/2)$  with radius  $1/2$ , and we recover the result of Isakov [\[2007\]](#) for domains where the inaccessible part is part of a sphere. The corresponding results for the remaining two limiting Carleman weights do not seem so easy to state and we omit them.

**3E. Extension of Kenig, Sjöstrand, and Uhlmann’s result.** Now let  $\Omega \subset \mathbb{R}^3$  be a bounded domain with smooth boundary, assume that  $0$  is not in the convex hull of  $\bar{\Omega}$ , and let  $\varphi(x) = \log|x|$ . Define

$$\partial\Omega_\pm = \{x \in \partial\Omega : \pm\partial_\nu\varphi(x) > 0\}, \quad \partial\Omega_{\tan} = \{x \in \partial\Omega : \partial_\nu\varphi(x) = 0\}.$$

It was proved in [\[Kenig et al. 2007\]](#) that whenever  $\Gamma_D$  is a neighborhood of  $\partial\Omega_- \cup \partial\Omega_{\tan}$  and  $\Gamma_N$  is a neighborhood of  $\partial\Omega_+ \cup \partial\Omega_{\tan}$ , we have

$$C_{q_1}^{\Gamma_D, \Gamma_N} = C_{q_2}^{\Gamma_D, \Gamma_N} \implies q_1 = q_2.$$

In particular,  $\Gamma_D$  and  $\Gamma_N$  always need to overlap. This result is a consequence of the reduction given above for the logarithmic weight, [Theorem 2.1](#) (the special case where  $E = \partial\Omega_0$ , so that  $\Gamma_i = \emptyset$ ), and injectivity of the ray transform. If  $\partial\Omega_{\tan}$  has zero measure in  $\partial\Omega$ , then [Theorem 2.3](#) allows us to improve this result: we have

$$C_{q_1}^{\partial\Omega_-, \partial\Omega_+} = C_{q_2}^{\partial\Omega_-, \partial\Omega_+} \implies q_1 = q_2.$$

In this case, the sets where Dirichlet and Neumann data are measured are disjoint, but their union covers all of  $\partial\Omega$  except for a set of measure zero. The result remains true if the roles of  $\partial\Omega_+$  and  $\partial\Omega_-$  are changed.

**3F. Extension of Isakov’s result.** According to [\[Isakov 2007\]](#), the condition  $C_{q_1}^{\Gamma, \Gamma} = C_{q_2}^{\Gamma, \Gamma}$  implies  $q_1 = q_2$  in  $\Omega$  if  $\Omega \subset \{x_3 > 0\}$  and  $\Gamma_i \subset \{x_3 = 0\}$ , or if  $\Omega \subset B$  for some ball  $B$  and  $\Gamma_i \subset \partial B$ . We have already recovered these results in [Sections 3A](#) and [3D](#), since in these cases the local injectivity set  $O$  is so large that the result  $q_1 = q_2$  holds in all of  $\Omega$ . Of course, the results above also extend [\[Isakov 2007\]](#), since we can conclude at least local uniqueness for potentials when the inaccessible part of the boundary satisfies a (conformal) flatness condition in only one direction, such as being part of a cylindrical set, a conical set, or a surface of revolution.

We also get global uniqueness if the local injectivity set  $O$  is sufficiently large. For instance, if

$$\Omega \subset \mathbb{R} \times \Omega_0, \quad \Gamma_i \subset \mathbb{R} \times (\partial\Omega_0 \setminus E),$$

where  $\Omega_0$  is a bounded domain with smooth boundary and  $E$  is a nonempty open subset of  $\partial\Omega_0$ , and if the lines in  $\mathbb{R}^2$  that have  $\partial\Omega_0 \setminus E$  on one side cover a dense subset of  $\Omega_0$ , then  $q_1 = q_2$  in  $\Omega$ . One example of this situation is if

$$\Omega \subset \mathbb{R} \times \{(x_2, x_3) : x_3 > \eta(x_2)\}, \quad \Gamma_i \subset \mathbb{R} \times \{(x_2, x_3) : x_3 = \eta(x_2)\},$$

where  $\eta : \mathbb{R} \rightarrow \mathbb{R}$  is a smooth concave function.

#### 4. Carleman estimate

Let  $(M, g)$  be a CTA manifold, so  $(M, g)$  is compact with boundary and

$$(M, g) \Subset (\mathbb{R} \times M_0, g), \quad g = c(e \oplus g_0).$$

Here  $(M_0, g_0)$  is any compact  $(n - 1)$ -dimensional manifold with boundary. We wish to prove a Carleman estimate with boundary terms for the conjugated operator  $e^{\varphi/h}(-\Delta_g)e^{-\varphi/h}$  in  $M$ , where  $\varphi$  is the limiting Carleman weight  $\varphi(x) = x_1$  or  $\varphi(x) = -x_1$ , and  $h > 0$  is small. Following [Kenig et al. 2007], it is useful to consider a slightly modified weight

$$\varphi_\varepsilon = \varphi + hf_\varepsilon$$

where  $f_\varepsilon$  is a smooth real-valued function in  $M$  depending on a small parameter  $\varepsilon$ , with  $\varepsilon$  independent of  $h$ . The convexity of  $f_\varepsilon$  will lead to improved lower bounds in terms of the  $L^2(M)$  norms of  $u$  and  $h\nabla u$ . On the other hand, the sign of  $\partial_\nu \varphi_\varepsilon$  in the boundary term of the Carleman estimate will allow us to control functions on different parts of the boundary. Of special interest is the set  $\partial M_{\text{tan}}$ , where  $\partial_\nu \varphi = 0$ , and in this set we have

$$\partial_\nu \varphi_\varepsilon|_{\partial M_{\text{tan}}} = h\partial_\nu f_\varepsilon.$$

We would like to have  $\partial_\nu f_\varepsilon < 0$  on  $\partial M_{\text{tan}}$ . It is not easy to find a global convex function  $f_\varepsilon$  satisfying the last condition for a general set  $\partial M_{\text{tan}}$ . However, splitting  $f_\varepsilon$  into a convex part whose normal derivative vanishes on  $\partial M_{\text{tan}}$  and another part which ensures the correct sign on  $\partial M_{\text{tan}}$  will give the required result. We will use semiclassical conventions in the next proof; see [Ferreira 2009a, Section 4; Zworski 2012] for more details. We also write  $(v, w) = (v, w)_{L^2(M)}$ ,  $\|v\| = \|v\|_{L^2(M)}$ , and for  $\Gamma \subset \partial M$  we write  $(v, w)_\Gamma = (v, w)_{L^2(\Gamma)}$ .

**Proposition 4.1.** *Let  $(M, g)$  be as above, let  $\varphi(x) = \pm x_1$ , and let  $\kappa$  be a smooth real-valued function in  $M$  so that  $\partial_\nu \kappa = -1$  on  $\partial M$ . Also let  $q \in L^\infty(M)$ . There are constants  $\varepsilon, C_0, h_0 > 0$  with  $h_0 \leq \varepsilon/2 \leq 1$  such that, for the weight*

$$\varphi_\varepsilon = \varphi + \frac{h}{\varepsilon} \frac{\varphi^2}{2} + h\kappa,$$

where  $0 < h \leq h_0$ , one has

$$\frac{h^3}{C_0} (|\partial_v \varphi_\varepsilon| \partial_v u, \partial_v u)_{\partial M_-(\varphi_\varepsilon)} + \frac{h^2}{C_0} (\|u\|^2 + \|h \nabla u\|^2) \\ \leq \|e^{\varphi/h} (-h^2 \Delta_g + h^2 q)(e^{-\varphi/h} u)\|^2 + h^3 (|\partial_v \varphi_\varepsilon| \partial_v u, \partial_v u)_{\partial M_+(\varphi_\varepsilon)}$$

for any  $u \in C^\infty(M)$  with  $u|_{\partial M} = 0$ .

*Proof.* Since  $\varphi(x) = \pm x_1$  is a limiting Carleman weight in a manifold strictly containing  $M$ , the computations in the proof of [Ferreira 2009a, Theorem 4.1] apply and we can follow that proof. First of all, note that

$$c^{(n+2)/4} (-\Delta_g + q)u = (-\Delta_{c^{-1}g} + q_c)(c^{(n-2)/4}u),$$

where  $q_c = cq + c^{(n+2)/4} \Delta_g (c^{-(n-2)/4})$ . Thus, by replacing  $q$  with another potential, we may assume that  $c = 1$  so that  $g = e \oplus g_0$  and  $\varphi$  is a distance function in the  $g$  metric, that is,  $|\nabla_g \varphi|_g = 1$ .

Let  $P_0 = -h^2 \Delta_g$  and  $P_{0,\varphi_\varepsilon} = e^{\varphi_\varepsilon/h} P_0 e^{-\varphi_\varepsilon/h}$ . Then  $P_{0,\varphi_\varepsilon} = A + iB$ , where  $A$  and  $B$  are the formally self-adjoint operators

$$A = -h^2 \Delta_g - |\nabla \varphi_\varepsilon|^2, \quad B = -2i \langle \nabla \varphi_\varepsilon, h \nabla \cdot \rangle - ih \Delta_g \varphi_\varepsilon.$$

Assume  $u \in C^\infty(M)$  and  $u|_{\partial M} = 0$ . We have

$$\|P_{0,\varphi_\varepsilon} u\|^2 = ((A + iB)u, (A + iB)u) \\ = \|Au\|^2 + \|Bu\|^2 + i(Bu, Au) - i(Au, Bu) \\ = \|Au\|^2 + \|Bu\|^2 + (i[A, B]u, u) - ih^2 (Bu, \partial_v u)_{\partial M} \\ = \|Au\|^2 + \|Bu\|^2 + (i[A, B]u, u) - 2h^3 ((\partial_v \varphi_\varepsilon) \partial_v u, \partial_v u)_{\partial M}.$$

Define

$$\tilde{\varphi}_\varepsilon(x) = \varphi + \frac{h}{\varepsilon} \frac{\varphi^2}{2}.$$

Thus  $\varphi_\varepsilon = \tilde{\varphi}_\varepsilon + h\kappa$ . Let

$$\tilde{A} = -h^2 \Delta - |\nabla \tilde{\varphi}_\varepsilon|^2, \quad \tilde{B} = -2i \langle \nabla \tilde{\varphi}_\varepsilon, h \nabla \cdot \rangle - ih \Delta \tilde{\varphi}_\varepsilon.$$

Since  $\Delta \varphi_\varepsilon = \Delta \tilde{\varphi}_\varepsilon + h \Delta \kappa$  and  $\nabla \varphi_\varepsilon = \nabla \tilde{\varphi}_\varepsilon + h \nabla \kappa$ , we have

$$A = \tilde{A} + A_e, \quad A_e = -h^2 |\nabla \kappa|^2 - 2h \langle \nabla \tilde{\varphi}_\varepsilon, \nabla \kappa \rangle, \\ B = \tilde{B} + B_e, \quad B_e = -2ih \langle \nabla \kappa, h \nabla \cdot \rangle - ih^2 \Delta \kappa.$$

Consequently,

$$i[A, B] = i[\tilde{A}, \tilde{B}] + i[\tilde{A}, B_e] + i[A_e, \tilde{B}] + i[A_e, B_e].$$

Recall from [Ferreira 2009a, p. 143] that

$$i[\tilde{A}, \tilde{B}] = \frac{4h^2}{\varepsilon} \left(1 + \frac{h}{\varepsilon} \varphi\right)^2 + h \tilde{B} \beta \tilde{B} + h^2 R,$$



where  $\beta = (h/\varepsilon)(1 + (h/\varepsilon)\varphi)^{-2}$  and  $R$  is a first order semiclassical differential operator whose coefficients are uniformly bounded with respect to  $h$  and  $\varepsilon$  if we assume that  $h/\varepsilon \leq 1/2$ . Consider now

$$i[\tilde{A}, B_e] = i[-h^2 \Delta - |\nabla \tilde{\varphi}_\varepsilon|^2, -2ih \langle \nabla \kappa, h \nabla \cdot \rangle - ih^2 \Delta \kappa].$$

It is clear that this equals  $h^2 Q$ , where  $Q$  is a second order semiclassical differential operator whose coefficients are uniformly bounded in  $h$  and  $\varepsilon$ . The terms  $i[A_e, \tilde{B}]$  and  $i[A_e, B_e]$  are better. We thus have

$$i[A, B] = \frac{4h^2}{\varepsilon} \left(1 + \frac{h}{\varepsilon} \varphi\right)^2 + h \tilde{B} \beta \tilde{B} + h^2 Q$$

for some  $Q$  as described above. It follows that

$$(i[A, B]u, u) = \frac{4h^2}{\varepsilon} \|(1 + h\varphi/\varepsilon)u\|^2 + h(\tilde{B}\beta\tilde{B}u, u) + h^2(Qu, u).$$

We will choose  $h_0$  so small that  $|h\varphi/\varepsilon| \leq 1/2$  in  $M$  for  $h \leq h_0$ . Since  $u|_{\partial M} = 0$ , integration by parts gives

$$|h(\tilde{B}\beta\tilde{B}u, u)| \leq C_1 \frac{h^2}{\varepsilon} \|\tilde{B}u\|^2,$$

Similarly,

$$|h^2(Qu, u)| \leq C_2 h^2 (\|u\|^2 + \|h\nabla u\|^2).$$

Putting this information together, we get

$$(i[A, B]u, u) \geq \frac{h^2}{\varepsilon} \|u\|^2 - C_1 \frac{h^2}{\varepsilon} \|\tilde{B}u\|^2 - C_2 h^2 (\|u\|^2 + \|h\nabla u\|^2).$$

Next we revisit the term  $\|Au\|^2$ . Let  $K$  be a positive constant whose value will be specified later. Since  $u|_{\partial M} = 0$ , integration by parts and Young's inequality give that

$$\begin{aligned} h^2 \|h\nabla u\|^2 &= h^2 (-h^2 \Delta u, u) = h^2 (Au, u) + h^2 (|\nabla \varphi_\varepsilon|^2 u, u) \\ &\leq \frac{1}{2K} \|Au\|^2 + \frac{Kh^4}{2} \|u\|^2 + C_3 h^2 \|u\|^2, \end{aligned}$$

or

$$\|Au\|^2 \geq 2Kh^2 \|h\nabla u\|^2 - K^2 h^4 \|u\|^2 - 2KC_3 h^2 \|u\|^2.$$

Also recall that  $B - \tilde{B} = B_e = -2ih \langle \nabla \kappa, h \nabla \cdot \rangle - ih^2 \Delta \kappa$ . Thus,

$$\|(B - \tilde{B})u\|^2 \leq C_4 h^2 (\|u\|^2 + \|h\nabla u\|^2).$$

Hence

$$\|\tilde{B}u\|^2 \leq 2\|Bu\|^2 + 2C_4 h^2 (\|u\|^2 + \|h\nabla u\|^2)$$

and

$$\|Bu\|^2 \geq \frac{1}{2} \|\tilde{B}u\|^2 - C_4 h^2 (\|u\|^2 + \|h\nabla u\|^2).$$

Putting our estimates together, we obtain

$$\begin{aligned} \|P_{0,\varphi_\varepsilon} u\|^2 &\geq 2Kh^2 \|h\nabla u\|^2 - K^2 h^4 \|u\|^2 - 2KC_3 h^2 \|u\|^2 + \frac{1}{2} \|\tilde{B}u\|^2 - C_4 h^2 (\|u\|^2 + \|h\nabla u\|^2) \\ &\quad + \frac{h^2}{\varepsilon} \|u\|^2 - C_1 \frac{h^2}{\varepsilon} \|\tilde{B}u\|^2 - C_2 h^2 (\|u\|^2 + \|h\nabla u\|^2) - 2h^3 ((\partial_\nu \varphi_\varepsilon) \partial_\nu u, \partial_\nu u)_{\partial M}. \end{aligned}$$

At this point, we choose  $h_0$  so small that

$$C_1 h_0^2 / \varepsilon \leq \frac{1}{4}.$$

We also make the choice

$$K = \frac{1}{\alpha\varepsilon},$$

where  $\alpha$  is to be determined. Then, for  $h \leq h_0$ ,

$$\begin{aligned} \|P_{0,\varphi_\varepsilon} u\|^2 &\geq \frac{h^2}{\varepsilon} \left( \|u\|^2 + \frac{2}{\alpha} \|h\nabla u\|^2 \right) - (C_2 + C_4) h^2 (\|u\|^2 + \|h\nabla u\|^2) \\ &\quad - \frac{h^2}{\varepsilon} \frac{h^2}{\alpha^2 \varepsilon} \|u\|^2 - \frac{h^2}{\varepsilon} \frac{2C_3}{\alpha} \|u\|^2 + \frac{1}{4} \|\tilde{B}u\|^2 - 2h^3 ((\partial_\nu \varphi_\varepsilon) \partial_\nu u, \partial_\nu u)_{\partial M}. \end{aligned}$$

Choose first  $\alpha = 4C_3$ . It follows that

$$\|P_{0,\varphi_\varepsilon} u\|^2 \geq \frac{h^2}{2\varepsilon} \left( 1 - 2\varepsilon(C_2 + C_4) - \frac{2h^2}{\alpha^2 \varepsilon} \right) \|u\|^2 + \frac{h^2}{\varepsilon} \left( \frac{2}{\alpha} - \varepsilon(C_2 + C_4) \right) \|h\nabla u\|^2 - 2h^3 ((\partial_\nu \varphi_\varepsilon) \partial_\nu u, \partial_\nu u)_{\partial M}.$$

Next choose  $\varepsilon$  so that

$$\varepsilon = \min \left\{ \frac{1}{4(C_2 + C_4)}, \frac{1}{\alpha(C_2 + C_4)} \right\}.$$

Finally, choose  $h_0$  so it satisfies the restrictions made earlier, i.e.,  $h_0 \leq \frac{\varepsilon}{2}$ ,  $h_0 \max_{x \in M} |\varphi| \leq \frac{\varepsilon}{2}$ , and  $h_0^2 \leq \frac{\varepsilon}{4C_1}$ , and additionally

$$\frac{2h_0^2}{\alpha^2 \varepsilon} \leq \frac{1}{4}.$$

With these choices, we have

$$\|P_{0,\varphi_\varepsilon} u\|^2 \geq \frac{h^2}{8\varepsilon} \|u\|^2 + \frac{h^2}{\alpha\varepsilon} \|h\nabla u\|^2 - 2h^3 ((\partial_\nu \varphi_\varepsilon) \partial_\nu u, \partial_\nu u)_{\partial M}.$$

Adding a potential gives

$$\|P_{0,\varphi_\varepsilon} u\|^2 \leq 2\|(P_{0,\varphi_\varepsilon} + h^2 q)u\|^2 + 2h^4 \|q\|_{L^\infty(M)}^2 \|u\|^2.$$

Choosing an even smaller value of  $h_0$  depending on  $\|q\|_{L^\infty(M)}$  if necessary, we obtain for  $0 < h \leq h_0$  that

$$\|(P_{0,\varphi_\varepsilon} + h^2 q)u\|^2 \geq \frac{h^2}{C_0} (\|u\|^2 + \|h\nabla u\|^2) - 2h^3 ((\partial_\nu \varphi_\varepsilon) \partial_\nu u, \partial_\nu u)_{\partial M}.$$

Finally, we replace  $u$  by  $e^{\varphi^2/2\varepsilon+\kappa} u$ , where  $u \in C^\infty(\bar{\Omega})$  and  $u|_{\partial\Omega} = 0$ , and use the fact that

$$1/C \leq e^{\varphi^2/2\varepsilon+\kappa} \leq C, \quad |\nabla(e^{\varphi^2/2\varepsilon+\kappa})| \leq C \quad \text{on } M.$$

The required estimate follows. □

We now pass from  $\varphi_\varepsilon$  to  $\varphi$  in the boundary terms of the previous result, making use of the special properties of  $\varphi_\varepsilon$  on  $\partial M$ . Note that the factor  $h^4$  in the boundary term on  $\{x \in \partial M : -\delta < \partial_\nu \varphi(x) < h/3\}$  below is weaker than the factor  $h^3$  in the other boundary terms. This follows from the fact that  $\partial_\nu \varphi_\varepsilon = h\partial_\nu \kappa = -h$  in the set where  $\partial_\nu \varphi$  vanishes, so one only has the weak lower bound.

**Proposition 4.2.** *Let  $(M, g)$  be as above, let  $q \in L^\infty(M)$ , and let  $\varphi(x) = \pm x_1$ . There exist constants  $C_0, h_0 > 0$  such that, whenever  $0 < h \leq h_0$  and  $\delta > 0$ , one has*

$$\frac{\delta h^3}{C_0} \|\partial_\nu u\|_{L^2(\{\partial_\nu \varphi \leq -\delta\})}^2 + \frac{h^4}{C_0} \|\partial_\nu u\|_{L^2(\{-\delta < \partial_\nu \varphi < h/3\})}^2 + \frac{h^2}{C_0} (\|u\|^2 + \|h\nabla u\|^2) \\ \leq \|e^{\varphi/h}(-h^2\Delta_g + h^2q)(e^{-\varphi/h}u)\|^2 + h^3 \|\partial_\nu u\|_{L^2(\{\partial_\nu \varphi \geq h/3\})}^2$$

for any  $u \in C^\infty(M)$  with  $u|_{\partial M} = 0$ .

*Proof.* Note that

$$\partial_\nu \varphi_\varepsilon = \left(1 + \frac{h}{\varepsilon}\varphi\right)\partial_\nu \varphi + h\partial_\nu \kappa = \left(1 + \frac{h}{\varepsilon}\varphi\right)\partial_\nu \varphi - h.$$

We choose  $h_0$  so small that whenever  $h \leq h_0$ , one has, for  $x \in M$ ,

$$\frac{1}{2} \leq 1 + \frac{h}{\varepsilon}\varphi(x) \leq \frac{3}{2}.$$

On the set where  $\partial_\nu \varphi(x) \leq -\delta$ , we have

$$|\partial_\nu \varphi_\varepsilon| \geq \delta/2.$$

If  $-\delta < \partial_\nu \varphi < h/3$ , we use the estimate

$$|\partial_\nu \varphi_\varepsilon| \geq h/2.$$

Moreover,  $|\partial_\nu \varphi_\varepsilon| \leq C_0$  on  $\partial M$ . Since  $\{\partial_\nu \varphi < h/3\} \subset \{\partial_\nu \varphi_\varepsilon < 0\}$  and  $\{\partial_\nu \varphi_\varepsilon \geq 0\} \subset \{\partial_\nu \varphi \geq h/3\}$ , the result follows from [Proposition 4.1](#) after replacing  $C_0$  by some larger constant.  $\square$

We can now obtain a solvability result from the previous Carleman estimate in a standard way by duality; see [\[Bukhgeim and Uhlmann 2002; Kenig et al. 2007; Nachman and Street 2010\]](#). There is a slight technical complication, since the solution will be in  $L^2$  but not in  $H^1$ . To remedy this, we will work with the space

$$H_{\Delta_g}(M) = \{u \in L^2(M) : \Delta_g u \in L^2(M)\}$$

with norm  $\|u\|_{H_\Delta} = \|u\|_{L^2} + \|\Delta u\|_{L^2}$ . As in [\[Bukhgeim and Uhlmann 2002\]](#), we see that  $H_\Delta(M)$  is a Hilbert space having  $C^\infty(M)$  as a dense subset, and there is a well defined bounded trace operator from  $H_\Delta(M)$  to  $H^{-1/2}(\partial M)$  and a normal derivative operator from  $H_\Delta(M)$  to  $H^{-3/2}(\partial M)$ . We also recall that if  $u \in H_\Delta(M)$  and  $u|_{\partial M} \in H^{3/2}(\partial M)$ , then  $u \in H^2(M)$ .

**Proposition 4.3.** *Let  $(M, g)$  be as above, let  $q \in L^\infty(M)$ , and let  $\varphi(x) = \pm x_1$ . There exist constants  $C_0, \tau_0 > 0$  such that when  $\tau \geq \tau_0$  and  $\delta > 0$ , for any  $f \in L^2(M)$  and  $f_- \in L^2(S_- \cup S_0)$  there exists  $u \in L^2(M)$  satisfying  $e^{\tau\varphi}u \in H_{\Delta_g}(M)$  and  $e^{\tau\varphi}u|_{\partial M} \in L^2(\partial M)$  such that*

$$e^{-\tau\varphi}(-\Delta_g + q)(e^{\tau\varphi}u) = f \text{ in } M, \quad e^{\tau\varphi}u|_{S_- \cup S_0} = e^{\tau\varphi}f_-,$$

and

$$\|u\|_{L^2(M)} \leq C_0(\tau^{-1}\|f\|_{L^2(M)} + (\delta\tau)^{-1/2}\|f_-|_{S_-}\|_{L^2(S_-)} + \|f_-|_{S_0}\|_{L^2(S_0)}).$$

Here  $S_\pm$  and  $S_0$  are the subsets of  $\partial M$  defined by

$$S_- = \{\partial_\nu\varphi \leq -\delta\}, \quad S_0 = \{-\delta < \partial_\nu\varphi < 1/(3\tau)\}, \quad S_+ = \{\partial_\nu\varphi \geq 1/(3\tau)\}.$$

*Proof.* Write  $Lv = e^{\tau\varphi}(-\Delta_g + \bar{q})(e^{-\tau\varphi}v)$  and  $\tau = 1/h$ ,  $\tau_0 = 1/h_0$ . We rewrite the Carleman estimate of Proposition 4.2 as

$$(\delta\tau)^{1/2}\|\partial_\nu v\|_{L^2(S_-)} + \|\partial_\nu v\|_{L^2(S_0)} + \tau\|v\| + \|\nabla v\| \leq C_0\|Lv\| + C_0\tau^{1/2}\|\partial_\nu v\|_{L^2(S_+)}.$$

This is valid for any  $\delta > 0$ , provided that  $\tau \geq \tau_0$  and  $v \in C^\infty(M)$  with  $v|_{\partial M} = 0$ .

Consider the following subspace of  $L^2(M) \times L^2(S_+)$ :

$$X = \{(Lv, \partial_\nu v|_{S_+}) : v \in C^\infty(M), v|_{\partial M} = 0\}.$$

Any element of  $X$  is uniquely represented as  $(Lv, \partial_\nu v|_{S_+})$ , where  $v|_{\partial M} = 0$  by the Carleman estimate. Define a linear functional  $l : X \rightarrow \mathbb{C}$  by

$$l(Lv, \partial_\nu v|_{S_+}) = (v, f)_{L^2(M)} - (\partial_\nu v, f_-)_{L^2(S_- \cup S_0)}.$$

By the Carleman estimate, we have

$$\begin{aligned} |l(Lv, \partial_\nu v|_{S_+})| &\leq \|v\|\|f\| + \|\partial_\nu v\|_{L^2(S_-)}\|f_-|_{L^2(S_-)} + \|\partial_\nu v\|_{L^2(S_0)}\|f_-|_{L^2(S_0)} \\ &\leq C_0(\tau^{-1}\|f\| + (\delta\tau)^{-1/2}\|f_-|_{L^2(S_-)} + \|f_-|_{L^2(S_0)}) \times (\|Lv\| + \tau^{1/2}\|\partial_\nu v\|_{L^2(S_+)}). \end{aligned}$$

The Hahn–Banach theorem implies that  $l$  extends to a continuous linear functional

$$\bar{l} : L^2(M) \times \tau^{-1/2}L^2(S_+) \rightarrow \mathbb{C}$$

such that

$$\|\bar{l}\| \leq C_0(\tau^{-1}\|f\| + (\delta\tau)^{-1/2}\|f_-|_{L^2(S_-)} + \|f_-|_{L^2(S_0)}).$$

By the Riesz representation theorem, there exist functions  $u \in L^2(M)$  and  $u_+ \in L^2(S_+)$  satisfying  $\bar{l}(w, w_+) = (w, u)_{L^2(M)} + (w_+, u_+)_{L^2(S_+)}$ . Moreover,

$$\|u\|_{L^2(M)} + \tau^{-1/2}\|u_+\|_{L^2(S_+)} \leq C_0(\tau^{-1}\|f\| + (\delta\tau)^{-1/2}\|f_-|_{L^2(S_-)} + \|f_-|_{L^2(S_0)}).$$

If  $v \in C^\infty(M)$  and  $v|_{\partial M} = 0$ , we have

$$(Lv, u)_{L^2(M)} + (\partial_\nu v, u_+)_{L^2(S_+)} = (v, f)_{L^2(M)} - (\partial_\nu v, f_-)_{L^2(S_- \cup S_0)}.$$

Choosing  $v$  compactly supported in  $M^{\text{int}}$ , it follows that  $L^*u = f$ , or

$$e^{-\tau\varphi}(-\Delta_g + q)(e^{\tau\varphi}u) = f \quad \text{in } M.$$

Furthermore,  $e^{\tau\varphi}u \in H_\Delta(M)$ .

If  $w, v \in C^\infty(M)$  with  $v|_{\partial M} = 0$ , an integration by parts gives

$$(Lv, w) = -(e^{-\tau\varphi}\partial_\nu v, e^{\tau\varphi}w)_{L^2(\partial M)} + (v, L^*w).$$

Given our solution  $u$ , we choose  $u_j \in C^\infty(M)$  so that  $e^{\tau\varphi}u_j \rightarrow e^{\tau\varphi}u$  in  $H_\Delta(M)$ . Applying the above formula with  $w = u_j$  and taking the limit, we see that

$$(Lv, u) = -(e^{-\tau\varphi}\partial_\nu v, e^{\tau\varphi}u)_{L^2(\partial M)} + (v, L^*u)$$

for  $v \in C^\infty(M)$  with  $v|_{\partial M} = 0$ . Combining this with (4-1), using that  $L^*u = f$  gives

$$(\partial_\nu v, f_-)_{L^2(S_- \cup S_0)} + (\partial_\nu v, u_+)_{L^2(S_+)} = (e^{-\tau\varphi}\partial_\nu v, e^{\tau\varphi}u)_{L^2(\partial M)}.$$

Since  $\partial_\nu v$  can be chosen arbitrarily, it follows that  $e^{\tau\varphi}u|_{S_- \cup S_0} = e^{\tau\varphi}f_-$  and  $e^{\tau\varphi}u|_{S_+} = e^{\tau\varphi}u_+$ . We also see that  $e^{\tau\varphi}u|_{\partial M} \in L^2(\partial M)$ . □

### 5. Reflection approach

In the previous section, we employed Carleman estimates and duality to obtain a solvability result (Proposition 4.3) that will be used to produce correction terms in complex geometrical optics solutions with prescribed behavior on parts of the boundary. In this section we give an alternative approach to the construction of correction terms vanishing on parts of the boundary. The method is based on a reflection argument. We extend the method of [Isakov 2007], which dealt with inaccessible parts that are part of a hyperplane, to the case of inaccessible parts that are part of the graph of a function independent of one of the variables. The results are less general than the ones in Section 4, and, for simplicity, will only be stated for domains in  $\mathbb{R}^3$  with Euclidean metric, but on the other hand, the method is constructive and is based on direct Fourier arguments in the spirit of [Kenig et al. 2011a; Kenig et al. 2011b].

Let  $\Omega \subset \mathbb{R}^3$  be a bounded domain with smooth boundary, and assume that

$$\Omega \subset \mathbb{R} \times \{(x_2, x_3) : x_3 > \eta(x_2)\},$$

where  $\eta : \mathbb{R} \rightarrow \mathbb{R}$  is a smooth function. Also assume that  $\Gamma_0$  is a closed subset of  $\partial\Omega$  such that

$$\Gamma_0 \subset \mathbb{R} \times \{(x_2, x_3) : x_3 = \eta(x_2)\}.$$

We will show that if one has access to suitable amplitudes of complex geometrical optics solutions that vanish on  $\Gamma_0$ , it is possible to produce correction terms that also vanish on  $\Gamma_0$ .

**Proposition 5.1.** *Let  $\Omega$  and  $\Gamma_0$  be as above, and let  $q \in L^\infty(\Omega)$ . There are  $C_0, \tau_0 > 0$  such that, for any  $\tau$  with  $|\tau| \geq \tau_0$  and for any  $m \in H^2(\Omega)$  with  $m|_{\Gamma_0} = 0$ , the equation  $(-\Delta + q)u = 0$  in  $\Omega$  has a solution  $u \in H^2(\Omega)$  of the form*

$$u = e^{-\tau x_1}(m + r)$$

such that  $r|_{\Gamma_0} = 0$  and

$$\|r\|_{L^2(\Omega)} \leq \frac{C_0}{|\tau|} \|e^{\tau x_1}(-\Delta + q)(e^{-\tau x_1}m)\|_{L^2(\Omega)}.$$

The proof involves a reflection argument that reduces the construction of the correction term to the problem of solving a conjugated equation with anisotropic metric,

$$e^{\tau x_1}(-\Delta_{\hat{g}} + \hat{q})(e^{-\tau x_1} \hat{r}) = \hat{f} \quad \text{in } \mathbb{R} \times \widehat{\Omega}_0,$$

where  $\widehat{\Omega}_0 \subset \mathbb{R}^2$  is a bounded open set and  $\hat{g}$  is a metric of the form

$$\hat{g}(y_1, y') = \begin{pmatrix} 1 & 0 \\ 0 & \hat{g}_0(y') \end{pmatrix},$$

and where  $g_0$  is smooth for  $y_3 \neq 0$  but only Lipschitz continuous across  $\{y_3 = 0\}$ . In three and higher dimensions, it is not known how to handle equations of this type with general Lipschitz coefficients in the second order part (the case of  $C^1$  coefficients, and also Lipschitz coefficients with a smallness condition, is considered in [Haberman and Tataru 2013]). However, in our case, the singularity of  $\hat{g}$  only appears in the lower right block  $\hat{g}_0$ , and this turns out not to be a problem.

The following is an analogue of [Kenig et al. 2011a, Proposition 4.1], the main difference being that the transversal metric is only Lipschitz. (With correct definitions, one could easily deal with  $L^\infty$  transversal metrics as well, but then the solution would only be in  $H^1_{-\delta}(T)$ .) Here we write  $(x_1, x')$  for coordinates in  $T = \mathbb{R} \times M_0$ , and for  $\delta \in \mathbb{R}$  we consider the spaces

$$\|f\|_{L^2_\delta(T)} = \|\langle x_1 \rangle^\delta f\|_{L^2(T)}, \quad \|f\|_{H^1_\delta(T)} = \|f\|_{L^2_\delta(T)} + \|df\|_{L^2_\delta(T)}$$

with  $\langle t \rangle = (1 + t^2)^{1/2}$ , and similarly for  $H^2_\delta(T)$ . We also write  $\text{Spec}(-\Delta_{g_0})$  for the set of Dirichlet eigenvalues of the Laplace–Beltrami operator  $-\Delta_{g_0}$  in  $(M_0, g_0)$ .

**Proposition 5.2.** *Let  $T = \mathbb{R} \times M_0$  with metric  $g = e \oplus g_0$ , where  $(M_0, g_0)$  is a compact oriented manifold with smooth boundary and  $g_0$  is a Lipschitz continuous Riemannian metric on  $M_0$ . Given any  $q \in L^\infty_{\text{comp}}(T)$  and any  $\delta > 1/2$ , there are constants  $C_0, \tau_0 > 0$  such that whenever*

$$|\tau| \geq \tau_0 \quad \text{and} \quad \tau^2 \notin \text{Spec}(-\Delta_{g_0}),$$

the equation

$$e^{\tau x_1}(-\Delta_g + q)(e^{-\tau x_1} r) = f \quad \text{in } T$$

has a unique solution  $r \in H^1_{-\delta}(T)$  with  $r|_{\partial T} = 0$  for any  $f \in L^2_\delta(T)$ . Moreover,  $r \in H^2_{-\delta}(T)$ , and one has the bounds

$$\|r\|_{L^2_{-\delta}(T)} \leq \frac{C_0}{|\tau|} \|f\|_{L^2_\delta(T)}, \quad \|r\|_{H^1_{-\delta}(T)} \leq C_0 \|f\|_{L^2_\delta(T)}.$$

*Proof.* The proof is almost exactly the same as the proof of [Kenig et al. 2011a, Proposition 4.1], and we only give the main idea. Since  $\Delta_g = \partial_{x_1}^2 + \Delta_{g_0}$ , the equation that we need to solve is

$$(-\partial_{x_1}^2 + 2\tau \partial_{x_1} - \Delta_{g_0} - \tau^2 + q)r = f \quad \text{in } T.$$

It is enough to consider  $q = 0$ . The standard argument based on weak solutions shows that even when  $g_0$  has very little regularity, there is an orthonormal basis of  $L^2(M_0)$  consisting of Dirichlet eigenfunctions

of  $-\Delta_{g_0}$ ,

$$-\Delta_{g_0}\varphi_l = \lambda_l\varphi_l \text{ in } M_0, \quad \varphi_l \in H_0^1(M_0),$$

where  $0 < \lambda_1 \leq \lambda_2 \leq \lambda_3 \leq \dots \rightarrow \infty$  are the Dirichlet eigenvalues of  $-\Delta_{g_0}$  in  $M_0$ .

Considering the partial Fourier expansions

$$r(x_1, x') = \sum_{l=1}^{\infty} \tilde{r}(x_1, l)\varphi_l(x'), \quad f(x_1, x') = \sum_{l=1}^{\infty} \tilde{f}(x_1, l)\varphi_l(x'),$$

it is enough to solve

$$(-\partial_{x_1}^2 + 2\tau\partial_{x_1} + \lambda_l - \tau^2)\tilde{r}(\cdot, l) = \tilde{f}(\cdot, l) \quad \text{in } \mathbb{R} \text{ for all } l.$$

The condition  $\tau^2 \notin \text{Spec}(-\Delta_{g_0})$  allows us to solve these ordinary differential equations by the Fourier transform as in [Kenig et al. 2011a, Section 4], and the estimates given there imply that one obtains a unique solution  $r \in H_{-\delta}^1(T)$  with  $r|_{\partial T} = 0$  satisfying the required bounds. Elliptic  $H^2$  regularity also works with Lipschitz  $g_0$ , and the argument in [Kenig et al. 2011a, Section 4] gives that  $r \in H_{-\delta}^2(T)$ .  $\square$

*Proof of Proposition 5.1.* We begin by flattening  $\Gamma_0$  via the map

$$\Phi : \mathbb{R}^3 \rightarrow \mathbb{R}^3, \quad (x_1, x_2, x_3) \mapsto (x_1, x_2, x_3 - \eta(x_2)).$$

Let  $\tilde{\Omega} = \Phi(\Omega)$ , write  $y$  for coordinates in  $\tilde{\Omega}$ , and let  $R$  be the reflection

$$R(y_1, y_2, y_3) = (y_1, y_2, -y_3).$$

Note that  $\tilde{\Omega} \subset \{y_3 > 0\}$ . Consider the reflected domain  $\tilde{\Omega}^* = R(\tilde{\Omega})$ , so  $\tilde{\Omega}^* \subset \{y_3 < 0\}$ , and let  $U$  the double domain  $\tilde{\Omega} \cup \Phi(\Gamma_0)^{\text{int}} \cup \tilde{\Omega}^*$ .

Let  $\Psi = \Phi^{-1}$ , let  $\tilde{g} = \Psi^*e$  be the metric in  $\tilde{\Omega}$  that is the pullback of the Euclidean metric in  $\Omega$ , let  $\tilde{q} = \Psi^*q$ , and let  $\tilde{m} = \Psi^*m$ . In the double domain  $U$ , we use even reflection to define the quantities

$$\hat{g} = \begin{cases} \tilde{g} & \text{if } y_3 > 0, \\ R^*\tilde{g} & \text{if } y_3 < 0, \end{cases} \quad \hat{q} = \begin{cases} \tilde{q} & \text{if } y_3 > 0, \\ R^*\tilde{q} & \text{if } y_3 < 0, \end{cases}$$

and odd reflection to define the amplitude

$$\hat{m} = \begin{cases} \frac{1}{2}\tilde{m} & \text{if } y_3 > 0, \\ -\frac{1}{2}R^*\tilde{m} & \text{if } y_3 < 0. \end{cases}$$

Since the flattening map  $\Phi$  leaves  $x_1$  intact, we have

$$\hat{g}(y_1, y') = \begin{pmatrix} 1 & 0 \\ 0 & \hat{g}_0(y') \end{pmatrix},$$

where  $\hat{g}_0$  is a Lipschitz continuous metric only depending on  $y_2$  and  $y_3$ . (In fact,  $\tilde{g}$  and  $\tilde{g}_0$  are well defined in  $\{y_3 > 0\}$  by the flattening map  $\Phi$  and the Euclidean metric in  $\{x_3 > \eta(x_2)\}$ .) Also,  $\hat{q} \in L^\infty(U)$ , and  $\hat{m} \in H^2(U)$  by the boundary condition  $m|_{\Gamma_0} = 0$  and by the properties of odd reflection.

We wish to find  $\hat{r} \in H^1(U)$  satisfying

$$e^{\tau x_1}(-\Delta_{\hat{g}} + \hat{q})(e^{-\tau x_1} \hat{r}) = \hat{f},$$

where  $\hat{f} = -e^{\tau x_1}(-\Delta_{\hat{g}} + \hat{q})(e^{-\tau x_1} \hat{m})$ . Now

$$\begin{aligned} \|\hat{f}\|_{L^2(U)} &= \|\hat{f}\|_{L^2(\tilde{\Omega})} + \|\hat{f}\|_{L^2(\tilde{\Omega}^*)} \\ &= \|\Psi^*(e^{\tau x_1}(-\Delta + q)(e^{-\tau x_1} m))\|_{L^2(\tilde{\Omega})} + \|R^*\Psi^*(e^{\tau x_1}(-\Delta + q)(e^{-\tau x_1} m))\|_{L^2(\tilde{\Omega}^*)} \\ &\leq C\|e^{\tau x_1}(-\Delta + q)(e^{-\tau x_1} m)\|_{L^2(\Omega)}. \end{aligned}$$

Choose a bounded open set  $\widehat{\Omega}_0 \subset \mathbb{R}^2$  such that

$$U \Subset \mathbb{R} \times \widehat{\Omega}_0,$$

and let  $\hat{g}_0$  be the metric in  $\widehat{\Omega}_0$  that is the even extension of  $\tilde{g}_0$  from  $\{y_3 > 0\}$  to  $\widehat{\Omega}_0$ . Then  $\hat{g}_0$  is smooth for  $y_3 \neq 0$  and Lipschitz continuous across  $\{y_3 = 0\}$ . Extending  $\hat{g}$  to  $\mathbb{R} \times \widehat{\Omega}_0$  using the block structure and extending  $\hat{q}$  and  $\hat{f}$  by zero to  $\mathbb{R} \times \widehat{\Omega}_0$ , it is enough to find a solution  $\hat{r} \in H_{loc}^2(\mathbb{R} \times \widehat{\Omega}_0)$  of the equation

$$e^{\tau x_1}(-\Delta_{\hat{g}} + \hat{q})(e^{-\tau x_1} \hat{r}) = \hat{f} \quad \text{in } \mathbb{R} \times \widehat{\Omega}_0. \quad (5-1)$$

Such a solution may be found by [Proposition 5.2](#), and denoting by  $\hat{r}$  its restriction to  $U$ , we have

$$\|\hat{r}\|_{L^2(U)} \leq \frac{C}{|\tau|} \|\hat{f}\|_{L^2(U)}.$$

Now define

$$\hat{u} = e^{-\tau x_1}(\hat{m} + \hat{r}) \quad \text{in } U$$

and

$$\tilde{u} = \hat{u} - R^*\hat{u} \quad \text{in } \tilde{\Omega}.$$

Then  $(-\Delta_{\hat{g}} + \hat{q})\hat{u} = 0$  in  $U$ , and  $(-\Delta_{\tilde{g}} + \tilde{q})\tilde{u} = 0$  in  $\tilde{\Omega}$  by the definition of  $\hat{g}$  and  $\hat{q}$  and using that  $\hat{u} \in H^2(U)$ . We also have

$$\tilde{u} = e^{-\tau x_1}(\hat{m} - R^*\hat{m} + \hat{r} - R^*\hat{r}) \quad \text{in } \tilde{\Omega}.$$

But here  $\hat{m} - R^*\hat{m}|_{\tilde{\Omega}} = \tilde{m}$  by the definition of  $\hat{m}$ . Consequently, if we define  $u = \Phi^*\tilde{u}$ , then  $(-\Delta + q)u = 0$  in  $\Omega$  and

$$u = e^{-\tau x_1}(m + r) \quad \text{in } \Omega,$$

where  $r = \Phi^*(\hat{r} - R^*\hat{r})$  satisfies

$$\|r\|_{L^2(\Omega)} \leq C\|\hat{r}\|_{L^2(U)} \leq \frac{C}{|\tau|} \|\hat{f}\|_{L^2(U)} \leq \frac{C}{|\tau|} \|e^{\tau x_1}(-\Delta + q)(e^{-\tau x_1} m)\|_{L^2(\Omega)}. \quad \square$$

Note how the odd reflection of the amplitude  $m$  in the proof ensured that the solution obtained by reflection is not the zero solution. We also remark that under certain conditions, the arguments in [Sections 6 and 7](#) allow to construct amplitudes  $m$  vanishing on a part  $\Gamma_0$  as above.



### 6. Local uniqueness on simple manifolds

In this section we prove Theorems 2.1–2.3. In these results the transversal manifold is assumed to be simple and we only use nonreflected geodesics. This case already illustrates the main features of the approach, and we can use a quasimode construction that is much easier than the Gaussian beam one used for nonsimple transversal manifolds and reflected geodesics.

The first observation is the usual integral identity.

**Proposition 6.1.** *If  $\Gamma_D, \Gamma_N \subset \partial M$  are open and if  $C_{g,q_1}^{\Gamma_D, \Gamma_N} = C_{g,q_2}^{\Gamma_D, \Gamma_N}$ , then*

$$\int_M (q_1 - q_2)u_1u_2 dV_g = 0$$

for any  $u_j \in H_{\Delta_g}(M)$  satisfying  $(-\Delta_g + q_j)u_j = 0$  in  $M$  and

$$\text{supp}(u_1|_{\partial M}) \subset \Gamma_D, \quad \text{supp}(u_2|_{\partial M}) \subset \Gamma_N.$$

*Proof.* Let  $u_j$  be as stated. Since  $C_{g,q_1}^{\Gamma_D, \Gamma_N} = C_{g,q_2}^{\Gamma_D, \Gamma_N}$ , there is a function  $\tilde{u}_2 \in H_{\Delta}(M)$  with  $(-\Delta + q_2)\tilde{u}_2 = 0$  in  $M$ ,  $\text{supp}(\tilde{u}_2|_{\partial M}) \subset \Gamma_D$ , and

$$(u_1|_{\Gamma_D}, \partial_\nu u_1|_{\Gamma_N}) = (\tilde{u}_2|_{\Gamma_D}, \partial_\nu \tilde{u}_2|_{\Gamma_N}).$$

Using that  $u_1, u_2$ , and  $\tilde{u}_2$  are solutions, we have

$$\begin{aligned} \int_M (q_1 - q_2)u_1u_2 dV &= \int_M [(\Delta u_1)u_2 - u_1(\Delta u_2)] dV \\ &= \int_M [(\Delta(u_1 - \tilde{u}_2))u_2 - (u_1 - \tilde{u}_2)(\Delta u_2)] dV. \end{aligned}$$

Now  $u_1 - \tilde{u}_2|_{\partial M} = 0$ , so in fact  $u_1 - \tilde{u}_2 \in H^2(M)$  by the properties of the space  $H_{\Delta}(M)$ . Recall also that  $C^\infty(M)$  is dense in  $H_{\Delta}(M)$  and that  $u_2|_{\partial M} \in H^{-1/2}(\partial M)$  and  $\partial_\nu u_2|_{\partial M} \in H^{-3/2}(\partial M)$ . These facts make it possible to integrate by parts, and we obtain that

$$\int_M (q_1 - q_2)u_1u_2 dV = \int_{\partial M} [(\partial_\nu(u_1 - \tilde{u}_2))u_2 - (u_1 - \tilde{u}_2)(\partial_\nu u_2)] dS$$

in the weak sense. The last expression vanishes since  $\partial_\nu(u_1 - \tilde{u}_2)|_{\Gamma_N} = 0$  and  $\text{supp}(u_2|_{\partial M}) \subset \Gamma_N$ .  $\square$

The next result will be used to pass from the metric  $g = c(e \oplus g_0)$  to the slightly simpler metric  $\tilde{g} = e \oplus g_0$ .

**Lemma 6.2.** *Let  $c$  be a smooth positive function in  $M$ . Then  $u \in H_{\Delta_g}(M)$  satisfies  $(-\Delta_g + q)u = 0$  in  $M$  if and only if  $\tilde{u} \in H_{\Delta_{\tilde{g}}}(M)$  satisfies  $(-\Delta_{\tilde{g}} + \tilde{q})\tilde{u} = 0$  in  $M$ , where*

$$\tilde{g} = c^{-1}g, \quad \tilde{u} = c^{(n-2)/4}u, \quad \tilde{q} = c(q - c^{(n-2)/4}\Delta_g(c^{-(n-2)/4})).$$

*Proof.* This follows from the identity for  $v \in C^\infty(M)$ ,

$$c^{(n+2)/4}(-\Delta_g + q)(c^{-(n-2)/4}v) = (-\Delta_{c^{-1}g} + c(q - c^{(n-2)/4}\Delta_g(c^{-(n-2)/4})))v,$$

upon approximating  $u$  or  $\tilde{u}$  by smooth functions. □

*Proof of Theorem 2.1.* Let  $\tilde{g} = e \oplus g_0$  and  $\tilde{q}_j = c(q_j - c^{(n-2)/4}\Delta_g(c^{-(n-2)/4}))$ . Let  $\lambda$  be a fixed real number, and consider the complex frequency

$$s = \tau + i\lambda,$$

where  $\tau > 0$  will be large. We look for solutions

$$\begin{aligned} \tilde{u}_1 &= e^{-sx_1}(v_s(x') + r_1), \\ \tilde{u}_2 &= e^{sx_1}(v_s(x') + r_2) \end{aligned}$$

of the equations  $(-\Delta_{\tilde{g}} + \tilde{q}_1)\tilde{u}_1 = 0$ ,  $(-\Delta_{\tilde{g}} + \tilde{q}_2)\tilde{u}_2 = 0$  in  $M$ . Here  $v_s$  will be a quasimode for the Laplacian in  $(M_0, g_0)$  that concentrates near the given geodesic  $\gamma$ . Next we will construct a suitable solution  $\tilde{u}_1$ , and the case of  $\tilde{u}_2$  will be analogous.

Since  $\Delta_{\tilde{g}} = \partial_1^2 + \Delta_{g_0}$ , the function  $\tilde{u}_1$  is a solution if and only if

$$e^{sx_1}(-\Delta_{\tilde{g}} + \tilde{q}_1)(e^{-sx_1}r_1) = -(-\Delta_{g_0} + \tilde{q}_1 - s^2)v_s(x') \quad \text{in } M. \tag{6-1}$$

We want to choose  $v_s \in C^\infty(M_0)$  to satisfy

$$\|v_s\|_{L^2(M_0)} = O(1), \quad \|(-\Delta_{g_0} - s^2)v_s\|_{L^2(M_0)} = O(1) \tag{6-2}$$

as  $\tau \rightarrow \infty$ . Looking for  $v_s$  in the form

$$v_s = e^{is\psi}a,$$

where  $\psi, a \in C^\infty(M_0)$ , a direct computation shows that

$$(-\Delta_{g_0} - s^2)v_s = e^{is\psi}(s^2[|d\psi|_{g_0}^2 - 1]a - is[2\langle d\psi, d\cdot \rangle_{g_0} + \Delta_{g_0}\psi]a - \Delta_{g_0}a).$$

Since  $(M_0, g_0)$  is simple, it is easy to find  $\psi$  and  $a$  so that the expressions in brackets will vanish and that the resulting quasimode  $v_s$  will concentrate near the geodesic  $\gamma$ . To do this, let  $(\widehat{M}_0, g_0)$  be a simple manifold that is slightly larger than  $(M_0, g_0)$ , extend  $\gamma$  as a geodesic in  $\widehat{M}_0$ , and choose  $\varepsilon > 0$  such that  $\gamma|_{(-2\varepsilon, 0) \cup (L, L+2\varepsilon)}$  stays in  $\widehat{M}_0 \setminus M_0$  (this is possible since  $\gamma$  is nontangential). Let  $\omega = \gamma(-\varepsilon) \in \widehat{M}_0 \setminus M_0$ , and let  $(r, \theta)$  be polar normal coordinates in  $(M_0, g_0)$  with center  $\omega$ . Then  $\gamma$  corresponds to the curve  $r \mapsto (r, \theta_0)$  for some fixed  $\theta_0 \in S^{n-2}$ . We will choose

$$\begin{aligned} \psi(r, \theta) &= r, \\ a(r, \theta) &= |g_0(r, \theta)|^{-1/4}b(\theta), \end{aligned}$$

where  $|g_0|$  is the determinant of  $g_0$ , and  $b$  is a fixed function in  $C^\infty(S^{n-2})$  that is supported so close to  $\theta_0$  such that  $v_s|_{\partial M_0 \setminus E} = 0$ . With these choices, we have, as in [Ferreira 2009a],

$$(-\Delta_{g_0} - s^2)v_s = -e^{is\psi}\Delta_{g_0}a.$$

Thus  $v_s$  satisfies the estimates (6-2), and also the estimate

$$\|v_s\|_{L^\infty(M_0)} = O(1).$$

We now go back to (6-1) and look for a solution in the form  $r_1 = e^{i\lambda x_1} r'_1$  where  $r'_1$  satisfies

$$e^{\tau x_1}(-\Delta_{\tilde{g}} + \tilde{q}_1)(e^{-\tau x_1} r'_1) = f \quad \text{in } M \tag{6-3}$$

with

$$f = -e^{-i\lambda x_1}(-\Delta_{g_0} + \tilde{q}_1 - s^2)v_s(x').$$

We also want to arrange that  $\text{supp}(\tilde{u}_1|_{\partial M}) \subset \Gamma_D$ , where  $\Gamma_D \supset \partial M_- \cup \Gamma_a$ . For this purpose, let  $\delta > 0$  be a small number to be fixed later, let  $S_\pm$  and  $S_0$  be the sets in Proposition 4.3 with Carleman weight  $\varphi(x) = -x_1$ , define

$$\begin{aligned} V^\delta &= \{x \in S_- \cup S_0 : \text{dist}_{\partial M}(x, \Gamma_i) < \delta \text{ or } x \in \partial M_+\}, \\ \Gamma_a^\delta &= (S_- \cup S_0) \setminus V_\delta, \end{aligned}$$

and impose the boundary condition

$$e^{\tau\varphi} r'_1|_{S_- \cup S_0} = e^{\tau\varphi} f_-, \tag{6-4}$$

where

$$f_- = \begin{cases} -e^{-i\lambda x_1} v_s(x') & \text{on } V^\delta, \\ 0 & \text{on } \Gamma_a^\delta. \end{cases}$$

Note that  $\partial M_+ \cup \partial M_{\text{tan}}$  (these sets refer to the weight  $x_1$ ) is in the interior of  $S_- \cup S_0$  in  $\partial M$ .

We have seen that  $\|f\|_{L^2(M)} = O(1)$  as  $\tau \rightarrow \infty$ . We also have

$$f_-|_{\partial M_{\text{tan}}} = 0,$$

since  $f_-|_{\Gamma_a^\delta \cap \partial M_{\text{tan}}} = 0$  by definition and  $f_-|_{\partial M_{\text{tan}} \cap V_\delta} = 0$  for sufficiently small  $\delta > 0$  by the construction of  $v_s$  and using that  $\Gamma_i \subset \mathbb{R} \times (\partial M_0 \setminus E)$ . Since  $\|f_-\|_{L^\infty(S_- \cup S_0)} \lesssim 1$ , we have

$$\|f_-\|_{L^2(S_-)} \lesssim \sigma(\{\partial_\nu x_1 \geq \delta\})$$

and

$$\|f_-\|_{L^2(S_0)} \lesssim \sigma(\{-1/(3\tau) < \partial_\nu x_1 < 0\} \cup \{0 < \partial_\nu x_1 < \delta\}),$$

where  $\sigma$  is the surface measure on  $\partial M$ . It follows from Proposition 4.3 that (6-3) has a solution  $r'_1$  satisfying the boundary condition (6-4), and having the estimate

$$\|r'_1\|_{L^2(M)} \lesssim \tau^{-1} + (\delta\tau)^{-1/2} \sigma(\{\partial_\nu x_1 \geq \delta\}) + \sigma(\{-1/(3\tau) < \partial_\nu x_1 < 0\}) + \sigma(\{0 < \partial_\nu x_1 < \delta\}).$$

The implied constants in the previous inequality are independent of  $\tau$  and  $\delta$ . By the basic properties of measures, for some constant  $C_0 > 0$ , we have

$$\|r'_1\|_{L^2(M)} \leq C_0[\tau^{-1} + (\delta\tau)^{-1/2} + o_{\tau \rightarrow \infty}(1) + o_{\delta \rightarrow 0}(1)].$$

Given  $\varepsilon > 0$ , we first choose  $\delta$  so that  $C_0 o_{\delta \rightarrow 0}(1) \leq \varepsilon/2$ . After this, we choose  $\tau > 0$  so large that  $C_0(\tau^{-1} + (\delta\tau)^{-1/2} + o_{\tau \rightarrow \infty}(1)) \leq \varepsilon/2$ . This shows that

$$\lim_{\tau \rightarrow \infty} \|r'_1(\cdot; \tau)\|_{L^2(M)} = 0.$$

Choosing  $r'_1$  as described above and choosing  $r_1 = e^{i\lambda x_1} r'_1$ , we have produced a solution  $\tilde{u}_1 \in H_{\Delta_{\tilde{g}}}(M)$  of the equation  $(-\Delta_{\tilde{g}} + \tilde{q}_1)\tilde{u}_1 = 0$  in  $M$ , having the form

$$\tilde{u}_1 = e^{-sx_1}(v_s(x') + r_1)$$

and satisfying

$$\text{supp}(\tilde{u}_1|_{\partial M}) \subset \Gamma_D$$

and  $\|r_1\|_{L^2(M)} = o(1)$  as  $\tau \rightarrow \infty$ . Repeating this construction for the Carleman weight  $\varphi(x) = x_1$ , we obtain a solution  $\tilde{u}_2 \in H_{\Delta_{\tilde{g}}}(M)$  of the equation  $(-\Delta_{\tilde{g}} + \tilde{q}_2)\tilde{u}_2 = 0$  in  $M$ , having the form

$$\tilde{u}_2 = e^{sx_1}(v_s(x') + r_2)$$

and satisfying

$$\text{supp}(\tilde{u}_2|_{\partial M}) \subset \Gamma_N$$

and  $\|r_2\|_{L^2(M)} = o(1)$  as  $\tau \rightarrow \infty$ .

Writing  $u_j = c^{-(n-2)/4}\tilde{u}_j$ , [Lemma 6.2](#) shows that  $u_j \in H_{\Delta_g}(M)$  are solutions of  $(-\Delta_g + q_1)u_1 = 0$  and  $(-\Delta_g + \bar{q}_2)u_2 = 0$  in  $M$ . Then [Proposition 6.1](#) implies that

$$\int_M (q_1 - q_2)u_1\bar{u}_2 dV_g = 0.$$

We extend  $q_1 - q_2$  by zero to  $\mathbb{R} \times M_0$ . Inserting the expressions for  $u_j$ , and using that  $dV_g = c^{n/2} dx_1 dV_{g_0}(x')$ , we obtain

$$\int_{M_0} \int_{-\infty}^{\infty} (q_1 - q_2)ce^{-2i\lambda x_1}(|v_s(x')|^2 + v_s\bar{r}_2 + \bar{v}_s r_1 + r_1\bar{r}_2) dx_1 dV_{g_0}(x') = 0.$$

Since  $\|r_j\|_{L^2(M)} = o(1)$  as  $\tau \rightarrow \infty$  and since  $dV_{g_0} = |g_0|^{1/2} dr d\theta$  in the  $(r, \theta)$  coordinates, it follows that

$$\int_{S^{n-2}} \int_0^{\infty} e^{-2\lambda r} (c(q_1 - q_2))^{\wedge}(2\lambda, r, \theta)|b(\theta)|^2 dr d\theta = 0.$$

Varying  $b$  in  $C^\infty(S^{n-2})$  so that the support of  $b$  is very close to  $\theta_0$ , this implies that

$$\int_0^{\infty} e^{-2\lambda r} (c(q_1 - q_2))^{\wedge}(2\lambda, r, \theta_0) dr = 0.$$

Since  $\gamma$  was the curve  $r \mapsto (r, \theta)$ , this shows the result. □

*Proof of Theorem 2.2.* Suppose that the local ray transform is injective on  $O$  and  $O \cap \partial M \subset E$ . By Theorem 2.1, we know that

$$\int_0^L e^{-2\lambda t} (c(q_1 - q_2))^{\wedge}(2\lambda, \gamma(t)) dt = 0 \tag{6-5}$$

for any nontangential geodesic  $\gamma$  in  $O$ . Setting  $\lambda = 0$  and using local injectivity of the ray transform, we obtain that

$$(c(q_1 - q_2))^{\wedge}(0, \cdot) = 0 \quad \text{in } O.$$

Going back to (6-5) and differentiating this identity with respect to  $\lambda$ , and then setting  $\lambda = 0$  and using the vanishing of  $(c(q_1 - q_2))^{\wedge}(0, \cdot)$  on  $O$ , it follows that

$$\int_0^L \frac{\partial}{\partial \lambda} [(c(q_1 - q_2))^{\wedge}](0, \gamma(t)) dt = 0 \quad \text{in } O$$

for any nontangential geodesic in  $O$ . Local uniqueness for the ray transform again implies that

$$\frac{\partial}{\partial \lambda} [(c(q_1 - q_2))^{\wedge}](0, \cdot) = 0 \quad \text{in } O.$$

Iterating this argument by taking higher order derivatives of (6-5) shows that

$$\left(\frac{\partial}{\partial \lambda}\right)^k [(c(q_1 - q_2))^{\wedge}](0, \cdot) = 0 \quad \text{in } O$$

for any  $k$ . Since  $c(q_1 - q_2)$  is compactly supported in  $x_1$ , its Fourier transform is analytic and we have

$$(c(q_1 - q_2))^{\wedge}(\lambda, \cdot) = 0 \quad \text{in } O \text{ for all } \lambda \in \mathbb{R}.$$

Inverting the Fourier transform and using that  $c$  is positive, we obtain that  $q_1 = q_2$  in  $M \cap (\mathbb{R} \times O)$ .  $\square$

*Proof of Theorem 2.3.* Since  $(M, g)$  is admissible, we may assume that

$$(M, g) \subset (\mathbb{R} \times M_0, g), \quad g = c(e \oplus g_0),$$

where  $(M_0, g_0)$  is simple. The argument is very similar to the proof of Theorem 2.1, and we only indicate the required changes. Up to the formula (6-3), the only change is that there is no restriction on  $b \in C^\infty(S^{n-2})$  (we do not require  $v_s$  to vanish on any part of the boundary). The function  $r'_1$  is obtained as a solution of (6-3), but this time we want  $\text{supp}(\tilde{u}_1|_{\partial M}) \subset \partial M_-$ . Fix  $\delta > 0$ . The boundary condition for  $\tilde{u}_1$  is (6-4), where  $f_-$  is chosen to be

$$f_- = -e^{-i\lambda x_1} v_s(x') \quad \text{on } S_- \cup S_0.$$

We use Proposition 4.3 to solve for  $r'_1$ . We have  $\|f\|_{L^2(M)} = O(1)$ , and the bound  $\|f_-\|_{L^\infty} \lesssim 1$  implies

$$\|f_-\|_{L^2(S_-)} \lesssim \sigma(\{\partial_\nu x_1 \geq \delta\})$$

and

$$\|f_-\|_{L^2(S_0)} \lesssim \sigma(\{-1/(3\tau) < \partial_\nu x_1 < 0\}) + \sigma(\partial M_{\text{tan}}) + \sigma(\{0 < \partial_\nu x_1 < \delta\}).$$

Now we use that

$$\sigma(\partial M_{\tan}) = 0.$$

This shows that we obtain the same estimate for  $r'_1$  as before:

$$\|r'_1\|_{L^2(M)} \leq C_0[\tau^{-1} + (\delta\tau)^{-1/2} + o_{\tau \rightarrow \infty}(1) + o_{\delta \rightarrow 0}(1)].$$

We can now continue as in the proof of [Theorem 2.1](#) to conclude that

$$\int_0^L e^{-2\lambda t} (c(q_1 - q_2))^{\wedge}(2\lambda, \gamma(t)) dt = 0$$

for any  $\lambda \in \mathbb{R}$  and for any nontangential geodesic in  $(M_0, g_0)$ . The geodesic ray transform (with zero attenuation) is injective in  $(M_0, g_0)$  [[Sharafutdinov 1994](#)]. Following the proof of [Theorem 2.2](#), but now using all the nontangential geodesics in  $(M_0, g_0)$ , shows that  $q_1 = q_2$  in  $M$ . □

### 7. Quasimodes concentrating near broken rays

In this section, to simplify notation, we write  $(M, g)$  instead of  $(M_0, g_0)$  and we assume that  $(M, g)$  is a compact oriented Riemannian manifold having smooth boundary and  $\dim(M) = m \geq 2$ . Suppose that  $E$  is a nonempty open subset of  $\partial M$ , and let  $R = \partial M \setminus E$ . We think of  $E$  as the observation set where geodesics can enter and exit, and  $R$  is the reflecting set. In the Calderón problem with partial data, we are led to consider attenuated broken ray transforms, where one integrates a function on  $M$  over broken geodesic rays that enter  $M$  at some point of  $E$ , reflect nontangentially at points of  $R$ , and then exit  $M$  at some point of  $E$ . The reflections will obey the law of geometric optics, so that a geodesic hitting the boundary in direction  $v$  will be continued by the geodesic in the reflected direction  $\hat{v} = v - 2\langle v, \nu \rangle \nu$ .

Given a slightly complex frequency  $s = \tau + i\lambda$ , we will construct corresponding quasimodes, or approximate eigenfunctions, that concentrate near a fixed nontangential broken ray.

**Proposition 7.1.** *Let  $\gamma : [0, L] \rightarrow M$  be a nontangential broken ray with endpoints on  $E$ , and let  $\lambda$  be a fixed real number. For any  $K > 0$ , there is a family  $\{v_s : s = \tau + i\lambda, \tau \geq 1\}$  in  $C^\infty(M)$  such that, as  $\tau \rightarrow \infty$ ,*

$$\|(-\Delta_g - s^2)v_s\|_{L^2(M)} = O(\tau^{-K}), \quad \|v_s\|_{L^2(M)} = O(1),$$

the boundary values of  $v_s$  satisfy

$$\|v_s\|_{L^2(R)} = O(\tau^{-K}), \quad \|v_s\|_{L^2(\partial M)} = O(1),$$

and, for any  $\psi \in C(M)$ ,

$$\int_M |v_{\tau+i\lambda}|^2 \psi dV_g \rightarrow \int_0^L e^{-2\lambda t} \psi(\gamma(t)) dt \quad \text{as } \tau \rightarrow \infty.$$

Let us begin by proving this result in the special case  $E = \partial M$ , so that  $R = \emptyset$  and one does not need to worry about reflected rays. The next three preparatory lemmas describe a modified Fermi coordinate system that is very useful in this construction.

**Lemma 7.2.** *Let  $(\widehat{M}, g)$  be a compact manifold without boundary, and let  $\gamma : (a, b) \rightarrow \widehat{M}$  be a unit speed geodesic segment that has no loops. There are only finitely many times  $t \in (a, b)$  such that  $\gamma$  intersects itself at  $\gamma(t)$ .*

*Proof.* Since  $\gamma$  has no loops,  $(\gamma(t), \dot{\gamma}(t)) = (\gamma(t'), \dot{\gamma}(t'))$  implies  $t = t'$ . The first observation is that  $\gamma$  can only self-intersect transversally, since  $(\gamma(t), \dot{\gamma}(t)) = (\gamma(t'), -\dot{\gamma}(t'))$  also implies  $t = t'$  (if this would happen for  $t < t'$ , then, by uniqueness of geodesics,  $\dot{\gamma}((t + t')/2) = -\dot{\gamma}((t + t')/2)$ , which is impossible). Next note that if  $r$  is smaller than the injectivity radius of  $(\widehat{M}, g)$ , any two geodesic segments of length  $\leq r$  can intersect transversally in at most one point (locally geodesics are close to straight lines). Partitioning  $(a, b)$  in disjoint intervals  $\{J_k\}_{k=1}^K$  of length  $\leq r$ , we have an injective map

$$\{(t, t') \in (a, b)^2 \mid t < t' \text{ and } \gamma(t) = \gamma(t')\} \mapsto \{(k, l) \in \{1, \dots, K\}^2 : t \in J_k, t' \in J_l\}.$$

Consequently,  $\gamma$  can only self-intersect finitely many times. □

**Lemma 7.3.** *Let  $F$  be a  $C^1$  map from a neighborhood of  $(a, b) \times \{0\}$  in  $\mathbb{R}^n$  into a smooth manifold such that  $F|_{(a,b) \times \{0\}}$  is injective and  $DF(t, 0)$  is invertible for  $t \in (a, b)$ . If  $[a_0, b_0]$  is a closed subinterval of  $(a, b)$ , then  $F$  is a  $C^1$  diffeomorphism in some neighborhood of  $[a_0, b_0] \times \{0\}$  in  $\mathbb{R}^n$ .*

*Proof.* For any  $t \in [a_0, b_0]$ , the inverse function theorem implies that there is  $\varepsilon_t > 0$  such that  $F|_{(t-3\varepsilon_t, t+3\varepsilon_t) \times B_{3\varepsilon_t}(0)}$  is a  $C^1$  diffeomorphism. Since  $[a_0, b_0]$  is covered by the intervals  $(t - \varepsilon_t, t + \varepsilon_t)$ , by compactness we have  $[a_0, b_0] \subset \bigcup_{j=1}^N (t_j - \varepsilon_j, t_j + \varepsilon_j)$ , where  $F|_{(t_j-3\varepsilon_j, t_j+3\varepsilon_j) \times B_{3\varepsilon_j}(0)}$  is bijective. We can further assume (upon throwing away or shrinking some intervals if necessary) that the intervals  $I_j = (t_j - \varepsilon_j, t_j + \varepsilon_j)$  satisfy  $\bar{I}_j \cap \bar{I}_k = \emptyset$  unless  $|j - k| \leq 1$ . Since  $\gamma(t) = F(t, 0)$  is injective, we also have  $\gamma(\bar{I}_j) \cap \gamma(\bar{I}_k) = \emptyset$  unless  $|j - k| \leq 1$ .

Fix a Riemannian metric in the target manifold, and define

$$\delta = \inf \{\text{dist}(\gamma(\bar{I}_j), \gamma(\bar{I}_k)) : |j - k| \geq 2\} > 0.$$

Let  $U_j = I_j \times B_\varepsilon(0)$ , where  $\varepsilon < \min\{\varepsilon_1, \dots, \varepsilon_N\}$  is chosen so small that  $F(U_j) \subset \{q : \text{dist}(q, \gamma(\bar{I}_j)) < \delta/2\}$ . Then  $F(U_j) \cap F(U_k) = \emptyset$  unless  $|j - k| \leq 1$ . Define

$$U = \bigcup_{j=1}^N U_j.$$

To show that  $F|_U$  is a  $C^1$  diffeomorphism, it is enough to check injectivity. If  $F(t, y) = F(t', y')$  for  $(t, y), (t', y') \in U$ , then, necessarily,  $(t, y) \in U_j, (t', y') \in U_k$ , where  $|j - k| \leq 1$ . We may assume that  $\varepsilon_j \geq \varepsilon_k$ . Since  $F|_{(t_j-3\varepsilon_j, t_j+3\varepsilon_j) \times B_{3\varepsilon_j}(0)}$  is bijective, we obtain  $(t, y) = (t', y')$ . □

**Lemma 7.4.** *Let  $(\widehat{M}, g)$  be a compact manifold without boundary, and assume that  $\gamma : (a, b) \rightarrow \widehat{M}$  is a unit speed geodesic segment with no loops. Given a closed subinterval  $[a_0, b_0]$  of  $(a, b)$  such that  $\gamma|_{[a_0, b_0]}$  self-intersects only at times  $t_j$  with  $a_0 < t_1 < \dots < t_N < b_0$  (set  $t_0 = a_0$  and  $t_{N+1} = b_0$ ), there is an open cover  $\{(U_j, \varphi_j)\}_{j=0}^{N+1}$  of  $\gamma([a_0, b_0])$  consisting of coordinate neighborhoods having the following properties.*

- (1)  $\varphi_j(U_j) = I_j \times B$ , where  $I_j$  are open intervals and  $B = B(0, \delta)$  is an open ball in  $\mathbb{R}^{n-1}$  where  $\delta$  can be taken arbitrarily small.
- (2)  $\varphi_j(\gamma(t)) = (t, 0)$  for  $t \in I_j$ .
- (3)  $t_j$  only belongs to  $I_j$  and  $\bar{I}_j \cap \bar{I}_k = \emptyset$  unless  $|j - k| \leq 1$ .
- (4)  $\varphi_j = \varphi_k$  on  $\varphi_j^{-1}((I_j \cap I_k) \times B)$ .

Further, if  $S$  is a hypersurface through  $\gamma(a_0)$  that is transversal to  $\dot{\gamma}(a_0)$ , one can arrange for the map  $y \mapsto \varphi_0^{-1}(a_0, y)$  to parametrize  $S$  near  $\gamma(a_0)$ .

*Proof.* We will use modified Fermi coordinates, constructed as follows. Let  $\{v_1, \dots, v_{n-1}\}$  be an orthonormal set of vectors in  $T_{\gamma(a_0)}\widehat{M}$  such that  $\{\dot{\gamma}(a_0), v_1, \dots, v_{n-1}\}$  is a basis. (The case where  $\{\dot{\gamma}(a_0), v_1, \dots, v_{n-1}\}$  is an orthonormal basis corresponds to the usual Fermi coordinates.) Let  $E_\alpha(t)$  be the parallel transport of  $v_\alpha$  along the geodesic  $\gamma$ . Since  $\dot{\gamma}(t)$  is also parallel along  $\gamma$ , the set  $\{\dot{\gamma}(t), E_1(t), \dots, E_{n-1}(t)\}$  is a basis of  $T_{\gamma(t)}\widehat{M}$  for  $t \in (a, b)$ .

Define the function

$$F : (a, b) \times \mathbb{R}^{n-1} \rightarrow \widehat{M}, \quad F(t, y) = \exp_{\gamma(t)}(y^\alpha E_\alpha(t)).$$

Here  $\exp$  is the exponential map in  $(\widehat{M}, g)$  and  $\alpha, \beta$  run from 1 to  $n - 1$ . Then  $F(t, 0) = \gamma(t)$  and (with  $e_\alpha$  the  $\alpha$ -th coordinate vector)

$$\frac{\partial}{\partial s} F(t, se_\alpha) \Big|_{s=0} = E_\alpha(t), \quad \frac{\partial}{\partial t} F(t, 0) = \dot{\gamma}(t).$$

Thus  $F$  is a  $C^\infty$  map near  $(a, b) \times \{0\}$  such that  $DF(t, 0)$  is invertible for  $t \in (a, b)$ .

In the case where  $\gamma$  does not self-intersect,  $F|_{(a,b) \times \{0\}}$  is injective and Lemma 7.3 implies the existence of a single coordinate neighborhood of  $\gamma([a_0, b_0])$  so that (1) and (2) are satisfied (then (3) and (4) are void). In the general case, by Lemma 7.2 the geodesic segment  $\gamma|_{[a_0, b_0]}$  only self-intersects at finitely many times  $t_j$  with  $a_0 < t_1 < \dots < t_N < b_0$ . For some sufficiently small  $\delta$ ,  $\gamma$  is injective on the intervals  $(a, t_1 - \delta)$ ,  $(t_1 - 2\delta, t_2 - \delta)$ ,  $\dots$ ,  $(t_N - 2\delta, b)$  and each interval intersects at most two of the others. Restricting the map  $F$  above to suitable neighborhoods corresponding to these intervals (or slightly smaller ones) and using Lemma 7.3, we obtain the required coordinate charts with  $\varphi_j = F^{-1}|_{U_j}$ .

Let  $S$  be a hypersurface transversal to  $\dot{\gamma}(a_0)$ , and choose some parametrization  $y \mapsto q(y)$  of  $S$  near  $\gamma(a_0)$  satisfying  $(\partial/\partial s)q(se_\alpha) = v_\alpha$ . We will form a new chart  $(\tilde{U}_0, \tilde{\varphi}_0)$  by modifying  $(U_0, \varphi_0)$  so that  $y \mapsto \tilde{\varphi}_0^{-1}(a_0, y)$  parametrizes  $S$  near  $\gamma(a_0)$ .

We may assume that  $a_0 = 0$ , and write  $F_0 = \varphi_0^{-1}$ ,  $\tilde{F}_0 = \tilde{\varphi}_0^{-1}$ . It is enough to choose  $\tilde{F}_0 = F_0 \circ \Phi$ , where  $\Phi$  is a diffeomorphism near  $\bar{I}_0 \times B$  such that

$$\begin{aligned} \Phi(t, 0) &= (t, 0), \\ \Phi(0, y) &= F_0^{-1}(q(y)), \\ \Phi(t, y) &= (t, y) \quad \text{for } t > c \text{ with suitable } c > 0. \end{aligned}$$



Write the components of  $\tilde{q} = F_0^{-1} \circ q$  as Taylor series

$$\tilde{q}^j(y) = \tilde{q}^j(0) + \nabla \tilde{q}^j(0) \cdot y + H^j(y)y \cdot y,$$

where  $H^j$  are smooth matrices, and  $j = 0, \dots, n - 1$  ( $t$  is the 0-th variable). The properties of  $q$  imply that

$$\tilde{q}^j(0) = 0, \quad \partial_\beta \tilde{q}^0(0) = 0, \quad \partial_\beta \tilde{q}^\alpha(0) = \delta_\beta^\alpha.$$

We look for  $\Phi$  in the form

$$\Phi^j(t, y) = f^j(t) + a^j(t) \cdot y + R^j(t, y)y \cdot y$$

for some smooth functions  $f^j$ , vectors  $a^j$ , and matrices  $R^j$ . The conditions for  $\Phi$  motivate the choices

$$f^0(t) = t, \quad f^\alpha(t) = 0, \quad a_\beta^0(t) = 0, \quad a_\beta^\alpha(t) = \delta_\beta^\alpha.$$

We choose  $R^j(t, y)$  to be a smooth matrix with  $R^j(0, y) = H^j(y)$  and  $R^j(t, y) = 0$  for  $t > c$ . Then  $D\Phi(t, 0) = \text{Id}$ , and Lemma 7.3 ensures that  $\Phi$  is a diffeomorphism near  $\bar{I}_0 \times B$ , possibly after decreasing  $B$ . □

The next result gives the construction of (nonreflected) Gaussian beam quasimodes associated with a finite length geodesic segment that enters and exits the domain nontangentially. To prepare for the reflected case, we also consider the possibility of prescribing the boundary values of the quasimode at least up to high order at a point. Recall that if  $f$  is a smooth function having a critical point at  $p$ , the Hessian of  $f$  at  $p$  is the quadratic form

$$\text{Hess}_p(f)(\dot{\eta}(0), \dot{\eta}(0)) = (f \circ \eta)''(0),$$

where  $\eta$  is any smooth curve with  $\eta(0) = p$ .

**Proposition 7.5.** *Let  $\gamma : [0, L] \rightarrow M$  be any unit speed geodesic in  $(M, g)$  such that  $\gamma(0), \gamma(L) \in \partial M$ ,  $\dot{\gamma}(0)$  and  $\dot{\gamma}(L)$  are nontangential, and  $\gamma(t) \in M^{\text{int}}$  for  $0 < t < L$ . Also let  $\lambda$  be a fixed real number. For any  $K > 0$  there is a family  $\{v_s : s = \tau + i\lambda, \tau \geq 1\}$  in  $C^\infty(M)$  such that, as  $\tau \rightarrow \infty$ ,*

$$\|(-\Delta_g - s^2)v_s\|_{L^2(M)} = O(\tau^{-K}), \quad \|v_s\|_{L^2(M)} = O(1), \quad \|v_s\|_{L^2(\partial M)} = O(1),$$

and, for any  $\psi \in C(M)$ ,

$$\int_M |v_{\tau+i\lambda}|^2 \psi \, dV_g \rightarrow \int_0^L e^{-2\lambda t} \psi(\gamma(t)) \, dt \quad \text{as } \tau \rightarrow \infty. \tag{7-1}$$

Given any neighborhood of  $\gamma([0, L])$ , one can arrange for each  $v_s$  to be supported in this neighborhood, and away from the points where  $\gamma$  self-intersects one has  $v_s = e^{is\Theta} a$  where  $\Theta$  and  $a$  are smooth complex functions with

$$d\Theta(\dot{\gamma}(t)) = \dot{\gamma}(t)^\flat, \quad a(\gamma(t)) \neq 0 \text{ for } \tau \text{ large.}$$

If  $\gamma$  does not self-intersect at  $\gamma(0)$ , the  $K$ -th order jets of  $\Theta|_{\partial M}$  and  $a|_{\partial M}$  can be prescribed freely at  $\gamma(0)$  except for the following restrictions:  $d\Theta(\gamma(0)) = \dot{\gamma}(0)^b$ , the Hessian of  $\text{Im}(\Theta|_{\partial M})$  at  $\gamma(0)$  is positive definite, and  $a(\gamma(0)) \neq 0$ .

*Proof.* We embed  $(M, g)$  in a compact manifold  $(\widehat{M}, g)$  without boundary and extend  $\gamma$  as a unit speed geodesic in  $\widehat{M}$ . Choose  $\varepsilon > 0$  so that  $\gamma(t)$  lies in  $\widehat{M} \setminus M$  and has no self-intersections in the interval  $t \in [-2\varepsilon, 0) \cup (L, L + 2\varepsilon]$ . We will construct a Gaussian beam quasimode in a neighborhood of  $\gamma([-\varepsilon, L + \varepsilon])$ .

Fix a point  $p_0 = \gamma(t_0)$  on  $\gamma([-\varepsilon, L + \varepsilon])$  and let  $(t, y)$  be any local coordinates near  $p_0$ , defined in  $U = \{(t, y) : t \in I, |y| < \delta\}$  for some open interval  $I$  containing  $t_0$ , such that  $p_0$  corresponds to  $(t_0, 0)$  and the geodesic near  $p_0$  is given by  $\Gamma = \{(t, 0) : t \in I\}$ . Write  $x = (t, y)$ , where  $x_1 = t$  and  $(x_2, \dots, x_m) = y$ . We seek to find a quasimode  $v_s$  concentrated near  $\Gamma$ , having the form

$$v_s = e^{is\Theta} a,$$

where  $s = \tau + i\lambda$ , and  $\Theta$  and  $a$  are smooth complex functions near  $\Gamma$  with  $a$  supported in  $\{|y| < \delta/2\}$ .

We compute

$$(-\Delta - s^2)v_s = f,$$

where

$$f = e^{is\Theta} (s^2[\langle d\Theta, d\Theta \rangle - 1]a - is[2\langle d\Theta, da \rangle + (\Delta\Theta)a] - \Delta a).$$

We first choose  $\Theta$  so that

$$\langle d\Theta, d\Theta \rangle = 1 \quad \text{to } N\text{-th order on } \Gamma. \tag{7-2}$$

In fact, we look for  $\Theta$  of the form  $\Theta = \sum_{j=0}^N \Theta_j$  where

$$\Theta_j(t, y) = \sum_{|\gamma|=j} \frac{\Theta_{j,\gamma}(t)}{\gamma!} y^\gamma.$$

We also write  $g^{jk} = \sum_{l=0}^N g_l^{jk} + g_{N+1}^{jk}$ , where

$$g_l^{jk}(t, y) = \sum_{|\beta|=l} \frac{g_{l,\beta}^{jk}(t)}{\beta!} y^\beta, \quad g_{N+1}^{jk} = O(|y|^{N+1}).$$

Set  $g_l^{jk} = 0$  for  $l \geq N + 2$ .

With the understanding that  $j, k$  run from 1 to  $m$  and  $\alpha, \beta$  run from 2 to  $m$ , the main part of the argument will consist of finding suitable  $\Theta_0, \Theta_1$ , and  $\Theta_2$  in the following form:

$$\Theta_0(t) \text{ real-valued, } \quad \Theta_1(t) = \xi_\alpha(t)y^\alpha \text{ with } \xi_\alpha(t) \text{ real-valued, } \quad \Theta_2(t) = \frac{1}{2}H_{\alpha\beta}(t)y^\alpha y^\beta,$$

where  $H(t) = (H_{\alpha\beta}(t))$  is a complex symmetric matrix,  $H_{\alpha\beta} = H_{\beta\alpha}$ , such that  $\text{Im}(H(t))$  is positive definite for all  $t$ . We also write

$$\xi_1(t) = \partial_t \Theta_0(t).$$

Since  $\partial_t \Theta_0 = \xi_1$  and  $\partial_\alpha \Theta_1 = \xi_\alpha$ , we compute

$$\begin{aligned} &g^{jk} \partial_j \Theta \partial_k \Theta - 1 \\ &= g^{11} (\partial_t \Theta_0 + \partial_t \Theta_1 + \dots) (\partial_t \Theta_0 + \partial_t \Theta_1 + \dots) + 2g^{1\alpha} (\partial_t \Theta_0 + \partial_t \Theta_1 + \dots) (\partial_\alpha \Theta_1 + \partial_\alpha \Theta_2 + \dots) \\ &\quad + g^{\alpha\beta} (\partial_\alpha \Theta_1 + \partial_\alpha \Theta_2 + \dots) (\partial_\beta \Theta_1 + \partial_\beta \Theta_2 + \dots) - 1 \\ &= g^{jk} \xi_j \xi_k + 2g^{11} \xi_1 (\partial_t \Theta_1 + \dots) + 2g^{1\alpha} \xi_1 (\partial_\alpha \Theta_2 + \dots) \\ &\quad + 2g^{1\alpha} \xi_\alpha (\partial_t \Theta_1 + \dots) + 2g^{\alpha\beta} \xi_\alpha (\partial_\beta \Theta_2 + \dots) + g^{11} (\partial_t \Theta_1 + \dots) (\partial_t \Theta_1 + \dots) \\ &\quad + 2g^{1\alpha} (\partial_t \Theta_1 + \dots) (\partial_\alpha \Theta_2 + \dots) + g^{\alpha\beta} (\partial_\alpha \Theta_2 + \dots) (\partial_\beta \Theta_2 + \dots) - 1 \\ &= g^{jk} \xi_j \xi_k + 2g^{1k} \xi_k (\partial_t \Theta_1 + \dots) + 2g^{\alpha k} \xi_k (\partial_\alpha \Theta_2 + \dots) + g^{11} (\partial_t \Theta_1 + \dots) (\partial_t \Theta_1 + \dots) \\ &\quad + 2g^{1\alpha} (\partial_t \Theta_1 + \dots) (\partial_\alpha \Theta_2 + \dots) + g^{\alpha\beta} (\partial_\alpha \Theta_2 + \dots) (\partial_\beta \Theta_2 + \dots) - 1. \end{aligned}$$

Writing  $g^{jk} = g_0^{jk} + g_1^{jk} + \dots$  and grouping like powers of  $y$ , we obtain

$$\begin{aligned} &g^{jk} \partial_j \Theta \partial_k \Theta - 1 \\ &= [g_0^{jk} \xi_j \xi_k - 1] + [g_1^{jk} \xi_j \xi_k + 2g_0^{1k} \xi_k \dot{\xi}_\beta y^\beta + 2g_0^{\alpha k} \xi_k H_{\alpha\beta} y^\beta] + (g_2^{jk} + \dots) \xi_j \xi_k + 2g_0^{1k} \xi_k (\partial_t \Theta_2 + \dots) \\ &\quad + 2(g_1^{1k} + \dots) \xi_k (\partial_t \Theta_1 + \dots) + 2g_0^{\alpha k} \xi_k (\partial_\alpha \Theta_3 + \dots) + 2(g_1^{\alpha k} + \dots) \xi_k (\partial_\alpha \Theta_2 + \dots) \\ &\quad + g^{11} (\partial_t \Theta_1 + \dots) (\partial_t \Theta_1 + \dots) + 2g^{1\alpha} (\partial_t \Theta_1 + \dots) (\partial_\alpha \Theta_2 + \dots) \\ &\quad + g^{\alpha\beta} (\partial_\alpha \Theta_2 + \dots) (\partial_\beta \Theta_2 + \dots). \quad (7-3) \end{aligned}$$

We can make the two expressions in brackets vanish by choosing  $\xi(t)$  to be part of the solution  $(x(t), \xi(t))$  of the cogeodesic flow with Hamiltonian  $h(x, \xi) = \frac{1}{2} g^{jk}(x) \xi_j \xi_k$ ,

$$\begin{aligned} \dot{x}^j(t) &= \partial_{\xi_j} h(x(t), \xi(t)), \\ \dot{\xi}_j(t) &= -\partial_{x^j} h(x(t), \xi(t)). \end{aligned}$$

There is a unique solution with  $x(t_0) = p_0$  and  $\xi(t_0) = \dot{\gamma}(t_0)^b$  (here we raise and lower indices with respect to the metric  $g$ ). It follows that  $x(t)$  is the unit speed geodesic  $t \mapsto (t, 0)$ , and  $\xi^j(t) = \dot{x}^j(t)$ . Then  $g_0^{jk} \xi_j \xi_k = 1$ , and with our choice of coordinates  $\xi^1 = 1$  and  $\xi^\alpha = 0$  so that also

$$g_0^{1k} \xi_k = 1, \quad g_0^{\alpha k} \xi_k = 0.$$

We further have

$$\dot{\xi}_\beta y^\beta = -\frac{1}{2} \partial_{x_\beta} g^{jk}(t, 0) \xi_j \xi_k y^\beta = -\frac{1}{2} g_1^{jk} \xi_j \xi_k.$$

Noting that  $\partial_1$  has unit length, we have

$$\xi_1 = g_{1k}(t, 0) \xi^k = 1.$$

Since  $\xi_\alpha = g_{\alpha k}(t, 0) \xi^k = g_{\alpha 1}(t, 0)$ , we can therefore choose

$$\begin{aligned} \Theta_0(t) &= t, \\ \Theta_1(t) &= g_{\alpha 1}(t, 0) y^\alpha. \end{aligned}$$

Using these choices and the facts above, in (7-3), the expressions in brackets will indeed vanish, and one obtains

$$\begin{aligned}
 &g^{jk} \partial_j \Theta \partial_k \Theta - 1 \\
 &= (g_2^{jk} + \dots) \xi_j \xi_k + 2(\partial_t \Theta_2 + \dots) + 2(g_1^{1k} + \dots) \xi_k (\partial_t \Theta_1 + \dots) + 2(g_1^{\alpha k} + \dots) \xi_k (\partial_\alpha \Theta_2 + \dots) \\
 &\quad + g^{11} (\partial_t \Theta_1 + \dots) (\partial_t \Theta_1 + \dots) + 2g^{1\alpha} (\partial_t \Theta_1 + \dots) (\partial_\alpha \Theta_2 + \dots) + g^{\alpha\beta} (\partial_\alpha \Theta_2 + \dots) (\partial_\beta \Theta_2 + \dots) \\
 &= (g_2^{jk} \xi_j \xi_k + 2\partial_t \Theta_2 + 2g_1^{\alpha k} \xi_k \partial_\alpha \Theta_2 + 2g_0^{1\alpha} \partial_t \Theta_1 \partial_\alpha \Theta_2 + g_0^{\alpha\beta} \partial_\alpha \Theta_2 \partial_\beta \Theta_2 + 2g_1^{1k} \xi_k \partial_t \Theta_1 + g_0^{11} (\partial_t \Theta_1)^2) \\
 &\quad + \sum_{p=3}^N \left( g_p^{jk} \xi_j \xi_k + 2\partial_t \Theta_p + 2g_1^{\alpha k} \xi_k \partial_\alpha \Theta_p + 2g_0^{1\alpha} \partial_t \Theta_1 \partial_\alpha \Theta_p + 2g_0^{\alpha\beta} \partial_\alpha \Theta_2 \partial_\beta \Theta_p \right. \\
 &\quad + 2 \sum_{l=1}^{p-1} g_{p-l}^{1k} \xi_k \partial_t \Theta_l + 2 \sum_{l=2}^{p-1} g_{p-l+1}^{\alpha k} \xi_k \partial_\alpha \Theta_l + \sum_{l=0}^{p-2} g_l^{11} \sum_{\substack{r+s=p-l \\ 1 \leq r, s < p}} \partial_t \Theta_r \partial_t \Theta_s \\
 &\quad \left. + \sum_{l=0}^{p-2} g_l^{1\alpha} \sum_{\substack{r+s=p-l+1 \\ 1 \leq r < p \\ 2 \leq s < p}} \partial_t \Theta_r \partial_\alpha \Theta_s + \sum_{l=0}^{p-2} g_l^{\alpha\beta} \sum_{\substack{r+s=p-l+2 \\ 2 \leq r, s < p}} \partial_\alpha \Theta_r \partial_\beta \Theta_s \right) + O(|y|^{N+1}).
 \end{aligned}$$

We want to choose  $\Theta_2$  so that the first term in brackets vanishes. Recalling that we are looking for  $\Theta_2$  in the form  $\Theta_2(t, y) = \frac{1}{2} H_{\alpha\beta}(t) y^\alpha y^\beta$ , where  $H(t)$  is a smooth complex symmetric matrix; it follows that  $H$  should satisfy the matrix equation

$$\dot{H}_{\alpha\beta} y^\alpha y^\beta + 2g_1^{\gamma k} \xi_k H_{\gamma\beta} y^\beta + 2g_0^{1\gamma} \partial_t \Theta_1 H_{\gamma\beta} y^\beta + g_0^{\gamma\delta} H_{\gamma\alpha} H_{\delta\beta} y^\alpha y^\beta = F_{\alpha\beta} y^\alpha y^\beta,$$

where  $F(t)$  is a real-valued smooth symmetric matrix. This can be further written as the matrix Riccati equation

$$\dot{H} + BH + HB^t + HCH = F,$$

where  $B(t)$  and  $C(t)$  are real smooth matrices and  $C$  is symmetric. More precisely, since  $g_1^{jk} = \partial_\alpha g^{jk}(t, 0) y^\alpha$ , we have

$$B_\alpha^\gamma = \partial_\alpha g^{\gamma k}(t, 0) \xi_k + g_0^{1\gamma} \xi_\alpha, \quad C^{\gamma\delta} = g_0^{\gamma\delta}. \tag{7-4}$$

Choosing  $H(t_0) = H_0$ , where  $H_0$  is a complex symmetric matrix with  $\text{Im}(H_0)$  positive definite, it follows that the Riccati equation has a unique smooth complex symmetric solution  $H(t)$  with  $\text{Im}(H(t))$  positive definite; see [Katchalov et al. 2001]. This completes the construction of  $\Theta_2$ . From the lower order terms we can find  $\Theta_3, \dots, \Theta_N$  successively by solving linear first order ODEs on  $\Gamma$  with prescribed initial conditions at  $t_0$ . In this way, we obtain a smooth  $\Theta$  satisfying (7-2).

The next step is to find  $a$  such that

$$s[2\langle d\Theta, da \rangle + (\Delta\Theta)a] - i\Delta a = 0 \quad \text{to } N\text{-th order on } \Gamma.$$

We look for  $a$  in the form

$$a = \tau^{(m-1)/4} (a_0 + s^{-1} a_{-1} + \dots + s^{-N} a_{-N}) \chi(y/\delta'),$$

where  $\chi$  is a smooth function with  $\chi = 1$  for  $|y| \leq 1/4$  and  $\chi = 0$  for  $|y| \geq 1/2$ . Writing  $\eta = \Delta\Theta$ , it is sufficient to determine  $a_j$  so that

$$\begin{aligned} 2\langle d\Theta, da_0 \rangle + \eta a_0 &= 0 \quad \text{to } N\text{-th order on } \Gamma, \\ 2\langle d\Theta, da_{-1} \rangle + \eta a_{-1} - i\Delta a_0 &= 0 \quad \text{to } N\text{-th order on } \Gamma, \\ &\vdots \\ 2\langle d\Theta, da_{-N} \rangle + \eta a_{-N} - i\Delta a_{-(N-1)} &= 0 \quad \text{to } N\text{-th order on } \Gamma. \end{aligned}$$

Consider  $a_0 = a_{00} + \dots + a_{0N}$ , where  $a_{0j}(t, y)$  is a polynomial of order  $j$  in  $y$ , and similarly let  $\eta = \eta_0 + \dots + \eta_N$ . We compute

$$\begin{aligned} 2\langle d\Theta, da_0 \rangle + \eta a_0 &= 2(g_0^{11} + \dots)(\partial_t \Theta_0 + \dots)(\partial_t a_{00} + \dots) + 2(g_0^{1\alpha} + \dots)(\partial_t \Theta_0 + \dots)(\partial_\alpha a_{01} + \dots) \\ &\quad + 2(g_0^{1\alpha} + \dots)(\partial_\alpha \Theta_1 + \dots)(\partial_t a_{00} + \dots) + 2(g_0^{\alpha\beta} + \dots)(\partial_\beta \Theta_1 + \dots)(\partial_\alpha a_{01} + \dots) \\ &\quad + (\eta_0 + \eta_1 + \dots)(a_{00} + a_{01} + \dots). \end{aligned}$$

Recalling that  $\partial_t \Theta_0 = \xi_1 = 1$  and  $\partial_\alpha \Theta_1 = \xi_\alpha$ , where  $g_0^{1j} \xi_j = 1$  and  $g_0^{\alpha j} \xi_j = 0$ , we obtain

$$\begin{aligned} 2\langle d\Theta, da_0 \rangle + \eta a_0 &= 2[g_0^{11} \xi_1 + g_0^{11}(\partial_t \Theta_1 + \dots) + (g_1^{11} + \dots)(\partial_t \Theta_0 + \dots) + g_0^{1\alpha} \xi_\alpha + g_0^{1\alpha}(\partial_\alpha \Theta_2 + \dots) + (g_1^{1\alpha} + \dots)(\partial_\alpha \Theta_1 + \dots)] \\ &\quad \times (\partial_t a_{00} + \dots) \\ &\quad + 2[g_0^{1\alpha}(\partial_t \Theta_1 + \dots) + (g_1^{1\alpha} + \dots)(\partial_t \Theta_0 + \dots) + g_0^{\alpha\beta}(\partial_\beta \Theta_2 + \dots) + (g_1^{\alpha\beta} + \dots)(\partial_\beta \Theta_1 + \dots)] \\ &\quad \times (\partial_\alpha a_{01} + \dots) + (\eta_0 + \eta_1 + \dots)(a_{00} + a_{01} + \dots) \\ &= [2\partial_t a_{00} + \eta_0 a_{00}] + \sum_{p=1}^N [2\partial_t a_{0p} + q_p^{\alpha\beta} y^\beta \partial_\alpha a_{0p} + \eta_0 a_{0p} + F_p] + O(|y|^{N+1}), \end{aligned}$$

where  $q_p^{\alpha\beta}(t)$  are smooth functions only depending on  $g$  and  $\Theta$ , and  $F_p(t, y)$  is a polynomial of degree  $p$  in  $y$  that only depends on  $g, \Theta, \eta$ , and  $a_{00}, \dots, a_{0,p-1}$ .

We want to choose  $a_{00}$  so that the first term in brackets vanishes, that is,

$$\partial_t a_{00} + t \frac{1}{2} \eta_0 a_{00} = 0.$$

This has the solution

$$a_{00}(t) = c_0 e^{-(1/2) \int_{t_0}^t \eta_0(s) ds}, \quad a_{00}(t_0) = c_0.$$

We obtain  $a_{01}, \dots, a_{0N}$  successively by solving linear first order ODEs with prescribed initial conditions at  $t_0$ . The functions  $a_1, \dots, a_N$  may be determined in a similar way so that the required equations are satisfied to  $N$ -th order on  $\Gamma$ . This completes the construction of  $a$ .

To review what has been achieved so far, we have constructed a function  $v_s = e^{is\Theta}a$  in  $U$ , where

$$\begin{aligned} \Theta(t, y) &= t + \xi_\alpha(t)y^\alpha + \frac{1}{2}H_{\alpha\beta}(t)y^\alpha y^\beta + \tilde{\Theta}, \\ a(t, y) &= \tau^{(m-1)/4}(a_0 + s^{-1}a_{-1} + \dots + s^{-N}a_{-N})\chi(y/\delta'), \\ a_0(t, 0) &= c_0 e^{-(1/2)\int_{t_0}^t \eta_0(s) ds}. \end{aligned}$$

Here  $\tilde{\Theta} = O(|y|^3)$  and  $\tilde{\Theta}$  and each  $a_j$  are independent of  $\tau$ . Also,  $f = (-\Delta - s^2)v_s$  has the form

$$f = e^{is\Theta}\tau^{(m-1)/4}(s^2h_2a + sh_1 + \dots + s^{-(N-1)}h_{-(N-1)} - s^{-N}\Delta a_{-N})\chi(y/\delta') + e^{is\Theta}\tau^{(m-1)/4}sb\tilde{\chi}(y/\delta'),$$

where, for each  $j$ , one has  $h_j = 0$  to  $N$ -th order on  $\Gamma$ ,  $b$  vanishes near  $\Gamma$ , and  $\tilde{\chi}$  is a smooth function with  $\tilde{\chi} = 0$  for  $|y| \geq 1/2$ . We also note that  $d\Theta(\gamma(t)) = \dot{\gamma}(t)^b$  and  $\text{Hess}_{\gamma(t_0)}(\text{Im}(\Theta)|_{\{t=t_0\}}) = \text{Im}(H(t_0))$ .

To prove the norm estimates for  $v_s$  in  $U$ , note that

$$|e^{is\Theta}| = e^{-\lambda \text{Re } \Theta} e^{-\tau \text{Im } \Theta} = e^{-\lambda t} e^{-(1/2)\tau \text{Im}(H(t))y \cdot y} e^{-\lambda O(|y|)} e^{-\tau O(|y|^3)}.$$

Here  $\text{Im}(H(t))y \cdot y \geq c|y|^2$  for  $(t, y) \in U$ , where  $c > 0$  depends on  $H_0$ ,  $g$ , and  $I$ . By decreasing  $\delta'$  if necessary, this shows the following bound when  $t$  in a fixed compact set:

$$|v_s(t, y)| \lesssim \tau^{(m-1)/4} e^{-(1/4)c\tau|y|^2} \chi(y/\delta').$$

Integrating the square of this over  $U$ , we get, as  $\tau \rightarrow \infty$ ,

$$\|v_s\|_{L^2(U)} \lesssim \|\tau^{(m-1)/4} e^{-(1/4)c\tau|y|^2}\|_{L^2(U)} = O(1).$$

Similarly, we have

$$\begin{aligned} \|(-\Delta - s^2)v_s\|_{L^2(U)} &\lesssim \|\tau^{(m-1)/4} e^{-(1/4)c\tau|y|^2} (\tau^2|y|^{N+1} + \tau^{-N})\|_{L^2(U)} \\ &= O(\tau^{(3-N)/2}). \end{aligned}$$

The norm estimates for  $v_s$  in  $U$  follow upon replacing  $N$  by  $2K + 3$ .

For the  $L^2(\partial M)$  estimate, if  $U$  contains a boundary point  $x_0 = (t_0, 0) \in \partial M$ , by assumption  $(\partial/\partial t)|_{x_0}$  is transversal to  $\partial M$ . If  $\rho$  is a boundary defining function for  $M$ , so  $\partial M$  is given as the zero set  $\rho(t, y) = 0$  near  $x_0$  and  $\nabla\rho = -\nu$  on  $\partial M$ , then  $(\partial\rho/\partial t)(x_0) \neq 0$  and, by the implicit function theorem, there is a smooth function  $y \mapsto t(y)$  near 0 such that  $\partial M$  is given by  $\{(t(y), y) : |y| < r_0\}$  near  $x_0$ . The bound for  $v_s$  given above implies that, for  $\delta'$  small,

$$\|v_s\|_{L^2(\partial M \cap U)}^2 = \int_{|y| < r_0} |v_s(t(y), y)| dS(y) \lesssim \int_{|y| < r_0} \tau^{(m-1)/2} e^{-(1/2)c\tau|y|^2} dy = O(1).$$

At this point we can construct the quasimode  $v_s$  in  $M$  from the corresponding quasimodes defined on small pieces. Let  $\gamma([-\varepsilon, L + \varepsilon])$  be covered by open sets  $U^{(0)}, \dots, U^{(N+1)}$  as in Lemma 7.4, and note that each  $U^{(j)}$  corresponds to  $I_j \times B(0, \delta)$  in the  $(t, y)$  coordinates. Suppose first that  $\gamma$  does not self-intersect at time  $t = 0$ . We find a quasimode  $v^{(0)} = e^{is\Theta^{(0)}}a^{(0)}$  in  $U^{(0)}$  by the above procedure, with some fixed initial conditions at  $t = 0$  for the ODEs determining  $\Theta^{(0)}$  and  $a^{(0)}$ . Choose some  $t'_0$  with  $\gamma(t'_0) \in U^{(0)} \cap U^{(1)}$ , and construct a quasimode  $v^{(1)} = e^{is\Theta^{(1)}}a^{(1)}$  in  $U^{(1)}$  by choosing the initial conditions for the ODEs for  $\Theta^{(1)}$  and  $a^{(1)}$  at  $t'_0$  to be the corresponding values of  $\Theta^{(0)}$  and  $a^{(0)}$  at  $t'_0$ . Continuing

in this way, we obtain  $v^{(2)}, \dots, v^{(N+1)}$ . If  $\gamma$  self-intersects at  $t = 0$ , we start the construction from  $v^{(1)}$  fixing initial conditions for the ODEs at  $t = 0$ , and find  $v^{(0)}$  by going backward.

Let  $\{\tilde{\chi}_j(t)\}$  be a partition of unity near  $[-\varepsilon, L + \varepsilon]$  corresponding to the cover  $\{I_j\}$ , let  $\chi_j(t, y) = \tilde{\chi}_j(t)$  on  $U^{(j)}$ , and define

$$v_s = \sum_{j=0}^{N+1} \chi_j v^{(j)}.$$

Then  $v_s$  is smooth in  $\widehat{M}$  and it is supported in a small neighborhood of  $\gamma([-\varepsilon, L + \varepsilon])$ . The important point is that since the ODEs for the phase functions and amplitudes have the same initial data in  $U^{(j)}$  and in  $U^{(j+1)}$ , and since the local coordinates  $\varphi_j$  and  $\varphi_{j+1}$  coincide on  $\varphi_j^{-1}((I_j \cap I_{j+1}) \times B)$ , one actually has  $v^{(j)} = v^{(j+1)}$  in  $\varphi_j^{-1}((I_j \cap I_{j+1}) \times B)$ . Letting  $p_1, \dots, p_R$  be the points where  $\gamma$  intersects itself, we choose an open cover of  $\text{supp}(v_s) \cap M$ ,

$$\text{supp}(v_s) \cap M \subset \left( \bigcup_{r=1}^R V_r \right) \cup \left( \bigcup_{j=0}^{N+1} (W_{j,0} \cup W_{j,1}) \right),$$

where  $V_r$  are small neighborhoods of the points  $p_r$  and  $W_{j,0}, W_{j,1} \subset U^{(j)}$ , such that

$$v_s|_{V_r} = \sum_{\gamma(t_j)=p_r} v^{(j)} \quad \text{and} \quad v_s|_{W_{j,l}} = v^{(j+l)}.$$

Since  $v_s$  is a finite sum of the  $v^{(j)}$  in each case, the  $L^2(M)$  bounds for  $v_s$  and  $(-\Delta - s^2)v_s$  and the  $L^2(\partial M)$  bounds for  $v_s$  follow from corresponding bounds for the  $v^{(j)}$ . The form of  $v_s$  near points where  $\gamma$  does not self-intersect and the possibility to prescribe the  $K$ -th order jets of  $\Theta|_{\partial M}$  and  $a|_{\partial M}$  at  $\gamma(0)$  follow from the construction and [Lemma 7.4](#).

To conclude the proof, using a partition of unity, it is enough to verify the limit (7-1) for any  $\psi$  supported in one of the sets  $V_r \cap M$  or  $W_{j,l} \cap M$ . Further, we can choose the sets  $V_r$  to be so small that the real part of  $d\Theta^{(j)} - d\Theta^{(k)}$  is nonvanishing near  $V_r$  if  $\gamma(t_j) = \gamma(t_k) = p_r$  but  $j \neq k$ . This follows since

$$\text{Re}(d\Theta^{(j)} - d\Theta^{(k)})(p_r) = \dot{\gamma}(t_j)^b - \dot{\gamma}(t_k)^b \neq 0.$$

Here we may need to decrease  $\delta$  so that we still have an open cover.

Consider first the case where  $\psi \in C_c(W_{j,l} \cap M)$ . Here the support of  $\psi$  may reach  $\partial M$ , and we extend  $\psi$  by zero outside of  $W_{j,l} \cap M$ . Suppose that  $v_s = e^{is\Theta} a$ , where  $\Theta = t + \xi_\alpha y^\alpha + \frac{1}{2}H(t)y \cdot y + O(|y|^3)$  and  $a = \tau^{(m-1)/4}(a_0 + O(\tau^{-1}))\chi(y/\delta')$ , and let  $\rho = |g|^{1/2}$ . Then

$$\begin{aligned} & \int_M |v_{\tau+i\lambda}|^2 \psi \, dV_g \\ &= \int_{-\infty}^{\infty} \int_{\mathbb{R}^{m-1}} e^{-2\lambda t} e^{-\tau \text{Im}(H(t))y \cdot y} e^{\tau O(|y|^3)} e^{O(|y|)} \tau^{(m-1)/2} (|a_0|^2 + O(\tau^{-1})) \chi(y/\delta')^2 \psi \rho \, dt \, dy \\ &= \int_{-\infty}^{\infty} e^{-2\lambda t} \int_{\mathbb{R}^{m-1}} e^{-\text{Im}(H(t))y \cdot y} e^{\tau^{-1/2} O(|y|^3)} e^{\tau^{-1/2} O(|y|)} (|a_0(t, \tau^{-1/2}y)|^2 \\ & \quad + O(\tau^{-1})) \chi(y/\tau^{1/2}\delta')^2 \psi(t, \tau^{-1/2}y) \rho(t, \tau^{-1/2}y) \, dt \, dy. \end{aligned}$$

Since  $\text{Im}(H(t))$  is positive definite and  $\delta'$  is sufficiently small, the term  $e^{-\text{Im}(H(t))y \cdot y}$  dominates the other exponentials and one obtains

$$\lim_{\tau \rightarrow \infty} \int_M |v_{\tau+i\lambda}|^2 \psi \, dV_g = \int_0^L e^{-2\lambda t} \left( \int_{\mathbb{R}^{m-1}} e^{-\text{Im}(H(t))y \cdot y} \, dy \right) |a_0(t, 0)|^2 \psi(t, 0) \rho(t, 0) \, dt.$$

Evaluating the integral over  $y$  gives

$$\lim_{\tau \rightarrow \infty} \int_M |v_{\tau+i\lambda}|^2 \psi \, dV_g = C_m \int_0^L e^{-2\lambda t} \frac{|a_0(t, 0)|^2 \rho(t, 0)}{\sqrt{\det \text{Im}(H(t))}} \psi(t, 0) \, dt.$$

We will prove below that

$$\frac{|a_0(t, 0)|^2 \rho(t, 0)}{\sqrt{\det \text{Im}(H(t))}} = \text{const.} \tag{7-5}$$

The limit (7-1) will follow upon dividing the family  $\{v_s\}$  by a suitable constant.

If  $\psi \in C_c(V_r \cap M)$  (again  $\text{supp}(\psi)$  may extend up to  $\partial M$ ), we have

$$v_s|_{V_r} = \sum_{\gamma(t_j)=p_r} v^{(j)},$$

so that on  $V_r$

$$|v_s|^2 = \sum_{\gamma(t_j)=p_r} |v^{(j)}|^2 + \sum_{\substack{\gamma(t_j)=\gamma(t_k)=p_r \\ j \neq k}} v^{(j)} \overline{v^{(k)}}.$$

We arranged earlier for  $\text{Re}(d\Theta^{(j)} - d\Theta^{(k)})$  to be nonvanishing near  $V_r$  if  $\gamma(t_j) = \gamma(t_k) = p_r$  but  $j \neq k$ . Thus the cross terms give rise to terms of the form

$$\int_{V_r \cap M} v^{(j)} \overline{v^{(k)}} \psi \, dV = \int_{V_r \cap M} e^{i\tau\varphi} w^{(j)} \overline{w^{(k)}} \psi \, dV,$$

where  $\varphi = \text{Re}(\Theta^{(j)} - \Theta^{(k)})$  has nonvanishing gradient in  $V_r$ , and  $w^{(l)} = e^{is \text{Im}(\Theta^{(l)})} e^{-\lambda \text{Re}(\Phi^{(l)})} a^{(l)}$ . We wish to prove that

$$\lim_{\tau \rightarrow \infty} \int_{V_r \cap M} e^{i\tau\varphi} w^{(j)} \overline{w^{(k)}} \psi \, dV = 0, \quad j \neq k, \tag{7-6}$$

showing that the cross terms vanish in the limit and the previous computation for  $|v^{(l)}|^2$  shows the limit (7-1) also when  $\psi$  is supported in some  $V_r \cap M$ . To show (7-6), let  $\varepsilon > 0$  and decompose  $\psi = \psi_1 + \psi_2$ , where  $\psi_1 \in C_c^\infty(V_r \cap M)$  and  $\|\psi_2\|_{L^\infty(V_r \cap M)} \leq \varepsilon$ . Then

$$\left| \int_{V_r \cap M} e^{i\tau\varphi} w^{(j)} \overline{w^{(k)}} \psi_2 \, dV \right| \lesssim \|w^{(j)}\|_{L^2} \|w^{(k)}\|_{L^2} \|\psi_2\|_{L^\infty} \lesssim \varepsilon,$$

since  $\|w^{(l)}\|_{L^2} \lesssim \|v^{(l)}\|_{L^2} \lesssim 1$ . For the smooth part  $\psi_1$ , we employ a nonstationary phase argument and integrate by parts using that

$$e^{i\tau\varphi} = \frac{1}{i\tau} L(e^{i\tau\varphi}), \quad Lw = \langle |d\varphi|^{-2} d\varphi, dw \rangle.$$



This gives

$$\int_{V_r \cap M} e^{i\tau\varphi} w^{(j)} \overline{w^{(k)}} \psi_1 dV = \int_{\partial M} \frac{\partial_v \varphi}{i\tau |d\varphi|^2} v^{(j)} \overline{v^{(k)}} \psi_1 dS + \frac{1}{i\tau} \int_{V_r \cap M} e^{i\tau\varphi} L^t(w^{(j)} \overline{w^{(k)}}) \psi_1 dV.$$

Since  $\|v^{(l)}\|_{L^2(\partial M)} = O(1)$ , the boundary term can be made arbitrarily small as  $\tau \rightarrow \infty$ . As for the last term, the worst behavior is when the transpose  $L^t$  acts on  $e^{is \operatorname{Im}(\Theta^{(l)})}$ , and these terms have bounds of the form

$$\| |d(\operatorname{Im}(\Theta^{(j)}))| v^{(j)} \|_{L^2} \|v^{(k)}\|_{L^2} \|\psi_1\|_{L^\infty}.$$

Here  $|d(\operatorname{Im}(\Theta^{(j)}))| \lesssim |y|$  if  $(t, y)$  are coordinates along the geodesic segment corresponding to  $v^{(j)}$ , and the computation above for  $\|v^{(j)}\|_{L^2}$  shows that

$$\| |d(\operatorname{Im}(\Theta^{(j)}))| v^{(j)} \|_{L^2} \|v^{(k)}\|_{L^2} \|\psi_1\|_{L^\infty} \lesssim \tau^{-1/2}.$$

This finishes the proof of (7-6) and also of (7-1).

It remains to show (7-5). We have

$$|a_0(t, 0)|^2 \rho(t, 0) = |c_0|^2 e^{-\int_{t_0}^t \operatorname{Re}(\eta_0)(s) ds} |g(t, 0)|^{1/2}.$$

Note that  $\eta_0(t)$  is given by

$$\begin{aligned} \eta_0(t) &= \Delta \Theta(t, 0) \\ &= (g^{jk} \partial_{jk} \Theta + \partial_j g^{jk} \partial_k \Theta + |g|^{-1/2} \partial_j (|g|^{1/2}) g^{jk} \partial_k \Theta)(t, 0) \\ &= g^{11} \partial_t^2 \Theta_0 + 2g^{1\alpha} \partial_{t\alpha} \Theta_1 + g^{\alpha\beta} \partial_{\alpha\beta} \Theta_2 + \partial_j g^{j1} \partial_t \Theta_0 + \partial_j g^{j\alpha} \partial_\alpha \Theta_1 + \frac{1}{2} \partial_j (\log |g|) (g^{j1} \partial_t \Theta_0 + g^{j\alpha} \partial_\alpha \Theta_1) \\ &= 2g^{1\alpha} \dot{\xi}_\alpha + g^{\alpha\beta} H_{\alpha\beta} + (\partial_j g^{jk}) \xi_k + \frac{1}{2} \partial_j (\log |g|) g^{jk} \xi_k. \end{aligned}$$

The conditions  $g^{jk} \xi_k = \delta_1^j$  and  $g^{1\alpha} \dot{\xi}_\alpha = g^{1k} \dot{\xi}_k = -(\partial_t g^{1k}) \xi_k$  at  $(t, 0)$ , together with the general fact that  $\partial_t (\log |g|) = -g_{jk} \partial_t g^{jk}$ , imply that

$$\eta_0(t) = g^{1\alpha} \dot{\xi}_\alpha + g^{\alpha\beta} H_{\alpha\beta} + (\partial_\alpha g^{\alpha k}) \xi_k + \frac{1}{2} \partial_t (\log |g|).$$

Recalling the definition of the  $B$  and  $C$  matrices in (7-4), this says precisely that

$$\begin{aligned} \eta_0(t) &= B_\alpha^\alpha + C^{\alpha\gamma} H_{\gamma\alpha} + \frac{1}{2} \partial_t (\log |g|) \\ &= \operatorname{tr}(B(t) + C(t)H(t)) + \frac{1}{2} \partial_t (\log |g|). \end{aligned}$$

Consequently,  $|a_0(t, 0)|^2 \rho(t, 0) = c_0' e^{-\int_{t_0}^t \operatorname{tr}(B(s)+C(s) \operatorname{Re}(H(s))) ds}$ . On the other hand, by [Katchalov et al. 2001, Lemma 2.58], solutions of the matrix Riccati equation have the property that

$$\det \operatorname{Im}(H(t)) = \det \operatorname{Im}(H(t_0)) e^{-2 \int_{t_0}^t \operatorname{tr}(B(s)+C(s) \operatorname{Re}(H(s))) ds}.$$

This proves the result. □

The proof of Proposition 7.1 now follows quickly from the way we have set up the previous result.

*Proof of Proposition 7.1.* Let  $\gamma : [0, L] \rightarrow M$  be a nontangential broken ray with endpoints on  $E$ , and let  $0 < t_1 < \dots < t_N < L$  be the times of reflection. Let  $v_s^{(0)}$  be a Gaussian beam quasimode as in Proposition 7.5 associated with the geodesic  $\gamma|_{[0, t_1]}$ . We will construct another Gaussian beam quasimode  $v_s^{(1)}$  associated with  $\gamma|_{[t_1, t_2]}$  such that  $v_s^{(0)} - v_s^{(1)}|_{\partial M}$  will be small near  $\gamma(t_1)$ .

In fact, by Proposition 7.5 we have  $v_s^{(j)} = e^{is\Theta^{(j)}} a^{(j)}$  near  $\gamma(t_1)$ , and we can choose the  $K$ -th order jet of  $\Theta^{(1)}|_{\partial M}$  at  $\gamma(t_1)$  to be equal to that of  $\Theta^{(0)}|_{\partial M}$  with the following exception: we always have

$$\begin{aligned} d(\Theta^{(0)})|_{\gamma(t_1)} &= \dot{\gamma}(t_1-)^b, \\ d(\Theta^{(1)})|_{\gamma(t_1)} &= \dot{\gamma}(t_1+)^b. \end{aligned}$$

It follows that

$$\begin{aligned} d(\Theta^{(0)}|_{\partial M})|_{\gamma(t_1)} &= \dot{\gamma}(t_1-)^b_{\text{tan}}, \\ d(\Theta^{(1)}|_{\partial M})|_{\gamma(t_1)} &= \dot{\gamma}(t_1+)^b_{\text{tan}}, \end{aligned}$$

where we have taken the projections to the cotangent space of  $\partial M$  at  $\gamma(t_1)$ . But by the rule that the angle of incidence equals angle of reflection,  $\dot{\gamma}(t_1-)^b_{\text{tan}}$  equals  $\dot{\gamma}(t_1+)^b_{\text{tan}}$ . Thus the  $K$ -th order jets of  $\Theta^{(0)}|_{\partial M}$  and  $\Theta^{(1)}|_{\partial M}$  actually coincide at  $\gamma(t_1)$ , and by Proposition 7.5 we can also arrange for the  $K$ -th order jets of  $a^{(0)}|_{\partial M}$  and  $a^{(1)}|_{\partial M}$  to coincide at  $\gamma(t_1)$ .

Write  $f_s = v_s^{(0)} - v_s^{(1)}|_{\partial M}$ , and let  $(t, y)$  be coordinates near  $\gamma(t_1)$  such that  $\partial M$  is parametrized by  $y \mapsto (t_1, y)$  and  $\gamma(t_1)$  corresponds to  $(t_1, 0)$ . Recall that  $v_s^{(j)}$  are supported in small tubular neighborhoods of the corresponding geodesic segments. By the above considerations and the construction of  $\Theta^{(j)}$  and  $a^{(j)}$ , and dropping the variable  $t_1$  from the notations, the restrictions of  $\Theta^{(j)}$  and  $a^{(j)}$  to  $\partial M$  satisfy

$$\Theta^{(j)}(y) = \Theta(y) + \Xi^{(j)}(y), \quad a^{(j)}(y) = a(y) + b^{(j)}(y),$$

where  $\Theta$  is a polynomial of order  $K$ ,  $a = \tau^{(m-1)/4} \tilde{a} \chi(y/\delta')$ , where  $\tilde{a}$  is a polynomial of order  $K$ , and  $|\Xi^{(j)}(y)| \leq C|y|^{K+1}$  and  $|b^{(j)}(y)| \leq C\tau^{(m-1)/4}|y|^{K+1} \chi(y/\delta')$  on  $\text{supp}(\chi(\cdot/\delta'))$ , where  $\chi$  is a cutoff function and  $\delta'$  is a constant independent of  $\tau$  that can be chosen as small as we want (these initially depend on  $j$ , but since there are finitely many reflections, we can choose them independently of  $j$ ). Here  $\Theta$  and  $\Xi^{(j)}$  are independent of  $\tau$ , and  $a$  and  $b^{(j)}$  are mildly  $\tau$ -dependent and satisfy uniform bounds with respect to  $\tau$ . Then

$$f_s = e^{is\Theta} ((e^{is\Xi^{(0)}} - e^{is\Xi^{(1)}})a + e^{is\Xi^{(0)}}b^{(0)} - e^{is\Xi^{(1)}}b^{(1)}).$$

We have

$$e^{is\Xi^{(0)}} - e^{is\Xi^{(1)}} = is(\Xi^{(0)} - \Xi^{(1)}) \int_0^1 e^{is(r\Xi^{(0)} + (1-r)\Xi^{(1)})} dr$$

and, consequently, near  $y = 0$ ,

$$|e^{is\Xi^{(0)}} - e^{is\Xi^{(1)}}| \leq C\tau|y|^{K+1} e^{C\tau|y|^{K+1}}.$$

Thus, near  $y = 0$ ,

$$|f_s(y)| \leq C\tau^{(m-1)/4} e^{-\tau \text{Im}(\Theta)} \tau|y|^{K+1} e^{C\tau|y|^{K+1}} \chi(y/\delta').$$

Using that the Hessian of  $\text{Im}(\Theta)$  at 0 is positive definite and choosing  $\delta'$  sufficiently small, we have

$$|f_s(y)| \leq C\tau^{(m-1)/4} e^{-c\tau|y|^2} \tau|y|^{K+1} \chi(y/\delta').$$

Integrating the square of  $|f_s|$  over  $\mathbb{R}^{m-1}$  and changing  $y$  to  $\tau^{-1/2}y$ , we obtain

$$\|f_s\|_{L^2(R_1)} = O(\tau^{-(K-1)/2}),$$

where  $R_1$  is a small neighborhood of  $\gamma(t_1)$  on  $\partial M$  containing the set of interest.

Repeating this construction for the other points of reflection, we end up with a quasimode

$$v_s = \sum_{j=0}^N (-1)^j v_s^{(j)}$$

that is supported in a small neighborhood of the broken ray  $\gamma$ . Since all points of reflection are distinct, we can arrange that the quasimode satisfies

$$\|v_s|_R\|_{L^2(R)} = O(\tau^{-(K-1)/2}).$$

It also satisfies

$$\|(-\Delta - s^2)v_s\|_{L^2(M)} = O(\tau^{-K}), \quad \|v_s\|_{L^2(M)} = O(1).$$

Replacing  $K$  by  $2K + 1$ , we have proved all the other statements in the proposition except for the expression of the limit measure. To do this, we consider the finitely many points where the full broken ray  $\gamma$  self-intersects or reflects, and decompose the terms  $v_s^{(j)}$  as in the proof of Proposition 7.5 to parts living in small neighborhoods of the self-intersection and reflection points and parts away from these points. Now all self-intersection points are in the interior or on  $E$  and all self-intersections must be transversal, and also all reflections are transversal. Consequently, when forming  $|v_s|^2$ , the cross terms arising from different parts living near the same self-intersection or reflection point contribute an  $o(1)$  term by nonstationary phase as in the proof of Proposition 7.5. Thus the limit measure of  $|v_s|^2 dV_g$  is indeed the measure  $e^{-2\lambda t} \delta_\gamma$ , where  $\delta_\gamma$  is the delta function of the broken ray  $\gamma$ .  $\square$

### 8. Recovering the broken ray transform

In this section we give the proof of Theorem 2.4 concerning the recovery of integrals over broken rays.

*Proof of Theorem 2.4.* The proof is very similar to the proof of Theorem 2.1, except that we use reflected Gaussian beam quasimodes instead of WKB type quasimodes. Let  $\gamma : [0, L] \rightarrow M_0$  be a nontangential broken ray with endpoints on  $E$ , and let  $\lambda > 0$ . Also let  $\tilde{g} = e \oplus g_0$  and  $\tilde{q}_j = c(q_j - c^{(n-2)/4} \Delta_g(c^{-(n-2)/4}))$ . Consider the complex frequency

$$s = \tau + i\lambda,$$

where  $\tau > 0$  will be large. We look for solutions

$$\begin{aligned} \tilde{u}_1 &= e^{-sx_1} (v_s(x') + r_1), \\ \tilde{u}_2 &= e^{sx_1} (v_s(x') + r_2) \end{aligned}$$

of the equations  $(-\Delta_{\tilde{g}} + \tilde{q}_1)\tilde{u}_1 = 0$ ,  $(-\Delta_{\tilde{g}} + \tilde{q}_2)\tilde{u}_2 = 0$  in  $M$ . Here  $v_s \in C^\infty(M_0)$  is the quasimode constructed in Proposition 7.1 that concentrates near the given broken ray  $\gamma$  and is small on  $\partial M_0 \setminus E$ .

Since  $\Delta_{\tilde{g}} = \partial_1^2 + \Delta_{g_0}$ , the function  $\tilde{u}_1$  is a solution if and only if

$$e^{sx_1}(-\Delta_{\tilde{g}} + \tilde{q}_1)(e^{-sx_1}r_1) = -(-\Delta_{g_0} + \tilde{q}_1 - s^2)v_s(x') \quad \text{in } M.$$

We look for a solution in the form  $r_1 = e^{i\lambda x_1}r'_1$  where  $r'_1$  satisfies

$$e^{\tau x_1}(-\Delta_{\tilde{g}} + \tilde{q}_1)(e^{-\tau x_1}r'_1) = f \quad \text{in } M$$

with

$$f = -e^{-i\lambda x_1}(-\Delta_{g_0} + \tilde{q}_1 - s^2)v_s(x').$$

To arrange that  $\tilde{u}_1|_{\Gamma_i} = 0$ , fix some small  $\delta > 0$ , let  $S_\pm$  and  $S_0$  be the sets in Proposition 4.3 with Carleman weight  $\varphi(x) = -x_1$ , and consider the boundary condition

$$e^{\tau\varphi}r'_1|_{S_- \cup S_0} = e^{\tau\varphi}f_-,$$

where

$$f_- = \begin{cases} -e^{-i\lambda x_1}v_s(x') & \text{on } \Gamma_i, \\ 0 & \text{on } (S_- \cup S_0) \setminus \Gamma_i. \end{cases}$$

For any fixed  $K > 0$ , by Proposition 7.1 and by the condition that  $\Gamma_i \subset \mathbb{R} \times (\partial M_0 \setminus E)$ , we may assume that the following bounds are valid:

$$\|f\|_{L^2(M)} = O(1), \quad \|f_-\|_{L^2(S_-)} = 0, \quad \|f_-\|_{L^2(S_0)} = O(\tau^{-K}).$$

It follows from Proposition 4.3 that there is a solution  $r'_1$  satisfying the above boundary condition and having the estimate

$$\|r'_1\|_{L^2(M)} = O(\tau^{-1}).$$

Choosing  $r'_1$  as described above and choosing  $r_1 = e^{i\lambda x_1}r'_1$ , we have produced a solution  $\tilde{u}_1 \in H_{\Delta_{\tilde{g}}}(M)$  of the equation  $(-\Delta_{\tilde{g}} + \tilde{q}_1)\tilde{u}_1 = 0$  in  $M$ , having the form

$$\tilde{u}_1 = e^{-sx_1}(v_s(x') + r_1)$$

and satisfying

$$\text{supp}(\tilde{u}_1|_{\partial M}) \subset \overline{\partial M_+ \cup \partial M_- \cup \Gamma_a}$$

and  $\|r_1\|_{L^2(M)} = O(\tau^{-1})$  as  $\tau \rightarrow \infty$ . Repeating this construction for the Carleman weight  $\varphi(x) = x_1$ , we obtain a solution  $\tilde{u}_2 \in H_{\Delta_{\tilde{g}}}(M)$  of the equation  $(-\Delta_{\tilde{g}} + \tilde{q}_2)\tilde{u}_2 = 0$  in  $M$ , having the form

$$\tilde{u}_2 = e^{sx_1}(v_s(x') + r_2)$$

satisfying the same support condition and bound for  $\|r_2\|_{L^2(M)}$ .

Writing  $u_j = c^{-(n-2)/4}\tilde{u}_j$ , Lemma 6.2 shows that  $u_j \in H_{\Delta_g}(M)$  are solutions of  $(-\Delta_g + q_1)u_1 = 0$  and  $(-\Delta_g + \bar{q}_2)u_2 = 0$  in  $M$ . Then Proposition 6.1 implies that

$$\int_M (q_1 - q_2)u_1\bar{u}_2 dV_g = 0.$$

We extend  $q_1 - q_2$  by zero to  $\mathbb{R} \times M_0$ . Inserting the expressions for  $u_j$  and using the equality  $dV_g = c^{n/2} dx_1 dV_{g_0}(x')$ , we obtain

$$\int_{M_0} \int_{-\infty}^{\infty} (q_1 - q_2) c e^{-2i\lambda x_1} (|v_s(x')|^2 + v_s \bar{r}_2 + \bar{v}_s r_1 + r_1 \bar{r}_2) dx_1 dV_{g_0}(x') = 0.$$

Since  $\|r_j\|_{L^2(M)} = O(\tau^{-1})$  as  $\tau \rightarrow \infty$ , Proposition 7.1 implies that

$$\int_0^L e^{-2\lambda t} (c(q_1 - q_2))^{\wedge}(2\lambda, \gamma(t)) dt = 0.$$

This concludes the proof. □

### Acknowledgements

Kenig is partly supported by the NSF. Salo is supported in part by the Academy of Finland and an ERC Starting Grant. Salo thanks David dos Santos Ferreira, Yaroslav Kurylev, and Matti Lassas for several helpful discussions and for allowing the authors to use arguments from [Ferreira 2013], which was in preparation simultaneously with this manuscript and involved Gaussian beam quasimodes and a reduction from the attenuated ray transform to the usual ray transform. Salo also thanks the Department of Mathematics of the University of Chicago, where part of this work was carried out.

### References

- [Albin et al. 2013] P. Albin, C. Guillarmou, L. Tzou, and G. Uhlmann, “Inverse boundary problems for systems in two dimensions”, *Ann. Henri Poincaré* **14**:6 (2013), 1551–1571. MR 3085925 Zbl 06195029
- [Alessandrini and Kim 2012] G. Alessandrini and K. Kim, “Single-logarithmic stability for the Calderón problem with local data”, *J. Inverse Ill-Posed Probl.* **20**:4 (2012), 389–400. MR 2984474
- [Ammari and Uhlmann 2004] H. Ammari and G. Uhlmann, “Reconstruction of the potential from partial Cauchy data for the Schrödinger equation”, *Indiana Univ. Math. J.* **53**:1 (2004), 169–183. MR 2005h:35351 Zbl 1051.35103
- [Astala and Päivärinta 2006] K. Astala and L. Päivärinta, “Calderón’s inverse conductivity problem in the plane”, *Ann. of Math.* (2) **163**:1 (2006), 265–299. MR 2007b:30019 Zbl 1111.35004
- [Astala et al. 2005] K. Astala, L. Päivärinta, and M. Lassas, “Calderón’s inverse problem for anisotropic conductivity in the plane”, *Comm. Partial Differential Equations* **30**:1-3 (2005), 207–224. MR 2005k:35421 Zbl 1129.35483
- [Bukhgeim 2008] A. L. Bukhgeim, “Recovering a potential from Cauchy data in the two-dimensional case”, *J. Inverse Ill-Posed Probl.* **16**:1 (2008), 19–33. MR 2008m:30049 Zbl 1142.30018
- [Bukhgeim and Uhlmann 2002] A. L. Bukhgeim and G. Uhlmann, “Recovering a potential from partial Cauchy data”, *Comm. Partial Differential Equations* **27**:3-4 (2002), 653–668. MR 2003d:35262 Zbl 0998.35063
- [Caro et al. 2009] P. Caro, P. Ola, and M. Salo, “Inverse boundary value problem for Maxwell equations with local data”, *Comm. Partial Differential Equations* **34**:10-12 (2009), 1425–1464. MR 2010m:35558 Zbl 1185.35321
- [Chung 2012] F. J.-H. Chung, *A partial data result for the magnetic Schrödinger inverse problem*, Ph.D. Thesis, The University of Chicago, 2012, Available at <http://search.proquest.com/docview/1032548281>. MR 3054984
- [Chung et al. 2013] F. Chung, M. Salo, and L. Tzou, “Partial data inverse problems for the Hodge Laplacian”, preprint, 2013. arXiv 1310.4616
- [Eskin 2004] G. Eskin, “Inverse boundary value problems in domains with several obstacles”, *Inverse Problems* **20**:5 (2004), 1497–1516. MR 2005h:35358 Zbl 1074.35086

- [Ferreira 2007] D. dos Santos Ferreira, C. E. Kenig, J. Sjöstrand, and G. Uhlmann, “Determining a magnetic Schrödinger operator from partial Cauchy data”, *Comm. Math. Phys.* **271**:2 (2007), 467–488. MR 2008a:35044 Zbl 1148.35096
- [Ferreira 2009a] D. dos Santos Ferreira, C. E. Kenig, M. Salo, and G. Uhlmann, “Limiting Carleman weights and anisotropic inverse problems”, *Invent. Math.* **178**:1 (2009), 119–171. MR 2010h:58033 Zbl 1181.35327
- [Ferreira 2009b] D. dos Santos Ferreira, C. E. Kenig, J. Sjöstrand, and G. Uhlmann, “On the linearized local Calderón problem”, *Math. Res. Lett.* **16**:6 (2009), 955–970. MR 2011d:35515 Zbl 1198.31003
- [Ferreira 2013] D. dos Santos Ferreira, Y. Kurylev, M. Lassas, and M. Salo, “The Calderon problem in transversally anisotropic geometries”, preprint, 2013. arXiv 1305.1273
- [Florescu et al. 2011] L. Florescu, V. A. Markel, and J. C. Schotland, “Inversion formulas for the broken-ray Radon transform”, *Inverse Problems* **27**:2 (2011), 025002, 13. MR 2011h:45019 Zbl 1211.65168
- [Guillarmou and Tzou 2011a] C. Guillarmou and L. Tzou, “Calderón inverse problem with partial data on Riemann surfaces”, *Duke Math. J.* **158**:1 (2011), 83–120. MR 2012f:35574 Zbl 1222.35212
- [Guillarmou and Tzou 2011b] C. Guillarmou and L. Tzou, “Identification of a connection from Cauchy data on a Riemann surface with boundary”, *Geom. Funct. Anal.* **21**:2 (2011), 393–418. MR 2012k:58027 Zbl 1260.58011
- [Guillarmou and Tzou 2013] C. Guillarmou and L. Tzou, “The Calderón inverse problem in two dimensions”, pp. 119–166 in *Inverse problems and applications: inside out II*, edited by G. Uhlmann, Cambridge University Press, 2013.
- [Haberman and Tataru 2013] B. Haberman and D. Tataru, “Uniqueness in Calderón’s problem with Lipschitz conductivities”, *Duke Math. J.* **162**:3 (2013), 496–516. MR 3024091 Zbl 1260.35251
- [Heck and Wang 2006] H. Heck and J.-N. Wang, “Stability estimates for the inverse boundary value problem by partial Cauchy data”, *Inverse Problems* **22**:5 (2006), 1787–1796. MR 2007g:35270 Zbl 1106.35133
- [Helgason 1999] S. Helgason, *The Radon transform*, 2nd ed., Progress in Mathematics **5**, Birkhäuser, Boston, MA, 1999. MR 2000m:44003 Zbl 0932.43011
- [Hubenthal 2013a] M. Hubenthal, “The broken ray transform on the square”, preprint, 2013. arXiv 1302.6193
- [Hubenthal 2013b] M. Hubenthal, “The broken ray transform in  $n$  dimensions”, preprint, 2013. arXiv 1310.7156
- [Hyvönen et al. 2012] N. Hyvönen, P. Piiroinen, and O. Seiskari, “Point measurements for a Neumann-to-Dirichlet map and the Calderón problem in the plane”, *SIAM J. Math. Anal.* **44**:5 (2012), 3526–3536. MR 3023421 Zbl 1257.35197
- [Ilmavirta 2013a] J. Ilmavirta, “Broken ray tomography in the disc”, *Inverse Problems* **29**:3 (2013), 035008, 17. MR 3040563 Zbl 06158616
- [Ilmavirta 2013b] J. Ilmavirta, “A reflection approach to the broken ray transform”, preprint, 2013. arXiv 1306.0341
- [Ilmavirta 2013c] J. Ilmavirta, “Boundary reconstruction for the broken ray transform”, preprint, 2013. arXiv 1310.2025
- [Imanuvilov and Yamamoto 2012a] O. Y. Imanuvilov and M. Yamamoto, “Inverse boundary value problem for Schrödinger equation in two dimensions”, *SIAM J. Math. Anal.* **44**:3 (2012), 1333–1339. MR 2982714 Zbl 1273.35316
- [Imanuvilov and Yamamoto 2012b] O. Y. Imanuvilov and M. Yamamoto, “Inverse problem by Cauchy data on an arbitrary sub-boundary for systems of elliptic equations”, *Inverse Problems* **28**:9 (2012), 095015, 30. MR 2972464 Zbl 1250.35184
- [Imanuvilov and Yamamoto 2013a] O. Y. Imanuvilov and M. Yamamoto, “Inverse boundary value problem for the Schrödinger equation in a cylindrical domain by partial boundary data”, *Inverse Problems* **29**:4 (2013), 045002, 8. MR 3041540 Zbl 1273.35317
- [Imanuvilov and Yamamoto 2013b] O. Y. Imanuvilov and M. Yamamoto, “Uniqueness for Inverse Boundary Value Problems by Dirichlet-to-Neumann Map on Subboundaries”, *Milan J. Math.* **81**:2 (2013), 187–258. MR 3129784 Zbl 06236315
- [Imanuvilov et al. 2010] O. Y. Imanuvilov, G. Uhlmann, and M. Yamamoto, “The Calderón problem with partial data in two dimensions”, *J. Amer. Math. Soc.* **23**:3 (2010), 655–691. MR 2012c:35472 Zbl 1201.35183
- [Imanuvilov et al. 2011a] O. Y. Imanuvilov, G. Uhlmann, and M. Yamamoto, “Determination of second-order elliptic operators in two dimensions from partial Cauchy data”, *Proc. Natl. Acad. Sci. USA* **108**:2 (2011), 467–472. MR 2012a:35364 Zbl 1256.35203
- [Imanuvilov et al. 2011b] O. Y. Imanuvilov, G. Uhlmann, and M. Yamamoto, “Inverse boundary value problem by measuring Dirichlet data and Neumann data on disjoint sets”, *Inverse Problems* **27**:8 (2011), 085007, 26. MR 2012c:78002 Zbl 1222.35213

- [Isakov 2007] V. Isakov, “On uniqueness in the inverse conductivity problem with local data”, *Inverse Probl. Imaging* **1**:1 (2007), 95–105. [MR 2007m:35273](#) [Zbl 1125.35113](#)
- [Katchalov et al. 2001] A. Katchalov, Y. Kurylev, and M. Lassas, *Inverse boundary spectral problems*, Monographs and Surveys in Pure and Applied Mathematics **123**, Chapman & Hall/CRC, Boca Raton, FL, 2001. [MR 2003e:58045](#) [Zbl 1037.35098](#)
- [Kenig and Salo 2013] C. Kenig and M. Salo, “Recent progress in the Calderón problem with partial data”, preprint, 2013. [arXiv 1302.4218](#)
- [Kenig et al. 2007] C. E. Kenig, J. Sjöstrand, and G. Uhlmann, “The Calderón problem with partial data”, *Ann. of Math. (2)* **165**:2 (2007), 567–591. [MR 2008k:35498](#) [Zbl 1127.35079](#)
- [Kenig et al. 2011a] C. E. Kenig, M. Salo, and G. Uhlmann, “Inverse problems for the anisotropic Maxwell equations”, *Duke Math. J.* **157**:2 (2011), 369–419. [MR 2012d:35408](#) [Zbl 1226.35086](#)
- [Kenig et al. 2011b] C. E. Kenig, M. Salo, and G. Uhlmann, “Reconstructions from boundary measurements on admissible manifolds”, *Inverse Probl. Imaging* **5**:4 (2011), 859–877. [MR 2012k:58038](#) [Zbl 1237.35160](#)
- [Knudsen 2006] K. Knudsen, “The Calderón problem with partial data for less smooth conductivities”, *Comm. Partial Differential Equations* **31**:1-3 (2006), 57–71. [MR 2006k:35303](#) [Zbl 1091.35116](#)
- [Knudsen and Salo 2007] K. Knudsen and M. Salo, “Determining nonsmooth first order terms from partial boundary measurements”, *Inverse Probl. Imaging* **1**:2 (2007), 349–369. [MR 2008k:35500](#) [Zbl 1122.35152](#)
- [Kohn and Vogelius 1984] R. Kohn and M. Vogelius, “Determining conductivity by boundary measurements”, *Comm. Pure Appl. Math.* **37**:3 (1984), 289–298. [MR 85f:80008](#) [Zbl 0595.35092](#)
- [Krishnan 2009] V. P. Krishnan, “A support theorem for the geodesic ray transform on functions”, *J. Fourier Anal. Appl.* **15**:4 (2009), 515–520. [MR 2010i:53146](#) [Zbl 1186.53089](#)
- [Krupchyk et al. 2012] K. Krupchyk, M. Lassas, and G. Uhlmann, “Inverse problems with partial data for a magnetic Schrödinger operator in an infinite slab and on a bounded domain”, *Comm. Math. Phys.* **312**:1 (2012), 87–126. [MR 2914058](#) [Zbl 1238.35188](#)
- [Lassas and Oksanen 2010] M. Lassas and L. Oksanen, “An inverse problem for a wave equation with sources and observations on disjoint sets”, *Inverse Problems* **26**:8 (2010), 085012, 19. [MR 2011m:35422](#) [Zbl 1197.35323](#)
- [Lassas and Oksanen 2012] M. Lassas and L. Oksanen, “Inverse problem for the Riemannian wave equation with Dirichlet data and Neumann data on disjoint sets”, preprint, 2012. [arXiv 1208.2105](#)
- [Li and Uhlmann 2010] X. Li and G. Uhlmann, “Inverse problems with partial data in a slab”, *Inverse Probl. Imaging* **4**:3 (2010), 449–462. [MR 2011f:35361](#) [Zbl 1200.35331](#)
- [Lozev 2013] K. Lozev, “Mathematical modeling and solution of the tomography problem in domains with reflecting obstacles”, *J. Phys.: Conf. Ser.* **410** (2013).
- [Nachman 1996] A. I. Nachman, “Global uniqueness for a two-dimensional inverse boundary value problem”, *Ann. of Math. (2)* **143**:1 (1996), 71–96. [MR 96k:35189](#) [Zbl 0857.35135](#)
- [Nachman and Street 2010] A. Nachman and B. Street, “Reconstruction in the Calderón problem with partial data”, *Comm. Partial Differential Equations* **35**:2 (2010), 375–390. [MR 2012b:35368](#) [Zbl 1186.35242](#)
- [Salo and Tzou 2010] M. Salo and L. Tzou, “Inverse problems with partial data for a Dirac system: a Carleman estimate approach”, *Adv. Math.* **225**:1 (2010), 487–513. [MR 2011g:35432](#) [Zbl 1197.35329](#)
- [Sharafutdinov 1994] V. A. Sharafutdinov, *Integral geometry of tensor fields*, VSP, Utrecht, 1994. [MR 97h:53077](#)
- [Sylvester and Uhlmann 1987] J. Sylvester and G. Uhlmann, “A global uniqueness theorem for an inverse boundary value problem”, *Ann. of Math. (2)* **125**:1 (1987), 153–169. [MR 88b:35205](#) [Zbl 0625.35078](#)
- [Sylvester and Uhlmann 1988] J. Sylvester and G. Uhlmann, “Inverse boundary value problems at the boundary—continuous dependence”, *Comm. Pure Appl. Math.* **41**:2 (1988), 197–219. [MR 89f:35213](#) [Zbl 0632.35074](#)
- [Uhlmann and Vasy 2012] G. Uhlmann and A. Vasy, “The inverse problem for the local geodesic ray transform”, preprint, 2012. [arXiv 1210.2084](#)
- [Zhang 2012] G. Zhang, “Uniqueness in the Calderón problem with partial data for less smooth conductivities”, *Inverse Problems* **28**:10 (2012), 105008, 18. [MR 2987911](#) [Zbl 1256.35207](#)
- [Zworski 2012] M. Zworski, *Semiclassical analysis*, Graduate Studies in Mathematics **138**, American Mathematical Society, Providence, RI, 2012. [MR 2952218](#) [Zbl 1252.58001](#)

Received 6 May 2013. Accepted 13 Nov 2013.

CARLOS KENIG: [cek@math.uchicago.edu](mailto:cek@math.uchicago.edu)

*Department of Mathematics, University of Chicago, 5734 S. University Avenue, Chicago, Illinois 60637, United States*

MIKKO SALO: [mikko.j.salo@jyu.fi](mailto:mikko.j.salo@jyu.fi)

*Department of Mathematics and Statistics, University of Jyväskylä, FI-40014 Jyväskylä, Finland*



# Analysis & PDE

[msp.org/apde](http://msp.org/apde)

## EDITORS

EDITOR-IN-CHIEF

Maciej Zworski  
[zworski@math.berkeley.edu](mailto:zworski@math.berkeley.edu)

University of California  
Berkeley, USA

## BOARD OF EDITORS

Nicolas Burq	Université Paris-Sud 11, France <a href="mailto:nicolas.burq@math.u-psud.fr">nicolas.burq@math.u-psud.fr</a>	Yuval Peres	University of California, Berkeley, USA <a href="mailto:peres@stat.berkeley.edu">peres@stat.berkeley.edu</a>
Sun-Yung Alice Chang	Princeton University, USA <a href="mailto:chang@math.princeton.edu">chang@math.princeton.edu</a>	Gilles Pisier	Texas A&M University, and Paris 6 <a href="mailto:pisier@math.tamu.edu">pisier@math.tamu.edu</a>
Michael Christ	University of California, Berkeley, USA <a href="mailto:mchrist@math.berkeley.edu">mchrist@math.berkeley.edu</a>	Tristan Rivière	ETH, Switzerland <a href="mailto:riviere@math.ethz.ch">riviere@math.ethz.ch</a>
Charles Fefferman	Princeton University, USA <a href="mailto:cf@math.princeton.edu">cf@math.princeton.edu</a>	Igor Rodnianski	Princeton University, USA <a href="mailto:irod@math.princeton.edu">irod@math.princeton.edu</a>
Ursula Hamenstaedt	Universität Bonn, Germany <a href="mailto:ursula@math.uni-bonn.de">ursula@math.uni-bonn.de</a>	Wilhelm Schlag	University of Chicago, USA <a href="mailto:schlag@math.uchicago.edu">schlag@math.uchicago.edu</a>
Vaughan Jones	U.C. Berkeley & Vanderbilt University <a href="mailto:vaughan.f.jones@vanderbilt.edu">vaughan.f.jones@vanderbilt.edu</a>	Sylvia Serfaty	New York University, USA <a href="mailto:serfaty@cims.nyu.edu">serfaty@cims.nyu.edu</a>
Herbert Koch	Universität Bonn, Germany <a href="mailto:koch@math.uni-bonn.de">koch@math.uni-bonn.de</a>	Yum-Tong Siu	Harvard University, USA <a href="mailto:siu@math.harvard.edu">siu@math.harvard.edu</a>
Izabella Laba	University of British Columbia, Canada <a href="mailto:ilaba@math.ubc.ca">ilaba@math.ubc.ca</a>	Terence Tao	University of California, Los Angeles, USA <a href="mailto:tao@math.ucla.edu">tao@math.ucla.edu</a>
Gilles Lebeau	Université de Nice Sophia Antipolis, France <a href="mailto:lebeau@unice.fr">lebeau@unice.fr</a>	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA <a href="mailto:met@math.unc.edu">met@math.unc.edu</a>
László Lempert	Purdue University, USA <a href="mailto:lempert@math.purdue.edu">lempert@math.purdue.edu</a>	Gunther Uhlmann	University of Washington, USA <a href="mailto:gunther@math.washington.edu">gunther@math.washington.edu</a>
Richard B. Melrose	Massachusetts Institute of Technology, USA <a href="mailto:rbm@math.mit.edu">rbm@math.mit.edu</a>	András Vasy	Stanford University, USA <a href="mailto:andras@math.stanford.edu">andras@math.stanford.edu</a>
Frank Merle	Université de Cergy-Pontoise, France <a href="mailto:Frank.Merle@u-cergy.fr">Frank.Merle@u-cergy.fr</a>	Dan Virgil Voiculescu	University of California, Berkeley, USA <a href="mailto:dvv@math.berkeley.edu">dvv@math.berkeley.edu</a>
William Minicozzi II	Johns Hopkins University, USA <a href="mailto:minicozz@math.jhu.edu">minicozz@math.jhu.edu</a>	Steven Zelditch	Northwestern University, USA <a href="mailto:zelditch@math.northwestern.edu">zelditch@math.northwestern.edu</a>
Werner Müller	Universität Bonn, Germany <a href="mailto:mueller@math.uni-bonn.de">mueller@math.uni-bonn.de</a>		

## PRODUCTION

[production@msp.org](mailto:production@msp.org)

Silvio Levy, Scientific Editor

---

See inside back cover or [msp.org/apde](http://msp.org/apde) for submission instructions.

---

The subscription price for 2013 is US \$160/year for the electronic version, and \$310/year (+\$35, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

---

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

---

APDE peer review and production are managed by EditFLOW<sup>®</sup> from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2013 Mathematical Sciences Publishers

# ANALYSIS & PDE

Volume 6 No. 8 2013

---

- $L^p$  and Schauder estimates for nonvariational operators structured on Hörmander vector fields with drift 1793  
MARCO BRAMANTI and MAOCHUN ZHU
- Strichartz estimates for Schrödinger equations with variable coefficients and unbounded potentials 1857  
HARUYA MIZUTANI
- Uniformity of harmonic map heat flow at infinite time 1899  
LONGZHI LIN
- A rotational approach to triple point obstructions 1923  
NOAH SNYDER
- On the energy subcritical, nonlinear wave equation in  $\mathbb{R}^3$  with radial data 1929  
RUIPENG SHEN
- Global well-posedness for the nonlinear Schrödinger equation with derivative in energy space 1989  
YIFEI WU
- The Calderón problem with partial data on manifolds and applications 2003  
CARLOS KENIG and MIKKO SALO