

**This is an electronic reprint of the original article.  
This reprint *may differ* from the original in pagination and typographic detail.**

**Author(s):** Tuikka, Matti; Haukka, Matti

**Title:** Crystal structure of the pyridine–diiodine (1/1) adduct

**Year:** 2015

**Version:**

**Please cite the original version:**

Tuikka, M., & Haukka, M. (2015). Crystal structure of the pyridine–diiodine (1/1) adduct. *Acta Crystallographica Section E : Crystallographic Communications*, 71(7), o463. <https://doi.org/10.1107/S2056989015010518>

All material supplied via JYX is protected by copyright and other intellectual property rights, and duplication or sale of all or part of any of the repository collections is not permitted, except that material may be duplicated by you for your research use or educational purposes in electronic or print form. You must obtain permission for any other use. Electronic or print copies may not be offered, whether for sale or otherwise to anyone who is not an authorised user.



# Crystal structure of the pyridine–diiodine (1/1) adduct

Matti Tuikka and Matti Haukka\*

University of Jyväskylä, Department of Chemistry, P.O. Box 35, FI-40014 University of Jyväskylä, Finland. \*Correspondence e-mail: matti.o.haukka@ju.fi

Received 17 April 2015; accepted 1 June 2015

Edited by C. Rizzoli, Università degli Studi di Parma, Italy


In the title adduct,  $C_5H_5N \cdot I_2$ , the N—I distance [2.424 (8) Å] is remarkably shorter than the sum of the van der Waals radii. The line through the I atoms forms an angle of 78.39 (16)° with the normal to the pyridine ring.

**Keywords:** pyridine; diiodine; halogen bonding; crystal structure.

**CCDC reference:** 1404151

## 1. Related literature

For the structure of the pyridine– $I_2$  1:2 adduct, see: Hassel & Hope (1961). For the crystal structures of pyridine with ICl and IBr, see: Rømming (1972); Dahl *et al.* (1967). For van der Waals radii, see: Bondi (1964). For the I—I distance of iodine, see: Buontempo *et al.* (1997). For I—I⋯N angles in halogen bonding, see: Desiraju *et al.* (2013).



## 2. Experimental

### 2.1. Crystal data

|                      |                                 |
|----------------------|---------------------------------|
| $C_5H_5N \cdot I_2$  | $c = 20.1953$ (13) Å            |
| $M_r = 332.90$       | $\beta = 98.468$ (3)°           |
| Monoclinic, $P2_1/c$ | $V = 801.16$ (8) Å <sup>3</sup> |
| $a = 9.2432$ (6) Å   | $Z = 4$                         |
| $b = 4.3392$ (2) Å   | Mo $K\alpha$ radiation          |

$\mu = 7.76$  mm<sup>-1</sup>  
 $T = 120$  K

$0.09 \times 0.07 \times 0.02$  mm

### 2.2. Data collection

Bruker KAPPA APEX II CCD diffractometer  
Absorption correction: numerical (SADABS; Bruker, 2012)  
 $T_{\min} = 0.574$ ,  $T_{\max} = 0.902$

6585 measured reflections  
1853 independent reflections  
1437 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.062$

### 2.3. Refinement

$R[F^2 > 2\sigma(F^2)] = 0.049$   
 $wR(F^2) = 0.091$   
 $S = 1.07$   
1853 reflections

73 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\max} = 1.11$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -1.26$  e Å<sup>-3</sup>

Data collection: Collect (Nonius, 2000); cell refinement: DENZO/SCALEPACK (Otwinowski & Minor, 1997); data reduction: DENZO/SCALEPACK; program(s) used to solve structure: SUPERFLIP (Palatinus & Chapuis, 2007; Palatinus & van der Lee, 2008; Palatinus *et al.*, 2012); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: OLEX2 (Dolomanov *et al.*, 2009); software used to prepare material for publication: OLEX2.

## Acknowledgements

Financial support provided by the Academy of Finland (project No. 129171) is gratefully acknowledged.

Supporting information for this paper is available from the IUCr electronic archives (Reference: RZ5157).

## References

- Bondi, A. (1964). *J. Phys. Chem.* **68**, 441–451.  
Bruker (2012). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.  
Buontempo, U., DiCiccio, A., Filippini, A., Nardone, M. & Postorino, P. (1997). *J. Chem. Phys.* **107**, 5720–5726.  
Dahl, T., Hassel, O. & Sky, K. (1967). *Acta Chem. Scand.* **21**, 592–593.  
Desiraju, G. R., Ho, P. S., Kloos, L., Legon, A. C., Marquardt, R., Metrangola, P., Politzer, P., Resnati, G. & Rissanen, K. (2013). *Pure Appl. Chem.* **85**, 1711–1713.  
Dolomanov, O. V., Bourhis, L. J., Gildea, R. J., Howard, J. A. K. & Puschmann, H. (2009). *J. Appl. Cryst.* **42**, 339–341.  
Hassel, O. & Hope, H. (1961). *Acta Chem. Scand.* **15**, 407–416.  
Nonius (2000). *COLLECT*. Nonius BV, Delft, The Netherlands.  
Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.  
Palatinus, L. & Chapuis, G. (2007). *J. Appl. Cryst.* **40**, 786–790.  
Palatinus, L., Prathapa, S. J. & van Smaalen, S. (2012). *J. Appl. Cryst.* **45**, 575–580.  
Palatinus, L. & van der Lee, A. (2008). *J. Appl. Cryst.* **41**, 975–984.  
Rømming, C. (1972). *Acta Chem. Scand.* **26**, 1555–1560.  
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

## supporting information

*Acta Cryst.* (2015). E71, o463 [doi:10.1107/S2056989015010518]

## Crystal structure of the pyridine–diiodine (1/1) adduct

Matti Tuikka and Matti Haukka

### S1. Comment


Diiodine is capable to act as halogen bond donor and form stable halogen bonds with Lewis bases, such as pyridine, due to the strong charge transfer. In the case of the pyridine-I<sub>2</sub> 1:2 adduct (Hassel & Hope, 1961), the interaction eventually results in the heterolytic cleavage of I<sub>2</sub> and formation of [py<sub>2</sub>I]<sup>+</sup> I<sub>3</sub><sup>-</sup> ion pairs. Although the crystal structures involving pyridine and interhalogens ICl and IBr are known (Rømming, 1972; Dahl *et al.*, 1967), the title pyI<sub>2</sub> 1:1 adduct has not been reported earlier. The N1—I1 distance in pyI<sub>2</sub> (2.425 (8) Å) is remarkably shorter than the sum of the van der Waals radii of iodine and nitrogen (3.53 Å; Bondi, 1964). The I—I distance (2.8043 (9) Å) is significantly longer than that observed in free diiodine in solid state (2.715 Å; Buontempo *et al.*, 1997). The I—I⋯N angle is approximately linear (176.44 (18)°) as expected in halogen bonds (Desiraju *et al.*, 2013).

### S2. Experimental

The title compound was synthesized by dissolving iodine (200 mg) in ethanol (5 ml) and adding pyridine (1 ml) into this solution. The solution was left to evaporate under ambient conditions and after a couple of days light yellow crystals were formed.

### S3. Refinement

All H atoms were positioned geometrically and refined using a riding model with C—H = 0.95 Å and with  $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$ .



**Figure 1**

The molecular structure of the title compound, with 50% probability displacement ellipsoids for non-H atoms.

## Pyridine–diiodine (1/1)

## Crystal data

C<sub>5</sub>H<sub>5</sub>N·I<sub>2</sub> $M_r = 332.90$ Monoclinic,  $P2_1/c$  $a = 9.2432$  (6) Å $b = 4.3392$  (2) Å $c = 20.1953$  (13) Å $\beta = 98.468$  (3)° $V = 801.16$  (8) Å<sup>3</sup> $Z = 4$  $F(000) = 592$  $D_x = 2.760$  Mg m<sup>-3</sup>Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 1865 reflections

 $\theta = 1.0$ – $27.5^\circ$  $\mu = 7.76$  mm<sup>-1</sup> $T = 120$  K

Plate, clear light yellow

 $0.09 \times 0.07 \times 0.02$  mm

## Data collection

Bruker KAPPA APEX II CCD  
diffractometer

Radiation source: fine-focus sealed tube

Curved graphite crystal monochromator

Detector resolution: 16 pixels mm<sup>-1</sup> $\varphi$  scans and  $\omega$  scans with  $\kappa$  offset

Absorption correction: numerical

(SADABS; Bruker, 2012)

 $T_{\min} = 0.574$ ,  $T_{\max} = 0.902$ 

6585 measured reflections

1853 independent reflections

1437 reflections with  $I > 2\sigma(I)$  $R_{\text{int}} = 0.062$  $\theta_{\max} = 27.6^\circ$ ,  $\theta_{\min} = 2.2^\circ$  $h = -11 \rightarrow 11$  $k = -5 \rightarrow 5$  $l = -26 \rightarrow 25$ 

## Refinement

Refinement on  $F^2$ 

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.049$  $wR(F^2) = 0.091$  $S = 1.07$ 

1853 reflections

73 parameters

0 restraints

Primary atom site location: iterative

Hydrogen site location: inferred from  
neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.0153P)^2 + 9.3396P]$ where  $P = (F_o^2 + 2F_c^2)/3$  $(\Delta/\sigma)_{\max} = 0.001$  $\Delta\rho_{\max} = 1.11$  e Å<sup>-3</sup> $\Delta\rho_{\min} = -1.26$  e Å<sup>-3</sup>

## Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å<sup>2</sup>)

|    | <i>x</i>    | <i>y</i>     | <i>z</i>    | $U_{\text{iso}}^*/U_{\text{eq}}$ |
|----|-------------|--------------|-------------|----------------------------------|
| I1 | 0.27234 (6) | 0.59684 (13) | 0.54587 (3) | 0.02083 (16)                     |
| I2 | 0.32645 (6) | 0.35101 (14) | 0.67558 (3) | 0.02480 (18)                     |
| N1 | 0.2243 (7)  | 0.8407 (18)  | 0.4368 (4)  | 0.0243 (18)                      |
| C5 | 0.3349 (9)  | 0.933 (2)    | 0.4053 (4)  | 0.0207 (19)                      |
| H5 | 0.4323      | 0.8743       | 0.4223      | 0.025*                           |
| C3 | 0.1668 (10) | 1.198 (2)    | 0.3214 (5)  | 0.028 (2)                        |
| H3 | 0.1471      | 1.3178       | 0.2818      | 0.033*                           |
| C1 | 0.0849 (10) | 0.921 (2)    | 0.4121 (5)  | 0.030 (2)                        |
| H1 | 0.0069      | 0.8529       | 0.4342      | 0.035*                           |

|    |             |           |            |           |
|----|-------------|-----------|------------|-----------|
| C2 | 0.0549 (10) | 1.098 (2) | 0.3558 (5) | 0.030 (2) |
| H2 | -0.0434     | 1.1533    | 0.3399     | 0.036*    |
| C4 | 0.3076 (10) | 1.112 (2) | 0.3480 (5) | 0.030 (2) |
| H4 | 0.3872      | 1.1783    | 0.3266     | 0.036*    |

*Atomic displacement parameters (Å<sup>2</sup>)*

|    | $U^{11}$   | $U^{22}$   | $U^{33}$   | $U^{12}$    | $U^{13}$   | $U^{23}$    |
|----|------------|------------|------------|-------------|------------|-------------|
| I1 | 0.0189 (3) | 0.0224 (3) | 0.0212 (3) | -0.0010 (2) | 0.0027 (2) | -0.0016 (3) |
| I2 | 0.0253 (3) | 0.0267 (3) | 0.0222 (4) | 0.0029 (2)  | 0.0031 (3) | 0.0011 (3)  |
| N1 | 0.016 (4)  | 0.032 (4)  | 0.024 (4)  | -0.003 (3)  | 0.004 (3)  | -0.007 (4)  |
| C5 | 0.017 (4)  | 0.031 (5)  | 0.013 (5)  | -0.001 (4)  | -0.001 (3) | 0.000 (4)   |
| C3 | 0.031 (5)  | 0.035 (6)  | 0.014 (5)  | -0.002 (4)  | -0.003 (4) | 0.004 (4)   |
| C1 | 0.019 (5)  | 0.039 (6)  | 0.031 (6)  | -0.009 (4)  | 0.007 (4)  | 0.005 (5)   |
| C2 | 0.014 (4)  | 0.049 (7)  | 0.026 (6)  | -0.002 (4)  | -0.003 (4) | 0.002 (5)   |
| C4 | 0.025 (5)  | 0.042 (6)  | 0.022 (6)  | -0.006 (4)  | 0.005 (4)  | 0.003 (5)   |

*Geometric parameters (Å, °)*

|             |             |             |            |
|-------------|-------------|-------------|------------|
| I1—I2       | 2.8043 (9)  | C5—C4       | 1.388 (13) |
| I1—N1       | 2.425 (8)   | C3—C2       | 1.397 (12) |
| N1—C5       | 1.342 (10)  | C3—C4       | 1.383 (13) |
| N1—C1       | 1.357 (12)  | C1—C2       | 1.364 (14) |
| N1—I1—I2    | 176.44 (18) | C4—C3—C2    | 116.6 (9)  |
| C5—N1—I1    | 120.7 (6)   | N1—C1—C2    | 121.1 (8)  |
| C5—N1—C1    | 119.8 (8)   | C1—C2—C3    | 120.9 (9)  |
| C1—N1—I1    | 118.9 (6)   | C3—C4—C5    | 121.2 (8)  |
| N1—C5—C4    | 120.3 (8)   |             |            |
| I1—N1—C5—C4 | -170.3 (7)  | C5—N1—C1—C2 | -0.8 (15)  |
| I1—N1—C1—C2 | 170.4 (8)   | C1—N1—C5—C4 | 0.8 (14)   |
| N1—C5—C4—C3 | -0.9 (15)   | C2—C3—C4—C5 | 1.0 (15)   |
| N1—C1—C2—C3 | 1.0 (16)    | C4—C3—C2—C1 | -1.0 (15)  |