

**This is an electronic reprint of the original article.
This reprint *may differ* from the original in pagination and typographic detail.**

Author(s): Kuokkanen, Riitta; Lappalainen, Raimo; Repo-Tiihonen, E.; Tiihonen, J.; Aho-Mustonen, K.

Title: Cognitive Insight, Clinical Insight, and Reasoning in Schizophrenia : A Pilot Study in a Forensic Setting

Year: 2016

Version:

Please cite the original version:

Kuokkanen, R., Lappalainen, R., Repo-Tiihonen, E., Tiihonen, J., & Aho-Mustonen, K. (2016). Cognitive Insight, Clinical Insight, and Reasoning in Schizophrenia : A Pilot Study in a Forensic Setting. *Journal of Forensic Psychology Practice*, 16(4), 253-267. <https://doi.org/10.1080/15228932.2016.1192337>

All material supplied via JYX is protected by copyright and other intellectual property rights, and duplication or sale of all or part of any of the repository collections is not permitted, except that material may be duplicated by you for your research use or educational purposes in electronic or print form. You must obtain permission for any other use. Electronic or print copies may not be offered, whether for sale or otherwise to anyone who is not an authorised user.

TABLE 1 Demographic Information for Study Participants

Characteristic	Mean (<i>SD</i>)	Range (%)
Age (years)	43.55 (12.24)	19-67
GAF*	17.39 (5.56)	10-31
Duration of illness (years)	16.45 (9.48)	3-37
Number of hospitalizations*	13.06 (10.75)	1-37
Duration of current admission (years)*	8.03 (6.52)	0.92-21.58
Education, highest completed, <i>n</i> *		
No formal education	1	(5%)
Elementary school	10	(50%)
Secondary education	7	(35%)
Diagnosis, <i>n</i>		
Paranoid schizophrenia	15	(75%)
Hebephrenic schizophrenia	1	(5%)
Undifferentiated schizophrenia	4	(20%)
Number of patients with		
comorbid substance abuse	11	(55%)
comorbid personality disorder	6	(30%)
History of violence*	18	(90%)

Note. *SD* = standard deviation; GAF = Global Assessment of Functioning

* Values missing from two forensic patients (*n*=18)

TABLE 2 Mean Values, Standard Deviations, Ranges, and Confidence Intervals for Means

Characteristic	Mean (<i>SD</i>)	Range	95% CI
PANSS P1 Delusions	2.20 (1.44)	1-5	1.53; 2.87
PANSS P6 Suspiciousness	3.00 (1.30)	1-5	2.39; 3.61
PANSS G12 Lack of judgment & insight	4.50 (1.15)	2-6	3.96; 5.04
PSYRATS Total	7.45 (5.51)	3-21	4.87; 10.03
PSYRATS Conviction	2.00 (1.75)	0-4	1.18; 2.82
PSYRATS Amount of distress	0.65 (1.27)	0-4	0.06; 1.24
PSYRATS Intensity of distress	0.50 (1.10)	0-4	-0.01; 1.01
BCIS Composite Index	6.15 (7.14)	-5-17	2.81; 9.49
BCIS Self-reflectiveness	15.30 (5.98)	6-27	12.50; 18.10
BCIS Self-certainty	9.15 (4.17)	0-16	7.20; 11.10
DTD	2.15 (1.76)	1-7	1.33; 2.97

Note. *SD* = standard deviation; *CI* = confidence interval; PANSS = Positive and Negative Syndrome Scale; PSYRATS = Psychotic Symptoms Rating Scales; BCIS = Beck Cognitive Insight Scale; DTD = draws-to-decision

TABLE 3

	DTD	BCIS SR	SC	CI	PANSS P1	P6	G12	PSYRATS Total	Amount of distress	Intensity of distress	Conviction
DTD											
BCIS SR		-.120	-.234	.054	-.293	-.113	-.344*	-.246	-.281*	-.300*	-.173
BCIS SC			-.055	.645***	.167	.144	-.231	.117	.050	.087	-.013
BCIS CI				-.445**	-.207	-.125	.319	-.117	-.044	-.060	-.113
PANSS P1					.187	.156	-.363	.094	.069	.100	-.020
PANSS P6						.373*	.213	.620***	.287	.327*	.600***
PANSS G12							.313	.400*	.188	.220	.500**
PSYRATS Total								.163	.013	.007	.233
Amount of distress									.475***	.533***	.673***
Intensity of distress										.513***	.240
Conviction											.253

Note. Significant values are shown in bold. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ (2-sided). DTD = draws-to-decision; BCIS = Beck Cognitive Insight Scale; SR = self-reflectiveness; SC = self-certainty; CI = composite index; PANSS = Positive and Negative Syndrome Scale; P1 = delusions item; P6 = suspiciousness item; G12 = lack of judgment & insight item; PSYRATS = Psychotic Symptoms Rating Scales