

**This is an electronic reprint of the original article.  
This reprint *may differ* from the original in pagination and typographic detail.**

**Author(s):** Soini, Anne; Gubbels, Jessica; Sääkslahti, Arja; Villberg, Jari; Kremers, Stef; Kann, Dave Van; Mehtälä, Anette; De Vries, Nanne; Poskiparta, Marita

**Title:** A Comparison of Physical Activity Levels in Childcare Contexts among Finnish and Dutch 3-Year-Olds

**Year:** 2016

**Version:**

**Please cite the original version:**

Soini, A., Gubbels, J., Sääkslahti, A., Villberg, J., Kremers, S., Kann, D. V., Mehtälä, A., De Vries, N., & Poskiparta, M. (2016). A Comparison of Physical Activity Levels in Childcare Contexts among Finnish and Dutch 3-Year-Olds. *European Early Childhood Education Research Journal*, 24(5). <https://doi.org/10.1080/1350293X.2016.1213569>

All material supplied via JYX is protected by copyright and other intellectual property rights, and duplication or sale of all or part of any of the repository collections is not permitted, except that material may be duplicated by you for your research use or educational purposes in electronic or print form. You must obtain permission for any other use. Electronic or print copies may not be offered, whether for sale or otherwise to anyone who is not an authorised user.

1

2

3

4

5

6

7

8

**A Comparison of Physical Activity Levels in Childcare Contexts among Finnish**

9

**and Dutch 3-Year-Olds**

10

## PHYSICAL ACTIVITY AT CHILDCARE

11 Abstract

12 To determine Finnish and Dutch 3-year-old preschool children's physical activity (PA) levels and  
13 how levels vary across gender, location, time of day and social contexts in both countries.

14 A modified version of the Observational System for Recording Physical Activity in Children-  
15 Preschool (OSRAC-P) was used to measure children's PA levels and contextual variables (e.g., group  
16 composition, prompts) of children attending childcare centres in Finland and the Netherlands. In total,  
17 90 Finnish children (46 boys) and 97 Dutch children (46 boys) were observed. Three-level linear  
18 regression analyses with cross level interactions were used to assess differences between the countries  
19 in the association between the context variables and PA. During the observations, the present sample  
20 of children was mostly sedentary in nature. Outdoor location and prompts (both positive and negative)  
21 were associated with higher levels of activity in both countries. Non-solitary playing was associated  
22 with higher activity levels in the Netherlands, but not in Finland, whereas child-initiated play was  
23 positively associated with Finnish but not Dutch children's PA levels. Finnish children were more  
24 active in the morning compared to the afternoon, while in the Dutch children PA was unaffected by  
25 time. The present findings indicate that better understanding of the contextual factors and interactions  
26 in children's PA behaviours across the two countries could help in planning childcare interventions to  
27 increase the PA levels of preschool children.

28 Keywords: physical activity; direct observation; childcare centre; country comparison; preschool  
29 children

### 30 **Introduction**

31 Nearly 70% of 3-year-old children in Europe attend childcare or early education services,  
32 whereas in other continents the numbers are much lower (OECD Family database 2008).  
33 There is some evidence that childcare centres influence children's physical activity (PA)  
34 levels (Finn, Johannsen, and Specker 2002; Pate et al. 2004; Strong et al. 2005; Pate et al.  
35 2008). Finn, Johannsen, and Specker (2002) concluded that among children attending  
36 childcare, the childcare centre was the strongest predictor of activity levels, with more than  
37 50% of the daily PA performed during childcare hours. However, various studies have  
38 indicated that PA levels are generally very low among preschool children during their time in  
39 childcare settings (Oliver, Schofield, and Kolt 2007; Pate et al. 2008; Brown et al. 2009-II;  
40 Reilly 2010; Gubbels et al. 2011). A U.S. study by Brown et al. (2009-II) reported 94% of the  
41 indoor activities observed to be sedentary. A smaller but nevertheless substantial proportion  
42 (59%) of the indoor activities observed among Dutch children also classified as sedentary  
43 (Gubbels et al. 2011), indicating that sedentariness might be more pronounced in U.S. pre-  
44 schoolers.

45 In a light of the socio-ecological framework, Hinkley et al. (2008) reviewed correlates  
46 of preschool children's PA level and showed that children's demographic and biological  
47 characteristics, as well as their physical and social environments are significantly associated  
48 with a children's total PA. Various studies have found links between physical environment  
49 factors in the childcare setting (e.g., the ground surface, playground markings, open space,  
50 play equipment) and children's PA (Zask et al. 2001; Dowda et al. 2004; Boldemann et al.  
51 2006; Ridgers et al. 2007; Bower et al. 2008; Cardon et al. 2008; Hannon & Brown 2008;  
52 Cosco et al. 2010; Gubbels, Van Kann, and Jansen 2012). Furthermore, boys have been found  
53 to be more active than girls (Oliver, Schofield, and Kolt 2007; Hinkley et al. 2008; Pate et al.  
54 2008), children are more active outdoors (Finn, Johannsen, and Specker 2002; Pate et al.

## PHYSICAL ACTIVITY AT CHILDCARE

55 2004; Boldeman et al. 2006; Hinkley et al. 2008), and children's PA increases in warm  
56 seasons and decreases in the colder seasons (Carson and Spence 2010).

57 Social environment factors, such as positive prompts by staff members (Brown et al.  
58 2009-I) and peers (Gubbels et al. 2011), have been shown to be associated with children's  
59 increased PA intensity. Despite these positive associations, teachers and peers rarely prompt  
60 children to increase their PA. Moreover, teacher-initiated play was negatively associated with  
61 children's levels of PA (Brown et al. 2009-II). European studies have also indicated negative  
62 relationships between the presence of more childcare staff and peers (Cardon et al. 2008) or  
63 direct involvement of staff and peers in children's play (Gubbels et al. 2011), and children's  
64 activity levels.

65 Currently, there is a lack of observational studies on pre-school children in Europe.  
66 Such observational studies provide valuable information on the context of PA behaviours,  
67 which cannot be derived from studies using accelerometer or other objective measurements  
68 of PA. Such contextual information can inform preventive interventions promoting PA in  
69 childcare, and have implications for both practitioners and researchers. The childcare setting  
70 holds great potential to make an important contribution to the welfare and health of young  
71 children by promoting PA (Pate et al. 2004; Bower et al. 2008; Hinkley et al. 2008; Pate et al.  
72 2008; Ward 2010). Furthermore, positive patterns of engagement in PA demonstrated during  
73 childhood are critical for the maintenance of a physically active lifestyle during both  
74 adolescence and adulthood (Strong et al. 2005).

75 The aim of this study was to determine existing PA levels among 3-year-old children  
76 and how these vary by gender, primary location (i.e. indoor vs. outdoor), time of day (i.e.  
77 morning vs. afternoon), social context (i.e. group composition, initiator of activity, prompts),  
78 outdoor temperatures and weather conditions during childcare in Finland and in the  
79 Netherlands.

80 **Methods**81 *Participants and Sampling*

82 **Finland.** Recruitment was performed in the city of Jyväskylä. A total of 14 childcare centres  
83 volunteered their participation in the study. In Finland, 3-year-olds in childcare are most  
84 commonly mixed in a group with 4- and 5-year-olds, while 1- and 2-year-olds are often  
85 grouped together. Finnish childcare centres in the present study provided care for an average  
86 of 74 children (SD = 20) in 4 groups (SD = 1). The mean number of staff members per centre  
87 was 16 (SD = 5). All the families of the 3-year-old children attending the childcare centres (N  
88 = 179) were invited to participate. One hundred and two (57%) parents provided informed  
89 consents. The children's PA data were collected between August and October 2010. Six  
90 children were not present and six children were attending for only half a day during the  
91 observations, and were therefore excluded from the final sample of 90 children (46 boys and  
92 44 girls) observed in Finland.

93 **The Netherlands.** A large Dutch childcare organization was approached to participate  
94 in the study, and gave consent to conduct the study. All 9 childcare centres in Maastricht run  
95 by this organization centres were approached and agreed to participate. In the Netherlands,  
96 childcare centres offer care for babies as young as 6 weeks to children up to four years, 3-  
97 year-olds are often grouped with 2-year-olds. In the present study, the Dutch childcare  
98 centres provided care for an average of 92 children (SD = 28) in 5 groups (SD = 1). The  
99 mean number of staff members per centre was 20 (SD = 6). All the parents of the 3-year-old  
100 children attending these childcare centres were informed about the study, and none of them  
101 refused to participate. Children's PA was observed in May and June 2008. A total of 97 3-  
102 year-olds (46 boys and 51 girls) were observed in the Netherlands.

103 The typical daily timetables in the Finnish and Dutch childcare centres are described  
104 in Table 1. The ethics committee of the University of Jyväskylä, and the Social Affairs and

## PHYSICAL ACTIVITY AT CHILDCARE

105 Health officer of the city of Jyväskylä approved the study. The Dutch study complied with  
106 the Dutch ‘Medical Research Involving Humans Act’.

107 INSERT Table 1 NEAR HERE (Table 1. Typical daily programmes in Finnish and Dutch  
108 childcare centres.)

### 109 *Measures and Procedures*

110 A translated and modified version of the Observational System for Recording Physical  
111 Activity in Children-Preschool Version (OSRAC-P; Brown et al. 2006) was used to measure  
112 children’s PA intensity, location, contexts, prompts and interactions. The design of the  
113 Finnish study was based on the Dutch study by Gubbels and colleagues (2011), with some  
114 minor adjustments to adapt to the Finnish situation. Two researchers in each country  
115 observed PA and contextual factors using a procedure in which an observation period of 15  
116 seconds was followed by 30 seconds for recording the observation. This procedure was  
117 repeated eight times for a total of six minutes per child. In Finland, each child was observed  
118 twice a day, both in the morning (between 8 a.m.-12 p.m.) *and* afternoon (2 p.m.-5 p.m.),  
119 yielding a total of sixteen observations per child. In the Netherlands, each child was observed  
120 either in the morning *or* afternoon, yielding a total of eight observations per child. The total  
121 number of single observations for both countries combined was thus 2,216 (1,440 in Finland;  
122 776 in the Netherlands). In both countries, children were randomly selected for observation  
123 and they were not observed during meals or rest times. To ensure accuracy of data, observer  
124 training (e.g., initial observer orientation, study of the observation manual and the  
125 memorization of codes and categories, direct in situ training session in field settings) is  
126 recommended (Brown et al. 2009-III). Therefore in both countries, before the measurements  
127 all field researchers were trained with the method by studying the instruments and  
128 background information, and subsequently observing children via videotape and ‘live’ in the  
129 childcare settings. Inter-observer agreement measures were done in both countries.

## PHYSICAL ACTIVITY AT CHILDCARE

130 Furthermore, to assess cross-country inter-observer agreement, a separate sample of  
131 children not included in the final study, was independently observed via videotape by one of  
132 the two researchers of each country. This resulted in cross-country coding of 305 observation  
133 intervals, i.e. 13.8% of the observation intervals in the main study (2216 intervals), sufficing  
134 the OSRAC-P norm of at least 12% independent coding (Brown et al. 2009-III).

135 PA intensity levels were measured on a five-point scale (1=stationary or motionless,  
136 2=stationary with limb or trunk movements, 3=slow or easy movements, 4=moderate  
137 movements, 5=fast movements) and reflected the highest intensity level reached by the child  
138 during each 15-second observation period. For the purpose of this study and further  
139 comparison, activity levels  $\leq 2$  were regarded as sedentary behaviour, activity level 3 as light  
140 PA and 4–5 levels as moderate-to-vigorous physical activity (MVPA) (Bower et al. 2006;  
141 Brown et al. 2009-II; Gubbels et al. 2011).

142 In addition to PA intensity, OSRAC-P scales assessing contextual variables primary  
143 location (inside vs. outdoor) and time of day (morning vs. afternoon) were used (Brown et al.  
144 2006). In addition, the following social OSRAC-P scales were assessed: group composition  
145 (i.e., solitary, one-to-one adult, one-to-one peer, group of children, group of children with  
146 adult(s)), initiator of activity (child or adult) and prompts (no prompts, staff member's/peers'  
147 positive prompts to increase PA or negative prompts to decrease PA). Children's gender was  
148 also recorded. Finally, weather conditions and outdoor temperatures were recorded per  
149 observation day.

### 150 *Statistical Analyses*

151 The scores of both observers on dichotomous variables (e.g., prompt by staff, yes/no) were  
152 combined by coding the variable as present (1) when one or both observers rated that variable  
153 as present, and coding it as absent (0) when both rated it as absent. For the continuous  
154 variables (e.g., activity intensity), the mean of the scores of both observers was calculated.



## PHYSICAL ACTIVITY AT CHILDCARE

155 Cohen's kappa was used to determine the inter-rater reliability (IRR) of the two observers for  
156 the observations of the OSRAC-P variables (i.e., activity intensity, type of activity, group  
157 composition, indoor and outdoor contexts, initiator of activity, prompts). The mean IRR of  
158 the variables assessed was 0.6 ( $SD = 0.2$ ;  $p < .001$ ) in Finland, and 0.7 ( $SD = 0.2$ ;  $p < .001$ ) in  
159 the Netherlands. The intraclass correlation used to determine cross-country IRR of two  
160 observers for the activity intensity variable was 0.7 ( $p < .001$ ).

161 Differences in the contextual variables, social context and gender between Finnish  
162 and Dutch children were examined using chi-square tests. Differences in mean activity  
163 intensity between the Finnish and Dutch children were examined using independent sample  $t$ -  
164 tests.

165 The association between gender (boy vs. girl), primary location (outdoor vs. indoor),  
166 time of day (afternoon vs. morning), group composition (non-solitary vs. solitary), initiator of  
167 activity (child vs. adult), prompts (no prompt vs. positive or negative prompt), temperature  
168 (linear) and weather condition (rain vs. sunny, with a clear sky or cloudy and dry) as  
169 independent, and mean PA intensity levels as dependent variables was examined using three-  
170 level linear regression, with cross-level interaction (MLR; with measurement level, child  
171 level, centre level) to examine differences between countries in these associations. All  
172 analyses were performed using SPSS 18.0 and STATA 12. In all analyses,  $P$ -values  $< .05$   
173 were considered statistically significant.

### 174 **Results**

#### 175 ***Temperature and Weather Conditions***

176 The mean outdoor temperature was 12.5°C during the Finnish observations (range: 2°C-20°C)  
177 and 20.5°C during Dutch observations (range: 14°C-26°C). This difference was significant ( $p$   
178  $< .001$ ). Most of the time weather was cloudy and dry (53%) or sunny with a clear sky (33%);  
179 the least prevalent was rain (14%). Differences between the countries in temperature

## PHYSICAL ACTIVITY AT CHILDCARE

180 categories and weather conditions, and in sedentary and MVPA associated with these  
181 categories are described in Table 2.

182 INSERT Table 2 NEAR HERE (Table 2. Differences between countries in percentages of  
183 observations observed and levels of sedentary activities and MVPA (N = 187).

### 184 *Contextual Variables and Activity Levels*

185 According to Chi-Square tests, children's indoor PA levels were mostly sedentary in nature:  
186 79% of total intervals recorded as sedentary, and only 3% coded as MVPA. Outdoors, 53%  
187 PA observations were classified as sedentary behaviour, whereas 10% were classified as  
188 MVPA. The initiators of the activities were most frequently children (81%). Staff and peers  
189 rarely prompted children to increase or decrease their PA: no prompts were recorded in 81%  
190 of all observations. Cross-country differences were more pronounced in the social and  
191 weather-related variables than in the non-social context variables. The Finnish children spent  
192 significantly more time in sedentary-level activities and less time in MVPA compared to the  
193 Dutch children in most of the observed categories (see Table 2).

### 194 *Associations Between Observed Contexts and PA*

195 The results of the multi-level linear regression are presented in Table 3. With respect to the  
196 primary location of the observations, children were significantly more active outdoors than  
197 indoors (activity intensity 2.65 vs. 2.18, respectively;  $p < .001$ ). In the Netherlands, an  
198 outdoor location had a stronger positive influence on children's activity levels than in Finland  
199 ( $p < .001$ ), although the influence was significant in both countries. All prompts (both  
200 positive and negative) were associated with an increase in the children's PA level in both  
201 countries. Non-solitary play was associated with higher activity levels in the Netherlands,  
202 whereas in Finland child-initiated play was positively associated with the children's PA  
203 levels. Finally, the Finnish children were less active in the afternoon compared to the  
204 morning, while the Dutch children's PA levels were unaffected by time of day.

## PHYSICAL ACTIVITY AT CHILDCARE

205 INSERT Table 3 NEAR HERE (Table 3. Children's mean PA intensity levels in Finland and in  
206 the Netherlands: three level linear regression.)

### 207 **Discussion**

208 The present study applied direct observation to compare Finnish and Dutch 3-year-old  
209 children's PA behaviour during childcare. In line with previous findings (Oliver, Schofield,  
210 and Kolt 2007; Pate et al. 2008; Brown et al. 2009-II; Gubbels et al. 2011), the present  
211 sample of children engaged in sedentary activities in 79% of indoor and 53% of outdoor  
212 observations. The results further showed significant differences between the countries in  
213 children's PA levels: the Finnish children spent significantly more time in sedentary-level  
214 activities and less time in MVPA compared to the Dutch children.

215 Several contextual variables had a differential influence on PA depending on the  
216 country. Surprisingly, this mainly concerned non-social influences: time of day and location,  
217 in addition to group composition. The significant interaction between country and time of day  
218 showed that the Finnish children were less active in the afternoon compared to mornings,  
219 while the Dutch children's PA levels were unaffected by time of day. The activities planned  
220 during an average childcare day provide little explanation for this difference, as they were  
221 very similar in both countries. The only difference worth noting is that in Finland fixed time  
222 for indoor and outdoor playing was scheduled, while in the Netherlands afternoons were  
223 spent flexibly, playing either indoors or outdoors, depending on, for instance, the weather and  
224 opportunities that arose during the day.

225 In the present study, temperature and weather conditions varied significantly between  
226 the two countries. However, we found no significant association between the temperature or  
227 weather variables and children's PA behaviour, and the absence of significant interactions  
228 between country and these variables indicates that these variables did not explain the  
229 differences in PA levels between the two countries. In line with this, Finn, Johannsen, and

## PHYSICAL ACTIVITY AT CHILDCARE

230 Specker (2002) found no seasonal variations in children's PA levels. Baranowski et al. (1993)  
231 indicated in their observational study that differences in children's PA were more related to  
232 time spent outdoors than to season or weather conditions. To date few seasonal studies have  
233 been conducted among preschool-aged children, and their findings have been conflicting  
234 (Carson and Spence 2010).

235 Consistent with previous findings (Pate et al. 2004; Boldeman et al. 2006; Hinkley et  
236 al. 2008; Brown et al. 2009-II), the present sample of children was physically more active  
237 outdoors than indoors. Outdoor location had a stronger positive influence on PA in the Dutch  
238 than Finnish children. Outdoor play in Dutch childcare centres is mainly unstructured, while  
239 in Finland it often involves teacher-arranged and teacher-initiated activities such as hiking. In  
240 general, children tend to be less active the more staff members there are present or involved  
241 with children's play (Cardon et al. 2008; Brown et al. 2009-II; Gubbels et al. 2011). Also in  
242 the present study, children were more sedentary when staff members were involved in  
243 children's activities, or when the initiator of the play was an adult. Perhaps a more  
244 unstructured and flexible approach is beneficial for raising children's PA levels. A Belgian  
245 study reported significant increases in children's objectively measured PA intensity during  
246 preschool recess and times of unstructured free plays that were taken as an opportunity to be  
247 physically active (Verbestel et al. 2011).

248 It is a matter for concern that the majority of the observations did not include any oral  
249 prompting by staff or peers, despite the fact that positive prompts were positively associated  
250 with time spent in MVPA. Other studies have confirmed that positive prompts by staff  
251 (Brown et al. 2009-I) and peers (Gubbels et al. 2011) have positive associations with  
252 children's PA intensity.

253 The differences between the two countries in children's PA levels were pronounced,  
254 although the differences in the social context, such as group sizes and numbers of children

## PHYSICAL ACTIVITY AT CHILDCARE

255 and staff members in the current childcare centres, were not. However, one notable difference  
256 between the countries was revealed in the children's typical childcare attendance. If children  
257 attend childcare once or twice per week, as is typical in the Netherlands (European  
258 Commission's Expert Group on Gender and Employment Issues [EGGE] 2009), this factor  
259 alone may positively influence their level of PA. For Finnish children, who often attend  
260 childcare five days a week (OECD Family database 2008), childcare is part of their normal  
261 daily routine and may not exert any particular influence on their PA. Another explanation for  
262 the cross-country difference may be found in group membership: Finnish 3-year-olds were  
263 grouped with 4- to 5-year-old children, whereas Dutch 3-year-old children were often  
264 grouped with 2-year-olds. The Finnish observers may have unintentionally underrated the 3-  
265 year-olds' behaviour when this was observed against the backdrop of the older, more skilled  
266 children present, while in the Netherlands the opposite may have occurred: the observers may  
267 have overrated the PA of the Dutch 3-year-olds by unwittingly comparing them with the 2-  
268 year-olds. The sufficiently high cross-country IRR contradicts this explanation, though. A  
269 final explanation might lie in differences in the physical childcare environment in the two  
270 countries, which was not taken into account in the study. The physical environment has  
271 previously been shown to significantly influence the activity levels of children in childcare  
272 (e.g., Cosco et al. 2010; Gubbels, Van Kann, and Jansen 2012).

273 A strength of this study is that PA levels were directly observed using the OSRAC-P  
274 method in two European countries. Observations were made in a total of 23 childcare centres  
275 indoors and outdoors, and during both mornings and afternoons, thereby covering regular  
276 childcare attendance times in both Finland and the Netherlands. The childcare daily schedules  
277 and outdoor times in both countries were very similar, and no major differences were  
278 observed in programmes. The benefit of the observation format was that it recorded not only  
279 the intensity of activity, but also *where, how and in what kind of interaction* the activity was

## PHYSICAL ACTIVITY AT CHILDCARE

280 being engaged. Moreover, the OSRAC-P has been shown to be a valid and reliable tool for  
281 measuring PA among preschool-aged children (Brown et al. 2006; Trost 2007; Pate, O'Neill,  
282 and Mitchell 2010).

283         This study has limitations that should be noted. The generalizability of the findings  
284 could be limited by the fact that in both countries all the participating childcare centres and  
285 children were located in the same city, though, the samples represented the general  
286 population structure. In Finland the number of observations was higher than in the  
287 Netherlands, however, comparisons were made between mean values, therefore it was  
288 possible to include all existing Finnish observations to provide more accurate data without  
289 affecting the final results. Furthermore, the direct observation method is based on subjective  
290 assessments. Indeed, the observers in the two countries were not the same, which might have  
291 influenced the findings, possibly explaining the systematically higher PA levels found in the  
292 Dutch compared to Finnish children. However, the interobserver reliabilities in both  
293 countries, as well as cross-country indicated appropriate levels of agreement.

### 294 **Conclusion**

295         In light of the growing concern about children's PA behaviour, this study indicated  
296 that interventions at the childcare level are needed to increase the PA levels of both Finnish  
297 and Dutch 3-year-old preschool children. The findings emphasized that the childcare setting  
298 itself may play an important part in promoting positive patterns of health behaviour during  
299 early childhood. Interventions should focus on increasing children's outdoor time, free play,  
300 positive prompting and encouragement by staff members. A better understanding of the  
301 interactions between and within different contextual factors in children's behaviours should  
302 help to identify ways of promoting participation in PA during childcare attendance.

### 303 ***Summary Bullets***

- 304         ▪ The present sample of children engaged in sedentary activities in 79% of indoor and

## PHYSICAL ACTIVITY AT CHILDCARE

305           53% of outdoor observations.

- 306           ▪ Children were more sedentary when staff members were involved in children's  
307           activities, or when the initiator of the play was an adult.
- 308           ▪ The majority of the observations did not include any oral prompting by staff or peers,  
309           despite the fact that positive prompts were positively associated with time spent in  
310           MVPA.
- 311           ▪ The Finnish children spent significantly more time in sedentary-level activities and less  
312           time in MVPA compared to the Dutch children.
- 313           ▪ The present findings indicate that better understanding of the contextual factors and  
314           interactions in children's PA behaviours across the two countries could help in  
315           planning childcare interventions to increase the PA levels of preschool children.

### 316 *Acknowledgements*

317 We appreciate the cooperation of the Finnish and Dutch childcare centres, children and their  
318 parents who participated in this study.

### 319 **Conflict of interest**

320 The Authors state that they have no conflicts of interest.

321

322 **References**

- 323 Baranowski, T., W. O. Thompson, R. H. DuRant, J. Baranowski, and J. Puhl. 1993.  
324 Observations on physical activity in physical locations: age, gender, ethnicity, and  
325 month effects. *Research Quarterly for Exercise and Sport* 64: 127–133.
- 326 Boldeman, C., M. Blennow, H. Dal, F. Mårtensson, A. Raustorp, K. Yuen, and U Wester.  
327 2006. Impact of preschool environment upon children’s physical activity and sun  
328 exposure. *Preventive Medicine* 42: 301–308.
- 329 Bower, J. K., D. P. Hales, D. F. Tate, D. A Rubin, S. E. Benjamin, and D. S. Ward. 2008. The  
330 childcare environment and children’s physical activity. *American Journal of Preventive  
331 Medicine* 34: 23–29.
- 332 Brown, W.H., H.S. Googe, K.L. McIver, and M . J. Rathel. 2009-I. Effects of teacher-  
333 encouraged physical activity on preschool playgrounds. *Journal of Early Intervention*  
334 31: 126–145.
- 335 Brown, W. H., K. A., Pfeiffer, K. L. McIver, M. Dowda, C. L. Addy, and R. R. Pate. 2009-II.  
336 Social and environmental factors associated with preschoolers’ nonsedentary physical  
337 activity. *Child Development* 80: 45–58.
- 338 Brown, W. H., K. A., Pfeiffer, K. L. McIver, M. Dowda, M. J. C. A. Almeida, and R. R. Pate.  
339 2006. Assessing preschool children’s physical activity: The observational system for  
340 recording physical activity in children-preschool version. *Research Quarterly for  
341 Exercise and Sport* 77: 167–176.
- 342 Brown, W. H., K. A. Pfeiffer, K. L. McIver, M. Dowda, M. J. Almeida, and R. R. Pate. 2009-  
343 III. Observational System for Recording Physical Activity in Children-Preschool  
344 (OSRAC-P). OSRAC-P Training Manual for Observers Columbia: University of South  
345 Carolina.
- 346 Cardon, G., E. Cauwenberghe, V. Labarque, L. Haerens, and I. De Bourdeaudhuij. 2008. The  
347 contribution of preschool playground factors in explaining children’s physical activity  
348 during recess. *International Journal of Behavioral Nutrition and Physical Activity*, 5.
- 349 Carson, V., and J. C. Spence. 2010. Seasonal variation in physical activity among children  
350 and adolescents: a review. *Pediatric Exercise Science* 22: 81–92.
- 351 Cosco, N. G., R. C. Moore, and M. Z. Islam. 2010. Behavior mapping: A method for linking  
352 preschool physical activity and outdoor design. *Medicine & Science in Sports &  
353 Exercise* 42: 513–519.
- 354 Dowda, M., R. R. Pate, S. G. Trost, M. J. C. A. Almeida, and J. R. Sirard. 2004. Influences of  
355 preschool policies and practices on children’s physical activity. *Journal of Community  
356 Health* 29: 183–196.
- 357 European Commission’s Expert Group on Gender and Employment Issues (EGGE) 2009.  
358 The provision of childcare services. A comparative review of 30 European countries.  
359 Luxemburg: Office for Official Publications of the European Communities.
- 360 Finn, K. J., N. Johannsen, and B. Specker. 2002. Factors associated with physical activity in  
361 preschool children. *The Journal of Pediatrics* 140: 81–85.
- 362 Gubbels, J. S., S. P. J. Kremers, D. H. H. Van Kann, A. Stafleu, M. J. J. M. Candel, P. C.  
363 Dagnelie, C. Thijs, and N. K. de Vries. 2011. Interaction between physical environment,  
364 social environment, and child characteristics in determining physical activity at child  
365 care. *Health Psychology* 30: 84–90.
- 366 Gubbels, J. S., D. H. H. Van Kann, and M. W. J. Jansen. 2012. Play equipment, physical  
367 activity opportunities, and children’s activity levels at childcare. *Journal of  
368 Environmental and Public Health*, doi:10.1155/2012/326520.



## PHYSICAL ACTIVITY AT CHILDCARE

- 369 Hannon, J. C., and B. B. Brown. 2008. Increasing preschoolers' physical activity intensities:  
370 An activity-friendly preschool playground intervention. *Preventive Medicine* 46: 532–  
371 536.
- 372 Hinkley, T., D. Crawford, J. Salmon, A. D. Okely, and K. Hesketh. 2008. Preschool children  
373 and physical activity. *American Journal of Preventive Medicine* 34: 435–441.
- 374 OECD Family database 2008. Last updated June 2011. Retrieved June 2, 2012 from  
375 <http://www.oecd.org/els/social/family/> database.
- 376 Oliver, M., G. M. Schofield, and G. S. Kolt. 2007. Physical activity in preschoolers.  
377 Understanding prevalence and measurement issues. *Sports Medicine* 37: 1045–1070.
- 378 Pate, R. R., J. R. O'Neill, and J. Mitchell. 2010. Measurement of physical activity in  
379 preschool children. *Medicine & Science in Sports & Exercise* 42: 508–512.
- 380 Pate, R. R., K. McIver, M. Dowda, W. H. Brown, and C. Addy. 2008. Directly observed  
381 physical activity levels in preschool children. *Journal of School Health* 78: 438–444.
- 382 Pate, R. R., K. A. Pfeiffer, S. G. Trost, P. Ziegler, and M. Dowda. 2004. Physical activity  
383 among children attending preschools. *Pediatrics* 114: 1258–1263.
- 384 Reilly, J. J. 2010. Low levels of objectively measured physical activity in preschoolers in  
385 child care. *Medicine & Science in Sports & Exercise* 42: 502–507.
- 386 Ridgers, N. D., G. Stratton, S. J. Fairclough, and W. R. Twisk. 2007. Long-term effects of a  
387 playground markings and physical structures on children's recess physical activity  
388 levels. *Preventive Medicine* 44: 393–397.
- 389 Strong, W. B., R. M. Malina, C. R. Blimkie, S. R. Daniels, R. K. Dishman, and B. Gutin.  
390 2005. Evidence based physical activity for school-age youth. *Journal of Pediatrics* 146:  
391 732–737.
- 392 Trost, S. G. 2007. Measurement of physical activity in children and adolescents. *American*  
393 *Journal of Lifestyle Medicine* 1: 299–314.
- 394 Ward, D. S. 2010. Physical activity in young children: The role of child care. *Medicine &*  
395 *Science in Sports & Exercise* 42: 499–501.
- 396 Verbestel, V., E. Van Cauwenberghe, V. De Coen, L. Maes, I. De Bourdeaudhuij, and G.  
397 Cardon. 2011. Within- and between-day variability of objectively measured physical  
398 activity in pre-schoolers. *Pediatric Exercise Science* 23: 366–378.
- 399 Zask, A., E. van Beurden, L. Barnett, L. O. Brooks, and U. C. Dietrich. 2001. Active school  
400 playgrounds-myth or reality? Results of the "move it groove it" project. *Preventive*  
401 *Medicine* 33: 402–408.