

**This is an electronic reprint of the original article.
This reprint *may differ* from the original in pagination and typographic detail.**

Author(s): Kärnä, Minna; Lahtinen, Manu; Valkonen, Jussi

Title: 1,1,4,4-Tetramethylpiperazinediium dibromide

Year: 2009

Version:

Please cite the original version:

Kärnä, M., Lahtinen, M., & Valkonen, J. (2009). 1,1,4,4-Tetramethylpiperazinediium dibromide. *Acta Crystallographica, section E*, 65(11), o2952.

All material supplied via JYX is protected by copyright and other intellectual property rights, and duplication or sale of all or part of any of the repository collections is not permitted, except that material may be duplicated by you for your research use or educational purposes in electronic or print form. You must obtain permission for any other use. Electronic or print copies may not be offered, whether for sale or otherwise to anyone who is not an authorised user.

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

1,1,4,4-Tetramethylpiperazinediium dibromide

Minna Kärnä, Manu Lahtinen* and Jussi Valkonen

University of Jyväskylä, Department of Chemistry, PO Box 35, FIN-40014 JY, Finland
Correspondence e-mail: manu.lahtinen@jyu.fi


Received 23 October 2009; accepted 28 October 2009

Key indicators: single-crystal X-ray study; $T = 123$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å;
R factor = 0.025; wR factor = 0.061; data-to-parameter ratio = 14.4.

A small quantity of the title compound, $\text{C}_8\text{H}_{20}\text{N}_2^{2+} \cdot 2\text{Br}^-$, was formed as a by-product in a reaction between a diamine and an alkyl bromide. The asymmetric unit contains half of a centrosymmetric dication and a bromide anion. In the crystal, weak intermolecular $\text{C}-\text{H} \cdots \text{Br}$ hydrogen bonds consolidate the crystal packing.

Related literature

For a possible synthetic route, see Creighton & Taylor (1987). For related structures, see; Linden *et al.* (1999, 2002); Guo *et al.* (2007).



Experimental

Crystal data

 $\text{C}_8\text{H}_{20}\text{N}_2^{2+} \cdot 2\text{Br}^-$ $M_r = 304.08$ Monoclinic, $P2_1/c$ $a = 5.8769$ (12) Å $b = 8.4584$ (17) Å $c = 11.200$ (2) Å $\beta = 92.79$ (3)° $V = 556.07$ (19) Å³ $Z = 2$ Mo $K\alpha$ radiation $\mu = 7.25$ mm⁻¹ $T = 123$ K $0.24 \times 0.16 \times 0.16$ mm

Data collection

Bruker Kappa APEXII
diffractometerAbsorption correction: multi-scan
(SADABS; Sheldrick, 2004) $T_{\min} = 0.296$, $T_{\max} = 0.390$

5032 measured reflections

1370 independent reflections

1223 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.032$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.025$ $wR(F^2) = 0.061$ $S = 1.11$

1370 reflections

95 parameters

All H-atom parameters refined

 $\Delta\rho_{\text{max}} = 0.38$ e Å⁻³ $\Delta\rho_{\text{min}} = -0.70$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
$\text{C2}-\text{H2A} \cdots \text{Br1}^{\text{i}}$	0.96 (3)	2.88 (4)	3.816 (4)	165 (2)
$\text{C2}-\text{H2B} \cdots \text{Br1}$	0.93 (3)	2.92 (4)	3.820 (4)	163 (2)
$\text{C2}-\text{H2C} \cdots \text{Br1}^{\text{ii}}$	0.98 (3)	2.83 (4)	3.770 (3)	161 (2)
$\text{C3}-\text{H3B} \cdots \text{Br1}^{\text{iii}}$	1.00 (3)	2.84 (3)	3.806 (4)	163 (2)
$\text{C4}-\text{H4A} \cdots \text{Br1}^{\text{iv}}$	0.93 (3)	2.92 (2)	3.566 (2)	127 (2)
$\text{C4}-\text{H4B} \cdots \text{Br1}^{\text{ii}}$	0.97 (3)	2.86 (5)	3.787 (2)	159 (2)

Symmetry codes: (i) $-x + 1, -y, -z + 1$; (ii) $x, -y + \frac{1}{2}, z - \frac{1}{2}$; (iii) $-x + 1, y + \frac{1}{2}, -z + \frac{3}{2}$; (iv) $x - 1, -y + \frac{1}{2}, z - \frac{1}{2}$.

Data collection: COLLECT (Nonius, 1999); cell refinement: DENZO-SMN (Otwinowski & Minor, 1997; Otwinowski *et al.*, 2003); data reduction: DENZO-SMN; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: Mercury (Macrae *et al.*, 2006); software used to prepare material for publication: SHELXL97.

The authors thank the Inorganic Materials Chemistry Graduate Program and the Magnus Ehrnrooth Foundation for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CV2642).

References

- Creighton, J. L. & Taylor, M. J. (1987). *Can. J. Chem.* **65**, 2526–2528.
- Guo, H.-X., Wu, S.-Z., Cai, M.-S. & Yao, S.-S. (2007). *Acta Cryst.* **E63**, m2747.
- Linden, A., Nugent, K. W., Petridis, A. & James, B. D. (1999). *Inorg. Chim. Acta*, **285**, 122–128.
- Linden, A., Petridis, A. & James, B. D. (2002). *Acta Cryst.* **C58**, m53–m55.
- Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
- Nonius (1999). COLLECT. Nonius BV, The Netherlands.
- Otwinowski, Z., Borek, D., Majewski, W. & Minor, W. (2003). *Acta Cryst.* **A59**, 228–234.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Sheldrick, G. M. (2004). SADABS. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supporting information

Acta Cryst. (2009). E65, o2952 [doi:10.1107/S1600536809045139]

1,1,4,4-Tetramethylpiperazinediium dibromide

Minna Kärnä, Manu Lahtinen and Jussi Valkonen

S1. Comment

Low quantity of the title compound (Fig. 1) was formed as a byproduct in a synthesis between a tetramethylethylenediamine (TMEDA) and ethoxyethylbromide. Most probably residues of dibromoethane existed as an impurity on either of the starting materials as it is known that piperazinium can be formed by reacting TMEDA and 1,2-dibromoethane. The compound has been recrystallized from acetonitrile/methanol solvent and its crystal structure is reported here.

The asymmetric unit consists of one anion and half a cation. The C—H···Br distances vary from 2.826 (30) to 2.924 (20) Å. In the crystal, cations are packed columnary along *a* axis forming at the same time layers along *b* axis. The bromide anions are analogously packed between the cation layers. The structure is stabilized by weak intermolecular C—H···Br interactions. Cation conformation of this compound is similar to those previously reported tetraiodidocadmate and pentabromothallate salts.

S2. Experimental

The compound was a byproduct from a reaction between tetramethylethylenediamine and ethoxyethylbromide. Few crystals suitable for a single-crystal structure determination recrystallized from an acetonitrile-methanol solution.

S3. Refinement

All H atoms were located from the difference map and refined isotropically.




Figure 1

Left: The molecular structure of (I) showing 50% probability displacement ellipsoids and the atomic numbering [symmetry code: (i) $-x + 1, y + 1/2, -z + 3/2$]. Right: Spacefill presentation of location of eight bromides around a single dication. Six of the anions belong to the neighboring ion pairs.

1,1,4,4-Tetramethylpiperazinedium dibromide*Crystal data*

C₈H₂₀N₂²⁺·2Br⁻
M_r = 304.08
 Monoclinic, *P*2₁/*c*
a = 5.8769 (12) Å
b = 8.4584 (17) Å
c = 11.200 (2) Å
 β = 92.79 (3)°
V = 556.07 (19) Å³
Z = 2

F(000) = 304
D_x = 1.816 Mg m⁻³
 Mo *K*α radiation, λ = 0.71073 Å
 Cell parameters from 3094 reflections
 θ = 0.4–28.3°
 μ = 7.25 mm⁻¹
T = 123 K
 Block, colourless
 0.24 × 0.16 × 0.16 mm

Data collection

Bruker Kappa APEXII
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 φ and ω scans
 Absorption correction: multi-scan
 (*SADABS*; Sheldrick, 2004)
T_{min} = 0.296, *T_{max}* = 0.390

5032 measured reflections
 1370 independent reflections
 1223 reflections with *I* > 2σ(*I*)
R_{int} = 0.032
 θ_{\max} = 28.2°, θ_{\min} = 3.0°
h = -6→7
k = -11→10
l = -14→14

Refinement

Refinement on *F*²
 Least-squares matrix: full
R[*F*² > 2σ(*F*²)] = 0.025
wR(*F*²) = 0.061
S = 1.11
 1370 reflections
 95 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: difference Fourier map
 All H-atom parameters refined
 $w = 1/[\sigma^2(F_o^2) + (0.0256P)^2 + 0.322P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.38 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.70 \text{ e \AA}^{-3}$

Special details

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of *F*² against ALL reflections. The weighted *R*-factor *wR* and goodness of fit *S* are based on *F*², conventional *R*-factors *R* are based on *F*, with *F* set to zero for negative *F*². The threshold expression of *F*² > 2σ(*F*²) is used only for calculating *R*-factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. *R*-factors based on *F*² are statistically about twice as large as those based on *F*, and *R*-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	<i>U_{iso}</i> [*] / <i>U_{eq}</i>
N1	0.5241 (3)	0.3260 (2)	0.48286 (16)	0.0111 (4)
C2	0.6413 (5)	0.1983 (3)	0.4148 (2)	0.0146 (5)
C3	0.3697 (5)	0.2474 (3)	0.5693 (2)	0.0155 (5)
C4	0.3920 (4)	0.4292 (3)	0.3948 (2)	0.0120 (4)
C5	0.2945 (4)	0.5749 (3)	0.4538 (2)	0.0117 (5)

Br1	0.89763 (4)	0.01387 (3)	0.702200 (19)	0.01391 (10)
H2A	0.524 (5)	0.137 (3)	0.374 (2)	0.017 (7)*
H2B	0.720 (5)	0.138 (3)	0.473 (2)	0.013 (7)*
H2C	0.733 (4)	0.252 (3)	0.356 (2)	0.011 (6)*
H3A	0.462 (5)	0.186 (3)	0.620 (2)	0.019 (7)*
H3B	0.292 (5)	0.333 (3)	0.614 (2)	0.020 (7)*
H3C	0.270 (5)	0.190 (3)	0.524 (3)	0.021 (8)*
H4A	0.272 (4)	0.370 (3)	0.360 (2)	0.010 (6)*
H4B	0.491 (5)	0.456 (3)	0.331 (2)	0.010 (6)*
H5A	0.182 (5)	0.546 (3)	0.519 (2)	0.013 (7)*
H5B	0.220 (4)	0.637 (3)	0.394 (2)	0.005 (6)*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0101 (10)	0.0108 (9)	0.0122 (9)	-0.0002 (7)	0.0004 (7)	-0.0004 (7)
C2	0.0186 (14)	0.0101 (11)	0.0149 (11)	0.0019 (10)	0.0000 (10)	-0.0024 (9)
C3	0.0157 (14)	0.0142 (12)	0.0168 (12)	-0.0011 (10)	0.0034 (10)	0.0033 (9)
C4	0.0128 (12)	0.0105 (10)	0.0122 (10)	0.0004 (9)	-0.0029 (9)	-0.0010 (9)
C5	0.0116 (12)	0.0095 (10)	0.0137 (11)	0.0019 (9)	-0.0034 (9)	-0.0011 (9)
Br1	0.01243 (16)	0.01658 (15)	0.01261 (14)	0.00082 (9)	-0.00030 (9)	-0.00037 (8)

Geometric parameters (Å, °)

N1—C4	1.505 (3)	C3—H3B	1.01 (3)
N1—C5 ⁱ	1.506 (3)	C3—H3C	0.90 (3)
N1—C2	1.507 (3)	C4—C5	1.524 (3)
N1—C3	1.512 (3)	C4—H4A	0.93 (3)
C2—H2A	0.96 (3)	C4—H4B	0.97 (3)
C2—H2B	0.93 (3)	C5—N1 ⁱ	1.506 (3)
C2—H2C	0.99 (3)	C5—H5A	1.04 (3)
C3—H3A	0.93 (3)	C5—H5B	0.94 (2)
C4—N1—C5 ⁱ	108.49 (18)	N1—C3—H3C	105.5 (18)
C4—N1—C2	108.52 (17)	H3A—C3—H3C	113 (2)
C5 ⁱ —N1—C2	107.85 (18)	H3B—C3—H3C	112 (2)
C4—N1—C3	111.58 (19)	N1—C4—C5	112.19 (18)
C5 ⁱ —N1—C3	112.09 (17)	N1—C4—H4A	108.5 (16)
C2—N1—C3	108.19 (18)	C5—C4—H4A	108.6 (16)
N1—C2—H2A	107.1 (16)	N1—C4—H4B	108.0 (16)
N1—C2—H2B	105.1 (16)	C5—C4—H4B	112.5 (15)
H2A—C2—H2B	111 (2)	H4A—C4—H4B	107 (2)
N1—C2—H2C	106.7 (15)	N1 ⁱ —C5—C4	112.47 (19)
H2A—C2—H2C	109 (2)	N1 ⁱ —C5—H5A	104.9 (15)
H2B—C2—H2C	117 (2)	C4—C5—H5A	112.6 (15)
N1—C3—H3A	106.8 (18)	N1 ⁱ —C5—H5B	108.7 (15)
N1—C3—H3B	107.6 (15)	C4—C5—H5B	108.2 (14)
H3A—C3—H3B	112 (2)	H5A—C5—H5B	110 (2)

C5 ⁱ —N1—C4—C5	-55.1 (3)	C3—N1—C4—C5	68.8 (2)
C2—N1—C4—C5	-172.1 (2)	N1—C4—C5—N1 ⁱ	57.4 (3)

Symmetry code: (i) $-x+1, -y+1, -z+1$.

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
C2—H2A \cdots Br1 ⁱⁱ	0.96 (3)	2.88 (4)	3.816 (4)	165 (2)
C2—H2B \cdots Br1	0.93 (3)	2.92 (4)	3.820 (4)	163 (2)
C2—H2C \cdots Br1 ⁱⁱⁱ	0.98 (3)	2.83 (4)	3.770 (3)	161 (2)
C3—H3B \cdots Br1 ^{iv}	1.00 (3)	2.84 (3)	3.806 (4)	163 (2)
C4—H4A \cdots Br1 ^v	0.93 (3)	2.92 (2)	3.566 (2)	127 (2)
C4—H4B \cdots Br1 ⁱⁱⁱ	0.97 (3)	2.86 (5)	3.787 (2)	159 (2)

Symmetry codes: (ii) $-x+1, -y, -z+1$; (iii) $x, -y+1/2, z-1/2$; (iv) $-x+1, y+1/2, -z+3/2$; (v) $x-1, -y+1/2, z-1/2$.