

**This is an electronic reprint of the original article.
This reprint *may differ* from the original in pagination and typographic detail.**

Author(s): Villalta, Irene; Amor, Fernando; Galarza, Juan; Dupont, Simon; Ortega, Patrocinio;
Hefetz, Abraham; Dahbi, Abdallah; Cerdá, Xim; Boulay, Raphaël

Title: Origin and distribution of desert ants across the Gibraltar Straits

Year: 2017

Version:

Please cite the original version:

Villalta, I., Amor, F., Galarza, J., Dupont, S., Ortega, P., Hefetz, A., Dahbi, A., Cerdá, X.,
& Boulay, R. (2017). Origin and distribution of desert ants across the Gibraltar Straits.
Molecular Phylogenetics and Evolution, 118, 122-134.
<https://doi.org/10.1016/j.ympev.2017.09.026>

All material supplied via JYX is protected by copyright and other intellectual property rights, and duplication or sale of all or part of any of the repository collections is not permitted, except that material may be duplicated by you for your research use or educational purposes in electronic or print form. You must obtain permission for any other use. Electronic or print copies may not be offered, whether for sale or otherwise to anyone who is not an authorised user.

Accepted Manuscript

Origin and distribution of desert ants across the Gibraltar Straits

Irene Villalta, Fernando Amor, Juan A. Galarza, Simon Dupont, Patrocinio Ortega, Abraham Hefetz, Abdallah Dahbi, Xim Cerdá, Raphaël Boulay


PII: S1055-7903(17)30135-5
DOI: <https://doi.org/10.1016/j.ympev.2017.09.026>
Reference: YMPEV 5936

To appear in: *Molecular Phylogenetics and Evolution*

Received Date: 12 February 2017
Revised Date: 6 September 2017
Accepted Date: 30 September 2017

Please cite this article as: Villalta, I., Amor, F., Galarza, J.A., Dupont, S., Ortega, P., Hefetz, A., Dahbi, A., Cerdá, X., Boulay, R., Origin and distribution of desert ants across the Gibraltar Straits, *Molecular Phylogenetics and Evolution* (2017), doi: <https://doi.org/10.1016/j.ympev.2017.09.026>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Origin and distribution of desert ants across the Gibraltar Straits

Irene Villalta^{1,2,3}, Fernando Amor¹, Juan A. Galarza⁴, Simon Dupont², Patrocinio Ortega¹, Abraham Hefetz⁵, Abdallah Dahbi⁶, Xim Cerdá¹ and Raphaël Boulay^{2*}

1. Estación Biológica de Doñana, CSIC, Avenida Américo Vespucio 26, 41092 Sevilla, Spain
2. Institute of Insect Biology, Parc de Grandmont, 37200 Tours, France
3. Departamento de Ecología, Universidad de Granada, Avenida de la Fuente Nueva S/N 18071 Granada, Spain
4. Centre of Excellence in Biological Interactions, Dept. of Biological and Environmental Science, University of Jyväskylä, Finland
5. Department of Zoology, George S. Wise Faculty of Life Sciences, Tel Aviv University, Tel Aviv, Israel
6. Department of Biology, Cadi Ayyad University, Polydisciplinary Faculty of Safi, 46000 Safi, Morocco

*Corresponding author:

Raphaël Boulay

Institute of Insect Biology, Parc de Grandmont, 37200 Tours, France

Email: Raphael.boulay@univ-tours.fr

Tel. +33 (0) 698345172

Fax. +33 (0) 02 47 36 69 66

Keywords: ants; *Cataglyphis*; phylogeography; glaciations; thermophily; genital traits; cuticular hydrocarbons

Summary

The creation of geographic barriers has long been suspected to contribute to the formation of new species. We investigated the phylogeography of desert ants in the western Mediterranean basin in order to elucidate their mode of diversification. These insects which have a low dispersal capacity are recently becoming important model systems in evolutionary studies. We conducted an extensive sampling of species belonging to the *Cataglyphis albicans* group in the Iberian Peninsula (IP) and the northern Morocco (North Africa; NA). We then combined genetic, chemical and morphological analyses. The results suggest the existence of at least three and five clades in the IP and NA, respectively, whose delineation partially encompass current taxonomic classification. The three Iberian clades are monophyletic, but their origin in NA is uncertain (79% and 22% for Bayesian and Maximum Likelihood support, respectively). The estimation of divergence time suggests that a speciation process was initiated after the last reopening of the Gibraltar Straits ≈ 5.33 Ma. In the IP, the clades are parapatric and their formation may have been triggered by the fragmentation of a large population during the Pleistocene due to extended periods of glaciation. This scenario is supported by demographic analyses pointing at a recent expansion of Iberian populations that contrasts with the progressive contraction of the NA clades. Niche modeling reveals that this area, governed by favorable climatic conditions for desert ants, has recently increased in the IP and decreased in NA. Altogether, our data points at geoclimatic events as major determinants of species formation in desert ants, reinforcing the role of allopatric speciation.

INTRODUCTION

Speciation is a key determinant of the diversification of life and regional species richness. Theory predicts that the formation of new species mostly occurs in allopatry (Coyne and Orr, 2004; Mayr, 1963), although it has been suggested that sympatric speciation may occur at a more common rate than initially thought (Nosil, 2008). Allopatric speciation arises due to the absence of gene flow between populations separated by geographic barriers through vicariance or dispersal. The process of speciation may be accelerated by natural selection acting on diverging populations subject to different ecological conditions promoting the evolution of phenotypic differences (Nosil, 2012). However, niche conservatism—the tendency for new species to retain similar ecological requirements—may also limit phenotypic divergence in spite of reproductive isolation (Wiens et al., 2010). This might be particularly true in taxa in which reproductive and non-reproductive functions are completed by different individuals like social insects where worker morphological traits are often insufficient for species delineation (Schlick-Steiner et al., 2006; Seifert, 2009). The suppression of the geographic barriers may then allow secondary contacts between reproductively isolated populations that are phenotypically indistinguishable (Eyer et al., 2017; Jowers et al., 2014). Alternatively, populations of the same species distributed over large environmental gradients may show major phenotypic differences as a result of local adaptation and phenotypic plasticity, even though complete reproductive isolation is not achieved. In this context, phylogeographic analyses can provide useful information on the role of past and current geographic barriers on species formation and species delineation at a regional scale.

The Mediterranean basin is one of the world's main biodiversity hotspots (Myers et al., 2000). This extraordinary regional diversity partly results from the recurrent encounters of North-African and Eurasian biota as a consequence of geo-climatic events during the Cenozoic (Husemann et al., 2014). For example, the progressive closure of the Gibraltar Straits between North Africa (NA) and the Iberian Peninsula (IP), which culminated in the

creation of a land-bridge during the Messinian (Achalhi *et al.* 2016), may have allowed the migration of terrestrial species in both directions (Fig 1). The sudden re-opening of the Gibraltar Straits \approx 5.33 Ma, may then have initiated numerous processes of speciation through vicariance. This is supported by the fact that, despite being only 14 km wide, the current Gibraltar Straits is an effective barrier to gene flow, even for organisms with good dispersal capacities (Casimiro-Soriguer *et al.*, 2010; Castella *et al.*, 2000; Garcia-Mudarra *et al.*, 2009; Pardo *et al.*, 2008; Salzburger *et al.*, 2002; Sanchez-Robles *et al.*, 2014). Moreover, measures of divergence between sister clades of snakes and lizards distributed on both sides of the Gibraltar Straits often coincide with a scenario of late Miocene vicariance (Albert *et al.*, 2007; Paulo *et al.*, 2008; Velo-Anton *et al.*, 2012). However, other studies suggested that divergence occurred much earlier (Gantenbein and Largiadèr 2003) or much later than this period (Brändli *et al.*, 2005; Carranza *et al.*, 2006a, 2006b; Jowers *et al.*, 2015). In some lizard species a first migration event from the IP to NA during the late Miocene was followed by a second migration event in the early Pleistocene (-2.3 to -2.7 Ma), perhaps through natural rafting (Kaliontzopoulou *et al.*, 2011).

Climatic oscillations during the Quaternary period may also have promoted important vicariance events. Extensive studies based on fossil records and mtDNA analyses have revealed that the IP served as a refugium for temperate species that migrated towards lower latitudes during Ice Ages (Hewitt, 1999, 2004; Taberlet *et al.*, 1992). The marked genetic structuring of extant populations has revealed that migrations towards these favorable areas led to the formation of a mosaic of fragmented populations that persisted in small refugia throughout the entire Iberian refugium (Abellán and Svenning, 2014; Centeno-Cuadros *et al.*, 2009; Ferrero *et al.*, 2011; Gomez and Lunt, 2007; Gonçalves *et al.*, 2015; Miraldo *et al.*, 2011). A similar pattern of small refugia occupying river valleys separated by high elevation habitats has been postulated in NA (Husemann *et al.*, 2014). In some cases, climate stability during long periods may have initiated allopatric speciation between the isolated lineages through drift and/or selection. Post-glacial range expansion may then have led to secondary contact between sister lineages (Gonçalves *et al.*, 2015; Igea *et al.*, 2013; Jowers *et al.*,

2014; Lefebvre et al., 2016; Miraldo et al., 2011).

Previous studies on the thermophilic ant genus *Cataglyphis* suggest that it constitutes an interesting group in which to trace the phylogeography of this complex region (Boulay et al. 2017; Jowers et al. 2013, 2014). These species occupy hot and xeric environments of Eurasia and NA, where workers benefit from unusual thermoresistance (they are able to withstand a ground temperature $> 70^{\circ}\text{C}$) thereby limiting predation and competition with other ants and with vertebrates (Cerdá and Retana, 2000; Wehner et al., 1992). The high species richness found in the Middle East suggests that they may have diversified in this region and dispersed east- and westward (Radchenko 1997; Ruano et al. 2011; Sanllorente et al. 2015). In the western Mediterranean basin, *Cataglyphis* species have been regrouped into 6-7 phylogenetic groups (Aron et al. 2016; Knaden et al. 2012; Radchenko 1997). In the present study, we were specifically interested in the phylogeography of the *albicans* group around the Gibraltar Straits in order to determine the role of geographic isolation as a driver of new species formation. This group is particularly useful for this study because it contains several species distributed in both the IP (n=4) and Morocco (n= 9; Cagniant 2009). Nevertheless, many taxonomic and phylogenetic issues remain unresolved in this group, due to the low morphological differentiation among the workers.

To determine the North-African origin of the Iberian desert ants in the *C. albicans* group we conducted an extensive sampling of *C. albicans* in Morocco, Spain, and Portugal and determined their phylogenetic relationships with DNA sequences. We used the fossil-based divergence time estimation between *Cataglyphis iberica* collected in northern Spain and various other species of ants (Moreau and Bell, 2013) in order to calibrate our phylogenetic tree. We employed microsatellite markers to determine the level of isolation between identified clades. We also used male morphological measurements and the composition of worker cuticular hydrocarbons to supplement clade delineation. Male morphology, particularly the copulatory organs, is often considered a good marker of reproductive isolation, since a small difference in shape may prevent copulation between different species (Masly 2012; Kubota et al., 2013; Wojcieszek and Simmons, 2013; Jowers

et al., 2014). Moreover, cuticular hydrocarbons have been shown to be species-specific and to constitute useful tools in the identification of cryptic ant species (Bagnères and Wicker-Thomas, 2010; Dahbi et al., 2008, 1996; Jowers et al., 2014). Finally, we combined demographic analyses and niche modeling to test the “refugium within refugium” hypothesis (Gomez and Lunt 2007) in NA and the IP.

Materials and methods

Sample collection

A total of 130 *Cataglyphis* colonies were sampled across the IP and northern Morocco between 2011 and 2015 (Table S1; average distance between sampled colonies: 484 km, range: 0.14 – 1870 km). From all colonies, we collected several workers that were immediately conserved in 96% ethanol. When available, males were also collected and stored in 70% ethanol for morphological analyses. Finally, some workers were brought alive to the laboratory for cuticular hydrocarbon (CHC) extraction (see below). Species identification was reviewed by ant systematist authority Prof. Henri Cagniant (formerly at the Paul Sabatier University of Toulouse). Voucher specimens are conserved at the Doñana Biological Station collection.

DNA extraction, amplification, and sequencing

Genomic DNA was extracted from the thorax and legs of one worker per colony using Wizard® Genomic DNA Purification Kit (Promega). A 503 bp portion of the mitochondrial cytochrome oxidase subunit I (COI), 604 bp of the long wavelength rhodopsin (LWRh), and 779 bp of the ribosomal RNA (28S) were amplified from each specimen using the primers summarized in Table 1. Polymerase chain reaction (PCR) amplifications were performed with the HotstarTaq® Master Mix Kit (Qiagen). PCR reactions were performed in a volume of 30 µl using 1 µl of extracted DNA, 1.5 µl of each primer (20mM), 15 µl HotStarTaq Master Mix and 11µl of Nuclease-free water. PCR amplification was conducted with an initial

denaturation step at 95 °C (15 min), followed by 35 cycles at 94 °C (45 sec), annealing temperature (1 min), and 72 °C (1 min 30 sec), and finally by an extension step at 72 °C (10 min). The PCR templates were purified with NucleoSpin® Gel and PCR Clean-up kit (Macherey-Nagel) and sequenced on an automated ABI 3730XL sequencer using Big Dye® Terminator cycle sequencing kit (Applied Biosystems, Foster City, California, USA) at the Genoscreen facility (<http://www.genoscreen.fr/>).

Sequence analysis and phylogenetic inference

All sequences were corrected, aligned, and analyzed in Geneious v. 8.0 (Kearse et al. 2012) using the MAFFT plugin (Kato et al., 2002) with the auto option for other parameters set as default and then refined manually. For the protein coding genes, the correct translation to amino acids was checked, and introns were considered as a single separate partition for LWRH. DnaSP v. 5.1 (Librado and Rozas, 2009) was used to retrieve haplotypes through the algorithms provided by PHASE (Stephens and Donnelly 2003) as well as to determine nucleotide and haplotype diversity, and neutrality tests of Tajima (D) and Fu and Li (D*). All sequences were deposited in GenBank (KY100127-KY100256, KY314267-KY314396, KY314626-KY314755).

Bayesian phylogenetic analyses on the concatenated DNA matrix were implemented in BEAST2 v. 2.3.0 (Bouckaert et al., 2014) for the joint estimation of the topology of the tree and the ages of diversification. Analyses were conducted applying the best fitting substitution model to each partition as estimated by PartitionFinder v. 1.1.1 (Lanfear et al., 2012), with the following parameters: linked branch length, models available in BEAST, BIC model selection, and greedy search algorithm. We applied a relaxed uncorrelated lognormal molecular clock and a Yule speciation process prior. *Formica wheeleri* (DQ353362, DQ353149, DQ353556), *Proformica nasuta* (DQ353311, DQ353195, DQ353581) DNA sequences retrieved from GenBank and *Cataglyphis tartessica* (KY314398, KY314397, KY314756) sequences obtained in this study were included as outgroups. To calibrate the tree, we used the divergence time estimated by Moreau and Bell (2013) between *Formica*

wheeleri, *Proformica nasuta*, and *C. iberica*. Analyses were run for 30 million generations, sampling one tree every 1000 generations.

A maximum clade credibility tree was obtained with TreeAnnotator v. 1.8.1 (Drummond *et al.* 2012) after checking effective sample size values (ESS > 200) using Tracer v. 1.6 (Rambaut and Drummond, 2013). Ten percent of the initial trees was discarded as burning fraction. Maximum likelihood analyses were performed using the Geneious RAxML v. 7.2 plugin with the GTR CAT model. The dataset was partitioned according to the best-fit partitioning schemes. Node support was obtained after 1000 bootstrap replicates.

Haplotype median joining networks were constructed for the three genes using Network v. 4.6 (Bandelt *et al.*, 1999) applying the MP correction (Polzin and Vahdati Daneshmand, 2003) and avoiding hypervariable sites. Finally, the relative similarity among the clades identified by Bayesian inference was also analyzed by Discriminant Analysis of Principal Components (DAPC) using Adegenet (Jombart *et al.*, 2010) in R (R Development Core Team, 2015).

To determine whether expansion/contraction processes had occurred on both sides of the Mediterranean, we pooled all the samples from the IP and all the samples from NA into two groups. For each group, we estimated effective population size changes through time according to the Bayesian Skyline Plot (BSP) method implemented in BEAST v. 2.3.0 (Bouckaert *et al.*, 2014), setting a strict molecular clock and partitions and substitution models as defined above. Chains were run for 30 million iterations, of which the first 25% was discarded as burn-in. Genealogies and model parameters were sampled every 1000 iterations. The results were analyzed and summarized as BSPs in Tracer v. 1.6 (Rambaut and Drummond, 2013).

Microsatellite analyses

We obtained microsatellite data for 10 loci from 129 of the 130 specimens (one specimen had to be discarded owing to amplification problems). The microsatellite markers were developed for *C. cursor* (Chéron *et al.*, 2011; Percy *et al.*, 2004). Genotyping was

performed using multiplex PCR reactions. Details of multiplex compositions, reaction conditions, and temperature profiles are given in the supplementary information S2. Fragment analysis was performed using an ABI 3730 XL (Applied Biosystems). Allele size determination and binning was conducted using the Microsatellite plugin in Geneious. We tested for the occurrence of null alleles and scoring errors at each locus with MICROCHECKER 2.2.3 software (Van Oosterhout et al. 2004). Allele richness and frequency, observed and expected heterozygosity, and Wright's F-statistics were computed using GENEPOP ON THE WEB (Raymond and Rousset 1995). Linkage disequilibrium and departures from Hardy–Weinberg Equilibrium (HWE) were tested using GenAEx v. 6.5.

The Bayesian clustering software TESS 2.3.1 (Durand et al., 2009) was used to infer the most likely number of populations (K) accounting for the spatial distribution of the samples. Models with and without admixture were evaluated using default parameters, setting K from 2 to 15. The optimal number of clusters was determined by plotting the Deviance Information Criterion (DIC) against K . The software package CLUMMP v. 1.1.2 (Jakobsson and Rosenberg, 2007) was used to evaluate the consistency of results across 15 independent runs using the full-search algorithm. For sequence data, we also analyzed genotypic differences among the clades using DAPC (Jombart et al. 2010).

CHC analyses

To compare the CHC composition of the main clades identified using sequence data, we analyzed 5-10 workers from 5 - 9 colonies per clade. Colonies collected from the same clade were at least 10 km apart. Before extraction, the workers were killed by placing them at -18°C for 20 min. The carcasses were immersed in 2 ml of hexane for 24 h. All the workers from one colony were pooled together. The CHC extracts were then stored at -18 °C until the analyses were conducted using a Gas Chromatograph coupled to a Mass Spectrometer (7890 GC System, 7000C GC/MS triple quad, Agilent technologies). Extracts were completely evaporated and re-suspended in 50 µl of GC quality Hexane (Sigma Aldrich). One microliter of each sample was injected into a 30m-long HP-5MS fused silica capillary

column (HP-5 MS UI, Agilent) with a splitless time of 2 min at 250°C and a Helium flow rate of 1 ml.min⁻¹. The temperature program was set from 100°C (2 min initial hold) to 320°C at 6°C.min⁻¹ (5 min of final hold). The cuticular components were identified using an electron impact of 70eV (scanning mode 40–550 amu). Fragments were analyzed with the Masshunter software (qualitative analysis B.06.00, Agilent technologies) and compared to public data. Quantification was achieved by calculating peak area relative to the total peak area.

Only compounds with an average relative quantity higher than 1% of the total CHC composition in at least one specimen per clade were used in the statistical analyses. Because the number of CHC identified in the samples was too large compared with the number of samples, we first regrouped the compounds according to their carbon chain length and number of methyl branching. Differences in profiles among the clades identified by BI were then analyzed statistically using Discriminant Analysis of Principal Components implemented in the Ade package in R (R Development Core Team 2015).

Male genitalia analyses

A total of 211 males were collected from 25 locations (range: 1 to 20 males per location). For each male, 15 measurements were taken with a millimetric ruler, a digital camera Canon 30D with MP-E 65 mm macro lens, and the software ImageJ (Abràmoff et al., 2004). The measurements comprised head length and width, inter-ocular distance, scape length, pronotum length and hind femur length, subgenital plate width and length, paramere length, penis valves width, length and distance between the apexes of their sclerotic appendices, volsella width, and length and lacinia length. Morphological differences among specimens were then analyzed through principal component analysis in R (R Development Core Team 2015).

Geographic distribution of the clades and niche contraction

Current clades' niche modeling was conducted using Maxent v. 3.3 (Philips *et al.* 2004, 2006). First, we retrieved 19 bioclimatic variables from WORLDCLIM online database (<http://www.worldclim.org/>) with a resolution of 2.5 arc-minute. These 19 variables (Table S3) present climate conditions for the period 1960-1990 in the study area and were used to train a model of current distribution of the identified clades. The same bioclimatic variables estimated for the Last Glacial Maximum through the Max Planck Institute for Meteorology new Earth system model (MPI-ESM), also recovered from WORLDCLIM, were used to predict the potential geographic distribution of each clade 0.025 Ma.

RESULTS

Phylogenetic analyses

The combined dataset for the 130 sequenced specimens consisted in 1890 bp. 23.7%, 4.2% and 2% of the COI, LWRh and 28S positions, respectively, were parsimony informative. Haplotype number and diversity were much greater for COI (97 and 0.99, respectively) than for both nuclear genes (31 and 0.87 and 25 and 0.79 for LWRh and 28S, respectively). The best evolution models are given in Table 2. The Bayesian analysis of the three concatenated genes resulted in a well-supported phylogram that allows the identification of eight clades (Fig. 2). Descriptive statistics for each clade and the results of neutrality tests are given in Table 3. None of the tests showed a significant departure from neutrality.

The Bayesian method suggested that the separation between *C. tartessica* and *C. albicans* group occurred 9.66 Ma (95% interval: 15.15 – 5.21Ma). Among the *albicans* group, the most ancient node (7.26 Ma; 95% interval: 11.34 – 4.04 Ma) separated a group of North-African specimens only (clades E to H) with another group composed of both Iberian (clades A, B and C) and North-African (clade D) specimens. The three Iberian clades were well-supported (BI support = 1) and showed a parapatric distribution (Fig. 2 and 3). The separation among these three clades may have occurred less than 3 Ma.

Clade D contained North African specimens identified as *C. otini* and *C. cubica*. The position of this clade as a sister lineage to the three Iberian clades was not strongly

supported (posterior probability = 0.7). Divergence time between clade D and the common ancestor of the Iberian clades was estimated as 6.17Ma (95% interval: 9.69 – 3.18Ma). In contrast to the IP, the NA clades had overlapping distributions (Fig 3).

Maximum Likelihood (Fig 4) and the Median Joining Network (Fig. S4) were consistent with Bayesian Inference. However, the position of clade D as a sister clade of the three Iberian clades was not supported by the Maximum Likelihood analysis (22%). Separate analyses conducted on each gene revealed a much greater structuring for the mtDNA than for both nuclear genes, probably due to the greater rate of evolution of the former. The DAPC analysis conducted on the sequence data also suggested a major segregation between the three Iberian clades and the five North-African clades along the first axis, which explained 61% of the variance (Fig 5a). The second axis explained 26% of the variance and separated clade D from the other clades.

The BSP method showed a constant population size followed by a recent contraction for the NA group (Fig 6). In contrast, the IP group revealed a constant increase in population size with a rapid recent expansion. These results, however, should be taken with caution as samples were pooled, and hence treated as a single species under uniform selection pressure throughout their distribution range. Nonetheless, they are informative in illustrating general trends of demographic processes occurring on both sides of the Mediterranean, particularly after the opening of the Gibraltar Straits.

Microsatellite analyses

The summarized statistics for the 129 typed specimens and 10 loci are given in Table 4. Details for each locus and the results of Hardy-Weinberg tests are given in Table S4. All loci were polymorphic with allele numbers ranging from 4.3 ± 0.7 in clade F to 15.2 ± 2.3 in clade C. There was evidence of linkage disequilibrium for only four out of 45 pairs of loci (Table S6). Observed and expected heterozygosity ranged between 0.47 ± 0.08 and 0.59 ± 0.10 , respectively in clade A to 0.62 ± 0.08 and 0.76 ± 0.09 , respectively in clade C. The fixation

indexes were relatively high in all the clades, suggesting a high level of genetic differentiation within them.

The analysis of microsatellite markers in TESS supported the existence of seven and five clusters with and without admixture, respectively (as indicated by the breaks in the plot of the Deviance Information Criteria against the number of clusters; Fig 7). Independently of the exact number of clusters, the first level of clustering separated the Iberian and North-African clades, suggesting that the Gibraltar Straits represented a major barrier to gene flow between the IP and NA. Another level of structuring within NA was shown by the well-defined clade D, which denoted a lack of gene flow with other North-African and Iberian clades. Moreover, clade E showed a major partitioning with specimens grouping either with clade G or F. A high level of genetic diversity was found in clade H. In contrast, a very low-level structuring existed within the IP. The analysis conducted in DAPC supported the separation between the IP and NA (Fig 5b). In addition, DAPC suggested a clear distinction between clade A and clades B and C. Among the NA clades, D clearly segregated from the others while E and G overlapped.

CHC analyses

A total of 46 CHC were identified on the ant cuticle (Table S7). All were alkanes that belonged to 22 categories based on their chain length (25 to 33) and number of methylations (0 to 3). The DAPC conducted on these 22 categories suggested that all the clades had well distinct profiles except E and F (Fig 5c). All the profiles were relatively common for ants and were dominated by C29 and C27 linear- and methyl-alkanes. Clades F and H differed from the others by having abundant long-chain alkanes (C31 for clade F and di- and trimethyl-C33 for clade H). In contrast, clade B was limited to trimethyl-C29 and lacked carbon chain longer than C30.

Male biometry

Only males from clades A to E could be collected in the field. The 15 measurements of male

genitalia are given in Table S8. The result of the DAPC conducted on these measurements supported morphological differences among the three Iberian clades (Fig 5d). In addition, males of clades A, B, and C clearly differed from the two NA clades (D and E). The variables that most separated among the five clades were the measurements of genitalia organs (i.e. Paramere length, Subgenital plate width, Volsella width; Table S8).

Geographic distribution of the clades and niche contraction

Despite low sample size for some clades, the use of 19 bioclimatic variables enabled a relatively good prediction of the clades' current distribution (Fig. 8). The distributions of favorable environmental conditions for the three Iberian clades showed a clear extension since the Last Glacial Maximum. This was particularly noticeable for clade C, for which favorable conditions 0.025 Ma were limited to three small areas in the east, west and south of the IP. In contrast, the distribution of favorable conditions for the NA clades showed evidences of a drastic reduction since the Last Glacial Maximum.

DISCUSSION

Molecular analyses reveal the existence of at least eight clades among the *C. albicans* group that currently occupies NA and the IP. These clades also show phenotypic differences. The three Iberian clades are monophyletic. Although their position with respect to the North-African clades is not strongly supported, their separation may have occurred after the last reopening of the Gibraltar Straits during the Miocene. In contrast to the NA clades, the IP clades have a paratric distribution. Moreover, both their population size and available niche show signs of expansion. This suggests that the formation of three independent clades in the IP may have been triggered by repeated glaciations along the Pleistocene. Recent warming may then have allowed secondary contacts between these new species.

The analysis of mitochondrial and nuclear sequences of specimens collected in the IP and NA resulted in a clear delineation of at least eight clades that are also characterized by chemical and/or morphological differences. Bayesian and Maximum Likelihood methods of

phylogenetic inference suggest a monophyly of the three Iberian clades. Their position with respect to the NA clades is uncertain. Hence, the supports for the node between clade D and the Iberian clades are weak. Moreover, the DAPC conducted on the sequences data highlights a clear discrimination between D and all the other clades. Finally, microsatellites analysis in TESS tends to associate clade D with the NA rather than with the IP clades. Nevertheless, all methods except the CHC analysis point at the uniqueness of clade D and the topologies of the trees are in agreement with recent phylogenies of the genus *Cataglyphis* that suggest that *C. iberica* and *C. rosenhaueri* derive from *C. cubica*—which composes clade D—(Aron et al., 2016; Cagniant, 2009). Moreover, the absence of species belonging to the *C. albicans* group in other regions of Europe and its commonness in the Middle East and Maghreb support the African origin of the Iberian clades. The alternative according to which migration from the middle East occurred through the North of the Mediterranean is more likely for the *C. cursor* group of which isolated populations are known in Greece, Italy, France and Spain but not in NA. Future analyses using a phylogenomic approach may help resolving these uncertainties (Blaimer et al., 2015). In addition, it would be interesting to obtain samples from other countries of NA to get a more precise picture of species diversification in this region.

The Bayesian tree calibrated using the divergence between *F. wheeleri*, *P. nasuta*, and *C. iberica* (Moreau and Bell 2013) dates the branching of clade D and the Iberian clades to 6.17 Ma (9.69 – 3.18 Ma). This is consistent with the ancestor of the Iberian clades having crossed the Gibraltar Straits through a stepping-stone dispersal process in the late Miocene. The progressive closure of the straits in the early Tortonian (~8 Ma; Fig. 1) led to the formation of temporal islands separated by ephemeral corridors that were probably small enough (<1 km) to permit dispersal of *Cataglyphis* queens and males. The reduction of water flow between the Atlantic and the Mediterranean culminated in the formation of a continuous land bridge in the early Messinian (6.1 Ma; Achalhi et al., 2016), which probably enhanced the northbound dispersal of African species. However, the fast reopening of the Gibraltar Straits 5.3 Ma may have rapidly limited gene flow and induced vicariance. A similar scenario

of late Miocene dispersal from NA towards the IP has been suggested for some terrestrial vertebrates (Albert et al., 2007; Kaliontzopoulou et al., 2011; Paulo et al., 2008). This may also apply to other desert ants, like the *C. altisquamis* and *C. emmae* groups that occur on both sides of the Gibraltar Straits. It is intriguing, however, that some other groups with apparently similar dispersal capacity are frequent in NA but are missing in IP. This is the case of the *C. bicolor* group and of the *C. albicans* clades E-H. One explanation could be that these clades were not yet present in NA in the late Miocene and thus clearly could not have crossed the Gibraltar Straits. Alternatively, they may have dispersed towards the IP but became extinct possibly during the Pleistocene glaciation, as suggested, for example, for *Malpolon* snakes (Carranza et al., 2006a).

The three Iberian clades of *C. albicans* have a parapatric distribution that contrasts with the sympatry of the North-African clades. A similar geographic pattern among phylogenetically-related species exists in amphibians (Gonçalves et al., 2015), reptiles (Albert et al., 2007), and also in desert ants of the *altisquamis* and *emmae* groups (Jowers et al., 2014; Tinaut, 1990). This may indicate a second, much more recent vicariance event induced by Pleistocene glaciation in the IP. During this period, large populations of thermophile species were fragmented into small isolated populations distributed in the warmest valleys and coastal regions. The further global warming that characterized the Holocene may then have allowed the recolonization of the entire IP, allowing secondary contacts between once isolated populations. This is consistent with demographic analyses suggesting the recent expansion of the Iberian populations of *C. albicans*. It also coincides with climate niche reconstruction, which shows that the potential distribution of the Iberian clades during the last glaciation maximum was limited to three or four refugia, similar to those reported for other animals and plants (Abellán and Svenning, 2014; Albert et al., 2007; Centeno-Cuadros et al., 2009; Ferrero et al., 2011; Gomez and Lunt, 2007; Gonçalves et al., 2015; Miraldo et al., 2011; Weiss and Ferrand, 2007). Although the output of the niche construction models should be interpreted cautiously, simulation methods have shown that Maxent results are robust to small sample size (Wisz et al., 2008). However, despite climatic

variables being particularly determinant for the ecology of desert ants (Amor et al., 2011; Boulay et al., 2017; Cerdá et al., 1997, 1998), the validation of the models requires data on temperature resistance and on individual and colony-level performance as a function of climate. Moreover, factors not accounted for by the models may affect ant distribution. For example, because *Cataglyphis* ants inhabit open habitats, variations in the proportion of forest ecosystems may thus have affected their distribution independently of temperature and precipitations.

Clade A is currently separated from clade C by rather high mountains that may prevent between-population gene flow. However, a potential contact zone exists along the Mediterranean coast where there is no geographic barrier apart from, perhaps, recent urbanization. Moreover, some of the collection points of clades B and C are separated by less than 10 km (C361 and C364), suggesting that a contact zone is likely to exist between these two clades. The processes that prevents the spatial co-occurrence of these recently diverged clades, despite the often common parapatric occurrence even among distant clades, is unclear. Even in the absence of evident habitat differences between sister clades, their prolonged isolation in distinct areas may promote the selection of cryptic adaptations to different micro-habitats. Once a secondary contact has been established, competition for resources may preclude the overlap of the previously isolated clades. For example, in *Cataglyphis* ants thermal preference and prey-size selection may induce high competition and could prevent the coexistence of sister clades. Competition may then be attenuated by the process of niche partitioning among more phylogenetically distant clades, allowing their sympatric distribution (Dillier and Wehner, 2004; Knaden and Wehner, 2006).

Important morphological and chemical differences exist among some of the identified clades. Morphological differences among males may strongly limit intra-clade mating and may be driven by a reinforcement-like process if potential hybrids at the contact zone have a lower fitness than their parents. Such a mechanism has already been proposed for *C. tartessica* and *C. floricola*, which occur in parapatry in southern Spain (Jowers et al., 2014). The selection forces that affect workers' cuticular chemistry are more complex to

disentangle. The primary function of CHC is that of resistance against desiccation. This function may be particularly important for thermophilous species that forage in extremely arid environments. Thus, we cannot exclude a possible convergence of the CHC profiles among clades E, F, and G due to their exploitation of similarly dry habitats. In addition, CHCs have acquired an important communication function and help in assessing whether encountered individuals are aliens or nestmates. To what extent this new function interferes with the clade-specific composition of CHC is unknown.

Our results also have important consequences for the taxonomy of this complex group of species. To date, it was considered that the IP *albicans* group comprises four species (*C. iberica*, *C. rosenhaueri*, *C. gadeai*, and *C. douwesi*; Cagniant 2009), but our phylogenetic trees recovered only three clades (A, B, and C). In addition, *C. iberica*, which was thought to occupy the great majority of the IP, appears to be polyphyletic: *C. iberica sensu stricto* is probably limited to the north-east of the IP, while black specimens distributed in the center and the south of the IP are more closely related to *C. rosenhaueri*, despite a marked difference in coloration. Our data also question whether *C. douwesi* and *C. rosenhaueri* are distinct species. Interestingly, although the analysis of microsatellite markers through DAPC suggests the existence of a gene flow between clades B and C, both clades have distinct chemotypes and male morphotypes, suggesting that they may constitute two different species. The taxonomic situation in NA is even more complex. Some species that have been described mostly based on differences of workers are probably para- or polyphyletic like the pairs *C. otini* and *C. cubica* and *C. espadaleri* and *C. theryi*. Therefore, the *C. albicans* group deserves a major taxonomic revision and a more intensive sampling in the remote regions of NA. In addition to extensive morphological measurements on all the castes, the use of chemical markers may help to characterize otherwise cryptic species (Bagnères and Wicker-Thomas, 2010; Dahbi et al., 2008, 1997).

In conclusion, our data point at dispersal and vicariance as two major mechanisms explaining the genetic divergence among *Cataglyphis* desert ants. This reaffirms the important role played by geographic isolation as a driver of new species formation, either

through drift or selection. By modulating gene flow among populations, past geo-climatic events have shaped the diversity of the western Mediterranean basin. Further analyses should be undertaken in order to determine the respective roles of genetic drift and natural selection in this speciation process. Such knowledge will help in predicting how future global warming might affect local and regional species richness in this major hotspot of biodiversity.

Acknowledgements

We thank David Aragonés (LAST- EBD) for Map creation, Naomi Paz for English editing assistance and Alberto Tinaut for interesting comments on a previous version of this manuscript. We also thank Elena Angulo and the Cerdá-Angulo family for their help during sample collection. This project was supported by grants PICS # 24698 from CNRS to RB, P12-RNM-2705 from Consejería de Economía y Conocimiento (Junta de Andalucía) to IV, RB and XC, Spanish Ministry of Economy and Competitiveness and FEDER (projects CGL2012-36181 and CGL2015-65807) to XC, RB, AH, AD and FA, and Spanish Ministry of Education (sabbatical stay SAB2010-0088) to AH. The University of Jyväskylä awarded an International mobility grant to JAG (893/13.00.04.01/2014).

References

- Abellán, P., Svenning, J.-C., 2014. Refugia within refugia—patterns in endemism and genetic divergence are linked to Late Quaternary climate stability in the Iberian Peninsula. *Biological Journal of the Linnean Society* 113, 13–28. doi:10.1111/bij.12309
- Abràmoff, M.D., Magalhães, P.J., Ram, S.J., 2004. Image processing with imageJ. *Biophotonics International*.
- Achalhi, M., Münch, P., Cornée, J.J., Azdimousa, A., Melinte-Dobrinescu, M., Quillévére, F., Drinia, H., Fauquette, S., Jiménez-Moreno, G., Merzeraud, G., Moussa, A. Ben, El Kharim, Y., Feddi, N., 2016. The late Miocene Mediterranean-Atlantic connections through the North Rifian Corridor: New insights from the Boudinar and Arbaa Taourirt basins (northeastern Rif, Morocco). *Palaeogeography, Palaeoclimatology,*

- Palaeoecology 459, 131–152. doi:10.1016/j.palaeo.2016.06.040
- Albert, E.M., Zardoya, R., García-París, M., 2007. Phylogeographical and speciation patterns in subterranean worm lizards of the genus *Blanus* (Amphisbaenia: Blanidae). *Molecular Ecology* 16, 1519–1531. doi:10.1111/j.1365-294X.2007.03248.x
- Amor, F., Ortega, P., Cerdá, X., Boulay, R., 2011. Solar Elevation Triggers Foraging Activity in a Thermophilic Ant. *Ethology* 117, 1031–1039. doi:10.1111/j.1439-0310.2011.01955.x
- Aron, S., Mardulyn, P., Leniaud, L., 2016. Evolution of reproductive traits in *Cataglyphis* desert ants: mating frequency, queen number and thelytoky. *Behavioral Ecology and Sociobiology* 70, 1367–1379.
- Bagnères, A.-G., Wicker-Thomas, C., 2010. Chemical taxonomy with hydrocarbons, in: Blomquist, G.J., Bagnères, A.G. (Eds.), *Insect Hydrocarbons: Biology, Biochemistry, and Chemical Ecology*. Cambridge University Press, Cambridge, pp. 121–161.
- Bandelt, H.J., Forster, P., Rohlf, A., 1999. Median-joining networks for inferring intraspecific phylogenies. *Molecular Biology and Evolution* 16, 37–48.
- Blaimer, B.B., Brady, S.G., Schultz, T.R., Lloyd, M.W., Fisher, B.L., Ward, P.S., 2015. Phylogenomic methods outperform traditional multi-locus approaches in resolving deep evolutionary history: a case study of formicine ants. *BMC Evolutionary Biology* 15, 271. doi:10.1186/s12862-015-0552-5
- Bouckaert, R., Heled, J., Kühnert, D., Vaughan, T., Wu, C.H., Xie, D., Suchard, M.A., Rambaut, A., Drummond, A.J., 2014. BEAST 2: A Software Platform for Bayesian Evolutionary Analysis. *PLoS Computational Biology* 10.
- Boulay, R., Aron, S., Cerdá, X., Doums, C., Graham, P., Hefetz, A., Monnin, T., 2017. Social Life in the desert: the case study of *Cataglyphis* ants. *Annual Review of Entomology* 62, in press.
- Brändli, L., Handley, L.J.L., Vogel, P., Perrin, N., 2005. Evolutionary history of the greater white-toothed shrew (*Crocidura russula*) inferred from analysis of mtDNA, Y, and X chromosome markers. *Molecular Phylogenetics and Evolution* 37, 832–844.

doi:10.1016/j.ympev.2005.06.019

- Cagniant, H., 2009. Le Genre *Cataglyphis* Foerster, 1850 au Maroc (Hyménoptères Formicidae). *Orsis* 24, 41–71.
- Carranza, S., Arnold, E.N., Pleguezuelos, J.M., 2006a. Phylogeny, biogeography, and evolution of two Mediterranean snakes, *Malpolon monspessulanus* and *Hemorrhois hippocrepis* (Squamata, Colubridae), using mtDNA sequences. *Molecular Phylogenetics and Evolution* 40, 532–546. doi:10.1016/j.ympev.2006.03.028
- Carranza, S., Harris, D.J., Arnold, E.N., Batista, V., Gonzalez De La Vega, J.P., 2006b. Phylogeography of the lacertid lizard, *Psammodromus algirus*, in Iberia and across the Strait of Gibraltar. *Journal of Biogeography* 33, 1279–1288. doi:10.1111/j.1365-2699.2006.01491.x
- Casimiro-Soriguer, R., Talavera, M., Balao, F., Terrab, A., Herrera, J., Talavera, S., 2010. Phylogeny and genetic structure of *Erophaca* (Leguminosae), a East-West Mediterranean disjunct genus from the Tertiary. *Molecular Phylogenetics and Evolution* 56, 441–450. doi:10.1016/j.ympev.2010.02.025
- Castella, V., Ruedi, M., Excoffier, L., Ibanez, C., Arlettaz, R., Hausser, J., 2000. Is the Gibraltar Strait a barrier to gene flow for the bat *Myotis myotis* (Chiroptera: Vespertilionidae)? *Molecular Ecology* 9, 1761–1772.
- Centeno-Cuadros, A., Delibes, M., Godoy, J.A., 2009. Phylogeography of Southern Water Vole (*Arvicola sapidus*): Evidence for refugia within the Iberian glacial refugia? *Molecular Ecology* 18, 3652–3667. doi:10.1111/j.1365-294X.2009.04297.x
- Cerdá, X., Retana, J., 2000. Alternative strategies by thermophilic ants to cope with extreme heat: individual versus colony level traits. *Oikos* 89, 155–163. doi:10.1034/j.1600-0706.2000.890117.x
- Cerdá, X., Retana, J., Cros, S., 1997. Thermal disruption of transitive hierarchies in Mediterranean ant communities. *Journal of Animal Ecology* 66, 363–374. doi:doi:10.2307/5982
- Cerdá, X., Retana, J., Manzaneda, A., 1998. The role of competition by dominants and

temperature in the foraging of subordinate species in mediterranean ant communities.

Oecologia 117, 404–412. doi:doi:10.1007/s004420050674

Chéron, B., Monnin, T., Fédérici, P., Doums, C., 2011. Variation in patriline reproductive success during queen production in orphaned colonies of the thelytokous ant

Cataglyphis cursor. Mol. Ecol. 20(8), 2011–2022. doi:doi:10.1111/j.1365-294X.2011.05075.x

Coyne, J.A., Orr, H.A., 2004. Speciation. Sinauer Associates Inc, Sunderland, MA.

Dahbi, A., Cerdá, X., Hefetz, A., Lenoir, A., 1997. Adult transport in the ant *Cataglyphis iberica*: a means to maintain a uniform colonial odour in a species with multiple nests.

Physiol. Entomol. 22, 13–19. doi:DOI:10.1111/j.1365-3032.1997.tb01135.x

Dahbi, A., Hefetz, A., Lenoir, A., 2008. Chemotaxonomy of some *Cataglyphis* ants from Morocco and Burkina Faso. Biochemical Systematics and Ecology 36, 564–572.

doi:10.1016/j.bse.2008.03.004

Dahbi, A., Lenoir, A., Tinaut, A., Taghizadeh, T., Francke, W., Hefetz, A., 1996. Chemistry of the postpharyngeal gland secretion and its implication for the phylogeny of

Iberian *Cataglyphis* species (Hymenoptera: Formicidae). Chemoecology 7, 163–171.

doi:10.1007/BF01266308

Dillier, F.-X., Wehner, R., 2004. Spatio-temporal patterns of colony distribution in monodomous and polydomous species of North African desert ants, genus *Cataglyphis*.

Insectes Sociaux 51, 186–196. doi:10.1007/s00040-003-0722-0

Durand, E., Chen, C., Francois, O., 2009. Tess version 2.3 - Reference Manual August 2009

*. Available at: memberstimc. imag. fr/Olivier. Francois/tess. html 1–30.

Eyer, P.A., Seltzer, R., Hefetz, A., 2017. Molecular Phylogenetics and Evolution An

integrative approach to untangling species delimitation in the *Cataglyphis bicolor* desert ant complex in Israel. Molecular Phylogenetics and Evolution 115, 128–139.

doi:10.1016/j.ympev.2017.07.024

Ferrero, M.E., Blanco-Aguiar, J.A., Loughheed, S.C., Sánchez-Barbudo, I., De Nova, P.J.G.,

Villafuerte, R., D??vila, J.A., 2011. Phylogeography and genetic structure of the red-

- legged partridge (*Alectoris rufa*): More evidence for refugia within the Iberian glacial refugium. *Molecular Ecology* 20, 2628–2642. doi:10.1111/j.1365-294X.2011.05111.x
- Gantenbein, B., Largiadèr, C.R., 2003. The phylogeographic importance of the Strait of Gibraltar as a gene flow barrier in terrestrial arthropods: A case study with the scorpion *Buthus occitanus* as model organism. *Molecular Phylogenetics and Evolution* 28, 119–130. doi:10.1016/S1055-7903(03)00031-9
- Garcia-Mudarra, J.L., Ibañez, C., Juste, J., 2009. The Straits of Gibraltar: Barrier or bridge to Ibero-Moroccan bat diversity? *Biological Journal of the Linnean Society* 96, 434–450. doi:10.1111/j.1095-8312.2008.01128.x
- Gomez, A., Lunt, D.H., 2007. Refugia within refugia: patterns of phylogeographic concordance in Iberian Peninsula., in: Weiss, S., Ferrand, N. (Eds.), *Phylogeography of Southern European Refugia*. Springer, pp. 155–188.
- Gonçalves, H., Maia-Carvalho, B., Sousa-Neves, T., Garcia-Paris, M., Sequeira, F., Ferrand, N., Martínez-Solano, I., Gonçalves, H., Maia-Carvalho, B., Sousa-Neves, T., García-París, M., Sequeira, F., Ferrand, N., Martínez-Solano, I., 2015a. Multi locus phylogeography of the common midwife toad, *Alytes obstetricans* (Anura, Alytidae): Contrasting patterns of lineage diversification and genetic structure in the Iberian refugium. *Molecular Phylogenetics and Evolution* 93, 363–379. doi:10.1016/j.ympev.2015.08.009
- Hewitt, G., 1999. Post-glacial re-colonization of European biota. *Biological Journal of the Linnean Society* 68, 87–112. doi:10.1111/j.1095-8312.1999.tb01160.x
- Hewitt, G.M., 2004. Genetic consequences of climatic oscillations in the Quaternary. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences* 359, 183–195; discussion 195. doi:10.1098/rstb.2003.1388
- Husemann, M., Schmitt, T., Zachos, F.E., Ulrich, W., Habel, J.C., 2014. Palaeartic biogeography revisited: Evidence for the existence of a North African refugium for Western Palaeartic biota. *Journal of Biogeography* 41, 81–94. doi:10.1111/jbi.12180
- Igea, J., Aymerich, P., Fernández-González, A., González-Esteban, J., Gómez, A., Alonso,

- R., Gosálbez, J., Castresana, J., 2013. Phylogeography and postglacial expansion of the endangered semi-aquatic mammal *Galemys pyrenaicus*. *BMC evolutionary biology* 13, 115. doi:10.1186/1471-2148-13-115
- Jakobsson, M., Rosenberg, N.A., 2007. CLUMPP: A cluster matching and permutation program for dealing with label switching and multimodality in analysis of population structure. *Bioinformatics* 23, 1801–1806. doi:10.1093/bioinformatics/btm233
- Jombart, T., Devillard, S., Balloux, F., 2010. Discriminant analysis of principal components: a new method for the analysis of genetically structured populations. *BMC Genetics* 11, 94. doi:doi: 10.1186/1471-2156-11-94
- Jowers, M.J., Amor, F., Ortega, P., Lenoir, A., Boulay, R.R., Cerdá, X., Galarza, J.A., 2014. Recent speciation and secondary contact in endemic ants. *Molecular Ecology* 23, 2529–2542. doi:10.1111/mec.12749
- Jowers, M.J., Leniaud, L., Cerdà, X., Alasaad, S., Caut, S., Amor, F., Aron, S., Boulay, R.R., 2013. Social and Population Structure in the Ant *Cataglyphis emmae*. *PLoS ONE* 8. doi:10.1371/journal.pone.0072941
- Jowers, M.J., Taheri, A., Reyes-Lopez, J., 2015. The ant *Anochetus ghilianii* (Hymenoptera, Formicidae), not a Tertiary relict, but an Iberian introduction from North Africa: Evidence from mtDNA analyses. *Systematics and Biodiversity* 13, 865–874. doi:10.1080/14772000.2015.1061065
- Kaliontzopoulou, A., Pinho, C., Harris, D.J., Carretero, M.A., 2011. When cryptic diversity blurs the picture: A cautionary tale from Iberian and North African *Podarcis* wall lizards. *Biological Journal of the Linnean Society* 103, 779–800. doi:10.1111/j.1095-8312.2011.01703.x
- Katoh, K., Misawa, K., Kuma, K., Miyata, T., 2002. MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic acids research* 30, 3059–3066. doi:10.1093/nar/gkf436
- Kearse, M., Moir, R., Wilson, A., Stones-Havas, S., Cheung, M., Sturrock, S., Buxton, S., Cooper, A., Markowitz, S., Duran, C., Thierer, T., Ashton, B., Meintjes, P., Drummond,

- A., 2012. Geneious Basic: An integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 28, 1647–1649. doi:10.1093/bioinformatics/bts199
- Knaden, M., Tinaut, A., Stökl, J., Cerdá, X., Wehner, R., 2012. Molecular phylogeny of the desert ant genus *Cataglyphis* (Hymenoptera: Formicidae). *Myrmecol. News* 16, 123–132.
- Knaden, M., Wehner, R., 2006. Fundamental difference in life history traits of two species of *Cataglyphis* ants. *Frontiers in zoology* 3, 21. doi:10.1186/1742-9994-3-21
- Lanfear, R., Calcott, B., Ho, S.Y.W., Guindon, S., 2012. PartitionFinder: Combined selection of partitioning schemes and substitution models for phylogenetic analyses. *Molecular Biology and Evolution* 29, 1695–1701. doi:10.1093/molbev/mss020
- Lefebvre, T., Vargo, E.L., Zimmermann, M., Dupont, S., Kutnik, M., Bagnères, A.G., 2016. Subterranean termite phylogeography reveals multiple postglacial colonization events in southwestern Europe. *Ecology and Evolution* 6, 5987–6004. doi:10.1002/ece3.2333
- Librado, P., Rozas, J., 2009. DnaSP v5: A software for comprehensive analysis of DNA polymorphism data. *Bioinformatics* 25, 1451–1452. doi:10.1093/bioinformatics/btp187
- Mayr, E., 1963. *Animal Species and Evolution*. Belknap, Cambridge.
- Miraldo, A., Hewitt, G.M., Paulo, O.S., Emerson, B.C., 2011a. Phylogeography and demographic history of *Lacerta lepida* in the Iberian Peninsula: multiple refugia, range expansions and secondary contact zones. *BMC evolutionary biology* 11, 170. doi:10.1186/1471-2148-11-170
- Miraldo, A., Hewitt, G.M., Paulo, O.S., Emerson, B.C., 2011b. Phylogeography and demographic history of *Lacerta lepida* in the Iberian Peninsula: multiple refugia, range expansions and secondary contact zones. *BMC evolutionary biology* 11, 170. doi:10.1186/1471-2148-11-170
- Moreau, C.S., Bell, C.D., 2013. Testing the museum versus cradle tropical biological diversity hypothesis: phylogeny, diversification, and ancestral biogeographic range evolution of the ants. *Evolution* 67, 2240–2257. doi:10.1111/evo.12105

- Myers, N., Mittermeier, R.A., Mittermeier, C.G., Fonseca, G.B.A., Kent, J., 2000. Biodiversity hotspots for conservation priorities. *Nature* 403, 72.
- Nosil, P., 2012. *Ecological Speciation*. Oxford University Press, Oxford, United Kingdom.
- Nosil, P., 2008. Speciation with gene flow could be common. *Molecular ecology* 17, 2006–2008. doi:10.1186/1742-9994-4-11.Schmitt
- Pardo, C., Cubas, P., Tahiri, H., 2008. Genetic variation and phylogeography of *Stauracanthus* (Fabaceae, Genisteae) from the Iberian Peninsula and northern Morocco assessed by chloroplast microsatellite (cpSSR) markers. *American Journal of Botany* 95, 98–109. doi:10.3732/ajb.95.1.98
- Paulo, O.S., Pinheiro, J., Miraldo, A., Bruford, M.W., Jordan, W.C., Nichols, R.A., 2008. The role of vicariance vs. dispersal in shaping genetic patterns in ocellated lizard species in the western Mediterranean. *Molecular Ecology* 17, 1535–1551. doi:10.1111/j.1365-294X.2008.03706.x
- Pearcy, M., Clemencet, J., Chameron, S., Aron, S., Doums, C., 2004. Characterization of nuclear DNA microsatellite markers in the ant *Cataglyphis cursor*. *Molecular Ecology Notes* 4, 642–644. doi:10.1111/j.1471-8286.2004.00759.x
- Philips, S.J., Andersen, R.P., Schapire, R.E., 2006. Maximum entropy modeling of species geographic distributions. *Ecological modelling* 190, 231–259.
- Philips, S.J., Dubik, M., Schapire, R.E., 2004. A maximum entropy approach to species distribution modeling, in: *Proceedings of the 21st International Conference on Machine Learning*. ACM Press, New York, pp. 655–662.
- Polzin, T., Vahdati Daneshmand, S., 2003. On Steiner trees and minimum spanning trees in hypergraphs. *Operations Research Letters* 31, 12–20.
- Radchnko, A.G., 1997. Review of ants of the genus *Cataglyphis* Foerster (Hymenoptera, Formicidae) of Asia. *Entomologicheskoe Obozrenie* 76, 424–442.
- R Development Core Team, 2015a. *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing Vienna Austria. doi:10.1038/sj.hdy.6800737

- R Development Core Team, 2015b. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing Vienna Austria.
doi:10.1038/sj.hdy.6800737
- Rambaut, A., Drummond, A., 2007. BEAUTi v1.4.2. Bayesian Evolutionary Analysis Utility.
- Rambaut, A., Drummond, A.J., 2013. Tracer v1.6. Available from
<http://tree.bio.ed.ac.uk/software/tracer/>.
- Ruano, F., Devers, S., Sanllorente, O., Errard, C., Tinaut, A., Lenoir, A., 2011. A geographical mosaic of coevolution in a slave-making host-parasite system. *Journal of Evolutionary Biology* 24, 1071–1079. doi:10.1111/j.1420-9101.2011.02238.x
- Salzburger, W., Martens, J., Sturmbauer, C., 2002. Paraphyly of the Blue Tit (*Parus caerulescens*) suggested from cytochrome b sequences. *Molecular Phylogenetics and Evolution* 24, 19–25.
- Sanchez-Robles, J.M., Balao, F., Terrab, A., Garcia-Castaño, J.L., Ortiz, M.A., Vela, E., Talavera, S., 2014. Phylogeography of SW Mediterranean firs: Different European origins for the North African *Abies* species. *Molecular Phylogenetics and Evolution* 79, 42–53. doi:10.1016/j.ympev.2014.06.005
- Sanllorente, O., Ruano, F., Tinaut, A., 2015. Large-scale population genetics of the mountain ant *Proformica longiseta* (Hymenoptera: Formicidae). *Population Ecology* 57, 637–648. doi:10.1007/s10144-015-0505-2
- Schlick-Steiner, B.C., Steiner, F.M., Moder, K., Seifert, B., Sanetra, M., Dyreson, E., Stauffer, C., Christian, E., 2006. A multidisciplinary approach reveals cryptic diversity in Western Palearctic *Tetramorium* ants (Hymenoptera: Formicidae). *Molecular Phylogenetics and Evolution* 40, 259–273. doi:<http://dx.doi.org/10.1016/j.ympev.2006.03.005>
- Seifert, B., 2009. Cryptic species in ants (Hymenoptera: Formicidae) revisited: we need a change in the alpha-taxonomic approach. *Myrmecological News* 12, 149–166.
- Taberlet, P., Fumagalli, L., Wust-Saucy, A.-G., Cosson, J.-F., 1992. Comparative phylogeography and postglacial colonization routes in Europe. *Molecular Ecology* 453–464.

- Tinaut, A., 1990. Situacion Taxonomica del genero *Cataglyphis* Förster, 1850 en la Peninsula iberica. III. El grupo de *C. velox* Santschi, 1929 y descripcion de *Cataglyphis humeya* sp. n. (Hymenoptera, Formicidae). *Eos* 66, 215–227.
- Velo-Anton, G., Godinho, R., Harris, D.J., Santos, X., Martinez-Freiria, F., Fahd, S., Larbes, S., Pleguezuelos, J.M., Brito, J.C., 2012. Deep evolutionary lineages in a Western Mediterranean snake (*Vipera latastei/monticola* group) and high genetic structuring in Southern Iberian populations. *Molecular Phylogenetics and Evolution* 65, 965–973. doi:10.1016/j.ympev.2012.08.016
- Wehner, R., Marsh, A.C., Wehner, S., 1992. Desert ants on a thermal tightrope. *Nature* 357, 586–587.
- Weiss, S., Ferrand, N., 2007. *Phylogeography of Southern European Refugia*. Springer, Dordrecht, The Netherlands.
- Wiens, J.J., Ackerly, D.D., Allen, A.P., Anacker, B.L., Buckley, L.B., Cornell, H. V., Damschen, E.I., Jonathan Davies, T., Grytnes, J.-A., Harrison, S.P., Hawkins, B.A., Holt, R.D., McCain, C.M., Stephens, P.R., 2010. Niche conservatism as an emerging principle in ecology and conservation biology. *Ecology Letters* 13, 1310–1324. doi:10.1111/j.1461-0248.2010.01515.x
- Wisz, M.S., Hijmans, R.J., Li, J., Peterson, A.T., Graham, C.H., Guisan, A., Elith, J., Dudík, M., Ferrier, S., Huettmann, F., Leathwick, J.R., Lehmann, A., Lohmann, L., Loiselle, B.A., Manion, G., Moritz, C., Nakamura, M., Nakazawa, Y., Overton, J.M., Phillips, S.J., Richardson, K.S., Scachetti-Pereira, R., Schapire, R.E., Soberón, J., Williams, S.E., Zimmermann, N.E., 2008. Effects of sample size on the performance of species distribution models. *Diversity and Distributions* 14, 763–773. doi:10.1111/j.1472-4642.2008.00482.x
- Wojcieszek J.M., Simmons L.W. (2013) Divergence in genital morphology may contribute to mechanical reproductive isolation in a millipede. *Ecology and Evolution* 3, 334-343.

Table 1: Primer information

Primer name	5'-3' primer sequence	Tm	Reference
LWRhF	AATTGCTATTAYGARACNTGGGT	52°C	^a
LWRhR	ATATGGAGTCCANGCCATRAACCA	52°C	^a
For28SVesp	AGAGAGAGTTCAAGAGTACGTG	54°C	^b
Rev28SVesp	GGAACCAGCTACTAGATGG	54°C	^b
COIFdg	ARTTAAGADCTTGYYGGATCATT	52°C	this study
COIRdg	RATGTTGRATAAGAATAGGRTCHC	52°C	this study

Tm: annealing temperature

^a Danforth et al., 2004

^b Hines et al., 2007

Table 2: Best model of evolution for each partition based on the Bayesian Information Criteria.

Partitions	Best model
CO1_2; LWRh_1; LWRh_2	HKY+I
LWRh_3	K80+G
CO1_1; LWRh intron	HKY+I+G
28S	TrN+I+G
CO1_3	TrN+G

Table 3: Measures of diversity of the 3 loci for the 8 major clades of the *albicans* group in the IP (clades A to C) and Morocco (clades D to H).

Clade	Locus	N	L	S	Nh	Hd	Pi	D	D*
A	LWRh	14	604	1	2	0.36	0.001	0.324	0.716
	28S	14	782	1	2	0.44	0.001	0.842	0.716
	COI	7	503	23	7	1.00	0.015	-1.236	-1.351
B	LWRh	22	604	3	3	0.33	0.001	-0.750	-0.159
	28S	22	782	2	3	0.26	0.001	-1.175	-0.629
	COI	11	503	25	10	0.98	0.017	-0.078	-0.200
C	LWRh	102	604	10	10	0.69	0.003	-0.409	-1.452
	28S	102	782	5	6	0.25	0.001	-1.533	1.035
	COI	51	503	45	33	0.97	0.011	-1.648	-1.172
D	LWRh	28	604	3	4	0.56	0.001	-0.547	-1.440
	28S	28	782	3	4	0.64	0.001	0.038	0.959
	COI	14	503	45	12	0.98	0.029	0.019	0.167
E	LWRh	46	604	4	3	0.46	0.002	0.108	-1.164
	28S	46	782	5	8	0.82	0.002	0.862	0.155
	COI	23	503	25	16	0.96	0.009	-1.234	-1.608
F	LWRh	8	604	1	2	0.43	0.001	0.334	0.888
	28S	8	782	1	2	0.43	0.001	0.334	0.888
	COI	4	503	0	1	0	na	na	na
G	LWRh	22	604	10	6	0.54	0.004	-0.945	0.428
	28S	22	782	4	4	0.46	0.001	-0.537	0.143
	COI	11	503	81	11	1.00	0.052	-0.798	-0.647
H	LWRh	16	604	10	7	0.79	0.003	-1.193	-1.667
	28S	16	782	2	3	0.70	0.001	1.262	0.907
	COI	8	503	52	8	1.00	0.041	-0.136	0.085

N: Number of sequences used; L : sequence length; S: Number of variable sites; Nh: Number of haplotypes; Hd : Haplotype diversity ; Pi: Nucleotide diversity ; D: Tajima's D test statistic; D*: Fu and Li's D* test statistic.

Table 4: Descriptive statistics obtained from the microsatellite analyses.

Clade	N	Na	PA	Ne	Ho	He	Fst
A	7	5.5±0.9	5	4.12±0.81	0.47±0.08	0.59±0.1	0.17±0.06
B	11	7.1±1.3	1	5.12±0.93	0.51±0.09	0.68±0.09	0.29±0.09
C	51	15.2±2.3	28	8.87±1.71	0.62±0.08	0.76±0.09	0.21±0.04
D	14	8.2±1.6	22	5.44±1.18	0.48±0.08	0.70±0.09	0.33±0.06
E	23	9.9±1.6	6	5.84±0.93	0.48±0.07	0.73±0.09	0.34±0.06
F	4	4.3±0.7	3	3.71±0.72	0.59±0.10	0.64±0.08	0.10±0.08
G	11	7.9±1.1	3	5.95±1.01	0.52±0.08	0.73±0.08	0.28±0.08
H	7	6.6±0.8	4	4.90±0.72	0.53±0.09	0.71±0.08	0.25±0.09

N: number of genotyped specimens; Na: Mean±SE number of alleles; PA: Number of private alleles; He: Mean±SE number of effective alleles Ho and He: Observed and expected Heterozygosity (± SE); Fst: Fixation index (±SE).

Figure captions

Figure 1: Palaeogeographic map of Gibraltar Straits region showing the progressive formation of a land bridge between the North of Africa and the Iberian Peninsula. A: Early Tortonian (< 8 Mya), B: Late Tortonian (8 – 7.2 Mya), C: Early Messinian (7.2 – 6.1 Mya), D: Late Messinian (>6.1 Mya) (modified from Achalhi *et al.* 2016).

Figure 2: Phylogram obtained by Bayesian Inference for 130 samples of *C. albicans* group from the Iberian Peninsula and North Africa. Probable species names are given on the right side of the tree together a picture of one specimen. The 8 clades are denominated by letters A to H. Branch length denote time in Million Years. Messinian Salinity Crisis is represented by a red line. The T and M arrows on the x axis indicate the beginning of the Tortonian and Messinian, respectively. Numbers near the main nodes give their support.

Figure 3: Geographic distribution of the 8 clades identified by Bayesian Inference.


Figure 4: Phylogram obtained by Maximum Likelihood for 130 samples of *C. albicans* group from the Iberian Peninsula and North Africa. Letters A to H correspond to clades obtained by Bayesian Inference (Fig. 1). Numbers near the main nodes give their percentage of support from the ML.

Figure 5: Linear Discriminant Analysis of principal components conducted on sequences (a), microsatellite data (b), cuticular hydrocarbons (c) and morphological data (d).

Figure 6: Bayesian skyline plots showing effective population size fluctuation through time for the Iberian (A) and North-African (B) groups. The centre line shows the median estimate; upper and lower confidence limits (95% posterior density) are indicated by filled colour.

Figure 7: Results of the population structure analysis in TESS using microsatellites. Models with and without admixture are presented and suggest a maximum of 7 and 5 clusters respectively (as indicated by the sharp reduction of the slope of the DIC).

Figure 8: Models of distribution of the most favorable climatic conditions for each of the 8 clades identified by Bayesian Inference. Maps on the right show the current distribution while maps on the left show the potential distribution during the Last Glacial Maximum. Darker brown colors denote higher probability of presence.











Figure 3:












ACCEPTED



- Nine clades of desert ants (albicans group) exist in the Iberian Peninsula and Morocco.
- Iberian species diverged from their african ancestor after the Gibraltar Straits last opening
- Several refugia of desert ants occurred in the Iberian Peninsula during the last Glaciation
- Iberian clades are parapatric and populations are in expansion

Graphical abstract

