

**This is an electronic reprint of the original article.
This reprint *may differ* from the original in pagination and typographic detail.**

Author(s): Mpamah, Promise; Taipale, Sami; Rissanen, Antti Juhani; Biasi, Christina; Nykänen, Hannu

Title: The impact of long-term water level draw-down on microbial biomass : A comparative study from two peatland sites with different nutrient status

Year: 2017

Version:

Please cite the original version:

Mpamah, P., Taipale, S., Rissanen, A. J., Biasi, C., & Nykänen, H. (2017). The impact of long-term water level draw-down on microbial biomass : A comparative study from two peatland sites with different nutrient status. *European Journal of Soil Biology*, 80, 59-68. <https://doi.org/10.1016/j.ejsobi.2017.04.005>

All material supplied via JYX is protected by copyright and other intellectual property rights, and duplication or sale of all or part of any of the repository collections is not permitted, except that material may be duplicated by you for your research use or educational purposes in electronic or print form. You must obtain permission for any other use. Electronic or print copies may not be offered, whether for sale or otherwise to anyone who is not an authorised user.

The impact of long-term water level draw-down on microbial biomass: a comparative study from two peatland sites with different nutrient status

Promise Anyalewachi Mpamah^{1*}, Sami Taipale^{1,2}, Antti Juhani Rissanen³, Christina Biasi² and Hannu Kalevi Nykänen^{1,2}

¹University of Jyväskylä, Department of Biological and Environmental Science, Jyväskylä, Finland.

²University of Eastern Finland, Department of Environmental Science, Finland

³Tampere University of Technology, Laboratory of Chemistry and Bioengineering, Tampere, Finland.

* Corresponding author; Promise Anyalewachi Mpamah

Tel: +358449730977; Email: promise.a.mpamah@jyu.fi

Sami Taipale; Email: sami.taipale@jyu.fi

Antti Juhani Rissanen; Email: antti.rissanen@tut.fi

Christina Biasi; Email: christina.biasi@uef.fi

Hannu Kalevi Nykänen; Email: hannu.nykanen@uef.fi

1 **ABSTRACT**

2 We examined the effects of long-term (51 years) drainage on peat microbial communities using
3 phospholipid fatty acid (PLFA) analysis. We analysed the peat profiles of natural and adjacent drained
4 fen and bog sites. Viable microbes (i.e. microbial PLFA) were present in relatively large amounts
5 even in the deepest peat layers of both peatland sites, a finding that warrants further investigation.
6 Microbial biomass was generally higher in the fen than in the bog. Microbial community structure
7 (indexed from PLFA) differed between the fen and bog sites and among depths. Although we did not
8 exclude other factors, the effect of drainage on the total microbial biomass and community structure
9 was not limited to the surface layers, but extended to the deepest layers of the fen and bog. Long-term
10 drainage increased the total microbial PLFA biomass in the surface, subsurface and bottom layers of
11 the fen, but decreased it in the surface and bottom layers of the bog site. Drainage also increased the
12 characteristic FAs of Gram-positive and Gram-negative bacteria in the surface and subsurface layers
13 of the fen, and decreased them in the bottom layers of the bog site. The characteristic fungal FA was
14 only reduced in the surface layers of the bog site by drainage. Thus, by affecting the microbial
15 community beyond the surface layers, long-term peatland water-level draw-down can alter the
16 microbial contribution to deeper peat organic matter stabilization. This suggests that long-term
17 drainage may have a more significant climate change effect than revealed by the surface layer
18 analyses alone.

19

20 **Key words:** Fen; Bog; PLFA; long-term drainage; microbial biomass; Microbial community
21 structure.

22

23 **1. INTRODUCTION**

24

25 Peatlands are crucial global carbon (C) stores [1,2], containing about 15 – 30% of all terrestrial
26 organic C (OC); equivalent to 455 Gt (10^{15} g) C [1]. Microbes are key actors (as catalysts) in all peat

27 biogeochemical processes, controlling the peat OC accumulation and decomposition [3]. They also
28 contribute to the peat C exchange via respiration and, upon cell death, necromass addition to the peat
29 soil organic matter (SOM), via the microbial carbon pump (MCP; [4]). Different microbial groups,
30 with complementary enzymatic activities and different responses to environmental variables, interact
31 in the peat C-cycling processes [5]. For example, the Gram-negative and Gram-positive bacteria are
32 mainly associated with the mineralization steps involving labile and more recalcitrant C materials,
33 respectively [6], and the exo-enzymatic capabilities of fungi make them important in the
34 decomposition of macromolecules and recalcitrant C materials [7,8]. Changes in climate factors, such
35 as hydrology, affect microbial community biomass and activity, both spatially and temporarily [9,10].

36 Models predict a warmer global climate (average temperature increases of about 4°C) up to the
37 year 2100 [11] and, under these scenarios, increased evapotranspiration due to increased temperatures
38 would lead to a lower water table (WT) in peatlands [12]. Persistent draw-down of the peatland WT
39 affects the niches of peatland microbes by increasing the thickness of the aerated surface layer [10].
40 The impacts of changed hydrology on the microbial community depends on the peatland type,
41 intensity of change and the extent of change in space and time [9,13-15]. While changes in microbial
42 niches may lead to increased diversity in the short-term, repeated replacement of specialist by
43 generalist microbes may lead to loss of diversity in the long-run [14]. Changes in plant species cover
44 following water level draw-down also modifies the influence of temperature and water content on
45 peat microbial activities [16]. Studies have suggested that drainage could increase or decrease total
46 microbial biomass or the biomass of some microbial groups, depending on the peatland type and
47 depth [9,13]. Jaatinen et al [13] showed that fungi and actinobacteria suffer from drainage in a
48 nutrient-rich fen, but that in a drained bog, while fungi either suffer or benefit, actinobacteria
49 abundance remains the same or increases. Fungi and bacteria generally benefit (undergo biomass
50 increase) from persistent drainage of wet mesotrophic fen sites, though actinobacteria suffer or show
51 only minor responses [9,15]. Changes in peat C accumulation and decomposition activities following
52 drainage have also been related to changes in the structure of below-ground microbial communities

53 [5,16]. Altered microbial diversity, due to drainage-induced changes in the quality and quantity of
54 OC inputs, coupled with better oxygen availability, could increase the rate of soil OC (SOC) cycling;
55 this leads to changes in the balance of peat-atmosphere C exchange [5,10,17].

56 Phospholipid-derived fatty acids (PLFA) are reliable quantitative biomarkers of viable microbes,
57 since they are short-lived and readily metabolized upon cell death. Microbial biomasses, community
58 structure and community responses to changing peat hydrology at different sites, have been studied
59 with PLFAs [9,13,18]. Differences in microbial community structures between peatlands and the
60 effects of treatments (e.g drainage) have also been analysed in several studies based on PLFAs alone
61 [13,18-21], and their results are similar to those obtained using other molecular methods [22].

62 To our knowledge, previous studies on the effects of drainage on microbial communities (like
63 those mentioned above), focused on the upper layers of peatlands (e.g. [13], [22]). However,
64 drainage-induced increases in oxygenation coupled with temperature changes in the surface layers
65 could prompt dissolved OC (DOC) release to deeper depths via the “enzymatic latch” process [23].
66 This increase in the flow and lability of DOC [24], coupled with deeper deposition of labile root
67 exudates by roots of vascular plants [25-27], could modify the biomass of microbial communities and
68 their composition in deeper peatland layers. Recent molecular evidence [17] and higher bulk peat
69 stable C isotope ($\delta^{13}\text{C}$) values, indicating peat degradation in the bottom of drained peat, supports this
70 view [17,28].

71 This study examined the effect of long-term drainage on microbial communities in depth profiles
72 from surface to bottom layers of two peatland sites differing in nutrient status. We compared the
73 biomasses and structures of the microbial communities (indexed by total and relative abundance of
74 PLFA, respectively) between the natural and drained sides of fen and bog, representing boreal
75 peatlands of different fertility after 51 years of water level draw-down. We also specifically
76 investigated the effects of drainage on some selected microbial groups. We hypothesised that long-
77 term WT decrease will (1) increase microbial biomass and (2) influence the microbial community

78 structure in the deep anoxic layers. We also hypothesized that (3) there will be higher microbial
79 biomass increase in the fen than in the bog site, due to long-term drainage.

80

81 **2. MATERIAL AND METHODS**

82

83 2.1. Study sites

84

85 The study was conducted at two peatland sites (one fen and one bog) within the Lakkasuo boreal
86 mire complex (61°47'N, 24°18'E, ca.150 m a.s.l.), in the Orivesi area in central Finland. At the
87 nearest weather station to the sites in Juupajoki Hyytiälä (61°85'N, 24°29'E) the mean annual
88 temperature was 3.5 °C and precipitation 711 mm for the period of 1981–2010 [29]. The sampling
89 year was wetter than this long-term mean, with whole year precipitation of 907 mm and an average
90 temperature of 3.2 °C. The mire complex comprises a large variety of typical Finnish mire site types
91 [30]. Part of the Lakkasuo peatland was ditch-drained in 1961 (51 years before sampling) so that there
92 are adjacent natural and drained sides of different fertility along a border ditch (Fig. S1). There were
93 differences in the original fertility, WT and vegetation composition between the natural ombrotrophic
94 cotton grass pine bog with *Sphagnum fuscum* hummocks (bog) and the natural minerotrophic tall
95 sedge fen (fen) sites sampled. Drainage caused marked changes to the hydrology, peat and vegetation
96 properties, carbon dioxide (CO₂) and CH₄ fluxes, especially at the drained fen ([28,30-33];
97 summarized in Table 1). For example, the six-month average WTs before the sampling date were -
98 8.0 and -34.9 cm for the natural and drained fens, respectively, whereas it was -12.0 and -16.4 cm for
99 the natural and drained bogs, respectively (Fig. 1). CO₂ fluxes increased in both sites whereas CH₄
100 fluxes ceased in the fen and were reduced by half in the bog after 30–32 years of drainage. Thus,
101 there is strong evidence for significant, long-term changes in peat characteristics and greenhouse gas
102 fluxes. The pH increased from the surface downwards in the natural and drained sides of both sites
103 (Fig. S2). In general, the bog site is more acidic than the fen site and this was confirmed by previously

104 reported pH values (Fig. S2 & S3). Although temperatures vary seasonally, the temperature in deep
105 peat is rather constant (~ 6 - 8 °C). The bulk densities (BD) at different depths of the drained and
106 natural sides are the same in the bog site, but different in the fen site (Fig. S2).

107

108 2.2. Soil sampling and water table level measurement

109

110 2.2.1. Initial soil sampling and water table measurement

111

112 In 2012 (November 22nd), three replicate sets of peat samples were collected from random points
113 within each site, located several meters apart along a 50 m boardwalk. Soil was sampled from 4–5
114 depths (0 – 25 cm, 25 – 50 cm, 50 – 100 cm and deepest 25 cm) starting at the surface and extending
115 to the deepest layer above the mineral soil. Using a Russian pattern side-cutting sampler (5 x 50 cm;
116 [34,35]), samples were collected in segments along the profile from both the drained and adjacent
117 natural sides. The samples were put into polyethylene bags, mixed and cooled immediately after
118 collection in a box with crushed ice, and later stored at -20 °C until analysis. Part of the samples were
119 oven dried and ground into a fine powder for analysis of their C and N content (Flash EA 1112
120 elemental analyser, Thermo Finnigan), with a certified birch leaf standard (Elementar Microanalysis,
121 UK) used as a reference. Continuous (3-hourly) WT level measurements were recorded with an
122 automatic WT-HR 64K logger (Fig. 1). The logger values were calibrated by manual measurements.

123

124 2.2.2. Additional peat properties measurements

125

126 Volumetric samples (from the same depths as the initial samples) were used for pH and
127 temperature measurements, as well as for bulk density determination (Fig. S2). Sampling was done
128 with a similar, but smaller, Russian pattern side-cutting sampler to that described above (5.2 * 50 cm;
129 half cylinder diameter * length) on 14 October, 2015. Sampling and depth measurements were started

130 under a living *Sphagnum* carpet. Samples were transferred from the sampler into plastic bags
131 (Aromata, Lidl Stiftung & Co, Neckarsulm, Germany) and were mixed in the bag before insertion of
132 a pH electrode coupled with a temperature sensor (WTW P3 pH/conductivity with electrode SenTix
133 41; Weilheim, Germany). Values of pH and temperature were recorded after one minute. To
134 determine bulk density, the samples in the bags were dried in the oven (Memmert, UM 500,
135 Schwabach, Germany) at 80°C until there was no change in the dry weight.

136

137 2.3. PLFA analysis

138

139 2.3.1. PLFA extraction and quantification

140

141 PLFA analysis was done following the protocol used by Tavi et al. [36] with some modifications.
142 Freeze-dried and mixed peat samples from each depth profile were weighed into 50 ml extraction
143 tubes (> 3 g dry weight of peat) using tools cleaned with methanol. Total lipids were extracted from
144 the samples using a 1:2:0.8 (vol:vol:vol) ratio of chloroform–methanol–50 mM phosphate buffer [37].
145 Tubes were closed under a nitrogen flow, mixed and shaken at 200 rpm overnight.
146 Dipentadecanoylphosphatidylcholine (C₃₈H₇₆NO₈P) (Larodan Fine Chemicals) was added as an
147 internal standard for quantification of PLFAs. After shaking for another five minutes, the samples
148 were centrifuged at 2500 rpm for 15 minutes. The volume of the supernatant was measured and
149 adjusted with chloroform and phosphate buffer to a ratio of 1:1:0.9 (vol:vol:vol) of chloroform-
150 methanol-phosphate buffer. Samples were centrifuged again (2500 rpm, five min.) and the lower
151 organic phase (total lipids) was evaporated to dryness. The total lipids were fractionated on a silicic
152 acid column (Agilent silica-based HF Bond Elut LRC-SI, 500 mg, Varian), into neutral, glyco-, and
153 phospholipids using 10, 20 and 10 ml of chloroform, acetone and methanol, respectively. The
154 phospholipids fraction was evaporated to dryness under nitrogen flow and methylated using the
155 protocol in Virtue et al. [38], but at 60–80 °C for two hours. Methylation standard nonadecanoic acid

156 (C₂₀H₄₀O₂) (Sigma-Aldrich) was added just before methylation and was used to quantify the
157 methylation efficiency. To collect methylated fatty acids (FAs), two ml of hexane/chloroform (4:1,
158 vol:vol) were added to the samples, after which the samples were vortexed and centrifuged at 2000
159 rpm for five minutes. The top organic layer was then transferred, dried (under a nitrogen stream) and
160 re-dissolved in a known volume of n-hexane.

161 The methylated FAs were analysed using an Agilent 6890 GC connected to an Agilent 5973 mass
162 selective detector. The methylated FAs were separated with a DB-5 fused silica capillary column (30
163 m x 0.25 mm x 0.25 µm), using helium as a carrier gas. The samples were injected by splitless
164 injection using the constant flow mode and using similar settings as Kaneko et al. [39]. The initial
165 oven temperature was 50 °C, and subsequently it was increased by 30 °C min⁻¹ to 140 °C and then by
166 5 °C min⁻¹ to 320 °C. This final temperature was held for 20 min leading to a total run time of 60 min.
167 Peaks were identified based on their relative retention times and mass spectra measured in SCAN
168 mode. The retention times of the peaks were also compared with the retention times of the fatty acid
169 methyl esters (FAME) in the standard mix (Supelco 37 component FAME mix). The internal standard
170 PC (Dipentadecanoylphosphatidylcholine) 15:0 was used for quantitative analysis. Dimethyl
171 disulphide (DMDS) adducts were prepared, analysed and used in the determination of the position of
172 double bonds in the monounsaturated FAMES [40]. The FA contents [µg g⁻¹ dry weight of soil (dw)
173 and % of PLFAs] were calculated. In order to account for variation in the peat compaction between
174 the drained and natural sides, between the fen and bog sites and among the different depth layers, the
175 amount of microbial FA in µg g⁻¹ dw was converted to g m⁻³ using the dry bulk densities (BD; Fig.
176 S2).

177

178 2.3.2. Characteristic FAs of Gram-negative and Gram-positive bacteria and fungi

179

180 PLFA biomarkers common among general microbial groups, such as Gram-negative bacteria,
181 Gram-positive bacteria and fungi, were selected and grouped. Iso- and anteiso branched PLFAs,

182 i14:0, i15:0, a15:0, a17:0 and i17:0, typical of Gram-positive bacteria [18,41,42] were grouped as
183 Branched FA (BrFA). 16 monounsaturated fatty acids (MUFAs), 16:1 ω 5, 16:1 ω 6c, 16:1 ω 7c and
184 16:1 ω 8c, as well as 18 MUFAs, 18:1 ω 5, 18:1 ω 6c, 18:1 ω 7c, 18:1 ω 7t and 18:1 ω 8c, both of which are
185 typical of Gram-negative bacteria [42], were grouped separately. The 18:2 ω 6 FA, typical of fungi,
186 [21] was studied as fungi FA.

187

188 2.4. Data analysis

189

190 The effect of drainage (at each site and depth layer) and depth (at each natural and drained side)
191 on the total microbial PLFA biomass, and the absolute amounts and relative abundances of the
192 selected microbial group FAs (16 MUFAs, 18 MUFAs, BrFA and fungi FA), were tested using
193 independent sample t-test and one-way ANOVA, respectively. Correlations of the total microbial
194 PLFA biomass with pH and BD were tested using Spearman correlation analysis. Multivariate
195 analyses of the PLFA profiles were based on Bray-Curtis dissimilarities calculated among samples
196 using $\log_{10}(x+1)$ transformed data of the relative abundances (% composition) of the PLFA. The data
197 were assessed graphically using non-metric multidimensional scaling (NMS) constrained to 2
198 ordination axes. NMS was done for the whole (natural and drained) fen and bog site data to depict
199 the overall patterns among sides, drainage and depth zones. Furthermore, the effect of depth and
200 drainage on PLFA profiles in both fen and bog were tested using 2-way permutational multivariate
201 analysis of variance (PERMANOVA) [43,44] with both factors as fixed factors. Mantel's test was
202 used to analyse correlations of the PLFA profiles with pH and BD. The Mantel correlation test result
203 is only meaningful when the result is positive, which shows correlation. The positive sign in our
204 correlation results means that, the larger the change in the pH or BD, the larger the change in microbial
205 community structure (PLFA structure). ANOVA was done using IBM SPSS Statistics version 23.
206 NMS and Mantel's tests were performed using PC-ORD version 6.0 ([45]; PC-ORD. Multivariate

207 analysis of ecological data. MjM Software, Glendon Beach, Oregon, USA). PerMANOVA was done
208 using FORTRAN program by Anderson [46].

209

210 **3. RESULTS**

211

212 3.1. Total microbial PLFA biomass and structure

213

214 We only analysed PLFAs between C10 and C20, which represents the main range for prokaryotic
215 PLFAs and a few other microbial groups, such as fungi. The most common FA in the samples was
216 C16:0 (a universal FA), which contributed (mean \pm SE) 10.0 ± 0.3 % to the whole depth PLFA profile
217 of the fen site (natural + drained) and 10.1 ± 0.4 % to the whole depth PLFA profile of the bog site
218 (natural + drained). The total microbial PLFA biomass (g m^{-3}) was higher in the natural fen than the
219 natural bog site at all the depths, except for the 25 – 50 cm depth, where it did not differ between the
220 fen and bog (Fig. 2A, Table S1). There was no correlation between pH and the total microbial PLFA
221 biomass either in the combined dataset of the fen and the bog, or the fen and the bog separately. In
222 all but the 50 – 100 cm depth of the fen site, total microbial PLFA biomass was higher in the drained
223 than the natural side. In the bog site, the total microbial PLFA biomass was only different between
224 the drained and natural side at the top and bottom layers, where the amount was smaller in the drained
225 side (Fig. 2A, Table S1).

226 There were also depth differences in the total microbial PLFA biomass in both the natural and
227 drained fen, but only in the natural side of the bog (Fig. 2A, Table S2). In both the natural and drained
228 fen, the amount of microbial PLFA biomass in the surface layer (0–25 cm) was higher than in the 25–
229 50 cm layer, but similar to that in the bottom layer. In the drained fen, the total microbial PLFA
230 biomass was also higher in the surface layer than in the 50–100 cm layer (Table S2). In the natural
231 bog, the amount of microbial PLFA biomass was higher in the surface layer than in the other depth
232 layers, which did not differ from one another (Table S2).

233 As visualised by NMS ordination, there were differences in the microbial community PLFA
234 compositions between the fen and bog sites (Fig. 3). NMS further suggests that the variation in the
235 microbial community structure was explained by both depth and drainage in both the fen and bog
236 sites (Fig. 3). This was confirmed by a two-way factorial (drainage and depth) analysis
237 (PERMANOVA), which showed that drainage and depth independently affected the microbial
238 community structure in both the fen and bog sites (Tables 2 & 3). There was correlation between pH
239 and the community structure in the combined dataset of the fen and bog (Mantel's test, $r = 0.42$, $p <$
240 0.001 , $n = 16$) as well as in the fen (Mantel's test, $r = 0.48$, $p < 0.05$, $n = 8$) and bog (Mantel's test, r
241 $= 0.52$, $p < 0.05$, $n = 8$) alone. Correlations between BD and community structure were also detected
242 in the combined dataset (Mantel's test, $r = 0.30$, $p < 0.01$, $n = 16$) as well as in the bog (Mantel's test,
243 $r = 0.57$, $p < 0.01$, $n = 8$), but not in the fen.

244

245 3.2. Characteristic FAs of Gram-negative and Gram-positive bacteria and fungi

246

247 The overall concentration of all the major microbial group PLFAs followed the same trend as the
248 total microbial PLFA, being higher in the fen than the bog site, especially in the drained side (Fig.
249 2B). They were also mostly highest in the surface layer than the deeper layers of both sites. The
250 amount (g m^{-3}) of 16 C monounsaturated fatty acids (16 MUFAs), which are characteristic of Gram-
251 negative bacteria, was higher in the top two layers (0 – 25 and 25 – 50 cm) of the drained side of the
252 fen than the natural side, but lower in the bottom layer (only) of the drained side of the bog than the
253 natural side (Fig. 2B). The relative contribution of 16 MUFAs to the total microbial PLFA (%
254 contribution) did not differ between the drained and the natural sides in either fen or bog except at
255 the 50 – 100 cm depth of the bog site (Fig. 4A). Depth only affected the amount of 16 MUFAs (g m^{-3})
256 in the drained fen and natural bog sides. In both cases, the amount (g m^{-3}) of 16 MUFAs was higher
257 in the surface (0 – 25 cm) layer than the other depth layers, which were similar to one another (Fig.
258 2B, Table S2). There was also no depth effect on the relative contribution of 16 MUFAs to the total

259 PLFAs except in the natural bog, where it was higher in the surface layer (0–25 cm) than at the other
260 depths, which were similar to one another (Fig. 4A, Table S3).

261 The amount (g m^{-3}) of 18 C monounsaturated fatty acids (18 MUFAs), which are also characteristic
262 of Gram-negative bacteria, was higher in the sub-surface layer (25 – 50 cm) of the drained fen side
263 compared to the natural fen side, and lower in the bottom layer of the drained bog side compared to
264 the natural bog side (Fig. 2C). The relative contribution of 18 MUFAs to the total microbial PLFA
265 (% contribution) did not differ between the drained and the natural sides of either fen or bog (Fig.
266 4B). There were depth differences in the amount of 18 MUFAs (g m^{-3}) in all sides except in the
267 drained bog. In the sides with depth differences, the amount (g m^{-3}) of 18 MUFAs was higher in the
268 surface (0 – 25 cm) layer than the other depth layers, which were similar to one another (Fig. 2C,
269 Table S2). There was also a depth effect on the relative contribution of 18 MUFAs to the total PLFAs
270 in all the sides, except the drained bog side (Fig. 4B, Table S3).

271 The amount (g m^{-3}) of terminally branched fatty acids (BrFAs), which are characteristic of Gram-
272 positive bacteria, was higher in the top two layers (0 – 25 and 25 – 50 cm) of the drained fen than the
273 natural fen side, but lower in the sub-surface (25 – 50 cm) and bottom layers of the drained bog than
274 the natural bog side (Fig. 2D). The relative contribution of BrFAs to the total microbial PLFA (%
275 contribution) did not differ between the drained and natural sides of either fen or bog (Fig. 4C). There
276 were only depth differences in the amount (g m^{-3}) of BrFAs in the drained sides of both the fen and
277 bog. The amount of BrFAs in the surface layer (0 – 25 cm) of the drained fen and the top two layers
278 (0 – 25 and 20 – 50 cm) of the drained bog sides were higher than in the other depth layers, which
279 were similar to each other (Fig. 2D, Table S2). There was no depth effect on the relative contribution
280 of BrFAs to the total PLFAs in any of the sides, except the natural bog side (Fig. 4C, Table S3).

281 The amount (g m^{-3}) of fatty acids characteristic of fungi (fungi FA) did not differ between the
282 drained and natural fen sides, but was lower in the surface layer (0 – 25 cm) of the drained bog side
283 than the natural bog side (Fig. 2E). Neither drainage nor depth affected the relative contribution of
284 fungal FA to the total microbial PLFA (% contribution) in either fen or bog sites (Fig. 4D, Table S3).

285 There were also no depth differences in the amount (g m^{-3}) of fungal FAs in any of the sides (Fig.
286 2E).

287

288 4. DISCUSSION

289

290 4.1. Biomass and community structure of microbes

291

292 The higher total microbial PLFA biomass in the natural fen compared to the bog is best explained
293 by the higher nutrient content and pH in the fen (Table 1 and Fig. S2) [5,13,14,19,47]. Concomitant
294 differences in vegetation cover affect microbes due to differences in soil structure and C substrate
295 availability. Biomass of fen vegetation is more easily decomposed than that of bog vegetation, which
296 consists largely of recalcitrant *Sphagnum* mosses [5,13,48]. The roots of sedges in fens provide better
297 soil stability and macro pore structure than those of *Sphagnum* mosses on the bogs. The higher
298 microbial PLFA biomass in the fen is also reflected in the higher CO_2 and CH_4 emissions from the
299 fen (natural side; Table 1) [28,49,50]. Differences in microbial biomasses were also accompanied by
300 differences in the microbial community structures of the fen and bog, which are due to similar reasons
301 as those stated above in addition to differences in the natural wetness of the sites [13,47].

302 There was viable microbial biomass in all the peat layers of both the fen and the bog sites (Fig. 2).
303 The drastic reduction in total microbial PLFA biomass from the surface layer to the 25–50 cm layer
304 in the fen but not the bog is likely due to depth related differences in the fertility and litter quality
305 [47] (Table S4). In general, the microbial biomass PLFA (g m^{-3}) in the fen increased with increasing
306 nitrogen (N %) and carbon (C %) content and decreasing C/N ratios, but this was not the case in the
307 bog site. There was no decrease in microbial biomass with depth (depth effect) in the bog site,
308 probably due to poor substrate quality (higher C/N ratio). Quite surprisingly, the total microbial PLFA
309 biomass did not differ between surface layers and bottom layers except in the natural bog. Although
310 PLFA-analysis detects viable cells and indicates changes in the potentially active microbial biomass,

311 it cannot separate the active and non-active cells [51]. We also acknowledge that the turnover rate of
312 PLFAs in the deep anoxic peat layers is unknown and could be considerably slower than in the oxic
313 and warmer surface peat layers. This means that detection of FAs in deeper peat layers may not be
314 indicative of similar cell activities as in the surface peat layers. However, the presence of potential
315 (enzymatic) prokaryotic microbial activity [52] and active microbial populations [53] have previously
316 been reported for deep peat layers (100 cm to 300 cm), which supports our finding of living microbes
317 in bottom peat layers. Our community structure analyses (% PLFA profiles, Table 3, Fig. 3; see also
318 4.3 below) also indicated differences in the microbial groups occupying different depths, possibly
319 due to community adaptations to depth-related changes in several factors, e.g. pH as shown here,
320 oxygen availability, alternative electron acceptors and substrates [5,54]. There is high temperature
321 variability at the surface, but low and stable temperature at the bottom (Fig. S2 and [31]), which
322 could also modify the community structure. Since there are large amounts of C stored in deeper peat
323 layers, the significant amount of microbial biomass in these layers may have implications for the
324 global C cycles, such as microbial-enhanced C flows from peat to mineral subsoil. There is an
325 estimated average $13.6 \text{ g m}^{-2} \text{ yr}^{-1}$ C input from peat into mineral subsoils [55], which may increase
326 with changes in deep peat microbial biomass and composition.

327

328 4.2. Effects of drainage on the biomass and community structure of microbes

329

330 Drainage increased total microbial PLFA biomass in the surface, subsurface and bottom layers of
331 the fen, supporting our first hypothesis, but decreased it in the surface and bottom layers of the bog
332 site, which is contrary to our first hypothesis. This is possibly due to differences in their nutrient
333 status, thickness of the aerated layer, available substrate and vegetation changes [13,22,47,56]. In
334 general, while drainage in the fen led to a succession towards a different ecosystem from the original,
335 in the bog site it led to smaller changes. Typical fen species, such as tall sedges, were succeeded by
336 spruce swamp and forest species, such as *Pinus sylvestris*, *Betula pubescens* and *Polytrichum*

337 *commune*. At the bog site, mosses and dwarf shrubs decreased while the forest species, mainly *Pinus*
338 *sylvestris*, increased and *Pleurozium schreberi* appeared (see also Table 1). Thirty-eight years after
339 drainage, tree stand volume increased from nothing in the natural fen to 111 m³ ha⁻¹ in the drained
340 fen side and from 5 in the natural bog to 16 m³ ha⁻¹ in the drained bog side [31]. The increased tree
341 growth in the drained fen led to greater evapotranspiration and further decrease in the WT. Although
342 the WT depth at the time of sampling was about the same in both sites due to heavy rainfall in the
343 previous weeks, the six-month mean WT depths before our sampling date were -8.0 and -34.9 cm at
344 the natural and drained fen, respectively and, -12.0 and -16.4 cm for the natural and drained bog,
345 respectively. The previously reported annual mean WT depths were also much lower in the drained
346 fen than in the drained bog for most parts of the year (Fig. 1 and Fig. S4). The much lower WT in the
347 drained fen therefore explains why there were drainage-induced changes in the total microbial PLFA
348 biomass in the surface and sub-surface layers of the fen, but only in the surface layer of the bog site.
349 Other reasons could be the originally lower substrate quality and availability in the bog, which has
350 become poorer after a few decades of drainage (Table S4) [5,13,48]. Our C/N ratio results showed
351 similar ratios along the profiles of both the natural and drained fen, suggesting that the increase in
352 microbial PLFA biomass was not due to differences in peat quality. However, the C/N ratios in the
353 bog profiles were significantly higher in the drained side, especially in the bottom layer. High C/N
354 ratio usually indicates low substrate quality, which could also explain the low microbial biomass in
355 the drained bog. We cannot conclusively explain the differences in C/N ratios between the natural
356 and drained bog sides, which are possibly caused by loss of N after drainage. Since drainage increases
357 N losses by leaching and plant uptake, it means that we are discussing both the direct and indirect
358 effects of drainage. We also note that there could have been natural differences in the peat qualities
359 of the bog site before drainage. However, we believe that our basic assumption that the original peat
360 quality was similar holds true, and that the differences we found can be associated to drainage. Hence,
361 in accordance with our third hypothesis, drainage increased the total microbial biomass more in the

362 fen than in the bog site. The overall effect of drainage on microbial biomass was reflected by the
363 increased CO₂ and decreased CH₄ efflux, especially in the drained fen (Table 1).

364 The total microbial PLFA biomass in the deepest (bottom) layers of both the fen and bog sites
365 were different between the natural and drained sides. This was due to changes in the flow and
366 constituents of the leachate water [24], deeper deposition of labile root exudates by vascular plants
367 and differences in the amount and quality of DOC reaching the bottom layers in the drained sides
368 [26,27]. Although the roots of vascular plants may not reach the bottom layers, their exudates will
369 get to microbes in the bottom layers via water movement at a faster rate than those deposited by the
370 shallow roots of mosses. The results of DOC isotopic analysis and tritium analysis of pore water, by
371 Charman et al [24], supported the downward movement of younger C via water movement in
372 ombrotrophic boreal peatland. The study also showed that CO₂ and CH₄ (based on ¹⁴C dating) from
373 the deeper peat layers was younger than that from the surrounding peat. They attributed this to the
374 transfer of DOC and gaseous C compounds to deeper peat via water movement, and the microbial
375 usage of younger C. Furthermore, they concluded that low hydraulic conductivity in peat may not be
376 a real limitation to water movement over a long time scale. Although mainly studied in a fen peat
377 site, Krüger et al. [28] also reported the effect of drainage in the deeper peat profiles, which showed
378 higher δ¹³C values (a qualitative indicator of peat degradation) in the drained than the natural side.
379 They also reported much older peat in the drained sides of both fen and bog. This also suggests an
380 effect of drainage on deeper peat microbial activity, as microbes use younger C that is more enriched
381 with ¹⁴C and DOC is a potential source of relatively younger C (leaving ¹⁴C depleted older peat
382 behind) [24]. By affecting the deep peat microbial processes [28] and biomass, long-term drainage
383 may have additional significant effects on the peat C balance, besides the enhanced C losses from
384 surface peat. Increased microbial biomass in the bottom peat layers due to drainage can either
385 contribute to the peat organic matter (OM) stabilization via the microbial C pump (MCP) process
386 ([4]; not studied in peatland yet), or enhance deep peat microbial degradation [28] and C flow to sub-
387 soil [55]. Further studies are needed to elaborate on this.

388

389 There were, however, no differences in the total microbial biomass in some subsurface layers (25
390 – 100 cm in bog and 50 -100 cm in fen) of the natural and drained sides. Inconsistent patterns in the
391 long-term drainage-induced changes, especially in the subsurface layers, have been previously
392 reported [13]. This is possibly because the different microbial communities at the different depths in
393 the pristine peatlands [5,13,54] have different sensitivities to changed hydrology and react differently
394 to drainage-induced changes [5,13,15,19]. Since the effect of drainage on microbial communities of
395 the deeper anoxic layers also depends on the movement of materials (e.g. DOC) from the surface to
396 deeper layers [24,26,27], accumulation of materials to concentrations that can cause significant
397 changes in microbial biomass is less probable in the intermediate layers than the deepest layers many
398 years after drainage [24].

399 Similar to in several previous studies, peat microbial community structure in this study was
400 indexed by the relative composition of microbial PLFAs [13,18-21,57]. According to our second
401 hypothesis, our results showed that long-term drainage affected the microbial community structures
402 in all the depth layers including the bottom (deepest) layers. Jaatinen et al. [13] and Urbanová and
403 Bárta [22] studied the effects of long-term drainage on microbial communities in the surface layers
404 (0 -30 cm) of peatlands. They also showed that long-term drainage and the resulting changes in
405 vegetation pattern altered the microbial community structures in different peatland types. Our own
406 measurements and those from previous studies showed that the pH in both the fen and bog sites were
407 mostly reduced by drainage in all the depth layers (Fig. S2 and S3). Therefore, the correlation between
408 the microbial community structure and pH in the natural and drained sides of both the fen and bog
409 sites (by Mantel's test) partly explains the changes in the microbial community structures at the
410 different depth layers [13]. Other reasons for the changes in microbial community structure include
411 changes in the anoxic/oxic condition of the surface layer and the available litter quality at different
412 depths, occasioned by the drainage-induced changes in the prevailing plant communities and
413 structures (see table 1 and paragraph 1 of 4.2) [16,19,56].

414

415 4.3. Characteristic FAs of Gram-negative and Gram-positive bacteria, and fungi

416

417 Similar to in previous studies [13-15], drainage and depth-induced effects on the biomass of the
418 FAs characteristic of our microbial groups, varied among the microbial groups and between the fen
419 and bog sites. The higher concentrations of all the FAs characteristic of the major microbial groups,
420 except the fungal-characteristic FA, in the natural and drained fen than the bog site was due to the
421 same reasons given for total PLFA biomass (in 4.1). The biomass of both the Gram-negative and
422 Gram-positive bacteria characteristic FAs were increased in the surface and subsurface layers of the
423 drained fen side, probably due to drainage-induced increases in aeration and substrate availability.
424 However, they were decreased in the bottom layer of the bog site, possibly due to low substrate
425 quality, which was revealed by the C/N ratio in the drained side. The fungal characteristic FA was
426 only reduced in the surface layer of the bog, likely due to low peat biomass quality and differences in
427 the sensitivities of different fungal species to drainage [15]. For example, Peltoniemi et al [15] showed
428 that basidiomycetes are more sensitive to drainage than ascomycetes, though they are within the same
429 fungal phyla represented by the same characteristic FA.

430 Drainage-induced effects on the relative abundance of our selected microbial group characteristic
431 FAs were not observed, except for 16 MUFAs in the 50 – 100 cm depth of the bog site. This was
432 probably due to the contrasting drainage responses of the different species in these groups [9,13, 15].
433 Kim et al. [14] also reported no differences in the diversity and composition of denitrifiers and
434 methanogens in any of their sites following a short-term drought. The depth-induced effects were
435 also inconsistent among the groups and between the fen and bog sites (Fig.4, Table S3). This was
436 probably because these microbial groups contained different species with contrasting depth
437 stratification patterns. Our 25 cm sample depth-range may also be too long to reveal the inconsistent
438 changes observed in the shorter sampling depth-ranges for specific microbial groups (e.g. [9,13,22]).
439 For example, Lin et al. [52] reported an increasing proportion of yeast (*Saccharomyces*) and a

440 reduction of the white-rot fungi (*Agaricomycotina*) with depth, although both of them are fungi. The
441 effect of drainage on the relative abundances of the mainly aerobic bacteria FAs (16 MUFAs) in the
442 50 – 100 cm layer of bog may be due to change from a permanently anoxic to at least episodic oxic
443 conditions in the subsurface layers (see Fig. 1; [13]). Using techniques with higher taxonomic
444 resolution than PLFA - analysis, e.g. next-generation sequencing of 16S rRNA gene amplicons, could
445 give better insight to the drainage- and depth-induced effects observed with the total microbial
446 community. Nevertheless, it is clear from our results that the depth-induced effects on the microbial
447 community structure vary between the two sites and between the treatments. This suggests that the
448 factors regulating depth-induced effects on specific microbial species (within the groups) differ
449 between sites.

450

451 **5. CONCLUSIONS**

452

453 Our study shows that the biomass and the structure of *in-situ* microbial communities differ between
454 the studied fen and bog sites. While long-term water level draw-down significantly increased the total
455 microbial biomass in the affected layers of the fen, it decreased the total microbial biomass in the
456 affected layers of the bog, which is likely due to the differences in their original nutrient statuses, the
457 level of WT depth reduction and substrate quality. There was a considerable amount of living
458 microbial biomass even in the deepest and oldest (3000-3400 years) layers of the studied peatlands.
459 Although drainage mostly affects microbial community in the surface layers, the effect of long-term
460 WT draw-down on deeper peat microbes is also important, due to the long-term storage of large
461 amounts of C in deeper peat layers. Drainage-induced increase in microbial biomass, measured in
462 this study for the fen site, can enhance microbial contribution to deeper peat OM stabilization via the
463 microbial C pump (MCP) or increase microbial peat degradation and C release in the bottom peat
464 layers. This suggests a more significant effect of drainage on climate change than revealed by surface
465 layer analyses alone. Further studies are needed to elaborate on this. Since peatland drainage has been

466 used to model climate change effects on peatlands [11], our results might be revealing climate change
467 effects on deep peat microbial biomasses.

468

469 **Acknowledgements**

470

471 We are grateful to Niina Tavi, Hanne Säppi and Juhani Tarhanen for assisting in the laboratory
472 during the PLFA extraction. Many thanks to Timo Penttilä (M.Sc) and Raija Laiho (PhD) for helping
473 out with the WT data. Thanks to James Blande (PhD) for checking the English text. We also thank
474 Prof. Marja Tirola for her effort in making sure that the MS is ready in good time. Thanks also to the
475 reviewers for their constructive comments. The study was supported by the Academy of Finland
476 (projects 136455 and 140964 to Hannu Nykänen, project 286642 to Antti J Rissanen and project
477 260797 to Marja Tirola and project 291691 to Christina Biasi). Christina Biasi further acknowledges
478 support from Nessling foundation.

479 **References**

480

- 481 [1] C. Immirzi, E. Maltby, R. Clymo, The global status of peatlands and their role in carbon cycling, (1992).
- 482 [2] J. Turunen, E. Tomppo, K. Tolonen, A. Reinikainen, Estimating carbon accumulation rates of undrained
483 mires in Finland—application to boreal and subarctic regions, *The Holocene*. 12 (2002) 69-80.
- 484 [3] A.S. Grandy, J.C. Neff, Molecular C dynamics downstream: the biochemical decomposition sequence
485 and its impact on soil organic matter structure and function, *Sci. Total Environ*. 404 (2008) 297-307.
- 486 [4] A. Miltner, P. Bombach, B. Schmidt-Brücken, M. Kästner, SOM genesis: microbial biomass as a
487 significant source, *Biogeochemistry*. 111 (2012) 41-55.
- 488 [5] R. Andersen, S. Chapman, R. Artz, Microbial communities in natural and disturbed peatlands: a review,
489 *Soil Biol. Biochem*. 57 (2013) 979-994.
- 490 [6] M.P. Waldrop, M.K. Firestone, Microbial community utilization of recalcitrant and simple carbon
491 compounds: impact of oak-woodland plant communities, *Oecologia*. 138 (2004) 275-284.
- 492 [7] M. Rabinovich, A. Bolobova, L. Vasil'chenko, Fungal decomposition of natural aromatic structures and
493 xenobiotics: a review, *Appl. Biochem. Microbiol*. 40 (2004) 1-17.

- 494 [8] A.M. Treonis, N.J. Ostle, A.W. Stott, R. Primrose, S.J. Grayston, P. Ineson, Identification of groups of
495 metabolically-active rhizosphere microorganisms by stable isotope probing of PLFAs, *Soil Biol. Biochem.*
496 36 (2004) 533-537.
- 497 [9] K. Jaatinen, R. Laiho, A. Vuorenmaa, U. Del Castillo, K. Minkkinen, T. Pennanen, T. Penttilä, H. Fritze,
498 Responses of aerobic microbial communities and soil respiration to water-level drawdown in a northern
499 boreal fen, *Environ. Microbiol.* 10 (2008) 339-353.
- 500 [10] M. Maljanen, B. Sigurdsson, J. Guðmundsson, H. Óskarsson, J. Huttunen, P. Martikainen, Greenhouse
501 gas balances of managed peatlands in the Nordic countries—present knowledge and gaps, *Biogeosciences.* 7
502 (2010) 2711-2738.
- 503 [11] Change, Intergovernmental Panel On Climate, *Climate change 2007: The physical science basis,*
504 *Agenda.* 6 (2007) 333.
- 505 [12] E. Gorham, Northern peatlands: role in the carbon cycle and probable responses to climatic warming,
506 *Ecol. Appl.* 1 (1991) 182-195.
- 507 [13] K. Jaatinen, H. Fritze, J. Laine, R. Laiho, Effects of short-and long-term water-level drawdown on the
508 populations and activity of aerobic decomposers in a boreal peatland, *Global Change Biol.* 13 (2007) 491-
509 510.
- 510 [14] S. Kim, S. Lee, C. Freeman, N. Fenner, H. Kang, Comparative analysis of soil microbial communities
511 and their responses to the short-term drought in bog, fen, and riparian wetlands, *Soil Biol. Biochem.* 40
512 (2008) 2874-2880.
- 513 [15] K. Peltoniemi, H. Fritze, R. Laiho, Response of fungal and actinobacterial communities to water-level
514 drawdown in boreal peatland sites, *Soil Biol. Biochem.* 41 (2009) 1902-1914.
- 515 [16] R. Laiho, Decomposition in peatlands: Reconciling seemingly contrasting results on the impacts of
516 lowered water levels, *Soil Biol. Biochem.* 38 (2006) 2011-2024.
- 517 [17] K. Yrjälä, T. Tuomivirta, H. Juottonen, A. Putkinen, K. Lappi, E. Tuitila, T. Penttilä, K. Minkkinen, J.
518 Laine, K. Peltoniemi, CH₄ production and oxidation processes in a boreal fen ecosystem after long-term
519 water table drawdown, *Global Change Biol.* 17 (2011) 1311-1320.
- 520 [18] I. Sundh, M. Nilsson, P. Borga, Variation in microbial community structure in two boreal peatlands as
521 determined by analysis of phospholipid fatty acid profiles, *Appl. Environ. Microbiol.* 63 (1997) 1476-1482.
- 522 [19] P. Borga, M. Nilsson, A. Tunlid, Bacterial communities in peat in relation to botanical composition as
523 revealed by phospholipid fatty acid analysis, *Soil Biol. Biochem.* 26 (1994) 841-848.
- 524 [20] R. Andersen, L. Grasset, M.N. Thormann, L. Rochefort, A. Francez, Changes in microbial community
525 structure and function following Sphagnum peatland restoration, *Soil Biol. Biochem.* 42 (2010) 291-301.
- 526 [21] Å Frostegård, E. Bååth, The use of phospholipid fatty acid analysis to estimate bacterial and fungal
527 biomass in soil, *Biol. Fertility Soils.* 22 (1996) 59-65.
- 528 [22] Z. Urbanová, J. Bárta, Effects of long-term drainage on microbial community composition vary between
529 peatland types, *Soil Biol. Biochem.* 92 (2016) 16-26.
- 530 [23] C. Freeman, N. Fenner, N. Ostle, H. Kang, D. Dowrick, B. Reynolds, M. Lock, D. Sleep, S. Hughes, J.
531 Hudson, Export of dissolved organic carbon from peatlands under elevated carbon dioxide levels, *Nature.*
532 430 (2004) 195-198.

- 533 [24] D.J. Charman, R. Aravena, C.L. Bryant, D.D. Harkness, Carbon isotopes in peat, DOC, CO₂, and CH₄
534 in a Holocene peatland on Dartmoor, southwest England, *Geology*. 27 (1999) 539-542.
- 535 [25] Y. Jia, L. Bakken, G. Breedveld, P. Aagaard, Å Frostegård, Organic compounds that reach subsoil may
536 threaten groundwater quality; effect of benzotriazole on degradation kinetics and microbial community
537 composition, *Soil Biol. Biochem.* 38 (2006) 2543-2556.
- 538 [26] H.K. Jokinen, O. Kiikkilä, H. Fritze, Exploring the mechanisms behind elevated microbial activity after
539 wood ash application, *Soil Biol. Biochem.* 38 (2006) 2285-2291.
- 540 [27] M.P. Laine, R. Strömmer, L. Arvola, DOC and CO₂-C releases from pristine and drained peat soils in
541 response to water table fluctuations: a mesocosm experiment, *Applied and Environmental Soil Science*. 2014
542 (2014).
- 543 [28] J.P. Krüger, C. Alewell, K. Minkkinen, S. Szidat, J. Leifeld, Calculating carbon changes in peat soils
544 drained for forestry with four different profile-based methods, *For. Ecol. Manage.* 381 (2016) 29-36.
- 545 [29] P. Pirinen, H. Simola, J. Aalto, J. Kaukoranta, P. Karlsson, R. Ruuhela, Tilastpja Suomen Ilmastosta
546 1981-2010, Ilmatieteen laitos, 2012.
- 547 [30] J. Laine, V. Komulainen, R. Laiho, K. Minkkinen, A. Rasinmäki, T. Sallantausta, S. Sarkkola, N. Silvan,
548 K. Tolonen, E. Tuittila, Lakkasuo: A Guide to Mire Ecosystem. Department of Forest Ecology, University of
549 Helsinki, 2004.
- 550 [31] K. Minkkinen, H. Vasander, S. Jauhiainen, M. Karsisto, J. Laine, Post-drainage changes in vegetation
551 composition and carbon balance in Lakkasuo mire, Central Finland, *Plant Soil*. 207 (1999) 107-120.
- 552 [32] H. Nykänen, J. Alm, J. Silvola, K. Tolonen, P.J. Martikainen, Methane fluxes on boreal peatlands of
553 different fertility and the effect of long-term experimental lowering of the water table on flux rates, *Global*
554 *Biogeochem. Cycles*. 12 (1998) 53-69.
- 555 [33] J. Silvola, J. Alm, U. Ahlholm, H. Nykanen, P.J. Martikainen, CO₂ fluxes from peat in boreal mires
556 under varying temperature and moisture conditions, *J. Ecol.* (1996) 219-228.
- 557 [34] I. Belokopytov, V. Beresnevich, Giktorf's peat borers, *Torfyanaya Promyshlennost*. 8 (1955) 9-10.
- 558 [35] F. De Vleeschouwer, F.M. Chambers, G.T. Swindles, Coring and sub-sampling of peatlands for
559 palaeoenvironmental research, *Mires and Peat*. 7 (2010).
- 560 [36] N.M. Tavi, P.J. Martikainen, K. Lokko, M. Kontro, B. Wild, A. Richter, C. Biasi, Linking microbial
561 community structure and allocation of plant-derived carbon in an organic agricultural soil using ¹³C₂ pulse-
562 chase labelling combined with ¹³C-PLFA profiling, *Soil Biol. Biochem.* 58 (2013) 207-215.
- 563 [37] E.G. Bligh, W.J. Dyer, A rapid method of total lipid extraction and purification, *Canadian journal of*
564 *biochemistry and physiology*. 37 (1959) 911-917.
- 565 [38] P. Virtue, P.D. Nichols, P.I. Boon, Simultaneous estimation of microbial phospholipid fatty acids and
566 diether lipids by capillary gas chromatography, *J. Microbiol. Methods*. 25 (1996) 177-185.
- 567 [39] M. Kaneko, H. Naraoka, Y. Takano, N. Ohkouchi, Distribution and isotopic signatures of archaeal lipid
568 biomarkers associated with gas hydrate occurrences on the northern Cascadia Margin, *Chem. Geol.* 343
569 (2013) 76-84.
- 570 [40] P. Nichols, B.K. Stulp, J.G. Jones, D.C. White, Comparison of fatty acid content and DNA homology of
571 the filamentous gliding bacteria *Vitreoscilla*, *Flexibacter*, *Filibacter*, *Arch. Microbiol.* 146 (1986) 1-6.

- 572 [41] W. O'leary, S. Wilkinson, Gram-positive bacteria, *Microbial lipids*. 1 (1988) 117-201.
- 573 [42] S. Taipale, R.I. Jones, M. Tiirola, Vertical diversity of bacteria in an oxygen-stratified humic lake,
574 evaluated using DNA and phospholipid analyses, *Aquat. Microb. Ecol.* 55 (2009) 1-16.
- 575 [43] M.J. Anderson, A new method for non-parametric multivariate analysis of variance, *Austral Ecol.* 26
576 (2001) 32-46.
- 577 [44] B.H. McArdle, M.J. Anderson, Fitting multivariate models to community data: a comment on distance-
578 based redundancy analysis, *Ecology*. 82 (2001) 290-297.
- 579 [45] B. McCune, M.J. Mefford, PC-ORD, in: *Anonymous Multivariate Analysis of Ecological Data*, Version
580 6.0.MjM Software ed., Glenden Beach, Oregon, U.S.A., 2011.
- 581 [46] M. Anderson, PERMANOVA: a FORTRAN computer program for permutation multifactor analysis of
582 variance, Department of Statistics, University of Auckland, Auckland, NZ. (2005).
- 583 [47] M.D. Preston, K.A. Smemo, J.W. McLaughlin, N. Basiliko, Peatland microbial communities and
584 decomposition processes in the James Bay lowlands, Canada, *Front. Microbiol.* 3 (2012) 70.
- 585 [48] I. Mandic-Mulec, L. Ausec, T. Danevčič, Š. Levičnik-Höfferle, V. Jerman, B. Kraigher, *Microbial
586 Community Structure and Function in Peat Soil*, *Food Technology and Biotechnology*. 52 (2014) 180-187.
- 587 [49] J. Krüger, J. Leifeld, C. Alewell, Degradation changes stable carbon isotope depth profiles in peat
588 lands, *Biogeosciences*. 11 (2014) 3369-3380.
- 589 [50] J. Krüger, J. Leifeld, S. Glatzel, S. Szidat, C. Alewell, Biogeochemical indicators of peatland
590 degradation—a case study of a temperate bog in northern Germany, *Biogeosciences*. 12 (2015) 2861-2871.
- 591 [51] E. Blagodatskaya, Y. Kuzyakov, Active microorganisms in soil: critical review of estimation criteria
592 and approaches, *Soil Biol. Biochem.* 67 (2013) 192-211.
- 593 [52] X. Lin, M.M. Tfaily, J.M. Steinweg, P. Chanton, K. Esson, Z.K. Yang, J.P. Chanton, W. Cooper, C.W.
594 Schadt, J.E. Kostka, Microbial community stratification linked to utilization of carbohydrates and
595 phosphorus limitation in a boreal peatland at Marcell Experimental Forest, Minnesota, USA, *Appl. Environ.
596 Microbiol.* 80 (2014) 3518-3530.
- 597 [53] A. Putkinen, H. Juottonen, S. Juutinen, E. Tuittila, H. Fritze, K. Yrjälä, Archaeal rRNA diversity and
598 methane production in deep boreal peat, *FEMS Microbiol. Ecol.* 70 (2009) 87-98.
- 599 [54] R.R. Artz, Microbial community structure and carbon substrate use in northern peatlands, *Carbon
600 Cycling in Northern Peatlands*. (2009) 111-129.
- 601 [55] J. Turunen, K. Tolonen, S. Tolvanen, M. Remes, J. Ronkainen, H. Jungner, Carbon accumulation in the
602 mineral subsoil of boreal mires, *Global Biogeochemical Cycles*. 13 (1999) 71-79.
- 603 [56] J. Laine, H. Vasander, R. Laiho, Long-term effects of water level drawdown on the vegetation of
604 drained pine mires in southern Finland, *J. Appl. Ecol.* (1995) 785-802.
- 605 [57] K. Peltoniemi, M. Pyrhönen, R. Laiho, M. Moilanen, H. Fritze, Microbial communities after wood ash
606 fertilization in a boreal drained peatland forest, *Eur. J. Soil Biol.* 76 (2016) 95-102.
- 607 [58] P.J. Martikainen, H. Nykänen, J. Alm, J. Silvola, Change in fluxes of carbon dioxide, methane and
608 nitrous oxide due to forest drainage of mire sites of different trophic, *Plant Soil*. 168 (1995) 571-577.

- 609 [59] J. Laine, K. Minkkinen, J. Sinisalo, I. Savolainen, P.J. Martikainen, Greenhouse impact of a mire after
610 drainage for forestry, Northern Forested Wetlands, Ecology and Management, edited by: Trettin, CC,
611 Jurgensen, MF, Grigal, DF, Gale, MR, and Jeglum, JK, CRC Lewis Publishers, Boca Raton, USA. (1996)
612 437-447.
- 613 [60] H. Juottonen, P.E. Galand, E. Tuittila, J. Laine, H. Fritze, K. Yrjälä, Methanogen communities and
614 Bacteria along an ecohydrological gradient in a northern raised bog complex, Environ. Microbiol. 7 (2005)
615 1547-1557.
- 616 [61] P.E. Galand, H. Fritze, R. Conrad, K. Yrjala, Pathways for methanogenesis and diversity of
617 methanogenic archaea in three boreal peatland ecosystems, Appl. Environ. Microbiol. 71 (2005) 2195-2198.
- 618 [62] K. Jaatinen, E. Tuittila, J. Laine, K. Yrjälä, H. Fritze, Methane-oxidizing bacteria in a Finnish raised
619 mire complex: effects of site fertility and drainage. Microbial ecology. 50 (2005) 429-439.
- 620 [63] H. Toberman, R. Laiho, C. Evans, R. Artz, N. Fenner, P. Straková, C. Freeman, Long-term drainage for
621 forestry inhibits extracellular phenol oxidase activity in Finnish boreal mire peat, Eur. J. Soil Sci. 61 (2010)
622 950-957.
- 623 [64] M. Pearson, T. Penttilä, L. Harjunpää, R. Laiho, J. Laine, T. Sarjala, K. Silvan, N. Silvan, effects of
624 temperature rise and water-table-level drawdown on greenhouse gas fluxes of boreal sedge fens. Boreal
625 Environ. Res. 20 (2015) 489 – 505.
- 626

627 **Captions for figures**

628 **Fig. 1.** Average daily water table (WT) depth in the sampled peatland sites from May to December
629 of the sampling year

630
631 **Fig. 2.** Mean (\pm SE, $n = 3$) amount (g m^{-3}) of (A) microbial total PLFA, (B) 16 MUFAs, (C) 18
632 MUFAs, (D) terminally branched FAs and (E) FA characteristic of fungi in all the depths and sites
633 sampled. Significant differences ($p < 0.05$) between the drained and natural sides at each site in each
634 depth following independent sample t-tests are denoted by an asterisk (*). For depth effects Fn = fen
635 natural, Fd = fen drained, Bn = bog natural and Bd = bog drained, while sites with significant depth
636 effects (1-ANOVA, $p < 0.05$) are highlighted in bold. Bottom = 135 – 160 and 105 – 130 for fen
637 natural and drained, 217 – 242 and 245 – 270 for bog natural and drained respectively.

638
639 **Fig. 3.** Non-metric multidimensional scaling (NMS) ordination of PLFA [$\log_{10}(x+1)$] of the relative
640 abundance of individual PLFAs]. Average (\pm SE, $n = 3$) NMS axes scores of the sites. Axes are
641 arbitrary; the closer the sample points are on the plot, the more similar they are in PLFA composition.
642 Depth 0–25 cm (\circ), 25–50 cm (Δ), 50–100 cm (\square) and bottom layers (\diamond). The drained and natural
643 sides of both the fen and bog sites differ significantly ($p < 0.05$) in a two-way factorial (drainage and
644 depth) analysis (PERMANOVA, see Table 2).

645
646 **Fig. 4.** Mean (\pm SE, $n = 3$) relative abundance (% contribution to total microbial PLFAs) of (A) 16
647 MUFAs, (B) 18 MUFAs, (C) terminally branched FAs and (D) FA characteristic of fungi at different
648 depths in the studied fen and bog sites. Significant differences ($p < 0.05$) between drained and natural
649 sides at each depth and for each microbial group following independent sample t-tests are denoted by
650 an asterisk (*). For depth effects Fn = fen natural, Fd = fen drained, Bn = bog natural and Bd = bog

651 drained, while sites with significant depth effects (1-ANOVA, $p < 0.05$) are highlighted in bold.
652 Bottom = 135 – 160 and 105 – 130 for fen natural and drained, 217 – 242 and 245 – 270 for bog
653 natural and drained respectively.

654

655 Table 1. General features of the study sites

Site Managements	Fen				Bog											
	Natural		Drained		Natural		Drained									
Peatland type ^(1,2)	Tall sedge fen		Tall sedge fen planted with scots pines		Cottongrass pine bog with <i>Sphagnum fuscum</i> hummocks		Cottongrass pine bog with <i>Sphagnum</i> <i>fuscum</i> hummocks									
Tree stand volume ⁽¹⁾ (m ⁻³ ha ⁻¹)	0		111		5		16									
Peat thickness ⁽¹⁾ (cm)	168		140		267		244									
CO ₂ flux ⁽³⁾ (g CO ₂ - C year ⁻¹)	188		356		164		236									
CH ₄ flux ⁽²⁾ (g CH ₄ - C year ⁻¹)	31.0		-0.0		4.8		2.7									
Peat bottom age ⁽¹⁾ (years)	3400				3000											
Peat constituent ⁽¹⁾	C (L, S, Er)				S (Er, L)											
Vegetation ^(2,4)	Ap Cl Cr Ev Ps Sa Sp Sf Bn				Pc Ps Ac Bp Ce Dc Psc Sa Psy Bp				Sa Ap Ev Ps Rc Sf En Psy Cs Dp Ev Psc Sr Psy							
C (%) (surface)	50.1±1.1				53.2±0.4				47.9±0.5				46.8±0.4			
N (%) (surface)	2.5±0.3				2.3±0.4				1.2±0.2				0.86±0.1			
P (μg g ⁻¹) ⁽¹⁾ (surface)	0.82				1.20				0.37				0.50			

656 Lakkasuo mire complex features are adopted and modified from ¹Minkkinen *et al.* (1999) [31], ²Nykänen *et al.* (1998) [32], ³Silvola *et al.* (1996) [33] or ⁴Laine *et al.*
657 (2004) [30]. Peat constituents: C = *Carex*, L = *Lignum*, S = *Sphagnum*, Er = *Eriophorum*. Vegetation cover: Ac, *Agrostitis capillaris*; Ap, *Andromeda polifolia*; Bp,
658 *Betula pubescent*; Bn, *Betula nana*; Ce, *Carex echinata*; Cl, *Carex lasiocarpa*; Cr, *Carex rostrate*; Cs, *Cladonia* sp.; Dc, *Dryopteris carthusiana*; Dp, *Dicranum*
659 *polysetum*; En, *Empetrum nigrum*; Ev, *Eriophorum vaginatum*; Pc, *Polytrichum commune*; Ps, *Polytrichum strictum*; Psc, *Pleurozium schreberi*; Psy, *Pinus*
660 *sylvestris*; Rc, *Rubus chamaemorus*; Sa, *Sphagnum angustifolium*; Sf, *Sphagnum fuscum*; Sp, *Shagnum papillosum*; Sr, *Sphagnum russowii*. Dominant tree species
661 marked with bold.

662 Table 2. Two-way factorial (drainage and depth) analysis (PERMANOVA) explaining the structural
 663 variation in the analysed microbial communities, between depths (0 –25 cm, 25 – 50 cm, 50 – 100
 664 cm, bottom) and between drained and natural sides in fen and bog sites. Analysis was done with log₁₀
 665 (x+1) -transformed PLFA relative abundance (% composition) data. Significant depth and drainage
 666 effects (p < 0.05) are indicated with '*’.

	df	Fen		Bog	
		Rel. abundance		Rel. abundance	
		<i>F</i>	<i>p</i>	<i>F</i>	<i>p</i>
Drainage	1	2.89	0.0194*	6.76	0.0001*
Depth	3	3.16	0.0005*	7.52	0.0001*
Interaction	3	1.08	0.3704	1.09	0.3854


667

668 **Table 3.** Difference in the structure of microbial community among depths in fen and bog sites, using
 669 the $\log_{10}(x+1)$ of PLFA relative abundance (% composition). Significant differences ($p < 0.05$)
 670 among depths from pair-wise analyses following 2-way (PERMANOVA; Table 2) analysis are shown
 671 with letters (different letter denotes significant differences among depths).

Depth (cm)	Fen	Bog
0 – 25	a	a
25 – 50	b	a
50 – 100	bc	b
bottom	c	c


672 Bottom = 135 – 160 and 105 – 130 for fen natural and drained, 217 – 242 and 245 – 270 for bog natural and
 673 drained respectively.

674




675 **Fig. 1.** Average daily water table (WT) depth in the sampled peatland sites from May to December
 676 of the sampling year.

677




678 **Fig. 2.** Mean (\pm SE, $n = 3$) amount (g m^{-3}) of (A) microbial total PLFA, (B) 16 MUFAs, (C) 18
679 MUFAs, (D) terminally branched FAs and (E) FA characteristic of fungi in all the depths and sites
680 sampled. Significant differences ($p < 0.05$) between the drained and natural sides at each site in each
681 depth following independent sample t-tests, are denoted by asterisk (*). For depth effects Fn = fen
682 natural, Fd = fen drained, Bn = bog natural and Bd = bog drained, while sites with significant depth
683 effects (1-ANOVA, $p < 0.05$) are highlighted in bold. Bottom = 135 – 160 and 105 – 130 for fen
684 natural and drained, 217 – 242 and 245 – 270 for bog natural and drained respectively.

685



686 **Fig. 3.** Non-metric multidimensional scaling (NMS) ordination of PLFA [$\log_{10}(x+1)$ of the relative
 687 abundance of individual PLFAs]. Average (\pm SE, $n = 3$) NMS axes scores of the sites. Axes are
 688 arbitrary; the closer the sample points are on the plot, the more similar they are in PLFA composition.
 689 Depth 0–25 cm (\circ), 25–50 cm (Δ), 50–100 cm (\square) and bottom layers (\diamond). The drained and natural
 690 sides of both the fen and bog sites differ significantly ($p < 0.05$) in a two-way factorial (drainage and
 691 depth) analysis (PERMANOVA, see Table 2).

692



693 **Fig. 4.** Mean (\pm SE, $n = 3$) relative abundance (% contribution to total microbial PLFAs) of (A) 16
694 MUFAs, (B) 18 MUFAs, (C) terminally branched FAs and (D) FA characteristic of fungi at different
695 depths in the studied fen and bog sites. Significant differences ($p < 0.05$) between drained and natural
696 sides in each depth and for each microbial group following independent sample t-tests are denoted by
697 an asterisk (*). For depth effects **Fn** = fen natural, **Fd** = fen drained, **Bn** = bog natural and **Bd** = bog
698 drained, while sites with significant depth effects (1-ANOVA, $p < 0.05$) are highlighted in bold.
699 Bottom = 135 – 160 and 105 – 130 for fen natural and drained, 217 – 242 and 245 – 270 for bog
700 natural and drained respectively.