

**This is an electronic reprint of the original article.
This reprint *may differ* from the original in pagination and typographic detail.**

Author(s): Aarnos, Hanna; Gélinas, Yves; Kasurinen, Ville; Gu, Yufei; Puupponen, Veli-Mikko;
Vähätalo, Anssi

Title: Photochemical mineralization of terrigenous DOC to dissolved inorganic carbon in ocean

Year: 2018

Version:

Please cite the original version:

Aarnos, H., Gélinas, Y., Kasurinen, V., Gu, Y., Puupponen, V.-M., & Vähätalo, A. (2018). Photochemical mineralization of terrigenous DOC to dissolved inorganic carbon in ocean. *Global Biogeochemical Cycles*, 32(2), 250-266.
<https://doi.org/10.1002/2017GB005698>

All material supplied via JYX is protected by copyright and other intellectual property rights, and duplication or sale of all or part of any of the repository collections is not permitted, except that material may be duplicated by you for your research use or educational purposes in electronic or print form. You must obtain permission for any other use. Electronic or print copies may not be offered, whether for sale or otherwise to anyone who is not an authorised user.

Photochemical mineralization of terrigenous DOC to dissolved inorganic carbon in ocean

Hanna Aarnos^{1,6}, Yves Gélinas², Ville Kasurinen^{3,4}, Yufei Gu¹, Veli-Mikko

Puupponen¹ and Anssi V. Vähätalo¹

Accepted Article

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1002/2017GB005698

Accepted Article

¹University of Jyväskylä, Department of
Biological and Environmental Science, P.O.
Box 35, 40014 Jyväskylä, Finland

²Geotop and Concordia University
Montreal, Department of Chemistry and
Biochemistry, Quebec, Canada

³University of Helsinki, Department of
Forest Sciences, P.O. Box 27, 00014
Helsinki, Finland

⁴University of Helsinki, Department of
Physics, P.O. Box 48, 00014 Helsinki,
Finland

⁶University of Helsinki, Department of
Environmental Sciences, P.O. Box 65, 00014
Helsinki, Finland

Abstract. When terrigenous dissolved organic carbon (tDOC) rich in chromophoric dissolved organic matter (tCDOM) enters the ocean, solar radiation mineralizes it partially into dissolved inorganic carbon (DIC). This study addresses the amount and the rates of DIC photoproduction from tDOC and the area of ocean required to photomineralize tDOC. We collected water samples from ten major rivers, mixed them with artificial seawater and irradiated them with simulated solar radiation to measure DIC photoproduction and the photobleaching of tCDOM. The linear relationship between DIC photoproduction and tCDOM photobleaching was used to estimate the amount of photoproduced DIC from the tCDOM fluxes of the study rivers. Solar radiation was estimated to mineralize $12.5 \pm 3.7 \text{ Tg C yr}^{-1}$ (ten rivers)⁻¹ or $18 \pm 8\%$ of tDOC flux. The irradiation experiments also approximated typical apparent spectral quantum yields for DIC photoproduction (ϕ_λ) over the entire lifetime of the tCDOM. Based on ϕ_λ s and the local solar irradiances in river plumes, the annual areal DIC photoproduction rates from tDOC were calculated to range from 52 ± 4 (Lena River) to $157 \pm 2 \text{ mmol C m}^{-2} \text{ yr}^{-1}$ (Mississippi River). When the amount of photoproduced DIC was divided by the areal rate, 9.6 ± 2.5 million km^2 of ocean was required for the photomineralization of tDOC from the study rivers. Extrapolation to the global tDOC flux yields 45 (31–58) Tg of photoproduced DIC yr^{-1} in the river plumes that cover 34 (25–43) million km^2 of the ocean.

1. Introduction

Riverine transport of terrigenous dissolved organic carbon (tDOC) from land to the ocean represents a major flux (248 Tg C yr⁻¹) in the carbon cycle [Hedges *et al.*, 1997; Bauer *et al.*, 2013; Cai, 2011]. Although riverine tDOC consists of thousands of compounds with a continuum of reactivities [Vähätalo *et al.*, 2010; Ward *et al.*, 2013; Riedel *et al.*, 2016], it can be conceptually divided into the biologically labile (20 ± 16%; mean ± SD) and the remaining non-labile (80 ± 16%; mean ± SD) fractions with different residence times in the ocean [Søndergaard and Middelboe, 1995; Holmes *et al.*, 2008; Fichot and Benner, 2014]. Close to river mouths, microbes mineralize labile riverine tDOC (Table 1) into dissolved inorganic carbon (DIC) within a couple of weeks [Søndergaard and Middelboe, 1995; Holmes *et al.*, 2008; Fichot and Benner, 2014]. The biologically non-labile tDOC (Table 1) which is rich in terrestrial chromophoric dissolved organic matter (tCDOM) has a longer residence time and is transported offshore [Vodacek *et al.*, 1997; Blough and Del Vecchio, 2002]. In the coastal ocean, tCDOM is primarily responsible for the absorption of solar ultraviolet and short-wavelength visible radiation that cause photochemical reactions [Johannessen *et al.*, 2003; Bélanger *et al.*, 2013; Taalba *et al.*, 2013]. These photochemical reactions are a major sink for non-labile tDOC and tCDOM [Kieber *et al.*, 1990; Miller and Zepp, 1995].

Solar radiation can efficiently reduce the tCDOM content of surface waters through photobleaching [Vodacek *et al.*, 1997; Vähätalo and Wetzel, 2004, 2008; Spencer *et al.*, 2009; Lalonde *et al.*, 2014; Riedel *et al.*, 2016]. As much as 70% of the tCDOM content of the mixed layer can be photobleached during the transport of tCDOM across the con-

tinental shelf over a single summer stratification period [Vodacek *et al.*, 1997]. Simulated solar radiation can simultaneously photobleach non-labile tCDOM and photomineralize 13–49% of non-labile tDOC from major rivers into DIC [Miller and Zepp, 1995; Lalonde *et al.*, 2014; Riedel *et al.*, 2016]. This photomineralizable fraction of tDOC (Table 1) can potentially be exploited to estimate the amount of tDOC that is photochemically mineralized to DIC upon complete photobleaching of tCDOM in the ocean. Although photochemical reactions have long been recognized as a major sink for tDOC [Kieber *et al.*, 1990; Miller and Zepp, 1995], the amount of photoproduct DIC from tDOC in the ocean is still poorly known.

The rates of photochemical transformation per unit of volume or area (Table 1) [Zhang *et al.*, 2006; White *et al.*, 2010; Aarnos *et al.*, 2012; Powers *et al.*, 2015]) and the photochemical reactivity of DOC (Table 1) [Zhang *et al.*, 2006; Bélanger *et al.*, 2006; White *et al.*, 2010; Aarnos *et al.*, 2012] decrease offshore. Despite this decrease, the amount of phototransformed tDOC within river plumes (Table 1) increases towards the distal part of the plumes [Vodacek *et al.*, 1997; Hernes and Benner, 2003; Medeiros *et al.*, 2015]. The increase results from the mixing of river water and seawater as (i) the mixing of tCDOM-rich river water with low-CDOM seawaters decreases the photochemical turnover time of tDOC per unit of area along offshore transport [Vodacek *et al.*, 1997], and (ii) the advection of tDOC increases the surface water area available for the photochemical transformations. Typically, only 3.7% of the total land area is available for photochemical transformation of tDOC [Verpoorter *et al.*, 2014], and this percentage is even lower for catchments dominated by rivers (e.g., the Amazon, Congo, Ganges-Brahmaputra, Mekong, Mississippi). For example, photochemical transformation of tDOC plays a minor role in the Amazon

River itself [Amon and Benner, 1996], but its importance is remarkable in the river plume [Medeiros et al., 2015].

Currently, the amount of phototransformed DOC has been estimated for arbitrary sections of coastal ocean [Miller et al., 2002; Bélanger et al., 2006; Fichot and Benner, 2014], and these estimates often include the phototransformation of both tDOC and marine DOC [Miller et al., 2002; Bélanger et al., 2006; Aarnos et al., 2012]. The distribution of tDOC in the ocean can be traced with different proxies (e.g., with optical, molecular and isotopic signatures) [Fichot and Benner, 2014; Medeiros et al., 2017], but discriminating between the photochemical transformation of tDOC and that of marine DOC is challenging. In the Baltic Sea for example, the estimated amount of photomineralized DOC exceeds the riverine flux of photoreactive tDOC to the coastal waters [Aarnos et al., 2012], suggesting that solar radiation photomineralizes both tDOC and marine DOC. To estimate the amount of phototransformed tDOC in an ocean, it is important to account for the phototransformation of tDOC across the entire river plume, and in particular at the distal parts of river plumes, which are recognized as hotspots for the photochemical transformation of tDOC [Vodacek et al., 1997; Hernes and Benner, 2003; Medeiros et al., 2015]. There is an important gap in knowledge regarding the location and rates of tDOC photomineralization and the area of ocean surface water required to photomineralize tDOC (Table 1).

In this study, we estimate the amount of photomineralized tDOC and the area of surface ocean required to oxidize the photoreactive fraction of riverine tDOC delivered from major rivers, assuming that photobleaching is the only sink for tCDOM. To obtain these estimates, we conducted laboratory irradiations with water samples collected from ten major

ivers and from the St Lawrence Estuary. We then calculated the amounts of photomineralized DIC, the areal rates and the area of the river plumes where photomineralization takes place.

2. Material and Methods

2.1. Sample collection

The large rivers selected for this study (Table 2) drain 23% of the land area on five continents, contribute to 33% of the freshwater discharge and 28% of the tDOC flux to the ocean [Cauwet, 2002; Milliman and Farnsworth, 2011; Raymond and Spencer, 2014]. The St. Lawrence Estuary was also sampled for the examination of photochemical transformation of DOC across its freshwater end to a salinity of 31 PSU (Table 2).

The samples from the St. Lawrence Estuary (1-L; Table 2) were collected with Niskin bottles at a depth of 3 m, filtered (precombusted 0.45 μm GF/F filter), stored in amber glass bottles and sent to Helsinki, Finland. For collection of river water samples (40-L), we shipped empty polyethylene containers (cleaned with detergent, 6% HCl, and ion-exchanged water, Millipore Milli-Q) to the local collaborators (see the acknowledgements).

The samples were collected mostly from the main stem of rivers close to the river mouth, where the seasonal fluctuations in the quality of tDOC are expected to be the smallest (Creed et al. 2013). The river water samples were collected during the season of high discharge, which is the most important contributor to the annual tDOC flux (Table 2).

The local collaborators filled the containers by immersing them below the surface in the center of the main stream, except for the Mississippi and the St. Lawrence River water samples that were collected with Niskin bottles at a depth of 3 m. The river water samples were shipped as collected, in the dark, with a 32 d median transport time between the

sampling sites and our laboratory in Helsinki. During transport (Table 2), indigenous microbes consumed the labile fraction of tDOC (e.g., 19% of DOC in the St. Lawrence River sample [Lalonde *et al.*, 2014]). After arrival, the Rio Negro and Rio Solimões samples were mixed in a 1:4 ratio to obtain a representative sample of the Amazon River. The two samples from the Congo River were mixed in the same manner (1:1 ratio). The river waters were filtered through 1- μm cartridges (a quick-rinsed membrane cartridge, Nuclepore, or the double layer of a track-etch polyester membrane filter cartridge, Graver Technologies) and kept in the dark at 5 °C before the photochemical experiments.

Aliquots of the same river water samples have been examined earlier (Tables S2, S3) to show that the chromophores of iron can contribute to the tCDOM (several percent of tCDOM in the Congo and Amazon Rivers for example) [Xiao *et al.*, 2013], and that dissolved black carbon (DBC) contributes about 10% to tDOC in all rivers [Jaffé *et al.*, 2013]. Photomineralization leads to the preferential removal of the ^{12}C -isotope from tDOC [Lalonde *et al.*, 2014] and, together with biodegradation, to a decrease in the contribution of DBC to tDOC to a level comparable to the concentrations measured in the deep ocean [Riedel *et al.*, 2016]. The DOM in the river water samples consisted of at least 7838 distinct molecular formulas, of which about 1/3 were detected in every waters sample [Wagner *et al.*, 2015].

2.2. Photochemical experiments with non-labile DOC

Irradiation experiments were conducted to estimate the typical photochemical reactivity of tDOC across its entire lifetime during transport from river mouths to the distal edge of river plumes. Our experimental design addressed three major hypotheses. (i) Microbes consume biologically labile tDOC close to river mouths and photochemical transformations

concern primarily biologically modified non-labile tDOC further offshore. Therefore, the irradiation experiments targeted non-labile tDOC. (ii) The mixing of tDOC with seawater may change the photochemical reactivity of tDOC [Minor *et al.*, 2006; White *et al.*, 2010] and flocculate some terrigenous components (e.g., iron) present in river water [Sholkovitz *et al.*, 1976; Asmala *et al.*, 2014]. Therefore, the river water samples were mixed with artificial DOC-free seawater and filtered prior to the irradiation experiments. (iii) The photochemical reactivity of tDOC decreases with the photobleaching of tCDOM along its offshore transport and becomes negligible upon complete photobleaching of tCDOM at the distal edges of river plumes. In order to approximate the average photochemical reactivity of tDOC, the irradiation times were selected to photobleach about half of the tCDOM.

For the photochemical experiments, 1- μm filtered river water was mixed (1:1 ratio) with artificial seawater (Kester [1967]) and further filtered through 0.2- μm membrane filters (Supor-200, Pall). In the mixtures, the relative abundance of the different ions was similar to seawater but the salinity (16 PSU) was about half that of seawater. The seven St. Lawrence Estuary samples were not mixed with artificial seawater but apart from this, they were treated in the same way as the other water samples. Immediately following filtration, the background DIC was removed in custom-made gas exchange bottles (500 mL) by acidifying the water samples to pH 3 with 1 M HCl and bubbling them with CO₂-free air [Miller and Zepp, 1995]. The pH of water samples was then adjusted with 0.1 M Na₂B₄O₇ buffer to ca. pH 7–8, as found in the ocean (Johannessen and Miller [2001]). Aliquots were closed without any headspace into ground-glass stoppered 12-mL tubes: two quartz tubes to be irradiated (one quartz tube for the St. Lawrence Estuary

samples) and two 12-mL glass tubes either wrapped in aluminum foil (dark control) or kept in the dark in cold (0 °C) water (initial).

The quartz and the aluminum foil-wrapped dark tubes were exposed to simulated solar radiation (Atlas Suntest CPS+ Solar simulator, 765 W m⁻²) in a water bath regulated to 21 °C with a Lauda RE 112 thermostat (Supporting Information). The dark controls were placed on the bottom of water bath, while the quartz tubes were immersed about 6 cm below the water surface and received simulated solar radiation corresponding to 23–25 days of the mean UV radiation received by the Earth's surface, when calculated from the Earth's annual global mean energy budget [*Kiehl and Trenberth, 1997*], and accounting for the spectral irradiance of Suntest CPS+ and the ASTM G173-03 (American Society for Testing and Materials Standard G-173-03 derived from SMARTS version 2.9.2) solar reference spectrum.

After the irradiation, the concentration of DIC was determined in the initial, dark control, and irradiated water by injecting 750 μL (7 injections per sample) into an acid bath and measuring the evolved CO₂ [*Salonen, 1981*]. The photochemical production of DIC was calculated as the difference between the DIC concentrations in the irradiated and the dark control samples. No significant difference was found between the DIC concentration in the initial samples and the dark controls, indicating negligible interference by dark chemical or biological processes on DIC photoproduction. The rest of the initial, dark and irradiated samples were stored in the dark at 5 °C for the measurement of CDOM.

2.3. Absorption spectrum for CDOM

The spectral absorption by CDOM was measured with a UV-VIS spectrophotometer (Shimadzu UV-2550) using a 5-cm quartz cuvette. The apparent absorbance spectrum of

samples and blanks (Millipore Milli-Q water) was obtained from two scans between 190 and 800 nm in 1-nm steps against an empty reference cell holder. The blank was subtracted from the sample and the resulting absorbance of CDOM ($A_{CDOM,\lambda}$) was converted to an absorption coefficient (a_λ) by $2.303 A_{CDOM,\lambda} / 0.05$ [Bricaud *et al.*, 1981]. To avoid potential bias resulting from scattering, the CDOM absorption spectra were corrected according to the equation proposed by Bricaud *et al.* [1981]: $a_{CDOM,\lambda} = a_\lambda - a_{700} (\lambda/700)$, where $a_{CDOM,\lambda}$ is the corrected CDOM absorption coefficient at a wavelength λ ; and a_λ and a_{700} refer to uncorrected CDOM absorption coefficients at λ and at 700 nm, respectively. The photobleaching of CDOM during the irradiation was calculated as the difference in absorption coefficients of CDOM at 300 nm ($a_{CDOM,300}$; m^{-1}) between the dark controls and irradiated waters.

2.4. Areal rates of DIC photoproduction based on apparent quantum yields (AQY)

For the determination of the AQY spectrum for DIC photoproduction (ϕ_λ), the measured DIC photoproduction was divided by the number of photons absorbed by CDOM during the irradiation (see Aarnos *et al.* [2012]; details in Supporting Information). We assumed an exponential increase in ϕ_λ with decreasing wavelength:

$$\phi_\lambda = ce^{-d\lambda} \quad (1)$$

where c ($\text{mol C mol photons}^{-1}$) and d (nm^{-1}) are positive constants, and λ is the wavelength (nm). The parameters c and d in Eq. 1 were iterated by relating the amount of photoproduct (DIC) to the number of photons absorbed by CDOM through an unconstrained nonlinear optimization (the `fminsearch` function of Matlab R2013a) as in [Vähätalo *et al.*, 2000, 2003, 2011; Vähätalo and Wetzel, 2004; Vähätalo and Zepp, 2005;

Vähätalo and Järvinen, 2007; Aarnos *et al.*, 2012], but in this work a Monte Carlo approach was used to assess the error in fittings and to select the best ϕ_λ from 1000 fittings ([Gu *et al.*, 2017], Supporting Information).

An areal DIC photoproduction rate in river plumes (pr ; mol C m⁻² d⁻¹) was calculated according to Vähätalo *et al.* [2000]:

$$pr = \int_{\lambda_{min}}^{\lambda_{max}} \phi_\lambda Q_\lambda (a_{CDOM,\lambda} a_{tot,\lambda}^{-1}) d\lambda \quad (2)$$

where ϕ_λ is the determined AQY spectrum for DIC photoproduction; Q_λ represents the spectrum of mean annual photon flux density at the river plumes (mol photons m⁻² d⁻¹ nm⁻¹; Figure S7); $a_{CDOM,\lambda} a_{tot,\lambda}^{-1}$ (dimensionless) is a contribution of riverine tCDOM to the total absorption coefficient $a_{tot,\lambda}$. $a_{CDOM,\lambda} a_{tot,\lambda}^{-1}$ was approximated to be 1 (see Discussion for the associated error). The integration through λ_{min} to λ_{max} accounts for the wavelengths (300–600 nm) contributing to the DIC photoproduction. Annual DIC photoproduction rates were computed by multiplying the daily rate of pr (Eq. 2) by 365. The Q_λ for each river was calculated according to the computed mean annual global radiation at the ocean in front of the rivers over the years 1984 to 2000 [Hatzianastassiou *et al.*, 2005] and the spectral reference solar irradiance (ASTMG173-03) from where the spectral form for Q_λ was adapted (details in Supporting Information).

2.5. The amount of photoproduced DIC from tCDOM flux

The annual amount of photoproduced DIC (g C yr⁻¹ river⁻¹) from the tCDOM flux of each river was estimated using a linear regression between DIC photoproduction (mol C m⁻³) and photobleaching of tCDOM (m⁻¹) at a wavelength of 300 nm measured during the irradiation experiments (Figure 1). The regression coefficient (mol C m⁻², Figure

1) quantifies the amount of photoproducted DIC per photobleached tCDOM at 300 nm.

When the regression coefficient is multiplied by the amount of photobleached tCDOM from each river ($\text{m}^2 \text{yr}^{-1} \text{river}^{-1}$), the product expresses the amount of photoproducted DIC from each river ($\text{g C yr}^{-1} \text{river}^{-1}$). Carbon fluxes are either given as moles or grams and include always adequate conversion between the units. Photobleaching was assumed to be the only sink for tCDOM discharged in the ocean [Nelson and Siegel, 2013; Vähätalo and Wetzel, 2008; Vodacek et al., 1995; Vähätalo and Wetzel, 2004]. We assume that the amount of photobleached tCDOM from each river equals the tCDOM flux of each river ($\text{m}^2 \text{yr}^{-1} \text{river}^{-1}$; see Discussion for the associated uncertainty).

The annual tCDOM fluxes for the Mississippi, St. Lawrence and Lena Rivers were taken from earlier studies [Stedmon et al., 2011; Spencer et al., 2013]. The published tCDOM fluxes reported at 350 nm or 375 nm were converted to the 300 nm wavelength using the following spectral slope coefficients, measured from our water samples: $16.4 \mu\text{m}^{-1}$ for the Lena River [Stedmon et al., 2011], $17.6 \mu\text{m}^{-1}$ for the Mississippi River, and $16.3 \mu\text{m}^{-1}$ for the St. Lawrence River. For the other rivers, tCDOM fluxes were calculated by multiplying the measured absorption coefficient at 300 nm ($a_{CDOM,300}$) with the annual water discharge [Milliman and Farnsworth, 2011].

2.6. Area needed for photomineralization of annual riverine tDOC flux in the ocean

To estimate the area of ocean surface waters required to photochemically mineralize the photoreactive fraction of riverine tDOC discharged by each river, the amount of photoproducted DIC from the tCDOM flux of each river ($\text{g C yr}^{-1} \text{river}^{-1}$) was divided by the corresponding areal DIC photoproduction rate ($\text{mol C m}^{-2} \text{yr}^{-1}$). The areas were

converted into semicircles. The radius of a semicircle approximates the distance from the river mouth required to photomineralize photoreactive tDOC flux, but without accounting for local geography or currents that disperse tDOC.

3. Results

3.1. Photochemical production of DIC and photobleaching of tCDOM

The irradiations of river water samples (Table 2) mixed with artificial seawater produced DIC from terrigenous DOC (tDOC) and bleached terrigenous CDOM (tCDOM; Rivers in Table 3), at rates that varied between $1.9 \pm 0.1 \text{ mmol C m}^{-3} \text{ 43 h}^{-1}$ (Ganges-Brahmaputra River; mean \pm SD) and $42.5 \pm 0.3 \text{ mmol C m}^{-3} \text{ 39 h}^{-1}$ (Congo River) (Table 3). The irradiations photobleached on average $45 \pm 3\%$ of the tCDOM at 300 nm ($a_{CDOM,300}$; Table 3). When expressed as an absorption coefficient at 300 nm, photobleaching ranged from $0.54 \pm 0.04 \text{ m}^{-1}$ (Ganges-Brahmaputra River) to $11.56 \pm 0.05 \text{ m}^{-1}$ (Congo River; Table 3).

When sterile-filtered water samples collected from the St. Lawrence Estuary (Table 2) were exposed to the simulated solar radiation, DIC photoproduction and photobleaching were typically highest for the freshwater end-member and decreased with increasing salinities (Table 3). In both the St. Lawrence Estuary and the river water samples mixed with artificial seawater (Table 3), the DIC photoproduction was linearly dependent on tCDOM photobleaching despite large differences in the CDOM contents of water samples (Figure 1). The regression coefficient was $0.00402 \pm 0.00018 \text{ mol C m}^{-2}$ (\pm standard error) and R^2 was 0.96 for the river water samples (Figure 1)

3.2. The amount of photoproducted DIC from the tCDOM flux of each river

The annual amount of photoproducted DIC from tCDOM flux in each river plume ($\text{Tg C yr}^{-1} \text{ river}^{-1}$) was estimated from the tCDOM fluxes (Table 4) and the regression coefficient between DIC photoproduction and tCDOM photobleaching (Figure 1). The photoproduction of DIC of the Amazon ($6.98 \pm 1.08 \text{ Tg C yr}^{-1}$) and the Congo ($3.04 \pm 0.49 \text{ Tg C yr}^{-1}$) Rivers contributed over 80% of the total annual amount of photoproducted DIC for the 10 rivers examined ($12.48 \pm 2.10 \text{ Tg C yr}^{-1} \text{ 10 rivers}^{-1}$; Table 4). The photomineralized fraction of riverine tDOC flux ranged from $3 \pm 2\%$ to $38 \pm 7\%$ among the studied rivers and the weighted average was $18 \pm 4\%$ for the total flux of the 10 rivers examined (Table 4).

3.3. Apparent quantum yields

The photoproducted DIC (Table 3) was normalized against the photons absorbed by tCDOM for the determination of the spectral AQY for DIC photoproduction (ϕ_λ ; Eq. 1; Table 5). Table 5 reports the parameters c and d of Eq. 1 for the calculations of spectral ϕ_λ illustrated in Figure 2. The ϕ_λ was highest for the Mississippi River tDOC, and lowest for the Ganges-Brahmaputra River tDOC, whether taken from the spectra (Figure 2) or the AQYs at 330 nm (ϕ_{330} , Table 5). The value of ϕ_{330} was similar in the St. Lawrence River water mixed with artificial seawater and its freshwater end-member when accounting for the coefficient of variation related to the determination of ϕ_λ (Table 5). In the St. Lawrence Estuary, the ϕ_{330} s decreased with increasing salinity to about half of the value for the freshwater end-member (ϕ_{330} s in Table 5 and Figure 3).

3.4. Annual areal DIC photoproduction rates

To calculate the annual areal DIC photoproduction rate in a river plume (pr in Table 5 and Figure 4), ϕ_{λ} was multiplied by the local annual solar photon flux density spectrum (Eq. 2; Figure S7). These rates approximate the areal DIC photoproduction from tDOC at sites where half of the tCDOM has been photobleached. Similar to $\phi_{\lambda s}$, the areal DIC photoproduction rate from tDOC was highest for the Mississippi River (157 ± 2 mmol C m^{-2} yr^{-1} ; Table 5, Figure 4). The lowest rate (52 ± 4 mmol C m^{-2} yr^{-1}) was calculated for the tDOC discharged by Lena River to the Laptev Sea, where the intensity of solar irradiation is low (Table 5, Figure 4, Figure S7).

3.5. Area needed for the photomineralization of tCDOM flux in the ocean

A division of the annual amount of photoproduced DIC from the tCDOM flux discharged by each river (Tg C yr^{-1} river $^{-1}$; Table 4) by the corresponding areal DIC photoproduction rate (mol C m^{-2} yr^{-1} ; Table 5, Figure 4) provides an estimate of the ocean surface area required to completely photomineralize the photomineralizable fraction of tDOC (Table 6). The area was largest for the Amazon River (4.3 ± 1.1 million km^2) and smallest for the St. Lawrence River ($76\,000 \pm 19\,000$ km^2 ; Table 6). To estimate how far offshore the plume of riverine tCDOM extends until its photoreactive components are photochemically mineralized to DIC, we assumed there were semicircle-shaped river plumes, whose radii correspond to the distance required to completely photomineralize the photoreactive tDOC. These distances ranged from 221 ± 54 km (St. Lawrence River) to 1652 ± 422 km (Amazon River; Table 6). Although the results in Table 6 represent a simplified view of the photomineralization and dispersion of tDOC, they indicate that the photomineralization of tDOC fluxes from large rivers takes place mostly within a few

hundreds kilometers (383 ± 160 km; median \pm SD) from the river mouths for most rivers studied in this work, except for rivers with the two largest discharges (the Amazon and Congo Rivers), and the Arctic River (Lena River).

4. Discussion

The discussion estimates first the annual amount of photoproducted DIC from the tDOC fluxes (4.1.–4.6.). The second theme assesses the annual areal rates of DIC photoproduction in river plumes (4.7.). At the end, the discussion evaluates the area of ocean needed to photomineralize tDOC (4.8.).

4.1. The amount of photochemically mineralized tDOC in the ocean

This study estimates that photochemical reactions mineralize 12.5 ± 2 Tg C yr⁻¹, or $18 \pm 4\%$, of the tDOC flux in the plumes of the ten rivers (Table 4). The uncertainty of estimates reported in Table 4 is derived from the propagation of the analytical errors (Table 5) and the regression statistics (Figure 1). Below we evaluate other uncertainties that are associated with our calculations and provide an estimate for the annual DIC photoproduction from tDOC in the global ocean.

4.2. Photobleaching of tCDOM — a good proxy for DIC photoproduction

In this study, the amounts of photochemically mineralized tDOC in river plumes (Table 4) are estimated using the regression between DIC photoproduction and photobleaching (Figure 1). This relationship has high R^2 (0.96) and the standard error of regression coefficient is small (4.5%; Figure 1). In comparison, the coefficient of variation (29 %) and the standard error ($21 \mu\text{mol C mol photons} = 9.2\%$) is larger for the mean ϕ_{330} ($230 \mu\text{mol C mol photons}$) determined for the same ten samples (Table 5). This comparison suggests

that DIC photoproduction from tDOC is more closely related to the photobleaching of tCDOM than to the amount of photons (or energy) absorbed by tCDOM, which is used to calculate the AQYs. Photobleaching is tightly linked to the photoproduction of CO₂ (this study) and other low molecular weight carbonyl compounds [Kieber *et al.*, 1990; Miller and Moran, 1997; Bertilsson and Tranvik, 2000] indicating that the photoproduction of carbonyl compounds is mechanistically coupled to photobleaching. As conjugated carbonyl groups are chromophores in DOM [Korshin *et al.*, 1997; Chen *et al.*, 2002], the photochemical scission of carbonyl compounds (such as DIC, CO, and organic acids) represents a CDOM photobleaching pathway. Photobleaching and DIC photoproduction associate with the loss of aromatic groups [Riedel *et al.*, 2016], which are the most important chromophores in DOM [Korshin *et al.*, 1997; Chen *et al.*, 2002]. For example, solar radiation can mineralize the carbon atoms of aromatic rings to DIC, which is a direct evidence for a simultaneous photobleaching and DIC photoproduction [Vähätalo *et al.*, 1999]. The mechanistic coupling between the photoproduction of carbonyl compounds and the photochemical destruction of chromophoric moieties in DOM makes photobleaching a useful proxy for the photoproduction of DIC and other carbonyl compounds.

4.3. Photobleaching as a sink of tCDOM

Our estimate for the amount of photomineralized tDOC is based on the assumption that photobleaching is the only sink for riverine tCDOM. Experimental irradiations can completely photobleach tCDOM from the rivers examined in this study [Miller and Zepp, 1995; Spencer *et al.*, 2009; Lalonde *et al.*, 2014; Riedel *et al.*, 2016]. Photobleaching is also the primary sink of CDOM in the mixed layer of surface ocean waters [Vodacek *et al.*, 1997; Nelson *et al.*, 1998; Vähätalo and Wetzel, 2004; Shank *et al.*, 2010; Swan *et al.*, 2012].

A fraction of the tCDOM flux can also be transported below the mixed layer to remain in dark waters above a pycnocline for a short (<1 yr) period of times [Vodacek *et al.*, 1997; Fichot and Benner, 2014], but the seasonal deepening of mixed layer eventually re-introduces this CDOM into the mixed layer where it is eventually photobleached [Nelson *et al.*, 1998]. While about 1.7 Tg C yr^{-1} of tDOC discharged by Arctic rivers (or 0.7% of global tDOC flux) is exported into the North Atlantic Deep Water and is therefore shielded from photobleaching for hundreds of years [Benner *et al.*, 2005], the remaining 99.3% is susceptible to photobleaching, likely within the first year in the ocean surface waters. Indeed, tCDOM can be rapidly transported (within weeks or months) to the edge of river plumes [Coles *et al.*, 2013; Fichot and Benner, 2014] where photobleaching of CDOM can be completed within days-weeks [Vodacek *et al.*, 1997; Nelson *et al.*, 1998; Swan *et al.*, 2012].

Other competing sinks for the non-labile tDOC and tCDOM pools also exist, in addition to photobleaching. Biodegradation of non-labile tDOC can also reduce the abundance of tCDOM and thus cause biobleaching. Microbes can biobleach tCDOM in fresh waters [Vähätalo and Wetzel, 2004, 2008; Koehler *et al.*, 2012], but the biobleaching rates decrease upon mixing of tCDOM with seawater (Figures. 4–5 in Vähätalo and Wetzel [2004]). In the ocean, microbial metabolism is often considered to be as a source rather than a sink of CDOM [Jørgensen *et al.*, 2014]. In the deep dark ocean, the estimated turnover time of CDOM is 634 years [Catalá *et al.*, 2015], indicating very slow biobleaching rates. Thus, it is not always clear whether microbial metabolism is a sink or a source of tCDOM in marine waters, but it is likely that microbes biobleach tCDOM when they consume non-labile tDOC rich in tCDOM [Vähätalo and Wetzel, 2004, 2008; Koehler *et al.*, 2012].

Microbes mineralized 12–25% of non-labile tDOC from our river water samples during one year (Table S3). If this consumption of tDOC is associated with biobleaching [Koehler *et al.*, 2012], it may cause a systematic reduction (roughly estimated at about <20%) in the amount of non-labile tCDOM to be photobleached and thus in the DIC production from tDOC in the ocean.

4.4. tCDOM fluxes

Our DIC production estimates from tDOC in river plumes also depend on tCDOM fluxes. For most rivers, tCDOM fluxes were estimated by multiplying the amount of non-labile tCDOM measured in this study ($a_{CDOM,300}$; Table 4) and the annual water discharges (Table 4 from Milliman and Farnsworth [2011]). For the Lena, Mississippi and St. Lawrence Rivers, we used published tCDOM fluxes. If we would have treated these rivers like the others, our simple multiplication method would have given a collective tCDOM flux of $20.7 \text{ Tm}^2 \text{ yr}^{-1}$, which corresponds to 56% of the literature based estimate for these rivers ($37.2 \text{ Tm}^2 \text{ yr}^{-1}$ [Stedmon *et al.*, 2011; Spencer *et al.*, 2013]. tCDOM fluxes are not available for the Amazon and Congo Rivers, but the absorption coefficients of CDOM at 300 nm ($a_{CDOM,300}$) measured in this study can be compared to those obtained in previous work. For the Congo River, our $a_{CDOM,300}$ is 48.4 m^{-1} (Table 4), corresponding to 53% of the value of 92.09 m^{-1} reported by [Spencer *et al.*, 2009]. For the Amazon River, we used $a_{CDOM,300}$ value of 22.9 m^{-1} , which is 8.5% larger than the 21.1 m^{-1} value measured at the mouth of the river [Cao *et al.*, 2016]. A use of $a_{CDOM,300}$ estimates from literature [Spencer *et al.*, 2009; Cao *et al.*, 2016] would have lead to a a closer calculated DIC production from tDOC for the Congo ($5.78 \text{ Tg C yr}^{-1}$) and Amazon ($6.43 \text{ Tg C yr}^{-1}$) Rivers compared to the values listed in Table 4. The comparison of our rough estimates

of tCDOM fluxes to these available in the literature [*Stedmon et al.*, 2011; *Spencer et al.*, 2013, 2009; *Cao et al.*, 2016] yields differences that range from a 47% underestimation to a 9% overestimation in tCDOM fluxes and thus in the amounts of photoproduct DIC.

These differences in our tCDOM fluxes may come from natural interannual or seasonal variability in tCDOM fluxes. It is likely that biobleaching decreased the $a_{CDOM,300}$ values during the transportation of the samples. If this biobleaching was similar to the loss of tDOC during the transportation ([*Lalonde et al.*, 2014]), the amounts of photochemically produced DIC from tDOC are underestimated by about <20%. The impact of biobleaching on our estimated DIC production rates from tDOC (Table 4) is likely low, because the overestimate caused by biobleaching in the ocean is probably compensated by the underestimate derived from biobleaching during the transportation of our samples.

4.5. Uncertainty and global estimate for the amount of photochemically mineralized tDOC in the ocean

In this study, the uncertainty in tCDOM fluxes exceeds the analytical and statistical error reported in Table 4. We estimate that the uncertainty in tCDOM fluxes increases the uncertainty in the calculated amounts of tDOC photomineralized by about $\pm 30\%$ when examined collectively for all rivers, but uncertainty for individual rivers can be even larger. Accounting for this uncertainty, the amount of photochemically mineralized tDOC amounts to $12.5 \pm 3.7 \text{ Tg C yr}^{-1}$, or $18 \pm 8 \%$ of the tDOC flux from the ten rivers.

As the tCDOM fluxes represent the largest source of uncertainty in our estimate, we also used another tCDOM flux-independent method to estimate DIC photoproduction from non-labile tDOC. The photomineralizable fraction of tDOC was measured in a previous work for the same river water samples examined in this study [*Lalonde et al.*, 2014;

Riedel et al., 2016]. The photomineralizable fraction of non-labile tDOC ranged from 16% to 49% in the different river water samples in a laboratory study (Table S2). If the same fraction of non-labile tDOC is photochemically mineralized in the ocean, the DIC photoproduction from non-labile tDOC in the ten rivers amounts to 18.7–19.7 Tg C yr⁻¹ (Table S2). Correcting for the competing biobleaching of tCDOM in the ocean (20% yr⁻¹; *Riedel et al.* [2016]), the DIC photoproduction from the non-labile tDOC of ten rivers decreases to 15–15.8 Tg C yr⁻¹. These estimates are larger but within the uncertainty of the estimate based on tCDOM fluxes (12.5 ± 3.7 Tg C yr⁻¹). The possible differences between the results obtained using the two methods may arise from differences in experimental conditions. In this study, (i) the non-labile tDOC was transferred into a seawater matrix, (ii) irradiated at 21 °C, (iii) the mineralization of tDOC was measured as the production of DIC and related to photobleaching of $a_{CDOM,300}$. Finally, (iv) the latter relationship was multiplied by the tCDOM flux to obtain 12.5 Tg C yr⁻¹ DIC photoproduction from non-labile tDOC (Table 6). *Lalonde et al.* [2014] and *Riedel et al.* [2016], (i) used filtered river water directly, (ii) irradiated at 25°C, (iii) measured the mineralization as the loss of DOC, (iv) which was directly related to the non-labile tDOC flux (Table S2). The potential explanations for the differences in the estimates (Table S2 and Table 4) are (i) matrix effects (river water vs. seawater), (ii) the difference in the irradiation temperature, (iii) potential irradiation-induced flocculation of tDOC in *Lalonde et al.* [2014] and *Riedel et al.* [2016] vs. true mineralization to CO₂ in this study, (iv) an error in the estimated tCDOM flux in this study.

To upscale our results for the ten rivers to global fluxes, we first evaluate the global representativeness of the rivers studied in this work. The volume averaged concentration

of DOC in the rivers examined in this study is 5.6 mg C L^{-1} ($= 6.91 \times 10^{13} \text{ g C yr}^{-1} / 1.226 \times 10^{13} \text{ m}^3 \text{ yr}^{-1}$ from Table 4), similar to the mean DOC concentration (5.75 mg C L^{-1}) in the world's rivers [Meybeck, 1982] and to the mean DOC concentration (5.7 mg C L^{-1}) measured in 7500 lakes distributed over six continents [Sobek *et al.*, 2007]. Four of the rivers examined in this study (Amazon, Congo, Paraná and Lena Rivers) account for the largest tDOC fluxes to the ocean [Raymond and Spencer, 2014]. In the same study, the Mississippi and Ganges-Brahmaputra Rivers are also listed among the top ten rivers with the highest tDOC fluxes. The others (Yangtze, Mekong and Danube Rivers) are within the top 20. The rivers examined in this study span from high to low latitudes and drain major biomes on five continents, excluding Australia and Antarctica, which contribute little to the global tDOC flux. As the rivers examined in this study are responsible for 28% of tDOC flux to the global coastal ocean ($61.1 \text{ Tg C yr}^{-1}$ (Table 4) vs. 246 Tg C yr^{-1} [Cai, 2011]), we upscaled the cumulative results from our study rivers by multiplying them by 3.56 ($= 0.28^{-1}$) to estimate the global fluxes, leading to a DIC photoproduction from non-labile tDOC in the global ocean of 45 (31–58) Tg C yr^{-1} .

4.6. Photochemical mineralization of tDOC in the ocean vs. other global tDOC fluxes

This study estimates that the amount of tDOC mineralized by solar radiation is higher in the ocean ($45 \text{ (31–58) Tg C yr}^{-1}$) than in inland waters ($24 \text{ (13–35) Tg C yr}^{-1}$; Koehler *et al.* [2014]; Figure 5, Supporting Information). Photochemical reactions mineralize a larger percentage of imported tDOC in the ocean ($18\% = 45/246 \text{ Tg C yr}^{-1}$) than in the inland waters ($3\% = 24/794 \text{ Tg C yr}^{-1}$; Figure 5, Supporting Information). Earlier studies [Søndergaard and Middelboe, 1995; Holmes *et al.*, 2008] reported that $20 \pm 16\%$ of riverine

tDOC (or 49 (41–57) Tg C yr⁻¹; Figure 5) consists of labile tDOC and is biologically mineralized within a few weeks in the proximity of river mouths. Based on earlier one-year-long bioassays [Riedel *et al.*, 2016], 44 (25–49) Tg C yr⁻¹ of non-labile tDOC is expected to be mineralized biologically within its first year in the ocean (semilabile in Figure 5, Table S3, Supporting Information). Based on the ratios of photoproducted labile DOC to photoproducted DIC [Vähätalo *et al.*, 2003; Pullin *et al.*, 2004; Cory *et al.*, 2014], sunlight-induced reactions transform 36 (9–65) Tg C yr⁻¹ of non-labile tDOC into labile forms (Figure 5, Supporting Information). Thus, we estimate that photochemical reactions mineralize a total of 81 Tg C yr⁻¹, either directly (45 Tg C yr⁻¹) or through the production of labile forms (36 Tg C yr⁻¹; Figure 5). This amount of photochemical mineralization corresponds to 87% of tDOC mineralized by microbes during the first-year residence of tDOC in the ocean (49 + 44 = 93 Tg C yr⁻¹; Figure 5). The corresponding value is 67% for tDOC in the Mississippi River plume during its 2–3 month residence time on the Louisiana shelf, where ca. 50% of tDOC is mineralized [Fichot and Benner, 2014]. Our budget indicates that 71% of tDOC ((93 + 81)/246; Figure 5) is mineralized during its first year in the ocean. In Figure 5, refractory tDOC (72 (17–140) Tg C yr⁻¹ or 29% of the annual tDOC flux to the ocean) refers to a fraction of tDOC having a residence time of more than one year in the ocean. This refractory tDOC is either slowly metabolized by microbes, transformed by indirect photochemical reactions [Mopper *et al.*, 1991] and/or transported (1.7 Tg C yr⁻¹) to the deep ocean [Benner *et al.*, 2005].

4.7. The annual areal rates of DIC photoproduction from tDOC in river plumes

Our annual areal rates calculated based on Eq. 2 (Table 5, Figure 4) are meant to provide an average value for the DIC photoproduction rate ($\text{mol C m}^{-2} \text{ yr}^{-1}$) from tDOC over the entire lifetime of tCDOM in river plumes. The rate estimates are based on ϕ_{λ} s. Therefore, we first discuss about the ϕ_{λ} and later about the rates themselves.

Our spectral AQY for DIC photoproduction from tDOC approximates an average photoreactivity of tDOC from its entrance to estuary to its complete removal by photobleaching. This differs from earlier studies, which have characterized ϕ_{λ} at the site of sample collection [Vähätalo *et al.*, 2000; Vähätalo and Wetzel, 2004; Koehler *et al.*, 2014; Groeneveld *et al.*, 2016; Koehler *et al.*, 2016; Gao and Zepp, 1998; Bélanger *et al.*, 2006; White *et al.*, 2010; Johannessen and Miller, 2001; Johannessen *et al.*, 2007; Aarnos *et al.*, 2012; Reader and Miller, 2012; Powers *et al.*, 2015]. Our experimental irradiations photobleached about half of tCDOM and characterize the photoreactivity of tDOC at those sites of river plumes, where solar radiation has photobleached about half of tCDOM. These sites would be for example just before the shelf break in Mid-Atlantic Bight [Vodacek *et al.*, 1997] and offshore from the shelf break in the plume of Mississippi River [Fichot and Benner, 2014]. Thus, our ϕ_{λ} characterize tDOC at the distal parts of river plumes that are considered as the most important sites of photochemical breakdown of tDOC [Vodacek *et al.*, 1997; Hernes and Benner, 2003; Medeiros *et al.*, 2015].

Our ϕ_{330} values (range of 129 to 335 $\mu\text{mol C mol photons}^{-1}$) fall into the lower range of reported ϕ_{330} s in lakes [Vähätalo *et al.*, 2000; Vähätalo and Wetzel, 2004; Koehler *et al.*, 2014; Groeneveld *et al.*, 2016; Koehler *et al.*, 2016], rivers and estuaries [Gao and Zepp,

1998; *Bélanger et al.*, 2006; *White et al.*, 2010], as well as coastal ocean [*Johannessen and Miller*, 2001; *Johannessen et al.*, 2007; *Aarnos et al.*, 2012; *Reader and Miller*, 2012; *Powers et al.*, 2015]. A possible explanation for our low values is the long irradiation time. Because ϕ_{λ} s decrease with cumulative irradiation doses [*Andrews et al.*, 2000; *Vähätalo and Wetzel*, 2004], our ϕ_{λ} s do not correspond to the highest ϕ_{λ} s for tDOC upon its discharge into an estuary. For example, our ϕ_{330} for tDOC from the Mississippi River (335 $\mu\text{mol C mol photons}^{-1}$) is expectedly lower than the corresponding pooled mean of 514 $\mu\text{mol C mol photons}^{-1}$ for DOC collected from the Gulf of Mexico close to the mouth of the Mississippi River and irradiated for a shorter time period (6 h; *Powers et al.* [2015]) compared to the present study (39 h).

Our median ϕ_{330} s for riverine tDOC is 234 $\mu\text{mol C mol photons}^{-1}$ and about half of the 479 $\mu\text{mol C mol photons}^{-1}$ used to estimate the amount of DIC photoproduction in inland waters [*Koehler et al.*, 2014]. High ϕ_{330} s correlate positively with CDOM, the specific UV absorption coefficient and the aromaticity of tDOC [*Koehler et al.*, 2016], which are typically highest for tDOC imported freshly from land to inland waters without earlier exposure to solar radiation [*Salonen and Vähätalo*, 1994; *Xiao et al.*, 2013, 2015]. Additionally, ferric iron associated to tDOC and acidity increase ϕ_{330} s in freshwaters [*Gu et al.*, 2017]. The mixing of the river waters (1:1) with artificial seawater followed by filtering, as done in this study, transfers tDOC from a freshwater to a seawater matrix, increases pH to the slightly alkaline values found in the ocean, and potentially flocculates photoreactive components such as iron [*Minor et al.*, 2006]. Our ϕ_{λ} for tDOC from the St. Lawrence River is independent of the matrix (freshwater or seawater) used for its determination. This is in contrast to a decrease in DIC photoproduction after mixing

swamp water with seawater [Minor *et al.*, 2006]. The difference may be explained by the difference in pH and the concentration of iron: swamp water had a pH of 4 and an Fe concentration of $29 \mu\text{mol L}^{-1}$ [Minor *et al.*, 2006], compared to a pH of 7.6 and $0.08 \mu\text{mol Fe L}^{-1}$ for the St. Lawrence River water [Xiao *et al.*, 2013]. It remains unclear whether the seawater matrix used in this study changed ϕ_λ compared to a freshwater matrix in the other rivers as this was not experimentally addressed, but nevertheless our experimental design addressed those potential changes.

In this study, ϕ_λ s decrease with salinity in the St. Lawrence Estuary, as in the Beaufort Sea [Bélanger *et al.*, 2006], in the Baltic Sea [Aarnos *et al.*, 2012], and in the St. Lawrence Estuary for the photoproduction of CO [Zhang *et al.*, 2006]. The dependence of ϕ_λ s on salinity has not always been observed (Powers *et al.* [2015]), or the dependence has included an abrupt change (White *et al.* [2010]). These earlier studies have been conducted in relatively small regions close to river mouths and ϕ_λ s have been determined with short irradiance times (8 h, White *et al.* [2010]; 6 h, Powers *et al.* [2015]). These ϕ_λ s possibly trace (i) the complex mixing of riverine tDOC and marine DOC close to river mouths, (ii) describe, e.g., the flocculation of iron at a certain salinity threshold or (iii) may also include additional (more local) sources of DOC such as coastal wetlands or sediment pore waters [White *et al.*, 2010; Powers *et al.*, 2015]. This study and observations in long transects (> few hundreds km) across coastal regions where tDOC is more or less conservatively mixed with marine DOC [Vodacek *et al.*, 1997; Bélanger *et al.*, 2006; Zhang *et al.*, 2006; Aarnos *et al.*, 2012] indicate two end-members in terms of photoreactivity: 1) tDOC at the mouth of river with higher ϕ_λ and 2) marine DOC of open ocean with lower ϕ_λ . The mixing of these two end-members together with a decrease in ϕ_λ due to

the cumulative exposure of tDOC to solar irradiance can explain the decrease of ϕ_λ with increasing salinity over large spatial scales across the entire span of river plumes.

In this study, the calculated areal rates of DIC photoproduction depend also on the $a_{CDOM} a_{tot}^{-1}$ (Eq. 2) in the ocean. When tCDOM has lost half of its original absorption in the ocean, it is still expected to dominate the absorption of photolytic solar radiation in river plumes [Johannessen *et al.*, 2003; Bélanger *et al.*, 2013; Taalba *et al.*, 2013], but it does not absorb all photolytic solar radiation as approximated in Eq. 2. For example, in the river plume of Mackenzie River, the $a_{CDOM} a_{tot}^{-1}$ at 330 nm (Eq. 2) is ca. 0.9 [Taalba *et al.*, 2013; Bélanger *et al.*, 2013]. In the Mid Atlantic Bight and in the Bering Sea, the ratio of CDOM to the attenuation coefficient at the UVR ranges from 0.86 to 0.97 in coastal to offshore water outside turbid bays [Johannessen *et al.*, 2003]. Spectral $a_{CDOM} a_{tot}^{-1}$ are poorly available for the river plumes examined in this study and therefore our simplification ($a_{CDOM} a_{tot}^{-1} = 1$) likely overestimates the areal DIC photoproduction rates by <10–15% [Johannessen *et al.*, 2003; Bélanger *et al.*, 2013; Taalba *et al.*, 2013].

In a strict sense, our DIC photoproduction rates refer to those produced from tDOC in the matrix of seawater, and are not directly comparable to those determined from samples collected from the coastal ocean that also include marine DOC [Bélanger *et al.*, 2006; Fichot and Benner, 2014; Powers *et al.*, 2015]. However, tDOC was likely the primary source for photoproduced DIC in the samples collected from the plume of the Mississippi River [Fichot and Benner, 2014; Powers *et al.*, 2015] and river-influenced coastal Arctic Ocean corresponding to the latitudes of the Lena River plume [Bélanger *et al.*, 2006]. In the plume of the Mississippi River, our areal DIC photoproduction rate from tDOC (160 mmol C m⁻² yr⁻¹) is between the two earlier estimates: 73 mmol C m⁻² yr⁻¹ (0.054 Tg

C yr⁻¹ /62,068 km²); *Fichot and Benner* [2014] and 296 mmol C m⁻² yr⁻¹ [*Powers et al.*, 2015]. Our estimate for the areal DIC photoproduction from tDOC in the Laptev Sea without sea ice (52 mmol C m⁻² yr⁻¹) is similar to the corresponding estimates in the Mackenzie Shelf (76 mmol C m⁻² yr⁻¹) and Canada Basin (53 mmol C m⁻² yr⁻¹; calculated from Figure 7 in *Bélanger et al.* [2006]). Based on this comparison, our areal rates of DIC photoproduction agree well with those reported earlier.

4.8. The area of ocean required for the photochemical mineralization of tDOC to DIC

For the calculation of the area required for the photomineralization of tDOC, the annual amount of photomineralized tDOC was divided by an annual average photomineralization rate of tDOC in each river plume. The calculation thus estimates an annual average area required to photochemically mineralize the photoreactive part of tDOC flux and photobleach the entire tCDOM flux completely. After a complete photobleaching of tCDOM, a river plume becomes indistinguishable from the low-CDOM background of the open ocean.

This study estimates that the complete photomineralization of photoreactive tDOC from the ten rivers requires ocean surface area of 9.6 ± 2.5 millions km² (Table 6). Assuming that the rivers studied here are representative of the entire global tDOC flux, a simple upscaling (by a factor of 3.56) indicates that 34 (25–43) millions km² of ocean surface are required for the photomineralization of global tDOC flux. This area is less than 12% of the total ocean surface area (362 million km²; *Eakins and Sharman* [2010]), and slightly more than the surface area over the coastal shelves (24 millions km²; *Bauer et al.* [2013]), but 5–9-times larger than the total area covered by lakes (5 millions km²; *Verpoorter et*

al. [2014]). When our estimate for the photomineralization area of tDOC in the ocean is divided by the length of coastline (356 000 km excluding Antarctica without notable rivers; *Brinkhoff* [2017]), the photomineralization of tDOC is expected to take place on average within a distance of 97 ± 35 km from the coastline.

4.9. Synopsis

This study provides the first global estimate for the amount of tDOC mineralized photochemically to DIC in the ocean (45 (31 – 58) Tg C yr^{-1}) and the area (34 (25 – 43) million km^2) required for this process. Our study estimates that the amount of photomineralized tDOC is larger in the ocean than in inland waters (Figure 5). Our tDOC budget estimates that 71% of riverine tDOC is mineralized through photochemical and biological reactions within one year from its discharge to the ocean (Figure 5). Nearly half (47%) of this mineralization of tDOC is estimated to be caused by the photochemical production of labile DOC and DIC (Figure 5). In order to improve our estimates, the photomineralization areas (Table 6) could be quantitatively compared to the distribution of CDOM in the coastal ocean obtained with remote sensing methodology. In addition a better knowledge of tCDOM fluxes should put tighter constraints on the amounts of photomineralized tDOC in the ocean.

Acknowledgments. We thank all the scientists who contributed to the sampling for the Big River project: E. M. Paolucci (Paraná), D. Musibono (Congo), A. Shantz (Mekong), S. R. Khan (Ganges-Brahmaputra), Q. Huang (Yangtze), W. Schneider and B. Heim (Lena), A. Rivas and C. E. Rezende (Amazon), E. Petrescu (Danube), and H. Reader (Mississippi). We are grateful to Nikos Hatzianastassiou for the data of global radiation. We thank Patricia Medeiros for providing the unpublished a_{CDOM} data from

the Amazon River plume. David J. Kieber and Roger I. Jones are thanked for commenting on the manuscript. This work was supported by the Academy of Finland (grant no. 111761), Academy of Finland Center of Excellence programme (grant no. 307331) and by a personal grant of Maa- ja vesiteknikan Tuki ry (H. A.). Data is provided in the figures and the tables; the method and the data needed for determination of apparent quantum yield spectrum is described in an SI file.

Accepted Article

Citations in Supporting Information [*Bertilsson et al.*, 1999] [*Gao and Zepp*, 1998]
[*Jankowski et al.*, 2000] [*Lignell et al.*, 2008] [*Regnier et al.*, 2013] [*Rossel et al.*, 2013]

References

Aarnos, H., P. Ylöstalo, and A. V. Vähätalo (2012), Seasonal phototransformation of dissolved organic matter to ammonium, dissolved inorganic carbon, and labile substrates supporting bacterial biomass across the Baltic Sea, *Journal of Geophysical Research*, *117*(G1), 1–14, doi:10.1029/2010JG001633.

Amon, R. M. W. and R. Benner (1996), Photochemical and microbial consumption of dissolved organic carbon and dissolved oxygen in the Amazon River system, *Geochimica et Cosmochimica Acta*, *60*(10), 1783–1792, doi:10.1016/0016-7037(96)00055-5.

Andrews, S. S., S. Caron, and O. C. Zarioul (2000), Photochemical oxygen consumption in marine waters: a major sink for colored dissolved organic matter, *Limnology and Oceanography*, *45*(2), 267–277, doi:10.4319/lo.2000.45.2.0267.

Asmala, E., D. G. Bowers, R. Autio, H. Kaartokallio, and D. N. Thomas (2014), Qualitative changes of riverine dissolved organic matter at low salinities due to flocculation, *Journal of Geophysical Research: Biogeosciences*, *119*(10), 1919–1933, doi:10.1002/2014JG002722.

Bauer, J. E., W.-J. Cai, P. A. Raymond, T. S. Bianchi, C. S. Hopkinson, and P. A. G. Regnier (2013), The changing carbon cycle of the coastal ocean, *Nature*, *504*, 61–70, doi:10.1038/nature12857.

Bélanger, S., H. Xie, N. Krotkov, P. Larouche, W. F. Vincent, and M. Babin (2006), Photomineralization of terrigenous dissolved organic matter in Arctic coastal waters

from 1979 to 2003: Interannual variability and implications of climate change, *Global Biogeochemical Cycles*, 20(4), GB4005, doi:10.1029/2006GB002708.

Bélangier, S., S. A. Cizmeli, J. Ehn, A. Matsuoka, D. Doxaran, S. Hooker, and M. Babin (2013), Light absorption and partitioning in Arctic Ocean surface waters: Impact of multiyear ice melting, *Biogeosciences*, 10(10), 6433–6452, doi:10.5194/bg-10-6433-2013.

Benner, R., P. Louchouart, and R. M. W. Amon (2005), Terrigenous dissolved organic matter in the Arctic Ocean and its transport to surface and deep waters of the North Atlantic *Global Biogeochemical Cycles*, 19(2), 1–11, doi:10.1029/2004GB002398.

Bertilsson, S., and L. Tranvik (2000), Photochemical transformation of dissolved organic matter in lakes, *Limnology and Oceanography*, 45(4), 753–762, doi:10.4319/lo.2000.45.4.0753.

Bertilsson, S., R. Stepanauskas, R. Cuadros-Hansson, W. Graneli, J. Wikner, and L. Tranvik (1999), Photochemically induced changes in bioavailable carbon and nitrogen pools in a boreal watershed, *Aquatic Microbial Ecology*, 19, 47–56, doi:10.3354/ame019047.

Blough, N. V., and R. Del Vecchio (2002), *Biogeochemistry of Marine Dissolved Organic Matter*, 509–546, Elsevier, doi:10.1016/B978-012323841-2/50012-9.

Bricaud, A., A. Morel, and L. Prieur (1981), Absorption by dissolved organic matter of thesea (yellow substance) in the UV and visible domains, *Limnology and Oceanography*, 26(1), 43–53, doi:10.4319/lo.1981.26.1.0043.

Brinkhoff, T. (2017), *CIA World Factbook*, Coastline lengths.

Cai, W.-J. (2011), Estuarine and Coastal Ocean Carbon Paradox: CO₂ Sinks or Sites of Terrestrial Carbon Incineration?, *Annual Review of Marine Science*, 3(1), 123–145, doi:10.1146/annurev-marine-120709-142723.

- Cao, F., P. M. Medeiros, and W. L. Miller (2016), Optical characterization of dissolved organic matter in the Amazon River plume and the Adjacent Ocean: Examining the relative role of mixing, photochemistry, and microbial alterations, *Marine Chemistry*, *186*, 178–188, doi:10.1016/j.marchem.2016.09.007.
- Catalá, T. S. et al. (2015), Water mass age and aging driving chromophoric dissolved organic matter in the dark global ocean, *Global Biogeochemical Cycles*, *29*(7), 917–934, doi:10.1002/2014GB005048.
- Cauwet, G. (2002), DOM in the Coastal Zone, in *Biogeochemistry of Marine Dissolved Organic Matter*, edited by D. A. Hansell and C. A. Carlson, chap. 12, pp. 579–609, Academic Press, San Diego, doi:10.1016/B978-012323841-2/50014-2..
- Chen, J., B. Gu, E. J. LeBoeuf, H. Pan, and S. Dai (2002), Spectroscopic characterization of the structural and functional properties of natural organic matter fractions, *Chemosphere*, *48*(1), 59–68, doi:10.1016/S0045-6535(02)00041-3.
- Coles, V. J., M. T. Brooks, J. Hopkins, M. R. Stukel, P. L. Yager, and R. R. Hood (2013), The pathways and properties of the Amazon river plume in the tropical North Atlantic Ocean *Journal of Geophysical Research: Oceans*, *118*(12), 6894–6913, doi:10.1002/2013JC008981.
- Cory, R. M., C. P. Ward, B. C. Crump, and G. W. Kling (2014), Sunlight controls water column processing of carbon in arctic fresh waters, *Science*, *345*(6199), 925–928, doi:10.1126/science.1253119.
- Coynel, A., P. Seyler, H. Etcheber, M. Meybeck, and D. Orange (2005), Spatial and seasonal dynamics of total suspended sediment and organic carbon species in the Congo River, *Global Biogeochemical Cycles*, *19*(4), 1–17, doi:10.1029/2004GB002335.

Eakins, B., and G. Sharman (2010), Volumes of the World's Oceans from ETOPO1.

Fichot, C. G., and R. Benner (2014), The fate of terrigenous dissolved organic carbon in a river-influenced ocean margin, *Global Biogeochemical Cycles*, *28*, 300–318, doi:10.1002/2013GB004670.

Gao, H., and R. G. Zepp (1998), Factors Influencing Photoreactions of Dissolved Organic Matter in a Coastal River of the Southeastern United States, *Environmental Science & Technology*, *32*(19), 2940–2946, doi:10.1021/es9803660.

Groeneveld, M. M., L. J. Tranvik, and B. Koehler (2016), Photochemical mineralisation in a boreal brown water lake: considerable temporal variability and minor contribution to carbon dioxide production, *Biogeosciences*, *13*, 3931–3943, doi:10.5194/bg-13-3931-2016.

Gu, Y., A. Lensu, S. Perämäki, A. Ojala, and A. V. Vähätalo (2017), Iron and pH Regulating the Photochemical Mineralization of Dissolved Organic Carbon *ACS Omega*, *2*(5), 1905–1914, doi:10.1021/acsomega.7b00453.

Gustafsson, Ö, A. Widerlund, P. S. Andersson, J. Ingri, P. Roos, and A. Ledin (2000), Colloid dynamics and transport of major elements through a boreal river - Brackish bay mixing zone, *Marine Chemistry*, *71*(1-2), 1–21, doi:10.1016/S0304-4203(00)00035-9.

Hatzianastassiou, N., C. Matsoukas, A. Fotiadi, K. G. Pavlakis, E. Drakakis, D. Hatzidimitriou, and I. Vardavas (2005), Global distribution of Earth's surface shortwave radiation budget, *Atmospheric Chemistry and Physics Discussions*, *5*(4), 4545–4597, doi:10.5194/acpd-5-4545-2005.

Hedges, J. I., R. G. Keil, and R. Benner (1997), What happens to terrestrial organic matter in the ocean?, *Organic Geochemistry*, *27*(56), 195–212, doi:10.1016/S0146-

6380(97)00066-1.

Hernes, P. J., and R. Benner (2003), Photochemical and microbial degradation of dissolved lignin phenols: Implications for the fate of terrigenous dissolved organic matter in marine environments, *Journal of Geophysical Research*, *108*(c9), 3291, doi:10.1029/2002JC001421.

Holmes, R. M., J. W. McClelland, P. A. Raymond, B. B. Frazer, B. J. Peterson, and M. Stieglitz (2008), Lability of DOC transported by Alaskan rivers to the Arctic Ocean, *Geophysical Research Letters*, *35*(3), 3–7, doi:10.1029/2007GL032837.

Jaffé, R., Y. Ding, J. Niggemann, A. V. Vähätalo, A. Stubbins, R. G. M. Spencer, J. Campbell, and T. Dittmar (2013), Global charcoal mobilization from soils via dissolution and riverine transport to the oceans, *Science*, *340*(6130), 345–347, doi:10.1126/science.1231476.

Jankowski, J. J., D. J. Kieber, K. Mopper, and P. J. Neale (2000), Development and Intercalibration of Ultraviolet Solar Actinometers, *Photochemistry and Photobiology*, *71*(4), 431–440, doi:10.1562/0031-8655(2000)0710431DAIOUS2.0.CO2.

Johannessen, S. C., and W. L. Miller (2001), Quantum yield for the photochemical production of dissolved inorganic carbon in seawater, *Marine Chemistry*, *76*(4), 271–283, doi:10.1016/S0304-4203(01)00067-6.

Johannessen, S. C., W. Miller, and J. Cullen (2003), Calculation of UV attenuation and colored dissolved organic matter absorption spectra from measurements of ocean color, *Journal of Geophysical Research*, *108*(C9, 3301), 1–13, doi:10.1029/2000JC000514.

Johannessen, S. C., M. A. Peña, and M. L. Quenneville (2007), Photochemical production of carbon dioxide during a coastal phytoplankton bloom, *Estuarine, Coastal and Shelf*

Science, 73(1-2), 236–242, doi:10.1016/j.ecss.2007.01.006.

Jørgensen, L., C. A. Stedmon, M. A. Granskog, and M. Middelboe (2014), Tracing the long-term microbial production of recalcitrant fluorescent dissolved organic matter in seawater, *Geophysical Research Letters*, 41(7), 2481–2488, doi:10.1002/2014GL059428.

Kester, D. R., I. W. Duedall, D. N. Connors, and R. Pytkowic (1967), Preparation of artificial seawater, *Limnology and Oceanography*, 12(1), 176–179, doi:10.4319/lo.1967.12.1.0176.

Kieber, R. J., X. Zhou, and K. Mopper (1990), Formation of carbonyl compounds from UV-induced photodegradation of humic substances in natural waters: Fate of riverine carbon in the sea, *Limnology and Oceanography*, 35(7), 1503–1515, doi:10.4319/lo.1990.35.7.1503.

Kiehl, J. T., and K. E. Trenberth (1997), Earth's annual global mean energy budget, *Bulletin of the American Meteorological Society*, 78(2), 197–208, doi:10.1175/1520-0477(1997)078<0197:EAGMEB>2.0.CO;2.

Koehler, B., E. von Wachenfeldt, D. Kothawala, and L. J. Tranvik (2012), Reactivity continuum of dissolved organic carbon decomposition in lake water, *Journal of Geophysical Research*, 117, G01024, doi:10.1029/2011JG001793.

Koehler, B., T. Landelius, G. A. Weyhenmeyer, N. Machida, and L. J. Tranvik (2014), Sunlight-induced carbon dioxide emissions from inland waters, *Global Biogeochemical Cycles*, 28(7), 696–711, doi:10.1002/2014GB004850.

Koehler, B., E. Broman, and L. J. Tranvik (2016), Apparent quantum yield of photochemical dissolved organic carbon mineralization in lakes, *Limnology and Oceanography*, 61(6), 2207–2221, doi:10.1002/lno.10366.

Korshin, G. V., C.-W. Li, and M. M. Benjamin (1997), Organic Matter Through UV Spectroscopy, *Water Research*, 31(7), 1787–1795, doi:10.1016/S0043-1354(97)00006-.

Kuivikko, M., T. Kotiaho, K. Hartonen, A. Tanskanen, and A. V. Vähätalo (2007), Modeled direct photolytic decomposition of polybrominated diphenyl ethers in the Baltic Sea and the Atlantic Ocean, *Environmental Science & Technology*, 41(20), 7016–7021, doi:10.1021/es070422+.

Lalonde, K., A. V. Vähätalo, and Y. Gélinas, Revisiting the disappearance of terrestrial dissolved organic matter in the ocean: a $\delta^{13}\text{C}$ study, *Biogeosciences*, 11(13), 3707–3719, doi:10.5194/bg-11-3707-2014.

Lignell, R., L. Hoikkala, and T. Lahtinen (2008), Effects of inorganic nutrients, glucose and solar radiation on bacterial growth and exploitation of dissolved organic carbon and nitrogen in the northern Baltic Sea, *Aquatic Microbial Ecology*, 51, 209–221, doi:10.3354/ame01202.

Medeiros, P., M. Seidel, L. C. Powers, T. Dittmar, D. Hansell, and W. Miller (2015), Dissolved organic matter composition and photochemical transformations in the northern North Pacific Ocean, *Geophysical Research Letters*, 42, 863–870, doi:10.1002/2014GL062663.

Medeiros, P. M., L. Babcock-Adams, M. Seidel, R. M. Castelao, D. Di Iorio, J. T. Hollibaugh, and T. Dittmar (2017), Export of terrigenous dissolved organic matter in a broad continental shelf, *Limnology and Oceanography*, 62(4), 1718–1731, doi:10.1002/lno.10528.

Meybeck, M., Carbon, nitrogen, and phosphorus transport by world rivers, *American Journal of Science*, 282(4), 401–450, doi:10.2475/ajs.282.4.401.

Miller, W. L., and M. A. Moran (1997), Interaction of photochemical and microbial processes in the degradation of refractory dissolved organic matter from a coastal marine environment, *Limnology and Oceanography*, *42*(6), 1317–1324, doi:10.4319/lo.1997.42.6.1317.

Miller, W. L., and R. G. Zepp (1995), Photochemical production of dissolved inorganic carbon from terrestrial organic matter: Significance to the oceanic organic carbon cycle, *Geophysical Research Letters*, *22*(4), 417–420, doi:10.1029/94GL03344.

Miller, W. L., M. A. Moran, and W. M. Sheldon (2002), Determination of apparent quantum yield spectra for the formation of biologically labile photoproducts, *Limnology and Oceanography*, *47*(2), 343–352, doi:10.4319/lo.2002.47.2.0343.

Milliman, J. D., and K. L. Farnsworth (2011), River discharge to the coastal ocean: a global synthesis, vol. 24, 143–144 pp., Cambridge University Press, Cambridge, doi:10.1017/CBO9780511781247.

Minor, E. C., J. Pothén, B. J. Dalzell, H. Abdulla, and K. Mopper (2006), Effects of salinity changes on the photodegradation and ultraviolet-visible absorbance of terrestrial dissolved organic matter, *Limnology and Oceanography*, *51*(5), 2181–2186, doi:10.4319/lo.2006.51.5.2181.

Mopper, K., X. Zhou, R. J. Kieber, D. J. Kieber, R. J. Sikorski, R. D. Jones, and Gratzel (1991), Photochemical degradation of dissolved organic carbon and its impact on the oceanic carbon cycle, *Nature*, *353*(6339), 60–62, doi:10.1038/353060a0.

Nelson, N. B., and D. A. Siegel (2013), The global distribution and dynamics of chromophoric dissolved organic matter, *Annual Review of Marine Science*, (5), 447–476, doi:10.1146/annurev-marine-120710-100751.

Nelson, N. B., D. A. Siegel, and a. F. Michaels (1998), Seasonal dynamics of colored dissolved material in the Sargasso Sea, *Deep-Sea Research Part I: Oceanographic Research Papers*, 45(6), 931–957, doi:10.1016/S0967-0637(97)00106-4.

Powers, L. C., and W. L. Miller (2015), Photochemical production of CO and CO₂ in the Northern Gulf of Mexico: Estimates and challenges for quantifying the impact of photochemistry on carbon cycles, *Marine Chemistry*, 171, 21–35, doi:10.1016/j.marchem.2015.02.004.

Pullin, M. J., S. Bertilsson, J. V. Goldstone, and B. M. Voelker (2004), Effects of sunlight and hydroxyl radical on dissolved organic matter: Bacterial growth efficiency and production of carboxylic acids and other substrates, *Limnology and Oceanography*, 49(6), 2011–2022, doi:10.4319/lo.2004.49.6.2011.

Raymond, P. A., and R. G. M. Spencer (2014), Riverine DOM, *Biogeochemistry of Marine Dissolved Organic Matter: Second Edition*, pp. 509–533, doi:10.1016/B978-0-12-405940-5.00011-X.

Reader, H. E., and W. L. Miller (2012), Variability of carbon monoxide and carbon dioxide apparent quantum yield spectra in three coastal estuaries of the South Atlantic Bight, *Biogeosciences*, 9(11), 4279–4294, doi:10.5194/bg-9-4279-2012.

Regnier, P., et al. (2013), Anthropogenic perturbation of the carbon fluxes from land to ocean, *Nature Geoscience*, 6, 597–607, doi:10.1038/ngeo1830.

Riedel, T., M. Zark, A. V. Vähätalo, J. Niggemann, R. G. M. Spencer, P. J. Hernes, and T. Dittmar (2016), Molecular Signatures of Biogeochemical Transformations in Dissolved Organic Matter from Ten World Rivers, *Frontiers in Earth Science*, 4(85), 1–16, doi:10.3389/feart.2016.00085.

Rossel, P. E., A. V. Vähätalo, M. Witt, and T. Dittmar (2013), Molecular composition of dissolved organic matter from a wetland plant (*Juncus effusus*) after photochemical and microbial decomposition (1.25 yr): Common features with deep sea dissolved organic matter, *Organic Geochemistry*, *60*, 62–71, doi:10.1016/j.orggeochem.2013.04.013.

Salonen, K. (1981), Rapid and precise determination of total inorganic carbon and some gases in aqueous solutions, *Water Research*, *15*(4), 403–406, doi:10.1016/0043-1354(81)90049-X.

Salonen, K., and A. Vähätalo (1994), Photochemical mineralisation of dissolved organic matter in lake Skjervatjern, *Environment International*, *20*(3), 307–312, doi:10.1016/0160-4120(94)90114-7.

Shank, G. C., R. G. Zepp, A. Vähätalo, R. Lee, and E. Bartels (2010), Photobleaching kinetics of chromophoric dissolved organic matter derived from mangrove leaf litter and floating Sargassum colonies, *Marine Chemistry*, *119*(1-4), 162–171, doi:10.1016/j.marchem.2010.01.003.

Sholkovitz, E. R. (1976), Flocculation of dissolved organic and inorganic matter during the mixing of river water and seawater, *Geochimica et Cosmochimica Acta*, *40*(7), 831–845, doi:10.1016/0016-7037(76)90035-1.

Sobek, S., L. Tranvik, and Y. Prairie (2007), Patterns and Regulation of Dissolved Organic Carbon: An analysis of 7,500 Widely Distributed Lakes, *Limnology and Oceanography*, *52*(3), 1208–1219, doi:10.4319/lo.2007.52.3.1208.

Søndergaard, M., and M. Middelboe (1995), A cross-system analysis of labile dissolved organic carbon, *Marine Ecology Progress Series*, *118*, 283–294, doi:10.3354/meps118283.

Spencer, R. G. M., et al. (2009), Photochemical degradation of dissolved organic matter and dissolved lignin phenols from the Congo River, *Journal of Geophysical Research: Biogeosciences*, *114*(3), 1–12, doi:10.1029/2009JG000968.

Spencer, R. G. M., G. R. Aiken, M. M. Dornblaser, K. D. Butler, R. M. Holmes, G. Fiske, P. J. Mann, and A. Stubbins (2013), Chromophoric dissolved organic matter export from U.S. rivers, *Geophysical Research Letters*, *40*(8), 1575–1579, doi:10.1002/grl.50357.

Stedmon, C., R. Amon, A. Rinehart, and S. Walker (2011), The supply and characteristics of colored dissolved organic matter (CDOM) in the Arctic Ocean: Pan Arctic trends and differences, *Marine Chemistry*, *124*(1-4), 108–118, doi:10.1016/j.marchem.2010.12.007.

Swan, C. M., N. B. Nelson, D. A. Siegel, and T. S. Kostadinov (2012), The effect of surface irradiance on the absorption spectrum of chromophoric dissolved organic matter in the global ocean, *Deep Sea Research Part I: Oceanographic Research Papers*, *63*, 52–64, doi:10.1016/j.dsr.2012.01.008.

Taalba, A., H. Xie, M. G. Scarratt, S. Belanger, and M. Levasseur (2013), Photooxidation of dimethylsulde (DMS) in the Canadian Arctic, *Biogeosciences*, *10*(11), 6793–6806, doi:10.5194/bg-10-6793-2013.

Vähätalo, A., and M. Järvinen (2007), Photochemically produced bioavailable nitrogen from biologically recalcitrant dissolved organic matter stimulates production of a nitrogen-limited microbial food web in the Baltic Sea *Limnology and Oceanography*, *52*(1), 132–143, doi:10.4319/lo.2007.52.1.0132.

Vähätalo, A., M. Salkinoja-salonen, P. Taalas, and K. Salonen (2000), Spectrum of the quantum yield for photochemical mineralization of dissolved organic carbon in a humic lake, *Limnology and Oceanography*, *45*(3), 664–676, doi:10.4319/lo.2000.45.3.0664.

Vähätalo, A. V., and R. G. Wetzel (2004), Photochemical and microbial decomposition of chromophoric dissolved organic matter during long (months - years) exposures, *Marine Chemistry*, 89(1-4), 313–326, doi:10.1016/j.marchem.2004.03.010.

Vähätalo, A. V., and R. G. Zepp (2005), Photochemical Mineralization of Dissolved Organic Nitrogen to Ammonium in the Baltic Sea, *Environmental Science & Technology*, 39(18), 6985–6992, doi:10.1021/es050142z.

Vähätalo, A. V., K. Salonen, M. Salkinoja-Salonen, and A. Hatakka (1999), Photochemical mineralization of synthetic lignin in lake water indicates enhanced turnover of aromatic organic matter under solar radiation, *Biodegradation*, 10(6), 415–420, doi:10.1023/A:1008322520299.

Vähätalo, A. V., K. Salonen, U. Münster, M. Järvinen, and R. G. Wetzel (2003), Photochemical transformation of allochthonous organic matter provides bioavailable nutrients in a humic lake, *Archiv für Hydrobiologie*, 156(3), 287–314, doi:10.1127/0003-9136/2003/0156-0287.

Vähätalo, A. V., H. Aarnos, and S. Mäntyniemi (2010), Biodegradability continuum and biodegradation kinetics of natural organic matter described by the beta distribution, *Biogeochemistry*, 100(1-3), 227–240, doi:10.1007/s10533-010-9419-4.

Vähätalo, A. V., H. Aarnos, L. Hoikkala, and R. Lignell (2011), Photochemical transformation of terrestrial dissolved organic matter supports hetero- and autotrophic production in coastal waters *Marine Ecology Progress Series*, 423, 1–14, doi:10.3354/meps09010.

Vähätalo, A. V. A., and R. R. G. Wetzel (2008), Long-term photochemical and microbial decomposition of wetland-derived dissolved organic matter with alter-

ation of $^{13}\text{C}:^{12}\text{C}$ mass ratio, *Limnology and Oceanography*, *53*(4), 1387–1392, doi:10.4319/lo.2008.53.4.1387.

Verpoorter, C., T. Kutser, D. A. Seekell, and L. J. Tranvik (2014), A global inventory of lakes based on high-resolution satellite imagery, *Geophysical Research Letters*, *41*(18), 6396–6402, doi:10.1002/2014GL060641.

Vodacek, A., F. E. Hoge, R. N. Swift, J. K. Yungel, E. T. Peltzer, and N. V. Blough (1995), The use of in situ and airborne fluorescence measurements to determine UV absorption coefficients and DOC concentrations in surface waters, *Limnology and Oceanography*, *40*(2), 411–415, doi:10.4319/lo.1995.40.2.0411.

Vodacek, A., N. V. Blough, M. D. Degrandpre, E. T. Peltzer, and R. K. Nelson (1997), Seasonal variation of CDOM and DOC in the Middle Atlantic Bight: Terrestrial inputs and photooxidation, *Limnology and Oceanography*, *42*(4), 674–686, doi:10.4319/lo.1997.42.4.0674.

Wagner, S., T. Riedel, J. Niggemann, A. V. Vähätalo, T. Dittmar, and R. Jaffe (2015), Linking the Molecular Signature of Heteroatomic Dissolved Organic Matter To Watershed Characteristics in World Rivers, *Environmental Science & Technology*, *49*(23), 13798–13806, doi:10.1021/acs.est.5b00525.

Ward, N. D., R. G. Keil, P. M. Medeiros, D. C. Brito, A. C. Cunha, T. Dittmar, P. L. Yager, A. V. Krusche, and J. E. Richey (2013), Degradation of terrestrially derived macromolecules in the Amazon River, *Nature Geoscience*, *6*(7), 530–533, doi:10.1038/ngeo1817.

White, E. M., D. J. Kieber, J. Sherrard, W. L. Miller, and K. Mopper (2010), Carbon dioxide and carbon monoxide photoproduction quantum yields in the Delaware Estuary,

Marine Chemistry, 118(1-2), 11–21, doi:10.1016/j.marchem.2009.10.001.

Xiao, Y.-H., T. Sara-Aho, H. Hartikainen, and A. V. Vähätalo (2013), Contribution of ferric iron to light absorption by chromophoric dissolved organic matter, *Limnology and Oceanography*, 58(2), 653–662, doi:10.4319/lo.2013.58.2.0653.

Xiao, Y.-H., A. Räike, H. Hartikainen, and A. V. Vähätalo (2015), Iron as a source of color in river waters, *Science of The Total Environment*, 536, 914–923, doi:10.1016/j.scitotenv.2015.06.092.

Zhang, Y., H. Xie, G. Chen, S. Lawrence, and G. Chen (2006), Factors affecting the efficiency of carbon monoxide photoproduction in the St. Lawrence estuarine system (Canada), *Environmental Science & Technology*, 40(24), 7771–7777, doi:10.1021/es0615268.

Accepted Article

Table 1. Terms related to tDOC in this study.

Term	Definition
Labile	a fraction that can be mineralized by microbes with few weeks (unitless).
Non-labile	a fraction that remains after microbial mineralization of labile fraction (unitless).
Photomineralizable fraction of tDOC	the fraction of tDOC that can be mineralized by direct photochemical reactions upon complete photobleaching of tCDOM (unitless).
Photochemical reactivity of tDOC	the tendency of tDOC to undergo photochemical mineralization (unit, mol C mol absorbed photons ⁻¹).
Volumetric photomineralization rate	the amount of photomineralized tDOC per volume of water at a defined depth and unit of time (unit, mol C m ⁻³ yr ⁻¹).
Areal photomineralization rate	the amount of photomineralized tDOC per area of water surface and unit of time (unit, mol C m ⁻² yr ⁻¹).
The amount of photomineralized tDOC	the amount of photoproduct DIC from tDOC discharged by a river per unit of time (unit, g C river ⁻¹ yr ⁻¹).
Area required for mineralization of photomineralizable tDOC	the area of ocean required to complete the photomineralization of tDOC discharged by a river (unit, m ² river ⁻¹).
The distance for photomineralization of tDOC	approximates how far offshore tDOC is transported from river mouths until the photomineralization is completed (unit, m).

Accepted Article

Table 2. The coordinates and the dates for the water samples collected from the rivers and the St. Lawrence Estuary (PSU refers to the practical salinity unit) as well as the transport time between the sampling and the filtration in Helsinki.

	Sample	Latitude	Longitude	Sampling date	Transport time (days)
Rivers	Rio Negro ^a	03°07'59"S	59°54'09"W	03 Jun 10	82
	Rio Solimões ^a	03°07'58"S	59°54'04"W	03 Jun 10	82
	Congo ^b	04°18'18"S	15°28'32"E	01 May 09	25
	Congo ^b	04°20'18"S	15°11'13"E	02 May 09	24
	Danube	45°13'38"N	28°44'05"E	19 Apr 10	21
	Ganges-Brahmaputra	23°34'12"N	90°10'53"E	01 Oct 09	58
	Lena	71°54'14"N	127°15'16"E	16 Aug 09	155
	Mekong	11°33'28"N	104°56'53"E	21 Aug 09	33
	Mississippi	29°02'20"N	89°19'20"W	22 Apr 09	34
	Paraná	34°18'07"S	58°32'47"W	29 Mar 09	18
	St. Lawrence	46°54'45"N	70°52'32"W	12 Jun 09	31
	Yangtze	31°45'49"N	121°02'22"E	07 Sep 09	16
	Salinity (PSU)				
St. Lawrence	0.1	46°54'46"N	70°52'32"W	12 Jun 09	27
Estuary	5.3	47°10'49"N	70°37'16"W	12 Jun 09	27
	20	47°30'59"N	70°10'53"W	13 Jun 09	26
	22	47°55'16"N	69°46'38"W	13 Jun 09	26
	28.7	48°15'54"N	69°21'08"W	10 Jun 09	29
	27.9	48°42'06"N	68°39'07"W	10 Jun 09	29
	30.7	49°25'22"N	66°19'29"W	09 Jun 09	30

^a The river samples from Rio Negro and Rio Solimões were compiled as an Amazon River sample with the mixing ratio of 1:4, respectively.

^b The Congo River samples (twice 20-L) from two sampling locations were compiled as one Congo River sample.

Table 3. DIC photoproduction (mean \pm SD*) and the photochemical bleaching of CDOM during the experimental irradiations. Photobleaching is expressed as a change in the absorption coefficients at wavelength of 300 nm (m^{-1} ; $a_{CDOM,300,dark} - a_{CDOM,300,irradiated}$, mean \pm SD) and as relative differences (%; $[a_{CDOM,300,dark} - a_{CDOM,300,irradiated}] a_{CDOM,300,dark}^{-1}$) in the river samples and in the St. Lawrence Estuary samples.

	Sample	DIC photoproduction (mmol C m^{-3})	Photobleaching of CDOM (m^{-1})	%
Rivers	I Irradiation (39.1 h)			
	Congo	42.5 ± 0.3	11.56 ± 0.05	55 ± 0
	Mekong	3.7 ± 0.2	1.01 ± 0.13	34 ± 4
	Mississippi	16.3 ± 0.2	3.28 ± 0.37	45 ± 5
	Paraná	6.0 ± 0.3	2.03 ± 0.26	43 ± 5
	St. Lawrence	11.7 ± 0.2	3.00 ± 0.15	48 ± 2
	II Irradiation (43.3 h)			
	Amazon	29.4 ± 0.7	6.79 ± 0.60	52 ± 5
	Danube	7.2 ± 0.9	1.94 ± 0.09	45 ± 2
	Ganges-Brahmaputra	1.9 ± 0.1	0.54 ± 0.04	41 ± 3
	Lena	25.8 ± 1.2	4.99 ± 0.43	45 ± 4
Yangtze	4.4 ± 0.5	1.24 ± 0.05	41 ± 2	
St. Lawrence Estuary	III Irradiation (41.4 h)			
	0.1 PSU	24.1 ± 0.5	4.27 ± 0.16	34 ± 1
	5.3 PSU	20.3 ± 0.7	5.73 ± 0.20	49 ± 2
	20.0 PSU	9.1 ± 0.5	3.14 ± 0.04	44 ± 1
	22.0 PSU	6.9 ± 0.7	1.71 ± 0.38	33 ± 7
	28.7 PSU	4.3 ± 0.8	1.81 ± 0.49	43 ± 12
	27.9 PSU	5.9 ± 0.4	1.87 ± 0.40	36 ± 8
	30.7 PSU	2.7 ± 0.5	0.54 ± 0.09	24 ± 4

* In rivers, the error is the SD of two replicates. In the St. Lawrence Estuary, SD is calculated as the cumulative error in the replicated determinations of DIC ($n = 7$) and CDOM ($n = 2$). The data was derived with three separate 31.6, 41.4 or 43.3 hour-long irradiations.

Table 4. The water discharge, absorption coefficient at 300 nm ($a_{CDOM,300}$), the fluxes of tDOC and tCDOM, the amount of photoproducted DIC from tCDOM flux (DIC prod), and the fraction of photomineralized tDOC flux (DIC prod:tDOC flux).

River	Discharge ^a (km ³ yr ⁻¹)	$a_{CDOM,300}$ ^b (m ⁻¹)	tDOC flux ^c (Tg C yr ⁻¹)	tCDOM flux ^d (Tm ² yr ⁻¹)	DIC prod. ^e (Tg C yr ⁻¹)	DIC prod. : tDOC flux (%)
Amazon	6 300	22.9 ± 0.02	37.50	144.55 ± 0.15	6.98 ± 1.08	19 ± 3
Congo	1 300	48.4 ± 0.29	10.15	62.92 ± 0.38	3.04 ± 0.49	30 ± 5
Danube	210	7.6 ± 0.03	0.59	1.59 ± 0.01	0.08 ± 0.02	13 ± 3
Ganges-Brahm.*	1 120	2.3 ± 0.07	3.60	2.60 ± 0.08	0.13 ± 0.06	3 ± 2
Lena	520	21.0 ± 0.02	3.60	28.24 ± 0.02	1.36 ± 0.24	38 ± 7
Mekong	550	4.9 ± 0.01	0.87	2.67 ± 0.00	0.13 ± 0.04	15 ± 4
Mississippi	490	12.7 ± 0.05	3.50	5.06 ± 0.02	0.24 ± 0.05	7 ± 1
Parana	530	8.9 ± 0.07	5.90	4.71 ± 0.04	0.23 ± 0.05	4 ± 1
St. Lawrence	340	12.5 ± 0.14	1.55	1.43 ± 0.02	0.07 ± 0.01	4 ± 1
Yangtze	900	5.3 ± 0.24	1.80	4.74 ± 0.22	0.23 ± 0.07	13 ± 4
Total	12 260		69.06	258.52 ± 0.93	12.48 ± 2.10	18 ± 4 ^f

* Ganges-Brahmaputra

^a Milliman and Farnsworth [2011].

^b measured (SD = measurement uncertainty) from the river waters upon the arrival to our laboratory (Table 2), except a mean of our measurement ($24.9 \pm 0.02 \text{ m}^{-1}$) and 21.1 m^{-1} by Cao et al. 2016 for the Amazon River

^c Cauwet [2002], but updated for Amazon River by Coynel et al. [2005] and calculated for Mekong as the product of water discharge and DOC measured (1.58 mg C L^{-1}) upon the arrival of the water sample.

^d calculated by multiplying $a_{CDOM,300}$ with discharge.

^e calculated by multiplying the regression coefficient ($0.00402 \text{ mol C m}^{-2}$, Figure 1) with tCDOM flux.

^f A discharge weighted average.

Table 5. Apparent quantum yields (AQY) for DIC photoproduction (ϕ_λ) and the annual DIC photoproduction rates pr (mmol C m⁻² yr⁻¹, mean \pm SD*). The parameters c and d of AQY spectra (ϕ_λ ; Eq. 1) for DIC photoproduction, ϕ at 330 nm (ϕ_{330} ; μ mol C mol photons⁻¹). The DIC photoproduction rates in river plumes and in the St. Lawrence Estuary are calculated (Eq. 2) according to ϕ_λ and the local solar radiation (Figure S7 in Supporting Information).

Rivers	ϕ_λ				DIC photo- production rate
	c (mol C mol photons ⁻¹)	d (nm ⁻¹)	ϕ_{330} (μ mol C mol photons ⁻¹)	CV% ^a	pr (mmol C m ⁻² yr ⁻¹)
Amazon	1.4037	0.0259	272	13	136 \pm 14
Congo	1.4180	0.0258	286	1	133 \pm 1
Danube	1.3951	0.0266	218	20	71 \pm 14
Ganges-Brahmaputra	1.3730	0.0281	129	12	54 \pm 7
Lena	1.3875	0.0257	289	9	52 \pm 4 ^c
Mekong	1.3710	0.0273	165	3	76 \pm 2
Mississippi	1.4101	0.0253	335	1	157 \pm 2
Parana	1.3804	0.0271	179	14	70 \pm 7
St. Lawrence	1.3922	0.0261	250	5	75 \pm 3
Yangtze	1.3235	0.0271	172	40	54 \pm 16
St. Lawrence Estuary	c	d	ϕ_{330}	CV% ^b	pr
0.1 PSU	1.3967	0.0264	231	16	68 \pm 11
5.3 PSU	1.3731	0.0264	226	12	67 \pm 8
20.0 PSU	1.3860	0.0273	168	11	47 \pm 5
22.0 PSU	1.3752	0.0278	140	13	38 \pm 5
28.7 PSU	1.3733	0.0282	126	19	34 \pm 7
27.9 PSU	1.3557	0.0278	142	9	39 \pm 4
30.7 PSU	1.3799	0.0284	116	14	31 \pm 4

* SD of replicates (Rivers) or uncertainty of determination (Estuary)

^a CV% describes the coefficient of variability for two replicates

^b CV% calculated as in *Aarnos et al.* [2012] as a propagation of error in the DIC photoproduction, the incident photon flux density and the CDOM used for the determination of AQY.

^c assumes ice free conditions

Table 6. The area (A ; km²) and distance (r ; km) required for the photomineralization of photoreactive tDOC flux in coastal ocean (mean \pm SD*).

River	Photomineralization area ^a for tDOC (10 ³ km ²)	Photomineralization distance ^b for tDOC (km)
Amazon	4 286 \pm 1 096	1 652 \pm 422
Congo	1 899 \pm 318	1 099 \pm 184
Danube	91 \pm 38	240 \pm 100
Ganges-Brahmaputra	194 \pm 115	351 \pm 209
Lena	2 186 \pm 533	1 180 \pm 288
Mekong	141 \pm 46	299 \pm 99
Mississippi	130 \pm 26	287 \pm 58
Parana	270 \pm 89	415 \pm 136
St. Lawrence	76 \pm 19	221 \pm 54
Yangtze	352 \pm 209	474 \pm 282
Total	9 624 \pm 2 489 Median	383 \pm 160

* SD is calculated as propagation of uncertainty related to the areal DIC photoproduction rate (pr in Table 5) and to the amount of photoproduced DIC from tCDOM flux (DIC prod. in Table 4).

^a Calculated by dividing the annual amount of photoproduced DIC from tCDOM flux (Table 4) with the areal DIC photoproduction rate (pr ; Table 5).

^b $r = (2A/\pi)^{0.5}$ assuming that tCDOM flux spreads uniformly into a semicircle shaped area around the river mouth.




Figure 1. The dependence of DIC photoproduction on the tCDOM photobleaching expressed as a change in the absorption coefficient at 300 nm. The blue diamonds show the measurements from river waters mixed with artificial seawater and open diamonds show data from the salinity transect in the St. Lawrence Estuary (Table 3). The line shows a linear regression, DIC photoproduction = $4.02 \times$ photobleaching, fitted on the river water data ($R^2 = 0.96$; $n = 18$). The regression coefficient (\pm SE) $0.00402 (\pm 0.00018)$ mol C m⁻² is significantly different from zero ($p < 0.001$, $t_{\text{Stat}_{1,9}} = 22.3$). The linear regression equation without intercept was selected, because the model with intercept did not improve the coefficient of determination ($R^2 = 0.96$) and the intercept 0.63 was not significantly different from zero ($p = 0.66$, $t_{\text{Stat}_{1,8}} = 0.45$). The model fitted to all data (Rivers and Estuary) was DIC photoproduction = 4.00 photobleaching ($p < 0.001$, $t_{\text{Stat}_{1,16}} = 23.0$, $R^2 = 0.93$).




Figure 2. The apparent quantum yield spectra for DIC photoproduction (ϕ_λ) in river waters mixed with artificial seawater (Table 5; note the logarithmic scale on y-axis). The examined rivers are listed in the order of ϕ_λ from the largest to the smallest.




Figure 3. The apparent quantum yields (ϕ_{330} ; $\mu\text{mol C mol photons}^{-1}$ at 330 nm) and the annual areal DIC photoproduction rates (pr ; $\text{mmol C m}^{-2} \text{ yr}^{-1}$) along the salinity transect in the St. Lawrence Estuary. The lines refer to linear fit ($R^2 = 0.95$) between salinity and ϕ_{330} or pr . Error bars calculated as described in Table 5.




Figure 4. The annual areal DIC photoproduction rates from tDOC in river plumes (pr in Table 5). The rates are arranged order of increasing $\phi_{\lambda S}$ (as in Figure 2). Error bars describe the standard deviation of pr ($n = 2$). * = Ganges-Brahmaputra River.




Figure 5. Global fluxes of tDOC across the aquatic continuum from land to the ocean. The transportation and transformations of tDOC are shown with solid and dotted arrows, respectively. The numbers refer to the fluxes of tDOC (Tg C yr^{-1}). The best estimates and their range of uncertainty given in brackets are explained in the Supporting Information. Transport: 794 (647–950) from land to inland waters, 246 (168–800) from inland waters to the ocean; Photochemistry: 24 (15–35) DIC photoproduction in inland waters, 45 (31–58) DIC photoproduction in the ocean, 36 (9–65) photoproduction of labile tDOC in the ocean; Microbial mineralization in the ocean: 49 (41–57) labile tDOC, 44 (25–49) semilabile tDOC; Others processes: 548 (401–708) total loss in inland waters, 72 (17–140) refractory tDOC that survived photochemical and microbial mineralization during its first year in the ocean.