

**This is an electronic reprint of the original article.  
This reprint *may differ* from the original in pagination and typographic detail.**

**Author(s):** Hämäläinen, Jarmo; Landi, Nicole; Loberg, Otto; Lohvansuu, Kaisa; Pugh, Kenneth; Leppänen, Paavo H.T.

**Title:** Brain event-related potentials to phoneme contrasts and their correlation to reading skills in school-age children

**Year:** 2018

**Version:**

**Please cite the original version:**

Hämäläinen, J., Landi, N., Loberg, O., Lohvansuu, K., Pugh, K., & Leppänen, P. H. (2018). Brain event-related potentials to phoneme contrasts and their correlation to reading skills in school-age children. *International Journal of Behavioral Development*, 42(3), 357-372. <https://doi.org/10.1177/0165025417728582>

All material supplied via JYX is protected by copyright and other intellectual property rights, and duplication or sale of all or part of any of the repository collections is not permitted, except that material may be duplicated by you for your research use or educational purposes in electronic or print form. You must obtain permission for any other use. Electronic or print copies may not be offered, whether for sale or otherwise to anyone who is not an authorised user.

## **Brain event-related potentials to phoneme contrasts and their correlation to reading skills in school-age children**

Jarmo A. Hämäläinen<sup>\*,1</sup>, Nicole Landi<sup>2,3,4</sup>, Otto Loberg<sup>1</sup>, Kaisa Lohvansuu<sup>1</sup>, Kenneth Pugh<sup>2,3,5</sup>, Paavo H.T. Leppänen<sup>1</sup>

<sup>1</sup> Department of Psychology, 40014 University of Jyväskylä, Finland

<sup>2</sup> Haskins Laboratories, New Haven, Connecticut 06511

<sup>3</sup> Department of Psychology, University of Connecticut, Storrs, Connecticut 06269-1020

<sup>4</sup> Yale University Child Study Center, New Haven, Connecticut 06520

<sup>5</sup> Department of Diagnostic Radiology, Yale University School of Medicine, New Haven, Connecticut 06520-8042

\*Corresponding author

Jarmo A. Hämäläinen

Department of Psychology

P.O. Box 35

40014 University of Jyväskylä

Finland

jarmo.a.hamalainen@jyu.fi

tel. +358408053490

## **Abstract**

Development of reading skills has been shown to be tightly linked to phonological processing skills and to some extent to speech perception abilities. Although speech perception is also known to play a role in reading development, it is not clear which processes underlie this connection. Using event-related potentials (ERPs) we investigated the speech processing mechanisms for common and uncommon sound contrasts (/ba/-/da/-/ga/ and /ata/-/at:a/) with respect to the native language of school-age children in Finland and the United States. In addition, a comprehensive behavioral test battery of reading and phonological processing was administered. ERPs revealed that the children could discriminate between the speech sound contrasts (place of articulation and phoneme length) regardless of their native language. No differences emerged between the Finnish and US children in their change detection responses. However, the brain responses to the phoneme length contrast correlated robustly with reading scores in the US children with larger responses being linked to poorer reading skills. Finnish children also showed correlations between the reading and phonological measures and ERP responses, but the pattern of results was not as clear as for the US children. The results indicate that speech perception is linked to reading skills and this link is more robust for uncommon speech sound contrasts.

**Keywords:** children, cross-linguistic, dyslexia, EEG, event-related potentials, phonology, reading, speech

## Introduction

Development of reading skills has been shown to be tightly linked to facility with phonological processing skills (e.g. Anthony & Francis, 2005; Bradley & Bryant, 1978; Goswami & Bryant, 1990; Melby-Lervåg, Lyster, & Hulme, 2012; Wagner & Torgesen, 1987). Reading skills have also been previously linked to speech perception abilities (e.g., McBride-Chang, 1995; Mody et al., 1997). Here we set out to investigate the relationship between speech sound processing, using brain event-related potentials (ERPs), and reading skills in two different languages, English and Finnish.

Speech perception is known to play a role in reading development, but it is not clear which processes underlie this connection. Deficient speech processing measured with ERPs as well as discrimination and categorization tasks is related to reading failure in individuals with dyslexia (e.g., Hämäläinen et al., 2009; Leppänen et al., 2002; McBride-Chang, 1995; Richardson et al., 2003; Schulte-Körne et al., 1998), and atypical specialization to phoneme contrasts of one's own native language is related to deficient reading skills (Serniclaes et al., 2004). It is also known that speech perception abilities change during the life course as a result of spoken language exposure (Kuhl et al., 2008).

For studying speech representations in the brain, the electroencephalogram (EEG) technique offers an objective measure at any age (for reviews, see Bishop, 2007; Conboy et al., 2008; Friederici, 2005; Kuhl, 2004; Schulte-Körne & Bruder, 2010). Two event-related potentials (ERPs) derived from EEG have been shown to index auditory and speech sound discrimination accuracy: mismatch negativity (MMN) and late discriminative negativity (LDN) (Cheour et al., 2001; Näätänen et al., 2010). The amplitude of MMN to speech stimuli is also modulated by longer-term experience with speech sound representations, leading to larger responses for native contrasts as compared to non-native contrasts (Kirmse et al., 2008; Näätänen et al., 1997; Winkler et al., 1999). The attenuated MMN to non-native contrasts is likely due to top-down processes such as tuning of native speech sound categories and the consequent drop of discrimination accuracy for within-category speech contrasts (Näätänen et al., 1997; Kuhl et al., 2008).

There is less clear picture on how native and non-native speech sounds affect the LDN. Several alternatives for the functional significance of this ERP component has been put forward, for example that it would reflect preattentive cognitive evaluation of the stimuli (Ceponiene et al., 2004; Jakoby et al., 2011) or the formation of memory representations of the stimuli (Barry et al., 2009). The LDN seems to be related to the success in learning a foreign language (Jakoby et al., 2011; Shestakova et al., 2003) and the capacity to learn new words as reflected by a non-word repetition task (Barry et al., 2009). These studies indicate that LDN reflects an important processing stage in speech perception.

The strength of these ERP responses also differs between children with dyslexia and typically developing individuals (e.g., Hämäläinen et al., 2013; Lohvansuu et al., 2014; Maurer, Bucher, Brem, & Brandeis, 2003; Schulte-Körne, Deimel, Bartling, &

Remschmidt, 1998). Also other ERP responses which are generated as a consequence of afferent activation and the related processing, have been found to differentiate between individuals with reading problems and with typical reading skills (e.g., Helenius et al., 2002; Hämäläinen et al., 2013; Lohvansuu et al., 2014). For example, in children, the obligatory ERPs form a series of responses termed P1, N2 (or N250) and N4 named after their latency and order (e.g., Ceponiene et al., 2005). Moreover, infants at high familial risk for dyslexia already show abnormal brain responses to the contrasts of their native language (Leppänen et al., 1999; 2002; van Zuijen et al., 2013) as well as to speech sounds occurring rarely in the mother tongue (Guttorm et al., 2001; 2005; 2010). Furthermore, among typically developing children, correlations between neural responses to speech, reading and phonological skills have been observed (Bonte et al., 2007; Espy et al., 2004; Kuuluvainen et al., 2016; Parviainen et al., 2011).

The first aim of this study was therefore to examine how longer-term exposure to native language affects the brain responses generated by detection of deviant speech sounds embedded in a stream of repeated speech sounds in both English speaking and Finnish speaking school aged children. We hypothesized that language-group differences would emerge particularly for the phoneme length contrast. The phoneme length contrast is not semantically distinctive in English whereas it is in Finnish. Supporting our hypothesis, earlier studies have observed amplitude differences in MMN responses to phoneme length changes in Finnish and German speaking adults (e.g., Kirmse et al., 2008). For the other speech sound contrasts we used, i.e. consonant-vowel syllables with place-of-articulation changes (see the methods), the change-detection ERP responses were expected to be larger for sounds that are common than sounds that are uncommon in the Finnish language, although to a lesser degree due to English language exposure of almost all Finnish children.

The second aim of the study was to examine whether the ERP responses to speech sounds that are common or uncommon in one's native language were associated with reading skills or skills that are highly predictive of reading accuracy and speed (i.e., phonological awareness, rapid naming, verbal short-term memory) as previous studies have shown such associations for native speech sound processing (e.g., Bonte et al., 2007; Kuuluvainen et al., 2016).

## **Methods**

### *Participants*

The Finnish children were recruited at the Central Finland area via the local day-care centers, schools and a learning disability clinic. Altogether 41 children (20 girls, 21 boys) participated in the EEG study at the age of 10.3 - 12.5 years (at the 4<sup>th</sup> grade in school). They had been screened for exclusion criteria (learning disabilities other than dyslexia, neurological disorders, medication, head injuries, hearing problems). There were 4 children with a diagnosis of dyslexia and additional nine children had reading scores below -1.25 standard deviations (SD) at the 2<sup>nd</sup> grade (however, at the 4<sup>th</sup> grade

only five of them had reading scores below -1.25 SDs). These children were included in the final sample of 38 children with successful EEG data acquisition.

The children from the USA were recruited from the greater New Haven CT region via local advertisements. Altogether 76 children participated in the EEG study at the age of 4.9 - 12.2 years (kindergarten to 5th grade in school) who had been screened for exclusion criteria (neurological disorders, medication, head injuries, hearing problems). Some children were excluded based on the diagnosis of attention deficit disorder (9), minor brain dysfunction (6), and specific language impairment (3). There were 6 children with a diagnosis of dyslexia who were included in the final sample. Good EEG data were obtained for the experiment with English speech sounds for 54 children and for the experiment with Finnish speech sounds for 44 children.

### *Behavioral measures*

Finland: All behavioral assessments were conducted in June-November, while at the end of 4<sup>th</sup> grade and start of 5<sup>th</sup> grade in two testing sessions, to characterize the reading level, phonological skills and verbal working memory of the children.

*Working memory.* Series of numbers both forward and backward from the Wechsler Intelligence Scale for Children - Third edition (WISC-III) was used (Wechsler, 1991).

*Reading in Finnish.* Six reading tests were used. Standardized test of *word list reading* (Lukilasse; Häyrynen et al., 1999), number of correctly read words in 45 sec was used as the score; *non-word list reading* based on TOWRE (Torgesen et al., 1999), number of correctly read non-words in 45 sec was used as the score; *standardized test of sentence reading* where the children had to read a sentence and match it with the related correct picture (Lindeman, 1998), number of read sentences in 120 sec was used as the score; *text reading* (Puolakanaho et al., 2008), number of correctly read words in 1 minute was used as the score; *pseudoword text reading* (Eklund et al., 2015), number of correctly read words and total reading time were used as the scores; *lexical decision task* where the children silently read words and had to decide whether the word had a meaning or not, number of correct decisions was used as the score.

*Writing in Finnish.* Writing 4-syllabic words from dictation, number of correct items out of 10 was used as the score; Writing 4-syllabic pseudowords from dictation, number of correct items out of 12 was used as the score.

*Phonological processing.* *Phoneme deletion task* requiring children to delete a specified phoneme from 1-3 syllabic nonwords, number of correct items out of 18 was used as the score; *phoneme length perception* where the child heard through headphones two non-words and had to decide whether they were the same or different (Hämäläinen et al., 2009), number of correct items out of 22 was used as the score; *non-word repetition*

*task* from the Neuropsychological test battery (NEPSY; Korkman, Kirk, Kemp, 1998), number of correct items out of 16 was used as the score.

*Rapid naming.* Rapid automatized naming task (RAN: Objects Letters (Denckla & Rudel, 1976). Total matrix completion time (in seconds) was used as a measure.

The United States: All behavioral assessments were conducted in year round, and children participated in two testing sessions to characterize reading level, phonological skills and verbal working memory. Standard scores from all tests were used in the analyses.

The Comprehensive Test of Phonological Achievement (*CTOPP*) (Wagner et al., 1999), subtests of Phonological Awareness (*Elision*, *Blending Words*, and *Blending Nonwords*) were administered to determine awareness and access to the speech sound structure. Subtests of Phonological Memory (*Memory for Digits* and *Nonword Repetition*) and Rapid Naming were also administered to assess phonological encoding in working memory and speed of lexical retrieval.

The Tests of Word Reading Efficiency (*TOWRE*) (*Torgesen et al., 1999*) is comprised of two subtests requiring the speeded reading of real English words (*Sight Word Efficiency*) and of pseudowords (*Phonemic Decoding Efficiency*). For both subtests, the items are ordered from easiest to most difficult, and the examinee is asked to read as many items as possible in 45 seconds. Total score across the subtests was also calculated.

The Gray Oral Reading Test–3 (*GORT-3*) (Widerholdt, 1992) was used as an assessment of reading proficiency. Participants read aloud stories of increasing difficulty, followed by answering questions about the stories. This test measures oral reading ability in the domains of accuracy in terms of word pronunciation, fluency, and comprehension.

The Peabody Picture Vocabulary Test-III (*PPVT-III*) was used to measure lexical/vocabulary skills (Dunn & Dunn, 1997). In the PPVT, stimulus words are provided for which an individual must select the corresponding picture from a field of four. The PPVT-IV is arranged to provide words of increasing difficulty.

The WASI (Psychological Corporation, 1999) is administered as a measure of IQ. The four subtests of the WASI—Vocabulary, Block Design, Similarities, and Matrix Reasoning are used to measure various facets of intelligence, including verbal knowledge, spatial reasoning, and visual information processing. Vocabulary and Similarities subtests compose Verbal IQ. Block Design and Matrix Reasoning subtests make up the Performance IQ.

### *Stimuli and procedure*

Three passive oddball experiments were run for all children with English (/ba/, /da/, /ga/), Finnish (/ata/, intermediate /atta/ and long /atta/), and Taiwan Mandarin (/fau/ with three different lexical tones) speech stimuli (see Table 1 for details of the English

and Finnish stimuli). Place-of-articulation contrast was used in the English experiment with the consonant-vowel syllable /ba/ as the standard sound and /da/ and /ga/ syllables as the deviant sounds, and stop consonant (silent gap) length contrast was used in the Finnish experiment with the pseudoword /ata/ as the standard sound (short consonant length) and an intermediate /atta/ and long /atta/ sounds as the deviant sounds (8.3 % probability for each deviant in each experiment). The long /atta/ sound was clearly categorized as having a long consonant length, whereas the intermediate /atta/ was in the long category but close to the category border (Richardson et al., 2003). The experiments were run in the same order (English, Finnish, Chinese) for all children. The experiment with the Taiwan Mandarin stimuli was not carried out to all children due to fatigue and is therefore not reported here. In each experiment the standard sound was repeated 1000 times and two deviants were repeated each 125 times. The stimuli were presented in a pseudorandom order such that there were always at least 2 standard stimulus presentations between each deviant stimulus.

Stimuli were presented at a comfortable hearing level through a loudspeaker situated approximately 80 cm above the participant with 75 - 82 dB(C). The onset-to-onset stimulus onset asynchrony (SOA) was 1220 ms in each experiment which caused the offset-to-onset inter-stimulus interval (ISI) to vary according to the length of the stimuli.

During the experiment the participants sat in an armchair watching a muted movie and were asked not to pay attention to the presented speech stimuli. The ERP measurement lasted altogether ca. 1.5 hours. Breaks were provided when necessary.

**Table 1. Details of the speech stimuli used in the experiments.**

	English			Finnish		
	/ba/	/da/	/ga/	/ata/	intermediate /atta/	long /atta/
Stimulus type	std	dev	dev	std	dev	dev
Contrast	POA	POA	POA	gap duration (95 ms)	gap duration (155 ms)	gap duration (255 ms)
Total duration (ms)	251	251	251	298	359	440

Note. std = standard stimulus, dev = deviant stimulus, POA = place-of-articulation






Figure 1. Waveforms and spectrograms of the English (above) and Finnish (below) stimuli used in the experiments.

#### *EEG acquisition*

In both sites, the EEG data were collected with an Electric Geodesics Inc. (EGI) EEG-system and NetStation 4.2 software (<http://www.egi.com/>). Ag-AgCl electrodes with the EGI 128-channel Hydrocel sensor net were used with Cz as the reference channel during recording. The sampling rate was 500 Hz. The EEG was filtered online with a highpass filter of 0.1 and a lowpass filter of 200 Hz. Electrode impedances were pursued to be set below 50 k $\Omega$  in the beginning of the experiment. During the experiment, the quality of the data was monitored and the electrode impedances were adjusted when necessary.

#### *Data analysis*

The EEG data was analyzed using BESA Research 6.0 (BESA GmbH, Gräfelfing, Germany). Channels showing continuously bad data were interpolated using the spherical spline method (3.0 channels on average for the Finnish sample, 5.3 for the US sample). The EEG data was offline filtered using zero-phase 0.5 Hz (12dB/Oct) highpass and 30 Hz (24 dB/Oct) lowpass. ICA (Infomax on 20-120 sec time window containing at least 2 blinks) was used for correcting eye blink artifacts in the data. For averaging the epoch length was -200 - 1020 ms with 200 ms prestimulus baseline. Artifact rejection criteria were 175  $\mu$ V (maximum minus minimum amplitude) within the whole epoch and 75  $\mu$ V for fast transient amplitude changes. See Table 2 for accepted number of epochs after the artifact rejection. The groups differed in the number of epochs (US children having less epochs after artifact rejection than Finnish children; all  $ps < 0.006$ ) when tested with independent samples  $t$ -tests.

**Table 2. Average epoch numbers (SDs) included in ERP averaging after artifact rejection for the Finnish (FI) and US samples.**

	PreDa std	PreGa std	Da dev	Ga dev	Pre int /atta/ std	Pre long /atta/ std	Int /atta/ dev	Long /atta/ dev
FI	100.3 (18.8)	101.0 (19.6)	102.5 (18.3)	100.0 (19.5)	94.2 (12.8)	93.0 (14.3)	94.5 (13.9)	92.1 (14.3)
US	73.2 (22.5)	72.4 (21.9)	73.0 (21.9)	72.3 (22.9)	80.0 (20.5)	80.6 (21.3)	80.5 (20.7)	80.9 (19.8)

Note. Std = standard sound, Dev = deviant sound, Int = Intermediate. PreDa std = Standard sound preceding the deviant /da/ sound.

Due to the large number of behavioral reading measures in Finnish, Principal axis factoring (based on correlation matrix, Varimax rotation, and Bartlett factor scores) was used for the six reading variables yielding two factors: reading accuracy and reading speed explaining 49 % and 29 % of the total variance, respectively. In addition, the scores of the two writing tasks were summed, and the time (in seconds) in the two RAN tasks was averaged.

ERPs to the deviant stimuli and the pre-deviant standard stimuli were examined. When examining the responses to the pre-deviant standard stimuli only those results similar to both pre-deviant standards are reported for each experiment. The responses to the pre-deviant standard stimuli were examined separately to have comparable signal-to-noise ratio (same number of trials) to the deviant stimulus responses. To test the effect of native language exposure, the ERP responses were compared between the Finnish and US samples in BESA Statistics 2.0 using non-parametric permutation statistics and clustering (time points and electrodes) that is based on initial independent samples *t*-tests. All analyses used a channel neighbor distance of 3 cm, and time window 0 - 998 ms. The clustering (time points and electrodes) is used to control for Type I error (see Maris & Oostenveld, 2007). The number of permutations was 1000. To control for the effects of age, it was entered as a covariate into BESA Statistics 2.0 using permutation statistics based on one-way ANOVA (between groups comparison). For completeness, we report if the group effects were affected by the age covariate. To examine the effects of speech sound discrimination on reading skills the ERP data (each channel and each time point for each type of stimulus) was correlated with the behavioral measures of reading and reading related skills in BESA Statistics 2.0. The correlations were also corrected for multiple comparisons of channels and time points using permutation statistics and data clustering as implemented in BESA Statistics 2.0. Therefore, the time windows and channel clusters were data-driven and not defined a priori.

## Results

### *Cognitive skills*

Descriptive statistics of the cognitive skill measures are presented in Tables 3 and 4.

**Table 3. Descriptive statistics of the cognitive skill measures for the Finnish children (N=38).**

	Mean	Standard deviation	Range
Age (years)	11.08	0.51	10.3 - 12.5
Digit span, raw score	11.6	2.8	8 - 21
Word list reading (Lukilasse), correct items	45.9	12.0	19 - 71
Text reading, correct items	119.1	4.8	104 - 124
Non-word text reading, correct items	43.6	11.4	19-66
Sentence reading (ALLU), correct items	37.8	10.8	14-57
Lexical decision, correct items	143.8	4.5	131-150
Writing, words, correct items	5.7	2.5	0-10
Writing, non-words, correct items	8.3	2.1	3-12
Phoneme deletion, correct items	14.8	3.2	4-18
Non-word repetition, correct items	9.7	2.0	4-14
RAN, objects, time (sec)	49.6	8.8	33-72
RAN, letters, time (sec)	28.9	7.4	16-52
Phoneme length perception, correct items	16.9	2.6	11-21

**Table 4. Descriptive statistics of the cognitive skill measures for the US children.**

	Mean	Standard deviation	Range
Age (years), N=54	7.97	1.67	4.9-11.2
CTOPP, phonological awareness, N=46	107.4	15.6	85-151
CTOPP, phonological memory, N=48	96.3	9.9	79-127
CTOPP, RAN, N=21	102.6	12.6	85-124
GORT, accuracy, N=42	9.1	4.3	1-17
GORT, comprehension, N=42	11.8	3.4	5-19
GORT, fluency, N=42	10.1	4.3	1-18
TOWRE, total, N=43	107.6	20.3	66-146
TOWRE, phonemic	106.8	16.9	79-140
TOWRE, sight	105.9	18.1	59-138
PPVT, N=52	113.4	11.1	95-139
WASI, pIQ, N=43	109.6	14.2	79-138
WASI, vIQ, N=44	110.7	13.7	61-151

*ERP results*

**Differences between responses to the deviant and standard stimuli**

To ascertain that the children were able to detect the differences between the stimuli, the responses to the deviant and standard stimuli were compared. Differences were found for all conditions and in both samples, except between those to the deviant /da/ and standard /ba/ in the US sample. See below details of the stimulus differences in Tables 5 and 6, and Figures 2 and 3 for the ERP waveforms, and Figures 4 and 5 for the topographies.

**Table 5. Summary of the statistical differences between the responses to the deviant and standard stimuli in the Finnish sample.**

	Cluster n:o	p-value	Topography	Time window (ms)	ERP components
da vs. ba	1	<0.0001	P	238-400	N250/MMN
	2	<0.005	T – FC	416-738	LDN
	3	<0.009	left PO	502-750	LDN
	4	<0.042	left FC	260-386	N250
ga vs. ba	1	<0.0001	FC	274-950	MMN+LDN
	2	<0.0001	P	400-876	LDN
	3	<0.042	right T	252-440	N250
int atta vs. ata	1	<0.0001	FC	556-826	N250/LDN to second syllable
	2	<0.0001	FC	254-514	Atta4 larger N250, P1_2
	3	<0.003	PO	500-798	LDN more positive for atta4
	4	<0.004	FC	174-286	larger N250 for atta4
	5	<0.004	PO/left T	318-474	earlier N250 for ata1
	6	<0.014	PO/right T	198-280	larger N250 for atta4
long atta vs. ata	1	<0.0001	FC until 592 ms then PO	362-998	P1_2 and N250_2 for atta8
	2	<0.0001	FC	560-998	N250_2 for atta8
	3	<0.002	PO	366-542	P1_2 for atta8
	4	<0.009	FC	212-364	larger N250 for atta8
	5	<0.014	PO	174-370	larger N250 for atta8


Note. F = Frontal, C = Central, P = Parietal, T = Temporal, O = Occipital scalp areas, int = intermediate.

**Table 6. Summary of the statistical differences between the responses to the deviant and standard stimuli in the US sample.**

	Cluster n:o	p-value	Topography	Time window (ms)	ERP components
da vs. ba	ns	ns	Ns	ns	
ga vs. ba	1	<0.0001	PO – FC	80-492; 494-998	P1, N250, LDN
	2	<0.027	F	102-476	P1
	3	<0.036	left T	270-736	LDN
int atta vs. ata	1	<0.001	FC	488-876	N250 to 2nd syllable in atta
	2	<0.003	left TO	488-994	N250 to 2nd syllable in atta
	3	<0.006	FC	340-476	N250 to 2nd syllable in ata
	4	<0.020	PO	364-470	N250 to 2nd syllable in ata
	5	<0.012	FC	182-364	larger N250 for atta
long atta vs. ata	1	<0.0001	FC until 372 ms then PO	212- 546	larger N250 for atta and P1 to 2 <sup>nd</sup> syllable in atta
	2	<0.0001	FC	330-534	N250 to 2nd syllable in ata
	3	<0.002	FC	540-956	N250 to 2nd syllable in atta
	4	<0.004	PO	528-868	N250 to 2nd syllable in atta
	5	<0.038	PO	224-374	larger N250 for atta

Note. F = Frontal, C = Central, P = Parietal, T = Temporal, O = Occipital scalp areas.

US



Finnish

— Standard /ba/    — Deviant /da/    — Deviant /ga/





Figure 2. ERP waveforms in response to the English stimuli in the US sample of children (above; N=54) and in the Finnish sample of children (below; N=38). Black line is the response to the standard /ba/ stimulus, blue line the deviant /da/ stimulus, and red line the deviant /ga/ stimulus. ERP montage has been transformed to the standard 10-10 electrode positions and re-referenced to the average reference. Horizontal line marks 100 ms and vertical line marks 1  $\mu$ V, negative voltages are plotted up.

US



Finnish

— Standard /ata/    — Deviant intermediate /atta/    — Deviant long /atta/




Figure 3. ERP waveforms in response to the Finnish stimuli in the US sample of children (above; N=44) and in the Finnish sample of children (below; N=38). Black line is the response to the standard /ata/ stimulus, blue line to the deviant intermediate /atta/ stimulus, and red line the deviant long /atta/ stimulus. ERP montage has been transformed to the standard 10-10 electrode positions and re-referenced to the average reference. Horizontal line marks 100 ms and vertical line marks 1  $\mu$ V, negative voltages are plotted up.






Figure 4. Topographic maps of the voltages in the English speech sound experiment. Time points correspond to: 298 ms = MMN, 460 ms = N4, 686 ms = LDN. The contour lines represent 0.2  $\mu\text{V}$  per line, red is positive voltages, blue negative voltages.




Figure 5. Topographic maps of the voltages in the Finnish speech sound experiment. The components correspond to time points: P1 = 160 ms, first N2 = 288/274/328 ms, second N2 = 452/520/588 ms for standard ata, deviant intermediate atta, and deviant long atta, respectively. The contour lines represent 0.5  $\mu\text{V}$  per line, red is positive voltages, blue negative voltages.

### Differences between Finnish and US samples

To test if there are language specific differences in the neural responses, we compared ERP amplitudes between the groups. First, the group differences were examined for the responses to the standard and deviant sounds separately in order to see processing differences at the level of exogenous responses. There were general group differences that resulted from larger responses starting with the N250 for all stimuli in the Finnish

children compared to the US children, see Table 7 for a summary of these differences. When age was used as a covariate the significant amplitude differences between the groups remained.

Second, the group differences were examined for the difference waves (response to the deviant minus response to the standard) in order to examine processing related to change detection mechanisms that have been linked to discrimination abilities in previous literature (e.g., Näätänen et al., 2010). When examining the difference waves, the cluster-based test showed one cluster where the Finnish and the US children differed for the response to the deviant /ga/ sound. This difference was most prominent at parieto-occipital electrodes at 576 - 886 ms ( $p < 0.043$ ). The US children had larger negative going responses at the parietal electrodes than the Finnish children while Finnish children had larger positive going responses at the occipital electrodes. When age was entered as a covariate to the analysis this group difference became non-significant. Therefore, it is not discussed further.

**Table 7. Summary of the statistical differences between the ERP responses of the Finnish and the US children.**

	Cluster n:o	p-value	Topography	Time window (ms)	ERP components
<b>Ba-Da-Ga</b>					
da	1	<0.0001	TO	190-712	N250 and N4
	2	<0.002	FC	340-714	N4/LDN
	3	<0.024	FC	194-330	N250
ga	1	<0.002	PO	90-520	P1 and N4
	2	<0.017	FC	118-334	N250
	3	<0.019	FC	344-660	N4/LDN
preda std	1	<0.006	FC	178-694	N250 and N4
	2	<0.042	left T	196-676	N250 and N4
prega std	1	<0.022	FC	180-406	N250
	2	<0.015	right T	180-724	N250 and N4
diff da	1	ns			
diff ga	1	<0.043	PO	576-886	LDN
<b>Ata-Atta</b>					
int atta	1	<0.0001	FC	154-998	N250 for 1st syllable, P1, 250 for 2nd syllable

	2	<0.003	TO	184-998	N250 for 1st syllable, P1, 250 for 2nd syllable
long atta	1	<0.005	FC	386-998	P1, N250 for 2nd syllable
	2	<0.012	FC	84-366	P1, N250 for 1st syllable
	3	<0.019	TO	180-364	P1, N250 for 1st syllable
	4	<0.036	TO	638-998	N250/LDN for 2nd syllable
pre int atta std	1	<0.0001	FC	182-892	N250 for 1st syllable, P1, N250 for 2nd syllable
	2	<0.0001	TO	168-988	N250 for 1st syllable, P1, N250 for 2nd syllable
pre long atta std	1	<0.0001	FC	138-704	N250 for 1st syllable, P1, N250 for 2nd syllable
diff int atta	1	ns			
diff long atta	1	ns			

Note. F = Frontal, C = Central, P = Parietal, T = Temporal, O = Occipital scalp areas, std = standard, int = intermediate, diff = difference wave.

### Correlations

The second goal of the study was to examine if the brain responses to common and uncommon speech sound contrasts would be linked to reading abilities. Below the associations between the amplitudes of exogenous and change detection responses are correlated with the reading measures.

*Correlations to common speech sound stimuli.* In the Finnish sample the ERP amplitudes for intermediate and long /atta/s showed significant correlations with phonological measures (digit span, phoneme deletion, non-word repetition) and reading accuracy, as shown in Table 8. Larger negative amplitudes were associated with better performance in the tasks. However, the scalp areas and time windows with largest correlations varied depending on the variables. There were no significant correlations between the ERP amplitudes and cognitive skill measures in the US sample for the common speech sound stimuli (/ba/, /da/, /ga/).

*Correlations to uncommon speech sound stimuli.* In the Finnish sample the response to the deviant /ga/ sound correlated with reading speed, accuracy, and phoneme deletion scores. For reading speed the correlation indicated faster reading speed with larger response, for reading accuracy better accuracy with smaller (more negative) amplitudes, for phoneme deletion better phoneme deletion skill with larger positive amplitudes. There was also a correlation between the response to the deviant /da/ sound and reading speed indicating faster reading speed with larger response. When difference waves were examined response to the deviant /ga/ showed correlation with RAN indicating slow rapid naming with large positive voltages. Table 8 shows again that the scalp areas and time windows with maximal correlations vary depending on the behavioral measure.

**Table 8. Summary of the correlations between the ERP and cognitive skill measures in the Finnish sample.**

ERP stimuli	Cognitive measure	Topography	Time window (ms)	p-value
Ga	Reading speed	right FC	664-998	<0.004
Ga	Reading accuracy	FC	2-250	<0.026
Ga	Phoneme deletion	PO	200-370	<0.046
Da	Reading speed	PO	480-722	<0.036
Int Atta	Digit span	FC	372-816	<0.049
Diff Ga	RAN, time	TO	434-662	<0.030
Diff int Atta	Phoneme deletion	right PO	200-466	<0.019
Diff int Atta	Non-word repetition	C	308-746	<0.010
Diff long Atta	Reading accuracy	FC	514-788	p<0.020

F = Frontal, C = Central, P = Parietal, T = Temporal, O = Occipital scalp areas, Int = intermediate, Diff = Difference wave.

In contrast, in the US sample systematic and robust correlations were found between the responses to the uncommon speech sounds (/ata/, intermediate and long /atta/s) and reading skills (Table 9). Poor performance in GORT reading accuracy, fluency and comprehension as well as TOWRE reading tasks was linked to larger ERP amplitudes for all variables. The time windows where the correlations were observed encompassed all ERP components starting from P1 generated by the first syllable of the stimulus at the fronto-central channels and at the parieto-occipital channels starting from P1 generated by the second syllable of the stimulus.

The difference waves showed similar correlation pattern for GORT reading accuracy, fluency and comprehension: the larger the amplitude the poorer the reading starting from the MMN time window or even earlier. Correlations were also found between the difference wave amplitudes and TOWRE sight word and phonemic reading skills. Difference wave for the intermediate /atta/ correlated significantly with TOWRE sight word reading and had three correlation clusters: parieto-occipital channels at 280-554 ms, the more negative the voltage the better the reading score; parieto-occipital channels at 658-998 ms, the more negative the voltage the better the reading score; left fronto-central channels at 324-748 ms, the more positive the voltage the better the reading score. Difference wave for the long deviant /atta/ correlated significantly with TOWRE sight word reading score showing a variable topography for the maximal correlations depending on the latency: starting at the fronto-central channels at 464 ms changing to parieto-occipital channels at 800 ms and continuing until 998 ms, the more positive the voltage the better the reading score.

See Figures 6 and 7 for examples of the correlation coefficient topographies between the ERP measures and cognitive skill test scores.

**Table 9. Summary of the correlations between the ERP and cognitive skill measures in the US sample.**

ERP stimuli	Cognitive measure	Topography	Time window (ms)	p-value
Int Atta	GORT accuracy (N=34)	FC	156-998	<0.008
Int Atta	GORT accuracy (N=34)	PO	412-816	<0.019
Int Atta	GORT fluency (N=34)	FC	136-998	<0.014
Int Atta	GORT fluency (N=34)	PO	412-744	<0.045
Int Atta	GORT comprehension (N=34)	FC	136-998	<0.017
Int Atta	GORT comprehension (N=34)	PO	412-744	<0.047
Long Atta	GORT accuracy (N=34)	FC	120-998	<0.001
Long Atta	GORT fluency (N=34)	FC	124-998	<0.001
Long Atta	GORT comprehension (N=34)	FC	158-998	<0.001
Long Atta	GORT comprehension (N=34)	right T	408-910	<0.048
Int Atta	TOWRE phonemic (N=35)	FC	186-826	<0.013
Long Atta	TOWRE phonemic (N=35)	FC	166-872	<0.001
Long Atta	TOWRE phonemic (N=35)	right PO	158-468	<0.046
Int Atta	TOWRE sight word (N=35)	right PO	228-998	<0.001
Int Atta	TOWRE sight word (N=35)	FC	182-990	<0.002
Long Atta	TOWRE sight word (N=35)	FC	70-430	<0.005
Long Atta	TOWRE sight word (N=35)	FC	448-902	<0.010
Long Atta	TOWRE sight word (N=35)	right PO	116-604	<0.018
Long Atta	TOWRE sight word (N=35)	right T	564-996	<0.036
Diff Int Atta	GORT accuracy (N=34)	FC	242-998	<0.003
Diff Int Atta	GORT accuracy (N=34)	PO	240-662	<0.003
Diff Int Atta	GORT fluency (N=34)	FC	2-998	<0.002
Diff Int Atta	GORT fluency (N=34)	PO	240-662	<0.024
Diff Int Atta	GORT comprehension (N=34)	FC	2-998	<0.002
Diff Int Atta	GORT comprehension (N=34)	PO	40-672	<0.003
Diff Long Atta	GORT accuracy (N=34)	FC - PO	320-998	<0.005
Diff Long Atta	GORT accuracy (N=34)	F - FC - O	34-354	<0.036
Diff Long Atta	GORT fluency (N=34)	FC	314-998	<0.005
Diff Long Atta	GORT fluency (N=34)	PO	36-354	<0.036
Diff Long Atta	GORT comprehension (N=34)	FC	298-998	<0.005
Diff Long Atta	GORT comprehension (N=34)	PO	408-998	<0.003
Diff Int Atta	TOWRE sight word (N=35)	PO	280-554	<0.013
Diff Int Atta	TOWRE sight word (N=35)	PO	658-998	<0.015
Diff Int Atta	TOWRE sight word (N=35)	Left FC	324-748	<0.030
Diff Long Atta	TOWRE sight word (N=35)	FC - PO	464-998	<0.002

F = Frontal, C = Central, P = Parietal, T = Temporal, O = Occipital scalp areas, Int = intermediate, Diff = Difference wave.




Figure 6. Topographic distribution of the correlation coefficient values between the deviant /ga/ stimulus and cognitive measures in the Finnish sample (N=38). Channel clusters with statistically significant values are marked with boxes in the topography. In the topography plot, red color denotes positive coefficients and blue negative coefficients. Topography is plotted at the highest coefficient values (time point indicated in parenthesis after the cognitive measure). Left bottom: Time windows associated with the channel cluster are marked with red and blue on the ERP waveform from the channel with maximal correlation. Right bottom: correlation coefficients for each time point at the maximal channel.




Figure 7. Topographic distribution of the correlation coefficient values between the intermediate deviant /atta/ stimulus and reading test scores, for TOWRE (N=35), and for GORT (N=34) in the US sample. Channel clusters with statistically significant values (1<sup>st</sup> cluster at left column, 2<sup>nd</sup> cluster at right column) are marked with boxes. In the topography plot, red denotes positive coefficients and blue negative coefficients. Topography is plotted at the highest coefficient values (time point indicated in parenthesis after the reading test name). Left bottom: Time windows associated with the channel cluster are marked with red and blue on the ERP waveform from the channel with maximal correlation. Right bottom: correlation coefficients for each time point at the maximal channel.

## Discussion

We set out to examine whether exposure to one's native language would result in differential brain responses in school-age children, and whether these responses would be associated with reading skills as predicted by research on dyslexia and phonological processing. The Finnish and the US children differed in ERP amplitude for all stimulus types with the Finnish sample having larger responses. However, after including age as a covariate in our model, no group differences were observed for the difference waves for the Finnish or English stimuli indicating that long-term language exposure did not cause robust differences in ERP responses related to detection of speech sound changes. Critically, associations between the ERPs and reading measures in both the Finnish and the US children were found, mainly to the ERPs elicited by the speech sounds that were uncommon in the native language.

Exposure to native language was hypothesized to enhance the processing of the speech sounds common in each language. This was not found in the current study. There might be two possible explanations for the lack of clear group differences for change detection responses. First, the Finnish children had studied English at school already for two years and have most likely been exposed to English in the environment. In addition, although the /ba-/da-/ga/ contrasts do not form semantically distinctive minimal word pairs in Finnish, all of these stop consonants are part of the Finnish phonology. Second, the Finnish speech sound contrast with a stop consonant length change (as in /ata-/atta/) involves a shift in the major energy peak in the stimulus. This leads to large differences in afferent activation and obligatory responses between the stimuli and therefore makes the examination of the difference wave between the short /ata/ and the long /atta/ sounds difficult to interpret. The heavy overlap with the obligatory responses could diminish the native language effect on the discriminatory ERP responses.

We also hypothesized that the ERP responses elicited by speech processing would be associated with phonological skills and reading skills based on theories on impaired reading skills (e.g., Elbro, 1998; Elbro & Jensen, 2005; Espy et al., 2004; Mody et al., 1997). In the current study, the correlations between the ERP responses and cognitive test scores, mainly reading scores, were most robust in the US sample and for the ERP responses to the stimulus contrasts that are uncommon to English (phoneme length). The larger the responses to the uncommon speech stimulus the poorer were the reading skills. This would be in line with theories on longer-term phonological representations being important for reading acquisition and development (e.g., Vellutino et al., 2004) and with the effects of perceptual narrowing to native speech sounds during the first year of life (Tsao et al., 2004; Kuhl et al., 2008). Less efficient perceptual narrowing to the native speech sounds has been shown to be correlated with poorer later language skills (Tsao et al., 2004; Kuhl et al., 2008) and increased discrimination accuracy of non-native speech sound contrasts to be linked with reading problems (Serniclaes et al., 2004). Therefore, in the current study, it is likely that the larger ERP



responses to the speech sound contrasts uncommon to English would reflect poorly developed longer-term phonological representations in children.

The Finnish sample showed also association between the ERP measures and phonological measures for both types of speech stimuli (common and uncommon to Finnish). Interestingly the correlation between the cognitive measures and the ERP amplitudes to the phonemic length contrasts was opposite in the Finnish sample compared to the US sample. This suggests that better change detection of native language features (i.e., larger ERP amplitude) would be linked to better reading and reading-related cognitive skills whereas better change detection of non-native language features would be linked to poorer reading skills. On the other hand, in the Finnish children the direction of the association for the voiced stop consonant contrasts that are uncommon in the Finnish language was opposite to that found for the US children: the Finnish children showed better cognitive performance with larger change detection responses for /ba/-/ga/ contrast. The time window of the correlation seemed to correspond to that of MMN (phoneme deletion) and LDN (reading speed), but the topography pattern of the correlations were not typical for the MMN response. For the Finnish children, at the age of 12 years there could already be many intervening variables, for example exposure to English language via TV, music, internet, gaming and school, affecting the ERP amplitudes that could obscure the effects of the English stimuli as uncommon sounds and therefore also affect the associations with cognitive skill measures. Also, the English stimulus contrasts used are a part of the Finnish phonology, though they occur relatively rarely, and therefore might not be processed as non-native. These could have had an effect on the different correlation pattern between the US and Finnish samples.

The orthographic differences between Finnish and English could also affect the strength of the associations between the ERP measures and reading skills. The reading processes in the transparent Finnish language could rely less on phonological processes than in the opaque English language (Ziegler et al., 2010). This is reflected, for example, in previous results showing stronger association between phonological skills and reading in opaque orthographies than in transparent orthographies particularly at later school age (e.g., Georgiou et al., 2008; Ziegler et al., 2010). It is possible that the ERP measures are more closely linked to phonological abilities and therefore stronger associations between ERP measures and reading were found in the English speaking children.

Most previous studies that have found associations between ERP measures and reading or reading-related skills have examined individuals with dyslexia and/or using sounds from only one language (e.g., Hämäläinen et al., 2013; Lohvansuu et al., 2014; Schulte-Körne et al., 1998). Other studies examining typically developing children have also found associations between ERP responses to non-linguistic sounds measured in young children and reading skills at school-age (Espy et al., 2004) and ERP responses to native speech sounds, phonological skills and prereading skills in kindergarten children (Kuuluvainen et al., 2016). Also, associations between infant ERPs to non-

speech sounds and later language skills have been observed in both typically developing children and children at risk for language problems (Choudhury & Benasich, 2011). These studies suggest that the associations between ERP responses and reading related skills are more readily observable for native speech sound contrasts earlier in development than at school-age. This could be due to the ongoing changes of the phonological representations that can be larger at younger ages than in the older school-age children.

The majority of the significant correlations were found for the deviant sounds in wide time windows encompassing both the obligatory P1 and N250 responses as well as the change detection responses MMN and LDN. However, no associations were found between the ERP responses to the standard sounds and cognitive skill measures. This suggests that general level encoding, not just change detection and sound discrimination related processes, of the rarely presented speech sounds is associated with reading skills. This was somewhat unexpected because previous studies suggest that particularly the MMN and LDN responses would be sensitive to exposure to different languages (e.g., Jakoby et al., 2011; Näätänen et al., 1997; Shestakova et al., 2003; Winkler et al., 1999). However, most previous studies have not examined the associations between MMN, LDN and reading ability as continuous variables or have examined only the effect of language exposure, and not reading skills, on these ERPs. Previous studies on ERP responses in individuals with dyslexia, on the other hand, have also shown associations between obligatory N250 responses and reading or pre-reading skills in children (Hämäläinen et al., 2013; Hämäläinen et al., 2015) and N1 response and reading in adults (Helenius et al., 2002).

In order to specifically examine the contribution of MMN and LDN in the correlations, the obligatory responses should be controlled for using reversed standard and deviant probabilities or a mixture of stimuli occurring with equal probabilities to the deviant stimuli in the oddball experiment (e.g., Jacobsen & Schröger, 2001; Lohvansuu et al., 2013; Schröger & Wolff, 1996). Due to time limitations and endurance of the child participants such control experiments were not carried out in the current study.

An additional interesting finding was the larger responses of the Finnish children compared to the US children for all of the stimuli. This enhancement was particularly prominent at the N250 time window while it was not present at the P1 time window. This indicates that the cause for the larger responses is not related to technical issues in the EEG measurements but to differences in the two samples. The cause for the larger responses cannot, however, be solved based on the variables available from the current datasets.

There were two differences between the samples that could have affected the results. First, the Finnish children were older than the US children. However, age covariate did not eliminate the group differences, and therefore it is unlikely to be the primary cause for the larger ERPs in the US children. Second, despite the same analysis pipeline for both of the datasets there were more trials left in the Finnish data than in the US data. Poorer signal-to-noise ratio usually leads to larger ERP responses, but here the

Finnish children had larger responses and slightly better signal-to-noise ratio based on the trial numbers and therefore this is an unlikely explanation for the results.

Overall, our results show that processing of uncommon speech sound contrasts with respect to native language is associated with reading skills. This is in line with earlier studies on young children showing that less efficient specialization to the native language can be associated with poorer language skills (Tsao et al., 2004; Kuhl et al., 2008) and with reading problems (Serniclaes et al., 2004). Our results also support the link between speech perception, phonological skills and reading skills particularly in opaque orthographies.

### **Funding**

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This study was supported by the Academy of Finland (profiling action “Multilete” #292 466), European Union H2020MSCA-ITN-2014-ETN Programme, “Advancing brain research in children’s developmental neurocognitive disorders”—project (ChildBrain, #641652), NIH grants: P01 HD HD001994 “Nature and acquisition of the speech code and reading”, PI: C. Fowler; R01 HD 48830 Neurobiological Foundations of Reading (Dis)ability, PI: K. Pugh R03 HD053409 & R03 HD053409 “Neurocognitive development in RD children with/without general cognitive deficits”, PI: N. Landi.

### **References**

- Adams, M. J. (1994). *Beginning to read: Thinking and learning about print*. MIT press.
- Anthony, J. L., & Francis, D. J. (2005). Development of phonological awareness. *Current Directions in Psychological Science, 14*, 255-259.
- Barry, J. G., Hardiman, M. J., & Bishop, D. V. (2009). Mismatch response to polysyllabic nonwords: A neurophysiological signature of language learning capacity. *PloS one, 4*, e6270.
- Best, C. C., & McRoberts, G. W. (2003). Infant perception of non-native consonant contrasts that adults assimilate in different ways. *Language and Speech, 46*, 183-216.
- Bishop, D. V. M. (2007). Using mismatch negativity to study central auditory processing in developmental language and literacy impairments: where are we, and where should we be going?. *Psychological Bulletin, 133*, 651.
- Bonte, M. L., Poelmans, H., & Blomert, L. (2007). Deviant neurophysiological responses to phonological regularities in speech in dyslexic children. *Neuropsychologia, 45*, 1427–1437.
- Bradley, L., & Bryant, P. (1978). Difficulties in auditory organisation as a possible cause of reading backwardness. *Nature, 271*, 746–747.
- Čeponienė, R., Alku, P., Westerfield, M., Torki, M., & Townsend, J. (2005). ERPs differentiate syllable and nonphonetic sound processing in children and adults. *Psychophysiology, 42*, 391-406.

- Cheour, M., Korpilahti, P., Martynova, O., & Lang, A. H. (2001). Mismatch negativity and late discriminative negativity in investigating speech perception and learning in children and infants. *Audiology and Neurotology*, *6*, 2-11.
- Choudhury, N., & Benasich, A. A. (2011). Maturation of auditory evoked potentials from 6 to 48 months: prediction to 3 and 4 year language and cognitive abilities. *Clinical Neurophysiology*, *122*, 320-338.
- Conboy, B. T., Rivera-Gaxiola, M., Silva-Pereyra, J., & Kuhl, P. K. (2008). Event-related potential studies of early language processing at the phoneme, word, and sentence levels. *Early Language Development*, *5*, 23-64.
- Denckla, M. B., & Rudel, R. G. (1976). Rapid 'automatized' naming (RAN): Dyslexia differentiated from other learning disabilities. *Neuropsychologia*, *14*, 471-479.
- Eklund, K., Torppa, M., Aro, M., Leppänen, P. H., & Lyytinen, H. (2015). Literacy Skill Development of Children With Familial Risk for Dyslexia Through Grades 2, 3, and 8. *Journal of Educational Psychology*, *107*, 126-140.
- Elbro, C. (1998). When reading is "readn" or somthn. Distinctness of phonological representations of lexical items in normal and disabled readers. *Scandinavian Journal of Psychology*, *39*, 149-153.
- Elbro, C., & Jensen, M. N. (2005). Quality of phonological representations, verbal learning, and phoneme awareness in dyslexic and normal readers. *Scandinavian Journal of Psychology*, *46*, 375-384.
- Espy, K. A., Molfese, D. L., Molfese, V. J., & Modglin, A. (2004). Development of auditory event-related potentials in young children and relations to word-level reading abilities at age 8 years. *Annals of Dyslexia*, *54*, 9-38.
- Friederici, A. D. (2005). Neurophysiological markers of early language acquisition: from syllables to sentences. *Trends in Cognitive Sciences*, *9*, 481-488.
- Georgiou, G. K., Parrila, R., & Papadopoulos, T. C. (2008). Predictors of word decoding and reading fluency across languages varying in orthographic consistency. *Journal of Educational Psychology*, *100*(3), 566.
- Goswami, U., & Bryant, P. (1990). *Phonological skills and learning to read*. Hove: Lawrence Erlbaum.
- Guttorm, T.K., Leppänen, P.H.T., Hämäläinen, J.A., Eklund, K.M., & Lyytinen, H. (2010). Newborn event-related potentials predict poorer pre-reading skills in children at risk for dyslexia. *Journal of Learning Disabilities*, *43*, 391-401.
- Guttorm, T. K., Leppänen, P. H. T., Poikkeus, A. M., Eklund, K. M., Lyytinen, P., & Lyytinen, H. (2005). Brain event-related potentials (ERPs) measured at birth predict later language development in children with and without familial risk for dyslexia. *Cortex*, *41*, 291-303.
- Guttorm, T. K., Leppänen, P. H. T., Richardson, U., & Lyytinen, H. (2001). Event-related potentials and consonant differentiation in newborns with familial risk for dyslexia. *Journal of Learning Disabilities*, *34*, 534-544.

- Helenius, P., Salmelin, R., Richardson, U., Leinonen, S., & Lyytinen, H. (2002). Abnormal auditory cortical activation in dyslexia 100 msec after speech onset. *Journal of Cognitive Neuroscience*, *14*, 603-617.
- Hämäläinen, J. A., Leppänen, P. H. T., Eklund, K., Thomson, J., Richardson, U., Guttorm, T. K., ... & Lyytinen, H. (2009). Common variance in amplitude envelope perception tasks and their impact on phoneme duration perception and reading and spelling in Finnish children with reading disabilities. *Applied Psycholinguistics*, *30*, 511–530.
- Hämäläinen, J. A., Guttorm, T. K., Richardson, U., Alku, P., Lyytinen, H., & Leppänen, P. H. T. (2013). Auditory Event-Related Potentials Measured in Kindergarten Predict Later Reading Problems at School Age. *Developmental neuropsychology*, *38*, 550–566.
- Hämäläinen, J. A., Lohvansuu, K., Ervast, L., & Leppänen, P. H. (2015). Event-related potentials to tones show differences between children with multiple risk factors for dyslexia and control children before the onset of formal reading instruction. *International Journal of Psychophysiology*, *95*, 101-112.
- Häyrynen, T., Serenius-Sirve, S., & Korkman, M. (1999). *Lukilasse [Lukilasse-test battery for screening reading, spelling, and arithmetics]*. Helsinki: Psykologien Kustannus.
- Helenius, P., Salmelin, R., Service, E., Connolly, J.F., Leinonen, S., Lyytinen, H. (2002). Cortical activation during spoken-word segmentation in nonreading-impaired and dyslexic adults. *Journal of Neuroscience*, *22*, 2936–2944.
- Jacobsen, T., & Schröger, E. (2001). Is there pre-attentive memory-based comparison of pitch? *Psychophysiology*, *38*, 723-727.
- Jakoby, H., Goldstein, A., & Faust, M. (2011). Electrophysiological correlates of speech perception mechanisms and individual differences in second language attainment. *Psychophysiology*, *48*, 1517-1531.
- Kirmse, U., Ylinen, S., Tervaniemi, M., Vainio, M., Schröger, E., & Jacobsen, T. (2008). Modulation of the mismatch negativity (MMN) to vowel duration changes in native speakers of Finnish and German as a result of language experience. *International Journal of Psychophysiology*, *67*, 131–143.
- Korkman, M., Kirk, U., & Kemp, S. (1998). *NEPSY - lasten neuropsykologinen tutkimus*. Psykologien Kustannus Oy, Helsinki.
- Kuhl, P. K. (2004). Early language acquisition: cracking the speech code. *Nature Reviews Neuroscience*, *5*, 831–843.
- Kuhl, P. K., Tsao, F. M., & Liu, H. M. (2003). Foreign-language experience in infancy: Effects of short-term exposure and social interaction on phonetic learning. *Proceedings of the National Academy of Sciences*, *100*, 9096–9101.
- Kuhl, P. K., Conboy, B. T., Coffey-Corina, S., Padden, D., Rivera-Gaxiola, M., & Nelson, T. (2008). Phonetic learning as a pathway to language: new data and native language magnet theory expanded (NLM-e). *Philosophical Transactions of the Royal Society of London B: Biological Sciences*, *363*, 979–1000.

- Kuhl, P. K. (2010). Brain mechanisms in early language acquisition. *Neuron*, 67, 713–727.
- Kuuluvainen, S., Alku, P., Makkonen, T., Lipsanen, J., & Kujala, T. (2016). Cortical speech and non-speech discrimination in relation to cognitive measures in preschool children. *European Journal of Neuroscience*, 43, 738–750.
- Landerl, K., Ramus, F., Moll, K., Lyytinen, H., Leppänen, P. H., Lohvansuu, K., ... & Kunze, S. (2013). Predictors of developmental dyslexia in European orthographies with varying complexity. *Journal of Child Psychology and Psychiatry*, 54, 686–694.
- Leppänen, P. H., Pihko, E., Eklund, K. M., & Lyytinen, H. (1999). Cortical responses of infants with and without a genetic risk for dyslexia: II. Group effects. *Neuroreport*, 10, 969–973.
- Leppänen, P. H. T., Richardson, U., Pihko, E., Eklund, K. M., Guttorm, T. K., Aro, M., & Lyytinen, H. (2002). Brain responses to changes in speech sound durations differ between infants with and without familial risk for dyslexia. *Developmental Neuropsychology*, 22, 407–422.
- Lindeman, J. (1998). *Allu-Ala-asteen lukutesti [Reading test for primary school]*. Turku, Finland: University of Turku.
- Lohvansuu, K., Hämäläinen, J. A., Tanskanen, A., Bartling, J., Bruder, J., Honbolygó, F., ... & Leppänen, P. H. (2013). Separating mismatch negativity (MMN) response from auditory obligatory brain responses in school-aged children. *Psychophysiology*, 50, 640–652.
- Lohvansuu, K., Hämäläinen, J. A., Tanskanen, A., Ervast, L., Heikkinen, E., Lyytinen, H., & Leppänen, P. H. T. (2014). Enhancement of brain event-related potentials to speech sounds is associated with compensated reading skills in dyslexic children with familial risk for dyslexia. *International Journal of Psychophysiology*, 94, 298–310.
- Maris, E., & Oostenveld, R. (2007). Nonparametric statistical testing of EEG-and MEG-data. *Journal of neuroscience methods*, 164, 177–190.
- Maurer, U., Bucher, K., Brem, S., & Brandeis, D. (2003). Altered responses to tone and phoneme mismatch in kindergartners at familial dyslexia risk. *Neuroreport*, 14, 2245–2250.
- McBride-Chang, C. (1995). Phonological processing, speech perception and reading disability: An integrative review. *Educational Psychologist*, 30, 109–121.
- Melby-Lervåg, M., Lyster, S. A. H., & Hulme, C. (2012). Phonological skills and their role in learning to read: a meta-analytic review. *Psychological Bulletin*, 138, 322.
- Mody, M., Studdert-Kennedy, M., & Brady, S. (1997). Speech perception deficits in poor readers: auditory processing or phonological coding? *Journal of Experimental Child Psychology*, 64, 199–231.
- Näätänen, R., Lehtokoski, A., Lennest, M., Luuki, A., Alliki, J., Sinkkonen, J., & Alho, K. (1997). Language-specific phoneme representations revealed by electric and magnetic brain responses. *Nature*, 385, 432–434.

- Näätänen, R., Astikainen, P., Ruusuvirta, T., & Huotilainen, M. (2010). Automatic auditory intelligence: An expression of the sensory–cognitive core of cognitive processes. *Brain research reviews*, *64*, 123–136.
- Parviainen, T., Helenius, P., Poskiparta, E., Niemi, P., & Salmelin, R. (2011). Speech perception in the child brain: Cortical timing and its relevance to literacy acquisition. *Human Brain Mapping*, *32*, 2193–2206.
- Puolakanaho, A., Ahonen, T., Aro, M., Eklund, K., Leppänen, P. H., Poikkeus, A. M., ... & Lyytinen, H. (2008). Developmental links of very early phonological and language skills to second grade reading outcomes: Strong to accuracy but only minor to fluency. *Journal of Learning Disabilities*, *41*, 353–70.
- Richardson, U., Leppänen, P. H., Leiwo, M., & Lyytinen, H. (2003). Speech perception of infants with high familial risk for dyslexia differ at the age of 6 months. *Developmental neuropsychology*, *23*, 385–397.
- Schulte-Körne, G., & Bruder, J. (2010). Clinical neurophysiology of visual and auditory processing in dyslexia: A review. *Clinical Neurophysiology*, *121*, 1794–1809.
- Schulte-Körne, G., Deimel, W., Bartling, J., & Remschmidt, H. (1998). Auditory processing and dyslexia: evidence for a specific speech processing deficit. *Neuroreport*, *9*, 337–340.
- Schröger, E., & Wolff, C. (1996). Mismatch response of the human brain to changes in sound location. *NeuroReport*, *7*, 3005–3008.
- Serniclaes, W., Van Heghe, S., Mousty, P., Carré, R., & Sprenger-Charolles, L. (2004). Allophonic mode of speech perception in dyslexia. *Journal of Experimental Child Psychology*, *87*, 336–361.
- Seymour, P. H., Aro, M., & Erskine, J. M. (2003). Foundation literacy acquisition in European orthographies. *British Journal of Psychology*, *94*, 143–174.
- Share, D. L. (2008). On the Anglocentricities of current reading research and practice: the perils of overreliance on an "outlier" orthography. *Psychological bulletin*, *134*, 584.
- Shestakova, A., Huotilainen, M., Ceponiene, R., & Cheour, M. (2003). Event-related potentials associated with second language learning in children. *Clinical Neurophysiology*, *114*, 1507–1512.
- Schulte-Körne, G., Deimel, W., Bartling, J., & Remschmidt, H. (1998). Auditory processing and dyslexia: evidence for a specific speech processing deficit. *Neuroreport*, *9*, 337–340.
- Torgeson, J. K., Wagner, R. K., & Rashotte, C. A. (1999). *Test of Word Reading Efficiency (TOWRE)*. Austin, TX: ProEd.
- Tsao, F. M., Liu, H. M., & Kuhl, P. K. (2004). Speech perception in infancy predicts language development in the second year of life: A longitudinal study. *Child development*, *75*, 1067–1084.
- Vellutino, F. R., Fletcher, J. M., Snowling, M. J., & Scanlon, D. M. (2004). Specific reading disability (dyslexia): what have we learned in the past four decades?. *Journal of child psychology and psychiatry*, *45*, 2–40.

- Wagner, R. K., & Torgesen, J. K. (1987). The nature of phonological processing and its causal role in the acquisition of reading skills. *Psychological Bulletin*, *101*, 192–212.
- Wagner, R. K., Torgesen, J. K., & Rashotte, C. A. (1999). *Comprehensive Test of Phonological Processing: CTOPP*. Austin, TX: Pro-Ed.
- Wechsler, D. (1991). *WISC-III: Wechsler intelligence scale for children: Manual*. Psychological Corporation.
- Winkler, I., Lehtokoski, A., Alku, P., Vainio, M., Czigler, I., Csepe, V., ... & Iivonen, A. (1999). Pre-attentive detection of vowel contrasts utilizes both phonetic and auditory memory representations. *Cognitive Brain Research*, *7*, 357–369.
- Ziegler, J. C., Bertrand, D., Tóth, D., Csépe, V., Reis, A., Faísca, L., ... & Blomert, L. (2010). Orthographic depth and its impact on universal predictors of reading: A cross-language investigation. *Psychological Science*, *21*, 551-559.
- van Zuijen, T. L., Plakas, A., Maassen, B. A., Maurits, N. M., & van der Leij, A. (2013). Infant ERPs separate children at risk of dyslexia who become good readers from those who become poor readers. *Developmental science*, *16*, 554–563.