

This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Henze, Miriam J.; Lind, Olle; Mappes, Johanna; Rojas Zuluaga, Bibiana; Kelber, Almut

Title: An aposematic colour-polymorphic moth seen through the eyes of conspecifics and predators : sensitivity and colour discrimination in a tiger moth

Year: 2018

Version: Accepted version (Final draft)

Copyright: © 2018 The Authors. Functional Ecology © 2018 British Ecological Society

Rights: In Copyright

Rights url: <http://rightsstatements.org/page/InC/1.0/?language=en>

Please cite the original version:

Henze, M. J., Lind, O., Mappes, J., Rojas Zuluaga, B., & Kelber, A. (2018). An aposematic colour-polymorphic moth seen through the eyes of conspecifics and predators : sensitivity and colour discrimination in a tiger moth. *Functional Ecology*, 32(7), 1797-1809.

<https://doi.org/10.1111/1365-2435.13100>

Functional Ecology

Article type : Research Article

Corresponding author mail id: almut.kelber@biol.lu.se

Section: Animal Morphology and Colouration

Editor: Dr Christine Miller

An aposematic colour-polymorphic moth seen through the eyes of conspecifics and predators - sensitivity and colour discrimination in a tiger moth

Miriam J. Henze^{1,2}, Olle Lind³, Johanna Mappes⁴, Bibiana Rojas⁴, Almut Kelber^{1*}

¹ Lund Vision Group, Department of Biology, Lund University, Sölvegatan 35, 22362 Lund, Sweden

² present address: Queensland Brain Institute, University of Queensland, St Lucia, QLD 4072, Australia,

³ Department of Philosophy, Cognitive Science, Lund University, Helgonavägen 3, 22362 Lund, Sweden.

⁴ Centre of Excellence in Biological Interactions, University of Jyväskylä, PO Box 35, FI 40001, Finland

*corresponding author, Orcid ID: 0000-0003-3937-2808

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/1365-2435.13100

This article is protected by copyright. All rights reserved.

Abstract

1. Although predation is commonly thought to exert the strongest selective pressure on colouration in aposematic species, sexual selection may also influence colouration. Specifically, polymorphism in aposematic species cannot be explained by natural selection alone.
2. Males of the aposematic wood tiger moth (*Arctia plantaginis*) are polymorphic for hindwing colouration throughout most of their range. In Scandinavia, they display either white or yellow hindwings. Female hindwing colouration varies continuously from bright orange to red. Redder females and yellow males suffer least from bird predation.
3. White males often have higher mating success than yellow males. Therefore, we ask whether females can discriminate the two male morphs by colour. Males approach females by following pheromone plumes from a distance, but search visually at short range. This raises the questions whether males discriminate female colouration and, in turn, whether female colouration is also sexually selected.
4. Using electroretinograms, we found significantly larger retinal responses in male than female *A. plantaginis*, but similar spectral sensitivities in both sexes, with peaks in the UV (349 nm), blue (457 nm), and green (521 nm) wavelength range.
5. According to colour vision models, conspecifics can discriminate white and yellow males as separate morphs, but not orange and red females. For moths and birds (*Cyanistes caeruleus*), white males are more conspicuous against green and brown backgrounds, mostly due to UV reflectivity, and red females are slightly more conspicuous than orange females.
6. The costly red colouration among females is likely selected by predator pressure, not by conspecifics, whereas male colour polymorphism is probably maintained, at least partly, by the opposing forces of predation pressure favouring yellow males, and female preference for white males. Whether or not the preference for white males is based on visual cues requires further testing.
7. The evolution of polymorphic aposematic animals can be better understood when the visual system of the species and their predators is taken into consideration.

Resumen

1. La depredación es la principal presión de selección en la evolución de la coloración aposemática, aunque selección sexual también puede ejercer una presión importante. Los polimorfismos en la coloración de especies aposemáticas no pueden ser explicados únicamente por selección natural.
2. En Escandinavia, los machos de la polilla aposemática *Arctia plantaginis* tienen una coloración polimórfica (blanca o amarilla) en sus alas traseras; las hembras presentan una variación continua entre naranja y rojo. Machos amarillos y hembras rojas sufren menor depredación por aves.
3. Los machos blancos suelen tener un mayor éxito reproductivo que los amarillos; aunque ambos se aproximan a las hembras siguiendo sus feromonas, a cortas distancias también utilizan pistas visuales para localizarlas. Por ello, nos preguntamos (1) si las hembras pueden discriminar entre los dos tipos de machos, y los machos entre los diferentes colores de las hembras; y (2) si la coloración de las hembras podría estar afectada por selección sexual.
4. Usando electroretinogramas y modelos de visión a color, encontramos: (1) mayor respuesta en la retina de los machos que de las hembras; (2) sensibilidad espectral similar en ambos sexos, con picos en las regiones UV (349 nm), azul (457 nm), y verde (521 nm); (3) que individuos de ambos sexos pueden discriminar entre machos blancos y amarillos, pero no entre hembras rojas y anaranjadas; y (4) que para las polillas, así como para un ave (*Cyanistes caeruleus*), los machos blancos son más conspicuos sobre sustratos verdes y cafés por su reflexión de UV. Asimismo, las hembras rojas son un poco más conspicuas que las hembras anaranjadas.
5. El color rojo de las hembras está probablemente bajo una fuerte presión de depredación, no de selección sexual; la coloración polimórfica de los machos parece mantenerse mediante un balance entre la presión de depredación favoreciendo a machos amarillos, y una preferencia por machos blancos por parte de las hembras. El papel del color en dicha preferencia aún se desconoce.
6. La evolución de polimorfismos en especies aposemáticas puede entenderse mejor teniendo en cuenta los sistemas visuales tanto de la especie en cuestión como de sus depredadores.

Key words: Arctiid moths, colour polymorphism, colour vision, natural selection, predator pressure, sexual selection, spectral sensitivity

Introduction

Predation is commonly advocated as the strongest selective pressure on the evolution of conspicuous colouration in aposematic organisms (Poulton 1887; Ruxton, Sheratt & Speed 2004). However, in aposematic species with variable colouration within a given population (colour polymorphism) this principle is challenged by the difficulty for predators to learn several colour morphs (Mallet & Joron 1999; Lindström et al. 2001; Endler & Mappes 2004). Previous studies have pointed at sexual selection as an alternative or additional selective pressure influencing the maintenance of colour variation (O'Donald & Majerus 1984; Jiggins et al. 2001; Crothers & Cummings 2013). This could be particularly relevant for species displaying sexual dichromatism, colour polymorphism in one of the sexes, or both (Maan & Cummings 2009; Nokelainen et al. 2012; Rojas & Endler 2013), as natural selection may favour one colour morph while sexual selection favours another (Nokelainen et al. 2012; Crothers & Cummings 2013). Moreover, each of the different morphs might exploit receiver biases or limitations in the receivers' perceptual systems, especially while searching for mates (Limeri & Morehouse 2014).

The wood tiger moth *Arctia plantaginis* (formerly *Parasemia plantaginis*; Rönkä et al. 2016) is an aposematic, colour-polymorphic, and sexually dimorphic arctiid moth with a wide distribution across the Holarctic. Adults of *A. plantaginis* are active and mate from mid-June until the end of July in most of their range. They are diurnal-crepuscular, meaning that females release pheromone towards the evening hours, whereas males already start flying during daytime (Conner 2009). Male flight activity is highest between 18h and 22h, and female calling and mating activity peak at 20h (own observations from the field, Gordon et al. 2015; Rojas et al. 2015). At this time of the year, nights in central and northern Europe are short; at the latitude of Helsinki (60°), the sun does not set before 22h, and nocturnal light levels are never reached.

Adult wood tiger moths have a conspicuous black-and-white pattern on the forewings and red, yellow, or white hindwings. In some populations, males are monomorphic, but in large parts of Europe two male colour morphs with either white or yellow hindwings co-occur (Hegna, Galarza & Mappes 2015). In Finnish populations, frequencies of yellow and white morphs vary from 60:40 to 25:75 (Nokelainen 2013). The hindwing colour of females, however, ranges continuously from orange to red within populations, except for Scotland, where all females are yellow, and Georgia, where all females are red (Lindstedt et al. 2011; Hegna et al. 2015; B. Rojas, personal observation).

Both adult males (Nokelainen et al. 2012; Rojas et al. 2017) and females (Lindstedt et al. 2010; Brain 2016) are unpalatable to bird predators, which learn the moths' hindwing colouration as a warning signal (Lindstedt et al. 2011; Nokelainen et al. 2012; Nokelainen et al. 2014; Rönkä et al. 2018).

Although the colouration of *A. plantaginis* has mostly been studied within the context of aposematism and predator-prey interactions (i.e. natural selection), considering birds as the main signal receivers (e.g. Nokelainen et al. 2012, 2014), wing colouration may also have a function in intraspecific communication (e.g. sexual selection). Experiments suggest that white males have a higher mating success than yellow males (Nokelainen et al. 2012; Gordon et al. 2015), particularly if males experience stress. However, these experiments did not reveal whether females based their choice on colour or on other properties related to the colour morph.

While females attract males from afar by pheromones, approaching males visually search for females at short range in the vegetation (B. Rojas & J. Mappes, personal observation). Males might be choosy about which female(s) to approach, and females may or may not mate with a male they have attracted (Gordon et al. 2015). Altogether, this raises the question whether wing colouration plays a role in mate detection and choice. Addressing this question requires that we know how tiger moths perceive their own wing colours.

In the present study, we investigated the eyes of *A. plantaginis* using histology electroretinograms (ERGs). We determined the spectral sensitivity of the retina, measured the wing reflectance and used a color vision model to estimate how well the moths can detect conspecifics against natural backgrounds, and discriminate between their colour morphs. We also assessed how well a natural

predator, the blue tit *Cyanistes caeruleus*, can detect the different colour morphs of the wood tiger moth. Our results allow predictions on the opposing selective forces that act on the colour morphs and maintain colour polymorphism within populations.

Methods

Animals

We investigated animals from a stock of wood tiger moths from southern and central Finland that has been kept at the University of Jyväskylä, Finland, since 2013, with wild individuals being added every year during the field season and crosses between brothers and sisters being systematically avoided. The frequency of Finnish populations. Frequencies of yellow and white morphs is kept to approximately 50:50. Larvae were raised under greenhouse conditions and fed dandelion (*Taraxacum sp.*) leaves *ad libitum* (for more details see Lindstedt et al. 2010). Rearing containers were checked and cleaned daily until the larvae pupated. Pupae were kept at 24 °C in transparent plastic boxes lined with paper towels. After the moths had eclosed, straightened their wings and hardened, they were subjected to experiments the same day or transferred to a dark chamber and stored at 8°C for later investigation. Moths used for electrophysiology originated from the 2nd, moths for anatomy from the 9th, and those for reflectance measurements from the 11th generation of the laboratory stock.

Eye histology

Three male and three female *A. plantaginis* were decapitated, and their heads dissected and fixated in 2% glutaraldehyde, 2% formaldehyde and 2% sucrose in 0.15 M sodium cacodylate buffer for 12–24 h. After repeated rinsing in buffer, the heads were embedded in epoxy resin. Vertical sections of 3 µm thickness were taken with a Reichert Ultracut microtome using glass knives. The sections were placed on a slide, dried on a hot plate, stained with toluidine blue, and photographed under a light microscope. Rhabdom lengths were measured from micrographs using Image J150 (NIH, Bethesda,

Maryland, USA), following the course of the rhabdom from the base of the crystalline cone to the basement membrane.

Electroretinogram (ERG) recordings

For electrophysiology, a male ($n = 8$) or female ($n = 19$) adult moth was mounted on a holder and placed in a light-tight, dark Faraday cage. To explore regional differences in the retina, we inserted a tungsten recording electrode either in the ventral ($n = 9$) or dorsal ($n = 5$) half of one compound eye.

The eye was illuminated via the central fibre of a light guide, whose angular position and distance was adjusted to maximize responses to flashes of 40 ms duration separated by pauses of 5 s. After dark adaptation for 30 minutes, the spectral sensitivity was measured up to four times (see Supplementary Methods in Supporting Information) presenting narrow-band spectral flashes of equal photon flux as described previously (Telles et al. 2014; Jakobsson et al. 2017). Preceding and following each spectral series, we determined a voltage response-intensity (V -log I) relationship using flashes of white light with increasing intensity I . Response characteristics of the dark-adapted retina were investigated 3 to 20 times for each individual, with 10 or more data sets for most animals recorded at different times of the day.

To isolate the contribution of short wavelength receptors to the ERG, we repeated the recording protocol with continuous adaptation light from one of four light emitting diodes (LEDs) presented through the peripheral fibres of the light guide in the order red, amber, green, and blue (see Supplementary Methods). Each LED was switched on 5 minutes before a series of recordings started, and operated to produce increasing intensities at the position of the eye in subsequent experiments. In between sets of experiments and in the end, we monitored the state of the animal by repeatedly recording spectral control series from the dark-adapted retina.

ERG analyses

ERGs were analysed by custom-made Matlab scripts (R2013b or R2015b, The MathWorks, 160 Natick, MA, USA). The response amplitude V was calculated as the absolute value of the potential

change from the baseline at stimulus onset to maximal hyperpolarization. To assess the stability of the recordings and to document the dynamic range and saturation level of responses, a Naka-Rushton function was fitted to each V -log I dataset based on a non-linear least-squares solution.

For the dark-adapted retina, we calculated the asymptotic maximal response V_{\max} , the intensity K that elicited a half-maximal response, and the slope of the Naka-Rushton function at K . We tested the effect of *sex* and *measurement number* (repetitions) on these three parameters using linear mixed-effects models with a nested design (*measurement number* nested within *individual*). To test for time dependence, we fitted linear regression models to the data, with the effect of *sex* adjusted for the *time of day* in hours after midnight. All statistical models were implemented in R v3.3.3. (R Core Team 2017).

Based on the Naka-Rushton function fitted to the V -log I datasets obtained before and after each spectral series, we converted response amplitudes for spectral flashes into normalized sensitivities as described in Telles et al. (2014). Templates (Govardovskii et al. 2000) were used to estimate the sensitivity maxima (λ_a) of receptor types contributing to the averaged sensitivity curves ($n = 4$) by a non-linear least squares approach (see Supplementary Methods). To get the best estimate for λ_a of a specific receptor type, we selected curves with minimal contributions from other receptors.

Reflectance measurements

Spectral reflectance $s(\lambda)$ of 16 white and 10 yellow males, as well as 8 females classified as orange and 10 females classified as red were measured in the range of 300 to 700 nm, with 1 nm resolution, using an Ocean Optics USB4000 spectrometer (Dunedin, FL, USA) connected to a PX-2 pulsed Xenon lamp (for further details see Lindstedt et al. 2011; Nokelainen et al. 2012). A Spectralon™ white reflectance standard (Labsphere, Congleton, UK) was used for calibration.

Model calculations

The quantum catch Q_i of photoreceptor i ($i = \text{UV, blue, green receptor}$) is given by:

$$Q_i = k_i \int_{300}^{700} R_i(\lambda) s(\lambda) I(\lambda) d\lambda \quad \text{Eq 1,}$$

where λ is wavelength, R is receptor sensitivity, s is the reflectance spectrum of the specimen, I is the illumination spectrum, and k is a scaling factor given by adaptation to the background spectrum s_b :

$$k_i = \frac{1}{\int_{300}^{700} R_i(\lambda) s_b(\lambda) I(\lambda) d\lambda} \quad \text{Eq 2.}$$

We estimated visual contrast using a receptor-noise limited (RNL) model of colour discrimination (Vorobyev and Osorio, 1998). In this model, we suppose that colour discrimination limits are set by receptor noise that propagates into higher neuronal levels via retinal opponent mechanisms. We assume a log-linear relationship between receptor quantum catch and receptor signals, $f = \ln(Q_i)$, so that the contrast Δf between two stimuli s_1 and s_2 , is

$$\Delta f = \ln \left(\frac{Q_i s_1}{Q_i s_2} \right) \quad \text{Eq 3.}$$

Each receptor mechanism is limited by a Weber fraction ω given by receptor noise:

$$\omega_i = \frac{e_i}{\sqrt{\eta_i}} \quad \text{Eq 4,}$$

where e is the noise in each single receptor and η is the proportion of each receptor channel in the retina. As receptor noise and spatial summation are unknown for *A. plantaginis*, we set noise in the green receptor channel to 0.1, the level measured in honeybees (Vorobyev et al. 2001). Assuming that each ommatidium has 1 UV, 1 blue and 7 green receptors, as in various other moths (Warrant, Kelber, & Kristensen 2003, Briscoe 2008; Jakobsson et al. 2017, but see Belušić, Šporar, & Meglič 2017), we set receptor the proportion to 1:1:7 (UV:blue:green receptor). Given the uncertainty of noise in moth photoreceptors, we focus on relative chromatic contrast and also test whether our conclusions hold even for a flat receptor abundance ratio of 1:1:1.

Colour contrast S is given by:

$$S^2 = \frac{\omega_{UV}^2(\Delta f_{green} - \Delta f_{blue})^2 + \omega_{blue}^2(\Delta f_{green} - \Delta f_{UV})^2 + \omega_{green}^2(\Delta f_{blue} - \Delta f_{UV})^2}{(\omega_{UV}\omega_{blue})^2 + (\omega_{UV}\omega_{green})^2 + (\omega_{blue}\omega_{green})^2} \quad \text{Eq 5.}$$

The unit for colour contrast is jnds (just noticeable differences), and 1 jnd is defined as the colour contrast at detection threshold. For estimating the contrast in the visual system of the blue tit (*Cyanistes caeruleus*), we used the model for tetrachromatic bird vision (given in Supplementary Methods).

We calculated colour contrast under standard daylight illumination (d65; Wyszecki and Stiles, 2000) using two background spectra corresponding to green and brown vegetation. Our conclusions do not change when assuming the illumination in a deciduous forest (Håstad, Victorsson & Ödeen 2005), daylight d75, or light spectra typical for sunset or twilight (Johnsen et al., 2006) instead (Fig. S3).

We determine the position of colour loci (position of colours in the colour space) in a chromaticity diagram according to the following axes:

$$X_1 = A(f_{green} - f_{blue})$$

$$X_2 = B(f_{UV} - (af_{green} + bf_{blue})) \quad \text{Eq 6,}$$

where the coefficients (A, B, a, b) scale the chromaticity diagram according to the receptor-noise limited model such that the Euclidean distance of 1 between two colour loci corresponds to 1 jnd (Eq. 5):

$$A = \sqrt{\frac{1}{(\omega_{blue})^2 + (\omega_{green})^2}} \quad \text{Eq 7,}$$

$$B = \sqrt{\frac{(\omega_{blue})^2 + (\omega_{green})^2}{(\omega_{UV}\omega_{blue})^2 + (\omega_{UV}\omega_{green})^2 + (\omega_{blue}\omega_{green})^2}} \quad \text{Eq 8,}$$

$$a = \frac{(\omega_{blue})^2}{(\omega_{blue})^2 + (\omega_{green})^2} \quad \text{Eq 9,}$$

$$b = \frac{(\omega_{green})^2}{(\omega_{blue})^2 + (\omega_{green})^2} \quad \text{Eq 10.}$$

Achromatic contrast was calculated as Michelson contrast (see Supplementary Methods).

Results

Eye anatomy

We inspected vertical sections of the compound eyes and optic lobes of male and female *Arctia plantaginis* under the light microscope. The light-adapted eyes (Fig. 1) are close to spherical, have a cornea of $\approx 18 \mu\text{m}$ thickness, a crystalline cone of about twice this length, and $76 \pm 8 \mu\text{m}$ long rhabdoms in females ($n = 10$), but $100 \pm 11 \mu\text{m}$ long rhabdoms in males ($n = 10$). As no clear zone could be detected, the eyes likely function as apposition eyes. The sections of the heads also revealed that the second optic neuropil, the lamina, is spatially not directly adjacent to the retina in *A. plantaginis* (Fig. 1).

Electroretinograms (ERGs)

ERGs from the compound eyes of *A. plantaginis* consisted of monophasic hyperpolarizations (Fig. 2A), indicating that we recorded photoreceptor responses only. The rather long distance between retina and lamina (see Fig. 1) is probably the reason why our electrode did not pick up signals of downstream neurons.

We observed striking differences in response strength between males ($n = 8$) and females ($n = 6$; Fig. 2; data available from the Dryad Digital Repository: <https://doi.org/10.5061/dryad.s46t627>). In the dark-adapted retina, the estimated maximal response V_{max} to flashes of white light was more than twice as large in males (mean \pm standard error: $18.6 \pm 1.0 \text{ mV}$) as in females ($6.5 \pm 1.1 \text{ mV}$; values are based on a linear mixed-effects model accounting for repeated measurements per *individual*). This difference of 12 mV was highly significant ($t_{(12)} = 8.86, p < 0.001$, Table 1). K , the intensity that elicits

a half-maximal response ($V_{\max}/2$) and marks the turning point of the Naka-Rushton function fitted to the response-intensity (V - $\log I$) relationship, did not differ significantly between the sexes.

However, the slope of the curve at K was significantly higher for males than females (mean difference 3.1 mV / $\log I$, $t_{(12)} = 7.69$, $p < 0.001$, Table 1 and Fig. 2). Regardless of the statistical model, *sex* had a robust effect on V_{\max} and on the *Slope* at K , but not on K itself (compare Tables 1 and S1). This was independent of the number of measurements or the time of day, at least for the 16 hours between 10 h and 2 h (the next day), for which we have recordings. *Measurement number* had a significant effect on K , but the p -value was high and the size of the effect minimal (mean difference -0.02 mV / $\log I$, $t_{(130)} = -2.10$, $p = 0.04$, Table 1).

The spectral sensitivity curves obtained under dark adaptation and adaptation to different intensities of red, yellow, green, or blue light were all in agreement with the assumption of three photoreceptor types expressing visual pigments with peak sensitivities in the UV, blue, and green range (for examples, see Fig. 3; data available from the Dryad Digital Repository:

<https://doi.org/10.5061/dryad.s46t627>). Adding another spectral receptor type in the models did not considerably improve the fit. Differences in the wavelengths of maximal sensitivity λ_{α} between males and females, and dorsal and ventral eye regions were within the limits of experimental accuracy. We therefore averaged all results for each receptor type leading to a λ_{α} -value of 521 ± 2 nm (mean \pm standard deviation, $n = 14$) for the green receptor, 456 ± 7 nm ($n = 14$) for the blue receptor, and 349 ± 2 nm ($n = 13$) for the UV receptor. Note that the peak sensitivities (λ_{α}) of the three receptors are not equally spaced across the spectrum. Instead, the sensitivity peak of the blue receptor is shifted to longer wavelengths (i. e., red-shifted) by 21 nm. For the dark-adapted eye, the sensitivity peak of the green receptor type was 11 to 12 times as high as the sensitivity peak of the UV and the blue receptor type, indicating a larger number of green receptors, likely also with bigger rhabdoms.

Wing colouration

As in previous studies on the species, we classified males after inspection by the human eye as belonging to the white or the yellow morph, and females as either orange or red. While we measured very similar reflectance curves from the forewings of all morphs, hindwings differed. In particular, the hindwings of white males reflected more UV than those of any other morph (Fig. 4; data available from the Dryad Digital Repository: <https://doi.org/10.5061/dryad.s46t627>).

We used the receptor sensitivities determined by ERGs (Fig. 5A) to estimate the colour contrast of white and coloured wing patches of *A. plantaginis* in daylight illumination (Fig. 5C) with the RNL model of colour discrimination (Vorobyev & Osorio, 1998). This model provides a quantitative measure of chromatic contrast while achromatic (or luminance) contrast is ignored. Because one of the input values, the noise level in each photoreceptor channel, is unknown for *A. plantaginis*, we can only give a rough estimate of absolute discrimination thresholds. Instead, we focus on relative colour contrast, which is unaffected by this uncertainty (Table S2, Fig. S2). We do not predict colour contrasts for black wing regions, assuming that the contrasts between black and either white or coloured areas are mostly detected by the achromatic channel.

We first address the discrimination between wing colours of different morphs, and plot all colours in the colour space described by the RNL model. We assume that the moths can discriminate distinct morphs by colour, if the average contrast between the colours of different morphs exceeds the contrast between colours within each morph, such that morph colours form non-overlapping clusters.

This is not the case for the forewing colours (Table 2, Fig. 6A). Hindwing colouration is more variable (Fig 6B) differing most strongly in the stimulation of the UV-receptor, which is evident from the oblique-vertical distribution of colour loci in colour space (Fig. 6B). The average contrast between hindwing colours of white and yellow males is higher than the contrast within each of these morphs (Table 2), and - with the exception of two outliers - the hindwing colour of the white morph

forms a distinct cluster in colour space, separated from all other morphs (Fig. 6B). Thus, we assume that *A. plantaginis* likely discriminates white and yellow males as two different colour morphs, just as we do, based on human vision. The orange and red hindwing colours of females, in contrast, do not form distinct clusters, but largely overlap (Table 2, Fig 6B) and build a continuum. While conspecifics will be able to discriminate certain orange and red individuals, they will not be able to categorize them visually as belonging to two morphs.

Next, we investigated the colour contrasts of the wings against different natural backgrounds (Fig. 5C). For the moths themselves, wing colours have very similar chromatic contrasts against green and brown backgrounds, and hindwings are more conspicuous than forewings (red symbols in Fig. 7). All achromatic contrasts are higher than 0.1 and thus easily discriminable (Fig. S1).

To understand whether the red-shift of the blue receptor of *A. plantaginis* enhances its ability to detect conspecifics, we compared colour discrimination of *A. plantaginis* with that of *Macroglossum stellatarum*, a moth with equally spaced receptor sensitivities (Fig. 5A, data from Telles et al. 2014) (white symbols in Fig. 7). Assuming the same noise levels, we found no significant differences between these two visual systems, neither in the discrimination of wing colours against green or brown backgrounds (Fig. 7), nor in the discrimination between wing patches (Table S3). All results hold true even if we assume different relative numbers of receptor types (Fig. S2) or different illumination spectra (Fig. S3).

Finally, we asked whether the blue tit, a predator with tetrachromatic colour vision (for receptor sensitivities see Fig. 5B), can detect specific morphs better than others. For this bird, colour contrast of all except the white hindwings is higher against the green than the brown background, and all hindwing colours are more conspicuous than forewing colours on both backgrounds (blue symbols in Fig. 7). Achromatic contrasts are also high (Fig. S1), but birds are generally thought to use chromatic rather than achromatic contrast.

Discussion

Classical approaches have invoked predation as the most representative selective pressure favouring aposematic colouration. However, this approach falls short to explain the intra-population variation in the colouration of some aposematic species, as this variation is expected to hinder predator learning and the subsequent avoidance of aposematic prey. Sexual selection has been suggested as a pressure capable of counterbalancing predation, thus allowing multiple morphs of an aposematic species to coexist. This could be the case in the polymorphic wood tiger morph, where one morph appears to be favoured by predators while the other morph seems to be preferred by females. In the present study we investigated whether the moths can visually discriminate the colour morphs of conspecifics, a precondition for sexual selection to play a role in the maintenance of this polymorphism. Our results indicate that male and female wood tiger moths have similar spectral sensitivities but, while both sexes can discriminate the two male colour morphs, they are unable to distinguish orange from red females. Furthermore, in the eyes of both conspecifics and predators, white males are more conspicuous against green and brown backgrounds, whereas red females are slightly more conspicuous than orange females. We discuss the implications of our findings below.

Eye anatomy and sensitivity

Wood tiger moths appear to have functional apposition compound eyes (Fig. 1), as the rhabdoms of the photoreceptors are in direct contact with the crystalline cones, without the clear zone that is typical for superposition eyes (Land & Nilsson 2002). As superposition eyes are the general rule for moths (Warrant et al. 2003) including Arctiidae (D.-E. Nilsson, personal communication), this result comes as a surprise, although similar cases have been reported (Warrant et al. 2003). Males have slightly longer rhabdoms than females, which should make their photoreceptors $\approx 20\%$ more sensitive. However, this can only partly explain why the signals recorded from the dark-adapted retina were more than 100% larger in males than females (Fig. 2). Further studies are required to understand the physiological basis of the observed sex-specific differences.

Our analyses suggest that the eyes of both sexes are optimized for similar light intensities, since the linear working range of male and female eyes is centred on the same intensity K (Table 1, Fig. 2).

However, the slope of the curves is steeper in males than females, indicating that intensity differences translate into larger response differences, which might increase contrast sensitivity.

Higher (absolute and contrast) sensitivity of male eyes fits well to the natural behaviour. Females, which carry a heavy weight of developed eggs when they eclose, mostly sit on the vegetation and emit pheromones until they mate, lay eggs and die. Their flight activity is limited and not important for survival or reproduction. Males instead fly to find a mate, and have to detect them visually in dense vegetation. For flight control and for detecting females, males need sensitive eyes.

Our results show sex differences in the response strength of the photoreceptors to flashes of white light but not in their spectral tuning (Fig. 3). We have found evidence for three spectral types of photoreceptors, with sensitivity peaks for UV, blue, and green light. More than three receptor types have been described in many butterflies (Arikawa & Stavenga 2014), yet rarely in moths (Langer, Hamann & Meinecke 1979; Warrant et al. 2003; Belušić et al. 2017; Jakobsson et al. 2017). We cannot exclude that we missed additional receptor types in *A. plantaginis*, but these receptors, if present, either have a spectral sensitivity close to those characterized here or do not contribute substantially to the ERG.

While the spectral sensitivity of the UV and green receptors of *A. plantaginis* is similar to those found in sphingid moths (*Manduca sexta*: White et al. 2003; *Macroglossum stellatarum*: Telles et al. 2014), the sensitivity of the blue receptor is red-shifted (Fig. 5A). It is normally assumed that equally-spaced photoreceptor sensitivities allow for optimal discrimination of general colour stimuli (e.g. Barlow 1982). However, our models comparing *A. plantaginis* with *M. stellatarum* revealed no significant differences in discriminability of colours, and thus no obvious effect of the red-shift of the blue receptor in *A. plantaginis* (Fig. 7).

Wing colouration in the eyes of conspecifics

Seen with the eyes of *A. plantaginis*, forewings do not differ between morphs, but white and yellow males have clearly distinct hindwing colouration (Table 2, Fig. 7). Mostly because of the stronger

UV-reflection of their hindwings, white males have a higher chromatic contrast than yellow males against green and brown backgrounds. Nokelainen et al. (2012) showed the opposite for a flat white background spectrum. Thus, yellow males have a higher chromatic contrast than white males on the white bark of a birch tree, but a lower contrast on green or brown vegetation. This could have implications for sexual selection in the species (Nokelainen et al. 2012; Gordon et al. 2015). Our results also show that orange and red females cannot be discriminated as distinct colour morphs (Table 2, Fig. 7). Therefore, we suggest that female hindwing colouration is of little importance for mate selection.

It is interesting to note that the high contrast of the white hindwings against green and brown backgrounds, and to the hindwing colours of other morphs mostly results from high UV reflection. Obviously, our anthropomorphic view and naming of colour morphs does not always conform with their appearance to conspecifics, and can be misleading when searching for communication signals.

Aposematic vs intraspecific signals

To a bird, white and yellow males of *A. plantaginis* have a similar contrast against a green background (blue symbols in Fig. 7). Blue tits hesitate much longer before attacking yellow than white males presented on a green background (Nokelainen et al. 2012), but the predation on both morphs depends on the local bird community (Nokelainen et al. 2014). We suggest that colour *per se*, and perhaps the black-and-yellow pattern, and *not* the contrast against the background determines the aposematic function of the hindwings. Alternatively, the recently observed strong pyrazine odour of yellow males, which is more effective than that of white males (Rojas et al. 2017), may explain the hesitation of birds when attacking yellow moths. By contrast, the strong UV reflection of the hindwings of white males, though making them conspicuous to both females and birds, is unlikely to contribute to avoidance learning. UV is not an efficient warning signal on its own (Lyytinen et al. 2001), and can even attract birds rather than stop them from attacking (Lyytinen, Lindström & Mappes 2004; Olofsson et al. 2010). Nevertheless, our results support the role of natural selection in

the evolution of hindwing colour in *A. plantaginis* males and suggest that sexual selection may play a role, too.

Our model calculations indicate that the variation in female hindwing colouration is perceived as a continuum by conspecifics; it is thus unlikely to contribute to mate choice. Predation, however, seems to represent a selection pressure on female colouration, as birds avoid red females more often than orange females (Lindstedt et al. 2011), and learn the red colour faster (Rönkä et al. 2018). While the red pigmentation warns birds about unprofitability (Lindstedt et al. 2011), it is costly and depends on larval diet, such that larvae reared on a diet high in iridoid glycosides become adults with paler (less red) wing pigmentation (Lindstedt et al. 2010).

Forewings, on the other hand, are highly conspicuous because of the high achromatic (luminance) contrast between white and black patches (Fig. S1). Naïve birds are more strongly attracted to patterns combining yellow or red with black-and-white than with grey patches. However, they learn to avoid the former much faster (Zylinski & Osorio 2013). The combination of black and white forewings with red or yellow hindwings may thus be an efficient learning cue for birds. In fact, birds readily avoid the forewing pattern of *A. plantaginis* (Hegna & Mappes 2014).

The interplay of natural and sexual selection in the evolution of visual signals manifests in different ways across taxa. Brightly coloured models of red postman butterflies (*Heliconius erato*) attracted more males *and* deterred predators more efficiently than achromatic models and models of nonlocal morphs (Finkbeiner, Briscoe & Reed 2014). This suggests that predator deterrence (aposematism) and mate choice work in the same direction, which may explain why only one morph occurs in the studied population of this *Heliconius* species (Finkbeiner et al. 2014). In contrast, in small cabbage white butterflies (*Pieris rapae*), females prefer males that are more contrasting against the background. These males are also most conspicuous to birds, and thus in greater risk of predation (Morehouse & Rutowski 2010), a situation resembling the balance between natural and sexual selection in the well-known case of Trinidadian guppies (Endler 1983).

Opposing sexual and natural selection alone cannot explain the maintenance of local polymorphism, as this would require exactly equal strength of both selective pressures. Thus, additional selective or

genetic factors are usually involved. In *A. plantaginis*, where yellow males, on average, have an advantage against predators while white males tend to have a mating advantage, gene flow and variable selection by predators seem to offer an explanation for polymorphism (Nokelainen et al. 2014; Galarza et al. 2014). Our results on *A. plantaginis* vision confirm that male selection for female colouration is unlikely, but a female may use chromatic cues for mate choice once a male has approached her, favouring the UV-reflecting white males. Whether females use visual cues in mate selection, or whether other traits explain mating success of white males better, requires further investigation.

Acknowledgements

We are grateful to K. Suisto and the greenhouse staff at the University of Jyväskylä for rearing the animals, to O. Nokelainen and C. Lindstedt for kindly providing reflectance measurements of *A. plantaginis*, to Carina Rasmussen for excellent help with histology, to Dan-E. Nilsson for insights into lepidopteran eye types, to Thomas Labhart, Eric Warrant and Dan-E. Nilsson for access to equipment, to Thomas Labhart for advice on data analyses and to Alan Ho for diligent assistance with statistics. We thank Alan Goldizen and three anonymous reviewers for critical comments on earlier versions of the manuscript. Funding from the Finnish Centre of Excellence in Biological Interactions (2100000256) to J. Mappes and B. Rojas, and from the Swedish Research Council to A. Kelber (2012-2212) and O. Lind (637-2013-388) is gratefully acknowledged.

Author contributions

JM, BR, MJH and AK planned the study. JM and BR provided animals, wing spectra and ecological knowledge on the study species, AK studied eye histology, MJH measured and analysed ERGs, OL calculated colour vision models. All authors discussed the results, contributed to writing text and preparing figures, and approved publication. No author has any conflict of interest.

Data accessibility

All original data used in this study that are not presented in the main text or the supplements, can be found on the Dryad Digital Repository: <https://doi.org/10.5061/dryad.s46t627>.

References

- Arikawa, K., Stavenga, D.G. (2014) Insect photopigments: photoreceptor spectral sensitivities and visual adaptations. In D. M. Hunt, M. W. Hankins, S. P. Collin & N. J. Marshall (Eds), *Evolution of visual and non-visual pigments* (pp. 137–162). New York: Springer. doi:10.1007/978-1-4614-4355-1.
- Barlow, H.B. (1982) What causes trichromacy? A theoretical analysis using comb-filtered spectra. *Vision Research*, **22**, 635–643.
- Belušić, G., Šporar, K., & Meglič, A. (2017) Extreme polarisation sensitivity in the retina of the corn borer moth *Ostrinia*. *Journal of Experimental Biology*, **220**, 2047–2056.
- Brain, M.A. (2016) Can warning signals be honest? Wing colouration and the strength of chemical defence in the female wood tiger moth. MSc Thesis, University of Jyväskylä, Finland.
- Briscoe, A.D. (2008) Reconstructing the ancestral butterfly eye: focus on the opsins. *Journal of Experimental Biology*, **211**, 1805–1813.
- Conner, W.E. (Ed.) (2009) Tiger Moths and Wolly Bears. Behaviour, Ecology and Evolution of the Arctiidae. Oxford: Oxford University Press.
- Crothers L.R. & Cummings, M.E. (2013) Warning signal brightness variation: sexual selection may work under the radar of natural selection in populations of a polytypic poison frog. *American Naturalist* **181**, E116-E124.
- Endler, J.A. (1983) Natural and sexual selection on color patterns in poeciliid fishes. *Environmental Biology of Fishes*, **9**, 173–190.
- Endler, J.A. & Mappes, J. (2004) Predator mixes and the conspicuousness of aposematic signals. *American Naturalist*, **163**, 532–547.
- Finkbeiner, S.D., Briscoe, A.D. & Reed, R.D. (2014) Warning signals are seductive: relative contributions of color and pattern to predator avoidance and mate attraction in *Heliconius* butterflies. *Evolution*, **68**, 3410–3420.

- Galarza, J.A., Nokelainen, O., Ashrafi, R., Hegna, R.H. & Mappes, J. (2014) Temporal relationship between genetic and warning signal variation in the aposematic wood tiger moth (*Parasemia plantaginis*). *Molecular Ecology*, **23**, 4939–4957.
- Gordon, S.P., Kokko, H., Rojas, B., Nokelainen, O. & Mappes, J. (2015) Colour polymorphism torn apart by opposing positive frequency-dependent selection, yet maintained in space. *Journal of Animal Ecology*, **84**, 1555–1564
- Govardovskii, V.I., Fyhrquist, N., Reuter, T., Kuzmin, D.G. & Donner, K. (2000) In search of the visual pigment template. *Visual Neuroscience*, **17**, 509–528
- Håstad, O., Victorsson, J. & Ödeen, A. (2005) Differences in color vision make passerines less conspicuous in the eyes of their predators. *Proceedings of the National Academy of Sciences of the United States of America*, **102**, 6391–6394.
- Hegna, J.R. & Mappes, J. (2014) Influences of geographic differentiation in the forewing warning signal of the wood tiger moth in Alaska. *Evolutionary Ecology*, **28**, 1003–1017.
- Hegna, R.H., Galarza, J. A. & Mappes, J. (2015) Global phylogeography and geographical variation in warning coloration of the wood tiger moth (*Parasemia plantaginis*). *Journal of Biogeography*, **42**, 1469–1481.
- Henze, M. J., Lind, O., Mappes, J., Rojas, B. & Kelber, A. (2018) Data from: An aposematic colour-polymorphic moth seen through the eyes of conspecifics and predators - sensitivity and colour discrimination in a tiger moth. Dryad Digital Repository. doi:10.5061/dryad.s46t627
- Jakobsson, J., Henze, M. J., Svensson, G.P., Lind, O., Anderbrant, O. (2017) Visual cues of oviposition sites and spectral sensitivity of *Cydia strobilella* L. *Journal of Insect Physiology*, **101**, 161–168.
- Jiggins, C.D., Naisbit, R.E., Coe, R.L., Mallet, J. (2001) Reproductive isolation caused by colour pattern mimicry. *Nature* **411**, 302–305.
- Land, M.F. & Nilsson, D.-E. (2002) Animal eyes. Oxford: Oxford University Press.
- Langer, H. & Hamann, B. & Meinecke C.-C. (1979) Tetrachromatic visual system in the moth *Spodoptera exempta* (Insecta: Noctuidae). *Journal of Comparative Physiology*, **129**, 235–239.

- Lindstedt, C., Reudler Talsma J. H., Ihalainen, E., Lindström, L., Mappes, J. (2010) Diet quality affects warning coloration indirectly: excretion costs in a generalist herbivore. *Evolution* 64:68-78.
- Lindstedt, C., Eager, H., Ihalainen, E., Kahilainen, A., Stevens, M. & Mappes, J. (2011) Direction and strength of selection by predators for the color of the aposematic wood tiger moth. *Behavioral Ecology*, **22**, 580–587.
- Limeri, L.B. & Morehouse, N.H. (2014) Sensory limitations and the maintenance of colour polymorphisms: viewing the 'alba' female polymorphism through the visual system of male *Colias* butterflies. *Functional Ecology*, **28**, 1197–1207.
- Lindström, L., Alatalo, R.V., Lyytinen, A. & Mappes, J. (2001) Strong antiapostatic selection against novel rare aposematic prey. *Proceedings of the National Academy of Sciences of the United States of America*, **98**, 9181–9184.
- Lyytinen, A., Alatalo, R.V., Lindström, L. & Mappes, J. (2001) Can ultraviolet cues function as aposematic signals? *Behavioral Ecology*, **12**, 65–70.
- Lyytinen, A., Lindström, L. & Mappes, J. (2004) Ultraviolet reflection and predation risk in diurnal and nocturnal Lepidoptera. *Behavioral Ecology*, **15**, 982–987.
- Maan, M.E. & Cummings, M.E. (2009) Sexual dimorphism and directional sexual selection on aposematic signals in a poison frog. *Proceedings of the National Academy of Sciences of the United States of America* 106, 19072–19077.
- Mallet, J. & Joron, M. (1999) Evolution of diversity in warning color and mimicry: polymorphisms, shifting balance, and speciation. *Annual Review of Ecology and Systematics*, **30**, 201–233.
- Morehouse, N.I. & Rutowski, R.L. (2010) In the eye of the beholders: female choice and avian predation risk associated with an exaggerated male butterfly color. *American Naturalist*, **176**, 768–784.
- Nokelainen, O. (2013) Many forms of the wood tiger moth (*Parasemia plantaginis*): selective heterogeneity favours polymorphic warning signals. PhD Thesis, University of Jyväskylä.
- Nokelainen, O., Hegna, R.H., Reudler, J.H., Lindstedt, C. & Mappes, J. (2012) Trade-off between warning signal efficacy and mating success in the wood tiger moth. *Proceedings of the Royal Society B-Biological Sciences*, **279**, 257–265.
- Nokelainen, O., Valkonen, J., Lindstedt, C. & Mappes, J. (2014) Changes in predator community structure shifts the efficacy of two warning signals in Arctiid moths. *Journal of Animal Ecology*, **83**, 598–605.

- O'Donald, P., Majerus, M.E.N. (1984) Polymorphism of melanic ladybirds maintained by frequency-dependent sexual selection. *Biological Journal of the Linnean Society* **23**, 101–111.
- Olofsson, M., Vallin, A., Jakobsson, S. & Wiklund, C. (2010) Marginal eyespots on butterfly wings deflect bird attacks under low light intensities with UV wavelengths. *PLoS ONE* **5**(5): e10798.
- Osorio, D., Ham, A.D., Gonda, Z. & Andrew, R.J. (2009) Sensory generalization and learning about novel colours by poultry chicks. *Quarterly Journal of Experimental Psychology (Hove)*, **62**, 1249–1256.
- Poulton, E.B. (1887) The experimental proof of the protective value of colour and markings in insects in reference to their vertebrate enemies. *Proceedings of the Zoological Society of London*, **55**, 191–274.
- R Core Team (2017) R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <http://www.R-project.org/>.
- Rojas, B., Burdfield-Steel, E., Pakkanen, H., Suisto, K., Maczka, M., Schulz, S., & Mappes, J. (2017) How to fight multiple enemies: target-specific chemical defences in an aposematic moth. *Proceedings of the Royal Society B-Biological Sciences*. In press.
- Rojas, B. & Endler, J.A. (2013) Sexual dimorphism and intra-populational colour pattern variation in the aposematic frog *Dendrobates tinctorius*. *Evolutionary Ecology* **27**, 739–753.
- Rojas, B., Gordon, S.P. & Mappes, J. (2015) Frequency-dependent flight activity in the colour polymorphic wood tiger moth. *Current Zoology*, **61**, 765–772.
- Rönkä, K., De Pasqual, C., Mappes, J., Gordon, S., Rojas, B. (2018) Colour alone matters: no predator generalization among morphs of an aposematic moth. *Animal Behaviour* **135**, 153–163.
- Rönkä, K., Mappes, J., Kaila, L. & Wahlberg, N. (2016) Putting Parasemia in its phylogenetic place: a molecular analysis of the subtribe Arctiina (Lepidoptera). *Systematic Entomology*, **41**, 844–853.
- Ruxton, G., Sherratt, T.N. & Speed, M.P. (2004) *Avoiding Attack*. Oxford University Press.
- Telles, F.J., Lind, O., Henze, M.J., Rodriguez-Girones, M.A., Goyret, J. & Kelber, A. (2014) Out of the blue: the spectral sensitivity of hummingbird hawkmoths. *Journal of Comparative Physiology A*, **200**, 537–546.
- Vorobyev, M., Brandt, R., Peitsch, D., Laughlin, S.B. & Menzel, R. (2001) Colour thresholds and receptor noise: behaviour and physiology compared. *Vision Research*, **41**, 639–653.

Vorobyev, M., Osorio, D. (1998) Receptor noise as a determinant of colour thresholds. *Proceedings of the Royal Society B-Biological Sciences*, **265**, 351–358.

Warrant, E.J., Kelber, A. & Kristensen, N.P. (2003) Eyes and vision. In N. P. Kristensen (Ed.) *Handbook of Zoology Vol. IV 36, Lepidoptera 2* (pp. 325–360). Berlin: De Gruyter.

White, R.H., Xu, H.H. Munch, T.A., Bennett, R.R. Grable, E.A. (2003) The retina of *Manduca sexta*: rhodopsin expression, the mosaic of green-, blue- and UV-sensitive photoreceptors, and regional specialization. *Journal of Experimental Biology* 206, 3337–3348.

Wyszecki, G. & Stiles, W.S. (2000) Color science: concepts and methods, quantitative data and formulae, 2nd Edn. New York: Wiley.

Zylinski, S., Osorio, D. (2013) Visual contrast and color in rapid learning of novel patterns by chicks. *Journal of Experimental Biology*, **216**, 4184–4189.

SUPPORTING INFORMATION

Additional supporting information may be found in the online version of this article.

Supplementary methods for ERGs and visual models

Table S1 Sex and time effects on ERGs

Table S2 Chromatic contrasts for 1:1:1 receptor proportions

Table S3 Chromatic contrasts for equally spaced receptors

Figure S1 Achromatic (Michelson) contrasts of wing colours

Figure S2 Chromatic contrasts for 1:1:1 receptor proportions

Figure S3 Effect of illumination on chromatic contrasts

Supplementary references

Table 1. Influence of sex and measurement number on response properties of the dark-adapted retina in the compound eyes of *A. plantuginis* tested by linear mixed-effects models. For a definition of V_{\max} , K and $Slope$ at K see Fig. 2.

	Mean difference	Standard error	95% confidence interval	t	df	p value
V_{\max} [mV]						
Sex	12.0	1.4	(9.1, 15.0)	8.87	12	<0.001
Measurement no.	0.1	0.1	(-0.2, 0.3)	0.54	130	0.59
K [log I]						
Sex	-0.1	0.2	(-0.5, 0.3)	-0.72	12	0.48
Measurement no.	-0.02	0.01	(-0.1, -0.001)	-2.10	130	0.04
$Slope$ at K [mV/ log I]						
Sex	3.1	0.4	(2.2, 4.0)	7.69	12	<0.001
Measurement no.	0.1	0.04	(-0.03, 0.1)	1.23	130	0.22

Table 2. Colour contrast (jnds) between patches of forewings and hindwings in male and female wood tiger moths. For each comparison, we give the median colour contrast with the 25th and 75th percentiles in brackets.

	Male forewing colour		Female forewing colour	
	White	Yellow	Orange	Red
White	0.9 (0.3 - 1.6)	1.0 (0.7-1.4)	Orange	1.1 (0.5-1.8)
Yellow		0.7 (0.3-1.2)	Red	0.5 (0.2-0.9)
	Male hindwing colour		Female hindwing colour	
	White	Yellow	Orange	Red
White	0.1 (0.1-0.3)	2.6 (1.8-3.3)	Orange	1.4 (0.6-2.7)
Yellow		0.9 (0.5-1.5)	Red	0.5 (0.2-0.8)

Figure captions

Fig. 1. Vertical sections of the compound eyes of female (top) and male (bottom) *Arctia plantaginis*. Left, enlarged region of the eye with c cornea, cc crystalline cone, rh rhabdom, b basement membrane, scale bar 50 μm . Right, eyes (R retina) and first optic neuropile, lamina (L), scale bar 100 μm .

Fig. 2. Sex differences in response strength of the dark-adapted retina in *Arctia plantaginis*.

(A) Electroretinograms (ERGs) from the compound eyes of a male and a female moth in response to 40 ms flashes of white light with increasing intensity (see B). For clarity, only every second ERG is shown. (B) Amplitudes V (potential changes from the baseline at stimulus onset to maximal hyperpolarization) plotted against intensity I for the datasets partially shown in A. We fitted a Naka-Rushton function to each dataset and estimated the asymptotic maximal response V_{max} , the intensity K that elicited a half-maximal response, and the *Slope* of the curve at K . The inset illustrates the range of data from all eight males and six females. (C) Sex differences of the parameters V_{max} , K , and *Slope* at K (mean +/- standard error) tested by linear mixed-effects models.

Fig. 3. Spectral sensitivities of male (left) and female (right) *Arctia plantaginis* derived from ERGs. For each individual, averaged measurements ($n = 4$) from one compound eye under different adaptation states were selected to estimate the wavelength of maximal sensitivity (λ_a) for three spectral types of photoreceptor. Example curves used to assess λ_a of the green receptors (A, B), λ_a of the UV receptors with λ_a of the green receptors predetermined (C, D), and λ_a of the blue receptors with λ_a of the green and UV receptors predetermined (E, F). (G, H) Examples, in which the blue receptors were suppressed by blue light, isolating the responses of the UV and green receptors.

Fig. 4. Reflectance of forewing (dashed lines) and hindwing (solid lines) colours of male and female *Arctia plantaginis*. Lines give averages and grey areas standard deviations. Data from 16 white males, 10 yellow males, 8 orange females, 10 red females.

Fig. 5. Data used in model calculations. (A) Spectral sensitivities of photoreceptors in the compound eyes of the wood tiger moth (*Arctia plantaginis*) and the hummingbird hawkmoth (*Macroglossum stellatarum*; data from Telles et al. 2014). The sensitivities for UV and green receptors based on the template (Govardovskii et al. 2000) are the same in both species, but the sensitivity of the blue receptors in *A. plantaginis* is red-shifted. (B) Spectral sensitivities of the four single cones in the blue tit (*Cyanistes caeruleus*; data from Hart et al. 2000). UVS, ultraviolet sensitive, SWS, short-wavelength sensitive, MWS, middle-wavelength sensitive, LWS, long-wavelength sensitive. (C) Reflectance spectra of green and brown vegetation and irradiance of daylight (d65; Wyszecki and Stiles 2000).

Fig 6. The loci of forewing (A) and hindwing (B) colours in the chromaticity diagram of *Arctia plantaginis*. Each data point represents the colour locus given by a reflectance spectrum illuminated by daylight d65. Receptors are adapted to a green background (green triangle), the colour locus of the brown background (brown triangle) is shown for comparison. The Euclidean distance between loci equals their contrast expressed in jnds (see Methods for details). Colour loci are visualized on specific opponent axes in the chromaticity diagram but the contrast between colours is set by receptor noise and independent of specific colour opponency (Vorobyev and Osorio 1998).

Fig 7. Colour contrast between wing patches and (A) a green, or (B) a brown background in *Macroglossum stellatarum* (white), *Arctia plantaginis* (red) and *Cyanistes caeruleus* (blue). Contrast was calculated for forewing and hindwing colours from 16 white and 10 yellow males, and 8 orange and 10 red females. Boxes show the 25th, 50th, and 75th percentiles of colour contrast with whiskers spanning the range of the distributions to a maximum of ± 1.5 times the interquartile range (75th-25th). Data points outside this interval are counted as outliers (crosses). The absolute threshold of detection is set to 1 jnd, however in the moths, this limit is uncertain as the receptor noise levels are not well known (see Methods).













