

JYX



**This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.**

**Author(s):** Vellis, Vyron

**Title:** Quasisymmetric extension on the real line

**Year:** 2018

**Version:** Accepted version (Final draft)

**Copyright:** © 2018 American Mathematical Society

**Rights:** In Copyright

**Rights url:** <http://rightsstatements.org/page/InC/1.0/?language=en>

**Please cite the original version:**

Vellis, V. (2018). Quasisymmetric extension on the real line. *Proceedings of the American Mathematical Society*, 146(6), 2435-2450. <https://doi.org/10.1090/proc/13346>

# QUASISYMMETRIC EXTENSION ON THE REAL LINE

VYRON VELLIS

ABSTRACT. We give a geometric characterization of the sets  $E \subset \mathbb{R}$  for which every quasisymmetric embedding  $f : E \rightarrow \mathbb{R}^n$  extends to a quasisymmetric embedding  $f : \mathbb{R} \rightarrow \mathbb{R}^N$  for some  $N \geq n$ .

## 1. INTRODUCTION

Suppose that  $E$  is a subset of a metric space  $X$  and  $f$  is a quasisymmetric embedding of  $E$  into some metric space  $Y$ . When is it possible to extend  $f$  to a quasisymmetric embedding of  $X$  into  $Y'$  for some metric space  $Y'$  containing  $Y$ ? Questions related to quasisymmetric extensions have been considered by Beurling and Ahlfors [3], Ahlfors [1, 2], Carleson [4], Tukia and Väisälä [11] and Kovalev and Onninen [7].

Tukia and Väisälä [12] showed that for  $M = \mathbb{R}^p, \mathbb{S}^p$ , any quasisymmetric mapping  $f : M \rightarrow \mathbb{R}^n$ , with  $n > p$ , extends to a quasisymmetric homeomorphism of  $\mathbb{R}^n$  when  $f$  is locally close to a similarity. Later, Väisälä [14] extended this result to all compact, co-dimension 1,  $C^1$  or piecewise linear manifolds  $M$  in  $\mathbb{R}^n$ .

In this article we are concerned with the case  $X = \mathbb{R}$  and  $Y = \mathbb{R}^n$ . Specifically, given a set  $E \subset \mathbb{R}$  and a quasisymmetric embedding  $f$  of  $E$  into  $\mathbb{R}^n$ , we ask when is it possible to extend  $f$  to a quasisymmetric embedding of  $\mathbb{R}$  into  $\mathbb{R}^N$  for some  $N \geq n$ . While any bi-Lipschitz embedding of a compact set  $E \subset \mathbb{R}$  into  $\mathbb{R}^n$  extends to a bi-Lipschitz embedding of  $\mathbb{R}$  into  $\mathbb{R}^N$  for some  $N \geq n$  [5], the same is not true for quasisymmetric embeddings. In fact, there exists  $E \subset \mathbb{R}$  and a quasisymmetric embedding  $f : E \rightarrow \mathbb{R}^n$  that can not be extended to a quasisymmetric embedding  $F : \mathbb{R} \rightarrow \mathbb{R}^N$  for any  $N$ ; see e.g. [6, p. 89]. Thus, more regularity for sets  $E$  should be assumed.

Following Trotsenko and Väisälä [10], a metric space  $X$  is termed  *$M$ -relatively connected* for some  $M > 1$  if, for any point  $x \in X$  and any  $r > 0$  with  $\overline{B}(x, r) \neq X$ , either  $\overline{B}(x, r) = \{x\}$  or  $\overline{B}(x, r) \setminus B(x, r/M) \neq \emptyset$ . A metric space  $X$  is called relatively connected if it is  $M$ -relatively connected for some  $M \geq 1$ .

With this terminology, our main theorem is stated as follows.

**Theorem 1.1.** *If  $E \subset \mathbb{R}$  is  $M$ -relatively connected and  $f : E \rightarrow \mathbb{R}^n$  is  $\eta$ -quasisymmetric then  $f$  extends to an  $\eta'$ -quasisymmetric embedding  $F : \mathbb{R} \rightarrow \mathbb{R}^{n+n_0}$  where  $n_0$  depends only on  $M$  and  $\eta$  while  $\eta'$  depends only on  $M, \eta$  and  $n$ .*

On the other hand, it follows from a theorem of Trotsenko and Väisälä [10] that if  $E \subset \mathbb{R}$  is not relatively connected, then there exists a quasisymmetric mapping

---

*Date:* August 24, 2018.

*2010 Mathematics Subject Classification.* Primary 30C65; Secondary 30L05.

*Key words and phrases.* quasisymmetric extension, relatively connected sets.

The author was supported by the Academy of Finland project 257482.

$f : E \rightarrow \mathbb{R}$  that admits no quasisymmetric extension  $F : \mathbb{R} \rightarrow \mathbb{R}^N$  for any  $N \geq 1$ ; see Corollary 2.5.

A subset  $E$  of a metric space  $X$  is said to have the *quasisymmetric extension property in  $X$*  if every quasisymmetric mapping  $f : E \rightarrow X$  that can be extended homeomorphically in  $X$  can also be extended quasisymmetrically in  $X$ . The question of characterizing such sets  $E$ , given a space  $X$ , poses formidable difficulties due to the topological complexity of  $X$ . For instance,  $\mathbb{S}^1$  and  $\mathbb{R}$  have the quasisymmetric extension property in  $\mathbb{R}^2$  [1], but it is unknown whether  $\mathbb{S}^n$  or  $\mathbb{R}^n$  have this property in  $\mathbb{R}^{n+1}$  when  $n \geq 2$ .

The sets  $E \subset \mathbb{R}$  that have the quasisymmetric extension property in  $\mathbb{R}$  are characterized by the relative connectedness.

**Theorem 1.2.** *A set  $E \subset \mathbb{R}$  has the quasisymmetric extension property in  $\mathbb{R}$  if and only if it is relatively connected.*

The arguments used in the proof of Theorem 1.2 apply verbatim in the case  $X = \mathbb{S}^1$  and  $E \subset \mathbb{S}^1$ . Thus, if  $X$  is quasisymmetric homeomorphic to either  $\mathbb{R}$  or  $\mathbb{S}^1$ , then a set  $E \subset X$  has the quasisymmetric extension property in  $X$  if and only if  $E$  is relatively connected.

In dimensions  $n \geq 2$ , however, Theorem 1.2 fails even for small sets such as the Cantor sets. In Section 5 we show that for each  $n \geq 2$ , there exists a relatively connected Cantor set  $E \subset \mathbb{R}^n$  and a bi-Lipschitz mapping  $f : E \rightarrow \mathbb{R}^n$  which admits a homeomorphic extension in  $\mathbb{R}^n$ , but not a quasisymmetric extension in  $\mathbb{R}^n$ ; see Remark 5.2.

**Acknowledgements.** The author wishes to thank Pekka Koskela for bringing this problem to his attention, Tuomo Ojala for various discussions and the anonymous referee for valuable comments on the manuscript.

## 2. PRELIMINARIES

In the following, given an open bounded interval  $I = (a, b) \subset \mathbb{R}$ , we denote by  $|I|$  its length  $b - a$ ; if  $I = \emptyset$  then  $|I| = 0$ . As usual,  $a \vee b$  and  $a \wedge b$  denote the maximum and minimum, respectively, of two real numbers  $a$  and  $b$ . Finally, for two points  $x, y \in \mathbb{R}^n$ , we denote by  $[x, y]$  the line segment in  $\mathbb{R}^n$  with endpoints  $x$  and  $y$ .

**2.1. Mappings.** A homeomorphism  $f : (X, d) \rightarrow (Y, d')$  between two metric spaces is called  *$L$ -bi-Lipschitz* for some  $L > 1$  if both  $f$  and  $f^{-1}$  are  $L$ -Lipschitz.

A mapping  $f : (X, d) \rightarrow (Y, d')$  is called  $\eta$ -quasisymmetric if there exists a homeomorphism  $\eta : [0, +\infty) \rightarrow [0, +\infty)$  such that for any  $x, a, b \in X$  with  $x \neq b$  we have

$$\frac{d'(f(x), f(a))}{d'(f(x), f(b))} \leq \eta \left( \frac{d(x, a)}{d(x, b)} \right).$$

It is a simple consequence of the definition that the composition of a similarity mapping of  $\mathbb{R}^n$  and an  $\eta$ -quasisymmetric mapping between sets of  $\mathbb{R}^n$  is  $\eta$ -quasisymmetric.

If  $f$  is  $\eta$ -quasisymmetric with  $\eta(t) = C(t^\alpha \vee t^{1/\alpha})$  for some  $\alpha \in (0, 1]$  and  $C > 0$  then  $f$  is termed *power quasisymmetric* and we say that  $f$  is  $(C, \alpha)$ -quasisymmetric. An important property of power quasisymmetric mappings is that they are bi-Hölder continuous on bounded sets [6, Corollary 11.5].

**Lemma 2.1.** *Suppose that  $(X, d)$  is a bounded metric space and  $f : (X, d) \rightarrow (Y, d')$  is  $(C, \alpha)$ -quasisymmetric. There exists  $C' > 1$  depending only on  $C, \alpha, \text{diam } X$  and  $\text{diam } f(X)$  such that for all  $x, y \in E$ ,*

$$(C')^{-1}d(x, y)^{1/\alpha} \leq d'(f(x), f(y)) \leq C'd(x, y)^\alpha.$$

For doubling connected metric spaces it is known that the quasisymmetric condition is equivalent to a weaker (but simpler) condition known in literature as *weak quasisymmetry*.

**Lemma 2.2** ([6, Theorem 10.19]). *Let  $I \subset \mathbb{R}$  be an interval and  $f : I \rightarrow \mathbb{R}^n$  be an embedding for which there exists  $H \geq 1$  such that for all  $x, y, z \in I$*

$$(1) \quad |x - y| \leq |x - z| \text{ implies } |f(x) - f(y)| \leq H|f(x) - f(z)|.$$

*Then  $f$  is  $\eta$ -quasisymmetric with  $\eta$  depending only on  $H$  and  $n$ .*

The next lemma is an immediate corollary to Lemma 2.2.

**Lemma 2.3.** *Let  $I_1, I_2$  be open bounded intervals and  $f : I_1 \cup I_2 \rightarrow \mathbb{R}$  be an embedding. Suppose that there exists  $C > 1$  such that  $|I|/|J| < C$  for all  $I, J \in \{I_1, I_2, I_1 \cap I_2\}$ . If  $f|_{I_1}$  and  $f|_{I_2}$  are  $\eta$ -quasisymmetric then  $f|(I_1 \cup I_2)$  is  $\eta'$ -quasisymmetric for some  $\eta'$  depending on  $\eta$  and  $C$ .*

*Proof.* If  $I_1 \subset I_2$  or  $I_2 \subset I_1$  there is nothing to prove. Suppose that  $I_1 = (a_1, b_1)$ ,  $I_2 = (a_2, b_2)$  with  $a_1 < a_2 < b_1 < b_2$  and denote by  $m$  the center of  $I_1 \cap I_2$ . We show that  $f|(I_1 \cup I_2)$  satisfies (1). Let  $x, y, z \in I_1 \cup I_2$  with  $|x - y| \leq |x - z|$ . Since  $f|_{I_j}$  is monotone for each  $j = 1, 2$ ,  $f|_{I_1 \cup I_2}$  is monotone and we may assume that either  $y < x < z$  or  $z < x < y$ . Assume the first; the second case is identical

If all three points are in the same  $I_j$  there is nothing to prove. Hence, we may assume that  $y \leq a_2$  and  $z \geq b_1$ .

$$\text{If } x \leq m \text{ then } |f(x) - f(y)| \leq \eta \left( \frac{|x-y|}{|x-b_1|} \right) |f(x) - f(z)| \leq \eta(2C) |f(x) - f(z)|.$$

If  $x \geq m$  then  $|f(x) - f(y)| = |f(x) - f(a_2)| + |f(a_2) - f(y)| \leq |f(x) - f(a_2)| (1 + \frac{|f(a_2) - f(a_1)|}{|f(a_2) - f(m)|}) \leq |f(x) - f(a_2)| (1 + \eta(\frac{|a_1 - a_2|}{|a_2 - m|})) \leq (1 + \eta(2C)) |f(x) - f(a_2)| \leq (1 + \eta(2C)) \eta \left( \frac{|x - a_2|}{|x - z|} \right) |f(x) - f(z)| \leq (1 + \eta(2C)) \eta(1) |f(x) - f(z)|$  where for the last inequality we used  $|x - a_2| \leq |x - y| \leq |x - z|$ .  $\square$

**2.2. Relatively connected sets.** Relatively connected sets were first introduced by Trotsenko and Väisälä [10] in the study of spaces for which every quasisymmetric mapping is power quasisymmetric. The definition given in [10] is equivalent to the one in Section 1 quantitatively [10, Theorem 4.11].

Relative connectedness is a weak form of the well known notion of uniform perfectness. A metric space  $X$  is *c-uniformly perfect* for some  $c > 1$  if for all  $x \in X$ ,  $\overline{B}(x, r) \neq X$  implies  $\overline{B}(x, r) \setminus B(x, r/c) \neq \emptyset$ . The difference between the two notions is that relatively connected sets allow isolated points. In particular, if  $E$  is  $c$ -uniformly perfect, then it is  $M$ -relatively connected for all  $M > c$ , and if  $E$  is  $M$ -relatively connected and has no isolated points, then it is  $(2M + 1)$ -uniformly perfect [10, Theorem 4.13].

The connection between relative connectedness and power quasisymmetric mappings is illustrated in the following theorem from [10].

**Theorem 2.4** ([10, Theorem 6.20]). *A subset  $E$  of a metric space  $X$  is relatively connected if and only if every quasisymmetric map  $f : E \rightarrow X$  is power quasisymmetric.*

The necessity of relative connectedness for extensions of quasisymmetric mappings on  $\mathbb{R}$  follows now as a corollary.

**Corollary 2.5.** *If  $E \subset \mathbb{R}$  is not relatively connected, then there exists a monotone quasisymmetric mapping  $f : E \rightarrow \mathbb{R}$  such that, for every metric space  $Y$  containing the Euclidean line  $\mathbb{R}$ , there exists no quasisymmetric extension  $F : \mathbb{R} \rightarrow Y$  of  $f$ .*

*Proof.* By [10, Theorem 6.6], there exists a quasisymmetric mapping  $f : E \rightarrow \mathbb{R}$  that is not power quasisymmetric. A close inspection in its proof reveals, moreover, that the mapping  $f$  is increasing. Let now  $Y$  be a metric space containing the Euclidean line  $\mathbb{R}$ . If there was a quasisymmetric extension  $F : \mathbb{R} \rightarrow Y$ , then, by Theorem 2.4,  $F$  would be power quasisymmetric. Thus,  $f$  would be power quasisymmetric which is a contradiction.  $\square$

**2.3. Relative distance.** Let  $E, F$  be two compact sets in a metric space  $(X, d)$  both of which contain at least two points. The *relative distance* of  $E$  and  $F$  is defined to be the quantity

$$d^*(E, F) = \frac{\text{dist}(E, F)}{\text{diam } E \wedge \text{diam } F}$$

where  $\text{dist}(E, F) = \min\{d(x, y) : x \in E, y \in F\}$ .

Note that if  $E, F \subset \mathbb{R}^n$  and  $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is a similarity then  $d^*(f(E), f(F)) = d^*(E, F)$ . In general, if  $f : E \cup F \rightarrow Y$  is  $\eta$ -quasisymmetric, then

$$(2) \quad \frac{1}{2}\phi(d^*(E, F)) \leq d^*(f(E), f(F)) \leq \eta(2d^*(E, F))$$

where  $\phi(t) = (\eta(t^{-1}))^{-1}$ ; see for example [13, p. 532].

The following remark ties the notions of uniform perfectness in  $\mathbb{R}$  and relative distance of sets in  $\mathbb{R}$ .

*Remark 2.6.* A closed set  $E \subset \mathbb{R}$  is  $c$ -uniformly perfect for some  $c \geq 1$  if and only if there exists  $C > 0$  such that for all bounded components  $I, J$  of  $\mathbb{R} \setminus E$ ,  $d^*(I, J) \geq C$ . The constants  $c$  and  $C$  are quantitatively related.

### 3. QUASISYMMETRIC EXTENSION ON $\mathbb{R}$

Suppose that  $E \subset \mathbb{R}$  is relatively connected and  $f : E \rightarrow \mathbb{R}^n$  is quasisymmetric. If  $E$  is a singleton then trivially  $f$  admits a quasisymmetric extension. Moreover, since quasisymmetric functions have a quasisymmetric extension to the closure of their domains, we may assume that  $E$  is closed.

In Section 3.1 we construct a quasisymmetric extension  $f_0 : E_0 \rightarrow \mathbb{R}^m$  of  $f$ , where  $E \subset E_0 \subset \mathbb{R}$  is a uniformly perfect set with no lower or upper bound and  $m$  is either  $n$  or  $n + 1$ . In Section 3.2, for some  $n_0 \in \mathbb{N}$  depending only on  $M$  and  $\eta$ , we construct a homeomorphic extension  $F_0 : \mathbb{R} \rightarrow \mathbb{R}^{n+n_0}$  of  $f_0$ . Finally, in Section 3.3 we construct a quasisymmetric extension  $F : \mathbb{R} \rightarrow F_0(\mathbb{R}) \subset \mathbb{R}^{n+n_0}$  of  $f_0$ .

For the rest,  $\mathbf{0}$  denotes the origin of  $\mathbb{R}^n$  and, for each  $i = 1, \dots, n$ ,  $\mathbf{e}_i$  denotes the vector in  $\mathbb{R}^n$  whose  $i$ -th coordinate is 1 and the rest are 0.

**3.1. Two preliminary extensions.** Throughout this section we assume that  $E$  is an  $M$ -relatively connected closed set and  $f$  is an  $\eta$ -quasisymmetric embedding of  $E$  into  $\mathbb{R}^n$  with  $\eta = C(t^\alpha \vee t^{1/\alpha})$ .

Suppose that  $E$  is bounded from above or bounded from below. Then one of the following cases applies.

*Case 1.* Suppose that  $E$  has a lower bound but no upper bound. Applying suitable similarities we may assume that  $1 \in E$ ,  $\min E = 0$  and  $f(0) = \mathbf{0}$ . Let  $C_0 = \max\{2, 1/\eta^{-1}(1/2)\}$ . Set  $a_0 = 0$  and, by relative connectedness, there exists a sequence  $\{a_k\}_{k \in \mathbb{N}} \subset E$  with  $a_1 = 1$  and  $a_k/a_{k-1} \in [C_0, MC_0]$ . Set  $\tilde{E} = E \cup \{-a_k\}_{k \in \mathbb{N}}$  and  $\tilde{f} : \tilde{E} \rightarrow \mathbb{R}^{n+1}$  with  $\tilde{f}|_E = f \times \{0\}$  and  $\tilde{f}(-a_k) = \{\mathbf{0}\} \times \{-|f(a_k)|\}$ .

*Case 2.* Suppose that  $E$  has an upper bound but no lower bound. Applying suitable similarities we may assume that  $1 \in E$ ,  $\max E = 0$  and  $f(0) = \mathbf{0}$ . We define  $\tilde{E}$  and  $\tilde{f}$  similarly to Case 1.

*Case 3.* Suppose that  $E$  is bounded. Applying suitable similarities, we may assume that  $\min E = 0$ ,  $\max E = 1$ ,  $\max_{x \in E} |f(x)| = 1$  and  $\text{diam } f(E) = 1$ . For any  $k \in \mathbb{Z}$  define  $\tilde{E}_k = \{2k + x : x \in E\}$ ,  $\tilde{E} = \bigcup_{k \in \mathbb{Z}} \tilde{E}_k$  and  $\tilde{f} : \tilde{E} \rightarrow \mathbb{R}^n$  with  $\tilde{f}(2k + x) = 2k\mathbf{e}_1 + f(x)$ . A similar extension in the case  $n = 1$  has been considered by Lehto and Virtanen in [8, II.7.2].

**Lemma 3.1.** *In each case,  $\tilde{E}$  is an  $\tilde{M}$ -relatively connected closed set and  $\tilde{f}$  is  $\tilde{\eta}$ -quasisymmetric with  $\tilde{M}$  and  $\tilde{\eta}$  depending only on  $M$  and  $\eta$ .*

*Proof.* We only prove the lemma for Case 1 and Case 3; the proof for Case 2 is similar to that of Case 1.

*Case 1.* Note first that  $\{-a_n\}_{n \in \mathbb{N}}$  is  $M_1$ -relatively connected for some  $M_1$  depending only on  $M$  and  $\eta$ . Let  $x \in \tilde{E}$  and  $r > 0$  such that  $\overline{B}(x, r) \cap \tilde{E} \neq \{x\}$ . If  $x \in E$  then  $\overline{B}(x, r) \cap E \neq \{x\}$  and  $(\overline{B}(x, r) \setminus B(x, r/M)) \cap \tilde{E} \neq \emptyset$ . If  $x = -a_n$ ,  $n \geq 1$ , then  $\overline{B}(x, r) \cap \{-a_n\}_{n \in \mathbb{N}} \neq \{x\}$  and  $(\overline{B}(x, r) \setminus B(x, r/M_1)) \cap \tilde{E} \neq \emptyset$ . Thus,  $\tilde{E}$  is  $(M \vee M_1)$ -relatively connected.

For the quasisymmetry of  $\tilde{f}$ , note first that  $\tilde{f}$  restricted on  $\{-a_n\}_{n \in \mathbb{N}}$  is  $C\eta$ -quasisymmetric for some  $C > 1$  depending only on  $\eta$ . Let  $x, y, z \in \tilde{E}$ . If all three of them are in  $E$  or in  $\tilde{E} \setminus E$  then the quasisymmetry of  $\tilde{f}$  follows trivially.

Assume first that  $x, z \in E$  and  $y = -a_n$  for some  $a_n \in E$ . Then,  $|\tilde{f}(y)| = |f(a_n)|$ ,  $|y| = |a_n|$  and

$$\begin{aligned} \frac{|\tilde{f}(x) - \tilde{f}(y)|}{|\tilde{f}(x) - \tilde{f}(z)|} &\leq 2 \frac{|f(x)|}{|f(x) - f(z)|} + \frac{|f(x) - f(a_n)|}{|f(x) - f(z)|} \\ &\leq 2\eta \left( \frac{|x|}{|x - z|} \right) + \eta \left( \frac{|x - a_n|}{|x - z|} \right) \leq 3\eta \left( \frac{|x - y|}{|x - z|} \right). \end{aligned}$$

We work similarly if  $x, z \in \{-a_n\}_{n \in \mathbb{N}}$  and  $y \in E$ .

Assume now that  $z \in E$  and  $y, x \notin E$ . Let  $n_0$  be the smallest integer  $n$  such that  $a_n \geq z$  and set  $\bar{z} = -a_{n_0}$ . Then, there exist constants  $C_1, C_2 > 1$  depending only on  $M, C$  and  $\alpha$  such that

$$\begin{aligned} \frac{|\tilde{f}(x) - \tilde{f}(y)|}{|\tilde{f}(x) - \tilde{f}(z)|} &\leq C_1 \min \left\{ \frac{|\tilde{f}(x) - \tilde{f}(y)|}{|\tilde{f}(x) - \tilde{f}(\bar{z})|}, \frac{|\tilde{f}(x) - \tilde{f}(y)|}{|\tilde{f}(x)|} \right\} \\ &\leq C_1 \min \left\{ \eta \left( \frac{|x - y|}{|x - \bar{z}|} \right), \eta \left( \frac{|x - y|}{|x|} \right) \right\} \leq C_2 \eta \left( \frac{|x - y|}{|x - z|} \right). \end{aligned}$$

We work similarly if  $z \in \{-a_n\}_{n \in \mathbb{N}}$  and  $x, y \in E$ .

*Case 3.* We first show that  $\tilde{E}$  is  $M'$ -relatively connected with  $M' = 8M$ . Let  $x \in \tilde{E}$  and  $r > 0$  such that  $\overline{B}(x, r) \cap \tilde{E} \neq \{x\}$ . Since  $\tilde{E}$  is unbounded,  $\tilde{E} \setminus \overline{B}(x, r) \neq \emptyset$ . By periodicity of  $\tilde{E}$ , we may assume that  $x \in E$ . If  $r \geq 4$  then  $\overline{B}(x, r) \setminus B(x, r/2)$  contains an interval of length 2 and therefore it contains points of  $\tilde{E}$ . Suppose now

that  $r < 4$ . Then,  $\tilde{E} \cap B(x, r/8) \subset E$  and  $E \setminus \overline{B}(x, r/8) \neq \emptyset$ . If  $E \cap \overline{B}(x, r/8) = \{x\}$  then  $\tilde{E} \cap \overline{B}(x, r/8) = \{x\}$  and the relative connectedness is satisfied with  $M' = 8$ . If  $E \cap \overline{B}(x, r/8) \neq \{x\}$  then, by the relative connectedness of  $E$ ,  $E \cap (\overline{B}(x, r) \setminus B(x, r/(8M))) \neq \emptyset$ .

We show now the second claim. Recall that by Theorem 2.4  $f$  is power quasymmetric. Let  $y, x, z \in \tilde{E}$  and assume  $y \in \tilde{E}_{n_1}$ ,  $x \in \tilde{E}_{n_2}$  and  $z \in \tilde{E}_{n_3}$  with  $n_1, n_2, n_3 \in \mathbb{Z}$ . If  $n_1 = n_2 = n_3$  the claim follows trivially. If  $n_1, n_2, n_3$  are all different then

$$\frac{|\tilde{f}(x) - \tilde{f}(y)|}{|\tilde{f}(x) - \tilde{f}(z)|} \leq \frac{2|n_2 - n_1| + 1}{2|n_3 - n_2| - 1} \leq 9 \frac{|n_2 - n_1|}{|n_3 - n_2| + 2} \leq 9 \frac{|x - y|}{|x - z|}.$$

If  $n_1 = n_2 \neq n_3$  then the second inequality in Lemma 2.1 gives

$$\frac{|\tilde{f}(x) - \tilde{f}(y)|}{|\tilde{f}(x) - \tilde{f}(z)|} \leq C' \frac{|x - y|^\alpha}{|n_3 - n_2|} \leq 3C' \left( \frac{|x - y|}{|x - z|} \right)^\alpha.$$

The remaining case  $n_1 \neq n_2 = n_3$  is treated similarly using the first inequality of Lemma 2.1.  $\square$

By Lemma 3.1 we may assume for the rest that  $E$  is a relatively connected closed set with no upper or lower bound. Hence, all components of  $\mathbb{R} \setminus E$  are bounded open intervals.

For the second extension, we treat the case when  $E$  has isolated points. For each isolated point  $x \in E$  let  $\pi(x) \in E$  be the closest point of  $E \setminus \{x\}$  to  $x$  and define

$$E_x = \overline{B}(x, |x - \pi(x)|/10)$$

and  $f_x : E_x \rightarrow \mathbb{R}^n$  with

$$f_x(y) = f(x) + \frac{1}{\eta(1)} \frac{|f(x) - f(\pi(x))|}{|x - \pi(x)|} (y - x) \mathbf{e}_1.$$

If  $x$  is an accumulation point of  $E$ , then set  $E_x = \{x\}$  and  $f_x : \{x\} \rightarrow \mathbb{R}$  with  $f_x(x) = f(x)$ . Finally, set  $\hat{E} = \bigcup_{x \in E} E_x$  and  $\hat{f} : \hat{E} \rightarrow \mathbb{R}$  with  $\hat{f}|_{E_x} = f_x$ . Similar extensions also appear in a paper of Semmes [9, Section 2].

*Remark 3.2.* Suppose that  $x \in E$  is an isolated point. Then,

$$4 \leq d^*(E_x, \hat{E} \setminus E_x) \leq 5 \quad \text{and} \quad 3 \leq d^*(\hat{f}(E_x), \hat{f}(\hat{E} \setminus E_x)) \leq 5\eta(1).$$

The first claim of Remark 3.2 is clear. For the upper bound of the second claim note that  $\text{dist}(\hat{f}(E_x), \hat{f}(\hat{E} \setminus E_x)) \leq |f(x) - f(\pi(x))| \leq 5\eta(1) \text{diam } \hat{f}(E_x)$ . For the lower bound, take points  $x' \in E_x$  and  $y' \in \hat{E} \setminus E_x$  and assume that  $y' \in E_y$ . Then,

$$(3) \quad \frac{|\hat{f}(x') - \hat{f}(y')|}{|\hat{f}(x') - \hat{f}(y')|} \leq \frac{1}{10\eta(1)} \eta \left( \frac{|x - \pi(x)|}{|x - y|} \right) \frac{|f(x) - \hat{f}(y)|}{\frac{4}{5}|f(x) - \hat{f}(y)|} \leq \frac{1}{8}.$$

Thus, if  $x'$  is an endpoint of  $E_x$ , (3) yields  $\text{dist}(\hat{f}(x'), \hat{f}(\hat{E} \setminus E_x)) \geq 4 \text{diam } \hat{f}(E_x)$ . Hence,  $\text{dist}(\hat{f}(E_x), \hat{f}(\hat{E} \setminus E_x)) \geq 3 \text{diam } \hat{f}(E_x)$  and the lower bound follows.

**Lemma 3.3.** *The set  $\hat{E}$  is closed and  $c$ -uniformly perfect and  $\hat{f} : \hat{E} \rightarrow \mathbb{R}^n$  is  $\hat{\eta}$ -quasisymmetric where  $c$  depends only on  $M$  and  $\hat{\eta}$  depends only on  $\eta$ .*

*Proof.* Clearly,  $\overline{E_x} \cap E_y = \emptyset$  for  $x, y \in E$  with  $x \neq y$ . To see that  $\hat{E}$  is closed, take  $y \in \hat{E}$ . If  $y \in \hat{E} \setminus E$  then  $y \in \overline{E_x}$  for some  $x \in E$  and, thus,  $y \in \hat{E}$ .

Since  $\hat{E}$  has no isolated points, we only need to show that  $\hat{E}$  is  $M'$ -relatively connected for some  $M'$  depending on  $M$ . Take  $x \in \hat{E}$  and  $r > 0$ . From the unboundedness of  $\hat{E}$  and the fact that  $\hat{E}$  has no isolated points, we have  $\{x\} \subsetneq \overline{B}(x, r) \cap \hat{E} \subsetneq \hat{E}$ . If  $x \in E$  is not isolated in  $E$ , then

$$\emptyset \neq E \cap (\overline{B}(x, r) \setminus B(x, r/M)) \subset \hat{E} \cap (\overline{B}(x, r) \setminus B(x, r/M)).$$

Suppose  $x \in E_z$  for some isolated point  $z$  in  $E$ . If  $r > 2M \operatorname{dist}(z, E \setminus \{z\})$  then  $\emptyset \neq (E \setminus \{z\}) \cap \overline{B}(z, r/2) \subset \hat{E} \cap \overline{B}(z, r/2)$ . Therefore,

$$\emptyset \neq E \cap (\overline{B}(z, r/2) \setminus B(z, (2M)^{-1}r)) \subset \hat{E} \cap (\overline{B}(z, r/2) \setminus B(z, (2M)^{-1}r)).$$

If  $r \leq 2M \operatorname{dist}(z, E \setminus \{z\})$  then  $(20M)^{-1}r \leq \frac{1}{10} \operatorname{dist}(z, E \setminus \{z\})$  and

$$\emptyset \neq E_z \cap (\overline{B}(x, r) \setminus B(x, (20M)^{-1}r)) \subset \hat{E} \cap (\overline{B}(x, r) \setminus B(x, (20M)^{-1}r)).$$

It remains to show that  $\hat{f}$  is quasisymmetric; then by Theorem 2.4  $\hat{f}$  will be power quasisymmetric. Let  $x, y, z \in \hat{E}$  be three distinct points with  $x \in E_{x'}$ ,  $y \in E_{y'}$  and  $z \in E_{z'}$  for some  $x', y', z' \in E$ . If  $x' = y' = z'$  then  $x, y, z$  are in an interval where  $\hat{f}$  is a similarity.

If  $x' \neq z'$  and  $x' = y'$  then, by Remark 3.2, the prerequisites of Lemma 2.29 in [9] are satisfied for  $A = E \setminus \{x'\}$ ,  $A^* = E \cup E_{x'}$  and  $H = \hat{f}|_{A^*}$  and  $\hat{f}|_{E \cup E_{x'}}$  is  $\eta'$ -quasisymmetric for some  $\eta'$  depending only on  $\eta$ . Hence,

$$\frac{|\hat{f}(x) - \hat{f}(y)|}{|\hat{f}(x) - \hat{f}(z)|} \leq C_1 \frac{|\hat{f}(x) - \hat{f}(y)|}{|\hat{f}(x) - \hat{f}(z')|} \leq C_1 \eta' \left( \frac{|x - y|}{|x - z'|} \right) \leq C_1 \eta' \left( C_2 \frac{|x - y|}{|x - z|} \right)$$

for some  $C_1, C_2 > 1$  depending only on  $\eta$ . Similarly for  $x' = z' \neq y'$ . If  $x', y', z'$  are distinct then by Remark 3.2,

$$\frac{|\hat{f}(x) - \hat{f}(y)|}{|\hat{f}(x) - \hat{f}(z)|} \leq C_3 \frac{|\hat{f}(x') - \hat{f}(y')|}{|\hat{f}(x') - \hat{f}(z')|} \leq C_3 \eta \left( \frac{|x' - y'|}{|x' - z'|} \right) \leq C_3 \eta \left( C_4 \frac{|x - y|}{|x - z|} \right)$$

for some constants  $C_3, C_4 > 1$  depending only on  $\eta$ . Thus,  $\hat{f}$  is quasisymmetric.  $\square$

**3.2. Bridges.** By Lemma 3.1 and Lemma 3.3, we may assume that  $E$  is a closed  $c$ -uniformly perfect set such that every component of  $\mathbb{R} \setminus E$  is a bounded open interval, and  $f : E \rightarrow \mathbb{R}^n$  is an  $\eta$ -quasisymmetric embedding.

In this section, for each component  $I$  of  $\mathbb{R} \setminus E$ , we construct a path in a higher dimensional space  $\mathbb{R}^N$ ,  $N \geq n$ , connecting the images of the endpoints of  $I$ . The union of these paths along with  $f(E)$  gives a homeomorphic extension  $F : \mathbb{R} \rightarrow \mathbb{R}^N$ .

For two points  $x, y \in \mathbb{R}^n \subset \mathbb{R}^k$  let  $T_k(x, y)$  be the equilateral triangle which contains the line segment  $[x, y]$  and lies on the 2-dimensional plane defined by the points  $x, y$  and  $\mathbf{e}_k$ . The *bridge* of  $x$  and  $y$  in dimension  $k$ , denoted by  $\mathcal{B}_k(x, y)$ , is the closure of  $T_k(a, b) \setminus [x, y]$ .

*Remark 3.4.* If  $z, a, b \in \mathbb{R}^n$  with  $|z - a| \leq |z - b|$  then, for all  $x \in \mathcal{B}_k(a, b)$ ,  $|z - x| \geq C^{-1}(|z - a| + |x - a|)$  for some universal  $C > 1$ .

*Remark 3.5.* Each bridge  $\mathcal{B}_k(x, y)$  is 4-bi-Lipschitz equivalent to a closed interval of  $\mathbb{R}$  of length  $|x - y|$ .



Using Remark 3.4 and triangle inequality, it is easy to verify that the relative distance of two bridges  $\mathcal{B}_k(x_1, y_1)$  and  $\mathcal{B}_m(x_2, y_2)$ , with  $k \neq m$ , is comparable to the relative distance of the sets  $\{x_1, y_1\}$  and  $\{x_2, y_2\}$ .

*Remark 3.6.* Let  $n, m_1, m_2 \in \mathbb{N}$  with  $n < m_1 \leq m_2$  and let  $x_1, y_1, x_2, y_2 \in \mathbb{R}^n$ . There exists a universal  $C_1 > 0$  such that

$$d^*(\mathcal{B}_{m_2}(x_1, y_1), \mathcal{B}_{m_1}(x_2, y_2)) \leq C_1 d^*(\{x_1, y_1\}, \{x_2, y_2\}).$$

On the other hand, there exist universal constants  $d_0 > 0$  and  $C_2 > 0$  such that  $d^*(\{x_1, y_1\}, \{x_2, y_2\}) \geq d_0$  implies

$$d^*(\{x_1, y_1\}, \{x_2, y_2\}) \leq C_2 d^*(\mathcal{B}_{m_2}(x_1, y_1), \mathcal{B}_{m_1}(x_2, y_2)).$$

For each component  $I$  of  $\mathbb{R} \setminus E$  we denote by  $a_I, b_I$  the endpoints of  $I$  with  $a_I < b_I$  and by  $m_I$  the center of  $I$ . We also write  $\mathcal{B}_k(I) = \mathcal{B}_k(f(a_I), f(b_I))$  where  $k > n$ . In general, two bridges  $\mathcal{B}_k(I)$  and  $\mathcal{B}_k(I')$ , with  $I \neq I'$ , may intersect. Therefore, more dimensions are needed to make sure that such an intersection will never happen. The next lemma allows us to use only a finite amount of dimensions for this purpose.

**Lemma 3.7.** *Let  $d > 0$ . If  $I_1, \dots, I_k$  are mutually disjoint closed intervals in  $\mathbb{R}$  with  $d^*(I_i, I_j) \leq d$  for all  $i, j = 1, \dots, k$ ,  $i \neq j$ , then  $k \leq 2d + 3$ .*

*Proof.* We may assume that if  $i \notin \{1, k\}$ ,  $x \in I_1$ ,  $y \in I_i$  and  $z \in I_k$  then  $x < y < z$ . Furthermore, applying a similarity we may assume that  $\text{dist}(I_1, I_k) = 1$ .

Since  $d^*(I_1, I_k) \leq d$ , we have  $\text{diam } I_1 \wedge \text{diam } I_k \geq d^{-1}$ . Since the intervals  $I_2, \dots, I_{k-1}$  are between  $I_1$  and  $I_k$ , there exists at least one  $j \in \{2, \dots, k-1\}$  such that  $\text{diam } I_j \leq \text{dist}(I_1, I_k)/(k-2) = (k-2)^{-1}$ . Thus,  $\text{dist}(I_1, I_j) \vee \text{dist}(I_k, I_j) \geq \frac{1}{2}(1 - \frac{1}{k-2})$ . If  $\text{diam } I_j \geq d^{-1}$  then  $k \leq d + 2$ . Otherwise,

$$d \geq d^*(I_1, I_j) \vee d^*(I_k, I_j) \geq \frac{\text{dist}(I_1, I_j) \vee \text{dist}(I_k, I_j)}{d^{-1} \wedge \text{diam } I_j} \geq \frac{1}{2}(k-3). \quad \square$$

Let now  $I_1, I_2, \dots$  be an enumeration of the components of  $\mathbb{R} \setminus E$ . By Remark 3.6 and (2), there exists  $C_0 > 0$  so that  $d^*(\overline{I_i}, \overline{I_j}) \geq C_0$  implies  $d^*(\mathcal{B}_m(I_i), \mathcal{B}_m(I_j)) \geq 1$  for all  $m > n$ . By Lemma 3.7, there exists  $n_0 \in \mathbb{N}$ , depending only on  $c$  and  $\eta$ , such that if distinct  $J_1, \dots, J_k \in \{I_1, I_2, \dots\}$  with  $d^*(J_i, J_j) < C_0$  for all  $i \neq j$  then  $k \leq n_0$ . Set  $N = n + n_0 + 1$ . Let  $\mathcal{B}_{n_{I_1}}(I_1)$  be the bridge with  $n_{I_1} = n + 1$ . Suppose that  $\mathcal{B}_{n_{I_1}}(I_1), \dots, \mathcal{B}_{n_{I_m}}(I_m)$  have been defined. Then, there exist at most  $n_0$  indices  $i_1, \dots, i_k$  in  $\{1, \dots, m\}$  such that  $d^*(I_{m+1}, I_{i_j}) < C_0$ . Pick  $n_{I_{m+1}} \in \{n+1, \dots, N\} \setminus \{n_{I_{i_1}}, \dots, n_{I_{i_k}}\}$  and define the bridge  $\mathcal{B}_{n_{I_{m+1}}}(I_{m+1})$ . Inductively, for each component  $I$  of  $\mathbb{R} \setminus E$  we obtain a bridge  $\mathcal{B}_{n_I}(I)$  with  $n_I \leq N$ .

**Corollary 3.8.** *Set  $I' = \{f(a_I), f(b_I)\}$  for any component  $I = (a_I, b_I)$  of  $\mathbb{R} \setminus E$ . Then, there exist  $C > 1$  depending only on  $c$  and  $\eta$  such that, for every two components  $I, J$  of  $\mathbb{R} \setminus E$  with  $I \neq J$ ,*

$$(C)^{-1} d^*(I', J') \leq d^*(\mathcal{B}_{n_I}(I), \mathcal{B}_{n_J}(J)) \leq C d^*(I', J')$$

and  $C^{-1} \text{dist}(I', J') \leq \text{dist}(\mathcal{B}_{n_I}(I), \mathcal{B}_{n_J}(J)) \leq C \text{dist}(I', J')$ .

**3.3. Reflected sets and functions.** As before, we assume that  $E$  is a closed  $c$ -uniformly perfect set such that every component of  $\mathbb{R} \setminus E$  is a bounded open interval, and  $f : E \rightarrow \mathbb{R}^N$  is an  $\eta$ -quasisymmetric embedding with  $N = n + n_0 + 1$ .

Recall from Section 3.2 that, given a component  $I = (a_I, b_I)$  of  $\mathbb{R} \setminus E$ , we denote by  $m_I$  the midpoint of  $I$ . Moreover, we denote by  $m_{\mathcal{B}(I)}$  the point in  $\mathcal{B}_{n_I}(I)$  such that  $\mathcal{B}_{n_I}(I) = [f(a_I), m_{\mathcal{B}(I)}] \cup [f(b_I), m_{\mathcal{B}(I)}]$ . Note that  $[f(a_I), m_{\mathcal{B}(I)}] \cap [f(b_I), m_{\mathcal{B}(I)}] = \{m_{\mathcal{B}(I)}\}$ .

Let  $I = (a_I, b_I)$  be a component of  $\mathbb{R} \setminus E$ . We define an increasing sequence in  $E$  converging to  $a_I$  as follows. Set  $\delta_0 = \min\{1/2, \eta^{-1}(1/2)\}$ . Since  $E$  is uniformly perfect, there exists  $a_0 \in E$ ,  $a_0 < a_I$  with  $|a_0 - a_I| \in [(2c)^{-1}|I|, 2^{-1}|I|]$ . Inductively, suppose that  $a_k$  has been defined. Since  $E$  is uniformly perfect, there exists  $a_{k+1} \in E \cap (a_k, a_I)$  such that

$$\frac{\delta_0}{c} \leq \frac{|a_{k+1} - a_I|}{|a_k - a_I|} \leq \delta_0.$$

Let  $a'_0 = m_I$  and for each  $k \geq 1$  let  $a'_k \in (a_I, m_I)$  with  $a'_k = 2a_I - a_k$ . Similarly we obtain sequences  $\{b_k\}_{k \geq 0} \subset E$  and  $\{b'_k\}_{k \geq 0} \subset [m_I, b_I]$  for the point  $b_I$ . In the following, two intervals  $[a'_{k+1}, a'_k]$  and  $[a'_k, a'_{k-1}]$  are called *neighbor intervals*. Similarly,  $[a'_1, m_I]$  is a neighbor of  $[m_I, b'_1]$  and for each  $k \in \mathbb{N}$ ,  $[b'_{k-1}, b'_k]$  is a neighbor of  $[b'_k, b'_{k+1}]$ .

We define now  $f_I : \bar{I} \rightarrow \mathcal{B}_{n_I}(I)$ . Set  $f_I(m_I) = m_{\mathcal{B}(I)}$  and for each  $k \geq 1$ , define  $f_I(a'_k) \in [f(a_I), m_{\mathcal{B}(I)}]$  and  $f_I(b'_k) \in [f(b_I), m_{\mathcal{B}(I)}]$  by

$$\frac{|f_I(a'_k) - f(a_I)|}{|f(a_k) - f(a_I)|} = 1 = \frac{|f_I(b'_k) - f(b_I)|}{|f(b_k) - f(b_I)|}.$$

On each interval  $[a'_{k+1}, a'_k]$  or  $[b'_k, b'_{k+1}]$  we extend  $f_I$  linearly. It follows from the choice of  $\delta_0$  that  $f_I$  is a homeomorphism.

Suppose that  $J_1, J_2 \subset I$  are neighbor intervals. Then, there exists constant  $C > 1$  depending only on  $\eta$  and  $c$  such that

$$(4) \quad C^{-1} \leq |J_1|/|J_2| < C \text{ and } C^{-1} \leq \text{diam } f_I(J_1)/\text{diam } f_I(J_2) < C.$$

Thus, by Lemma 2.3, Remark 3.5 and the linearity of  $f_I$  on each  $J_i$  the following remark can be easily verified.

*Remark 3.9.* Suppose that  $J_1, J_2, J_3 \subset I$  are consecutive neighbor intervals. Then, there exists  $\eta_1$  depending only on  $\eta$  and  $c$  such that  $f_I|(J_1 \cup J_2 \cup J_3)$  is  $\eta_1$ -quasisymmetric.

Note that  $f_I|\{a'_k\}_{k \geq 0}$  is  $\eta_2$ -quasisymmetric for some  $\eta_2$  depending only on  $\eta$  and  $c$ . We show in the next lemma that  $f_I$  is quasisymmetric.

**Lemma 3.10.** *Let  $I$  be a component of  $\mathbb{R} \setminus E$ . There exists  $\eta'$  depending only on  $\eta$  and  $c$  such that  $f_I$  is  $\eta'$ -quasisymmetric.*

*Proof.* By Remark 3.9,  $f_I|[a'_1, b'_1]$  is quasisymmetric. We show that  $f_I|[a_I, a'_0]$  is quasisymmetric and similar arguments apply for  $f_I|[b'_0, b_I]$ . Then, by Lemma 2.3 and Remark 3.5,  $f_I$  is  $\eta'$ -quasisymmetric with  $\eta'$  depending only on  $\eta$  and  $c$ . Recall that  $f_I|\{a'_k\}_{k \geq 0}$  is  $\eta_2$ -quasisymmetric with  $\eta_2$  depending only on  $\eta$  and  $c$ .

To show that  $f_I|[a_I, a'_0]$  is quasisymmetric, we apply Lemma 2.3. Let  $x, y, z$  be in  $[a_I, a'_0]$ , with  $x$  being between  $y$  and  $z$ , and  $|x - y| \leq |x - z|$ . Suppose  $x \in [a'_k, a'_{k-1}]$ .

Assume first that  $y < x < z$ . If  $z \geq a'_{k-2}$  then  $|f_I(x) - f_I(y)| \leq |f_I(a'_{k-1}) - f_I(a_I)| \leq \eta_2(2)|f_I(a'_{k-1}) - f_I(a'_{k-2})| \leq \eta_2(2)|f_I(x) - f_I(z)|$ . If  $z \leq a'_{k-2}$  and

$y \geq a'_{k+1}$  then the quasimmetry follows from Remark 3.9. If  $z \leq a'_{k-2}$  and  $y \leq a'_{k+1}$  then  $|x-z| \geq |x-y| \geq C^{-1}|a'_{k-1}-a'_k|$  and by Remark 3.9,  $|f_I(x)-f_I(y)| \leq |f_I(a'_{k-1})-f_I(a_I)| \leq \eta_2(2)|f_I(a'_k)-f_I(a'_{k-1})| \leq \eta_2(2)(|f_I(x)-f_I(a'_k)|+|f_I(x)-f_I(a'_{k-1})|) \leq 2\eta_2(2)\eta_1(C)|f_I(x)-f_I(z)|$ .

Assume now that  $z < x < y$ . Then, there exists  $m_0 \in \mathbb{N}$  depending only on  $c$  and  $\eta$  such that  $y \leq a'_{k-m}$  for some  $0 \leq m \leq m_0$ . If  $z \geq a'_{k+1}$  then we obtain quasimmetry by applying Lemma 2.3 at most  $m_0$  times. If  $z \leq a'_{k+1}$ , then  $|f_I(x)-f_I(y)| \leq |f_I(a'_k)-f_I(a'_{k-m})| \leq \eta_2(m_0C^{m_0})|f_I(a'_k)-f_I(a'_{k+1})| \leq \eta_2(m_0C^{m_0})|f_I(x)-f_I(z)|$  where  $C$  is as in (4).  $\square$

#### 4. PROOF OF MAIN RESULTS

We show Theorem 1.1 in this section. The proof of Theorem 1.2 is given in Section 4.3 and is a minor modification of that of Theorem 1.1.

Let  $N = n + n_0 + 1$  be as in Section 3.2. Define  $F : \mathbb{R} \rightarrow \mathbb{R}^N$  with  $F|_E = f$  and  $F|_I = f_I$  whenever  $I$  is a component of  $\mathbb{R} \setminus E$ . We show in Section 4.2 that  $F$  satisfies (1) and then, Lemma 2.2 concludes the proof of Theorem 1.1.

To limit the use of constants we write in the following  $u \lesssim v$  (resp.  $u \simeq v$ ) when the ratio  $u/v$  is bounded above (resp. bounded above and below) by a positive constant depending at most on  $\eta$  and  $c$ .

**4.1. A form of monotonicity.** For the proof of the quasimmetry of  $F$  we show first that  $F$  satisfies the following form of monotonicity.

**Lemma 4.1.** *Suppose that  $x_1, x_2, x_3 \in \mathbb{R}$  with  $x_1 < x_2 < x_3$ . Then,*

$$|F(x_2) - F(x_1)| \vee |F(x_3) - F(x_2)| \lesssim |F(x_3) - F(x_1)|.$$

First we make an observation. Let  $x, y \in \mathbb{R}$  with  $x < y$  that are not on the same component of  $\mathbb{R} \setminus E$ . Denote by  $x', y'$  the minimum and maximum, respectively, of  $E \cap [x, y]$ . By Corollary 3.8 and the quasimmetry of  $f$ ,

$$(5) \quad |F(x) - F(y)| \simeq |F(x) - F(x')| + |F(x') - F(y')| + |F(y') - F(y)|.$$

*Proof of Lemma 4.1.* Let  $x_1, x_2, x_3 \in \mathbb{R}$  with  $x_1 < x_2 < x_3$ . We only show that  $|F(x_2) - F(x_1)| \lesssim |F(x_3) - F(x_1)|$ ; the inequality  $|F(x_2) - F(x_1)| \lesssim |F(x_3) - F(x_1)|$  is similar.

If all three of them are in  $E$  or in the same component  $I$  of  $\mathbb{R} \setminus E$  then the claim follows from the quasimmetry of  $f$  and  $f_I$ . Therefore, we may assume that at least one of the  $x_1, x_2, x_3$  is in  $\mathbb{R} \setminus E$ .

*Case 1.* Suppose that there exists a component  $I$  of  $\mathbb{R} \setminus E$  that contains exactly two of the  $x_1, x_2, x_3$ . Assume, for instance that  $x_1, x_2 \in I$  and  $x_3 \notin I$ ; the case  $x_2, x_3 \in I$  is similar. Let  $x'_2$  and  $x'_3$  be the minimum and maximum, respectively, of  $E \cap [x_2, x_3]$ . By (5) and the quasimmetry of  $F$  on  $I$ ,  $|F(x_3) - F(x_1)| \gtrsim |F(x'_2) - F(x_1)| \gtrsim |F(x_2) - F(x_1)|$ .

*Case 2.* Suppose that there is no component of  $\mathbb{R} \setminus E$  containing two points from  $x_1, x_2, x_3$ . Let  $x'_1$  and  $x'_2$  be the minimum and maximum, respectively, of  $E \cap [x_1, x_2]$  and  $x''_2, x'_3$  be the minimum and maximum, respectively, of  $E \cap [x_2, x_3]$ . Applying (5) on  $x_1, x_3$  and quasimmetry on  $x'_1, x''_2, x'_3$ ,  $|F(x_3) - F(x_1)| \gtrsim |F(x'_2) - F(x'_2)| + |F(x'_2) - F(x'_1)| + |F(x'_1) - F(x_1)|$ . Applying quasimmetry on  $x'_2, x_2, x''_2$  and then (5) on  $x_1, x_2$ ,  $|F(x_3) - F(x_1)| \gtrsim |F(x_2) - F(x'_2)| + |F(x'_2) - F(x'_1)| + |F(x'_1) - F(x_1)| \gtrsim |F(x_2) - F(x_1)|$ .  $\square$

**4.2. Proof of Theorem 1.1.** Let  $x, y, z \in \mathbb{R}$  such that  $|x - y| \leq |x - z|$ . By Lemma 4.1, we may assume that  $x$  is between  $y$  and  $z$ . Without loss of generality we assume that  $y < x < z$ .

Since  $F|_E$  is already quasisymmetric, we may assume that at least one of the  $x, y, z$  is in  $\mathbb{R} \setminus E$ . The proof is divided in four cases.

For the first case, we use the following lemma that can easily be verified.

**Lemma 4.2.** *Let  $I = (a, b)$  be a component of  $\mathbb{R} \setminus E$ ,  $x_1 \in I$  and  $x_2 \in E$ .*

*Suppose  $x_1 < x_2$ . If  $|x_2 - b| > (4c)^{-1}|x_1 - b|$  set  $x'_1 = b$ . If  $|x_2 - b| \leq (4c)^{-1}|x_1 - b|$  and  $x_1 \leq m_I$  set  $x'_1 = b_0$ . If  $|x_2 - b| \leq (4c)^{-1}|x_1 - b|$  and  $x_1 \in [b'_{n+1}, b'_n]$  set  $x'_1 = b_{n+1}$ . In each case,  $|x_2 - x'_1| \simeq |x_2 - x_1|$  and  $|F(x_2) - F(x'_1)| \simeq |F(x_2) - F(x_1)|$ .*

*If  $x_2 < x_1$  replace  $b, b_0, b_{n+1}$  by  $a, a_0, a_n$ , respectively, and define  $x'_1$  similarly. The claim of the lemma holds in this case as well.*

*Case 1.* Suppose that exactly one of the  $x, y, z$  is in  $\mathbb{R} \setminus E$ .

*Case 1.1.* Assume that  $y \in \mathbb{R} \setminus E$  and  $x, z \in E$ . Let  $y'$  be as in Lemma 4.2 for the pair  $x_1 = y, x_2 = x$ . Then,  $|y' - x| \simeq |y - x| \lesssim |x - z|$  and

$$|F(y) - F(x)| \simeq |F(y') - F(x)| \lesssim |F(x) - F(z)|.$$

*Case 1.2.* Assume that  $z \in \mathbb{R} \setminus E$  and  $x, y \in E$ . We work as in Case 1.1.

*Case 1.3.* Assume that  $x \in \mathbb{R} \setminus E$  and  $y, z \in E$ . Let  $x'$  be the point defined in Lemma 4.2 for the pair  $x_1 = x, x_2 = z$ . Then,  $|y - x'| = |y - x| + |x - x'| \lesssim |x - z| \simeq |x' - z|$  and by Lemma 4.1,

$$|F(x) - F(y)| \lesssim |F(x') - F(y)| \lesssim |F(x') - F(z)| \simeq |F(x) - F(z)|.$$

*Case 2.* Suppose that exactly two of the  $x, y, z$  are in the same component of  $\mathbb{R} \setminus E$  and the third point is in  $E$ .

*Case 2.1.* Assume that  $x, y$  are in a component  $(a, b)$  of  $\mathbb{R} \setminus E$  and  $z \in E$ .

If  $|x - b| > |b - z|$  set  $z' = b$ . Note that  $|x - z| \simeq |x - z'|$  and, by quasisymmetry of  $F|(a, b)$  and Lemma 4.1,

$$|F(x) - F(y)| \lesssim |F(x) - F(z')| \lesssim |F(x) - F(z)|.$$

If  $|x - b| \leq |b - z|$  then set  $x' = b$ . Note that  $|x - y| \leq |x' - y| \lesssim |x - z| \simeq |x' - z|$ . By Lemma 4.1 and Case 1 for  $y, x', z$ ,

$$|F(x) - F(y)| \lesssim |F(x') - F(y)| \lesssim |F(x') - F(z)| \lesssim |F(x) - F(z)|.$$

*Case 2.2.* Assume that  $x, z$  are in a component  $(a, b)$  of  $\mathbb{R} \setminus E$  and  $y \in E$ . If  $|y - a| \leq |x - a|$  set  $y' = a$  and if  $|y - a| > |x - a|$  then set  $x' = a$ . In each case we work as in Case 2.1.

For the next two cases we use the following lemma.

**Lemma 4.3.** *Let  $(a_1, b_1), (a_2, b_2)$  be two components of  $\mathbb{R} \setminus E$  with  $b_1 < a_2$  and  $x_1 \in (a_1, b_1), x_2 \in (a_2, b_2)$ .*

*If  $|a_1 - b_1| \leq |a_2 - b_2|$  set  $x'_1 = b_1$ . Then,  $|x_1 - x_2| \simeq |x'_1 - x_2|$  and  $|F(x_2) - F(x_1)| \simeq |F(x_2) - F(x'_1)|$ .*

*If  $|a_1 - b_1| > |a_2 - b_2|$  set  $x'_2 = a_2$ . Then,  $|x_1 - x_2| \simeq |x_1 - x'_2|$  and  $|F(x_2) - F(x_1)| \simeq |F(x'_2) - F(x_1)|$ .*

*Proof.* Assume that  $|a_1 - b_1| \leq |a_2 - b_2|$ ; the case  $|a_2 - b_2| \leq |a_1 - b_1|$  is similar. By Remark 2.6,  $|x_1 - x_2| \simeq |x'_1 - x_2|$ . Moreover, by Lemma 4.1,

$$\begin{aligned} |F(x_2) - F(x'_1)| &\lesssim |F(x_2) - F(x_1)| \leq |F(x_2) - F(x'_1)| + |F(x'_1) - F(a_1)| \\ &\lesssim |F(x_2) - F(x'_1)| + |F(x'_1) - F(a_2)| \lesssim |F(x_2) - F(x'_1)|. \quad \square \end{aligned}$$

*Case 3.* Suppose that exactly two of the  $x, y, z$  are in  $\mathbb{R} \setminus E$  but in different components.

*Case 3.1.* Assume that  $y \in (a_1, b_1)$ ,  $x \in (a_2, b_2)$  and  $z \in E$  where for each  $i = 1, 2$ ,  $(a_i, b_i)$  is a component of  $\mathbb{R} \setminus E$  and  $b_1 < a_2$ .

If  $|a_1 - b_1| \leq |a_2 - b_2|$  then, by Lemma 4.3, setting  $y' = b_1$ , we have  $|x - y'| \simeq |x - y|$ ,  $|F(x) - F(y')| \simeq |F(x) - F(y)|$ . Now apply Case 1 for the points  $y', x, z$ .

If  $|a_2 - b_2| < |a_1 - b_1|$  then, by Lemma 4.3, setting  $x' = a_2$ , we have  $|x' - y| \simeq |x - y|$  and  $|F(x') - F(y)| \simeq |F(x) - F(y)|$ . Moreover,  $|x - z| \leq |x' - z| = |x - z| + |x - x'| \leq |x - z| + |x - y| \leq 2|x - z|$ . Thus,  $|x - z| \simeq |x' - z|$  and applying Case 1 for the points  $x', x, z$ , we have  $|F(z) - F(x)| \simeq |F(z) - F(x')|$ . Now apply Case 1 for  $y, x', z$ .

*Case 3.2.* Assume that  $x \in (a_1, b_1)$ ,  $z \in (a_2, b_2)$  and  $y \in E$  where for each  $i = 1, 2$ ,  $(a_i, b_i)$  is a component of  $\mathbb{R} \setminus E$  and  $b_1 < a_2$ .

If  $|a_1 - b_1| \leq |a_2 - b_2|$  then,  $|x' - z| \simeq |x - z|$ ,  $|F(x') - F(z)| \simeq |F(x) - F(z)|$ ,  $|y - x| \lesssim |y - x'| \lesssim |x' - z|$ ,  $|F(y) - F(x)| \lesssim |F(y) - F(x')| \lesssim |F(x') - F(z)|$  and we apply Case 1 for  $y, x', z$ .

If  $|a_2 - b_2| < |a_1 - b_1|$  then set  $z' = a_2$  and work as in Case 3.1.

*Case 3.3.* Assume that  $y \in (a_1, b_1)$ ,  $z \in (a_2, b_2)$  and  $x \in E$  where for each  $i = 1, 2$ ,  $(a_i, b_i)$  is a component of  $\mathbb{R} \setminus E$  and  $b_1 < a_2$ .

If  $|a_1 - b_1| \leq |a_2 - b_2|$  then set  $y' = a_1$ . Since  $|x - z| \simeq |x - y| + |x - z| \gtrsim |b_1 - a_2|$  we have that  $|x - y'| \simeq |x - y|$ . Moreover, by Lemma 4.1,  $|F(x) - F(y)| \lesssim |F(x) - F(y')|$  and we apply Case 1 for  $y', x, z$ .

If  $|a_2 - b_2| < |a_1 - b_1|$  then set  $z' = b_2$ . As before,  $|x - z| \simeq |x - z'|$ . Furthermore,  $|F(x) - F(z')| \simeq |F(x) - F(a_2)|$  when  $|x - a_2| > |a_2 - z|$  and  $|F(x) - F(z')| \simeq |F(b_2) - F(a_2)|$  when  $|x - a_2| \leq |a_2 - z|$ . In either case,  $|F(x) - F(z)| \simeq |F(x) - F(z')|$  and we apply Case 1 for the points  $y, x, z'$ .

*Case 4.* Suppose that  $y, x, z \in \mathbb{R} \setminus E$ . By Lemma 3.10, we may assume that either  $y$  or  $z$  is not in the same component as  $x$ .

*Case 4.1.* Assume that  $y \in (a_1, b_1)$  and  $x \in (a_2, b_2)$  where  $(a_i, b_i)$  are components of  $\mathbb{R} \setminus E$  and  $b_1 < a_2$ .

If  $|b_1 - a_1| \leq |b_2 - a_2|$  then set  $y' = b_1$  and, by Lemma 4.3,  $|x - y| \simeq |x - y'|$  and  $|F(x) - F(y)| \simeq |F(x) - F(y')|$ . Apply now Case 2 or Case 3 for the points  $y', x, z$ .

If  $|b_2 - a_2| < |b_1 - a_1|$  then set  $x' = a_2$  and, by Lemma 4.3,  $|x - y| \simeq |x' - y|$  and  $|F(x) - F(y)| \simeq |F(x') - F(y)|$ . As in Case 3.1,  $|x - z| \simeq |x' - z|$  and applying Case 2 or Case 3 for the points  $x', x, z$  we conclude that  $|F(x) - F(x')| \lesssim |F(x) - F(z)|$  which implies  $|F(x) - F(z)| \simeq |F(x') - F(z)|$ . Now apply Case 2 or Case 3 on the points  $y, x', z$ .

*Case 4.2.* Assume that  $x \in (a_1, b_1)$ ,  $z \in (a_2, b_2)$  where  $(a_i, b_i)$  are components of  $\mathbb{R} \setminus E$  and  $b_1 < a_2$ .

If  $|b_2 - a_2| \leq |b_1 - a_1|$  then set  $z' = a_2$  and work as in Case 4.1.

If  $|b_1 - a_1| < |b_2 - a_2|$  then set  $x' = b_1$  and, by Lemma 4.1 and Lemma 4.3,  $|x' - y| = |x - y| + |x - x'| \lesssim |x - z| \simeq |x' - z|$ ,  $|F(x') - F(z)| \simeq |F(x) - F(z)|$  and  $|F(x) - F(y)| \lesssim |F(x') - F(y)|$ . Apply now Case 2 or Case 3 for the points  $y, x', z$ .

**4.3. Proof of Theorem 1.2.** By Corollary 2.5 we only need to show the sufficiency in Theorem 1.2. The proof is a mild modification of the proof of Theorem 1.1. We only outline the steps.

Let  $E \subset \mathbb{R}$  be an  $M$ -relatively connected set and let  $f : E \rightarrow \mathbb{R}$  be a monotone  $\eta$ -quasisymmetric mapping. As before, we may assume that  $E$  is a closed set that contains at least two points and  $f$  is power quasisymmetric. Moreover, we may assume that  $f$  is increasing.

*Step 1.* First, we reduce the proof to the case that  $E$  has no lower or upper bound, as in Section 2. This time, however, in Case 1 and Case 2 we define  $\tilde{f}(-a_n) = -a_n$ , where  $\{a_n\} \subset E$  is as in Section 2. By Lemma 3.1,  $\tilde{E}$  is a closed relatively connected set and  $\tilde{f} : \tilde{E} \rightarrow \mathbb{R}$  is an increasing quasisymmetric embedding.

*Step 2.* We reduce the proof to the case that  $E$  has no isolated points. If  $E$  has isolated points, then define  $\hat{E}$  and  $\hat{f}$  as in Section 3.1. Since  $f(E) \subset \mathbb{R}$ , then  $\hat{f} : E \rightarrow \mathbb{R}$  and  $\hat{f}$  is increasing. By Lemma 3.3,  $\hat{E}$  is a uniformly perfect closed set and  $\hat{f}$  is quasisymmetric.

*Step 3.* Let  $I = (a, b)$  be a component of  $\mathbb{R} \setminus E$ . The bridge  $\mathcal{B}(f(a), f(b))$  in this case is simply the interval  $[f(a), f(b)]$ . The mapping  $f_I$  is defined as in Section 3.3. The rest of the proof is similar to that of Theorem 1.2.

## 5. THE QUASISYMMETRIC EXTENSION PROPERTY IN HIGHER DIMENSIONS

This paper was motivated by the following question: given a uniformly perfect Cantor set  $\mathcal{C}$  in  $\mathbb{R}^n$  and a quasisymmetric mapping  $f : \mathcal{C} \rightarrow \mathbb{R}^n$  that admits a homeomorphic extension on  $\mathbb{R}^n$ , is it always possible to extend  $f$  quasisymmetrically in  $\mathbb{R}^n$ ? While Theorem 1.2 shows that the answer is yes when  $n = 1$ , this is not the case when  $n \geq 2$ . In fact we show a slightly stronger statement.

**Theorem 5.1.** *For any  $n \geq 2$ , there exists a compact, countable, relatively connected set  $E \subset \mathbb{R}^n$  and a bi-Lipschitz mapping  $f : E \rightarrow \mathbb{R}^n$  that admits a homeomorphic but no quasisymmetric extension on  $\mathbb{R}^n$ .*

Before describing the construction we recall a definition. A domain  $\Omega \subset \mathbb{R}^n$  is a  $C$ -John domain if there exists  $C \geq 1$  such that for any two points  $x, y \in \Omega$ , there is a path  $\gamma \subset \Omega$  joining  $x, y$  such that  $\text{dist}(z, \partial\Omega) \leq C^{-1} \min\{|x - z|, |y - z|\}$  for all  $z \in \gamma$ . In this case, the arc  $\gamma$  is called a  $C$ -John arc. It is a simple consequence of quasisymmetry that quasisymmetric images of John arcs are John arcs quantitatively.

Fix now an integer  $n \geq 2$  and define  $h : \mathbb{R}^{n-1} \times \mathbb{R} \rightarrow \mathbb{R}^n$  with  $h(v, t) = (v, 2 - t)$ . Set  $Q_0 = Q'_0 = [-1, 1]^{n-1} \times [-1, 1]$  and for each  $k \in \mathbb{N}$  set

$$Q_k = [-4^{-k}, 4^{-k}]^{n-1} \times [2^{-k}, 2^{1-k}],$$

$h_k = h|_{Q_k}$  and  $Q'_k = h(Q_k)$ . For  $k = 0$  we set  $h_0 = \text{Id}$ . Define

$$U = \text{int}(Q_0 \setminus \bigcup_{k \in \mathbb{N}} Q_k), \quad U' = \text{int}(Q'_0 \cup \bigcup_{k \in \mathbb{N}} Q'_k)$$

and  $X = \partial U$ ,  $X' = \partial U'$ . Note that  $U$  is a  $C$ -John domain for some  $C \geq 1$ .

For each integer  $m \geq 0$  let  $\zeta_m : \mathbb{R}^n \rightarrow \mathbb{R}^n$  be a similarity that maps  $[-2, 2]^n$  onto  $[\frac{1}{2}4^{-m}, 4^{-m}] \times [-4^{-m-1}, 4^{-m-1}]^{n-1}$ . For each  $m, k \geq 0$  let  $Q_{m,k}, Q'_{m,k}, U_m, U'_m, X_m$  and  $X'_m$  be the images of  $Q_k, Q'_k, U, U', X$  and  $X'$ , respectively, under  $\zeta_m$ . Note that each  $U_m$  is  $C$ -John domain.

For each  $m, k \geq 0$  let  $E_{m,k}$  be a finite set on  $\partial Q_{m,k} \cap X_m$  such that

$$(6) \quad \text{dist}(x, E_{m,k}) < 8^{-k-m} \text{ for all } x \in \partial Q_{m,k} \cap X_m.$$

Let  $P_m = \zeta_m(0, \dots, 0, 0)$ ,  $P_m^* = \zeta_m(0, \dots, 0, -1/2)$  and  $P = (0, \dots, 0)$ . Set

$$E = \{P\} \cup \{P_m, P_m^*\}_{m \geq 0} \cup \bigcup_{m, k \geq 0} E_{m,k}.$$

Clearly,  $E$  is compact and countable. Moreover, by choosing the sets  $E_{m,k}$  to be relatively connected, we may assume that  $E$  is relatively connected.

Define  $f : E \rightarrow \mathbb{R}^n$  with  $f(P) = P$ ,  $f(P_m^*) = P_m^*$ ,  $f(P_m) = \zeta_m(0, \dots, 0, 2)$  and

$$f|_{E_{m,k}} = \zeta_m \circ h_k \circ \zeta_m^{-1}|_{E_{m,k}}.$$

Denote  $E'_{m,k} = f(E_{m,k})$  and  $E' = f(E)$ . It is easy to show that  $f$  is bi-Lipschitz and can be extended to a homeomorphism of  $\mathbb{R}^n$ . Let  $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$  be such an extension of  $f$ . We briefly describe why  $F$  can not be quasimetric; the details are left to the reader.

Assume that  $F$  is  $\eta$ -quasimetric. Fix  $m \in \mathbb{N}$  to be chosen later. Let  $x \in U_m$  with  $\text{dist}(x, X_m) = \text{dist}(x, E_{m,k}) = 4^{-m}4^{-k}$ . By quasimetricity, (6) and the fact that  $F|_{E_{m,k}}$  is an isometry, its image  $x' = F(x)$  satisfies  $c_1 4^{-m}4^{-k} \leq \text{dist}(x', E'_{m,k}) \leq c_2 4^{-m}4^{-k}$  for some  $0 < c_1 < c_2$  depending on  $\eta$ . We claim that if  $m$  is chosen big enough,  $x' \in U'_m$ . Indeed, let  $\gamma$  be a  $C$ -John arc connecting  $x$  and  $P_m^*$  in  $\mathbb{R}^n \setminus E$ . If  $x' \in \mathbb{R}^n \setminus U'_m$  then there would be a point  $z \in F(\gamma) \cap X'_m$ . If  $z \in \partial Q'_{m,l}$  then  $\text{dist}(z, E'_{m,l}) \leq 8^{-m-l} < 2^{-m} \min\{|z - x'|, |z - P_m^*|\}$  which contradicts the quasimetricity of  $F$  if  $m$  is sufficiently big.

Let now  $m$  be chosen as above. Let  $x, y \in U_m$  with

$$\text{dist}(x, X_m) = \text{dist}(x, E_{m,k}) = \text{dist}(y, X_m) = \text{dist}(y, E_{m,k}) = 4^{-m}4^{-k}$$

and with  $|x - y| = 4^{-m}2^{-k-1}$  where  $k$  is chosen later. Let  $a, b$  be the points in  $E_{m,k}$  closest to  $x, y$  respectively. By quasimetricity of  $F$ , (6) and the fact that  $F|_{E_{m,k}}$  is an isometry, there exist constants  $C_1, C_2 > 0$  depending only on  $\eta$  such that the images  $x', y'$  of  $x, y$  satisfy  $\text{dist}(x', E'_{m,k}), \text{dist}(y', E'_{m,k}) \leq C_1 4^{-m}4^{-k}$  and  $|x' - y'| \geq C_2 4^{-m}2^{-k}$ . Let  $\sigma$  be a  $C$ -John arc joining  $x$  and  $y$  in  $\mathbb{R}^n \setminus E$ . As before, we can show that  $\sigma$  is contained in  $U_m$  and its image  $\sigma'$  is contained in  $U'_m$ . Let  $z \in \sigma' \cap Q'_{m,k}$  such that  $|z - x'| = |z - y'|$ . Then,  $\min\{|z - x'|, |z - y'|\} \geq \frac{1}{2}C_2 2^{-k}4^{-m}$  while  $\text{dist}(z, E'_{m,k}) \leq \frac{1}{2}4^{-k}4^{-m}$  and the John condition for  $\sigma'$  fails if  $k$  is sufficiently big. The latter contradicts the quasimetricity of  $F$ .

*Remark 5.2.* Let  $\mathcal{C}$  be the standard ternary Cantor set in  $[-\frac{1}{2}, \frac{1}{2}]$ . If in the above construction we replace the finite sets  $E_{m,k}$  by uniformly perfect Cantor sets  $\mathcal{C}_{m,k}$  satisfying (6), and the points  $P_m^*$  by sets  $\mathcal{C}_m = \zeta_m(\mathcal{C} \times \{(0, \dots, 0, \frac{1}{2})\})$ , then we obtain a Cantor set

$$\mathcal{C} = \{P\} \cup \{P_m\}_{m \geq 0} \cup \bigcup_{m \geq 0} \mathcal{C}_m \cup \bigcup_{m, k \geq 0} \mathcal{C}_{m,k},$$

for which the mapping  $f$  defined as above is bi-Lipschitz and admits a homeomorphic extension on  $\mathbb{R}^n$  but no quasimetric extension on  $\mathbb{R}^n$ .



## REFERENCES

1. Lars V. Ahlfors, *Quasiconformal reflections*, Acta Math. **109** (1963), 291–301. MR 0154978 (27 #4921)
2. ———, *Extension of quasiconformal mappings from two to three dimensions*, Proc. Nat. Acad. Sci. U.S.A. **51** (1964), 768–771. MR 0167617 (29 #4889)
3. A. Beurling and L. Ahlfors, *The boundary correspondence under quasiconformal mappings*, Acta Math. **96** (1956), 125–142. MR 0086869 (19,258c)
4. Lennart Carleson, *The extension problem for quasiconformal mappings*, Contributions to analysis (a collection of papers dedicated to Lipman Bers), Academic Press, New York, 1974, pp. 39–47. MR 0377046 (51 #13220)
5. G. David and S. Semmes, *Singular integrals and rectifiable sets in  $\mathbf{R}^n$ : Beyond Lipschitz graphs*, Astérisque (1991), no. 193, 152. MR 1113517 (92j:42016)
6. Juha Heinonen, *Lectures on analysis on metric spaces*, Universitext, Springer-Verlag, New York, 2001. MR 1800917 (2002c:30028)
7. Leonid V. Kovalev and Jani Onninen, *An  $N$ -dimensional version of the Beurling-Ahlfors extension*, Ann. Acad. Sci. Fenn. Math. **36** (2011), no. 1, 321–329. MR 2797699 (2012b:30049)
8. O. Lehto and K. I. Virtanen, *Quasiconformal mappings in the plane*, second ed., Springer-Verlag, New York-Heidelberg, 1973, Translated from the German by K. W. Lucas, Die Grundlehren der mathematischen Wissenschaften, Band 126. MR 0344463 (49 #9202)
9. Stephen Semmes, *Quasisymmetry, measure and a question of Heinonen*, Rev. Mat. Iberoamericana **12** (1996), no. 3, 727–781. MR 1435482 (97k:30029)
10. D. A. Trotsenko and J. Väisälä, *Upper sets and quasisymmetric maps*, Ann. Acad. Sci. Fenn. Math. **24** (1999), no. 2, 465–488. MR 1724387 (2000m:30032)
11. P. Tukia and J. Väisälä, *Quasiconformal extension from dimension  $n$  to  $n + 1$* , Ann. of Math. (2) **115** (1982), no. 2, 331–348. MR 647809 (84i:30030)
12. ———, *Extension of embeddings close to isometries or similarities*, Ann. Acad. Sci. Fenn. Ser. A I Math. **9** (1984), 153–175. MR 752401 (85i:30048)
13. Jeremy Tyson, *Quasiconformality and quasisymmetry in metric measure spaces*, Ann. Acad. Sci. Fenn. Math. **23** (1998), no. 2, 525–548. MR 1642158
14. Jussi Väisälä, *Bi-Lipschitz and quasisymmetric extension properties*, Ann. Acad. Sci. Fenn. Ser. A I Math. **11** (1986), no. 2, 239–274. MR 853960 (88b:54012)

DEPARTMENT OF MATHEMATICS AND STATISTICS, P.O. BOX 35 (MAD), FI-40014 UNIVERSITY OF JYVÄSKYLÄ, JYVÄSKYLÄ, FINLAND

*Current address:* Department of Mathematics and Statistics, P.O. Box 35 (MaD), FI-40014 University of Jyväskylä, Jyväskylä, Finland

*E-mail address:* `vyron.v.vellis@jyu.fi`