

**This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.**

**Author(s):** Polku, Hannele; Mikkola, Tuija; Gagné, Jean-Pierre; Rantakokko, Merja; Portegijs, Erja; Rantanen, Taina; Viljanen, Anne

**Title:** Perceived Benefit From Hearing Aid Use and Life-Space Mobility Among Community-Dwelling Older Adults

**Year:** 2018

**Version:** Accepted version (Final draft)

**Copyright:** © The Author(s) 2016.

**Rights:** In Copyright

**Rights url:** <http://rightsstatements.org/page/InC/1.0/?language=en>

**Please cite the original version:**

Polku, H., Mikkola, T., Gagné, J.-P., Rantakokko, M., Portegijs, E., Rantanen, T., & Viljanen, A. (2018). Perceived Benefit From Hearing Aid Use and Life-Space Mobility Among Community-Dwelling Older Adults. *Journal of Aging and Health*, 30(3), 408-420.  
<https://doi.org/10.1177/0898264316680435>

## **Perceived Benefit from Hearing Aid Use and Life-Space Mobility among Community-Dwelling Older Adults**

### **ABSTRACT**

**Objectives:** To examine the association between perceived benefit from hearing aid (HA) use and life-space mobility among older adults.

**Methods:** Cross-sectional analysis of 76-to-91-year-old community-dwelling adults (n=702). Data on perceived hearing with and without a HA were obtained via postal questionnaire and data on life-space mobility (Life-Space Assessment, range 0-120) via phone interview.

**Results:** Participants who perceived more benefit from HA use, had a better life-space mobility score (mean 65, SD 2.6) than participants who had less benefit from using a HA (55, SD 3.2). Participants who benefitted more from HA use did not differ from those who did not have a HA (63, SD 0.9) in their life-space mobility score.

**Discussion:** Perceived benefit from HA use is associated with higher life-space mobility among community-dwelling older adults. Future studies are needed to examine whether use of an appropriate HA promotes life-space mobility among those with difficulties in hearing.

Key words: Hearing, hearing aid, life-space, aging, older people

## **Perceived Benefit from Hearing Aid Use and Life-Space Mobility among Community-Dwelling Older Adults**

Difficulty in hearing is a common health concern among older adults affecting about two-thirds of people aged 70 years and older (Lin, Thorpe, Gordon-Salant et al., 2011). Hearing difficulties can hamper interaction with other people and safe orientation in the environment (Arlinger, 2003; Helvik, Jacobsen & Hallberg, 2006), thus complicating engagement in daily life situations (Wallhagen, Strawbridge, Shema et al., 2001). Hearing aid (HA) use may improve hearing performance and ease the perceived negative consequences of hearing difficulties (Kochkin & Rogin, 2000), thereby supporting participation in society. However, while most HA users are likely to perceive benefit from HA use (Wong, Hickson & McPherson, 2003), fewer than half of older adults with impaired hearing report using them (Bainbridge & Ramachandran, 2014; Hartley, Rochtchina, Newall et al., 2010; Lin, Thorpe, Gordon-Salant et al., 2011). Among the main reasons for not using a HA are that it is perceived to provide poor benefit, device is found uncomfortable (Hartley, Rochtchina, Newall et al., 2010; McCormack & Fortnum, 2013) or stigmatizing (Gagné, Southall & Jennings, 2011) or, a hearing problem is not perceived severe enough for needing a HA (Gopinath, Schneider, Hartley et al., 2011; Knudsen, Oberg, Nielsen et al., 2010).

In the present study, older adults' participation in everyday life situations in society is examined through life-space mobility. Life-space mobility refers to the size of the spatial area a person moves through in everyday life, taking into account the frequency of moving and the assistance needed for movement, thus describing total mobility (Baker, Bodner & Allman, 2003). An individual with larger life-space has more opportunities to engage in desired activities in society (Kono, Kai, Sakato et al., 2004), while restrictions in life-space mobility may indicate more limited possibilities for participation (Rosso, Taylor, Tabb et al.,

2013). Maintenance of life-space mobility may help to sustain better quality of life (Rantakokko, Portegijs, Viljanen et al., 2016).

Previously, only two studies have investigated the association between hearing difficulties and life-space mobility among older adults, and the results of these studies are somewhat inconsistent (Allman, Baker, Maisiak et al., 2004; Polku, Mikkola, Rantakokko et al., 2015). In both studies, self-reported hearing difficulties were found to be associated with poorer life-space mobility in community-dwelling older adults, but in the study of Allman et al. (2004) the association was not found statistically significant when adjusted for other health conditions (Allman, Baker, Maisiak et al., 2004). Comparison between these results is difficult because the studies differ in their sample sizes and the methodologies used to measure hearing difficulties. Moreover, neither study investigated the effects of HA use on life-space mobility. Thus, the aim of the present study was to explore whether hearing aid use, and more specifically perceived benefit from HA use, is associated with life-space mobility among community-dwelling older adults. In this study, perceived benefit from HA use refers to the difference between unaided and aided self-rated hearing ability. We compared life-space mobility of older adults who benefit less from HA to those who benefit more from HA. In addition, we explored whether these groups differ from those who do not use HA.

## **METHODS**

### **Study design and participants**

This study is based on cross-sectional analyses of the data obtained from the second follow-up of the “Life-Space Mobility in Old Age” (LISPE) project, which is a two-year prospective cohort study of community-dwelling Finnish older adults. The study design and methods, including non-respondent analysis, have been reported in detail elsewhere (Rantanen, Portegijs, Viljanen et al., 2012). Briefly, at baseline, a random sample of 2550 older

community-dwelling persons between 75 and 90 years of age was obtained from the national population register and used as the basis for recruitment. After an information letter, the persons were contacted over the telephone to enquire about their willingness to participate. Inclusion criteria were being community-dwelling in the study area, and able to communicate (i.e. able to understand the questions and provide answers to them). In total, 848 eligible persons participated in the baseline interviews during spring 2012. Of these, 761 participated in the two-year follow-up, which consisted of a telephone interview and a postal questionnaire. If the participant was unable to answer the questions over the phone due to hearing problems, the possibility to take part in a face-to-face interview in their homes (n=3) or, to answer the study questions via a postal questionnaire (n=10) was offered and implemented. The 2-year follow-up postal questionnaire, which included data on HA use, was returned by 712 participants (Rantakokko, Portegijs, Viljanen et al., 2016). The LISPE project was approved by the Ethical Committee of the University of Jyväskylä. Participants were informed about the project and signed a written informed consent prior to the study.

### ***Life-Space mobility***

Life-space mobility was assessed via telephone interview using the University of Alabama at Birmingham Life-Space Assessment (LSA) questionnaire (Baker, Bodner & Allman, 2003). The LSA is based on self-report and contains 15 items. It measures mobility through different life-space levels (bedroom, other rooms in the home, outside home, neighborhood, town, beyond town) either by walking or using other forms of transportation, such as driving a car or using public transportation, during the four weeks preceding the assessment. Participants were asked how many days a week they attained each life-space level and whether they needed help from another person or from assistive devices. The life-space mobility score ranging from 0 to 120 (higher scores indicate better life-space mobility) was used in the analyses (Baker, Bodner & Allman, 2003).

### *Perceived hearing ability and perceived benefit from hearing aid use*

In the postal questionnaire the participants were asked "How is your hearing?" Specifically, participants were asked to assess their hearing by choosing the number on a scale from 0 to 10 that best corresponded to their perceived hearing ability. Higher scores indicated poorer performance. Participants, who reported having a HA, gave two answers: one assessing their hearing with, and the other without, their HA (Figure 1).

How is your hearing?

**A. Without hearing aid**

very good very poor

0 1 2 3 4 5 6 7 8 9 10

**B. With hearing aid**

very good very poor

0 1 2 3 4 5 6 7 8 9 10

*Perceived benefit from hearing aid use* was quantified by subtracting the aided hearing score from the unaided hearing score, a greater difference indicating greater perceived benefit. The median value of perceived benefit from HA use in the present sample was three, and was used as a cut point to categorize HA users. For the analyses, participants were divided into three groups: 1) More HA benefit (difference of 3 or more points between non-aided and aided hearing ability), 2) Less HA benefit (difference less than 3 points), and 3) No HA-group (participants who reported not having a HA). HA users were also asked whether they used their HA daily (yes/no), whether they had a HA only in one ear or in both ears and, with an open-ended question, to estimate how many hours per day on average they used their HA.

### **Potential confounders**

Factors previously found to be associated both life-space mobility decline and hearing difficulties were considered as potential confounders. *Age and gender* were derived from the population register. *Number of years of education* was used as a socioeconomic indicator. *Self-reported physician-diagnosed chronic conditions* were obtained from a list of 22 chronic conditions and with an open-ended question. Chronic conditions that could theoretically be linked to hearing difficulties and life-space mobility, namely diabetes, rheumatoid arthritis, cardiac, circulatory and neurological diseases, were chosen as potential covariates. *Cognitive functioning* was assessed using the Mini-Mental State Examination (MMSE) (Folstein, Folstein & McHugh, 1975). Depressive symptoms were assessed with the 20-item Centre for Epidemiological Studies Depression Scale (CES-D) (Radloff, 1977).

### **Statistical analyses**

Of those participants who had returned the postal questionnaire (n=712), 584 reported that they did not have a HA and, 127 reported having a HA (data on HA use were missing for one participant). Of these participants who reported having a HA, information on the perceived benefit from HA use was available for 118 participants. These participants were included in the analyses and thus the final sample consisted of 702 participants. Life-space mobility data were available for 699 of these participants. As life-space mobility data were missing for less than 1% of the participants, missing values were not imputed.


Characteristics of the participants are described using means and standard deviations (SD) or percentages. One-way ANOVA was used for continuous variables and chi-square tests for categorical variables to compare characteristics between the groups categorized according to HA use and perceived benefit. Analysis of covariance (ANCOVA) was used to test the association between perceived benefit from HA use and life-space mobility. Levene's test of equality of error variances indicated that the error variance of the dependent variable

(life-space mobility score) was equal across groups. Skewness and kurtosis of the life-space score distribution were acceptable in each group.

Of the potential covariates, only age and perceived hearing ability without a HA differed statistically significantly between the groups and were thus included in the analysis. A value of  $p < .05$  was chosen as the level of statistical significance. IBM SPSS version 22.0 (SPSS Inc. Chicago, IL) was used for the analyses.


## RESULTS

The mean age of the participants was 82 years (SD 4.2) and 63% were women. The mean life-space mobility score was 63 (SD 21.7), ranging from 6 to 120. Of the participants, 18% reported having a HA. The sample characteristics categorized according to perceived benefit from HA use are presented in Table 1. Among the participants who perceived more benefit from HA, 81 % reported using the HA daily and on average 9 hours/day. Among those who perceived less benefit from HA, 49% reported daily use and on average 6 hours/day. Of the 127 participants who reported having a HA, 5 participants (4%) reported having a HA in both ears (binaural fitting). Distribution of the unaided and aided hearing scores by categories of hearing aid use are shown in Figure 2.



The estimated marginal means for life-space mobility by categories of HA use are presented in Figure 3. In the crude model, the difference in the life-space mobility score was not statistically significant between participants who perceived more benefit from HA (mean 60, 95%CI 55-64) and those who perceived less benefit from HA (mean 51, 95%CI 44-

57; mean difference 8.7, 95%CI for difference -1.3 to 18.6). Participants who perceived more benefit from HA did not differ from no-HA group (64, 95%CI 62-66) in their life-space mobility score (mean difference -4.4, 95%CI for difference -10.6 to 1.9). However, no-HA group had significantly better life-space mobility score than those participants who perceived less benefit from HA (mean difference 13.0 , 95%CI for difference 4.8 to 21.3).



After further adjustment for age and perceived hearing ability without a HA (Figure 3, Model 2), the difference in life-space mobility score between those who perceived more benefit from HA and those who perceived less benefit from HA maintained at the same level but became statistically significant (65, 95%CI 60-70 vs. 55, 95%CI 49-61, respectively, mean difference 10.1, 95%CI for difference 0.8 to 19.4, p=.028). Participant who perceived more benefit from HA and no-HA group (63, 95%CI 61-65) did not differ in their life-space mobility score after this adjustment (mean difference 2.1, 95%CI for difference -4.8 to 8.9). The post-adjustment difference between no-HA group and participants who perceived less

benefit from HA was borderline significant (mean difference 8.0, 95%CI for difference -0.03 to 16.1,  $p=.051$ ).

## DISCUSSION

Our study showed that persons who perceived more benefit from HA use had a significantly higher life space mobility score than those who perceived less benefit and that their life-space mobility score was comparable to that of those without a HA.

The present study contributes novel information on perceived hearing difficulties and life-space mobility. To our knowledge, only two earlier studies have investigated this association (Allman, Baker, Maisiak et al., 2004; Polku, Mikkola, Rantakokko et al., 2015). However, their findings were conflicting and neither of them focused on HA use. Allman et al. (2004) reported that after adjustment for other health conditions, hearing difficulty did not predict decline in life-space mobility over an 18-month follow-up. Another previous study (Polku, Mikkola, Rantakokko et al., 2015) showed that life-space mobility was lower among people with hearing difficulties but declined at a similar rate over the two-year follow-up period compared to the people who did not have hearing difficulties. The results of the present study provides further evidence for the association of poor hearing and low life-space mobility, as the perceived benefit from the HA use was associated with higher life-space mobility.

There may be several explanations for the current result. For older adults, the need to interact with others is one of the main reasons for going outside the home (Gardner, 2014). However, older people with hearing difficulties can experience communication breakdowns and avoid situations in which they might experience feelings of frustration or being left out (Arlinger, 2003; Gopinath, Hickson, Schneider et al., 2012). Self-stigma means that some older adults hold the same negative stereotypes about their hearing loss as those held by the society. Persons who exhibit self-stigma associated with hearing loss are more likely to reduce their activities outside the home in order to minimize the chances of revealing their hearing loss to others (Gagné, Southall & Jennings, 2011). Even though a person would not

withdraw entirely from situations that are challenging for hearing, they may involve in them less frequently (Mikkola, Portegijs, Rantakokko et al. 2015; Polku, Mikkola, Rantakokko et al., 2015). Perceived benefit from the HA may facilitate participation in out-of-home situations, thereby supporting life-space mobility.

Hearing difficulties can also impair a person's ability to reliably detect and recognize acoustic environmental cues when moving around (Arlinger, 2003). These cues may be essential for safe orientation in space, e.g. in localizing sounds of the motor vehicles. Previous studies have shown that hearing impairment is associated with walking difficulties (Viljanen, Kaprio, Pyykkö et al., 2009), poorer postural balance (Viljanen, Kaprio, Pyykkö et al., 2009b) and fear of falling (Viljanen, Kulmala, Rantakokko et al., 2012). These difficulties together with impaired ability to detect environmental acoustic information may compromise safe mobility. Use of a HA may be hypothesized to reduce fear of falling and related activity restriction leading to increased out-of-home participation.

Perceived benefit from HA use is also related to the interaction between the person and his/her environment (Noble & Hetú, 1994). Listening situations differ in their auditory demands (Wong, Hickson & McPherson, 2003) and, for example, having a conversation in a group requires a lot more auditory processing than a conversation in a quiet room (Williger & Lang, 2014). A person who participates in activities that require a lot of communication is more likely to experience hearing difficulties as disturbing than a person who prefers solitary activities (Chang, Ho & Chou, 2009). Thus he or she may also be more likely to perceive benefit from HA use than the person who participates in fewer activities outside the home. It is probable that individuals with moderate or severe hearing impairment are more likely to use a HA and therefore, also more likely to experience more benefit from the use of a hearing aid (Knudsen, Oberg, Nielsen et al., 2010; Williger & Lang, 2014). For people with mild perceived hearing loss, the perceived gain from using a HA may be less. This

was also supported by the current study as those who perceived benefit from HA use had lower self-rated hearing without a HA.

The strengths of this study include the use of large population-based sample, which increases the generalizability of the results. As our cohort included participants with and without hearing difficulties, the associations observed in this study likely represent those prevalent in a similar-aged general population. The study also has its limitations. Because of the cross-sectional design, the causality of the associations between the variables investigated cannot be determined. That is, we cannot say that perceiving benefit from the use of HA is a cause or a consequence of higher life-space mobility. Although we aimed to control for the effects of potential confounders, the possibility of residual confounding variables induced by unmeasured factors cannot be ruled out. Also, a few persons were excluded from the study because they were not able to communicate due to hearing problems. Therefore, it is likely that persons with severe hearing impairment were under-represented and selection bias cannot be completely ruled out. Had these persons participated, the associations observed between hearing aid benefit and life-space mobility might have been stronger. Further studies with larger number of hearing aid users are needed to confirm the associations reported in this study. We were not able to use audiological equipment to quantify hearing status or to validate the appropriateness of the hearing aids fitted. However, previous studies support the validity of self-reported measures of hearing impairment and HA benefit (Strawbridge, Wallhagen, Shema et al., 2000). One advantage of self-report questionnaire is that they capture information about the everyday activities and situations (Kiely, Gopinath, Mitchell et al., 2011). Moreover, subjective experiences, such as perceived benefit can be evaluated only via self-reports. In the present study no information was collected concerning the types of HAs used by the participants. Only 4% of the HA users reported having two HAs and therefore it was not possible to compare the participants who were fitted with one HA (monaural fitting) and those

fitted with two HAs (binaural fitting). Future investigations of the effects of the HA use on life-space mobility should take into account the role of these variables.

In conclusion, our results indicate that perceived benefit from HA use is associated with better life-space mobility. The current result serves as a justification for future studies examining whether use of a proper HA will promote life-space mobility and participation among those with difficulties in hearing.

**REFERENCES**

- Allman, R. M., Baker, P. S., Maisiak, R. M., Sims, R. V., & Roseman, J. M. (2004). Racial similarities and differences in predictors of mobility change over eighteen months. *Journal of General Internal Medicine*, 19, 1118-1126.
- Arlinger, S. (2003). Negative consequences of uncorrected hearing loss-a review. *International Journal of Audiology*, 42(2 Suppl.), 17-20.
- Bainbridge, K. E., & Ramachandran, V. (2014). Hearing aid use among older United States adults: the National Health and Nutrition Examination Survey, 2005–2006 and 2009–2010. *Ear and hearing*, 35, 289-294.
- Baker, P. S., Bodner, E. V., & Allman, R. M. (2003). Measuring life-space mobility in community-dwelling older adults. *Journal of the American Geriatrics Society*, 51, 1610-1614.
- Chang, H.P., Ho, C.Y. & Chou, P. (2009). The factors associated with a self-perceived hearing handicap in elderly people with hearing impairment--results from a community-based study. *Ear and Hearing*, 30, 576-583.
- Folstein, M. F., Folstein, S. E., & McHugh, P. R. (1975). "Mini-mental state": a practical method for grading the cognitive state of patients for the clinician. *Journal of Psychiatric Research*, 12, 189-198.
- Gagné, J. P., Southall, K., & Jennings, M. B. (2011). Stigma and self-stigma associated with acquired hearing loss in adults. *Hearing Review*, 18, 16-22.
- Gardner, P. (2014). The role of social engagement and identity in community mobility among older adults aging in place. *Disability and rehabilitation*, 36, 1249-1257.
- Gopinath, B., Hickson, L., Schneider, J., McMahon, C. M., Burlutsky, G., Leeder, S. R., & Mitchell, P. (2012). Hearing-impaired adults are at increased risk of experiencing

emotional distress and social engagement restrictions five years later. *Age and Ageing*, 41, 618-623.

Gopinath, B., Schneider, J., Hartley, D., Teber, E., McMahon, C. M., Leeder, S. R., & Mitchell, P. (2011). Incidence and predictors of hearing aid use and ownership among older adults with hearing loss. *Annals of Epidemiology*, 21, 497-506.

Hartley, D., Rochtchina, E., Newall, P., Golding, M., & Mitchell, P. (2010). Use of hearing aids and assistive listening devices in an older Australian population. *Journal of the American Academy of Audiology*, 21, 642-653.

Helvik, A. S., Jacobsen, G. W., & Hallberg, L. R. (2006). Life consequences of hearing loss in terms of activity limitation and participation restriction. *Scandinavian Journal of Disability Research*, 8, 53-66.

Kiely, K. M., Gopinath, B., Mitchell, P., Browning, C. J., & Anstey, K. J. (2011). Evaluating a dichotomized measure of self-reported hearing loss against gold standard audiometry: prevalence estimates and age bias in a pooled national data set. *Journal of Aging and Health*. doi: 10.1177/0898264311425088

Knudsen, L. V., Öberg, M., Nielsen, C., Naylor, G., & Kramer, S. E. (2010). Factors influencing help seeking, hearing aid uptake, hearing aid use and satisfaction with hearing aids: A review of the literature. *Trends in Amplification*, 14, 127-154.

Kochkin, S. & Rogin, C. (2000). Quantifying the obvious: the impact of hearing instruments on quality of life. *The Hearing Review*, 7, 6-34.

Kono, A., Kai, I., Sakato, C., & Rubenstein, L. Z. (2004). Frequency of going outdoors: a predictor of functional and psychosocial change among ambulatory frail elders living at home. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*, 59, 275-280.

- Lin, F. R., Thorpe, R., Gordon-Salant, S., & Ferrucci, L. (2011). Hearing loss prevalence and risk factors among older adults in the United States. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*, 66, 582-590.
- McCormack, A., & Fortnum, H. (2013). Why do people fitted with hearing aids not wear them? *International Journal of Audiology*, 52, 360-368.
- Mikkola, T.M., Portegijs, E., Rantakokko, M., Gagne, J-P., Rantanen, T., & Viljanen A.(2015). Association of self-reported hearing difficulty to objective and perceived participation outside the home in older community-dwelling adults. *Journal of Aging and Health*, 27, 103-122.
- Noble, W., & Héту, R. (1994). An Ecological Approach to Disability and Handicap in Relation to Impaired Hearing: Original Paper. *Audiology*, 33, 117-126.
- Polku, H., Mikkola, T. M., Rantakokko, M., Portegijs, E., Törmäkangas, T., Rantanen, T., & Viljanen, A. (2015). Self-reported hearing difficulties and changes in life-space mobility among community-dwelling older adults: a Two-year follow-Up study. *BMC geriatrics*. doi: 10.1186/s12877-015-0119-8.
- Radloff, L. S. (1977). The CES-D scale a self-report depression scale for research in the general population. *Applied Psychological Measurement*, 1, 385-401.
- Rantakokko, M., Portegijs, E., Viljanen, A., Iwarsson, S., Kauppinen, M., & Rantanen, T. (2016). Changes in life-space mobility and quality of life among community-dwelling older people: a 2-year follow-up study. *Quality of Life Research*, 25, 1189-1197.
- Rantanen, T., Portegijs, E., Viljanen, A., Eronen, J., Saajanaho, M., Tsai, L. T., ... & Rantakokko, M. (2012). Individual and environmental factors underlying life space of older people—study protocol and design of a cohort study on life-space mobility in old age (LISPE). *BMC Public Health*. doi: 10.1186/1471-2458-12-1018.

- Rosso, A. L., Taylor, J. A., Tabb, L. P., & Michael, Y. L. (2013). Mobility, disability, and social engagement in older adults. *Journal of Aging and Health, 25*, 617-637.
- Strawbridge, W. J., Wallhagen, M. I., Shema, S. J., & Kaplan, G. A. (2000). Negative consequences of hearing impairment in Old Age: a longitudinal analysis. *The Gerontologist, 40*, 320-326.
- Viljanen, A., Kulmala, J., Rantakokko, M., Koskenvuo, M., Kaprio, J., & Rantanen, T. (2012). Fear of falling and coexisting sensory difficulties as predictors of mobility decline in older women. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences, 67*, 1230-1237.
- Viljanen, A., Kaprio, J., Pyykkö, I., Sorri, M., Koskenvuo, M., & Rantanen, T. (2009). Hearing acuity as a predictor of walking difficulties in older women. *Journal of the American Geriatrics Society, 57*, 2282-2286.
- Viljanen, A., Kaprio, J., Pyykkö, I., Sorri, M., Pajala, S., Kauppinen, M., ... & Rantanen, T. (2009b). Hearing as a predictor of falls and postural balance in older female twins. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences, 64*, 312-317.
- Wallhagen, M. I., Strawbridge, W. J., Shema, S. J., Kurata, J., & Kaplan, G. A. (2001). Comparative impact of hearing and vision impairment on subsequent functioning. *Journal of the American Geriatrics Society, 49*, 1086-1092.
- Williger, B. & Lang, F.R. (2014). Managing age-related hearing loss: how to use hearing aids efficiently - a mini-review. *Gerontology, 60*, 440-447.
- Wong, L. L., Hickson, L., & McPherson, B. (2003). Hearing aid satisfaction: what does research from the past 20 years say? *Trends in Amplification, 7*, 117-161.

Table 1: Characteristics of the Participants Categorized by Hearing Aid Use (n=702)

	<b>No hearing aid (n=584)</b>	<b>Less hearing aid benefit (n=41)</b>	<b>More hearing aid benefit (n=77)</b>	
	<b>Mean (SD)</b>	<b>Mean (SD)</b>	<b>Mean (SD)</b>	<b>p<sup>a</sup></b>
Age	81.3 (4.0)	83.3 (4.3)	83.9 (4.4)	<.001
Education in years	9.7 (4.2)	9.4 (4.2)	9.7 (4.4)	.872
MMSE score (range 0-30)	26.0 (2.7)	25.5 (3.2)	26.2 (2.3)	.820
CES-D score (range 0-60)	13.0 (7.7)	15.5(6.7)	13.8 (6.8)	.105
Perceived hearing ability				
without a hearing aid (range 0-10) <sup>c</sup>	2.9 (2.3)	5.8 (1.9)	7.3 (1.7)	<.001
Perceived hearing ability with a hearing aid (range 0-10) <sup>c</sup>	-	5.4(2.4)	2.7(1.4)	<.001
	<b>% (n)</b>	<b>% (n)</b>	<b>% (n)</b>	<b>p<sup>b</sup></b>
Women	63 (366)	61 (25)	65 (50)	.899
Cardiac diseases	40(233)	54 (22)	44(34)	.191
Circulatory diseases	65 (378)	73(30)	56 (34)	.146
Diabetes	17 (101)	17 (7)	20 (15)	.891
Neurological diseases	6 (34)	7 (3)	9 (7)	.517
Rheumatoid arthritis	5(30)	2 (1)	4 (3)	.679

Notes: CES-D = Centre for Epidemiological Studies Depression Scale, MMSE= Mini-Mental State

Examination

a= one-way ANOVA

b= Chi-Square test

c= Higher scores indicate poorer performance

**Legends for figures**

1. Assessment of Perceived Hearing Ability and Perceived Benefit from Hearing Aid Use.
2. Distribution of the unaided and aided hearing scores by categories of hearing aid use.  
Higher scores indicate poorer performance (0= very good, 10= very poor). Ends of the whiskers indicate the minimum and maximum values.
3. Estimated marginal means and 95% confidence intervals for life-space mobility (range 0-120) by categories of hearing aid use. Crude and adjusted scores. Model 1 is adjusted for age and Model 2 is adjusted for age and perceived hearing ability without a hearing aid.