

This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Kontro, Titta; Sarna, Seppo; Kaprio, Jaakko; Kujala, Urho

Title: Mortality and health-related habits in 900 Finnish former elite athletes and their brothers

Year: 2018

Version: Accepted version (Final draft)

Copyright: © the Authors & BMJ, 2018.

Rights: In Copyright

Rights url: <http://rightsstatements.org/page/InC/1.0/?language=en>

Please cite the original version:

Kontro, T., Sarna, S., Kaprio, J., & Kujala, U. (2018). Mortality and health-related habits in 900 Finnish former elite athletes and their brothers. *British Journal of Sports Medicine*, 52(2), 89-95. <https://doi.org/10.1136/bjsports-2017-098206>

Mortality and health-related habits in Finnish former elite athletes and their brothers.

Titta K. Kontro,¹ Seppo Sarna,² Jaakko Kaprio,^{2,3} Urho M. Kujala¹

¹Faculty of Sport and Health Sciences, University of Jyväskylä, Jyväskylä, Finland

²Department of Public Health, University of Helsinki, Helsinki, Finland

³Institute for Molecular Medicine Finland (FIMM), University of Helsinki, Helsinki, Finland

Correspondence to: Titta K. Kontro, Faculty of Sport and Health Sciences,
University of Jyväskylä, P.O. BOX 35, 40014 Jyväskylä, Finland;
titta.k.kontro@student.jyu.fi

Key words: ALCOHOL, CHRONIC DISEASE, COHORT STUDY, FORMER ATHLETE,
MORTALITY, PHYSICAL ACTIVITY, SMOKING.

Word count: 2983 (with text modifications and new references 3085)

ABSTRACT

Background: There is conflicting evidence on the associations between participation in vigorous sports, health habits, familial factors and subsequent mortality. We investigated all-cause mortality and health-related behavior among former elite athletes and their brothers.

Methods: The mortality of Finnish male former elite athletes, who had represented Finland between 1920 and 1965 (n=900) and their age-matched brothers (n=900) was followed from the time when athlete started an elite athlete career until December 31, 2015. The age-adjusted Hazard ratios (HRs) were calculated by a paired Cox proportional hazards model. In 2001 surviving participants (n=199 athletes and n=199 age-matched brothers) reported their self-rated health, physical activity, alcohol consumption and smoking habits in the questionnaire.

Results: During the total follow-up period 1296 deaths (72% of the cohort) occurred. The age-adjusted HRs for all-cause mortality in former athletes was 0.75 (95% CI 0.65 to 0.87, $p<0.001$) compared to their age-matched brothers. Median age at death was 79.9 years for endurance, 75.9 years for mixed sports, and 72.2 years for power sports athletes, and 77.5, 73.7 and 72.2 years for their age-matched brothers respectively. In 2001, compared to their brothers, former athletes smoked less ($p<0.001$), were more physically active ($p<0.05$), and rated their health more often as very good ($p<0.05$).

Conclusions: Former elite athletes are more physically active, smoke less, have better self-rated health and live longer than their brothers. Genetic differences between athletes and brothers, aerobic training for endurance elite sports and a healthier lifestyle may all contribute to reduced mortality.

Key words: ALCOHOL, CHRONIC DISEASE, COHORT STUDY, FORMER ATHLETE, MORTALITY, PHYSICAL ACTIVITY, SMOKING.

1 **INTRODUCTION**

2

3 Observational studies show that certain lifestyle habits, such as hazardous alcohol drinking,
4 cigarette smoking, and low physical activity are important predictors of morbidity and
5 mortality.¹ Furthermore, self-rated health (SRH) is a predictor of mortality,² and physical
6 activity (PA) is related to better SRH.³⁻⁴

7

8 Regular PA is associated with better functioning and well-being especially at older age,⁵⁻⁶ a
9 lower risk for morbidity,⁷⁻⁸ and all-cause mortality.⁹⁻¹² Former endurance and mixed sports
10 athletes have lower mortality than the general population,¹³ but previous studies report
11 inconsistent results for power sports athletes.¹⁴ However, very intense patterns of exercise
12 may increase risk for cardiovascular morbidity and mortality compared to light to moderate
13 amounts of exercise.^{9,15}

14

15 There is no irrefutable evidence on a causal relationship between physical activity in
16 adulthood and mortality, either in animal experiments, or randomized controlled intervention
17 studies with healthy individuals.¹⁶ Cardiorespiratory fitness (CRF) is as a strong, independent
18 predictor of all-cause and disease-specific mortality¹⁷ and high level of CRF is associated
19 with a lower risk of all-cause and cardiovascular disease (CVD) mortality, in both rats¹⁸ and
20 humans.^{19,20} It has previously been widely reported that former athletes have lower morbidity
21 and mortality compared to general population,¹² which may be due to their genetic
22 background, increased PA levels or other better health habits.^{7-8,13} Former athletes are more
23 physically active and smoke less than controls,¹² while former²¹⁻²² or current athletes consume
24 more alcohol than non-athletes.²³

25

26 In contrast to our earlier studies on a cohort of former elite athletes before,^{11,13} but we use
27 brothers as control rather than unrelated healthy men. We selected brothers as controls
28 because they are genetically related and generally share the same family environment in
29 childhood. Thus, they may serve to control for experiences and exposures that cannot be
30 assessed when using unrelated controls. The aim of this study was to investigate whether all-
31 cause mortality and the development of behavioral and biological risk factors differ between
32 athletes and their brothers in later life. This unique study reveals novel data on the
33 associations between PA, familial factors, health-related behaviour, and mortality.

34

35 **METHODS**

36

37 **Participants**

38

39 An original cohort of former elite athletes (n=2657) was formed by identifying men who had
40 represented Finland between 1920 and 1965 at least once at the Olympic Games, European or
41 World championships, or inter-country competitions.¹² The athlete's brothers (n=2674) has
42 been collected from local parish registry data in early 1980s.

43

44 We excluded 219 athletes and 1774 brothers due to missing or uncertain information on date
45 of birth, date of death or death before 1936. We also excluded brothers who had died before
46 the time when the athlete started an elite athlete career. After including only athletes who had
47 brothers with complete data, the final study population (n=2755) consisted of 900 former
48 male athletes and their 1855 brothers (Table 1 and Figure 1). Former elite athletes (n=900)
49 and their individually matched brothers who were closest in age (n=900) were included in the
50 primary mortality analysis (paired Cox regression model). The former male athletes were

51 classified according to sports and sports-specific aerobic fitness characteristics into the
52 following three groups:¹² endurance sports (n=217), mixed sports (n=307) and power sports
53 (n=376) (individual sports are given in Table 1 and Figure 1). We also did a sensitivity
54 analysis based on former elite athletes (n=900) and all of their brothers (n=1855) (Figure 1).

55

56 Socioeconomic status was defined by five occupation-based social class groups: upper white
57 collar (executive and professionals), lower white collar workers (clerical work and
58 equivalent), skilled workers, unskilled workers and farmers.²⁴ Data were collected partly from
59 the Central Population Registry of Finland and partly from questionnaires based on the
60 occupation in which they had practiced the longest. The occupational group distribution of
61 athletes differed from that of brothers ($p < 0.001$, χ^2 -test) (Supplementary Table 1).

62

63 The data consisted of time and cause of death until December 31, 2015. The main outcome
64 variable was total all-cause mortality, and it was analyzed by using Cox regression model.
65 Secondly, the distributions of cause-specific deaths were analyzed by using cross-tabulations.
66 Mortality data were collected from the National Death Register of Statistics Finland (causes
67 of death; http://www.stat.fi/til/ksyyt/2005/ksyyt_2005_2006-10-31_luo_002.html).

68

69 **Comparison of athletes and brothers: the questionnaire study 2001**

70

71 In 2001 a postal questionnaire was sent to surviving elite athletes and brothers resident in
72 Finland eliciting information on health, lifestyle and sociodemographic characteristics. In
73 2001, 460 of 900 former athletes and 392 of their 900 age-matched brothers were alive; there
74 were 199 athletes who with their 199 age-matched brothers responded to the questionnaire.
75 We also did sensitivity analysis for the athletes (n=199) and all of their brothers (n=322)

76 (Figure 1). Furthermore, we used population controls (n=416), who were the unrelated
77 controls in the present cohort.²⁵

78

79 **Questionnaire-based covariates**

80

81 Body mass index (BMI) was calculated as self-reported weight/height² (kg/m²). The volume
82 of PA (MET-hours/day) was based on responses to three structured questions on intensity,
83 duration and frequency of activity using a previously validated method.^{10,26} The participants
84 were classified into tertiles of PA. Sedentary time was assessed by continuous variable called
85 as sitting time (hours/day).

86

87 Based on a detailed smoking history, we define smoking status as never, former, current or
88 occasional smokers. Those who have never smoked more than 100 cigarettes lifetime were
89 defined as never smokers. Former smokers had smoked more than 100 cigarettes in their
90 lifetime, who had smoked regularly, but had not smoked during the last month. Current
91 smokers were regular (daily or almost daily) smokers at the time of data collection.²⁷

92 Occasional smokers were men who smoked no more than 2 cigarettes in a week²⁸ or had last
93 smoked 2-30 days ago.²⁹ Among current or ex-smokers, nicotine dependence was assessed
94 using the Heaviness of Smoking Index (HSI), i.e. 2 items of the Fagerström Test for Nicotine
95 Dependence (FTND): the number of cigarettes per day and the time to the first cigarette in the
96 morning.³⁰ The HSI sum score ranges from 0 – 6, reflecting the degree of physical
97 dependence, and accounts for about 80 % of the variance in the FTND.^{25,30}

98

99 Alcohol use was assessed by questions on total alcohol consumption and heavy drinking
100 occasions (HDO). Alcohol consumption based on quantity-frequency measures of beverage

101 use was converted into grams of pure alcohol per month as previously reported.³¹ Drinkers
102 were classified as abstainers and light (3 or fewer drinks per week), moderate (more than 3
103 but not more than 14 drinks per week) and heavy drinkers (on average more than 2 drinks a
104 day).³² Heavy drinking occasions (HDO) were asked by question on the frequency of
105 drinking 5 or more alcoholic drinks on a single occasion in the past 30 days.³³ A HDO
106 variable had three categories: no HDO, HDO 1-2 times a month and HDO at least weekly.

107
108 Self-rated health (SRH) was measured by asking the participants to rate their own health
109 status at this moment on a previously validated² 5-point Likert scale: 1) very good, 2) quite
110 good, 3) average, 4) quite poor and 5) poor. Self-reported current symptoms and a history of
111 physician-diagnosed chronic diseases were grouped as dichotomous variables, such as any
112 chronic disease, hypertension with medication, angina pectoris, myocardial infarct, diabetes,
113 kidney disease, gastric ulcer, asthma and osteoarthritis. A history of any sports injury- and
114 Achilles tendinitis- or total rupture – was queried.

115
116 Activities of Daily Living (ADL) and Instrumental Activities of Daily Living (IADL) were
117 assessed by a total of nine items.³⁴ These national survey questions were modified from
118 classification of functional capacity.³⁵⁻³⁶

119
120 Self-rated mood was defined by two partial scales (6 depression items and 6 anxiety items) of
121 the short stress symptom survey extracted from the Brief Symptom Inventory-53 (BSI-53).³⁷

122
123 **Statistical analysis**

124

125 The descriptive data are presented as the mean and standard deviation (SD) or 95%
126 confidence intervals (CI) if distributed normally; otherwise the results are shown as the
127 median. The differences in the distributions of the categorical variables among different
128 sports groups and brothers were examined using cross-tabulations with the Chi-square (χ^2) -
129 test.

130

131 The follow-up of all-cause mortality of each athlete and brother started from the time when
132 the athlete was an elite athlete, and continued until the date of death (outcome event), date of
133 emigration (censoring event) or end of follow-up on December 31, 2015. A paired Cox
134 proportional hazards model was used to calculate age-adjusted hazard ratios (HRs) with 95%
135 CIs for all-cause mortality of former athletes relative to their brothers. Post hoc analysis
136 taking into account the number of comparisons was used to compare statistical differences
137 between specific sports groups. We also adjusted the mortality analysis for family clustering
138 in the subgroup analysis based on all questionnaire 2001 respondents. The Cox regression
139 assumptions were tested by using Schoenfeld residuals (ph-test in Stata) and by plotting “log-
140 log plots”.

141

142 The 2001 questionnaire data was mainly analyzed using non-parametric Wilcoxon`s signed
143 rank test (a matched-pair analysis) for ordinal or continuous variables and McNemar`s test (a
144 matched-pair analysis) for nominal variables to analyze the differences between each athlete
145 and the age-matched brother. The Chi-square (χ^2) -test and The Mann-Whitney-U-test were
146 used to compare differences between athletes and all their brothers, because some of the
147 variables were not normally distributed and variances were not equal. Homogeneity of
148 variances were assessed using Levene`s test and normality using Kolmogorov-Smirnov`s test
149 ($p<0.05$).

150 P-values <0.05 were considered statistically significant. Statistical analyses were performed
151 using SPSS statistical software (version 24.0 for Windows; SPSS Inc., Chicago, IL) and Stata
152 14.0 (Stata Corp, College Station, Texas, USA).

153

154 **RESULTS**

155

156 **Mortality data**

157

158 The final sample for mortality analysis included 900 former elite athletes and their 900 age-
159 matched brothers. Mean age at inclusion was 24.8 years and mean year of inclusion was
160 1943, and median follow-up time was 52.0 years (ranging from 0.1 to 83.4 years). Study
161 group characteristics are shown in Table 1. During the total follow-up period 1296 deaths
162 occurred; 72.1% (n=649) of the athletes and 71.9% (n=647) of the brothers died. Natural
163 causes accounted for death in 92.2% (n=1085) of the participants. The most common specific
164 cause of death was ischemic heart disease (IHD), which occurred in 35.5% (n=418) of all men
165 (34.3% (n=219) athletes, 36.9% (n=199) brothers) (Supplementary Table 2).

166

167 Median age at death among the former athletes (75.9 years, 95% CI 75.1 to 76.7) was
168 statistically significantly higher than among brothers (73.0 years, 95% CI 72.0 to 74.0).
169 Median age at death among endurance (79.9 years, 95%CI 79.1 to 80.7) and mixed sports
170 athletes (75.9 years, 95% CI 75.1 to 76.7) was statistically significantly higher than among
171 power sports athletes (74.1 years, 95% CI 73.3 to 75.0) (Table 2).

172

173 The age-adjusted hazard ratios (HRs) for all-cause mortality in former athletes was 0.75
174 (95% CI 0.65 to 0.87, p<0.001) compared to their age-matched brothers (Table 2). In a

175 subgroup analysis, compared to their age-matched brothers HR for all-cause mortality was
176 0.61 (95% CI 0.45 to 0.82, $p=0.001$) for endurance sports athletes, 0.85 (95% CI 0.65 to 1.10,
177 $p=0.220$) for mixed sports athletes and 0.78 (95% CI 0.62 to 0.98, $p=0.033$) for power sports
178 athletes (Table 2). In the sensitivity analysis the age-adjusted HRs for all-cause mortality in
179 former athletes ($n=900$) was 0.76 (95% CI 0.67 to 0.86, $p<0.001$) compared to all their
180 brothers ($n=1855$) (Supplementary Table 3).

181

182 We found that in the individual-based non-pairwise analysis the assumptions of Cox
183 regression model were violated suggesting a lower mortality of athletes at younger ages but
184 less so at older ages. We divided the follow up time into two periods: age at entry to median
185 age at death and median age at death to December 31, 2015. The age-adjusted pairwise HRs
186 for all-cause mortality in former athletes was 0.68 (95% CI 0.57 to 0.82, $p<0.001$) compared
187 to their age-matched brothers in the first follow-up period and 0.74 (95% CI 0.57 to 0.98,
188 $p=0.03$) in the second period. The results of the sensitivity analysis were consistent with the
189 results of age-matched pair analysis, but there was no difference in mortality during the later
190 period (Supplementary Table 4).

191

192 We also calculated adjusted HRs for all-cause mortality for all questionnaire 2001
193 respondents until 2015. There were no differences in HRs between groups (Supplementary
194 Table 5).

195

196 **Questionnaire data**

197

198 There were no statistically significant differences in anthropometric data between former
199 athletes and brothers (Table 3). Mean MET-hours/month in 2001 were significantly higher

200 among former athletes than their age-matched brothers (4.4. vs. 3.2 MET-hours/day)
201 ($p<0.05$). More athletes (11.2%) than brothers (2.1%) participated in vigorous PA ($p<0.05$).
202 The athletes sat less than brothers ($p<0.001$) (Table 3).

203

204 Former athletes smoked less than their age-matched brothers ($p<0.001$), and among smokers
205 the athletes were less nicotine-dependent. No significant differences in alcohol use were
206 observed (Table 3). Former athletes self-reported their health as very good much more
207 frequently (20.6%) than their brothers (8.6%) ($p<0.05$). There were no statistically significant
208 differences in specific chronic diseases, such as hypertension, heart disease, diabetes or
209 asthma between former athletes and their brothers. However, brothers reported more chronic
210 disease than former athletes ($p<0.05$). Former athletes had more self-reported physician-
211 diagnosed sports injuries and Achilles tendinitis than their brothers ($p<0.05$) (Table 4). There
212 were no differences in mobility, physical or psychosocial functioning of daily living between
213 former athletes and their brothers, except travelling by public transport. Mean of anxiety score
214 was higher among brothers than among former athletes ($p<0.05$), and brothers felt themselves
215 more restless and excited than former athletes ($p<0.05$) (Table 4).

216

217 We also did sensitivity analysis based on former elite athletes ($n=199$), all of their brothers
218 ($n=322$) and population control ($n=416$) (Supplementary Tables 6 and 7). The results were
219 consistent with the results of age-matched pair analysis.

220

221 **DISCUSSION**

222

223 **Principal findings**

224 Former elite athletes survived 2-3 years longer than their brothers, with the largest difference
225 seen for endurance athletes and none for power athletes. However, there was no difference in
226 the risk of all-cause mortality between athletes and all of their brothers at older age. Overall
227 SRH and health-related lifestyle habits of former elite athletes were better than those of their
228 brothers. However, there were no significant differences in mobility, physical or psychosocial
229 functioning of daily living between former athletes and their brothers who responded to the
230 2001 questionnaire. The brothers showed evidence for worse mood than athletes.

231

232 **Strengths and limitations of our study**

233

234 This unique long-term follow-up study revealed novel data on associations between health-
235 related behaviour, and mortality among individuals from the same family, who share both
236 genes and childhood environment. Valid death register data covering all participants was the
237 strength of this study. Although our questions on PA,²⁶ smoking³⁸ use of alcohol³⁹ have been
238 previously validated, self-reported data on health-related behaviors is only based on an age-
239 matched respondents subgroup from 2001. We have neither exact data on level of PA across
240 the lifespan nor health at younger ages among brothers. A partly insufficient data of SES or
241 causes of death among brothers could be considered as a limitation. However, athletes and
242 their brothers are from same family and differences in SES may be caused by athletic career.
243 We do not exactly know how well the results can be generalized to non-athletes, women or
244 today`s athletes from different sports groups or cultures, such as NFL players⁴⁰⁻⁴¹ or U.S
245 basketball players⁴² who seem to have high prevalence of clinically significant cardiac
246 abnormalities.

247

248 **Comparisons with other studies**

249 Observational studies in general have found that regular PA is associated with a lower risk for
250 all-cause mortality,⁹⁻¹¹ but high amounts of exercise among poorly trained individuals may
251 increase risk for CVD morbidity and mortality compared to light to moderate amounts of
252 exercise.^{9,15} Studies of elite athletes, i.e, among well-trained sportsmen do not support that.
253 The pattern of training and frequency of high-intensity exercise needs to be taken into
254 account.

255

256 Our study extends previous findings that former athletes, especially former endurance sports
257 athletes, have higher longevity and lower mortality than the general population.¹²⁻¹³ In
258 general, runners have a 25%-40% reduced risk of premature mortality and live approximately
259 3 years longer than non-runners.⁴³ Although high lifelong exercise volume among surviving
260 athletes is associated with coronary artery calcifications, the atherosclerotic plaques of the
261 most active athletes have a more benign composition.⁴⁴ The evidence is mostly based on
262 observational follow-up studies, one limitation of which is the difficulty of controlling for
263 genetic or other selection bias. There is no an irrefutable evidence on a causal relationship
264 between PA in adulthood and mortality. Based on previous studies on twins and former elite
265 athletes^{8,10} and both animal and human findings, it could be proposed that some of the
266 association between high PA^{16-17,19} and low mortality is explained by familial or genetic
267 factors. Despite this, we see a lower mortality in former athletes, especially at younger age
268 compared to their brothers knowing the limitation that some of the brothers may have had
269 health concerns at the beginning of our follow-up.

270

271 Our findings support previous studies that have shown athletes were more physically active
272 than controls.^{12,22} But we also observed that brothers were more physically active and went in

273 for more vigorous activities than population controls (Supplementary Table 6), suggesting a
274 familial contribution to PA that probably originated prior to the elite athletes career.

275

276 Though brothers smoked more than former athletes, there was no difference in alcohol use.
277 Former athletes are known to smoke less than controls,^{12,22} and we find that brothers also
278 smoked less than population (Supplementary Table 6). It has been found that persistent
279 physical inactivity in adolescence relates to adult smoking, even after familial factors are
280 taken into account.⁴⁵

281

282 There were better SRH and less chronic diseases among former athletes in accordance with
283 healthy life expectancy. However, brothers had less chronic diseases than population controls
284 (Supplementary Table 7). There was no difference in disability, possibly because those with
285 highest disability may not be able to respond to questionnaire. As earlier discussed both
286 athletes and brothers were physically active, which could partly explain that no differences
287 were found in functioning. Observational studies have found that regular PA is associated
288 with better functioning and well-being especially at older age,⁵⁻⁶ and a lower morbidity.⁷⁻⁸

289

290 **Future directions**

291

292 Our results are based on men who were former athletes 50 or more years ago. So, it is
293 important to investigate the associations between PA, familial factors, health-related
294 behaviour, and mortality among today's athletes, non-athletes or women. Traditional and
295 genetically informative designs are needed to investigate how elite performance related to
296 future health outcomes.

297

298 **CONCLUSIONS**

299

300 Former elite athletes have lower premature mortality than their brothers, and correspondingly
301 better SRH and health-related lifestyle habits. Former endurance sports athletes and their
302 brothers had reduced mortality compared to power sports athletes and their brothers. In
303 conclusion, our findings support previous findings for a role of genetic or childhood family
304 factors in determining high aerobic fitness and reduced mortality.

305

What are the new findings?

- Former elite athletes survived 2-3 years longer than their brothers.
- There was no difference in the risk of all-cause mortality between athletes and all of their brothers at older age.
- Overall self-rated health and health-related lifestyle habits of former elite athletes were better than those of their brothers.

306 **Contributors**

307 SS, JK and UMK collected the data. TKK and UMK analyzed the data. TKK drafted the
308 manuscript. All authors contributed to study design, and the revision of the manuscript, and
309 accepted the final version.

310 **Funding**

311 This study was funded by departmental funds of the University of Helsinki and University of
312 Jyväskylä. Jaakko Kaprio was supported by the Academy of Finland (grants # 265240,
313 &263278). There was no conflict of interests.

314 **Competing interest**

315 None.

316 **Ethical approval**

317 This study was conducted according to good clinical and scientific practice and the
318 Declaration of Helsinki. The authors declare that the results of this study are presented

319 clearly, honestly, and without fabrication, falsification or inappropriate data manipulation.
320 Approval for register data collection was given by the Ministry of Social Affairs and Health
321 in Finland, and Statistics Finland. All the participants gave informed consent by returning the
322 questionnaires, which were accompanied by a cover letter explaining the purpose of the study.

323 **Data sharing**

324 The former athletes are well known persons in Finnish society; hence the data cannot be
325 openly shared. Researchers are encouraged to contact the authors and we will make every
326 effort to accommodate additional analyses.

327

328

329

330

331

332

333

334

335

336

337

338

339

340

341

342

343

344

345

346

347

348

349

350

351

352 **REFERENCES**

353

- 354 1. Lim SS, Vos T, Flaxman AD, et al. A comparative risk assessment of burden of
355 disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions,
356 1990-2010: a systematic analysis for the Global Burden of Disease Study 2010.
357 *Lancet* 2012;380,2224-60.
- 358 2. Idler EL, Benyamini Y. Self-rated health and mortality: a review of 27 community
359 studies. *J Health Soc Behav* 1997;38:21 – 37.
- 360 3. Abu-Omar K, Rütten A, Robine JM. Self-rated health and physical activity in the
361 European Union. *Soz Präventiv Med* 2004;49(4):235-42.
- 362 4. Södergren M, Sundquist J, Johansson SE, Sundquist K. Physical activity, exercise and
363 self-rated health: a population-based study from Sweden. *BMC Public Health*
364 2008;8:352.
- 365 5. King AC, Pruitt LA, Phillips W, Oka R, et al.
366 Comparative effects of two physical activity programs on measured and perceived
367 physical functioning and other health-related quality of life outcomes in older adults. *J*
368 *Gerontol A Biol Sci Med Sci* 2000 Feb;55(2):M74-83.
- 369 6. Paterson DH, Warburton DE. Physical activity and functional limitations in older
370 adults: a systematic review related to Canada's Physical Activity Guidelines. *Int J*
371 *Behav Nutr Phys Act* 2010;11(7):38. doi: 10.1186/1479-5868-7-38.
- 372 7. Gajewski AK, Poznanska A. Mortality of top athletes, actors and clergy in Poland:
373 1924-2000 follow-up study of the long-term effect of physical activity. *Eur J*
374 *Epidemiol* 2008;23(5):335-40.
- 375 8. Kujala UM, Marti P, Kaprio J, et al. Occurrence of chronic disease in former top-level
376 athletes. Predominance of benefits, risks or selection effects? *Sports Med*
377 2003;33(8),553-61.
- 378 9. Eijsvogels TM, Molossi S, Lee DC, et al. Exercise at the extremes: The amount of
379 exercise to reduce cardiovascular events. *J Am Coll Cardiol* 2016;26:67(3):316-29.
- 380 10. Kujala UM, Kaprio J, Sarna S, et al. Relationship of leisure-time physical activity and
381 mortality. The Finnish twin cohort. *JAMA* 1998;279:440-4.

- 382 11. Samitz G, Egger M, Zwahlen M. Domains of physical activity and all-cause mortality:
383 systematic review and dose-response meta-analysis of cohort studies. *Int J Epidemiol*
384 2011;40(5):1382–400.
- 385 12. Sarna S, Sahi T, Koskenvuo M, et al. Increased life expectancy of world class male
386 athletes. *Med Sci Sports Exerc* 1993;25(2):237–44.
- 387 13. Kettunen JA, Kujala UM, Kaprio J, et al. All-cause and disease-specific mortality
388 among male, former elite athletes: an average 50-year follow-up. *Br J Sports Med*
389 2015;49(13):893-7.
- 390 14. Clarke PM, Walter SJ, Hayen A, et al. Survival of the fittest: retrospective cohort
391 study of the longevity of Olympic medallists in the modern era. *Br J Sports Med*
392 2015;49(13):898-902.
- 393 15. Kim JH, Baggish AL. Strenuous exercise and cardiovascular disease outcomes. *Curr*
394 *Atheroscler Rep* 2017;19(1):1-8. doi: 10.1007/s11883-017-0636-3
- 395 16. Karvinen S, Waller K, Silvennoinen M, et al. Physical activity in adulthood: genes
396 and mortality. *Sci Rep* 2015;5:18259. doi: 10.1038/srep18259.
- 397 17. Harber MP, Kaminsky LA, Arena R, et al. Impact of cardiorespiratory fitness on all-
398 cause and disease-specific mortality: Advances since 2009. *Prog Cardiovasc Dis*
399 2017;60:11-20.
- 400 18. Koch LG, Kemi OJ, Qi N, et al. Intrinsic aerobic capacity sets a divide for aging and
401 longevity. *Circ Res* 2011;109(10):1162–72.
- 402 19. Lavie CJ, Arena R, Swift DL, et al. Exercise and the cardiovascular system: clinical
403 science and cardiovascular outcomes. *Circ Res* 2015;117(2):207-19.
- 404 20. Kodama S, Saito K, Tanaka S, et al. Cardiorespiratory fitness as a quantitative
405 predictor of all-cause mortality and cardiovascular events in healthy men and women:
406 a meta-analysis. *JAMA* 2009;301(19):2024–35.
- 407 21. Batista CHX, Soares JM. Is athletic background associated with a future lower
408 prevalence of risk factors for chronic disease? *J Exerc Sci Fit* 2014;12(2):47-54.
- 409 22. Kontro TK, Sarna S, Kaprio J, Kujala UM. Use of alcohol and alcohol-related
410 morbidity in Finnish former elite athletes. *Med Sci Sports Exerc* 2017;49(3):492-9.

- 411 23. Martens MP, Dams-O'Connor K, Beck NC. A systematic review of college student-
412 athlete drinking: prevalence rates, sport-related factors, and interventions. *J Subst*
413 *Abuse Treat.* 2006;31(3):305-16.
- 414 24. Finland Central Statistical Office. Alphabetical list of occupations and classification of
415 social class. Helsinki: Finland Central Statistical Office; 1972. (Paper version
416 available from authors).
- 417 25. Bäckmand H, Kujala U, Sarna S, et al. Former athletes' health-related lifestyle
418 behaviours and self-rated health in late adulthood. *Int J Sports Med* 2010;31(10):751-
419 8.
- 420 26. Waller K, Kaprio J, Kujala UM. Associations between long-term physical activity,
421 waist circumference and weight gain: a 30-year longitudinal twin study. *Int J Obesity*
422 2008;32(2):353-61.
- 423 27. Kaprio J, Koskenvuo MA. prospective study of psychological and socioeconomic
424 characteristics, health behaviour and morbidity in cigarette smokers prior to quitting
425 compared to persistent smokers and non-smokers. *J Clin Epidemiol* 1988;41:139-50.
- 426 28. Paavola M, Vartiainen E, Puska P. Smoking cessation between teenage years and
427 adulthood. *Health Educ Res* 2001;16:49-57.
- 428 29. Luoto R, Uutela A, Puska P. Occasional smoking increases total and cardiovascular
429 mortality among men. *Nicotine Tob Res* 2000;2:133-9.
- 430 30. Haddock CK, Lando H, Klesges RC, et al. A study of the psychometric and predictive
431 properties of the Fagerström Test for Nicotine Dependence in a population of young
432 smokers. *Nicotine Tob Res* 1999;1:59 – 66.
- 433 31. Romanov K, Rose RJ, Kaprio J, et al. Self-reported alcohol use: a longitudinal study
434 of 12954 adults. *Alcohol Alcohol Suppl.* 1987;1:619-23.
- 435 32. Järvenpää T, Rinne O, Koskenvuo M, et al. Binge drinking in midlife and dementia
436 risk. *Epidemiol* 2005;16(6):766-71.
- 437 33. Substance Abuse and Mental Health Services Administration. Results from the 2004
438 National Survey on Drug Use and Health: National Findings. Available online
439 at:<http://www.oas.samhsa.gov/NSDUH/2k4NSDUH/2k4results/2k4results.htm#fig7.3>
440 (accessed 1 Nov 2016).

- 441 34. Aromaa A, Klaukka T, Impivaara O, et al. Aikuisten työ- ja toimintakyky sekä
442 avuntarve. In: Aromaa A, Heliövaara M, Impivaara O et al (Eds.), Health, Functional
443 Limitations and Need for Care in Finland. Basic Results from the Mini-Finland Health
444 Survey. Helsinki and Turku; Publications of the Social Insurance Institution of Finland
445 AL 32;1989:355-66,751.
- 446 35. Katz S, Ford AB, Moskowitz RW, et al. Studies of illness in the aged. The index of
447 ADL: a standardized measure of biological and psychosocial function. *J Am Med*
448 *Assoc* 1963;185:914–19.
- 449 36. Lawton MP, Brody EM. Assessment of older people: self-maintaining and
450 instrumental activities of daily living. *Gerontologist* 1969;9:179–86.
- 451 37. Derogatis LR, Melisaratos N. The brief symptom inventory: an introductory report.
452 *Psychol Med* 1983;13:595–605.
- 453 38. Vartiainen E, Seppälä T, Lillsunde P, et al. Validation of self-reported smoking by
454 serum nicotine measurement in a community-based study. *J Epidemiol Community*
455 *Health* 2002;56(3):167-70.
- 456 39. Poikolainen K. Underestimation of recalled alcohol intake in relation to actual
457 consumption. *Br J Addict.* 1985;80:215-6.
- 458 40. Helzberg JH, Waeckerle JF, Camilo J, et al. Comparison of cardiovascular and
459 metabolic risk factors in professional baseball players versus professional football
460 players. *Am J Cardiol.* 2010 (1);106(5):664-7.
- 461 41. Selden MA, Helzberg JH, Waeckerle JF, et al. Cardiometabolic abnormalities in
462 current National Football League players. *Am J Cardiol.* 2009 (1);103(7):969-71.
- 463 42. Harmon KG, Asif IM, Maleszewski JJ, et al. Incidence and etiology of sudden cardiac
464 arrest and death in high school athletes in the United States. *Mayo Clin*
465 *Proc.* 2016;91(11):1493-1502.
- 466 43. Lee DC, Brellenthin AG, Thompson PD, Sui X, Lee IM, Lavie CJ. Running as a key
467 lifestyle medicine for longevity. *Prog Cardiovasc Dis* 2017;60:45-55.
- 468 44. Aengevaeren VL, Mosterd A, Braber TL, et al. Relationship between lifelong exercise
469 volume and coronary atherosclerosis in athletes. *Circulation.* 2017;136(2):138-148.
- 470 45. Kujala UM, Kaprio J, Rose RJ. Physical activity in adolescence and smoking in young
471 adulthood: a prospective twin cohort study. *Addiction* 2007;102(7):1151-7.

472

473

List of Figures

- **Figure 1.** Study profile

List of Tables

- **Table 1.** Number of participants at entry to study and still alive in 2001, and December 31, 2015.
- **Table 2.** Age-adjusted hazard ratios (HRs) for all-cause deaths from time of entry to December 31, 2015 among former athletes (n=900) compared to age-matched brothers (n=900), mean (SD) and median age at entry, median and mean age at death.
- **Table 3.** Demographic and lifestyle characteristics (physical activity, alcohol use and smoking) among former athletes (n=199) and brothers (n=199).
- **Table 4.** Self-rated health, diseases, functioning of daily living, mobility, self-rated mood and working status among former athletes (n=199) and brothers (n=199).

Table 1. Number of participants at entry to study and still alive in 2001, and December 31, 2015.

Sports groups	Participants at entry	Participants at entry included in statistical analysis		Participants alive in 2001		Questionnaire responders in 2001 included in statistical analysis		Participants alive in 2015	
	N	N	Mean age (SD)	N	Mean age (SD)	N	Mean age (SD)	N	Mean age (SD)
1. Endurance	341	217*	27.0 (4.1)	104	71.8 (8.4)	44	69.1 (7.4)	54	81.6 (5.8)
2. Mixed sports	941	307 [†]	23.0 (3.0)	177	67.4 (8.7)	80	66.0 (8.1)	106	78.0 (5.8)
3. Power sports	1142	376 [‡]	24.5 (3.7)	174	68.1 (7.3)	75	67.0 (6.6)	91	79.5 (5.5)
All athletes	2424	900	24.6 (3.9)	455	68.7 (8.3)	199	67.1 (7.5)	251	79.3 (5.8)
Brothers	2673	900	25.1 (6.6)	393	67.4 (9.6)	199	65.9 (8.6)	253	79.1 (8.0)
Total	5097	1800	24.8 (5.4)	848	68.1 (9.0)	398	66.6 (8.0)	504	79.2 (7.0)

Data are numbers at entry included statistical analysis.

* Long distance running 95, middle distance running 45, cross-country skiing 77.

[†] Soccer 75, ice hockey 33, basketball 22, high jump 17, pole vault 23, long jump 14, triple jump 15, hurdling 36, short-distance running 46, decathlon 26.

[‡] Weight-lifting 43, Boxing 95, wrestling 155, shot-put 18, discus 15, javelin 32, hammer 19.

Table 2. Age-adjusted hazard ratios (HRs) for all-cause deaths from time of entry to December 31, 2015 among former athletes (n=900) compared to age-matched brothers (n=900), mean (SD), median age at entry, median and mean age at death.

Sports groups	Mean age (SD) at entry	Median age at entry	Median age at death (95%CI)	Mean age at death (95%CI)	Age-adjusted HR (95%CI)	P-value*
Endurance (n=217)	27.0(4.1)	26.0	79.9(79.1-80.7)	77.5(75.7-79.3)	0.607(0.448-0.823)	0.001
Mixed sports (n=307)	23.0(3.0)	23.0	75.9(75.1-76.7)	73.3(72.0-75.4)	0.848(0.652-1.104)	0.220
Power sports (n=376)	24.5(3.7)	24.0	74.1(73.3-75.0)	72.2(70.7-73.3)	0.779(0.619-0.980)	0.033
All athletes (n=900)	24.6(3.9)	24.0	75.9(75.1-76.7)	74.0(73.0-75.0)	0.753(0.648-0.875)	<0.001
Brothers (n=900)	25.1(6.6)	25.0	73.0(72.0-74.0)	69.9(68.7-71.0)	1.000	-

*p-values for statistical differences between sports groups and brothers by a paired Cox regression analysis.

Table 3. Demographic and lifestyle characteristics (physical activity, alcohol use and smoking) among former athletes (n=199) and brothers (n=199).

	Former athletes (N=199)	Brothers (N=199)	P-value
Age (mean (SD)), years*	67.1(7.5)	65.9(8.6)	0.007 ^a
Marital status %(n)*			0.035 ^b
Single	1.0(2)	4.5(9)	
Married	69.7(138)	77.4(154)	
Remarried	6.6(13)	3.5(7)	
Cohabiting	8.1(16)	3.5(7)	
Divorced	8.1(16)	4.5(9)	
Widowed	6.6(13)	6.5(13)	
Anthropometric data			
Height, m, mean (95% CI)	175.9(174.8-177.0)	175.1(174.2-176.0)	0.13 ^a
Weight, kg, mean(95% CI)	81.4(79.6-83.3)	81.4(79.7-83.1)	0.94 ^a
Mean BMI, kg/m ² (95% CI)	26.2(25.8-26.7)	26.5(26.0-27.0)	0.41 ^a
Physical activity			
Mean MET-hours/day (95% CI)*	4.4(3.7-5.1)	3.2(2.7-3.6)	0.012 ^a
Physical Activity tertiles [†] % (n)			0.734 ^a
Less active	34.7(69)	37.7(75)	
Moderately active	38.2(76)	29.1(58)	
Highly active	27.1(54)	33.2(66)	
Types of physical activity % (n)			
No physical activity	3.6(7)	3.1(6)	0.001 ^a
Walking	45.2(89)	63.4(127)	
Walking and jogging	32.0(63)	20.6(40)	
Jogging	8.1(16)	10.8(21)	
Brisk running	11.2 (22)	2.1 (4)	
Sedentary lifestyle			
Mean sitting time hours/day (95% CI)*	5.3(4.9-5.7)	6.3(5.8-6.7)	<0.001
Smoking			
<u>Smoking status,% (n)*</u>			2.57e ^{-04a}
Current smoker	7.1(14)	13.9(27)	
Occasional smoker	1.0 (2)	2.1(4)	
Ex-smoker	25.9(51)	39.2(76)	
Never smoker	66.0(130)	44.8(87)	
<u>Heaviness of smoking index (HSI, range 0-6), mean (SD)*</u>			0.004
among current / ex-smokers	1.5(1.6)/1.4(1.8)	2.0(1.7)/1.3(1.5)	

Alcohol

Total alcohol consumption g/month (mean, 95% CI)	184.9(136.2-233.6)	239.3(137.0-341.6)	0.58 ^a
Type of alcohol drinker %(n)			0.062 ^a
Abstainer or light drinker	67.3(134)	60.3(120)	
Moderate drinker	21.1(42)	25.6(51)	
Heavy drinker	11.6(23)	14.1(28)	
Heavy drinking occasions (HDO) % (n)	65.8(129)	65.3(126)	0.70 ^b
At least once a week	10.2(20)	16.1(31)	
1-2 times a month	16.3(32)	18.1(35)	
No HDO	73.5(144)	65.8(127)	

* p<0.05 for statistical differences between athletes and age-matched brothers by ^{a)} Wilcoxon`s signed rank -test or ^{b)} McNemar`s test.

† Overall physical activity (MET-hours/day): MET-hours/day ≤ 2.3 are less active, 2.3< MET-hours/day ≤ 4.5 are moderately active) and MET-hours/day > 4.5 are highly active.

Table 4. Self-rated health, diseases, functioning of daily living, mobility, self-rated mood and working status among former athletes (n=199) and brothers (n=199).

	Former athletes (n=199)	Brothers (n=199)	P-value
Self-rated health (SRH) %(n)*			3.3e ^{-04a}
Very good	20.6(41)	8.6(17)	
Quite good	48.7(97)	48.0(95)	
Average	26.1(52)	34.8(69)	
Quite poor or poor	4.5(9)	8.6(17)	
Diseases %(n)			
Self-reported any chronic disease*	43.2(83)	52.8(102)	0.039 ^b
<u>Physician-diagnosed chronic disease</u>			
Hypertension with medication	33.7(67)	40.1(79)	0.12 ^b
Angina pectoris	17.2(25)	17.4(26)	0.13 ^b
Myocardial infarct	10.6(21)	11.1(22)	0.86 ^b
Diabetes	3.5(7)	6.1(12)	0.16 ^b
Kidney disease	4.0(5)	2.7(9)	0.74 ^b
Ulcer	14.8(20)	17.9(26)	0.84 ^b
Asthma	10.4(14)	14.7(21)	0.59 ^b
Osteoarthritis	43.5(70)	37.0(57)	0.32 ^b
Sports injuries %(n)			
Any sports injury*	26.3(52)	16.3(32)	0.021 ^b
Achilles tendinitis*	21.1(37)	6.1(10)	0.003 ^b
Achilles total rupture	4.9(8)	1.9(3)	0.21 ^b
Mobility %(n)			0.26 ^a
No restrictions	84.3(161)	79.4(154)	
No ancillaries	14.7(28)	19.1(37)	
With ancillaries	1.8(2)	1.5(3)	
Functioning of daily living[†]			
Physical (score range 5-20), mean (95% CI)	5.5(5.3-5.8)	5.8(5.5-6.1)	0.211 ^a
Physical, good functionality %(n)	98.0(195)	97.5(194)	0.74 ^b
Physical, poor functionality %(n)	2.0(4)	2.5(5)	
Psychosocial (score range 4-16), mean (95% CI)	5.2(4.9-5.2)	5.3(5.0-5.5)	0.74 ^a
Psychosocial, good functionality% (n)	98.5(196)	98.5(194)	1.000 ^b
Psychosocial, poor functionality% (n)	1.5(3)	1.5(3)	
Self-rated mood (BSI-53)[‡]			
Depression, mean (95% CI)	2.1(1.7-2.5)	2.6(2.1-3.1)	0.28 ^a
Depression, % (n) with high score (≥8)	6.5(13)	8.5(17)	0.43 ^b
Anxiety, mean (95% CI)*	2.1(1.7-2.4)	2.7(2.3-3.1)	0.048 ^a
Anxiety, %(n), with high score (≥7)	7.7(15)	10.1(20)	0.59 ^b

Working status %(n)			0.67 ^a
Employed	15.6(31)	19.6(39)	
On old age pension	57.8(115)	50.8(101)	
On disability pension	13.1(26)	18.6(37)	
Unemployed	4.0(8)	3.0(6)	
Other	9.5(19)	8.0(16)	

* p<0.05 for statistical differences between athletes and age-matched brothers by ^{a)} Wilcoxon`s signed rank -test or ^{b)} McNemar`s test.

†Functioning of daily living (Activities of daily living scales (Mini-Finland Health Survey):

Physical Two groups: 0= good functioning, score 1 or 2, 1= poor functioning, score 3 or 4 on any item

Psychological Two groups: 0= good functioning, score 1 or 2, 1= poor functioning, score 3 or 4 on any item

‡Self-rated mood (Shortened anxiety and depression scales of BSI-53):

Depression Two groups: 0= no depressive, score ≤7.99, 1= depressive, score ≥8

Anxiety Two groups: 0= no anxious, score ≤6.99, 1= anxious, score ≥7

Self-rated mood:

The participants were asked to rate their mood at this moment on a 5-point Likert scale of 0-4: never (0), seldom (1), sometimes (2), often (3), and very often (4). So, the range of the sum scores was 0–24. The highest decile of the outcome variable (depression or anxiety) was considered an *abnormal outcome*, with others classified as not affected for the purposes of analyzing presence of a possible mood disorder. However, the scales are not diagnostic of clinical disease.