

**This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.**

**Author(s):** Baltazar, Margarida; Västfjäll, Daniel; Asutay, Erkin; Koppel, Lina; Saarikallio, Suvi

**Title:** Is it me or the music? : Stress reduction and the role of regulation strategies and music

**Year:** 2019

**Version:** Published version

**Copyright:** © The Author(s) 2019

**Rights:** CC BY-NC 4.0

**Rights url:** <https://creativecommons.org/licenses/by-nc/4.0/>

**Please cite the original version:**

Baltazar, M., Västfjäll, D., Asutay, E., Koppel, L., & Saarikallio, S. (2019). Is it me or the music? : Stress reduction and the role of regulation strategies and music. *Music and Science*, 2, 1-16.  
<https://doi.org/10.1177/2059204319844161>

# Is it me or the music? Stress reduction and the role of regulation strategies and music

Music & Science  
Volume 2: 1–16

© The Author(s) 2019

Article reuse guidelines:

sagepub.com/journals-permissions

DOI: 10.1177/2059204319844161

journals.sagepub.com/home/mns



Margarida Baltazar<sup>1</sup>, Daniel Västfjäll<sup>2,3,4,5</sup>, Erkin Asutay<sup>2,3</sup>,  
Lina Koppel<sup>2,4</sup> and Suvi Saarikallio<sup>1</sup>

## Abstract

Music is a common resource for the regulation of emotions, moods, and stress. This study aimed at determining the individual and relative impact on stress reduction of two of the main factors involved in musical affect regulation: regulation strategies and music itself. The current study took place in an experimental setting and followed a factorial within-subjects design. First, the participants ( $n = 34$ ) filled in an online survey where they identified their self-perceived “adequate”/“inadequate” music examples for the purpose of reducing stress and self-perceived “adequate”/“inadequate” strategies for the same purpose. In the lab they went through a stress induction procedure and then were instructed to calm down. They did so by listening to their “adequate”/“inadequate” music and employing the “adequate”/“inadequate” strategy, depending on the experimental condition. The primary outcome measure was self-reported tension, complemented by self-reported energy and valence, skin conductance levels (SCL), startle blink amplitudes, and risk aversion. The results showed that both music and strategy had a strong significant effect on the self-reported tension. Additionally, music had strong significant effects on energy, valence, SCL, and risk aversion. Pairwise comparisons revealed that the condition “adequate strategy-adequate music” was consistently more beneficial for stress reduction than condition “inadequate strategy-inadequate music”. However, it did not outperform all the other conditions, nor did the “inadequate strategy-inadequate music” underperform all the others. Moreover, close inspection of the results showed a larger impact of music on the short-term outcomes of self-regulation in comparison to strategy. These findings suggest that successful affective regulation depends on the adequacy of the chosen strategies and music, but that music is more determinant for the affective outcomes in the short term. The results are discussed considering previous research and the implications for the understanding of musical affect regulation are explored.

## Keywords

Affect regulation, emotion regulation, music, musical mechanisms, relaxation, self-chosen music, strategies, stress, tension

Submission date: 17 April 2018; Acceptance date: 18 March 2019

## Introduction

Stress is integral to our daily life. If a friend arrives at our house and complains about being stressed we can immediately see the whole picture: the stressor (intense traffic), the symptoms (faster breathing, increased perspiration), the changes in behaviour (t-shirt dressed inside out), and the solution (mellow jazz). Indeed, stress responses are quite common, as all sorts of events can trigger them. Stress can be managed in a variety of ways. One typical way of regulating stress in daily life is through music listening. Music has been widely shown to be an effective tool for reducing stress (Pelletier, 2004), and it has been shown to support a

<sup>1</sup> Department of Music, Art and Culture Studies, University of Jyväskylä, Jyväskylä, Finland

<sup>2</sup> JEDI Lab, Division of Economics, Department of Management and Engineering, Linköping University, Linköping, Sweden

<sup>3</sup> Division of Psychology, Department of Behavioural Sciences and Learning, Linköping University, Linköping, Sweden

<sup>4</sup> Center for Social and Affective Neuroscience, Department of Clinical and Experimental Medicine, Linköping University, Linköping, Sweden

<sup>5</sup> Decision Research, Eugene, OR, USA

### Corresponding author:

Margarida Baltazar, Department of Music, Art and Culture Studies, University of Jyväskylä, Musica, P.O. Box 35, Seminaarinkatu 15, Jyväskylä, FI 40014, Finland.

Email: margarida.baltazar@jyu.fi



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 License (<http://www.creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on

the SAGE and Open Access pages (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).

broad range of generally adaptive affect regulatory strategies such as distraction and reappraisal (Baltazar & Saarikallio, 2016, 2017). Yet, it remains an open question whether the efficacy of music as a stress regulation resource can be better explained by the presence of these efficacious regulatory strategies per se, or whether there is something more music-specific. For example, we do not know yet whether our friend eventually relaxed because the mellow jazz served as a distraction (*strategy*) or because its slow tempo helped to modulate our friend's breathing (*music itself*).

### ***Stress: Implications for affect, physiology, and cognition***

Stress is an interesting response to examine in light of its manifestations and its regulation: on the one hand, it plays an adaptive homeostatic function by responding to a demand for change and preparing the organism to act (Selye, 1936), and on the other hand the responses it triggers can be experienced as highly unpleasant and, in the long-term, translate into poorer well-being and health (McEwen & Stellar, 1993; Schneiderman, Ironson, & Siegel, 2005). The impact of stress on the individual's well-being and health depends greatly on their ability to self-regulate the high tension and negative valence concomitant to stress responses (Lazarus, 1974; Shallcross, Troy, & Mauss, 2015).

As stress is a multidimensional phenomenon, it is inadequate to study it using a single measure (Lazarus, 1990). In the current study, self-reported tension is our main focus due to its close link to the affective experience of stress. Importantly, we complement our approach with self-reported energy and valence and with measures from two other modalities: physiology and cognition.

The valence of the affective experience triggered by a stressor depends on the appraisals generated. The same stressful situation can be appraised, for example, as an opportunity to show one's skills (thus triggering positive affect) or as a threat to one's image (thus triggering negative affect). In this work, we focus on responses triggered by situations that are perceived as a threat of harm, overwhelming, or difficult to cope with, and that lead to reports of tension, anger, fear, or anxiety (Lazarus, 2000; Lazarus & Folkman, 1984; Smith, Haynes, Lazarus, & Pope, 1993).

The physiological responses are activated by the hypothalamus, which acts as a central control system (Tsigos & Chrousos, 2002) and manages two major systems that are closely involved in stress: the hypothalamus-pituitary-adrenal (HPA) axis and the sympatho-adrenomedullary (SAM) system. As the first signals are sent from the SAM system, catecholamines such as epinephrine are released. This leads to a number of physiological changes: pulse and blood pressure increase, breathing accelerates, skin conductance levels rise, and overall alertness is increased (Mandrick, Peysakhovich, Rémy, Lepron, &

Causse, 2016). In turn, the endocrine system of the HPA axis prompts the production of cortisol, which mobilizes the body's energy supply and increases the availability of energy (Munck, Guyre, & Holbrook, 1984).

These endocrine changes impact the prefrontal cortex (Arnsten, 2009) and thereby hinder core executive functions such as working memory and cognitive flexibility (Shields, Sazma, & Yonelinas, 2016). In this study, we explore the impact of acute stress on one cognitive process relying on executive functions: decision-making under risk. A large body of research has shown that people tend to be risk-seeking in the loss domain (i.e. they risk losing more for the chance of not losing anything) and risk-averse in the gain domain (i.e. they settle for smaller gains instead of taking risky—and larger—gains) (Kahneman, 2003; Kahneman & Tversky, 1979). However, this pattern seems to change under specific affective states, including acute stress (for a review, see Starcke & Brand, 2012). Acute stress has been found to decrease risk-taking in gain situations and increase in loss situations (Porcelli & Delgado, 2009). According to dual-process approaches (slow, rational process vs. fast, intuitive process; Epstein, Pacini, Denes-Raj, & Heier, 1996; Tversky & Kahneman, 1983), this can be explained by an overall shift from cognitive and slow approaches to intuitive and fast reactions, as demanded by the emergency created by the stressor (for a review, see Yu, 2016). Based on this, we will use risky decision-making as an implicit measure of stress (Västfjäll, 2010).

### ***Music listening: A resource for reducing stress***

Music listening is one of the resources most commonly used for the self-regulation of affect (Van Goethem & Sloboda, 2011). Affect self-regulation can be defined as any attempt at maintaining, increasing, decreasing, or replacing any affective state,<sup>1</sup> whether positive or negative (Gross, 2015; Gross & Thompson, 2007). In the present work we will focus on the down-regulation of stress responses, i.e. the goal of relaxing and decreasing tension.

When faced with high levels of stress in their lives, people tend to listen to more music (Getz, Marks, & Roy, 2014). This is arguably a wise strategy since music has been shown to have a range of beneficial effects on stress reduction (Chanda & Levitin, 2013; Pelletier, 2004). Several psychological, neurological, and biochemical pathways are involved in processing and responding to music (Fancourt, Ockelford, & Belai, 2014) and some of these capture quite well the stress reduction effect of music. For example, music listening can decrease cortisol levels (Khalifa, Dalla Bella, Roy, Peretz, & Lupien, 2003; Linneemann, Ditzgen, Strahler, Doerr, & Nater, 2015), increase serotonin levels and activate brain areas involved in reward (Evers & Suhr, 2000; Menon & Levitin, 2005), and enable a faster recovery of the autonomic nervous system after

exposure to a stressor (Chafin, Roy, Gerin, & Christenfeld, 2004; Thoma et al., 2013).

While there is ample evidence for the efficacy of music listening for stress reduction, there is currently little knowledge about the underlying factors behind this efficacy. One line of research has investigated how music is able to induce emotions in the listener (e.g. Juslin & Västfjäll, 2008; Scherer & Coutinho, 2013). The BRECVEMA framework posits a number of psychological mechanisms that mediate between music itself and the emotional reaction, ranging from brain stem reflexes to associative memories (Juslin & Västfjäll, 2008). The experimental testing of this framework has revealed that the activation of different mechanisms translates into different emotional reactions (e.g. Juslin, Barradas, & Eerola, 2015). These psychological mechanisms help to understand how, for example, music that is perceived as peaceful facilitates stress reduction (e.g. Sandstorm & Russo, 2010). However, many of these mechanisms are highly dependent on the individual's previous experiences and set of attitudes/preferences. That is the case of *visual imagery*, *evaluative conditioning*, *episodic memory*, and *aesthetic judgement* (Juslin, 2013). Scherer and Coutinho's (2013) process approach explicitly highlights individual differences as one factor in emotion induction (the others being structural features, performance, and context). The authors summarize the listener features into musical expertise (including explicit training and implicit cultural exposure) and stable dispositions (including age, gender, prior experience . . .). In line with this proposition, numerous empirical studies have observed that preference and familiarity of music are influential mediators for music's effect on affective states (Jiang, Rickson, & Jiang, 2016; Liljeström, Juslin, & Västfjäll, 2012, Pereira et al., 2011; Tan, Yowler, Super, & Fratianne, 2012).

Another line of research has focused on music as a tool for affect regulation, showing that music is often (and successfully) used as a way of reducing subjective feelings of stress (DeNora, 1999; Skånland, 2011; Ter Bogt, Vieno, Doornwaard, Pastore, & Eijnden, 2017). Music supports affect regulation through a wide range of regulatory strategies, i.e. behavioural or cognitive tools used to reach an affective goal (for a review, see Baltazar & Saarikallio, 2016). Regulation strategies operate through the management of diverse components: attention (e.g. *diversion*, *rumination*), cognition (e.g. *perspective taking*, *memory recall*), subjective experience (e.g. *inhibition of feelings*, *sensation seeking*), physiology (e.g. *energization*, *relaxation*), and behaviour (e.g. *venting*, *expressive suppression*) (Baltazar, 2018; Koole, 2009; Parkinson & Totterdell, 1999). Although the use of a certain strategy is not, per se, beneficial or harmful, its continuous use might reflect on the individual's adaptation to the environment (Marik & Stegemann, 2016). The tendency to often use certain strategies such as distraction or rumination—which are known to differ in terms of their adaptiveness—while regulating through music, has been able to (at least partially) explain the impact of music listening on health and well-being

outcomes (Carlson et al., 2015; Chin & Rickard, 2014; Garrido & Schubert, 2013). These studies support the claim that in order to understand music's influence, one needs to go beyond musical attributes such as genre, lyrics, and valence, and explore the underlying strategic uses.

In sum, two major factors are involved in musical affect regulation: music itself and regulation strategies. It can be argued that stress reduction—as a goal of affect regulation—occurs in interplay between personal strategies of music use and affordances of music (Baltazar, in press; Baltazar & Saarikallio, 2017). The effects of these two factors have been studied separately; thus, their relative and combined impact is yet to be revealed.

### Aim of current study

This study aimed at assessing the individual and combined impact of the strategies employed and the music listened to on stress responses.

Based on the literature suggesting that successful self-regulation of affective states depends on the use of adaptive and effective strategies (Marik & Stegemann, 2016) and that different pieces of music foster different emotional processes in the individual (DeNora, 2000), the following hypotheses were formulated:

H1: Efficacious music will lead to a larger reduction in acute stress than ineffacious music.

H2: Efficacious regulatory strategies will lead to a larger reduction in acute stress than ineffacious strategies.

H3: The combination of efficacious regulatory strategy and efficacious music will lead to the largest reduction in acute stress, while the combination of ineffacious strategy and ineffacious music will lead to the smallest reduction.

The independent variables (music listened to and strategies employed) will be treated as categorical variables with two categories (efficacious and ineffacious). Given the deep influence that preferences, individual differences, and sense of control have on the outcomes of music (e.g. Liljeström et al., 2012, Pereira et al., 2011) and employed strategies (e.g. Bonanno & Burton, 2013; Doré, Silvers, & Ochsner, 2016; Tamir & Ford, 2012), the selection of music and strategies and their categorization will be performed by the participants.<sup>2</sup>

## Method

This study follows an experimental approach in laboratorial context. Ecological validity and inter-subject variability were addressed by using strategies and music stimuli chosen by the participants themselves. Therefore, two stages were required: online survey and experiment.

### Phase 1: Recruitment and online survey


The participants were recruited from the population of registered students and staff of Linköping University, Sweden.

The individuals that agreed to participate filled out an online survey in Qualtrics (12.2016, Provo, UT) with background information, questions on musical self-regulation, and pre-screen questions. The pre-screen questions helped to exclude participants who were taking anxiolytics, antidepressants, or pain relievers. Eligible participants were invited to take part in the experiment at the laboratory.

**Selection of musical examples and regulation strategies.** The participants were instructed to imagine a situation of acute stress with a specific affect regulation goal (specific instructions in Appendix A). Then they identified a musical piece they liked and were familiar with and that, according to their experience, was efficacious for calming down. They could name up to three examples. For each song, the participants rated several aspects of the music (e.g. memories, rhythm, lyrics) in terms of their contribution for the affective outcome (continuous sliding scale from *no contribution* to *very strong contribution*).<sup>3</sup> Then, the opposite question was also asked, i.e. which would be the songs they liked and were familiar with, but that hindered the goal of calming down. Again, they rated the relevance of each aspect of the music. Then, participants were asked to identify the most efficacious and the least efficacious regulation strategy for relaxing while listening to music. The presented musical aspects and strategies were retrieved from a literature review (Baltazar & Saarikallio, 2016) and a study exploring the simultaneous use of both (Baltazar & Saarikallio, 2017). The regulation strategies presented were major strategies, i.e. they represented a type of approach and may include more than one specific strategy (e.g. *cognitive work* includes *reappraisal* and *perspective taking*, amongst others). The items can be found in Appendix A.

## Phase 2: Experiment

**Participants.** Thirty-five participants (14 female, 21 male), between the ages of 19 and 44 ( $M = 23.71$ ,  $SD = 4.91$ ) took part in the experiment. The sample can be characterized as highly engaged in music, since a clear majority (88%) reported listening to music six or more times per week and no participant reported listening less than once per week. On average, they reported purposely listening to music 2.75 hours per day. Seventy-six percent of the participants had at some point played an instrument or sung, and 38% still did. Most of the participants had musical education at school, with the most common being until high-school level (44%), followed by elementary school (32%). Only two participants had formal musical education outside of the school curriculum—one at the conservatory and the other at bachelor level. One female participant was excluded due to an error in the lab protocol. Participants were paid 100 SEK (approximately 9.60 EUR / 10.75 USD) as a show-up compensation plus or minus the amount from one randomly selected decision (decision task described further in the paper). All participants gave written, informed consent in



**Figure 1.** Study design with two blocks: stress induction and musical stress regulation. A and B mark the measurement points. Note. Measures used in A and B: self-reported affective experience (tension, energy, valence), SCLs, and startle blink amplitudes. In B, the decision-making task was repeated. Each participant went through this sequence four times (one for each condition).

accordance with the Declaration of Helsinki.<sup>4</sup> The study was approved by the Regional Ethics Board for East Gothland.

**Design and procedure.** The present study implemented a within-subjects  $2 \times 2$  factorial design, with complete counterbalancing of conditions.<sup>5</sup> The two factors were regulation strategy and musical piece, both with two levels—efficacious and inefficacious. Note that the efficacy of the two factors corresponded to the participants' self-perceived efficacy. The efficacious and inefficacious levels will be labelled as *adequate* and *inadequate* from now on. Four experimental conditions resulted from the combinations between *strategy* (adequate—AS, inadequate—IS) and *music* (adequate—AM, inadequate—IM): AS-AM, AS-IM, IS-AM, and IS-IM.

Before starting the experiment, a setting-up period took place. Then, as depicted in Figure 1, the experiment had two blocks (stress induction and musical stress regulation) with measurement points after each one (A and B). Each participant went through this sequence four times, once per experimental condition. During the measurement points A and B, the degree and direction of change in affective states were assessed. Next, we will briefly describe the three stages forming the experiment (setting up, stress induction, and musical stress regulation).

**Setting up.** Upon arrival, participants were welcomed and informed about the study. Instructions were given in relation to the self-report scales, the importance of employing the indicated strategies throughout the study even if they did not seem suitable for the moment, and the overall sequence of the experiment. The apparatus for physiological measurement were set up and, after that, the pain threshold for heat was determined following Perini, Bergstrand, and Morrison's (2013) procedure, described in Koppel et al. (2017). The heat-induced pain was delivered using a 3 x 3 cm thermode placed on the participants' non-

dominant dorsal forearm (Thermal Stimulator Probe, Q-sense, Medoc).

Participants were seated in a comfortable chair in front of a computer screen (Philips 241E), one loudspeaker (Genelec 8010 AP-6), and a standard mouse. The experiment ran through Presentation (Neurobehavioral Systems, version 18.0). Participants followed the instructions on the screen and completed the tasks with a mouse.

**Stress induction.** Acute stress was induced through a combination of heat-based pain and a fast-paced decision-making task. This block had the duration of 60 seconds, during which the temperature varied between the participant's pain threshold and 2°C below it. Simultaneously, the participant performed a monetary decision task in which they chose between a safe and a risky option with a time limit of 9 seconds for each decision. Stress responses are expected to be activated by this task due to the highly unpleasant component of pain and to the exhausting and demanding component of repeated choices under time-pressure (Vohs et al., 2008). The task was adapted from Koppel et al. (2017) and is explained under "Outcome measures", since the participants' choices were analysed as their risk-taking preferences.

**Musical stress regulation.** Before each musical stress regulation block, the participants received instructions regarding the regulation strategy that they should use. Based on the answers in the survey and depending on the experimental condition, they were given a card with the description of a strategy. Two musical pieces per participant (one per condition) were downloaded to a smartphone through the streaming service Spotify.<sup>6</sup> With the purpose of facilitating the affective impact of the musical pieces, the participants also received a card instructing them to focus on a certain musical aspect while listening to music (e.g. "Let yourself be immersed by the *lyrics* and *rhythm* of this song").<sup>7</sup> The musical pieces were played through headphones (Beyerdynamic DT700). The volume was adjusted to each participant's preference at the beginning of the experiment and was kept fixed throughout the experiment. Only a three-minute clip was played to the participants, unless the audio file was just slightly over three minutes, in which case we preferred not to cut the piece. The three-minute length was chosen based on the typical length of pop songs (longer songs could be cut) and on the need to keep each section of the experiment short enough to accommodate several repetitions. Participants were informed that they would have three minutes of music listening, during which they should try to calm down by using the given strategy and focusing on the given musical aspects. The songs listened by the participants and the frequencies of the strategies chosen as *adequate* or *inadequate* can be found in Appendix B.

## Outcome measures

The outcome measures covered different modalities (subjective experience, physiology, and cognition) to better portray the changes in affective states. The experiential dimension was measured through self-reported affect. Self-reports were collected through three continuous visual analogue scales (VAS): *relaxed – tense* (tension arousal), *negative – positive* (valence), and *drowsy – alert* (energy arousal). Additional synonyms to the poles of the scales, as used in Schimmack & Grob (2000), were given before the start of the experiment.

The psychophysiological measures included electrodermal activity (EDA) and facial electromyography (EMG). EDA was registered in terms of skin conductance levels (SCLs). SCLs respond to arousal and consequently increase when experiencing acute stress responses (Witvliet & Vrana, 1995) and when exposed to emotionally arousing music (Khalfa, Peretz, Blondin, & Manon, 2002). The facial EMG registered the startle blink magnitude, which has been shown to increase when negatively valenced and high arousal affect is induced (Lang, Bradley, & Cuthbert, 1990). For recording EDA, surface Ag/AgCl electrodes (8 mm in diameter) were attached to medial phalanges of the index and middle fingers of the participants' non-dominant hand. As for startle responses, surface Ag/AgCl electrodes (4 mm in diameter) were placed on the right obicularis oculi muscle. The startle responses were elicited by acoustic probes (40 ms long white noise played at 100dB), which were presented six times. The recording of the EMG started with a silent period, during which the participants were instructed to keep their eyes open and fixate on the cross displayed at the centre of the screen. The first auditory probe was presented 10 to 15 seconds after the beginning of the startle period. Intervals between two consecutive startle probes were randomly set between 8 to 15 seconds. Both EDA and EMG were recorded through a BIOPAC MP150 system (amplifiers GSR100C and EMG100C, respectively). The data collection was controlled by AcqKnowledge software (version 4.1), which amplified and sampled the physiological signals at a rate of 1000 Hz.

The cognitive dimension of acute stress was approached through the choice patterns in risky gains and losses. The decision-making task was one element of the stress induction procedure, but since this task was repeated after the musical stress regulation block, the proportions of risky choices were kept as a self-regulation outcome (Koppel et al. 2017; Koppel et al., submitted). During the task, the participants made four choices regarding two types of decisions: risky gains and risky losses. The trials were time-locked to 9 seconds and were presented in random order. In the risky gains trials, participants had to choose whether they preferred to receive a smaller but safe amount of money (with 100% probability) or a larger but risky sum in a coin toss (with 50% probability). In the risky losses trials, participants chose between losing with certainty a

smaller sum of money (with 100% probability) and losing a larger sum of money in a coin toss (with 50% probability). The proportion of times the participants chose to toss the coin instead of accepting the secure (but less advantageous) gain or loss was used as a measure of risk-taking. Participants were informed beforehand that one of their decisions would be randomly picked at the end of the experiment to count for their final payment (either adding to or subtracting from the base amount).

Amongst these measures, the self-reported *tension arousal* was our primary outcome as it has the closest link to the experience of stress, while the other scales and measures were considered complementary information regarding the affective reactions. Overall, reduction in acute stress was operationalized as lower values in the tension self-report scale, lower skin conductance levels, smaller startle blink amplitudes, higher risk-taking in gains, and lower risk-taking in losses.

### Data analyses

Before the statistical analyses, we proceeded to some preliminary data treatment. The data from the self-reports was rescaled to the range of -1 (left pole of the VAS) to +1 (right pole of the VAS). Both EDA and EMG signals were inspected individually for possible artifacts. EMG signals were band-pass filtered from 20 to 480 Hz. The filtered signal was full-wave rectified and low-pass filtered at 40 Hz for smoothing. Startle responses were scored as the peak response within a 20–200 ms window following startle probe onset if the peak exceeded three times the mean activity during the 50 ms preceding the startle probe onset. Response amplitudes were calculated as the average startle response over the six probe presentations during each affect assessment session. Only the participants who reacted at least once during each trial were included in the analysis of EMG data ( $n = 23$ ). The raw EDA signals were resampled at 10 Hz and low-pass filtered at 1 Hz. To compute skin conductance levels (SCLs), we applied a 5-second moving average filter. Then the signal was averaged into time blocks. Each time block consisted of the mean activity for 10 seconds while at rest, between the self-report of affect and the startle blink test. The decision-making data was computed as the proportion of risky choices in the gain and loss trials.

The changes in the outcome measures were calculated as the difference between the levels after stress induction and the levels after musical stress regulation (points A and B in Figure 1). Following the recommendations of Braithwaite, Watson, Jones, and Rowe (2015), the physiological data were standardized after calculating the individual changes between measurement points.

Affective experience measured by self-report was a multidimensional variable following a normal distribution. Physiological and decision-making data, on the other hand, did not follow a normal distribution. Consequently,

factorial effects of *strategy* and *music* on the self-reported affect were investigated through a two-way repeated-measures MANOVA. The multivariate effects were tested by means of Wilks' lambda, after which univariate  $t$  tests were performed. The effects of the experimental factors on physiological and decision-making data were investigated by calculating first the marginal means for each factor and then analysing through Friedman and Wilcoxon signed rank tests. The planned pairwise comparisons between conditions were conducted through  $t$  tests for self-report and Wilcoxon signed ranks tests for the other variables.


Given the sensitivity of parametric tests to outliers, participants exhibiting responses in the VAS above or below 1.5 times the interquartile range were excluded from the analyses on that variable. After the exclusion of outliers, the Shapiro–Wilk and Levene's tests failed to reject the null hypotheses that the self-report data have equal variances and belong to a population with normal distribution.<sup>8</sup>

Relevant effects were examined through statistical significance testing ( $\alpha = .05$ ) and estimation of effect size (ES): Cohen's  $d$  for paired  $t$  tests (Cohen, 1988; Dunlap, Cortina, Vaslow, & Burke, 1996), partial eta squared— $\eta_p^2$ —for ANOVA and MANOVA (Cohen, 1977; Pallant, 2007), Kendall's  $W$  for Friedman tests (Kendall, 1938), and correlation— $r$ —for Wilcoxon tests (Rosenthal, 1994). Cohen's (1988) guidelines were used for the interpretation of effect sizes;  $\eta_p^2$  follows the scale .01 = small, .06 = moderate, and .14 = large,  $d$  follows the scale .2 = small, .5 = moderate, and .6 = large, while both  $r$  and  $W$  follow the scale used for correlations: .1 = small, .3 = moderate, and .5 = large.

## Results

### Stress manipulation check

In order to estimate the success of the stress induction, the scores obtained before and after the induction block throughout the four conditions were compared. Separate one-way ANOVAs were conducted on self-report ratings and statistically significant differences were found on the three scales: tension increased—*pre-stress*:  $M = -0.25$ ,  $SD = 0.38$ , *post-stress*:  $M = -0.15$ ,  $SD = 0.31$ ,  $F(1, 238) = 20.83$ ,  $p < .001$ ,  $\eta_p^2 = .08$ , energy increased—*pre-stress*:  $M = 0.11$ ,  $SD = 0.41$ , *post-stress*:  $M = 0.23$ ,  $SD = 0.30$ ,  $F(1, 238) = 7.10$ ,  $p = .008$ ,  $\eta_p^2 = .029$ , and valence became more negative—*pre-stress*:  $M = 0.38$ ,  $SD = 0.33$ , *post-stress*:  $M = 0.24$ ,  $SD = 0.29$ ,  $F(1, 246) = 12.52$ ,  $p < .001$ ,  $\eta_p^2 = .048$ . The SCLs were submitted to a Wilcoxon signed ranks test, which revealed higher levels after stress induction, *pre-stress*:  $M = 9.54$ ,  $SD = 4.31$ , *Mean rank* = 60.58, *post-stress*:  $M = 10.65$ ,  $SD = 4.41$ , *Mean rank* = 70.40,  $Z(135) = -5.70$ ,  $p < .001$ ,  $r = .49$ . Similarly, startle blink amplitudes were analysed through a Wilcoxon signed ranks test, but no manipulation effects were found, *pre-stress*:  $M = 0.05$ ,  $SD = 0.04$ , *Mean*



**Figure 2.** Affective, physiological, and cognitive changes in function of the experimental condition.

Note. The mean changes represent the mean difference between the moment after stress induction and the moment after stress reduction through music listening. Error bars represent the 95% CI. Mean ranks for measures analysed through Wilcoxon signed ranks tests: risky gains (AS-AM: 2.85, AS-IM: 2.38; IS-AM: 2.71; IS-IM: 2.06), risky losses (AS-AM: 2.65; AS-IM: 2.26; IS-AM: 2.54; IS-IM: 2.54), SCL (AS-AM: 2.29; AS-IM: 2.62; IS-AM: 2.18; IS-IM: 2.91). AS: adequate strategy, IS: inadequate strategy, AM: adequate music, IM: inadequate music. † $p < .10$ , \* $p < .05$ , \*\* $p < .01$ , \*\*\* $p < .001$ .

$rank = 52.96$ ,  $post-stress: M = 0.05$ ,  $SD = 0.03$ ,  $Mean rank = 51.94$ ,  $Z(104) = -0.94$ ,  $p = .35$ ,  $r = .092$ . Due to the lack of modulation of the blink amplitudes by the experimental manipulation, this measure was removed from further analyses.

### Influence of strategy and music on affective states

**Self-reported affect.** Figure 2 depicts the mean change in all the outcome measures across experimental conditions. The MANOVA conducted on self-reported affect revealed a main effect of both factors (*strategy*:  $F(3, 23) = 3.25$ ,  $p = .04$ ,  $\eta_p^2 = .30$ ; *music*:  $F(3, 23) = 6.73$ ,  $p = .002$ ,  $\eta_p^2 = .47$ ) and no statistically significant effect for the interaction:  $F(3,23) = 0.46$ ,  $p = .713$ ,  $\eta_p^2 = .06$ . As Table 1 shows, the univariate  $F$  tests revealed that the main effect of *strategy* was significant for the dependent variable tension with a large effect size, but not for energy or valence. This effect was expressed by a higher reduction in tension when using *adequate* strategies (mean difference = 0.12,  $std. error = 0.046$ ,  $p = .017$ ). The factor *music* showed a significant effect for all the three self-report scales: *adequate* music—compared to the *inadequate* music—had a larger contribution in reducing tension (mean difference =

0.21,  $std. error = .059$ ,  $p = .001$ ), increasing positive valence (mean difference = 0.17,  $std. error = 0.065$ ,  $p = .016$ ), and decreasing energy (mean difference = 0.13,  $std. error = 0.057$ ,  $p = .029$ ). Even though the effect size for the three scales was large ( $\eta_p^2 > .14$ ), the effects were larger for the reduction of tension than for valence and energy. There was no interaction effect between *strategy* and *music* on any of the affect dimensions.

### Physiological activity: Skin conductance levels

SCLs related to the factors *strategy* and *music* revealed statistically significant differences:  $\chi_F^2(3) = 10.20$ ,  $p = .016$ ,  $W = .10$ . As reported in Table 1, mean SCLs did not significantly vary in function of *strategy*, but exhibited statistically significant differences in function of *music*. The effect of *adequate* music in decreasing more successfully SCLs than *inadequate* music reached a moderate effect size.

### Cognitive processing: Risky choices

During the experiment, the decision task was performed eight times: four times as part of the stress induction task and four times in the post-regulatory measurement period.



**Table 1.** Marginal means  $\pm$  standard deviation of each level of factors *strategy* and *music* and factorial effects on each measure.

Measures	Strategy			Music			Effects	p value	Effect size
	Adequate	Inadequate	Inadequate	Adequate	Inadequate	Inadequate			
Tension	-0.30 $\pm$ 0.20	-0.18 $\pm$ 0.22	-0.18 $\pm$ 0.22	-0.35 $\pm$ 0.23	-0.14 $\pm$ 0.23	-0.14 $\pm$ 0.23	<b>Strategy:</b> $F(1, 25) = 6.52$ <b>Music:</b> $F(1, 25) = 12.84$	.017	$\eta_p^2 = .207$
Valence	0.15 $\pm$ 0.23	0.15 $\pm$ 0.27	0.15 $\pm$ 0.27	0.23 $\pm$ 0.23	0.06 $\pm$ 0.28	0.06 $\pm$ 0.28	<b>Strategy:</b> $F(1, 25) = 0.004$ <b>Music:</b> $F(1, 25) = 6.75$	.95	$\eta_p^2 = .339$
Energy	-0.17 $\pm$ 0.29	-0.10 $\pm$ 0.24	-0.10 $\pm$ 0.24	-0.20 $\pm$ 0.27	-0.07 $\pm$ 0.22	-0.07 $\pm$ 0.22	<b>Strategy:</b> $F(1, 25) = 1.27$ <b>Music:</b> $F(1, 25) = 5.39$	.016	$\eta_p^2 = .213$
Risky gains	0.13 $\pm$ 0.22 (2.72)	0.052 $\pm$ 0.26 (2.28)	0.052 $\pm$ 0.26 (2.28)	0.19 $\pm$ 0.24 (2.97)	-0.007 $\pm$ 0.26 (2.03)	-0.007 $\pm$ 0.26 (2.03)	<b>Strategy:</b> $F(1, 25) = -1.67$ <b>Music:</b> $Z(34) = -3.29$	.029	$\eta_p^2 = .177$
Risky losses	0.022 $\pm$ 0.25 (2.44)	0.037 $\pm$ 0.21 (2.56)	0.037 $\pm$ 0.21 (2.56)	0.052 $\pm$ 0.22 (2.60)	0.007 $\pm$ 0.23 (2.40)	0.007 $\pm$ 0.23 (2.40)	<b>Strategy:</b> $Z(34) = -0.077$ <b>Music:</b> $Z(34) = -2.80$	.11	$r = .20$
SCLs	0.029 $\pm$ 0.89 (2.50)	-0.029 $\pm$ 0.89 (2.50)	-0.029 $\pm$ 0.89 (2.50)	-0.41 $\pm$ 0.76 (2.00)	0.41 $\pm$ 0.76 (3.00)	0.41 $\pm$ 0.76 (3.00)		.001	$r = .40$

Note. The effects on self-reported affect (tension, valence, and energy) were computed jointly in a two-way within-subjects MANOVA, while the remaining effects were explored through Wilcoxon signed rank tests (for which mean ranks are provided in parentheses). The factors with statistically significant effects are marked in boldface. <sup>a</sup>The marginal means did not show statistically significant differences in the Friedman test, so no pairwise comparisons were made,  $\chi^2(3) = 1.59, p = .68, K = .016$ .

**Table 2.** Test statistics, p values, and effect sizes of pairwise comparisons between conditions.

Conditions	Tension (N = 29)		Valence (N = 30)		Energy (N = 29)	
	t	d	t	d	t	d
AS-AM—AS-IM	-1.97 <sup>†</sup>	.50	2.49*	.54	-1.13	ns
AS-AM—IS-AM	-1.22	ns	-0.60	ns	-0.52	ns
AS-AM—IS-IM	-3.90***	.92	1.99 <sup>†</sup>	.46	-2.24*	.49
AS-IM—IS-AM	0.96	ns	2.40*	.60	0.84	ns
IS-IM—AS-IM	-1.51	ns	0.11	ns	-0.84	ns
IS-IM—IS-AM	-2.93***	.71	2.53*	.59	-2.04*	.48
	SCLs (N = 34)		Gains (N = 34)		Losses (N = 34)	
	Z	r	Z	r	Z	r
AS-AM—AS-IM	-1.58	ns	-2.08*	.25	-1.09	ns
AS-AM—IS-AM	-0.45	ns	-0.53	ns	-0.63	ns
AS-AM—IS-IM	-1.77 <sup>†</sup>	.21	-3.12***	.38	-0.43	ns
AS-IM—IS-AM	-1.67 <sup>†</sup>	.20	-1.62	Ns	-1.16	ns
IS-IM—AS-IM	-0.44	ns	-1.43	Ns	-0.91	ns
IS-IM—IS-AM	-2.27*	.28	-2.42*	.29	-0.32	ns

Note. The first column shows the two conditions being compared. AS – adequate strategy, IS – inadequate strategy, AM – adequate music, IM – inadequate music. <sup>†</sup> $p < .10$ , \* $p < .05$ , \*\* $p < .01$ , \*\*\* $p < .001$

In comparison to the moment of stress induction, participants made more risky choices after listening to music and self-regulating in the gain trials:  $Z(34) = -2.34, p = .020, r = .21$ . However, no significant differences were found in the loss trials,  $Z(34) = -1.40, p = .18, r = .07$ . Friedman tests revealed that there were changes in decision-making as a function of condition in the gain trials, but not in the loss trials (Table 1).

Post-hoc analyses were performed on the marginal means of *strategy* and *music* to explore the effect of these factors. The Friedman test pointed at a significant effect of the experimental factors,  $\chi^2(3) = 19.42, p < .001, W = 0.19$ . The gain trials with *adequate* music playing registered greater proportions of risky choices than the ones with *inadequate* music, but no statistically significant differences between the *adequate* and *inadequate* strategy were registered (Table 1).

**Pairwise comparisons between conditions.** The mean changes registered after conditions AS-AM and IS-IM (Figure 2) were compared for each outcome measure across all conditions. The changes can be visualized in Figure 2 and the test statistics, p values, and effect sizes can be consulted in Table 2.

The condition AS-AM exhibited a higher success in stress reduction than the condition IS-IM as shown by a larger reduction of tension and energy, and a higher risk-taking behaviour in the gain trials. Also, there was a non-significant tendency for more positive affect ( $p = .056, d = .46$ ) and lower SCLs ( $p = .077, r = .21$ ). In comparison to

the condition AS-IM, AS-AM supported a larger (but only marginally significant,  $p = .059$ ) reduction of tension, a larger increase of positive affect, and a higher proportion of risky choices in gain trials. The condition IS-IM differed from condition IS-AM in terms of lower reduction in tension, lower increase of positive affect, lower decrease of energy, higher SCLs, and lower risk-taking in gain trials. There were no statistically significant pairwise comparisons in the risky choices in the loss domain.

## Discussion

While psychology and music research have consistently found beneficial effects of music on stress responses, the present study contributes to the existing literature by experimentally testing two of the underlying components of musical self-regulation: the specific music listened to and the employed regulatory strategy. We hypothesized that both factors would have a positive effect on stress reduction (H.1 and H.2) and that the combination of efficacious music and strategy would lead to the most successful reduction in stress while the combination of inefficacious music and strategy would lead to the least successful stress reduction (H.3). The results corroborated hypotheses 1 and 2, while not fully supporting hypothesis 3. No hypotheses were formulated regarding the relative impact of strategies and music; nevertheless, music seemed to have a more visible impact than strategies on the measures used in this study.

### Music's support for stress reduction (hypothesis 1)

As expected, music perceived as efficacious led to a larger reduction of acute stress than music perceived as inefficacious. This reduction was translated into lower tension levels, more positive affect, lower energy arousal, lower SCLs, and higher risk-taking in the gain trials. All the measured modalities (self-reported affect, physiological activity, and cognitive processing) were affected by the efficacy of music. These results provide solid support for hypothesis 1.

The changes in the self-report VAS portray a multi-dimensional modulation of affect in the direction of stress reduction. Aside from the predicted reduction of tension, our results suggest that the down-regulation of stress was also approached through creating a more positive/pleasant state and through decreasing the energy levels. These results are supported by previous literature that has already highlighted music's potential to support each one of these dimensions, albeit separately (tension reduction: Thayer, Newman, & McClain, 1994; increase of positive affect: Van Goethem & Sloboda, 2011; and decrease of energy levels: DeNora, 1999; Saarikallio, Baltazar, & Västfjäll, 2017).

Regarding the physiological dimension, SCLs were responsive to the stress manipulation and to music's efficacy, increasing after stress induction and decreasing

especially after the *adequate* music. Surprisingly, the startle blink amplitudes were not sensitive to the stress manipulation. The startle blink is an aversive response and it is expected to exhibit increased amplitudes in relation to negative valence (Lang et al., 1990). However, some authors have suggested that acute stress is a specific state that favours alertness and focused attention and that, consequently, does not increase the startle blink amplitudes during the experience of stress (Deuter et al., 2012; Pinkney, Wickens, Bamford, Baldwin, & Garner, 2014) but only after recovery (Herten et al., 2016).

Music's efficacy also influenced decisions in the gain trials of risk-taking. As expected, listening to the *adequate* music pieces was linked to increased risk-taking. Given that acute stress and anxiety have been reported as lowering risk-taking (Porcelli & Delgado, 2009), it can be concluded that the participants were under higher stress while listening to the *inadequate* music. Moreover, positive affective priming can lead to calculating probabilities in a more optimistic way or perceiving gains as more attractive, thus promoting risk-taking (for a review, see George & Dane, 2016). In our case, *adequate* music would have induced more positive affect than the *inadequate* music. Alternatively, the increase in risk-taking might be explained by the ongoing affect regulation. Andrade and Cohen (2007) posit that when individuals believe they can regulate their affect, they focus more on the gains of the risky activity and become more risk-taking. In our case, *adequate* music would have been perceived as adjuvant of self-regulation of stress, while *inadequate* music would have impeded the process. Both interpretations presented lead to the same conclusion: the *adequate* music was more supportive of stress reduction than the *inadequate* music. Even though research on music and decision-making is still slim, there is already some evidence for an increased subjective value of risky options while listening to preferred music (Halko & Kaustia, 2015) and to motivating music (Elvers & Steffens, 2017).

The loss trials registered stable behaviours across moments (stress induction and musical stress regulation) and across conditions. Since there were no trials before the stress induction, it is not possible to assess how this decision-making was impacted by the affect manipulation. However, one possible explanation for the lack of variation is that the procedure was not propitious for manipulating the loss domain. According to Pabst, Schoofs, Pawlikowski, Brand, and Wolf (2013), loss framing is slower and more complex; thus, depending on the time passed after the stressful stimulus, risk-taking might be increased, decreased, or unchanged.

### Strategic support for stress reduction (hypothesis 2)

Strategies perceived as efficacious showed a larger contribution to stress reduction than the ones perceived as inefficacious by leading to lower levels of self-reported *tension*

arousal. Given the relevance of tension for the experience of stress responses and the large effect size observed, we consider our results in line with H.2. The positive effect of the *adequate* strategy on experienced tension confirms previous claims that the use of adequate regulatory strategies while listening to music is responsible for a considerable part of its affective outcome (Chin & Rickard, 2014; Garrido & Schubert, 2013).

### Comparisons between conditions (hypothesis 3)

Hypothesis 3 could not be verified, since condition AS-AM was not significantly more successful in reducing stress than all the other conditions, and condition IS-IM was not significantly worse than all the other conditions. Yet, condition AS-AM consistently led to better results after stress induction than IS-IM.

Interestingly, the pairwise comparisons failed to reveal a statistically significant difference between conditions when music was *adequate* and strategy changed from *adequate* to *inadequate* (AS-AM v IS-AM), and when music was *inadequate* and strategy changed (AS-IM v IS-IM). In a similar vein, statistically significant differences were obtained when the employed strategy was the same and music changed from efficacious to inefficacious (AS-AM v AS-IM and IS-AM v IS-IM). This suggests that despite the overall main effect of the employed strategy, in the short term the selected music brings a larger impact to the listener's affective states. This phenomenon was not predicted and might explain why H. 3 could not be supported. Differences in the affective impact will be discussed next.

**Larger effect of music in the short term.** Despite the observed effects of both factors (music and regulatory strategies), their contribution to the participants' stress reduction was not equal in this experimental setup. While the regulatory strategies showed a significant effect in the principal outcome measure, music exhibited a larger effect size and, additionally, its effect was registered in complementary measures across the three dimensions. Also, in the pairwise comparisons music seemed to have a stronger impact, as it was a decisive factor when changing or maintaining the efficacy level.

We interpret the less visible role of regulation strategies in comparison to music as a product of several aspects:

1. *Automatic and non-deliberate nature of affect regulation.* One of the difficulties in experimentally studying affect self-regulation emerges from the fact that most of its processes do not happen at the conscious level (Bargh & Chartrand, 1999). Part of affect regulation happens automatically, triggered by sensory inputs, and based on previous experiences (Braunstein, Gross, & Ochsner, 2017; Koole, Webb, & Sheeran, 2015; Mauss, Bunge, & Gross,

2007). Given the organisms' drive to self-regulate, the participants might have taken unconscious and parallel efforts to reduce stress responses despite the instructions to use a specific inefficacious strategy.

2. *Music's immediacy and salience.* Music, as a highly emotionally-charged auditory stimulus, is particularly impactful (Panksepp & Bernatzky, 2002)—it activates the brain areas linked to emotion processing and induces neurochemical changes in the domains of reward, motivation, and pleasure (Chanda & Levitin, 2013). Music might have been, then, more salient and stimulating than the instructions regarding the strategy to employ.
3. *Temporal dynamics of strategies.* Regulatory strategies have varying onset times (Paul, Simon, Kniesche, Kathmann, & Endrass, 2013; Thiruchselvam, Blechert, Sheppes, Rydstrom, & Gross, 2011) and their effects can be sustained even after active regulation (Walter et al., 2009). It is possible that the strategies' effects (positive or negative) were still unfolding at the time of measurement.
4. *Not all emotions are the same.* The efficacy (and even adaptiveness) of strategies is relatively sensitive to varied aspects related to the origin and type of affective reaction. For instance, previous research has shown that reappraisal's positive effect depends on the origin of the emotion (McRae, Misra, Prasad, Pereira, & Gross, 2012) and on its intensity (Sheppes, Scheibe, Suri, & Gross, 2011). In our study, mismatches between the imagined affect (online survey) and experienced affect (laboratory) might have influenced the strategies' efficacy.

Moreover, it is possible to adopt a broader view on how music and strategies come together. Referring to Krueger's idea of "musical scaffolding" (2011, 2014), there is part of the affective work that is done in the music and by the music, as if we could transfer our internal processes to this external resource. During musical scaffolding, internal processes (such as regulatory strategies) are enhanced by the possibilities offered through music. It might be thus that, in practice, it is hard to separate the musical characteristics from the strategy, since all the components intertwine and become part of the same experience for the user. This interpretation is supported by the results obtained by Baltazar and Saarikallio (2017), which show deep connections between musical mechanisms and strategies alongside the dimensions of affective experience, cognition, and bodily reactions. Also, in the model suggested by Baltazar (in press), these same connections mediate the relationship between music engagement and affective outcomes. The intrinsically regulatory attributes of music mean that music offers wide affective possibilities beyond the regulatory strategy consciously applied.

## Health implications

A growing position defends that music's health impact does not depend solely on its overall features (e.g. genre) but mainly on the personal use that individuals make of it (McFerran, 2016; McFerran, Garrido, & Saarikallio, 2013; Saarikallio, Gold, & McFerran, 2015). Regulatory strategies have been one link between the resource (music) and health outcomes (e.g. Carlson et al., 2015; Chin & Rickard, 2014). However, the ability of choosing what music to listen to can be equally relevant. As our results show, for each individual there is certain music that facilitates or hinders a certain affective goal in a certain context, regardless of the strategy used. When, for some reason, the chosen music is not adequate for the situation, it can lead to undesired mood shifts or worsening of negative mood (McFerran, Garrido, O'Grady, Grocke, & Sawyer, 2015; Randall & Rickard, 2017). Importantly, studies with vulnerable or distressed groups indicate that not everyone has the same ability of regulating through music (Baker, Bor, & William, 2008; McFerran et al., 2015; Miranda & Claes, 2009). We suggest the term *musico-emotional competence* to encompass the skills necessary for picking the right song and coupling it with the right emotional processing (see also Saarikallio, 2017). While the emotional skills include, among others, skills related to the identification and regulation of emotions (Mayer & Salovey, 1997), the musical skills would include, among others, skills related to recognizing the emotions expressed by music and their impact on own affective states, selecting music in accordance with the affective goal, and matching the strategy to the music (Saarikallio, 2017).

In the current study, our healthy, adult participants could identify the adequate music and strategy for themselves. But to what degree can this competence be influenced by individual traits and in what situations would participants be unable to make the adequate regulatory and musical decisions? Further research with group comparisons (such as between ages, genders, psychological symptoms, etc.) is needed to better understand musico-emotional competence.

## Limitations

A few limitations should be considered when interpreting our results and planning future research. Some derive from the four points presented above in relation to the contrasting effects of music and regulatory strategy. That is, the lack of control of the strategies used by the participants due to automatic self-regulation, the potential differences in salience of the two factors (music played versus instruction to use a certain strategy), the relatively short time constrain for each regulation period, and the compromise of ecological validity for the sake of experimental control. Also, it should be noted that only one strategy was contemplated at a time; however, some studies have been showing that people use more than one strategy simultaneously or

sequentially (Aldao & Nolen-Hoeksema, 2013; Baltazar & Saarikallio, 2017). Investigating individual factors was not in the scope of this study; however it is likely that the sensitivity to the experimental manipulation and to the effects of the two factors, music and strategy, varies across groups in function of age (Cohrdes, Wrzus, Wald-Fuhrmann, & Riediger, 2018; Leipold & Loepthien, 2015), personality traits (Liljeström et al., 2012), reactivity to stress (Thoma, Scholz, Ehlert, & Nater, 2012), and profile of music use (ter Bogt, Mulder, Raaijmakers, & Gabhainn, 2011). In future studies, the individual factors could be approached by using larger samples with different groups; the contextual factors could be addressed by using methodologies that integrate them, such as *experience sampling method*; and the effect of regulatory strategies could be maximized by including a practice time before the experiment and by giving a longer period for self-regulation.

## Conclusion

If we had to answer the title question, we would be inclined to answer: *both*. Both the adequacy of the music and the adequacy of the strategy expressed themselves in the affective outcomes of stress reduction. Yet, given that music creates the platform where the process takes place, its adequacy has an immediate and decisive impact on the regulatory efforts. There are intricate and deep links between music and regulatory strategies (and, in the same line, affective goals) that are built on a matrix of associations, memories, individual traits, acoustic features, and emotion induction processes (Baltazar, in press; Baltazar & Saarikallio, 2017; DeNora, 1999; Juslin & Västfjäll, 2008). These links are at the foundation of musical affect self-regulation and, as in everything, solid and adequate foundations are needed for the construction of healthy structures. Our results highlight the relevance of musico-emotional skills by demonstrating the contrasting effects of using the “adequate” or “inadequate” music and the “adequate” or “inadequate” regulatory strategy.

## Acknowledgements

The authors wish to thank Kajsa Hansson, for her relentless help during recruitment and data collection, Dr. India Morrison, for generously sharing resources with us, Per Andersson, for volunteering his skin and ears during our testing phase, and all the participants who bravely came down to the laboratory.

## Authors' contributions

MB and DV conceived the study. MB, DV, EA, LK, and SS developed the design. EA coded the experiment script and handled the physiological data. LK assisted MB with the participant recruitment. MB collected and analysed the data. MB wrote the first draft of the manuscript, and all authors reviewed and edited the manuscript, and approved the final version of the manuscript.

### Declaration of conflicting interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

### Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This work was funded by the Finnish Cultural Foundation and the Science Council of the University of Jyväskylä to MB and by the Swedish Research Council to DV, EA, and LK.

### Supplemental material

Supplemental material for this article is available online.

### Notes

1. In this work, we use *affect* as an umbrella term encompassing all the evaluative—positive or negative—states (Juslin & Sloboda, 2010), including emotion, arousal, stress responses, motivational impulses, interpersonal stances, preferences, mood, attitudes, and affective style (Baltazar & Saarikallio, 2016).
2. NB: In this study, we are not interested in the efficacy of particular strategies or musical features, but on the measurement of the affective consequences of using either the adequate or inadequate strategy/music at the individual level. Therefore, data related to the chosen strategies and music examples will not be analysed in this work.
3. The rating of these aspects was a methodological choice. In the second phase they were used to guarantee that the same positive or negative effects were triggered regardless of the fit with the affective goal.
4. The informed consent form included information about the pain stimulation that would occur during the experiment. It was stated that a painful thermal stimulation would be applied on their forearm, that its level would be calibrated to their own pain threshold, and that they could experience some redness in the stimulated area that normally disappears in 1–2 hours. Participants were also informed that they had the right to withdraw from the study at any point and without any consequences. The information regarding the pain and their rights as participants was also expressed verbally before starting the experiment. After the data collection, there was a period for de-briefing and guaranteeing that the participants were not leaving the laboratory in an anxious or depressed mood.
5. I.e. the order of the conditions was randomized and every possible permutation of the four conditions was attributed to a participant at least once.
6. Some participants had given more than one example for each condition. Their first pick was always preferred, but in case it was not specific enough (a whole album, for example) or it was not available in Spotify, their second (or third) was used instead. Even when more than one song was available, only one per condition was used across the experiment to avoid potential confounds.

7. The limit of musical aspects was set to three since it was considered to be enough to produce the affective impact while not demanding too much from the participants. The cut-off point was a rating of relevance of 70 points out of 100. If a tie in ratings made it impossible for the researcher to automatically choose the top three mechanisms, the researcher asked the participant which was more relevant. From the list of nine aspects listed in the survey, two were excluded due to their low relevance for this purpose—music's genre/preference and familiarity with the music.
8. When outliers were excluded from the analysis, equivalent non-parametric tests were applied to the complete sample. As no differences were found in the results, these will not be reported.

### Peer review

John Sloboda, Guildhall School of Music and Drama, Research Department.  
Kimberly Moore, University of Miami.  
Eduardo Coutinho, University of Liverpool, Department of Music.  
One anonymous reviewer.

### References

- Aldao, A., Sheppes, G., & Gross, J. J. (2015). Emotion regulation flexibility. *Cognitive Therapy and Research*, *39*, 263–278.
- Aldao, A., & Nolen-Hoeksema, S. (2013). One versus many: Capturing the use of multiple emotion regulation strategies in response to an emotion-eliciting stimulus. *Cognition and Emotion*, *27*, 753–760.
- Andrade, E. B., & Cohen, J. B. (2007). Affect-based evaluation and regulation as mediators of behavior: The role of affect in risk taking, helping, and eating patterns. In K. D. Vohs, R. F. Baumeister, & G. Loewenstein (Eds.), *Do emotions help or hurt decision making? A hedgefoxian perspective* (pp. 35–68). New York, NY: Russell Sage.
- Arnsten, A. F. T. (2009). Stress signalling pathways that impair prefrontal cortex structure and function. *Nature Reviews Neuroscience*, *10*, 410–422.
- Baker, F., Bor, W., & William, B. (2008). Can music preference indicate mental health status in young people? *Australasian Psychiatry: Bulletin of Royal Australian and New Zealand College of Psychiatrists*, *16*, 284–288.
- Baltazar, M. (in press). Musical affect regulation in adolescents: A conceptual model. In K. McFerran, P. Derrington, & S. Saarikallio (Eds.), *Handbook of music and adolescence*. New York, NY: University of Oxford Press.
- Baltazar, M. (2018). *Where mind and music meet: Affect self-regulation through music*. Doctoral thesis, University of Jyväskylä. Retrieved from <http://urn.fi/URN:ISBN:978-951-39-7458-9>
- Baltazar, M., & Saarikallio, S. (2016). Toward a better understanding and conceptualization of affect self-regulation through music: A critical, integrative literature review. *Psychology of Music*, *44*, 1500–1521.

- Baltazar, M., & Saarikallio, S. (2017). Strategies and mechanisms in musical affect self-regulation: A new model. *Musicae Scientiae*. Advance online publication. <https://doi.org/10.1177/1029864917715061>
- Bargh, J. A., & Chartrand, T. L. (1999). The unbearable automaticity of being. *American Psychologist*, *54*, 462–479.
- Bonanno, G. A., & Burton, C. L. (2013). Regulatory flexibility: An individual differences perspective on coping and emotion regulation. *Perspectives on Psychological Science*, *8*, 591–612.
- Braithwaite, J., Watson, D., Jones, R., & Rowe, M. (2015). *A guide for analysing electrodermal activity (EDA) & skin conductance responses (SCRs) for psychological experiments, 2nd version*. Birmingham. doi:10.1017/S0142716405050034
- Braunstein, L. M., Gross, J. J., & Ochsner, K. N. (2017). Explicit and implicit emotion regulation: A multi-level framework. *Social Cognitive and Affective Neuroscience*, *20*(10), 1545–1557.
- Carlson, E., Saarikallio, S., Toiviainen, P., Bogert, B., Kliuchko, M., & Brattico, E. (2015). Maladaptive and adaptive emotion regulation through music: A behavioral and neuroimaging study of males and females. *Frontiers in Human Neuroscience*, *9*, 1–13.
- Chafin, S., Roy, M., Gerin, W., & Christenfeld, N. (2004). Music can facilitate blood pressure recovery from stress. *British Journal of Health Psychology*, *9*, 393–403.
- Chanda, M. L., & Levitin, D. J. (2013). The neurochemistry of music. *Trends in Cognitive Sciences*, *17*, 179–191.
- Chin, T., & Rickard, N. S. (2014). Emotion regulation strategy mediates both positive and negative relationships between music uses and well-being. *Psychology of Music*, *42*, 692–713.
- Cohen, J. (1977). F tests on means in the analysis of variance and covariance. *Statistical Power Analysis for the Behavioral Sciences*, 273–406. London: Academic Press.
- Cohen, J. (1988). *Statistical power analysis for the behavioral sciences*. Hillsdale, NJ: Erlbaum.
- Cohrdes, C., Wrzus, C., Wald-Fuhrmann, M., & Riediger, M. (2018). “The sound of affect”: Age differences in perceiving valence and arousal in music and their relation to music characteristics and momentary mood. *Musicae Scientiae*. Advance online publication.
- DeNora, T. (1999). Music as a technology of the self. *Poetics*, *27*, 31–56.
- DeNora, T. (2000). *Music in everyday life*. Cambridge, MA: Cambridge University Press.
- Deuter, C. E., Kuehl, L. K., Blumenthal, T. D., Schulz, A., Oitzl, M. S., & Schachinger, H. (2012). Effects of cold pressor stress on the human startle response. *PLoS One*, *7*, 1–7.
- Doré, B. P., Silvers, J. A., & Ochsner, K. N. (2016). Toward a personalized science of emotion regulation. *Social and Personality Psychology Compass*, *10*, 171–187.
- Dunlap, W. P., Cortina, J. M., Vaslow, J. B., & Burke, M. J. (1996). Meta-analysis of experiments with matched groups or repeated measures designs. *Psychological Methods*, *1*, 170–177.
- Elvers, P., & Steffens, J. (2017). The sound of success: Investigating cognitive and behavioral effects of motivational music in sports. *Frontiers in Psychology*, *8*, 2026.
- Epstein, S., Pacini, R., Denes-Raj, V., & Heier, H. (1996). Individual differences in intuitive–experiential and analytical–rational thinking styles. *Journal of Personality and Social Psychology*, *71*, 390–405.
- Evers, S., & Suhr, B. (2000). Changes of the neurotransmitter serotonin but not of hormones during short time music perception. *European Archives of Psychiatry and Clinical Neuroscience*, *250*, 144–147.
- Fancourt, D., Ockelford, A., & Belai, A. (2014). The psychoneuro-immunological effects of music: A systematic review and a new model. *Brain, Behavior, and Immunity*, *36*, 15–26.
- Garrido, S., & Schubert, E. (2013). Adaptive and maladaptive attraction to negative emotions in music. *Musicae Scientiae*, *17*, 147–166.
- George, J. M., & Dane, E. (2016). Affect, emotion, and decision making. *Organizational Behavior and Human Decision Processes*, *136*, 47–55.
- Getz, L. M., Marks, S., & Roy, M. (2014). The influence of stress, optimism, and music training on music uses and preferences. *Psychology of Music*, *42*, 71–85.
- Gross, J. J. (2015). Emotion regulation: Current status and future prospects. *Psychological Inquiry*, *26*, 1–26.
- Gross, J. J., & Thompson, R. A. (2007). Emotion regulation: Conceptual foundations. In J. J. Gross (Ed.), *Handbook of emotion regulation* (pp. 3–26). New York, NY: The Guilford Press.
- Halko, M. L., & Kaustia, M. (2015). Risk ON/Risk OFF: Risk-taking varies with subjectively preferred and disliked music. *PLoS One*, *10*, 1–16.
- Herten, N., Otto, T., Adolph, D., Pause, B. M., Kumsta, R., & Wolf, O. T. (2016). Enhanced startle responsivity 24 hours after acute stress exposure. *Behavioral Neuroscience*, *130*, 521–530.
- Jiang, J., Rickson, D., & Jiang, C. (2016). The mechanism of music for reducing psychological stress: Music preference as a mediator. *The Arts in Psychotherapy*, *48*, 62–68.
- Juslin, P. N. (2013). From everyday emotions to aesthetic emotions: Towards a unified theory of musical emotions. *Physics of Life Reviews*, *10*, 235266.
- Juslin, P. N., Barradas, G., & Eerola, T. (2015). From sound to significance: Exploring the mechanisms underlying emotional reactions to music. *The American Journal of Psychology*, *128*, 281–304.
- Juslin, P. N., & Sloboda, J. A. (2010). Introduction: Aims, organization, and terminology. In P. N. Juslin & S. J. A. (Eds.), *Handbook of music and emotion: Theory, research, applications* (pp. 3–12). New York, NY: Oxford University Press.
- Juslin, P. N., & Västfjäll, D. (2008). Emotional responses to music: The need to consider underlying mechanisms. *The Behavioral and Brain Sciences*, *31*, 559–621.
- Kahneman, D. (2003). A perspective on judgment and choice—Mapping bounded rationality. *American Psychologist*, *58*, 697–720.

- Kahneman, D., & Tversky, A. (1979). Prospect theory: An analysis of decision under risk. *Econometrica*, *47*, 263–292.
- Kendall, M. G. (1938). A new measure of rank correlation. *Biometrika*, *30*, 81–93.
- Khalfa, S., Dalla Bella, S., Roy, M., Peretz, I., & Lupien, S. J. (2003). Effects of relaxing music on salivary cortisol level after psychological stress. *Annals of the New York Academy of Sciences*, *999*, 374–376.
- Khalfa, S., Peretz, I., Blondin, J.-P., & Manon, R. (2002). Event-related skin conductance responses to musical emotions in humans. *Neuroscience Letters*, *328*, 145–149.
- Koole, S. L. (2009). The psychology of emotion regulation: An integrative review. *Cognition and Emotion*, *23*, 4–41.
- Koole, S. L., Webb, T. L., & Sheeran, P. L. (2015). Implicit emotion regulation: Feeling better without knowing why. *Current Opinion in Psychology*, *3*, 6–10.
- Koppel, L., Andersson, D., Morrison, I., Posadzy, K., Västfjäll, D., & Tinghög, G. (2017). The effect of acute pain on risky and intertemporal choice. *Experimental Economics*, *20*(4), 878–893.
- Koppel, L., Mederyd, A., Morrison, I., Tinghög, G., & Västfjäll, D. (submitted). The effect of pain and touch on risk taking depends on the perceived affective consequences.
- Kreutz, G., Ott, U., Teichmann, D., Osawa, P., & Vaitl, D. (2007). Using music to induce emotions: Influences of musical preference and absorption. *Psychology of Music*, *36*, 101–126.
- Krueger, J. W. (2011). Doing things with music. *Phenomenology and the Cognitive Sciences*, *10*, 1–22.
- Krueger, J. W. (2014). Affordances and the musically extended mind. *Frontiers in Psychology*, *4*, 1003–1015.
- Lang, P. J., Bradley, M. M., & Cuthbert, B. N. (1990). Emotion, attention, and the startle reflex. *Psychological Review*, *97*, 377–395.
- Lazarus, R. S. (1974). Psychological stress and coping in adaptation and illness. *The International Journal of Psychiatry in Medicine*, *5*, 321–333.
- Lazarus, R. S. (1990). Theory-based stress measurement. *Psychological Inquiry*, *1*, 3–13.
- Lazarus, R. S. (2000). How emotions influence performance in competitive sports. *The Sport Psychologist*, *14*, 229–252.
- Lazarus, R. S., & Folkman, S. (1984). *Stress, appraisal, and coping*. New York, NY: Springer.
- Leipold, B., & Loepthien, T. (2015). Music reception and emotional regulation in adolescence and adulthood. *Musicae Scientiae*, *19*, 111–128.
- Liljeström, S., Juslin, P. N., & Västfjäll, D. (2012). Experimental evidence of the roles of music choice, social context, and listener personality in emotional reactions to music. *Psychology of Music*, *41*, 579–599.
- Linnemann, A., Ditzen, B., Strahler, J., Doerr, J. M., & Nater, U. M. (2015). Music listening as a means of stress reduction in daily life. *Psychoneuroendocrinology*, *60*, 82–90.
- Mandrick, K., Peysakhovich, V., Rémy, F., Lepron, E., & Causse, M. (2016). Neural and psychophysiological correlates of human performance under stress and high mental workload. *Biological Psychology*, *121*, 62–73.
- Marik, M., & Stegemann, T. (2016). Introducing a new model of emotion dysregulation with implications for everyday use of music and music therapy. *Musicae Scientiae*, *20*, 5367.
- Mauss, I. B., Bunge, S. a., & Gross, J. J. (2007). Automatic emotion regulation. *Social and Personality Psychology Compass*, *1*, 146–167.
- Mayer, J. D., & Salovey, P. (1997). What is emotional intelligence? In P. Salovey & D. Sluyter (Eds.), *Emotional development and emotional intelligence: Implications for educators* (pp. 3–31). New York, NY: Basic Books.
- McEwen, B. S., & Stellar, E. (1993). Stress and the individual. *Archives of Internal Medicine*, *153*, 2093–2101.
- McFerran, K. S. (2016). Contextualising the relationship between music, emotions and the well-being of young people: A critical interpretive synthesis. *Musicae Scientiae*, *20*, 103–121.
- McFerran, K. S., Garrido, S., O’Grady, L., Grocke, D., & Sawyer, S. M. (2015). Examining the relationship between self-reported mood management and music preferences of Australian teenagers. *Nordic Journal of Music Therapy*, *24*, 187–203.
- McFerran, K. S., Garrido, S., & Saarikallio, S. (2013). A critical interpretive synthesis of the literature linking music and adolescent mental health. *Youth & Society*, *20*, 1–18.
- McRae, K., Misra, S., Prasad, A. K., Pereira, S. C., & Gross, J. J. (2012). Bottom-up and top-down emotion generation: Implications for emotion regulation. *Social Cognitive and Affective Neuroscience*, *7*, 253–262.
- Menon, V., & Levitin, D. J. (2005). The rewards of music listening: Response and physiological connectivity of the mesolimbic system. *NeuroImage*, *28*, 175–184.
- Miranda, D., & Claes, M. (2009). Music listening, coping, peer affiliation and depression in adolescence. *Psychology of Music*, *37*, 215–233.
- Munck, A., Guyre, P. M., & Holbrook, N. J. (1984). Physiological functions of glucocorticoids in stress and their relation to pharmacological actions. *Endocrine Reviews*, *5*, 25–44.
- Pabst, S., Schoofs, D., Pawlikowski, M., Brand, M., & Wolf, O. T. (2013). Paradoxical effects of stress and an executive task on decisions under risk. *Behavioral Neuroscience*, *127*, 369–379.
- Pallant, J. (2007). SPSS survival manual. *Journal of Advanced Nursing*, *36*, 478–478.
- Panksepp, J., & Bernatzky, G. (2002). Emotional sounds and the brain: The neuro-affective foundations of musical appreciation. *Behavioural Processes*, *60*, 133–155.
- Parkinson, B., & Totterdell, P. (1999). Classifying affect-regulation strategies. *Cognition and Emotion*, *13*, 277–303.
- Paul, S., Simon, D., Kniesche, R., Kathmann, N., & Endrass, T. (2013). Timing effects of antecedent- and response-focused emotion regulation strategies. *Biological Psychology*, *94*, 136–142.
- Pelletier, C. L. (2004). The effect of music on decreasing arousal due to stress: A meta-analysis. *Journal of Music Therapy*, *41*, 192–214.

- Pereira, C. S., Teixeira, J., Figueiredo, P., Xavier, J., Castro, S. L., & Brattico, E. (2011). Music and emotions in the brain: Familiarity matters. *PLoS One*, *6*, e27241.
- Perini, I., Bergstrand, S., & Morrison, I. (2013). Where pain meets action in the human brain. *The Journal of Neuroscience*, *33*(40), 15930–15939.
- Pinkney, V., Wickens, R., Bamford, S., Baldwin, D. S., & Garner, M. (2014). Defensive eye-blink startle responses in a human experimental model of anxiety. *Journal of Psychopharmacology*, *28*(9), 874–880.
- Porcelli, A. J., & Delgado, M. R. (2009). Acute stress modulates risk taking in financial decision making. *Psychological Science*, *20*, 278–283.
- Randall, W. M., & Rickard, N. S. (2017). Personal music listening: A model of emotional outcomes developed through mobile experience sampling. *Music Perception: An Interdisciplinary Journal*, *34*, 501–514.
- Rosenthal, R. (1994). Parametric measures of effect size. In H. Cooper & L. V. Hedges (Eds.), *The handbook of research synthesis* (pp. 231–244). New York, NY: Russell Sage Foundation.
- Saarikallio, S. (2017). Musical identity in fostering emotional health. In R. MacDonald & D. J. Hargreaves (Eds.), *Handbook of musical identities* (pp. 602–623). New York, NY: Oxford University Press.
- Saarikallio, S., Baltazar, M., & Västfjäll, D. (2017). Adolescents' musical relaxation: Understanding the related affective processing. *Nordic Journal of Music Therapy*, *26*, 376–389.
- Saarikallio, S., Gold, C., & McFerran, K. S. (2015). Development and validation of the Healthy-Unhealthy Music Scale. *Child and Adolescent Mental Health*, *4*, 210–217.
- Sandstorm, G. M., & Russo, F. A. (2010). Music hath charms: The effects of valence and arousal on recovery following an acute stressor. *Music & Medicine*, *2*, 137–143.
- Scherer, K. R., & Coutinho, E. (2013). How music creates emotion: A multifactorial process approach. In T. Cochrane, B. Fantini, & K. R. Scherer (Eds.), *The emotional power of music: Multifisciplinary perspectives on musical expression, arousal, and social control* (pp. 121–145). New York, NY: Oxford University Press.
- Schimmack, U., & Grob, A. (2000). Dimensional models of core affect: A quantitative comparison by means of structural equation modeling. *European Journal of Personality*, *14*, 325–345.
- Schneiderman, N., Ironson, G., & Siegel, S. D. (2005). Stress and health: Psychological, behavioral, and biological determinants. *Annual Review of Clinical Psychology*, *1*, 607–628.
- Selye, H. (1936). A syndrome produced by diverse nocuous agents. *Nature*, *138*, 32.
- Shallcross, A. J., Troy, A., & Mauss, I. B. (2015). Regulation of emotions under stress. In R. Scott & K. Stephen (Eds.), *Emerging trends in the social and behavioral sciences*. doi:10.1002/9781118900772.etrds0036.
- Sheppes, G., Scheibe, S., Suri, G., & Gross, J. J. (2011). Emotion-regulation choice. *Psychological Science*, *22*, 1391–1396.
- Shields, G. S., Sazma, M. A., & Yonelinas, A. P. (2016). The effects of acute stress on core executive functions: A meta-analysis and comparison with cortisol. *Neuroscience and Biobehavioral Reviews*, *68*, 651–668.
- Skånland, M. S. (2011). Use of mp3-players as a coping resource. *Music and Arts in Action*, *3*, 15–33.
- Smith, C. A., Haynes, K. N., Lazarus, R. S., & Pope, L. K. (1993). In search of the “hot” cognitions: Attributions, appraisals, and their relation to emotion. *Journal of Personality and Social Psychology*, *65*, 916–929.
- Starcke, K., & Brand, M. (2012). Decision making under stress: A selective review. *Neuroscience and Biobehavioral Reviews*, *36*, 1228–1248.
- Tamir, M., & Ford, B. Q. (2012). Should people pursue feelings that feel good or feelings that do good? Emotional preferences and well-being. *Emotion*, *12*, 1061–1070.
- Tan, X., Yowler, C. J., Super, D. M., & Fratianne, R. B. (2012). The interplay of preference, familiarity and psychophysical properties in defining relaxation music. *Journal of Music Therapy*, *49*, 150–179.
- ter Bogt, T. F. M., Mulder, J., Raaijmakers, Q. A. W., & Gabhainn, S. N. (2011). Moved by music: A typology of music listeners. *Psychology of Music*, *39*, 147–163.
- ter Bogt, T. F. M., Vieno, A., Doornwaard, S. M., Pastore, M., & van den Eijnden, R. J. J. M. (2017). “You’re not alone”: Music as a source of consolation among adolescents and young adults. *Psychology of Music*, *45*, 155–171.
- Thayer, R. E., Newman, J. R., & McClain, T. M. (1994). Self-regulation of mood: Strategies for changing a bad mood, raising energy, and reducing tension. *Journal of Personality and Social Psychology*, *67*, 910–925.
- Thiruchselvam, R., Blechert, J., Sheppes, G., Rydstrom, A., & Gross, J. J. (2011). The temporal dynamics of emotion regulation: An EEG study of distraction and reappraisal. *Biological Psychology*, *87*, 84–92.
- Thoma, M. V., La Marca, R., Brönnimann, R., Finkel, L., Ehlert, U., & Nater, U. M. (2013). The effect of music on the human stress response. *PLoS One*, *8*, e70156.
- Thoma, M. V., Scholz, U., Ehlert, U., & Nater, U. M. (2012). Listening to music and physiological and psychological functioning: The mediating role of emotion regulation and stress reactivity. *Psychology & Health*, *27*, 227–241.
- Tsigos, C., & Chrousos, G. P. (2002). Hypothalamic-pituitary-adrenal axis, neuroendocrine factors and stress. *Journal of Psychosomatic Research*, *53*, 865–871.
- Tversky, A., & Kahneman, D. (1983). Extensional vs. intuitive reasoning: The conjunction fallacy in probability judgment. *Psychological Review*, *90*, 293.
- Van Goethem, A., & Sloboda, J. A. (2011). The functions of music for affect regulation. *Musicae Scientiae*, *15*, 208–228.
- Västfjäll, D. (2010). Indirect perceptual, cognitive, and behavioural measures. In P. N. Juslin & J. A. Sloboda (Eds.), *Handbook of music and emotion: Theory, research, applications* (pp. 255–278). New York, NY: Oxford University Press.
- Vohs, K. D., Baumeister, R. F., Schmeichel, B. J., Twenge, J. M., Nelson, N. M., & Tice, D. M. (2008). Making choices impairs



- subsequent self-control: A limited-resource account of decision making, self-regulation, and active initiative. *Journal of Personality and Social Psychology*, *94*, 883–898.
- Walter, H., von Kalckreuth, A., Schardt, D., Stephan, A., Goschke, T., & Erk, S. (2009). The temporal dynamics of voluntary emotion regulation. *PLoS One*, *4*, e6726.
- Witvliet, C. V., & Vrana, S. R. (1995). Psychophysiological responses as indexes of affective dimensions. *Psychophysiology*, *32*, 436–443.
- Yu, R. (2016). Stress potentiates decision biases: A stress induced deliberation-to-intuition (SIDI) model. *Neurobiology of Stress*, *3*, 83–95.