JYX JYVÄSKYLÄN YLIOPISTO UNIVERSITY OF JYVÄSKYLÄ

This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Pirnes, Katariina; Kallio, Jouni; Siekkinen, Kirsti; Hakonen, Harto; Häkkinen, Arja; Tammelin, Tuija

Title: Test-retest repeatability of questionnaire for pain symptoms for school children aged 10–15 years

Year: 2019

Version: Accepted version (Final draft)

Copyright: © 2019 Scandinavian Association for the Study of Pain

Rights: In Copyright

Rights url: http://rightsstatements.org/page/InC/1.0/?language=en

Please cite the original version:

Pirnes, K., Kallio, J., Siekkinen, K., Hakonen, H., Häkkinen, A., & Tammelin, T. (2019). Test-retest repeatability of questionnaire for pain symptoms for school children aged 10–15 years. Scandinavian Journal of Pain, 19(3), 575-582. https://doi.org/10.1515/sjpain-2018-0338

Table 1. The prevalence of pain experienced once a week or more often in different body areas at first survey, for all children and separately for children in primary and lower secondary school.

Pain area	All (n=181)		Primary school (n=77)		Secondary school (n=104)	
	N	%	N	%	N	%
Head	60	29	13	16	47	37
Neck and shoulder	48	23	10	13	38	30
Upper extremities	17	8	4	5	13	10
Chest	12	6	3	4	9	7
Upper back	26	13	8	10	18	14
Lower back	27	13	9	11	18	14
Stomach	25	12	7	9	18	14
Buttocks	17	8	6	8	11	9
Lower extremities	26	13	7	9	19	15

Table 2. The test-retest reliability of the pain questions measured two weeks apart in all children, and separately for children in primary and lower secondary school. Trauma includes reported injury at first, second or both surveys.

Pain Area	N	Trauma N	ICC (5-point scale) Kappa (95% CI)	2-point scale Kappa (95% CI)	2-point scale Agreement (% same)
All					
Head	161	20	0.66 (0.56, 0.74)	0.70 (0.58, 0.83)	88
Neck and shoulder	168	13	0.67 (0.58, 0.75)	0.68 (0.55, 0.82)	89
Upper extremities	146	35	0.36 (0.21, 0.50)	0.41 (0.14, 0.68)	92
Chest	177	4	0.36 (0.23, 0.48)	0.35 (0.03, 0.66)	94
Upper back	170	11	0.40 (0.26, 0.52)	0.33 (0.10, 0.56)	89
Lower back	167	14	0.40 (0.26, 0.52)	0.36 (0.15, 0.58)	87
Stomach	174	7	0.49 (0.37, 0.59)	0.44 (0.23, 0.66)	90
Buttocks	173	8	0.38 (0.25, 0.50)	0.21 (-0.03, 0.46)	90
Lower extremities	109	72	0.62 (0.48, 0.72)	0.60 (0.33, 0.87)	94
Primary School					
Head	64	13	0.56 (0.36, 0.71)	0.66 (0.39, 0.93)	92
Neck and shoulder	71	6	0.59 (0.42, 0.72)	0.67 (0.40, 0.94)	93
Upper extremities	56	21	0.21 (-0.05, 0.45)	0.30 (-0.21, 0.80)	93
Chest	74	3	0.28 (0.06, 0.48)	0.49 (-0.13, 1.10)	97
Upper back	71	6	0.38 (0.16, 0.56)	0.47 (0.03, 0.91)	94
Lower back	73	4	0.58 (0.40, 0.71)	0.67 (0.40, 0.94)	93
Stomach	73	4	0.30 (0.08, 0.49)	0.36 (-0.04, 0.76)	92
Buttocks	73	4	0.44 (0.24, 0.61)	0.47 (0.03, 0.91)	95
Lower extremities	43	34	0.70 (0.51, 0.83)	0.79 (0.39, 1.19)	98
Secondary School					
Head	97	7	0.69 (0.57, 0.78)	0.69 (0.54, 0.84)	86
Neck and shoulder	97	7	0.70 (0.58, 0.79)	0.67 (0.51, 0.83)	87
Upper extremities	90	14	0.39 (0.20, 0.55)	0.45 (0.13, 0.77)	91
Chest	103	1	0.37 (0.19, 0.52)	0.29 (-0.07, 0.65)	92
Upper back	99	5	0.39 (0.21, 0.55)	0.26 (-0.01, 0.53)	85
Lower back	94	10	0.23 (0.03, 0.41)	0.11 (-0.14, 0.36)	83
Stomach	101	3	0.59 (0.44, 0.70)	0.47 (0.22, 0.73)	88
Buttocks	100	4	0.36 (0.18, 0.52)	0.07 (-0.17, 0.31)	87
Lower extremities	66	38	0.57 (0.38, 0.71)	0.52 (0.18, 0.86)	91