

**This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.**

**Author(s):** Suurnäkki, Suvi; Pulkkinen, Jani T.; Lindholm-Lehto, Petra C.; Tiirola, Marja; Aalto, Sanni L.

**Title:** The effect of peracetic acid on microbial community, water quality, nitrification and rainbow trout (*Oncorhynchus mykiss*) performance in recirculating aquaculture systems

**Year:** 2020

**Version:** Accepted version (Final draft)

**Copyright:** © 2019 Elsevier B.V.

**Rights:** CC BY-NC-ND 4.0

**Rights url:** <https://creativecommons.org/licenses/by-nc-nd/4.0/>

**Please cite the original version:**

Suurnäkki, S., Pulkkinen, J. T., Lindholm-Lehto, P. C., Tiirola, M., & Aalto, S. L. (2020). The effect of peracetic acid on microbial community, water quality, nitrification and rainbow trout (*Oncorhynchus mykiss*) performance in recirculating aquaculture systems. *Aquaculture*, 516, Article 734534. <https://doi.org/10.1016/j.aquaculture.2019.734534>

# Journal Pre-proof

The effect of peracetic acid on microbial community, water quality, nitrification and rainbow trout (*Oncorhynchus mykiss*) performance in recirculating aquaculture systems

Suvi Suurnäkki, Jani T. Pulkkinen, Petra C. Lindholm-Lehto, Marja Tiirola, Sanni L. Aalto

PII: S0044-8486(19)31591-1

DOI: <https://doi.org/10.1016/j.aquaculture.2019.734534>

Reference: AQUA 734534

To appear in: *Aquaculture*

Received Date: 26 June 2019


Revised Date: 20 September 2019

Accepted Date: 22 September 2019

Please cite this article as: Suurnäkki, S., Pulkkinen, J.T., Lindholm-Lehto, P.C., Tiirola, M., Aalto, S.L., The effect of peracetic acid on microbial community, water quality, nitrification and rainbow trout (*Oncorhynchus mykiss*) performance in recirculating aquaculture systems, *Aquaculture* (2019), doi: <https://doi.org/10.1016/j.aquaculture.2019.734534>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2019 Published by Elsevier B.V.



1 **The effect of peracetic acid on microbial community, water quality, nitrification and rainbow trout**  
2 **(*Oncorhynchus mykiss*) performance in recirculating aquaculture systems**

3 Suvi Suurnäkki<sup>1</sup>, Jani T. Pulkkinen<sup>1,2\*</sup>, Petra C. Lindholm-Lehto<sup>2</sup>, Marja Tiirola<sup>1</sup>, Sanni L. Aalto<sup>1,3</sup>

4 <sup>1</sup> Department of Biological and Environmental Science, University of Jyväskylä P.O. Box 35, Jyväskylä,  
5 Finland

6 <sup>2</sup> Natural Resources Institute Finland, Survontie 9A, 40500 Jyväskylä, Finland

7 <sup>3</sup> Department of Environmental and Biological Sciences, University of Eastern Finland, P.O. Box 1627,  
8 Kuopio, Finland

9 \*corresponding author: jani.t.pulkkinen@luke.fi, +358 29532 3297

10 **Highlights**

- 11 - The microbial community in biofilters was not affected by frequent peracetic acid (PAA)  
12 applications
- 13 - The highest PAA application frequency improved the water quality by decreasing total ammonium  
14 nitrogen values
- 15 - PAA applications disrupted nitrification, but the nitrifying community was capable of adapting
- 16 - Fish growth and feed conversion ratio were not significantly affected by the PAA application

17 **Abstract**

18 Microbial biofilters control water quality and enable the overall function of recirculation aquaculture  
19 systems (RAS). Changes in environmental conditions can affect the abundance and interactions of  
20 the diverse microbial populations of the biofilter, affecting nitrification of harmful ammonium and  
21 thus fish health. Here, we examined the effect of different application frequencies (0, 1, 2 and 4  
22 times per week) of a common disinfectant, peracetic acid (PAA, applied 1.1 mg l<sup>-1</sup> twice per day), on

23 biofilter microbial communities, focusing especially on nitrifying microbial groups and using a high  
24 throughput sequencing of 16S rRNA gene and quantitative PCR (qPCR). In addition, we measured  
25 biofilter nitrification rates, water quality parameters, and fish performance. Although PAA additions  
26 did not significantly change the overall microbial community composition or abundance, the  
27 abundance of ammonia-oxidizing bacteria (AOB) and nitrate-oxidizing bacteria (NOB) first decreased  
28 at the beginning of the experiment but increased in numbers towards the end of the experiment  
29 with frequent PAA applications. PAA application decreased the nitrification rate, but increased the  
30 water quality in terms of reduced ammonium levels. PAA application did not significantly affect fish  
31 growth, but higher mortality was observed with the highest PAA application level of 4 times per  
32 week. These results suggest that when applied before the fish tank, PAA can be used for temporary  
33 water quality improvement without disturbing microbial communities. However, the application  
34 frequency required for persistent water quality improvement caused increased mortality.

35 *Keywords:* 16S rRNA; biofiltration; comammox; nitrification genes; peracetic acid

## 36 **1. Introduction**

37 Aquaculture is one of the fastest growing food producing sectors (FAO, 2017). Recirculating  
38 aquaculture systems (RAS) have lower water consumption and lower nutrient discharge to the  
39 environment than the traditional flow-through or net pen farming, making them environmentally  
40 and economically sustainable fish-producing systems (Badiola et al., 2012). However, water quality  
41 management is crucial for successful RAS performance and fish health, where the bacterial  
42 community plays an important role (Blancheton et al., 2013; Rurangwa and Verdegem, 2015).

43 Biofilters play a key role in controlling water quality, hosting microbes that convert the produced  
44 ammonium into nitrate nitrogen through nitrification. Traditionally, the nitrification pathway is  
45 considered to consist of two steps driven by two different microbial groups: ammonia oxidizers  
46 (ammonia-oxidizing bacteria: AOB); and nitrite oxidizers (nitrite-oxidizing bacteria: NOB). Recently,  
47 ammonia-oxidizing archaea (AOA) have also been found in biofilters in RAS and aquariums (Bagchi et

48 al., 2014; Bartelme et al., 2017). The nitrification function of the biofilter depends on several water  
49 quality factors, e.g. pH, oxygen concentration, and ammonium levels (Chen et al., 2006). Nitrite-  
50 oxidizing NOBs are especially sensitive (Villaverde et al., 1997; Graham et al., 2007), which can cause  
51 nitrite accumulation after changes in the RAS operating conditions. However, recently, a group of  
52 NOB capable of complete ammonia oxidation (comammox), *Nitrospira*, has been identified and  
53 found to be quite abundant in RAS biofilters (Bartelme et al., 2017). In addition to nitrifying  
54 microbes, biofilters host a diverse heterotrophic microbial community (Bartelme et al., 2017; Rud et  
55 al., 2017), that can include harmful micro-organisms (Guttman and van Rijn, 2008; Schrader and  
56 Summerfelt, 2010) or microbes producing odorous metabolites (Lukassen et al., 2017; Lindholm-  
57 Lehto et al., 2019). Typically, organic matter accumulation (high C:N) in RAS favors heterotrophs,  
58 limiting oxygen availability and resulting in lower nitrification rates (Michaud et al., 2006).

59 Peracetic acid (PAA:  $\text{CH}_3\text{CO}_3\text{H}$ ) is a widely-used and efficient disinfecting agent in aquaculture and  
60 wastewater treatment plants (Kitis, 2004; Koivunen and Heinonen-Tanski, 2005; Lahnsteiner and  
61 Weismann, 2007). In aquaculture, PAA has been used for disinfection against pathogenic bacteria  
62 and other harmful micro-organisms, e.g. crayfish plague spores (Jussila et al., 2011), and for general  
63 disinfection to reduce bacterial and potentially bacterial by-product numbers (Kitis, 2004; Liu et al.,  
64 2017a). PAA has an optimal degradation time (2 mg L<sup>-1</sup> to 0 mg L<sup>-1</sup> in 6 hours) for RAS, and it has no  
65 toxic or harmful by-products (Pedersen et al., 2009, Liu et al., 2017a). The commercial PAA products,  
66 used in this and in previous studies (Pedersen et al., 2009, Pedersen et al., 2013, Liu et al., 2017a,  
67 Davidson et al., 2019), contain microbial disinfectant PAA, and hydrogen peroxide, which degrades  
68 the organic material from fish feces and feed, and acetic acid. Previous studies have shown that  
69 regular PAA additions of 1 mg L<sup>-1</sup> can decrease aerobic bacterial density in the rearing water (Liu et  
70 al., 2018), but in lower PAA concentrations of 0.1-0.3 mg L<sup>-1</sup>, bacteria remained unaffected (Davidson  
71 et al., 2019). Pulse applications remove bacterial biofilm in the tank wall more efficiently than  
72 continuous PAA applications (Liu et al., 2017a). A disadvantage of PAA is its tendency to increase  
73 organic matter content in the system due to acetic acid, causing regrowth of microbes and PAA

74 decays (Kitis, 2004; Pedersen et al., 2013). Indeed, continuous low concentration PAA applications  
75 can promote biofilm formation and microbial adaptation (Liu et al., 2017a).

76 Another challenge in PAA application is the potential disturbance to the crucial biofilter function.  
77 The PAA concentration cannot be too high, since a previous study has shown elevated nitrite levels,  
78 i.e. disturbed nitrification with 2 mg L<sup>-1</sup> and 3 mg L<sup>-1</sup> of PAA used, while this was not observed with 1  
79 mg L<sup>-1</sup> of PAA (Pedersen et al., 2009). Furthermore, semi-continuous PAA applications (0.3 mg L<sup>-1</sup>)  
80 should not affect nitrification negatively (Davidson et al., 2019). In addition to nitrification, fish  
81 welfare should be considered when using PAA or any other disinfectant or substance for water  
82 quality management in RAS. Fish tolerance to PAA depends on the species (Straus et al., 2018).  
83 Rainbow trout (*Oncorhynchus mykiss*) are sensitive to PAA, as the no-observed-effect concentration  
84 (NOEC) is 2.8 mg L<sup>-1</sup>, compared with the more tolerant channel catfish (*Ictalurus punctatus*) (NOEC  
85 4.0 mg L<sup>-1</sup>) and blue tilapia (*Oreochromis aureus*) (NOEC 5.8 mg L<sup>-1</sup>) (Straus et al., 2018). Exposed to  
86 PAA (up to 2 mg L<sup>-1</sup>), fish have shown stress responses measured by elevated cortisol levels in  
87 plasma (Gesto et al., 2018) and in rearing water (Liu et al., 2017b). In addition, increased mucus  
88 formation has been reported at similar PAA concentrations (Lindholm-Lehto et al., 2019). However,  
89 when PAA exposure has been continued, fish have been able to adapt during the treatment (Gesto  
90 et al., 2018; Liu et al., 2017b).

91 Although there are some previous studies of PAA and its effects in RAS, they have focused on  
92 general water quality and measured nitrification rates (Pedersen et al., 2009; Pedersen et al., 2013;  
93 Liu et al., 2017a; Davidson et al., 2019); little is known about the effects of disinfecting agents on the  
94 microbial communities in the RAS biofilter. Since PAA is widely used in RAS, this information is  
95 required to understand the stability of the biofilter function during pulse PAA applications and to  
96 improve the overall performance of RAS. In this study, the effect of three different pulse application  
97 frequencies of PAA (1.1 mg L<sup>-1</sup> applied twice per day) were studied with a control group. PAA were  
98 applied 1, 2 and 4 times per week to the pump sump on a replicated laboratory-scale RAS. We

99 studied the overall biofilter microbiological community composition, assessed with high throughput  
100 sequencing, and the total bacterial abundance and genetic nitrification potential (number of  
101 nitrification genes), assessed with quantitative PCR (qPCR). In addition, biofilter nitrification rates,  
102 water quality parameters, and the fish growth and feed conversion ratio were examined. We  
103 hypothesized that the highest application of PAA 4 times per week might have detrimental effects  
104 on microbial communities by disturbing the steady-state microbial community and thus disrupting  
105 nitrification. In addition, lower applications of PAA once a week were expected to improve water  
106 quality and decrease the microbial load in the fish tank, further leading to improved fish  
107 performance.

## 108 **2. Materials and methods**

### 109 **2.1 Experimental setup**

110 The experiment was performed at the Natural Resources Institute Finland (Luke) Laukaa fish farm,  
111 using an experimental RAS platform. The details of the research facility are described in more depth  
112 in Pulkkinen et al. (2018). Briefly, eight individual recirculating systems were used, each consisting of  
113 a 500 L bottom-drained plastic rearing tank, a feed collector unit, a 24 cm swirl separator (Eco-Trap  
114 Collector1, Pentair Aquatic Eco-Systems, Minneapolis, USA), a drum filter with 60  $\mu\text{m}$  filter panels  
115 (Hydrotech HDF501, Veolia, Paris, France), a 147 L fixed bed bioreactor (filled with 80 L of RK  
116 Biolements heavy, RK Plast A/S, Skive, Denmark), a trickling filter acting as a forced-ventilated  
117 cascade aeration column (Bio-Blok<sup>®</sup> 200, EXPO-NET Danmark A/S, Hjørring, Denmark), and a pump  
118 sump (Fig. 1). All RAS units had been used in previous experiments, and the biofilters were stabilized  
119 to full maturity. Water pH was adjusted to 7.2 in the pump sump with 20% NaOH (aq), using an  
120 automated system (Prominent, Heidelberg, Germany). Oxygen saturation was kept above 80% in the  
121 fish tanks, and water temperature around 16 °C. The relative water renewal rate was set to 500 L kg<sup>-1</sup>  
122 <sup>1</sup> feed, and the tank hydraulic retention time to 33 min.

123 In this thirteen week experiment, three different PAA application frequencies per week were used  
124 (1.1 mg PAA L<sup>-1</sup> in a fish tank dosed twice per day). The experiment is described in more detail in  
125 Lindholm-Lehto et al. (2019). Briefly, control tanks and three different PAA applications per week (1,  
126 2 and 4) applied in pump sumps were used in replicate tanks. The PAA applications were dosed by  
127 adding 4 mL of PAA solutions twice a day. A commercial PAA product was used, composed of 12-13%  
128 PAA, 19-23% hydrogen peroxide, and the remainder acetic acid (Bonsoxo 2901, Bang & Bonsomer,  
129 Helsinki, Finland).

## 130 **2.2 Fish and feeding**

131 Three weeks before PAA addition commenced, a total of 400 one-year-old rainbow trout  
132 (*Oncorhynchus mykiss*) (average weight 130 g) originating from the National JALO selective breeding  
133 programme (Natural Resources Institute Finland, Tervo, Finland) was divided into the eight RAS  
134 units. During the first two weeks after PAA addition commenced, mortality between 0-20% was  
135 observed in the RAS units, and fish were diagnosed with IPNV (Infectious Pancreatic Necrosis Virus),  
136 which may have caused the mortality. After three weeks, fish in poor condition were removed, and  
137 biomasses were equalized between the tanks (mean 12.5 kg m<sup>-3</sup>). Fish were weighed once during the  
138 experiment in week 8 by weighing the tank mass and counting the individuals. Feeding was evenly  
139 executed with a commercial feeding system (T Drum 2000, Arvo-Tec, Joroinen, Finland) 10-14 times  
140 per day in constant light. A 1:1 mixture of two commercial diets was used, Raisioaqua (Circuit Red 5  
141 mm, Raisio, Finland) and BioMar (Orbit 929 4.5 mm, Aarhus, Denmark). The crude protein and lipid  
142 contents of the diets were 43% and 42%, and 26% and 31%, respectively, as given by the  
143 manufacturer. Feed intake was monitored using a feed collector unit, and feeding was decreased if  
144 uneaten feed was observed. At the start, the feeding rate was set to 1.5 % bw d<sup>-1</sup>.

## 145 **2.3 Sampling for microbiome and qPCR analysis**

146 Biofilter biofilm samples were collected for sequencing and qPCR twice after 8 and 13 weeks of the  
147 experiment. Three sets of carrier media were collected from each bioreactor unit. In addition, water



148 and biofilm samples from the fish tank were collected for qPCR. Water samples, two per fish tank,  
149 were filtered using syringe filters (0.22  $\mu\text{m}$  Millipore Express<sup>®</sup> PLUS PES membrane). The tank biofilm  
150 was collected using detachable tank material blocks (HDPE), which were placed in the tanks at the  
151 beginning of the experiment. Prior to further analysis and DNA extraction, samples were frozen (- 20  
152  $^{\circ}\text{C}$ ) for at least 24 hours. To detach microbial biofilm from the biofilter and tank biofilm samples, 20  
153 mL of water was added to the samples, and they were sonicated for 4 min (Branson 1510). They  
154 were then freeze-dried (Alpha 1-4 LD plus, Christ). DNA extraction was performed for the freeze-  
155 dried materials using the DNeasy PowerLyzer<sup>™</sup>PowerSoil DNA Isolation Kit (Qiagen) in accordance  
156 with the manufacturer's instructions. PowerLyzer Homogenizer was applied once at 3,400 rpm for  
157 45 s during the extraction. The quantity of extracted DNA was measured with Qubit<sup>™</sup> dsDNA HS  
158 assay and Qubit 2.0 Fluorometer (Thermo Fischer Scientific). For sequencing and qPCR, three  
159 replicate samples from each biofilter unit or two water samples were pooled.

#### 160 **2.4 Microbial communities**

161 The microbial community composition was studied using Ion Torrent PGM next generation  
162 sequencing, targeting the V4 region of the 16S rRNA gene with primer pair 515F-Y  
163 (GTGYCAGCMGCCGCGGTAA; Parada et al., 2016) and 806R (GGACTACHVGGGTWTCTAAT; Caporaso  
164 et al., 2011). The 25  $\mu\text{l}$  PCR reaction consisted of Maxima SYBR Green/Fluorescein qPCR Master Mix  
165 (Thermo Fisher Scientific), 10 ng of template DNA, and 0.4  $\mu\text{M}$  of both primers. Thermal cycling  
166 consisted of 10 min initial denaturation at 95  $^{\circ}\text{C}$ , followed by 40 cycles at 95  $^{\circ}\text{C}$  for 30 s, 50  $^{\circ}\text{C}$  for 30 s  
167 and 72  $^{\circ}\text{C}$  for 60 s, followed by final elongation at 72  $^{\circ}\text{C}$  for 5 min. To add Ion Torrent PGM  
168 sequencing adapters and barcodes to the ends of the PCR product, one  $\mu\text{l}$  of the PCR product was  
169 used as a template in the second qPCR, where 10 cycles were performed using linker and fusion  
170 primers (0.04  $\mu\text{M}$  of M13\_515F-Y, 0.4  $\mu\text{M}$  of IonA\_IonXpressBarcode\_M13 and P1\_806R), with  
171 conditions otherwise identical to the first amplification (Mäki et al., 2016). Products were purified  
172 with the Agencourt AMPure XP purification system (Beckman Coulter Life Sciences, Indianapolis, IN,  
173 USA), quantified with Qubit<sup>™</sup> dsDNA HS assay, and pooled in equimolar quantities for sequencing on

174 Ion Torrent PGM using Ion PGM Hi-Q View OT2 Kit for emulsion PCR, PGM Hi-Q View Sequencing Kit  
175 for the sequencing reaction, and Ion 316 Chip v2 (all Life Sciences, Thermo Fisher Scientific). A 16S  
176 rRNA gene sequence analysis was done using mothur (version 1.39.5; Schloss et al., 2009), as in  
177 Aalto et al. (2018). The total number of sequences obtained was 294,464 and after subsampling,  
178 there were 13,245 sequences per sample. The sequences have been submitted to the NCBI  
179 Sequence Read Archive under BioProject PRJNA549384.

## 180 **2.5 Gene copy numbers and nitrification rates**

181 The abundance of 16S rRNA gene in the biofilter biofilm, tank water, and biofilm, as well as the  
182 nitrification genes in the biofilter biofilm, was quantified with qPCR. For the 16S rRNA gene, the  
183 primer pair 515F-Y and 806R was used and with the amplification protocol details mentioned above.

184 To quantify the abundance of the ammonia-oxidizing bacteria (AOB), nitrite-oxidizing bacteria (NOB),  
185 and complete ammonia-oxidizing (comammox) bacteria in the biofilter samples, qPCR  
186 quantifications were performed using published primer pairs for AOA, AOB, NOB, and comammox.  
187 The abundance of AOA was quantified using amoA<sub>archaea</sub> primers (Francis et al., 2005), and the  
188 abundance of AOB using amoA<sub>bac</sub> primers (Rotthauwe et al., 1997), with the protocols described by  
189 Aalto et al. (2018). For NOB abundances, the primers used were *nxrAF1/R2* (Poly et al., 2008; Wertz  
190 et al., 2008) and *nxB169F/638R* (Pester et al., 2014). Comammox *Nitrospira* clade A and clade B  
191 abundances were quantified with the comaA244F/659R and comaB224F/659R primers (Pjevac et al.,  
192 2017). All qPCR reactions included 10 ng of template DNA, forward and reverse primers, and 1x  
193 Maxima SYBR Green/Fluorescein Master Mix (Thermo Fischer) in a total volume of 25  $\mu$ l. The  
194 thermal conditions were as follows: initial denaturation 10 min at 95 °C, then 35 cycles at 95 °C for  
195 30 s, 52-59 °C for 30 s, and 72 °C for 30 s. Amplification efficiencies were between 83-97% for the  
196 qPCR assays. The quantification was performed using the CFX96 qPCR thermal cycler (Bio-Rad).

197 Bioreactor nitrification rates ( $\text{g NO}_x \text{ h}^{-1}$ ) were measured following Pulkkinen et al. (2018) at the end  
198 of the experiment. Briefly, 30 biofilter carrier media pieces were incubated with  $^{15}\text{NH}_4^+$  of 5  $\text{mg L}^{-1}$  for

199 3 hours, and the stable isotope composition of nitrite and nitrate was measured at the beginning  
200 and end of the incubation.

## 201 **2.6 Water quality measurements**

202 Water quality was monitored with an online monitoring system consisting of a spectrometer probe  
203 (spectro::lyser, s::can, Vienna, Austria), a carbon dioxide sensor (Franatech, Lüneburg, Germany), a  
204 pH probe (pH::lyser, s::can, Vienna, Austria) and an optical oxygen probe (oxi::lyser, s::can, Vienna,  
205 Austria) located in the fish tanks. The spectrometer probe measured turbidity, total suspended  
206 solids, and UV254 absorbance. Total ammonia nitrogen (TAN), nitrite, and nitrate were analyzed  
207 once a week from tank outlet water using a spectrophotometer (Procedure 8038 Nessler,  
208 LCK341/342 and LCK340 respectively, DS 3900, Hach, Loveland, USA). Alkalinity was analyzed once a  
209 week by titration with a standard method (ISO 9963-1:1994) (TitraLab AT1000, Hach, Loveland,  
210 USA).

## 211 **2.7 Statistical analyses**

212 The effects of PAA applications on water quality values, feed conversion ratio (FCR), specific growth  
213 rate (SGR), mortality and gene copy numbers and microbial community abundances were analyzed  
214 with linear regression analysis using SPSS Statistics software, version 25. Gene abundances were log-  
215 transformed before analysis to meet assumptions. The changes in the microbial community  
216 composition were assessed using the “vegan” (Oksanen et al., 2017) and “phyloseq” (McMurdie and  
217 Holmes 2013) packages in R (version 3.5.1).

## 218 **3. Results**

### 219 **3.1 Biofilter microbial community**

220 Based on the NMDS plot, the microbial community composition changed from week 8 to week 13 in  
221 all tanks. At week 8, communities in control units and those receiving four PAA applications per  
222 week were more similar than those with one or two PAA applications, which were more dispersed,

223 while at week 13, the communities were quite similar in all PAA application levels (Fig. 2). The  
224 dominant phylum across all samples was Proteobacteria (28-44% of sequences) (Fig. 3). The next  
225 most abundant OTUs were from phyla Gemmatimonadetes (5-19% of sequences) and Bacteroidetes  
226 (20-45% of sequences). Although the abundance of proteobacterial classes Deltaproteobacteria (8%  
227 vs 6-7% of all sequences) and Gammaproteobacteria (12% vs 7-10% of all sequences) decreased  
228 slightly with an increased PAA application rate (Fig. 3), the overall abundance of Proteobacteria was  
229 unaffected by the PAA application frequency (linear regression,  $P>0.05$ ). The species richness  
230 (Chao1) increased from week 8 to week 13 except in the other one-time PAA application per week  
231 unit. The diversity generally increased from week 8 to 13, and was higher in control units than in  
232 units with PAA applications (Suppl. Fig. 1).

233 When only nitrifying taxonomic groups were considered (Fig. 3/Suppl. Fig. 2), the number of  
234 sequences assigned to AOB (family Nitrosomonadaceae) appeared to decrease with the increasing  
235 PAA application frequencies at week 8 (117-235 reads, 0.7-1.3% of all sequences), but the difference  
236 was not statistically significant ( $P=0.052$ ). At week 13, the amount of AOB reads was very similar  
237 between PAA application frequencies (95-244 reads, 0.5-1.4% of all sequences;  $P=0.703$ ). The main  
238 genera were unclassified Nitrosomonadaceae (weeks 8, 13; 26-74% of AOB reads) and *Nitrosomonas*  
239 (week 13; 13-42% of AOB sequences). The number of sequences assigned to NOB/comammox  
240 (genera *Nitrospira*, *Nitrobacter*) was 0.4-3.5% of all sequences (79-623 reads) at week 8, while at  
241 week 13, it was 1.6-6.9% (284-1,232 reads). The PAA application frequency did not affect the  
242 number of NOB/comammox sequences ( $P>0.05$ ). *Nitrospira* was the dominant genus in all samples  
243 (90-100% of NOB sequences).

### 244 3.2 Microbial abundance and nitrification rates

245 The total amount of bacteria (16S rRNA gene copy numbers) in tank water was equal between the  
246 control and different PAA application frequencies at both weeks ( $P>0.05$ ). At week 8, the total  
247 amount of bacteria was  $2.7\pm 2.2\times 10^9$  gene copies  $\text{cm}^{-2}$  in tank biofilm and  $7.8\pm 6.3\times 10^{11}$  gene copies  $\text{g}^{-1}$

248 <sup>1</sup> dw in biofilters, while at week 13, there were  $1.4 \pm 1.5 \times 10^9$  gene copies  $\text{cm}^{-2}$  in tank biofilm and  
249  $9.4 \pm 5.9 \times 10^{10}$  gene copies  $\text{g}^{-1}$  dw in biofilters (Fig. 4). There was no significant relationship between  
250 total microbial abundances and PAA application frequencies neither in tank nor in biofilter biofilms.  
251 In biofilter samples, the copy numbers of nitrification genes were normalized against the total  
252 bacterial abundance. The relative abundance of the AOB gene was higher than the NOB or  
253 comammox genes, and ammonia-oxidizing archaea were not found in the samples. All the  
254 comammox *Nitrospira* were from clade A, while no comammox bacteria from clade B were observed  
255 in the samples. At week 8, the relative gene copy numbers of AOB and NOB appeared to decrease  
256 with an increasing PAA application rate, but the relationship was not significant ( $P=0.092$ ) for AOB  
257 and for NOB. The abundance of comammox bacteria remained similar in all systems ( $P>0.05$ ). At  
258 week 13, at the end of the experiment, the relative abundance of all three nitrifying groups was  
259 higher than at week 8. The AOB copy numbers were equal between the units, while NOB abundance  
260 was highest in the units receiving bi-weekly PAA application, and comammox abundance in the  
261 control units and those receiving PAA four times per week (Fig. 5). Nitrification rates ( $\text{g NO}_3 \text{ h}^{-1}$   
262  $\text{bioreactor}^{-1}$ ) were highest in tanks with one PAA application per week (Fig. 6). With PAA applications  
263 of two and four times per week, the nitrification rate decreased and was lowest, at  $0.09 \text{ g NO}_3 \text{ h}^{-1}$   
264  $\text{bioreactor}^{-1}$ , in tanks with four PAA applications per week (Fig. 6).

### 265 **3.3 Water quality and fish performance**

266 The turbidity and total suspended solids were lower in the 2 x PAA and 4 x PAA application rates but  
267 there was no significant relationship with PAA application frequency (Table 2, Suppl. Fig. 3;  $P=0.057$ ).  
268 Higher PAA application frequency significantly decreased TAN concentrations during the first part of  
269 the experiment ( $P<0.01$ , weeks 3-8) and during the whole experiment ( $P=0.026$ , weeks 3-13; Table  
270 2). PAA additions did not significantly increase  $\text{NO}_2\text{-N}$  concentrations ( $P>0.05$ ), and  $\text{NO}_3\text{-N}$   
271 concentrations were similar between the RAS units ( $P>0.05$ , Table 1). In addition, the PAA  
272 application frequencies did not significantly affect fish performance when measured by FCR and SGR

273 (Table 1), but the fish in the units receiving PAA four times per week had a 15% slower growth  
274 compared to fish in the control groups. However, mortality increased with increasing PAA  
275 application frequencies ( $P < 0.01$ , Table 1; Table 2).

#### 276 **4. Discussion**

277 In this study, we demonstrated the effect of different PAA application frequencies on biofilter  
278 microbiome function and composition, especially on the key nitrification microbes. We also  
279 examined the changes in biofilter nitrification rates, water quality, and fish performance in response  
280 to PAA application frequency. We observed that the overall microbial community composition  
281 remained quite stable, and nitrification bacteria did not substantially suffer from PAA applications,  
282 increasing in abundance during the experiment. More frequent PAA application decreased biofilter  
283 nitrification rates but yet could decrease the TAN values. PAA application frequencies did not  
284 significantly affect fish growth, but a higher mortality rate and increased slime formation were  
285 observed with the highest PAA application (four times per week).

286 Until this study, knowledge of the response of the biofilter microbiome to PAA applications has been  
287 scarce. In a few previous studies, a focus on the effect of a bi-weekly pulsed PAA addition of  $1 \text{ mg L}^{-1}$   
288 on RAS microbiology has been found to decrease the overall bacterial counts in the tank water and  
289 to nearly completely remove the biofilm in the fish tanks in RAS (Liu et al., 2017a, 2018), while  
290 enhanced biofilm formation has been observed under continuous PAA applications, which has been  
291 explained by the acetic acid and formed acetate feeding the heterotrophic microbial community (Liu  
292 et al., 2017a). Furthermore, when a prolonged low PAA dosage has been used, no effect on  
293 microbial counts has been observed, resulting in a minimum PAA threshold estimate of  $0.30 \text{ mg L}^{-1}$   
294 for disinfectant purposes in RAS (Davidson et al., 2019). In agreement, we observed that the  
295 abundance of tank water microbes, either free-living or attached to particles, was unaffected by a  
296 PAA addition. However, when weeks were inspected separately, we observed that bacterial  
297 abundance was higher, yet statistically insignificant, in the units receiving a PAA application one or

298 four times per week in both tank and biofilter biofilms at week 8. In the tanks at week 13, the PAA  
299 application rate had no overall effect on bacterial biofilm, because the bi-weekly PAA application  
300 rate was sufficient to decrease the bacterial biofilm, but the highest PAA application seemed to  
301 promote biofilm formation. In the biofilters, PAA had no clear effect on the biofilm bacterial  
302 abundance. It is likely that most of the PAA used was degraded before entering the biofilters, since it  
303 was added to the pump sump, explaining the minor beneficial and restricting effects of the PAA on  
304 the biofilm in the biofilters than in the tanks.

305 The dominating phylum was Proteobacteria in all RAS units throughout the experiment, while  
306 Bacteroidetes and Gemmatimonadetes were also relatively abundant. Previously, Proteobacteria  
307 and Bacteroidetes have found to be common phyla in RAS biofilter units in both freshwater and  
308 saltwater RAS, while Gemmatimonadetes were either absent or found in low abundance (Ruan et  
309 al., 2015; Gonzales-Silva et al., 2016; Bartelme et al., 2017; Rud et al., 2017), highlighting the unique  
310 nature of microbiomes in each biofilter and RAS unit (Blancheton et al., 2013). In previous studies,  
311 *Nitrospira* has been found to be quite abundant in RAS biofilters (Bartelme et al., 2017; Keuter et al.,  
312 2017), the group including both traditional nitrite-oxidizers and comammox bacteria. Here, we found  
313 the amount of *Nitrospira* increasing from week 8 to week 13 in all units. Overall, PAA application  
314 frequency did not significantly change the biofilm bacterial community composition or decrease the  
315 diversity or species richness, agreeing with previous findings on perturbations affecting free-living  
316 microbes rather than deep biofilm layers (Wietz et al., 2009; Schreier et al., 2010).

317 When nitrifying microbes are the sole focus, both qPCR and sequencing results demonstrated that  
318 PAA application frequency had a slightly, yet statistically insignificant, negative effect on the nitrifier  
319 abundance at week 8. At week 13, genetic nitrification potential (gene copy numbers) and the  
320 sequences associated with NOBs/comammox increased compared to week 8 in all units. This  
321 indicates that the possibly negative effect of PAA application on the nitrification microbes decreased  
322 towards the end of the experiments, probably because the nitrification community could adapt and

323 become less sensitive to PAA application. Previously, PAA additions have been found to only partially  
324 inhibit nitrification (Pedersen et al., 2009, Liu et al., 2017a), causing some accumulation of total  
325 ammonia nitrogen (TAN) and nitrate in RAS. Here, we measured nitrification rates using stable  
326 isotope incubations at the end of the experiment, and found higher PAA application frequency led to  
327 lower nitrification rates. However, we did not find the previously observed nitrite accumulation,  
328 suggesting that both ammonium oxidation and nitrite oxidation steps were suppressed, or that the  
329 abundant comammox bacteria were as or even more involved in ammonia oxidation than the  
330 traditional ammonia-oxidizers, releasing only nitrate as the end product.

331 In addition to changes in nitrification, the increased PAA application rate led to decreased turbidity  
332 and total suspended solids concentrations except in units with a PAA addition of twice a week,  
333 where values were closer to control units than other PAA application units. Decreased turbidity  
334 indicates that there were fewer large particles (Yao et al., 2014) because of the higher  $H_2O_2$   
335 degradation potential of organic matter. The observed decrease of total ammonia nitrogen with an  
336 increasing PAA addition was unexpected, because nitrification efficiency was lowest within the  
337 highest PAA addition. The direct oxidation of TAN is the likeliest explanation, because the slightly  
338 different daily feed consumption did not explain the difference in TAN concentrations between  
339 treatments. TAN values have previously been found to increase after a bi-weekly PAA addition due  
340 to lower nitrification rates (Liu et al., 2017a), and we also observed that the units with bi-weekly PAA  
341 applications behaved quite differently from those receiving one or four applications. There, we saw  
342 equal turbidity, TSS, TOC, TAN, and nitrate values with the control units, suggesting that this bi-  
343 weekly PAA addition promoted heterotrophic microbes more than removing them or improving  
344 water quality. However, one reason for the difference between bi-weekly and other PAA application  
345 frequencies is that the other 2x RAS unit (unit 2) showed a very different pattern in water quality  
346 and microbial results than the other systems. Furthermore, we saw that AOB and comammox  
347 microbes were less abundant in these systems, while the NOBs seemed to benefit there. In this  
348 experiment, PAA dosages were kept the same throughout the experiment, and feeding was



349 increased from the first (mean 102 g d<sup>-1</sup>) to the second (mean 127 g d<sup>-1</sup>) part of the experiment.  
350 Thus, the potential PAA effect time decreased during the experiment (Pedersen et al., 2013), which  
351 explained why the water quality differences between the treatments were higher in the first part of  
352 the experiment.

353 The PAA additions did not significantly affect fish performance in terms of feed conversion ratio or  
354 specific growth rate. However, the total biomass growth was 14% lower in the units where PAA was  
355 added four times per week compared to other units. In addition, mortality increased with the  
356 highest PAA addition, indicating that continued applications of PAA (1.1 mg L<sup>-1</sup> dosed twice per day),  
357 even below the reported no-observed-effect concentration (2.8 mg L<sup>-1</sup>) (Straus et al., 2018), were  
358 too high for the fish. High surface swimming was observed with the first additions of PAA, but this  
359 was not observed after a few weeks of additions, indicating the adaptation of fish to PAA as  
360 previously reported (Liu et al., 2017b; Gesto et al., 2018). The observed IPNV has low pathogenicity  
361 in Finland and mainly affects the first-feeding fry (Eriksson-Kallio et al., 2016). We thus concluded  
362 that it did not affect fish performance.

## 363 5. Conclusions

364 In conclusion, the microbial results suggest that the biofilter biofilm community is quite stable and  
365 less sensitive to PAA application, forming a “collaborome” in which heterotrophic microbes can  
366 support autotrophic nitrifiers rather than compete with them (Bartelme et al., 2017). Pulsed PAA  
367 applications disrupt nitrification, but the microbial community is capable of adapting and no long-  
368 term effect of PAA on inorganic nitrogen levels can be observed. Our study demonstrates that PAA  
369 application frequency has variable effects on microbes, water quality, and fish. Although the highest  
370 PAA application frequency (1.1 mg L<sup>-1</sup> twice a day, four times per week) improved water quality  
371 slightly by directly oxidizing TAN, and potentially turbidity, and TSS, it led to 50% lower biofilter  
372 nitrification rates compared to the control units, and increased fish mortality. On the other hand, a  
373 bi-weekly application did not improve water quality, but fish performance was better than with

374 other PAA application frequencies. Similarly, a one-time PAA application was too low to improve  
375 water quality, but did not interrupt nitrification or fish. The opposite chemical (i.e. direct oxidizing of  
376 TAN) and biological (i.e. decreasing nitrification thus increasing TAN) effects of PAA on water quality  
377 can complicate the interpretation of results. Based on these results, the continuous pulse  
378 applications of PAA are not a cost-efficient method for substantially improving water quality and  
379 controlling the microbial communities. However, if an accumulation of solids and/or total ammonia  
380 nitrogen is observed, PAA can be used in such cases to improve water quality.

### 381 **Acknowledgements**

382 We greatly appreciate the fish husbandry and water sampling of the Natural Resources Institute  
383 Finland Laukaa fish farm personnel. We gratefully acknowledge Lars-Flemming Pedersen for his  
384 helpful information about PAA dosages. This experiment was funded by the European Union through  
385 the European Maritime and Fisheries Fund (EMFF) and by the Ministry of Agriculture and Forestry of  
386 Finland.

### 387 **Author contributions**

388 All the authors planned the experiment. SS, JTP, and PL-L did the sampling and SS performed the  
389 DNA sequencing and qPCR measurements. SLA performed the statistical analyses of the microbial  
390 data, and JTP the water quality and fish performance analysis. All the authors contributed to  
391 manuscript writing.

392 **References**

- 393 Aalto, S.L, Saarenheimo, J., Mikkonen, A., Rissanen, A.J., Tirola, M., 2018. Resistant ammonia-  
394 oxidizing archaea endure, but adapting ammonia-oxidizing bacteria thrive in boreal lake sediments  
395 receiving nutrient-rich effluents. *Environ. Microbiol.* 20 (10), doi: 10.1111/1462-2920.14354.
- 396 Badiola, M., Mendiola, D., Bostock, J., 2012. Recirculating Aquaculture Systems (RAS) analysis: main  
397 issues on management and future challenges. *Aquacul. Eng.* 51, 26-35. doi:  
398 10.1016/j.aquaeng.2012.07.004
- 399 Bagchi, S., Vlaeminck, S. E., Sauder, L. A., Mosquera, M., Neufeld, J. D., Boon, N., 2014. Temporal and  
400 spatial stability of ammonia-oxidizing archaea and bacteria in aquarium biofilters. *PLoS One* 9(12),  
401 e113515. doi: 10.1371/journal.pone.0113515.
- 402 Bartelme, R.P., McLellan, S.L., Newton, R.J., 2017. Freshwater recirculating aquaculture system  
403 operations drive biofilter bacterial community shifts around a stable nitrifying consortium of  
404 ammonia-oxidizing *Archaea* and comammox *Nitrospira*. *Front. Microbiol.* 8, 101. doi:  
405 10.3389/fmic.2017.0010.
- 406 Blancheton, J.P., Attramadal, K.J.K., Michaud, L. d'Orbcastel, E.R., Vadstein, O., 2013. Insights into  
407 bacterial population in aquaculture systems and its implications. *Aquacult. Eng.* 53, 30-39.  
408 doi:10.1016/j.aquaeng.2012.11.009.
- 409 Caporaso, J.G., Lauber, C.L., Walters, W.A., Berg-Lyons, D., Lozupone, C.A., Turnbaugh, P.J, Fierer, N.,  
410 Knight, R., 2011. Global patterns of 16S rRNA diversity at a depth of millions of sequences per  
411 sample. *PNAS*, 108: 4516-4522. doi:10.1073/pnas.1000080107.
- 412 Chen, S., Ling, J., Blancheton, J.-P., 2006. Nitrification kinetics of biofilm as affected by water quality  
413 factors. *Aquacul. Eng.* 34(3), 179-197. doi: 10.1016/j.aquaeng.2005.09.004.
- 414 Davidson, J., Summerfelt, S., Straus, D.L., Schrader, K.K., Good, C., 2019. Evaluating the effects of  
415 prolonged peracetic acid dosing in water quality and rainbow trout *Oncorhynchus mykiss*

- 416 performance in recirculation aquaculture systems. *Aquacult. Eng.* 84, 117-127.  
417 doi:10.1016/j.aquaeng.2018.12.009.
- 418 Eriksson-Kallio, A.M., Holopainen, R., Viljamaa-Dirks, S, Venneström, P., Kuukka-Anttila, H., Koski, P.,  
419 Gadd, T. 2016. Infectious pancreatic necrosis virus (IPNV) strain with genetic properties associated  
420 with low pathogenicity at Finnish fish farms. *Dis. Aquat. Org.* 118: 21-30.
- 421 FAO, 2017. Food and Agriculture Organization (FAO). Fisheries and Aquaculture Statistics; FAO Year  
422 Book; FAO: Rome, Italy.
- 423 Francis, C.A., Roberts, K. J., Beman, J.M., Santoro, A.E., Oakley, B. B., 2005. Ubiquity and diversity of  
424 ammonia-oxidizing archaea in water columns and sediments of the ocean. *Proc. Natl. Acad. Sci. USA*  
425 102: 14683-14688. doi: 10.1073/pnas.0506625102.
- 426 Gesto, M., Liu, D., Pedersen, L-F., Meinelt, T., Straus, D.L., Jokumsen, A., 2018. Confirmation that  
427 pulse and continuous peracetic acid administration does not disrupt the acute stress response in  
428 rainbow trout. *Aquaculture* 492, 190-194. doi: 10.1016/j.aquaculture.2018.04.009.
- 429 Gonzalez-Silva, B.M., Jonassen, K.R., Bakke, I., Østgaard, K., Vadstein, O., 2016. Nitrification at  
430 different salinities: biofilm community composition and physiological plasticity. *Water Res.* 95, 48-  
431 58. doi: 10.1016/j.watres.2016.02.050.
- 432 Graham, D.W., Knapp, C.W., Van Vleck, E.S., Bloor, K., Lane, T.B., Graham, C. E., 2007. Experimental  
433 demonstration of chaotic instability in biological nitrification. *ISME J.* 1(5), 385. doi:  
434 10.1038/ismej.2007.45.
- 435 Guttman, L., van Rijn, J., 2008. Identification of conditions underlying production of geosmin and 2-  
436 methylisoborneol in a recirculating system. *Aquaculture* 279, 85-91. doi:  
437 10.1016/j.aquaculture.2008.03.047.

- 438 Jussila, J., Makkonen, J., Kokko H., 2011. Peracetic acid (PAA) treatment is an effective disinfectant  
439 against crayfish plague (*Aphanomyces astaci*) spores in aquaculture. *Aquaculture* 320, 37-42. doi:  
440 10.1016/j.aquaculture.2011.08.008.
- 441 Keuter, S., Beth, S., Quantz, G., Schulz, C., Spieck, E., 2017. Longterm monitoring of nitrification and  
442 nitrifying communities during biofilter activation of two marine recirculation aquaculture systems  
443 (RAS). *Int. J. Aquac. Fish Sci.* 3, 51-61. doi:10.17352/2455-8400.000029.
- 444 Kitis, M., 2004. Disinfection of wastewater with peracetic acid: a review. *Environ. Int.* 30, 47-55.  
445 doi:10.1016/S0160-4120(03)00147-8.
- 446 Koivunen, J., Heinonen-Tanski, H., 2005. Inactivation of enteric microorganisms with chemical  
447 disinfectants, UV irradiation and combined chemical/UV treatments. *Water Res.* 39 (8), 1519-1526.  
448 doi: 10.1016/j.watres.2005.01.021
- 449 Lahnsteiner, F., Weismann, T., 2007. Treatment of *Ichthyophthiriasis* in rainbow trout and common  
450 carp and alternative therapeutics. *J. Aquat. Anim. Health* 19, 186-194. doi: 10.1577/H07-002.1.
- 451 Lindholm-Lehto, P.C., Suurnäkki, S., Pulkkinen, J.T., Aalto, S.L., Tirola, M., Vielma, J., 2019. Effect of  
452 peracetic acid on levels of geosmin, 2-methylisoborneol, and their potential producers in a  
453 recirculating aquaculture system for rearing rainbow trout (*Oncorhynchus mykiss*). *Aquacult. Eng.* 85,  
454 56-64. doi:10.1016/j.aquaeng.2019.02.002.
- 455 Liu, D., Straus, D.L., Pedersen, L-F., Meinelt, T., 2017a. Pulse versus continuous peracetic acid  
456 applications: Effects on rainbow trout performance, biofilm formation and water quality. *Aquacult.*  
457 *Eng.* 77, 72-79. doi:10.1016/j.aquaeng.2017.03.004.
- 458 Liu, D., Pedersen, L-F., Straus, D.L., Kloas, W., Meinelt, T., 2017b. Alternative prophylaxis/disinfection  
459 in aquaculture – adaptable stress induced by peracetic acid at low concentration and its application  
460 strategy in RAS. *Aquaculture* 474, 82-85. doi: 10.1016/j.aquaculture.2017.03.027.

- 461 Liu, D., Straus, D.L, Pedersen, L-F., Meinelt, T., 2018. Periodic bacterial growth with peracetic acid in  
462 a recirculating aquaculture system and its long-term effect on fish health. *Aquaculture* 485, 154-159.  
463 doi:10.1016/j.aquaculture.2017.11.050.
- 464 Lukassen, M. B., Saunders, A. M., Sindilariu, P. D., Nielsen, J. L., 2017. Quantification of novel  
465 geosmin-producing bacteria in aquaculture systems. *Aquaculture* 479, 304-310. doi:  
466 10.1016/j.aquaculture.2017.06.004.
- 467 McMurdie, P.J., Holmes, S., 2013. phyloseq: An R Package for reproducible interactive analysis and  
468 graphics of microbiome census data. PLoS ONE. 8(4):e61217. doi: 10.1397./journal.pone.0061217.
- 469 Michaud, L., Blancheton, J. P., Bruni, V., Piedrahita, R., 2006. Effect of particulate organic carbon on  
470 heterotrophic bacterial populations and nitrification efficiency in biological filters. *Aquacult. Eng.* 34,  
471 224-233. doi: 10.1016/j.aquaeng.2005.07.005.
- 472 Mäki, A., Rissanen, A.J., Tirola, M., 2016. A practical method for barcoding and size-trimming PCR  
473 templates for amplicon sequencing. *Biotechniques* 60, 88-90. doi: 10.2144/000114380.
- 474 Oksanen, J., Blanchet, F.G., Kindt, R., Legendre, P., Minchin, P.R., O'hara, R.B., Simpson, G.L.,  
475 Solymos, P., Stevens, M.H.H., Wagner, H., 2017. vegan: Community Ecology Package. R package  
476 version 2.3–0. 2015.
- 477 Parada, A.E., Needham, D.M., Fuhrman, J.A., 2016. Every base matters: assessing small subunit rRNA  
478 primers for marine microbiomes with mock communities, time series and global field samples.  
479 *Environ. Microbiol.* 18, 1403-1414. doi: 10.1111/1462-2920.13023.
- 480 Pedersen, L-F., Pedersen, P.B., Nielsen, J.L., Nielsen, P.H., 2009. Peracetic acid degradation and  
481 effects on nitrification in recirculating aquaculture systems. *Aquaculture* 296, 246-254. doi:  
482 10.1016/j.aquaculture.2009.08.021.

- 483 Pedersen, L-F., Meinelt, T., Straus, D.L., 2013. Peracetic acid degradation in freshwater aquaculture  
484 systems and possible practical implications. *Aquacult. Eng.* 53, 65-71. doi:  
485 10.1016/j.aquaeng.2012.11.011.
- 486 Pester, M., Maixner, F., Berry, D., Rattei, T., Koch, H., Lückner, S., Nowka, B., Richter, A., Spieck, E.,  
487 Lebedeva, E., Loy, A., Wagner, M., Daims, H., 2014. *NxrB* encoding the beta subunit of nitrite  
488 oxidoreductase as functional and phylogenetic marker for nitrite-oxidizing *Nitrospira*. *Environ.*  
489 *Microbiol.* 16, 3055-3071. doi:10.1111/1462-2920.12300.
- 490 Pjevac, P., Schauburger, C., Poghosyan, L., Herbold, C.W., van Kessel, M.A.H.J., Daebeler, A.,  
491 Steinberger, M., Jetten, M.S.M, Lückner, S., Wagner, M., Daims, H., 2017. *AmoA*-targeted polymerase  
492 chain reaction primers for the specific detection and quantification of comammox *Nitrospira* in the  
493 environment. *Front. Microbiol.* 8, 1-11. doi:10.3389/fmicb.2017.01508.
- 494 Poly, F., Wertz, S., Brothier, E., Degrange, V., 2008. First exploration of *Nitrobacteria* diversity in soils  
495 by a PCR cloning-sequencing approach targeting functional gene *nxrA*. *FEMS Microbiol. Ecol.* 63, 132-  
496 140. doi:10.1111/j.1574-6941.2007.00404.x.
- 497 Pulkkinen, J., Kiuru, T., Aalto, S.L., Koskela, J., Vielma, J., 2018. Startup and effects of relative water  
498 renewal rate on water quality and growth of rainbow trout (*Oncorhynchus mykiss*) in a unique RAS  
499 research platform. *Aquacult. Eng.* 82, 38-45. doi: 10.1016/j.aquaeng.2018.06.003.
- 500 R Core Team, 2018. R: A language and environment for statistical computing. R Foundation for  
501 Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.
- 502 Rotthauwe, J. H., Witzel, K. P., Liesack, W., 1997. The ammonia monooxygenase structural gene  
503 *amoA* as a functional marker: molecular fine-scale analysis of natural ammonia-oxidizing  
504 populations. *Appl. Environ. Microbiol.* 63, 4704-4712.

- 505 Ruan, Y.J., Guo, X-S., Ye, Z-Y., Liu, Y., Zhu, S-M., 2015. Bacterial community analysis of different  
506 sections of a biofilter in full-scale marine recirculating aquaculture system. *N. Am. J. Aquacult.* 77,  
507 318-326. doi: 10.1080/15222055.2015.1017128.
- 508 Rud, I., Kolaveric, J., Holan, A.B., Berget, I., Calabrese, S., Terjesen, B.F., 2017. Deep-sequencing of  
509 the bacterial microbiota in commercial-scale recirculating and semi-closed aquaculture systems for  
510 Atlantic salmon post-smolt production. *Aquacult. Eng.* 78, 50-62. doi:  
511 10.1016/j.aquaeng.2016.10.003.
- 512 Rurangwa, E., Verdegem, M.C.J., 2015. Microorganisms in recirculating aquaculture systems and  
513 their management. *Rev. Aquacult.* 7, 117-130. doi: 10.1111/raq.12057.
- 514 Schloss, P.D., Wescott, S.L., Ryabin, T., Hall, J.R., Hartmann, M., Hollister, E.B., Lesniewski, R.A.,  
515 Oakley, B.B., Parks, D.H., Robinson, C.J., Sahl, J.W., Stres, B. Thallinger, G.G., Van Horn, D.J., Weber,  
516 C.F., 2009. Introducing mother: open-source, platform-independent, community-supported software  
517 for describing and comparing microbial communities. *Appl. Environ. Microbiol.* 75, 7537-7541. doi:  
518 10.1128/AEM.01541-09.
- 519 Schrader, K.K., Summerfelt, S.T., 2010. Distribution of off-flavor compounds and isolation of  
520 geosmin-producing bacteria in a series of water recirculating systems of Rainbow Trout culture. *N.*  
521 *Am. J. Aquacult.* 72 (1), 1-9. doi:10.1577/A09-009.1.
- 522 Schreier, H.J., Mirzoyan, N., Saito, K., 2010. Microbial diversity of biological filters in recirculating  
523 aquaculture systems. *Curr. Opin. Biotechnol.* 21, 318-325. doi: 10.1016/j.copbio.2010.03.011.
- 524 Straus, D.L., Meinelt, T., Liu, D., Pedersen, L-F., 2018. Toxicity of peracetic acid to fish: variation  
525 among species and impact of water chemistry. *J. World Aquacult. Soc.* 49, 715-724. doi:  
526 10.1111/jwas.12475.
- 527 Villaverde, S., Garcia-Encina, P. A., Fdz-Polanco, F., 1997. Influence of pH over nitrifying biofilm  
528 activity in submerged biofilters. *Water Res.* 31, 1180-1186. doi: 10.1016/S0043-1354(96)00376-4.



- 529 Wertz, S., Poly, F., Le Roux, X., Degrange, V. 2008. Development and application of a PCR-denaturing  
530 gradient gel electrophoresis tool to study the diversity of *Nitrobacter*-like nxrA sequences in soil.  
531 *FEMS Microbiol. Ecol.* 63, 261-271. Doi: 10.1111/j.1574-6941.2007.00416.x.
- 532 Wietz, M., Hall, M. R., Høj, L., 2009. Effects of seawater ozonation on biofilm development in  
533 aquaculture tanks. *Syst. Appl. Microbiol.* 32, 266-277. doi:10.1016/j.syamo.2009.04.001.
- 534 Yao, M., Nan, J., Chen, T., 2014. Effect of particle size distribution on turbidity under various water  
535 quality levels during flocculation processes. *Desalination*, 354, 116-124.

536 **Figures and tables**537 **Figure legends**

538 Figure 1. Schematic diagram of one RAS unit used in the experiment. Microbial samples were taken  
539 from the fish tank biofilm, water from fish tank and from the fixed bed bioreactor. PAA was dosed  
540 into the pump sump. FT = Fish tank, 1 = Swirl separator, 2 = drum filter, FBBR = Fixed bed bioreactor,  
541 TF = Trickling filter, 3 = Pump sump.

542 Figure 2. Non-metric multidimensional scaling (NMDS) of bacterial communities in RAS biofilters  
543 under different PAA application frequencies (control, 1, 2, and 4 applications per week) at weeks 8  
544 and 13.

545 Figure 3. The relative abundances of microbial phyla in biofilter biofilms under different PAA  
546 application frequencies (control, 1, 2, and 4 applications per week) at weeks 8 and 13.

547 Figure 4. The abundance of bacterial 16S rRNA gene in tank water, in tank biofilm and in biofilter  
548 biofilm under different PAA application frequencies (control, 1, 2, and 4 applications per week).  
549 Abundance denotes gene copies ml<sup>-1</sup> water for tank water samples, gene copies cm<sup>-2</sup> for tank biofilm  
550 samples, and gene copies g<sup>-1</sup> dw for biofilter biofilm samples.


551 Figure 5. The relative abundances of AOB (ammonia-oxidizing bacteria), NOB (nitrite-oxidizing  
552 bacteria) and comammox clade A genes in biofilter under different PAA application frequencies  
553 (control, 1, 2, and 4 applications per week).

554 Figure 6. The biofilter nitrification rates under different PAA application frequencies (control, 1, 2,  
555 and 4 applications per week) at week 13 at the end of the experiment.


556 **Table legends**

557 Table 1. The mean water quality parameters and observed fish performance under different PAA  
 558 application treatments (0, 1, 2, and 4 applications per week, PAA dosed  $1.1 \text{ mg L}^{-1}$  twice per day)  $\pm$   
 559 SD. FCR = Feed conversion ratio, SGR = Specific growth rate.

560 Table 2. The linear regression models on the interactions between water quality parameters and  
 561 different PAA application frequencies (0, 1, 2 and 4 applications per week). Data is shown, when  $p <$   
 562 0.1. TSS = total suspended solids, TAN = total ammonia nitrogen.




563  
 564 Figure 1. Schematic diagram of one RAS unit used in the experiment. Microbial samples were taken  
 565 from the fish tank biofilm, water from fish tank and from the fixed bed bioreactor. PAA was dosed  
 566 into the pump sump. FT = Fish tank, 1 = Swirl separator, 2 = drum filter, FBBR = Fixed bed bioreactor,  
 567 TF = Trickling filter, 3 = Pump sump.



568


569 Figure 2. Non-metric multidimensional scaling (NMDS) of bacterial communities in RAS biofilters  
570 under different PAA application frequencies (control, 1, 2, and 4 applications per week) at weeks 8  
571 and 13.



572


573 Figure 3. The relative abundances of microbial phyla in biofilter biofilms under different PAA

574 application frequencies (control, 1, 2, and 4 applications per week) at weeks 8 and 13.



575

576 Figure 4. The abundance of the bacterial 16S rRNA gene in tank water, in tank biofilm and in biofilter  
 577 biofilm under different PAA application frequencies (control, 1, 2, and 4 applications per week).  
 578 Abundance denotes gene copies  $\text{ml}^{-1}$  water for tank water samples, gene copies  $\text{cm}^{-2}$  for tank biofilm  
 579 samples, and gene copies  $\text{g}^{-1}$  dw for biofilter biofilm samples.




580

581 Figure 5. The relative abundances of AOB (ammonia-oxidizing bacteria), NOB (nitrite-oxidizing

582 bacteria), and comammox (clade A) genes in biofilter under different PAA application frequencies

583 (control, 1, 2, and 4 applications per week).



584

585 Figure 6. The biofilter nitrification rates under different PAA application frequencies (control, 1, 2,

586 and 4 applications per week) at week 13 at the end of the experiment.



587 Table 1. The mean water quality parameters and observed fish performance under different PAA application treatments (0, 1, 2, and 4 applications per  
 588 week, PAA dosed 1.1 mg L<sup>-1</sup> twice per day) ± SD. FCR = Feed conversion ratio, SGR = Specific growth rate.

589

Treatment	Mean of weeks 3-7 (±SD)				Mean of weeks 8-13 (±SD)				Mean of weeks 3-13 (±SD)			
	0	1	2	4	0	1	2	4	0	1	2	4
CO <sub>2</sub> (mg l <sup>-1</sup> )	8.9 ± 0.9	8.7 ± 0.4	7.4 ± 1.5	8.9 ± 0.5	11.0 ± 1.8	9.8 ± 0.6	8.9 ± 2.0	10.6 ± 0.8	9.8 ± 1.3	9.2 ± 0.5	8.0 ± 1.6	9.6 ± 0.8
UV254 (Abs m <sup>-1</sup> )	29.7 ± 4.2	26.5 ± 2.4	30.2 ± 1.1	24.0 ± 0.1	29.6 ± 5.7	27.4 ± 2.4	30.6 ± 0.2	26.4 ± 0.9	29.7 ± 4.8	26.9 ± 0.4	30.2 ± 0.9	25.0 ± 0.9
Turbidity (FTU)	2.0 ± 0.3	1.4 ± 0.2	1.8 ± 0.1	1.1 ± 0.3	1.8 ± 0.2	1.4 ± 0.1	1.9 ± 0.0	1.2 ± 0.1	1.9 ± 0.2	1.4 ± 0.1	1.9 ± 0.0	1.2 ± 0.1
TSS (mg l <sup>-1</sup> )	5.8 ± 0.4	3.9 ± 0.6	5.1 ± 0.5	3.0 ± 1.0	5.2 ± 0.3	3.9 ± 0.5	5.2 ± 0.1	3.1 ± 0.7	5.5 ± 0.4	3.9 ± 0.2	5.2 ± 0.2	3.0 ± 0.7
TAN (mg l <sup>-1</sup> )	0.93 ± 0.11	0.78 ± 0.08	0.78 ± 0.01	0.57 ± 0.01	0.80 ± 0.1	0.70 ± 0.15	0.80 ± 0.05	0.61 ± 0.04	0.83 ± 0.09	0.75 ± 0.05	0.78 ± 0.04	0.58 ± 0.04
NO <sub>2</sub> -N (mg l <sup>-1</sup> )	0.11 ± 0.01	0.12 ± 0.02	0.10 ± 0.01	0.10 ± 0.1	0.13 ± 0.02	0.17 ± 0.04	0.13 ± 0.04	0.10 ± 0.02	0.12 ± 0.01	0.15 ± 0.02	0.12 ± 0.03	0.10 ± 0.02
NO <sub>3</sub> -N (mg l <sup>-1</sup> )	54.6 ± 0.6	52.1 ± 4.2	56.8 ± 4.0	48.1 ± 4.6	60.2 ± 2.3	61.1 ± 9.7	65.5 ± 4.5	54.8 ± 3.5	55.5 ± 1.5	55.4 ± 7.4	57.5 ± 1.8	48.7 ± 3.5
Alkalinity (mg l <sup>-1</sup> )	62.3 ± 1.3	57.2 ± 2.3	42.7 ± 14.5	71.5 ± 2.6	63.1 ± 6.6	51.7 ± 1.6	45.9 ± 17.5	69.4 ± 3.3	62.1 ± 3.9	53.0 ± 1.4	44.3 ± 16.1	68.5 ± 3.3
pH	7.3 ± 0.0	7.3 ± 0.1	7.1 ± 0.1	7.3 ± 0.0	7.3 ± 0.1	7.3 ± 0.1	7.1 ± 0.1	7.4 ± 0.0	7.3 ± 0.0	7.2 ± 0.1	7.1 ± 0.1	7.4 ± 0.0
FCR	1.11 ± 0.1	0.92 ± 0.14	1.01 ± 0.14	1.21 ± 0.0	1.01 ± 0.05	1.12 ± 0.11	1.08 ± 0.01	1.11 ± 0.02	1.05 ± 0.02	1.02 ± 0.12	1.04 ± 0.07	1.16 ± 0.02
SGR (% d <sup>-1</sup> )	1.35 ± 0.18	1.32 ± 0.14	1.45 ± 0.06	1.28 ± 0.03	1.08 ± 0.05	1.04 ± 0.04	1.06 ± 0.0	1.03 ± 0.03	1.23 ± 0.07	1.19 ± 0.09	1.27 ± 0.04	1.17 ± 0.03
End weight (g)	303 ± 32	286 ± 22	308 ± 33	286 ± 9	428 ± 39	401 ± 36	431 ± 46	384 ± 12	428 ± 39	401 ± 36	431 ± 46	384 ± 12
Mortality (%)	2.9 ± 0.0	2.9 ± 2.9	2.9 ± 2.9	7.1 ± 1.4	2.9 ± 0.0	4.4 ± 1.3	4.4 ± 1.3	4.6 ± 0.0	5.7 ± 0.0	7.1 ± 1.4	7.1 ± 1.4	11.4 ± 0.0

590 Table 2. The linear regression models on the interactions between water quality parameters and  
 591 different PAA application frequencies (0, 1, 2, and 4 applications per week). Data is shown when  $p <$   
 592 0.1. TSS = total suspended solids, TAN = total ammonia nitrogen.

Variable	Week	Equation	R <sup>2</sup>	p value
Turbidity	3-8	$y = -2.465 x + 5.724$	0.46	0.064
TSS	3-8	$y = -0.800 x + 5.291$	0.46	0.067
TAN	3-8	$y = -8.709 x + 8.412$	0.73	0.007
TSS	8-13	$y = -1.008 x + 6.136$	0.42	0.079
Turbidity	3-13	$y = -2.900 x + 6.390$	0.48	0.057
TSS	3-13	$y = -0.945 x + 5.921$	0.48	0.057
TAN	3-13	$y = -10.370 x + 9.398$	0.59	0.026
Mortality	3-13	$y = 58.33 x - 2.917$	0.70	0.010

593

Conflict of Interest

The authors declare that there is no conflict of interest

Journal Pre-proof