

This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Ravantti, Janne J.; Laanto, Elina; Papponen, Petri; Sundberg, Lotta-Riina

Title: Complete Genome Sequence of Fish Pathogen *Flavobacterium columnare* Strain B185, Originating from Finland

Year: 2019

Version: Published version

Copyright: © 2019 Ravantti et al.

Rights: CC BY 4.0



Rights url: <https://creativecommons.org/licenses/by/4.0/>

Please cite the original version:

Ravantti, Janne J. ; Laanto, Elina; Papponen, Petri; Sundberg, Lotta-Riina (2019). Complete Genome Sequence of Fish Pathogen *Flavobacterium columnare* Strain B185, Originating from Finland. *Microbiology Resource Announcements*, 8 (49), e01285-19. DOI: 10.1128/MRA.01285-19



Complete Genome Sequence of Fish Pathogen *Flavobacterium columnare* Strain B185, Originating from Finland

 Janne J. Ravantti,^{a,b}  Elina Laanto,^{a,c} Petri Papponen,^c  Lotta-Riina Sundberg^c

^aMolecular and Integrative Biosciences Research Programme, Faculty of Biological and Environmental Sciences, University of Helsinki, Helsinki, Finland

^bApplied Tumor Genomics Research Program & Department of Medical Genetics, University of Helsinki, Helsinki, Finland

^cDepartment of Biological and Environmental Science and Nanoscience Center, University of Jyväskylä, Jyväskylä, Finland

ABSTRACT We report a complete genome sequence of a Finnish isolate of the fish pathogen *Flavobacterium columnare*. Using PacBio RS II sequencing technology, the complete circular genome of *F. columnare* strain B185 with 3,261,404 bp was obtained.

Flavobacterium columnare (*Bacteroidetes*) is an opportunistic fish pathogen affecting freshwater fish farming worldwide. As of January 2019, fewer than 20 genomes of *F. columnare* were available, and 5 were complete. We report a whole-genome sequence of *F. columnare* strain B185, isolated from Finland. *F. columnare* isolates have been traditionally divided into different genetic groups, and recently, four groups were assigned (1). All described Finnish isolates belonging to the genetic group I and have been further divided into five genotypes (A to G) (2), and strain B185 represents genotype G. Strain B185 has been used in several studies and confirmed as virulent (3, 4). This genome sequence will aid in resolving the genetics of virulence in *F. columnare* and in deciphering the genetic diversity and evolutionary origins of Finnish *F. columnare* strains.

Flavobacterium columnare strain B185 was isolated in 2008 from tank water at a fish farm in Central Finland during antibiotic treatment in a quarantine unit. The water sample was plated onto Shieh agar, and an *F. columnare* colony was isolated, purified, and further cultivated in Shieh liquid medium and stored at -80°C . Genomic DNA was extracted from an overnight (O/N) culture using a DNeasy blood and tissue kit (Qiagen) according to the manufacturer's instructions for Gram-negative cells. DNA was sequenced at the Institute of Biotechnology, University of Helsinki, Helsinki, Finland, using PacBio RS II technology with P4-C2 sequencing chemistry, generating 268,103 reads and an average read length of 3,962 bp with an N_{50} value of 6,100 bp. After the genome was assembled with the HGAP3 (SMRT portal) (5), a single contig remained. The Gap4 (6) program was then used to confirm that both ends contained overlapping sequence, indicating a circular genome. The overlaps were merged, and finally, the genome was rotated so that the starting point was at the gene *dnaA*. The final assembly yielded one scaffold with an average sequence coverage of 224.6 \times . The genome of *F. columnare* strain B185 consists of 3,261,404 bp with a GC content of 31.7%. No plasmids were detected. A total of 2,812 coding DNA sequences (CDSs) were predicted using NCBI's Prokaryotic Genome Annotation Pipeline (7). In all the analyses, default parameters were used except where otherwise noted. The genome contains 3 rRNA operons and 92 tRNA genes. Amino acid-level similarities (global alignment per translated open reading frames [ORFs] with 80% similarity cutoff) to other complete *F. columnare* genomes from genomic group I were 92% to strain TC 1691, 91% to strain Pf1, and 90% to strain ATCC 49512 and from genomic group II were 81% to strain C#2 and 80% to

Citation Ravantti JJ, Laanto E, Papponen P, Sundberg L-R. 2019. Complete genome sequence of fish pathogen *Flavobacterium columnare* strain B185, originating from Finland. *Microbiol Resour Announc* 8: e01285-19. <https://doi.org/10.1128/MRA.01285-19>.

Editor Julie C. Dunning Hotopp, University of Maryland School of Medicine

Copyright © 2019 Ravantti et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Lotta-Riina Sundberg, lotta-riina.sundberg@jyu.fi.

Received 11 October 2019

Accepted 7 November 2019

Published 5 December 2019

strain 94-081. Two functional CRISPR loci (type II-C and type VI-B) were identified earlier in an *F. columnare* genome (8) and are also present in strain B185. Interestingly, the II-C locus has a putative parDE type II toxin-antitoxin (TA) system between the *cas9* and the repeat region. A putative prophage of approximately 30 kbp in length was detected with HHPred (9) analysis for regions of annotated hypothetical proteins. Gene modules, e.g., for head and tail morphogenesis, were identified. In addition, we identified a putative ~40-kb genetic island rich in integrases and transposases.

Data availability. The complete genome sequence of *Flavobacterium columnare* strain B185 has been deposited in GenBank under the accession number [NZ_CP010992](https://ncbi.nlm.nih.gov/nucl/NZ_CP010992), and the raw sequence data were submitted to the SRA under accession number [PRJNA276168](https://www.ncbi.nlm.nih.gov/sra/PRJNA276168).

ACKNOWLEDGMENTS

This work was supported by Academy of Finland grant 314939 and the Jane and Aatos Erkkö Foundation. This work resulted from the BONUS Flavophage project supported by BONUS (Art 185), funded jointly by the European Union (EU) and the Academy of Finland.

We thank the DNA Sequencing and Genomics Unit (Institute of Biotechnology, University of Helsinki) for sequencing and genome assembly assistance.

REFERENCES

1. LaFrentz BR, García JC, Waldbieser GC, Evenhuis JP, Loch TP, Liles MR, Wong FS, Chang SF. 2018. Identification of four distinct phylogenetic groups in *Flavobacterium columnare* with fish host associations. *Front Microbiol* 9:452. <https://doi.org/10.3389/fmicb.2018.00452>.
2. Suomalainen L-R, Kunttu H, Valtonen ET, Hirvelä-Koski V, Tirola M. 2006. Molecular diversity and growth features of *Flavobacterium columnare* strains isolated in Finland. *Dis Aquat Organ* 70:55–61. <https://doi.org/10.3354/dao070055>.
3. Laanto E, Bamford JKH, Laakso J, Sundberg L-R. 2012. Phage-driven loss of virulence in a fish pathogenic bacterium. *PLoS One* 7:e53157. <https://doi.org/10.1371/journal.pone.0053157>.
4. Laanto E, Bamford JKH, Ravantti JJ, Sundberg L-R. 2015. The use of phage FCL-2 as an alternative to chemotherapy against columnaris disease in aquaculture. *Front Microbiol* 6:829.
5. Chin CS, Alexander DH, Marks P, Klammer AA, Drake J, Heiner C, Clum A, Copeland A, Huddleston J, Eichler EE, Turner SW, Korlach J. 2013. Non-hybrid, finished microbial genome assemblies from long-read SMRT sequencing data. *Nat Methods* 10:563–569. <https://doi.org/10.1038/nmeth.2474>.
6. Staden R, Judge DP, Bonfield JK. 2003. Managing sequencing projects in the GAP4 environment, p 327–344. *In* Krawetz SA, Womble DD (ed), *Introduction to bioinformatics: a theoretical and practical approach*. Human Press, Inc., Totowa, NJ.
7. Tatusova T, DiCuccio M, Badredin A, Chetverin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>.
8. Laanto E, Hoikkala V, Ravantti J, Sundberg L-R. 2017. Long-term genomic coevolution of host-parasite interaction in the natural environment. *Nat Commun* 8:111. <https://doi.org/10.1038/s41467-017-00158-7>.
9. Zimmermann L, Stephens A, Nam SZ, Rau D, Kübler J, Lozajic M, Gabler F, Söding J, Lupas AN, Alva VJ. 2018. A completely reimplemented MPI Bioinformatics Toolkit with a new HHPred server at its core. *Mol Biol* 50022-2836:30587–30589.