

This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Hicks, Jamie; Vasko, Petra; Goicoechea, Jose M.; Aldridge, Simon

Title: The Alumanyl Anion : A New Generation of Aluminium Nucleophile

Year: 2021

Version: Accepted version (Final draft)


Copyright: © 2020 WILEY

Rights: In Copyright

Rights url: <http://rightsstatements.org/page/InC/1.0/?language=en>

Please cite the original version:

Hicks, J., Vasko, P., Goicoechea, J. M., & Aldridge, S. (2021). The Alumanyl Anion : A New Generation of Aluminium Nucleophile. *Angewandte Chemie*, 60(4), 1702-1713.
<https://doi.org/10.1002/anie.202007530>



A Journal of the Gesellschaft Deutscher Chemiker

Angewandte Chemie

GDCh

International Edition

www.angewandte.org

Accepted Article

Title: The Aluminyl Anion: A New Generation of Aluminium Nucleophile

Authors: Jamie Hicks, Petra Vasko, Jose M. Goicoechea, and Simon Aldridge


This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). This work is currently citable by using the Digital Object Identifier (DOI) given below. The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

To be cited as: *Angew. Chem. Int. Ed.* 10.1002/anie.202007530

Link to VoR: <https://doi.org/10.1002/anie.202007530>

MINIREVIEW

The Alumanyl Anion: A New Generation of Aluminium Nucleophile

Jamie Hicks,^{[a]*} Petra Vasko,^[b] Jose M. Goicoechea^[c] and Simon Aldridge^[c]

MINIREVIEW

- [a] Dr J. Hicks
Research School of Chemistry
Australian National University
Sullivans Creek Road, Acton, 2601 (Australia)
E-mail: jamie.hicks@anu.edu.au
- [b] Dr P. Vasko
Department of Chemistry, Nanoscience Center
University of Jyväskylä
P. O. Box 35, Jyväskylä, FI-40014 (Finland)
- [c] Prof. J. M. Goicoechea, Prof. S. Aldridge
Inorganic Chemistry Laboratory, Department of Chemistry
University of Oxford
South Parks Road, Oxford, OX1 3QR (UK)

Supporting information for this article is given via a link at the end of the document.

Abstract: Trivalent aluminium compounds are well known for their reactivity as Lewis acids/electrophiles, a feature that is exploited in many pharmaceutical, industrial and laboratory-based reactions. Recently, a series of isolable aluminium(I) anions ('aluminyls') have been reported, which offer an alternative to this textbook description: these reagents behave as aluminium nucleophiles. This minireview covers the synthesis, structure and reactivity of aluminyl species reported to date, together with their associated metal complexes. The frontier orbitals of each of these species have been investigated using a common methodology to allow for a like-for-like comparison of their electronic structure and a means of rationalising (sometimes unprecedented) patterns of reactivity.

1. Introduction

Aluminium is the most abundant metal in the Earth's crust, accounting for >8% of its overall mass.¹ It is found in the sedimentary rock Bauxite, which is mined on a hundred million tonne scale annually. Bauxite is readily processed via the Bayer process to give alumina (Al₂O₃), the raw material for the production of aluminium metal.² In molecular chemistry, aluminium is most commonly found in the +3 oxidation state, with the +1 state which is favoured for the heavier Group 13 metals being strongly reducing.³ Hence, trivalent species constitute the vast majority of aluminium compounds used in chemical synthesis, such as trialkylaluminiums (typically used as alkylating reagents) and aluminium trihalides (widely used as Lewis acids).⁴ Low oxidation state aluminium compounds, i.e. those where the aluminium adopts the formal oxidation states +2, +1 or 0 are known, but are considerably less common.⁵

1.1. Aluminium(I) Complexes

Predating the work on aluminyl systems are the *neutral* aluminium compounds in the formal +1 oxidation state. The isolation of Al(I) species is challenging thermodynamically, and such species are typically highly reactive and prone to disproportionation.⁶ Binary Al(I) compounds such as AlH, AlX (X = halide) and Al₂O have been reported but are only stable for prolonged periods in the gas phase at low pressures.⁷ The development of cryochemical methods for the handling of metastable solutions of AlX (X = Cl, Br, I) proved to be instrumental in developing the chemistry of organo-aluminium(I) systems.^{7c-f} Accordingly, the first example of an isolable aluminium(I) complex - tetrameric (AlCp*)₄ **1** (Figure 1, Cp* =

C₅Me₅) - was reported in 1991 by Schnöckel and co-workers,⁸ from the reaction between a metastable solution of AlCl and MgCp*₂. An alternative synthesis using potassium metal reduction of Cp*AlCl₂ was later reported by Roesky and co-workers.⁹ In the solid state, **1** is centred around an Al₄ tetrahedron capped by four η⁵-Cp* ligands. The bulky Cp* ligands provide considerable kinetic stabilization, and contribute to its enhanced thermal tolerance compared to metastable AlX systems. In solution at elevated temperatures, **1** can undergo dissociation into monomeric AlCp* units, a factor which has been exploited in a significant part of its subsequently reported reaction chemistry.





Figure 1. The first examples of isolable aluminium(I) complexes (Cp* = C₅Me₅; Dipp = 2,6-*i*-Pr₂C₆H₃).

In 2000, Roesky and co-workers reported the first isolable monomeric aluminium(I) system, **2** (Figure 1), utilising a bulky β-diketiminato ligand.¹⁰ **2** features a two-coordinate aluminium(I) centre and can be regarded as an aluminium analogue of an NHC, with both a lone pair and empty *p*-orbital localized on the metal centre. The reactivity of these two landmark Al(I) systems (**1** and **2**) is vast and diverse and has recently been comprehensively reviewed by Ma, Yang, Roesky and co-workers.¹¹ The chemistry of **2**, in particular, is undergoing a recent renaissance, being a key reagent in a number of recent publications.¹² These recent reports include the activation of small molecules (CO, C₂H₄)^{12c,j}, reduction chemistry (including that of benzene),^{12b,e,i,k,m} cycloadditions^{12a,d,g} and various bond activations including (C–H and C–F).^{12c,f,h,l,n} The vast majority of the reactions of **1** and **2** are best described oxidative additions, although some evidence of nucleophilic behaviour has been reported (especially with electropositive metals/Lewis acids) and this has been exploited in the formation of Al–metal donor/acceptor bonds.¹¹ Since the initial reports of **1** and **2**, various other neutral aluminium(I) systems have been reported using an array of bulky ligands. These include the remarkable NHC-stabilised dialumene reported by Inoue and co-workers,¹³ the cyclic alkyl amino carbene (CAAC) supported Al(I) hydride complex¹⁴ and the monomeric ^tBu₃CpAl(I)¹⁵ reported by Braunschweig and co-workers and as well as a number of other Al(I) tetramers supported by bulky ancillary ligands.^{3,5}

MINIREVIEW


Jamie Hicks is originally from Chesterfield (UK). He received his MChem in 2011 from the University of Sheffield and his PhD in 2016 from Monash University (Prof. C. Jones). He then worked as a postdoctoral researcher at Monash University (Dr D. Turner & Prof. S. Batten) before moving to the University of Oxford to work under Profs. J. M. Goicoechea & S. Aldridge. Jamie is currently working at the Australian National University on an ARC funded DECRA fellowship.




Petra Vasko is originally from Mikkeli (Finland). She received her PhD in 2015 from the University of Jyväskylä under the joint supervision of Prof. Heikki M. Tuononen and Philip P. Power (UC Davis). In 2017, she relocated herself to Oxford to work as a postdoctoral researcher in the group of Prof. S. Aldridge. Currently, she is back at the University of Jyväskylä as an Academy of Finland funded postdoctoral researcher.




Jose Goicoechea is originally from Zaragoza (Spain) and is a Professor of Chemistry at the University of Oxford. He studied chemistry at Zaragoza (Spain) and obtained his Ph.D. from the University of Bath (Prof. M. Whittlesey). He has published over 100 papers and is a past winner of the RSC Main Group Chemistry Award (2018). His research interests include the chemistry of reactive compounds of the main-group elements for stoichiometric and catalytic bond forming reactions.



Simon Aldridge is originally from Shrewsbury (UK) and is Professor of Chemistry at the University of Oxford, and Director of the EPSRC/Oxford Centre for Doctoral Training in Inorganic Chemistry for Future Manufacturing. He has published more than 210 papers and is a past winner of the RSC's Main Group Chemistry (2010) and Frankland Awards (2018). His research interests include the development of p-block compounds with unusual electronic structure, and their applications in small molecule activation and catalysis.



was the gallyl anion, $[(^t\text{BuDAB})\text{Ga}]^-$ **3** (${}^t\text{BuDAB} = [({}^t\text{BuNCH})_2]^{2-}$) by Schmidbaur and co-workers over 20 years ago (Figure 2).¹⁷ This was followed by a diaryl substituted version reported by Jones and co-workers in 2002^{18a} and a BIAN stabilised version in 2008 by Fedushkin, Schumann and co-workers.^{18b} The reactivity of these gallyl anions have been thoroughly investigated, especially with regards to their coordination chemistry.¹⁹ However, very little has been reported with respect to small molecule activation with these systems.



Schmidbaur, *JACS* **1999** Yamashita, Nozaki, *Science* **2006** Coles, *Angew. Chem.* **2018**

Figure 2. The first reported gallyl, boryl and indyl anions (18-c-6 = 18-crown-6; crypt-222 = cryptand 222)

In 2006, Yamashita and Nozaki reported the groundbreaking boryl lithium complex, $[(^{\text{Dipp}}\text{DAB})\text{B}(\text{Li}(\text{DME}))_2]$ **4** (${}^{\text{Dipp}}\text{DAB} = [(\text{DippNCH})_2]^{2-}$), the first isolated example of a boryl anion (Figure 2).²⁰ Unlike the potassium gallyl **3**, which was initially isolated as an ion separated pair, **4** shows a strong electrostatic interaction between the boron lone pair and the lithium cation. Boryl system **4** exists as a dimer in the solid state (with two DME molecules bridging the two Li^+ cations), however a monomeric version of this complex has since been reported.²¹ To date, boryl anions featuring an array of chelating ligands have been reported,^{22,23} all of which exhibit strong interactions between the boron and the cation (either covalent or electrostatic); no charge-separated boryl anion has yet been reported.

The reactivity of the boryl anions has become an area of considerable interest over the last decade. They have been shown to be highly reactive reagents, which is partly due to their relatively small HOMO-LUMO energy gap. This defining feature allows them to activate a wide array of small molecules and/or substrates with strong σ - and π -bonds. In addition, their highly directional lone pair also makes them ideal ligands in coordination chemistry. The reactivity of boryl anions has recently been covered in a comprehensive review by Weber.²²

The heaviest member of the known group 13 element(I) anions is the indyl anion. Currently, only one example of an indyl anion is known, which was reported by Coles and co-workers in 2018.²⁴ The anion was reported as both an ion separated pair **5** (Figure 2) and as the more tightly bound lithium complex. Little reactivity of this indyl compound has yet been reported. However, preliminary studies with MeOTf and bulky metal halide complexes suggest nucleophilic behaviour,²⁴ whilst reactions with organic azides lead to oxidation of the indium.²⁵

2. Aluminyl Anions

2.1. Synthesis and Structure

Since the isolation of gallyl compound **3** in 1999, various research groups have targeted an aluminium analogue, in particular using similar diazabutadiene supporting ligands. In

1.2. Anionic Group 13 Centred Nucleophiles

Before our initial report of an anionic aluminium(I) complex (an 'aluminyl' anion)¹⁶ in 2018, analogous anions of the other group 13 elements, including boron, gallium and indium were already known in the literature. The first of these to be reported

MINIREVIEW

2002, Jones and co-workers reported the reduction of $(\text{Dipp})\text{AlI}_2$ with potassium metal.^{18a} This reaction, however, generates $(\text{Dipp})\text{AlH}_2$ and aluminium metal. The groups of Graves, Mair and Yang also reported reductions of similar diazabutadiene ligated aluminium halide complexes but no signs of an alumanyl anion were mentioned in any of these reports.²⁶

2.1.1 Diamido Alumanyl Systems

In 2018, we reported an alumanyl anion, $\text{K}_2[\text{Al}(\text{NON})]_2$, **6**, stabilised by the bulky xanthene-based diamido ligand $[\text{NON}]^{2-}$ (Figure 3).²⁷ **6** was prepared by the two-electron reduction of the neutral Al(III) precursor complex $(\text{NON})\text{AlI}$ with excess KC_8 . Interestingly, adding the reducing agent in the correct stoichiometry (i.e. 2 equiv. of KC_8) was found to give large amounts of the dialumane $[(\text{NON})\text{Al}]_2$, presumably by the radical coupling of two $(\text{NON})\text{Al}^\cdot$ units and/or by nucleophilic attack of the of the alumanyl on the $(\text{NON})\text{AlI}$ starting material (see section 2.3.). In contrast to the digallane $[(\text{tBuDAB})\text{Ga}]_2$ (which is readily reduced by potassium metal to give potassium gallyl complex **3**),¹⁷ $[(\text{NON})\text{Al}]_2$ does not react with potassium (or KC_8) even at elevated temperatures. This suggests the synthesis of **6** does not go through the dialumane as an intermediate in the reaction.




Figure 3. Structures of the diamido substituted alumanyl anion complexes 6-9.

The structure of **6** was confirmed by X-ray crystallography, which revealed the complex to exist as a dimer in the solid state. The dimer is held together by interactions between the flanking aryl groups on the NON ligands and the potassium cations. The aluminium centres are >6.6 Å apart, implying no significant interaction between them. It can also be shown by DOSY NMR experiments that the dimeric structure of **6** is retained in solution.²⁷ In **6**, there is a (relatively weak) interaction between the O atom incorporated into the backbone of the NON ligand and the aluminium centre (2.28 Å).²⁷

Subsequently, two further potassium alumanyl complexes featuring bidentate diamido ligands have been reported.²⁸ The first of these, $\text{K}_2[\text{Al}(\text{NON})]_2$ (**7**), was reported by Coles and co-workers in 2019,²⁹ and more recently $\text{K}_2[\text{Al}(\text{NCCN})]_2$ (**8**) was reported by Hill, McMullin and co-workers in 2020 (Figure 3).³⁰ Both of these complexes were synthesised using a similar synthetic route to **6**, i.e. by the two-electron reduction of the corresponding aluminium(III) iodide precursor with potassium.

Superficially, both **7** and **8** possess similar structures to **6**: both are dimeric in the solid state featuring interactions between the flanking aryl rings of the bulky diamido ligands and the potassium cations. **7** features a 6-membered aluminium-containing heterocycle, with a N-Al-N angle of $103.89(8)^\circ$,²⁹ while the 7-membered ring in **8** (not surprisingly) makes the corresponding unit somewhat wider ($108.84(9)^\circ$).³⁰ Both of these angles however are considerably smaller than that in **6** ($128.11(9)^\circ$).²⁷

To date, only two examples of truly 'naked' alumanyl anions have been reported, i.e. alumanyl anions that show no significant interaction with the cation. The first of these is the charge-separated complex $[\text{K}(\text{crypt-222})][\text{Al}(\text{NON})]$ **9** (Figure 3), reported by us in 2019.³¹ The second example is complex **11** discussed below. Complex **9** is prepared in straightforward fashion by the addition of cryptand-222 to **6** in THF. The solid-state structure of **9** confirms that the K^+ cations are encapsulated within the cryptand, and that the $[\text{Al}(\text{NON})]^-$ anion engages in no close interactions with the cation (all $\text{K}^+\cdots\text{Al}$ interactions >7 Å).

2.1.2. Alkyl Alumanyl Systems

Other than the series of diamido supported alumanyl anions discussed above, two further examples of alumanyl anions have recently been reported. These are the remarkable alkyl substituted alumanyl complexes $[\text{K}(\text{toluene})_2-\text{Al}\{\text{C}(\text{SiMe}_3)_2\text{CH}_2\}_2]$ **10** and $[\text{K}(12\text{-crown-4})][\text{Al}(\text{CAA})]$ **11** (CAA = cyclic alkyl amino), both reported earlier this year by the groups of Yamashita and Kinjo, respectively (Figure 4).^{31,32} Unlike the diamido supported alumanyl complexes **6-8**, which were all synthesised by reduction from their corresponding Al(III) iodide complexes, **10** and **11** were both synthesised by reduction of the corresponding Al-Al bonded dialumanes. This contrasts with the synthesis of **6**, for which reduction of the dialumane $[(\text{NON})\text{Al}]_2$ is not possible (at least under conditions that the alumanyl is stable).²⁷




Figure 4. Structures of alkyl substituted alumanyl complexes (12-c-4 = 12-crown-4).

The structures of **10** and **11** have been determined by X-ray crystallography and are monomeric in the solid state. Both feature 5-membered aluminium heterocycles, with similar C-Al-R angles ($R = \text{C}$, **10**; $R = \text{N}$, **11**); **11** being the slightly more acute at $86.8(1)^\circ$ compared with $90.40(5)^\circ$ for that in **10**.^{32,33} In the solid state, alumanyl **10** shows a significant interaction between the aluminium centre and the potassium cation ($\text{Al}-\text{K}$ 3.4549(5) Å). However, the $\text{Al}-\text{K}$ distance is considerably longer than the sum of the covalent radii of Al and K (3.24 Å)³⁴ suggesting an electrostatic interaction between the Al lone pair and cation, similar to that observed in boryl lithium **4**.²⁰ Attempts were made by the authors to synthesise a charge separated analogue by the addition of cryptand-222 or THF to **10**. However, both of these reactions led to immediate decomposition of the alumanyl anion.³² Compound **11** on the other

MINIREVIEW

hand is a *bone fide* charge-separated species, and only the second reported example of a 'naked' aluminyl anion. The K^+ cation in **11** is contained within two 12-crown-4 ligands and consequently has no close interactions with the aluminyl fragment (closest $Al \cdots K$ distance >6.6 Å).³³

2.2. Electronic Structure

For the purposes of like-for-like comparison, the electronic structures of all of the known aluminyl anions have been assessed by quantum chemical methods using the same level of theory as part of the present review (DFT-D3, PBE0, Def-TZVP – see SI for details). The calculations have been performed on the (full) monomeric anions, neglecting any stabilising/destabilising interactions with the counter-cations, and are summarised in Figure 5. All five anions possess relatively similar HOMO-LUMO energy gaps, varying by <0.3 eV (approx. 25 kJ mol⁻¹). More important in terms of reactivity is the energy separation between the occupied and vacant orbitals at the aluminium centre (i.e. the lone pair of electrons and the empty p_z -orbital). In all cases, the HOMO corresponds to a lone pair on aluminium. However, in every case bar the anionic component of **10**, the LUMO is a ligand-based orbital, with the empty Al p_z -orbital being higher in the orbital manifold: the LUMO+3 for the three diamido systems **6-8** and the LUMO+1 in **11** (see SI for more information). That said, the energy gap between the HOMO and the LUMO+n (LUMO+n corresponding to the empty Al centred p -orbital) is also similar in all cases, varying by a maximum of just over 50 kJ mol⁻¹ (from 3.42 eV, 330 kJ mol⁻¹ for **10** to 4.06 eV, 392 kJ mol⁻¹ for **7**).




Figure 5. Calculated HOMO, LUMO and LUMO+n energies of reported aluminyl anions at the DFT-D3, PBE0, Def-TZVP level of theory (LUMO+n indicates the orbital associated with the aluminium empty p -orbital). Orbital energies displayed in eV.

In general terms, many of the energetic trends shown in Figure 5 can be ascribed to structural factors familiar from carbene chemistry.³⁵ Thus, the lower lying nature of the HOMO

for diamido systems **6**, **7** and **8** compared to **10** and **11** (which feature alkyl substituents) can be ascribed to the more electronegative nature of the α -substituent. That the HOMO energies for **10** and **11** are so similar despite the narrower angle at aluminium and the presence of one σ electron-withdrawing nitrogen substituent in the latter is perhaps more surprising. The diamido aluminyl systems **7** and **8** are structurally very similar, with a slight difference being the wider N–Al–N angle in **8**, consistent with the larger heterocycle size ($108.84(9)^\circ$ *cf.* $103.89(8)^\circ$ for **7**). As such, the HOMO energy is marginally higher in the case of **8** (by 14 kJ mol⁻¹), consistent with higher p -character. Interestingly, however, the HOMO of **6**, which features both a much wider N–Al–N angle ($128.1(1)^\circ$) and potentially a weak additional donor/acceptor interaction from the xanthene oxygen is in a similar energy range.

In terms of the unoccupied orbitals, the differences between **10** and **11** also conform to established trends in carbene chemistry. Anion **10** is the only system where the Al p_z -orbital contributes to the LUMO; the analogous orbital in **11** is the LUMO+1. The difference in energy between these two orbitals is >30 kJ mol⁻¹, with that of **11** being the higher of the two, consistent with π -donation from the single amido substituent. For the diamido aluminyl systems **6-8**, the orbital characterized by significant Al p_z character lies higher in the orbital manifold (LUMO+3 in each case), with systems **7** and **8** being associated with almost identical energies. Superficially, the most surprising finding is the much lower lying nature of the corresponding orbital for xanthene-derived system **6**. On the face of it, this would imply relatively little elevation on the basis of Al–O σ^* character, and most significantly, much reduced N-to-Al π donation from the amido substituents (*cf.* **7** and **8**, see SI for molecular orbital images). The underlying origins for this are geometric in nature, and find precedent in the anti-Bredt carbene systems reported by Bertrand and co-workers.³⁶ In marked contrast to **7** and **8**, the amido substituents in **6** do not lie coplanar with the AlN_2 unit (Figure 6). The X-ray crystal structures of both dimeric $K_2[Al(NON)]_2$ and monomeric $[K(2.2.2\text{-crypt})][Al(NON)]$ reveal that the angles between the AlN_2 and NC_2 planes in these systems are typically of the order of $40\text{--}55^\circ$ (e.g. 53.7 and 50.9° for $[K(2.2.2\text{-crypt})][Al(NON)]$). Geometrically, this relates to the 'folded' non-planar geometry of the xanthene backbone required to accommodate the aluminium centre between two substituents in the 4- and 5-positions. As such, N-to-Al π -donation is severely restricted, as reflected in the significantly longer Al–N bonds measured for **6** compared to those in **7** and **8** (e.g. $2.022(1)/2.049(1)$ Å for $[K(2.2.2\text{-crypt})][Al(NON)]$, *cf.* $1.879(2)\text{--}1.896(2)$ Å for **7**).




Figure 6. Diagrammatic representation of the orbitals involved in the N-to-Al π -donation for anions **6** and **7**.

MINIREVIEW


In summary, dialkylaluminum compound **10**, not unexpectedly features the smallest energy gap between the HOMO and the Al p_z -orbital at (3.42 eV, 330 kJ mol⁻¹). The corresponding energy separation for alkyl amino aluminum **11** is higher (3.72 eV, 360 kJ mol⁻¹) due to elevation of the Al p_z -orbital, while those for the planar diamido systems **7** and **8** are higher still. Anion **7** features the largest energy separation (4.02 eV, 392 kJ mol⁻¹) with that of the related 7-membered system **8** being slightly smaller (3.90 eV, 376 kJ mol⁻¹) due to the effects of ring expansion on the HOMO. 'Base-stabilized' system **6** appears in reality to be destabilized by the constraints of the xanthene ligand backbone (with a corresponding energy separation of only 3.57 eV, 345 kJ mol⁻¹) due to geometrically restricted N-to-Al π donation. In terms of understanding reactivity patterns, it should be noted that only compounds **6** and **11** have so far been isolated as 'naked' aluminum anions. For ion pairs, interaction between the Al lone pair and the K⁺ cation would be expected to stabilise the HOMO and thereby widen the respective energy gaps.

2.3. Reactivity

2.3.1 Nucleophilic Substitution

One of the foremost points of novelty of aluminum anions is their ability to react as aluminium-centred nucleophiles. All of the reported aluminum anions have shown nucleophilic behaviour to some extent, with **6**, **8** and **10** showing evidence for nucleophilic substitution chemistry.^{27,30,32,37,38}


In our initial report we showed that **6** could undergo nucleophilic substitution reactions with carbon- and metal-centred electrophiles, as well as with a proton source.²⁷ The reaction of **6** with either MeI or MeOTf in toluene leads to an almost instantaneous reaction to give the Al–C bonded complex (NON)AlMe **12** with elimination of corresponding potassium salt (Scheme 1). Yamashita and co-workers have reported a similar reaction between **10** and MeOTf, yielding the aluminium methyl complex **13**.³² Both of these complexes were synthesised by nucleophilic attack on the electrophilic carbon centre of MeI or MeOTf by the aluminum. We also showed that a similar reaction can be performed using a bulky weak acid such as H₂(NON). This leads to formal nucleophilic attack on a proton (i.e. protonation) to give the aluminium hydride **14** (Scheme 1).



Scheme 1. Nucleophilic substitution reactions of **6** and **10** with carbon and hydrogen electrophiles.

Nucleophilic substitution reactions have also been successfully employed in the synthesis of Al–metal covalent bonds.^{27,30,37,38} This was first demonstrated in reactions between **6** and bulky Mg/Al iodide electrophiles,²⁷ which were

used to generate the corresponding Al–Al and Al–Mg bonded complexes **15** and **16** (Scheme 2). A small number of additional Al–metal bonded complexes have since been synthesised by this route, including those with Al–Ca (**18**)³⁰ and Al–Y (**19**)³⁸ Al–Au (**20** and **21**)³⁷ bonds (Scheme 2). It is noteworthy that all of these nucleophilic substitution reactions use electrophiles featuring weakly coordinating anions (i.e. OTf⁻, I⁻, BPh₄⁻), in order to promote salt elimination. Attempting these reactions with more coordinating anions (e.g. F⁻ or Cl⁻) tends to lead to the formation of 'ate' complexes (i.e. formal oxidative addition products) with the anion bound to the Al centre (see 2.3.2).



Scheme 2. Nucleophilic substitution reactions of aluminum anions to form the Al–metal bonded complexes **15**–**21**.

The Al–metal bonded heterobimetallic products **16**–**21** have also shown some interesting reactivity. These complexes can generally be divided into two groups depending on the polarisation of the Al–metal bond. If the aluminium centre is bound to a metal with an electronegativity lower than itself (i.e. <1.61 on the Pauling scale), then (superficially) the compound would be expected to be polarized in the sense Al(δ⁻)-M(δ⁺), with the reactivity profile of an aluminium-centred nucleophile (albeit moderated by the degree of polarity in the M–Al bond). At the other end of the spectrum, if aluminium is bound to a metal with an electronegativity greater than itself, then polarization would be expected in the sense Al(δ⁺)-M(δ⁻), consistent with a description as an aluminium electrophile bound to an electron-rich metal centre. Magnesium, calcium and yttrium all have electronegativities <1.61, and **16**–**19** might therefore be regarded as coordinated aluminum complexes.^{27,30,38} No reactivity of **16** or **19** has yet been reported, however unpublished work from our laboratory shows that **16** does in fact react as an aluminium centred nucleophile with various electrophiles including MeI. Reports on compounds **17** and **18** state that neither compound displays any observable reaction with benzene or toluene. **18** has been shown to react with both THF and cyclooctatetraene (COT);


MINIREVIEW

17 on the other hand shows no reactivity towards either of these substrates.³⁰

In contrast, gold has an electronegativity considerably greater than that of aluminium (2.54 on the Pauling scale). Accordingly, the Al–Au bonds in **20** and **21** show reversed polarization compared with the Al–metal bonds in **16–19**.³⁷ Reactions of **21** with the carbon centred electrophiles CO₂ and diisopropylcarbodiimide yield isolable Au–C bonded products. These reactions constitute the first examples of nucleophilic attack by a stable molecular gold complex.

2.3.2 Oxidative addition

As previously demonstrated for neutral Al(I) systems, the two-electron oxidation of Al(I) to Al(III) is typically a highly thermodynamically favourable process.¹¹ The situation appears to be little different for anionic alumanyl ions. As such, various oxidative addition reactions have been reported for alumanyl systems, in some cases involving the cleavage of very strong σ -bonds.^{27,31–33}



Scheme 3. Oxidative addition reactions of alumanyl anions **6**, **10** and **11**; tol = toluene.

In our initial work, we showed that **6** can oxidatively cleave the H–H bond of H₂ and a C–H bond in benzene under mild reaction conditions.²⁷ These reactions generate the formal oxidative addition products **22** and **23** (Scheme 3), demonstrating that the activation of strong σ -bonds by alumanyl anions is achievable. Furthermore, the reaction to yield **23** represented the first example of the oxidative addition of a benzene C–H bond at a single well-defined main-group metal centre. In more recent

work, we also showed that C–H activation of naphthalene was also possible.³¹

Since then, Yamashita and co-workers have shown that **10** can also C–H activate benzene at room temperature (RT) to give the formal oxidative addition product **24** (Scheme 3).³² The same group also reported reactions of **10** with C₆F₆ and benzyl chloride. The reaction with C₆F₆ yields a mixture of formal oxidative addition products **25a** and **25b**, showing mono- and di- activation of the C₆F₆, respectively. The ratio of these products was shown to vary depending on the molar ratio of C₆F₆ used. In contrast, the reaction of **10** with benzyl chloride did not give the expected C–Cl oxidative addition product, but a mixture of the corresponding benzylaluminate and chloroalumane. This result suggests that **10** is susceptible to a nucleophilic substitution at the chlorine atom that generates the chloroalumane and benzylpotassium.³²

Earlier this year, we showed that the activation of N–H bonds by an alumanyl anion was also possible, with the reaction between **6** and DippNH₂. This reaction gave the corresponding oxidative addition product **26** (Scheme 3).³⁹ Kinjo and co-workers followed this report by demonstrating that the N–H activation of ammonia and Si–H activation of phenylsilane are also possible with **11**.³³ These reactions led to the formal oxidative addition products **27** and **28** (Scheme 3). The group also showed reactivity with borane in the construction of an σ -aromatic AlB₂ ring and cleavage of a strained C–C bond in biphenylene to give the ring opened product **29**. The latter is the second example of C–C bond activation by a potassium alumanyl complex. The first reported C–C bond activation by an alumanyl anion was the remarkable ring opening of benzene reported by us in 2019.³¹ The ‘naked’ alumanyl **9** reacts cleanly with benzene at RT to give the 7-membered aluminium heterocycle **30** in high yields (Scheme 4). This reaction involves the oxidative cleavage of a C–C bond in benzene at a single aluminium centre. Remarkably, this reaction was also found to be reversible when heated. The ring opening of benzene to give **30** was calculated to be energetically favourable (–17 kJ mol^{–1}), even taking into account the loss of aromaticity in benzene.³¹ This has since been confirmed in a more comprehensive computational study by Fernández and co-workers reported earlier this year,⁴⁰ where the authors also investigated the effects incorporating electron withdrawing groups (EWGs) onto the ring and the C–C activation of the polycyclic aromatic systems naphthalene and anthracene. The study found that C–C activation of the polycyclic systems was unlikely, as the competing C–H activation pathway had a lower activation barrier (and more energetically favoured products). On the other hand, incorporating EWGs onto the benzene was found to favour the C–C activation both kinetically (by lowering the activation barrier) and thermodynamically. Unfortunately, this would be difficult to confirm experimentally as **9** would likely react with most of the common EWGs. Haloarenes, for example, are known to react with the related dimeric system **6** via formal C–I oxidative addition.



Scheme 4. Ring opening of benzene by the monomeric alumanyl anion **9**.


Compound **30** was found to react with 2 equivalents of the electrophile Me₂SnCl₂, yielding the *cis,cis,cis*-triene **31** and the

MINIREVIEW

corresponding dichloroaluminate.³¹ This reaction ultimately shows that alumanyl anions can be used to generate functionalized acyclic products from aromatic molecules.

2.3.3 Cycloaddition

Various cycloaddition reactions have been reported for alumanyl anions, including those with alkenes, alkynes and polyaromatics.^{29,31,41} The first of these to be reported was the reaction between **7** and cyclooctatriene (COT) by Coles and co-workers.²⁹ This reaction led to the two-electron reduction of the COT molecule to the essentially planar aromatic anion [COT]²⁻ dianion, which engages in η^2 -coordination to the Al centre in the product (**32**; Scheme 5). The addition of 18-crown-6 to this complex leads to its conversion to **33**, the product of a formal (1+4) cycloaddition addition product of COT to the aluminium centre (Scheme 5). Last year, we also reported that **6** can perform cycloaddition reactions, in this case with ethylene under slightly elevated pressure (2 bar).³¹ This gave the (1+2) cyclised product **34** (Scheme 5).



Scheme 5. Cycloaddition reactions between alumanyl anions **6**, **7** and **11** and various unsaturated substrates.


The most detailed study on cycloaddition reactions involving an alumanyl anion was reported earlier this year by Yamashita and co-workers.⁴¹ In this report, the authors showed that **11** can undergo various cycloaddition reactions with naphthalene **35**, anthracene **36**, diphenylacetylene **37** and 1,2-stilbene **38** (both *E* and *Z*). Compound **11** reacts with the polycyclic aromatic systems naphthalene and anthracene via a (1+4) cyclisation reaction at RT,

giving the aluma-norbornadiene derivatives **35** and **36**, respectively (Scheme 5). Interestingly **6** also undergoes a (1+4) cycloaddition reaction with naphthalene at RT, but at 80 °C yields C-H activation products,³¹ suggesting that the cycloaddition products of this type are kinetic products. Alumanyl complex **11** also undergoes (1+2) cycloaddition reactions with diphenylacetylene and (*E*)/(*Z*)-stilbenes, forming products **37** and **38** featuring AlC₂ 3-membered rings (Scheme 5). Interestingly, the reactions of **11** with both (*E*)- and (*Z*)-stilbenes gave exclusively the *trans*-cycloadduct **38**. DFT calculations revealed that the cycloaddition of **11** with (*Z*)-stilbene proceeds via a single transition state with a carbanion character, which results in the selectivity towards the *trans*-cycloadduct.⁴¹

2.3.4. Oxidation

Considering the highly reducing nature of the alumanyl anions, it comes as no surprise that these systems have been shown to react with a range of oxidising agents. However, by reversing the 'normal' reactivity of aluminium from electrophilic to nucleophilic, a range of previously inaccessible (in some cases highly reactive) Al(III) complexes can be accessed via this route.


In 2019, Coles and co-workers synthesised aluminium analogues of carbonyl and dioxirane moieties by oxidation with selenium.⁴² The 1:1 reaction between **7** and elemental selenium (using a catalytic amount of tri-*n*-butylphosphine) gave the aluminium selenide complex **39** in moderate yield (Scheme 6). The complex was found to be highly insoluble in common organic solvents due to an extended network of interactions between the potassium cation and the selenide and aryl groups in the solid state. This network could be broken with the addition of cryptand-222 to sequester the potassium cations to give a soluble monomeric form, **40** (Scheme 6). Complexes **39** and **40** represent the first examples of aluminium-selenium carbonyl analogues. DFT analysis performed on **40** found a Wiberg bond index of 1.39 for the Al–Se bond, implying significant multiple bond character, even though it is highly polarized.⁴² Addition of a further equivalent of selenium to **40** in THF yields complex **41**, which features an AlSe₂ three-membered ring (Scheme 6), which can be viewed as a heavy dioxirane analogue. The synthesis of **41** was also reported in a single step reaction by the addition of **7** to a suspension of selenium (2 equiv.) and cryptand-222 in THF. In a more recent account,⁴³ the group also reported the oxidation of **7** with sulphur, targeting the sulphur analogue of **39**. This reaction however did not give the targeted monosulfide but the 5-membered AlS₄ ring **42**.



Scheme 6. Oxidation of **7** with chalcogens.

MINIREVIEW


Later in 2019, two reports emerged almost simultaneously describing the reactivity of alumanyl anions with CO₂ and N₂O. These reports were by ourselves⁴⁴ and by Anker and Coles.⁴⁵ In our report,⁴⁴ we showed that **6** reacts rapidly with 1 atm. of CO₂ at RT, cleanly yielding the aluminium carbonate species **43** (Scheme 7). We also reported the analogous reaction between **6** and the phenyl isocyanate, which gave the isoelectronic aluminium carbamate complex **44**. Both reactions were shown to proceed via a two-step mechanism; first by initial reduction of the substrate (to give CO or PhNC) and a highly reactive aluminium oxide intermediate, which undergoes a (2+2) cycloaddition with a second equivalent of the substrate to yield the isolated products **43** and **44**. Similar reactions have been reported with other low oxidation state main group systems.⁴⁶ In an attempt to isolate the oxide intermediate, the oxidation of **6** with N₂O was investigated. However, the reaction between **6** and N₂O (1 atm.) in toluene did not give the targeted oxide species, but the rather unexpected *cis*-hyponitrite complex **45**. As with the syntheses of **43** and **44**, the reaction seems to proceed via a highly reactive oxide intermediate, which remarkably undergoes a (2+3) cycloaddition with a molecule of N₂O. This reaction demonstrates the first example of a *cis*-hyponitrite ligand being directly synthesised from N₂O. The reaction between **6** and N₂O in toluene was monitored by low temp. ¹H NMR, but even at -80 °C, no signs of an intermediate were observed. However, it was found that the intermediate could be trapped by the coordination of THF to the Al centre, by performing the reaction in THF at low temperature. The reaction between **6** and N₂O (1 atm.) <0 °C cleanly gives the THF-coordinated monoaluminoxane **46** in high yield (Scheme 7).⁴⁴



Scheme 7. Oxidation of **6** with CO₂, PhNCO and N₂O and subsequent reactivity of monoaluminoxane **46**.

Complex **46** was found to be stable in the solid state at RT but not in solution, decomposing over the course of several hours. **46** was found to react with CO₂, PhNCO and N₂O at RT to give the previously isolated products **43-45**, showing that the (more reactive) base-free monoaluminoxane is a viable intermediate in these reactions. Remarkably, we also found that due to the extremely polar Al-O bond, **46** could heterolytically cleave H₂ at RT to give the aluminium hydride/hydroxide species **47** (Scheme 7).

Anker and Coles reported a similar study,⁴⁵ showing that alumanyl **7** also reacts with 1 atm. of CO₂ to give the aluminium carbonate complex **48** (Scheme 8). However, in their case, they found that the monoaluminoxane intermediate **49** could be isolated without the coordination of a Lewis-base, directly from the reaction of **7** and N₂O (1 atm.). In an analogous fashion to **46**, the monoaluminoxane **49** was found to react with CO₂ and N₂O (in the presence of cryptand-222) in cycloaddition reactions to give **42** and the *cis*-hyponitrite complex **50**. Addition of cryptand-222 to **49** in the absence of CO₂ or N₂O, leads to intramolecular C-H activation of the ligand across the Al-O bond; C-H activation has not yet been reported for **46**.




Scheme 8. Oxidation of **7** with CO₂ and N₂O and subsequent reactivity of monoaluminoxane **49**.

In a more recent account,⁴³ the Coles group showed that an aluminium trithiocarbonate complex **51** (Scheme 8) could also be synthesised, by the addition of a hexane solution of **7** to a solution of CS₂ (2 equiv.) at RT. However, in this case the expected monosulfide intermediate could not be isolated (see Scheme 6). Interestingly though, the group found that performing the same reaction under slightly different conditions (4 equiv. 195 K), a different product could be isolated. This was the ethenetetrathiolate complex **52** (Scheme 8), which is synthesised by a C-C bond forming reaction between two CS₂ units. Remarkably, the synthesis of the lighter analogue, the ethenetetraolate complex **53** was also reported, by the reaction between CO and the monoaluminoxane **49**. This complex represents the first report of an ethenetetraolate ligand.

MINIREVIEW


The oxidation of **6** and **7** have also been investigated with regards to organic azides.^{39,47} Two recent reports have found that under certain conditions, these reactions can give isolable aluminium imide complexes with highly polarised (and highly reactive) Al–N bonds. Earlier this year, our group reported reactions of **6** with various azides using a variety of reaction conditions (Scheme 9).³⁹ The RT reaction between **6** and 2 equiv. of trimethylsilyl azide gave the aluminium tetrazine complex **54**, which is essentially isostructural to the *cis*-hyponitrite complex **45**.⁴⁴ It can be proposed that the reaction to give **54** proceeds via a similar mechanism to that to give **45**; i.e. firstly by reduction of one molecule of azide to give a highly reactive aluminium imide intermediate, which undergoes a (2+3) cycloaddition reaction with a further equivalent of the azide to give **54**. In an attempt to trap this azide intermediate, the 1:1 reaction between **6** and trimethylsilylazide was attempted in toluene at $-80\text{ }^{\circ}\text{C}$. This did not give the targeted imide but instead the C–H activated toluene product **55** (Scheme 9), demonstrating the highly basic nature of the aluminium imide intermediate. In an attempt to isolate a stable aluminium imide, the bulkier azide DippN₃ was sought. In a 1:1 reaction between **6** and DippN₃, the targeted aluminium imide **56** was isolated in high yields (Scheme 9). The compound was found to feature an extremely polar Al–N bond (Mulliken charges +0.67 Al and -0.16 N) with a Wiberg Bond Index (WBI) of 0.705.³⁹



Scheme 9. Oxidation of **6** with organic azides and the activation of H_2 and CO by the aluminium imide **56**.

The Al imide **56** was found to be extremely reactive with regards to small molecule activation, showing reactivity towards H_2 and CO under mild reaction conditions.³⁹ The reaction between **56** and H_2 led to heterolytic cleavage of the H_2 molecule to give **57** (Scheme 9), whilst the reaction with CO saw the C–C coupling of two CO molecules to give an unprecedented Al bound $[(\text{DippN})\text{CCO}_2]^{2-}$ fragment **58**.

In a report by Anker and Coles, a similar aluminium imide **59** was reported by the reaction between **7** and MesN_3 (Scheme 10).⁴⁷ This complex was found to react with CO_2 (1 atm.) at RT in a (2+2) cycloaddition reaction to give the carbamate complex **60**.



Scheme 10. Oxidation of **7** with MesN_3 to give the aluminium imide **59** and subsequent reactivity with CO_2 .

3. Summary and Outlook

Although the field of aluminyl chemistry is still in its infancy, the scope of transformations reported by these novel anions is both fascinating and diverse. In this minireview, we have covered the synthesis, structures and reactivity for the currently known series of aluminyl anions. Reactions including nucleophilic substitution, oxidative addition, cycloaddition and oxidations have already been reported, with highlights including the facile C–H and C–F activation, small molecule activation and the ring opening of aromatics. DFT calculations performed on the series of known anions revealed some interesting trends with regards to their electronic structures. Firstly, anion **6** has the lowest energy LUMO+n (empty Al *p*-orbital) of the series in addition to one of the lowest HOMO-LUMO gaps. **6** is also one of only two anions (the second being **11**) that has been isolated free from its cation, which would otherwise stabilise the HOMO. These findings are consistent with its reactivity, as the monomeric anion **6** appears to be the most reactive of the current series (the only reported anion to ring open benzene for example). The diamido anions **7** and **8** have similar electronic structures, with **8** having a slightly smaller HOMO-LUMO gap due to a more destabilised HOMO. The alkyl systems **10** and **11** feature the high energy HOMOs as expected, due to the more electropositive substituents bound to the Al centre. **10** was found to have the smallest HOMO-LUMO gap of the series, but the charge-separated anion does not appear to be stable.

We believe the chemistry highlighted in this report only scratches the surface for what is to come from this novel class of nucleophiles. The coordination chemistry of the anions is only just beginning and small molecule activation including N_2 and CO is also yet to be reported. The use of aluminyl anions in catalysis is also an area that we are currently investigating. The field is rapidly expanding and definitely ‘one to watch’ over the next few years.

Acknowledgements

JH would like to thank the Australian Research Council and the Australian Government for funding aspects of this work (DE190100524). PV thanks the Academy of Finland (Grant No. 314794) and Prof. Heikki M. Tuononen for continued support and computational resources. JMG and SA would also like to thank the Leverhulme Trust (F/08699/E) for funding aspects of this work.


Keywords: aluminyl • aluminium • nucleophile • organometallic • main group

MINIREVIEW

- [1] W. M. Haynes (Ed.) in *CRC Handbook of Chemistry and Physics*, 97th Edition, CRC Press, **2016**, pp. 14-17.
- [2] A. Bagshaw (Ed.) in *Bauxite to Alumina the Bayer Process: An Introductory Text*, Nedlands, **2017**.
- [3] S. Aldridge, A. J. Downs (Eds.) in *The Group 13 Metals Aluminium, Gallium, Indium and Thallium: Chemical Patterns and Peculiarities*, Wiley, Chichester, **2011**.
- [4] a) S. Saito in *Main Group Metals in Organic Synthesis*, (Eds.: H. Yamamoto, K. Oshima) Wiley-VCH, **2004**, pp. 189-306; b) P. A. Chaloner in *The Metal-Carbon Bond* (Ed.: F. R. Hartley), John Wiley & Sons Ltd., **1987**, pp. 411-472; c) T. Ooi, K. Maruoka in *Lewis Acids in Organic Synthesis*, (Ed.: H. Yamamoto) Wiley-VCH, **2000**, pp. 191-281.
- [5] a) M. S. Hill in *Low Oxidation State Chemistry. Encyclopedia of Inorganic and Bioinorganic Chemistry*, **2015**, 1-18; b) P. Bag, C. Weetman, S. Inoue, *Angew. Chem. Int. Ed.* **2018**, *57*, 14394-14413; c) A. Schnepf, H. Schnöckel, *Angew. Chem. Int. Ed.* **2002**, *41*, 3532-3552 and references therein.
- [6] a) W. Uhl, *Angew. Chem. Int. Ed.* **1993**, *32*, 1386-1397; (b) C. Dohmeier, D. Loos, H. Schnöckel, *Angew. Chem. Int. Ed.* **1996**, *35*, 129-149.
- [7] a) W. Klemm, E. Voss, K. Z. Geiersberger, *Z. Anorg. Allg. Chem.* **1948**, *256*, 15-24; b) P. J. Durrant, B. Durrant, *Introduction to Advanced Inorganic Chemistry*: Prentice Hall Press, London, **1972**; c) M. Mocker, C. Robl, H. Schnöckel, *Angew. Chem. Int. Ed.* **1994**, *33*, 1754-1755; d) A. Ecker, H. Schnöckel, *Z. Anorg. Allg. Chem.* **1996**, *622*, 149-152; e) H. J. Himmel, J. Bahlo, M. Haussmann, F. Kurth, G. Stösser, H. Schnöckel, *Inorg. Chem.* **2002**, *41*, 4952-4960; f) M. Tacke, H. Schnöckel, *Inorg. Chem.* **1989**, *28*, 2895-2896.
- [8] C. Dohmeier, C. Robl, M. Tacke, H. Schnöckel, *Angew. Chem. Int. Ed.* **1991**, *30*, 564-565.
- [9] S. Schulz, H. W. Roesky, H. J. Koch, G. M. Sheldrick, D. Stalke, A. Kuhn, *Angew. Chem. Int. Ed.* **1993**, *32*, 1729-1731.
- [10] C. Cui, H. W. Roesky, H.-G. Schmidt, M. Noltemeyer, H. Hao, F. Cimpoesu, *Angew. Chem. Int. Ed.* **2000**, *39*, 4274-4276.
- [11] Y. Lui, J. Li, X. Ma, Z. Yang, H. W. Roesky, *Coord. Chem. Rev.* **2018**, *374*, 387-415.
- [12] a) C. Bakewell, M. Garçon, R. Y. Kong, L. O'Hare, A. J. P. White, M. R. Crimmin, *Inorg. Chem.* **2020**, *59*, 4608-4616; b) A. Dmitrienko, J. F. Britten, D. Spasyuk, G. I. Nikonov, *Chem. Eur. J.* **2020**, *26*, 206-211; c) C. Bakewell, A. J. P. White, M. R. Crimmin, *Chem. Sci.* **2019**, *10*, 2452-2458; d) L. L. Liu, J. Zhou, L. L. Cao, D. W. Stephan, *J. Am. Chem. Soc.* **2019**, *141*, 16971-16982; e) S. Sinhababu, M. M. Siddiqui, S. K. Sarkar, A. Münch, R. Herbst-Irmer, A. George, P. Parameswaran, D. Stalke, H. W. Roesky, *Chem. Eur. J.* **2019**, *25*, 11422-11426; f) A. Paparo, C. D. Smith, C. Jones, *Angew. Chem. Int. Ed.* **2019**, *58*, 11459-11463; g) A. Koner, B. M. Gabidullin, Z. Kelemen, L. Nyulászi, G. I. Nikonov, R. Streubel, *Dalton Trans.* **2019**, *48*, 8248-8253; h) C. Bakewell, B. J. Ward, A. J. P. White, M. R. Crimmin, *Chem. Sci.* **2018**, *9*, 2348-2356; i) S. Brand, H. Elsen, J. Langer, W. A. Donaubaue, F. Hampel, S. Harder, *Angew. Chem. Int. Ed.* **2018**, *57*, 14169-14173; j) R. Y. Kong, M. R. Crimmin, *J. Am. Chem. Soc.* **2018**, *140*, 13614-13617; k) S. Sinhababu, S. Kundu, A. N. Paesch, R. Herbst-Irmer, D. Stalke, H. W. Roesky, *Eur. J. Inorg. Chem.* **2018**, 2237-2240; l) C. Bakewell, A. J. P. White, M. R. Crimmin, *Angew. Chem. Int. Ed.* **2018**, *57*, 6638-6642; m) L. Tuscher, C. Helling, C. Wölper, W. Frank, A. S. Nizovtsev, S. Schulz, *Chem. Eur. J.* **2018**, *24*, 3241-3250; n) T. N. Hopper, M. Garçon, A. J. P. White, M. R. Crimmin, *Chem. Sci.* **2018**, *9*, 5435-5440.
- [13] P. Bag, A. Porzelt, P. J. Altmann, S. Inoue, *J. Am. Chem. Soc.* **2017**, *139*, 14384-14387.
- [14] S. K. Møllerup, Y. Cui, F. Fantuzzi, P. Schmid, J. T. Goettel, G. Bélanger-Chabot, M. Arrowsmith, I. Krummenacher, Q. Ye, V. Engel, B. Engels, H. Braunschweig, *J. Am. Chem. Soc.* **2019**, *141*, 16954-16960.
- [15] A. Hofmann, C. Prankevicus, T. Tröster, H. Braunschweig, *Angew. Chem. Int. Ed.* **2019**, *58*, 3625-3629.
- [16] We use the term 'aluminyl' to refer to systems containing the formally anionic AlX_2^- fragment, based on well established literature precedent for the related derivatives of other group 13 elements: i.e. boryl, gallyl and indyl for the boron, gallium and indium analogues. Although the terms boranyl, aluminyl, gallanyl and indanyl are recommended by the IUPAC red book (R. M. Hartshorn, A. T. Hutton in *Nomenclature of Inorganic Chemistry, IUPAC Recommendations 2005, IUPAC Red Book*, (Eds.: N. G. Connelly, T. Damhus), RSC Publishing, **2005**) for the parent BH_2^- , AlH_2^- , GaH_2^- and InH_2^- systems, this convention has not been followed in the naming of previous boron, gallium or indium systems of the types BX_2^- , GaX_2^- and InX_2^- .
- [17] E. S. Schmidt, A. Jockisch, H. Schmidbauer, *J. Am. Chem. Soc.*, **1999**, *121*, 9758-9759.
- [18] a) R. J. Baker, R. D. Farley, C. Jones, M. Kloth, D. M. Murphy, *J. Chem. Soc., Dalton Trans.*, **2002**, 3844-3850; b) I. L. Fedushkin, A. N. Lukoyanov, G. K. Fukin, S. Y. Ketkov, M. Hummert, H. Schumann, *Chem. Eur. J.* **2008**, *14*, 8465-8468.
- [19] a) R. J. Baker, C. Jones, *Coord. Chem. Rev.* **2005**, *249*, 1857-1869; b) M. Assay, C. Jones, M. Driess, *Chem. Rev.* **2011**, *111*, 354-396.
- [20] Y. Segawa, M. Yamashita, K. Nozaki, *Science* **2006**, *314*, 113-115.
- [21] Y. Segawa, Y. Suzuki, M. Yamashita, K. Nozaki, *J. Am. Chem. Soc.* **2008**, *130*, 16069-16079.
- [22] L. Weber, *Eur. J. Inorg. Chem.* **2017**, 3461-3488.
- [23] a) W. Lu, H. Hu, Y. Li, R. Ganguly, R. Kinjo, *J. Am. Chem. Soc.* **2016**, *138*, 6650-6661; b) A.-F. Pécharman, A. L. Colebatch, M. S. Hill, C. L. McMullin, M. F. Mahon, C. Weetman, *Nat. Commun.* **2017**, *8*, 15022.
- [24] a) R. J. Schwamm, M. D. Anker, M. Lein, M. P. Coles, C. M. Fitchett, *Angew. Chem. Int. Ed.* **2018**, *57*, 5885-5887; b) M. D. Anker, Y. Altaf, M. Lein, M. P. Coles, *Dalton Trans.* **2019**, *48*, 16588-16594.
- [25] M. D. Anker, M. Lein, M. P. Coles, *Chem. Sci.*, **2019**, *10*, 1212-1218.
- [26] a) B. E. Cole, J. P. Wolbach, W. G. Dougherty, Jr., N. A. Piro, W. S. Kassel, C. R. Graves, *Inorg. Chem.* **2014**, *53*, 3899-3906; b) A. Hinchliffe, F. S. Mair, E. J. L. McInnes, R. G. Pritchard, J. E. Warren, *Dalton Trans.* **2008**, 222-233; c) Y. Zhao, Y. Liu, B. Wu, X.-J. Yang, *Dalton Trans.* **2015**, *44*, 13671-13680.
- [27] J. Hicks, P. Vasko, J. M. Goicoechea, S. Aldridge, *Nature* **2018**, *557*, 92-95.
- [28] A third example of a diamido aluminyl anion was reported by Harder and co-workers whilst this manuscript was under review - see S. Harder, S. Grams, J. Eyslein, J. Langer, C. Färber, *Angew. Chem. Int. Ed.* **2020** Accepted Articles (DOI: 10.1002/anie.202006693).
- [29] R. J. Schwamm, M. D. Anker, M. Lein, M. P. Coles, *Angew. Chem. Int. Ed.* **2019**, *58*, 1489-1493.
- [30] R. J. Schwamm, M. P. Coles, M. S. Hill, M. F. Mahon, C. L. McMullin, N. A. Rajabi, A. S. S. Wilson, *Angew. Chem. Int. Ed.* **2020**, *59*, 3928-3932.
- [31] J. Hicks, P. Vasko, J. M. Goicoechea, S. Aldridge, *J. Am. Chem. Soc.* **2019**, *141*, 11000-11003.
- [32] S. Kurumada, S. Takamori, M. Yamashita, *Nat. Chem.* **2020**, *12*, 36-39.
- [33] K. Koshino, R. Kinjo, *J. Am. Chem. Soc.* **2020**, *142*, 9057-9062.
- [34] B. Cordero, V. Gómez, A. E. Platero-Prats, M. Revés, J. Echeverría, E. Cremades, F. Barragán S. Alvarez, *Dalton Trans.* **2008**, 2832-2838.
- [35] D. Bourissou, O. Guerret, F. P. Gabbaï, G. Bertrand, *Chem. Rev.* **2000**, *100*, 39-92.
- [36] D. Martin, N. Lassauque, B. Donnadiou, G. Bertrand, *Angew. Chem. Int. Ed.* **2012**, *51*, 6172-6175.
- [37] J. Hicks, A. Mansikkamäki, P. Vasko, J. M. Goicoechea, S. Aldridge, *Nat. Chem.* **2019**, *11*, 237-241.
- [38] K. Sugita, M. Yamashita, *Chem. Eur. J.* **2020**, *26*, 4520-4523.
- [39] A. Heilmann, J. Hicks, P. Vasko, J. M. Goicoechea, S. Aldridge, *Angew. Chem. Int. Ed.* **2020**, *59*, 4897-4901.
- [40] J. J. Cabrera-Trujillo, I. Fernandez, *Chem. Eur. J.* **2020**, Accepted Articles (DOI: 10.1002/chem.202000921).
- [41] K. Sugita, R. Nakano, M. Yamashita, *Chem. Eur. J.* **2020**, *26*, 2174-2177.
- [42] M. Anker, M. P. Coles, *Angew. Chem. Int. Ed.* **2019**, *58*, 13452-13455.
- [43] M. Anker, C. L. McMullin, N. A. Rajabi, M. P. Coles, *Angew. Chem. Int. Ed.* **2020**, Accepted Articles (DOI: 10.1002/ange.202005301).
- [44] J. Hicks, A. Heilmann, P. Vasko, J. M. Goicoechea, S. Aldridge, *Angew. Chem. Int. Ed.* **2019**, *58*, 17265-17268.
- [45] M. Anker, M. P. Coles, *Angew. Chem. Int. Ed.* **2019**, *58*, 18261-18265.
- [46] a) D. Wendel, A. Porzelt, F. A. D. Herz, D. Sarkar, C. Jandl, S. Inoue, B. Rieger, *J. Am. Chem. Soc.* **2017**, *139*, 8134-8137; b) R. Lalrempuia, A. Stasch, C. Jones, *Chem. Sci.* **2013**, *4*, 4383-4388.
- [47] M. D. Anker, R. J. Schwamm, M. P. Coles, *Chem. Commun.* **2020**, *56*, 2288-2291.

MINIREVIEW

Entry for the Table of Contents



Nucleophilic Aluminium! A series of aluminium(I) anions (aka aluminyl anions) have recently been reported that remarkably show nucleophilic behaviour at aluminium. This minireview compares and contrasts the synthesis, structure, reactivity and electronic properties of these aluminyl anions whilst proposing future directions for the field.

Institute and/or researcher Twitter usernames: @HicksChem @ANUChemistry @GroupAldridge @OxfordChemistry