

This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Koivunen, Kaisa; Sillanpää, Elina; Munukka, Matti; Portegijs, Erja; Rantanen, Taina

Title: Cohort differences in maximal physical performance : a comparison of 75- and 80-year-old men and women born 28 years apart

Year: 2021

Version: Accepted version (Final draft)

Copyright: © 2020 the Authors

Rights: In Copyright

Rights url: <http://rightsstatements.org/page/InC/1.0/?language=en>

Please cite the original version:

Koivunen, K., Sillanpää, E., Munukka, M., Portegijs, E., & Rantanen, T. (2021). Cohort differences in maximal physical performance : a comparison of 75- and 80-year-old men and women born 28 years apart. *Journals of Gerontology Series A : Biological Sciences and Medical Sciences*, 76(7), 1251-1259. <https://doi.org/10.1093/gerona/glaa224>

Cohort differences in maximal physical performance: a comparison of 75- and 80-year-old men and women born 28 years apart

Kaisa Koivunen, MSc¹, Elina Sillanpää, PhD¹, Matti Munukka, PhD¹, Erja Portegijs PhD¹, Taina Rantanen, PhD¹

1. Faculty of Sport and Health Sciences and Gerontology Research Center, University of Jyväskylä, Finland

Address correspondence to:

Kaisa Koivunen, MSc,

Faculty of Sport and Health Sciences and Gerontology Research Center,

University of Jyväskylä, Finland, PO Box 35 (Viveca),

University of Jyväskylä, Jyväskylä FI-40014, Finland.

E-mail: kaisa.m.koivunen@jyu.fi

p. +358 503608818

Abstract

Background: Whether increased life expectancy is accompanied by increased functional capacity in older people at specific ages is unclear. We compared similar validated measures of maximal physical performance in two population-based older cohorts born and assessed 28 years apart.

Methods: Participants in the first cohort were born in 1910 and 1914 and were assessed at age 75 and 80 years, respectively (N=500, participation rate 77%). Participants in the second cohort were born in 1938 or 1939 and 1942 or 1943 and were assessed at age 75 and 80 years, respectively (N=726, participation rate 40%). Participants were recruited using a population register and all community-dwelling persons in the target area were eligible. Both cohorts were interviewed at home and examined at the research center with identical protocols. Maximal walking speed, maximal isometric grip and knee extension strength, lung function measurements; forced vital capacity (FVC) and forced expiratory volume in 1 second (FEV1) were assessed. Data on non-participation were systematically collected.

Results: Walking speed was on average 0.2-0.4 m/s faster in the later than earlier cohort. In grip strength, the improvements were 5-25%, and in knee extension strength 20-47%. In FVC, the improvements were 14-21% and in FEV1 0-14%.

Conclusions: The later cohort showed markedly and meaningfully higher results in the maximal functional capacity tests, suggesting that currently 75- and 80-year old people in Finland are living to older ages nowadays with better physical functioning.

Key words: Secular trends, Birth cohorts, Functional capacity

Introduction

The life expectancy of older people is increasing. In Finland, for example, a person aged 75 in 1989 could expect to live a further 9.7 years whereas a same-age individual in 2017 could expect a further 12.6 years¹. However, it is not clear whether a longer life accompanies improvements in functioning at specific ages. If functioning at specific ages is better than in the past, this could lead to a more positive outlook towards aging and contribute to projections on the needs of the work force and health and social care.

The earlier studies assessing cohort differences in the health and functioning of older people give an inconsistent picture. The prevalence of chronic conditions have been found to be stable or to increase among more recent than earlier cohorts of older people²⁻⁴. The results obtained from self-rated health and disability show improved^{2,5}, worsened⁶ and stable^{7,8} trends, depending on the study. These differences may stem from differences in the age groups studied, intervals between cohorts, indicators of disability and functioning, different trends between countries, and possibly from problems of comparability between recent and earlier cohorts. Moreover, the earlier studies were often based on self-report data. Apart from an individual's intrinsic capacity, self-assessments may be influenced by environmental circumstances, which may underlie the results⁵.

Compared to self-assessments, performance-based measures requiring maximum effort provide more explicit and standardized information on cohort differences in physical functioning. Muscle strength, walking speed and respiratory function tests are informative and widely used performance-based tests of functional capacity, that capture current and preceding lifetime influences on functioning and predict disability and mortality risk^{9,10}. Only a few studies have assessed cohort effects in performance-based maximum measures of physical performance, and the results have been mixed. For example, two studies reported an improved trend in hand grip strength^{11,12} whereas other studies noted no improvement^{5,13} or decline¹⁴ among the more

recent cohorts. In addition, a large population-based study assessing older adults from Germany, Sweden and Spain found contrasting trends in grip strength for different age groups. The results showed strong improvement for older adults aged 80 years and older, while younger older adults stagnated or even decreased¹⁵. In other studies, the later-born cohorts performed better in chair stand, walking speed and peak expiratory flow (PEF) tests^{13,16,17}. When interpreting these results, it is important to bear in mind that timing and the rate of age-related changes between organs or biological systems within and between individuals differ^{18,19}. For example, muscle strength declines earlier than walking speed and possibly precedes the decline in lung function^{20,21}. Incorporating multiple measures will build a more comprehensive picture of the changes accompanying aging.

The challenges in assessing cohort effects include ensuring the comparability of the assessment methods and populations studied. Researchers in Finland are in an exceptionally good position to meet these challenges. First, a population-based study conducted in our center 28 years ago with standardized maximum performance-based assessment methods provides us with a valid point of reference on the functioning of people born approximately one generation ago^{22,23}. Second, we can base recruitment on population registers, which reduces selection bias¹¹.

This study examined whether older adults born 28 years later have better physical performance compared to an earlier cohort measured at the same age. The factors underlying potential cohort differences are also investigated.

Methods

Study population

This study forms part of two projects conducted at the University of Jyväskylä, Finland. The dataset comprises the Evergreen cohort data collected in 1989-1990²⁴ and the Evergreen II cohort data collected in 2017-2018 as part of the *Active Ageing – Resilience and external support as modifiers of disablement outcome* (AGNES) project²⁵. For both projects, samples were drawn from the Finnish Population Register based on birth year and place of residence. All community-living 75- and 80-year-old residents of the city of Jyväskylä formed the target group. Members of the earlier cohort examined in 1989-1990 were born in 1910 and 1914 and members of the later cohort examined in 2017-2018 were born in 1938-1939 and 1942-1943.

Recruitment, participation and non-participation

The Evergreen and Evergreen II recruitment procedures are comparable. Recruitment was as inclusive as possible. All persons in the targeted age groups, who were living in the community in a non-institutional setting in the recruitment area and able to respond, and who consented to take part, were included.

The recruitment area, the City of Jyväskylä, had expanded since the first Evergreen project due to mergers with neighboring municipalities. However, we targeted people, whose addresses were within the previous city area or in similar adjacent areas, including urban areas and suburbs with apartment buildings and detached houses.

During the Evergreen study, in 1989-1990, participants were sent a letter informing them about the study and suggesting a time for a home interview. Those who declined were asked to give their reasons for non-attendance and the reasons were documented. Evergreen II participants in 2017-2018 were sent a letter informing them about the study after which we enquired about their willingness to take part by phone. For those willing to take part, the home interview was scheduled. During the phone call, those declining to take part were questioned on their reasons for non-participation. The study flow charts are shown in Figure 1. In Evergreen, 500 (77%) and in Evergreen II, 726 (40%) of those eligible participated in the home interviews and research center assessments. In both studies, self-rated health was examined with the question: “How would you yourself describe your health during the last year?” using a five-option response scale ranging from very poor to very good. For statistical analysis, we recoded the responses as good, moderate and poor. In the Evergreen study, the question was asked during the home interview while in the Evergreen II it was posed during the initial phone call.

Figure 1

Assessment procedure

The implementation and assessment methods in both projects were identical for all practical purposes. The interviews were conducted in the participants' homes and the physical tests in the Sport and Health Laboratory of the University of Jyväskylä. The measurement equipment and laboratory environment were similar for both cohorts.

Physical performance measurements

Ten-meter maximal walking speed was assessed in the laboratory corridor using a hand-held stopwatch. Five meters was allowed for acceleration and the participant was encouraged to continue a few meters past the finish line. Participants wore walking shoes or sneakers²⁵.

Maximal isometric hand grip strength and maximal isometric knee extension strength were measured in the Evergreen II cohort using an adjustable dynamometer chair (Good Strength; Metitur Oy, Palokka, Finland) and the result expressed in Newtons (N)²⁶. For the Evergreen cohort, we used the prototype of the Good Strength device, including similar strain gauge technology. In both cohorts, the measurements were performed identically with similar joint angles and instructions to the participant. The measurements were done on the side of the dominant hand in a sitting position with the lower back supported. Hand grip strength was measured using a dynamometer fixed to the arm of the chair. Knee extension strength was measured at an angle of 60 degrees from the fully extended leg towards flexion. After a practice trial, the test was performed at least three times with a one-minute inter-trial rest period until no further improvement occurred, and the highest value was recorded²⁵. The test-retest reliability of both tests is excellent. In the 80-year-olds, the Pearson correlation coefficients between measurements conducted one to two weeks apart were $r= 0.967$ for hand grip strength and $r= 0.965$ for knee extension strength²³.

Respiratory function in the Evergreen II cohort was assessed using spirometry (Medikro Pro spirometer, Medikro Oy, Kuopio, Finland) in a standing position with a nose clip. The forced vital capacity (FVC) maneuver was performed at least two times. Participants were instructed to inhale maximally and exhale as hard and as fast as possible and continue until there was no air left. The manoeuver was continued until the criterion of the ATS/ERS Taskforce²⁷ was met or a maximum of eight exhalations was performed. With the Evergreen cohort, respiratory function was assessed using comparable electronic spirometry (Medikro 202;

Medikro, Kuopio, Finland) in a standing position and three trials were allowed. In both studies, the highest volume of FVC and the forced expiratory volume in one second (FEV1) were recorded in liters. PEF was recorded in liters/second.

Covariates

Our analyses were not adjusted for any confounders. The age and gender groups were similar, and we concluded that differences in covariates between the cohorts were more likely to be factors underlying the cohort differences than confounders. To study these factors, we chose correlates of physical performance that differed between the cohorts, and that theoretically can be part of the mechanism leading to secular change. Years of full-time education is known to be associated with health and functional status²⁸, and was used to describe socio-economic position. Body size, especially height, affects muscle strength, walking speed and respiratory function^{29,30}. We measured height with a stadiometer in cm and body mass with a beam scale in kg. Health behavior was described with physical activity and smoking. Physical activity was assessed with a single validated self-report question with six response options ranging from mostly sitting and resting to regular strenuous exercise²⁶. For the statistical analysis, the responses were recoded as low, moderate and high. Smoking was classified as never vs. current/ former smoker.

Statistical analysis

To compare the current and earlier same-age cohorts, we used t-tests for continuous and chi-square test for categorical variables. We tested whether the cohort differences varied according to age and sex by adding birth cohort-by-sex and birth cohort-by-age interaction terms into the linear regression analyses comprising all the participants. Factors underlying the potential cohort differences were studied in a set of linear regression models. First, the models were fitted with each physical test as a dependent variable and birth

cohort as an independent variable. Subsequently, we run several models adding covariates one at a time to analyze which of them attenuates the cohort differences in physical performance. Self-rated health was not included in the models explaining cohort differences, as we believe that improved self-rated health is likely to be a result of better physical functioning and not an explanatory factor. To clarify the potential clinical significance of the cohort differences, hand grip strength cut points for increased risk for mobility limitation were determined separately for men and women based on the Finnish reference data (37kg for men and 21kg for women)³¹. According to our knowledge, cut-points for knee extension strength and gait speed that predict mobility limitations have not been analyzed based on nationally representative samples in Finland, and therefore are not available for our use. Finally, we did sensitivity analyses to assess the comparability of our cohorts based on the data available for non-participants. If the non-participants were comparable, this would suggest that the cohorts were also comparable and that the differences observed between the cohorts were less likely attributable to selection bias.

Results

In men and women and in both age groups, the number of years of education had doubled in the later- compared to earlier-born cohort (Table 1). In addition, the later-born cohort reported higher daily physical activity and better self-rated health compared to the earlier-born cohort. Among men, the proportion of ever smokers was lower in the later than earlier cohort. Among the 75-year-old women, the proportion of ever smokers was higher in the later than earlier cohort.

Mean grip strength, knee extension strength and walking speed were higher in the later- than earlier-born cohort (Table 2). In the respiratory function measures, the later-born cohort performed better in FVC and, among the 80-old men and women, in the FEV1 measures. The cohorts did not differ in PEF.

Grip strength below the validated cut-point for increased risk for mobility limitation (37 kg for men and 21 kg for women,³¹) was more evident in the earlier-born cohort. Among 75-year-old men, percentage of participants below the cut-point was 48% and 27% in the earlier and later-born cohort, respectively (between cohorts $p < 0.001$). Among 75-year-old women the proportions were 35% vs. 28% ($p = 0.052$). In 80-year-olds, 71% vs. 52% in men ($p = 0.012$) and 75% vs. 44% in women ($p < 0.001$), respectively, had values below the cut-points for increased risk for mobility limitation.

Table 2 shows the relative differences between the cohorts. The regression analyses comprising all participants showed significant cohort-by-sex interactions for grip strength ($p = 0.041$), FVC ($p = 0.015$) and FEV1 ($p = 0.008$), suggesting larger increases in the absolute values among men compared to women in these assessments. However, the relative improvements in walking speed, grip strength and knee extension strength were greater in the 80-year-old women than in men in either age group or the 75-year-old women.

Cohort-by-age interactions were significant for walking speed ($p = 0.035$) and grip strength ($p = 0.001$), suggesting larger increases in the absolute values among the 80-year-olds than 75-year-olds. The interaction was also significant for FEV1 ($p = 0.004$) in which only the 80-year-olds improved. Moreover, the relative percentile differences were larger among the 80-year-olds than 75-year-olds.

The linear regression models showed that the selected covariates did not fully explain the cohort differences in walking speed and muscle strength (Table 3 and 4). Better walking speed in the later cohort was partially explained by higher physical activity and longer education. The muscle strength differences in the later-born cohort were partially explained by their increased height, weight and physical activity level.

In general, the associations between birth cohort and respiratory functions attenuated after adjusting for body height and education (Table 3 and 4). The results suggest that increased body height in the later-born cohort explained a large part of the differences between the birth cohorts.

The participation rate in the later study was lower (see Figure 1). For this reason, we compared all knowledge available on non-participants. The most common reason for non-participation at both times was lack of interest or not having time to take part (eTable 1 in the Supplement). Poor health was slightly more common in the earlier cohort. In both studies, the proportions with unknown or other reasons for participation were practically identical. Information on self-rated health was available for 47% of the non-participants in the earlier cohort and for 73% of the non-participants in the later cohort. Self-rated health did not differ between the non-participants of the Evergreen and Evergreen II cohorts ($p=0.539$). The result did not change when the comparisons were made separately for age groups and sex. Overall, we observed no explicit differences between the non-participants of the earlier and later cohorts, suggesting the absence of systematic selection bias between the studies.

Discussion

We observed that the maximal physical performance of men and women aged 75 or 80 years assessed 28 years apart was markedly and meaningfully better in the later-born cohort. For grip strength, the improvements varied between 11 and 55 Newtons depending on age and sex. Inferring from a meta-analysis¹⁸ (10 Newton or 1 kg higher grip strength corresponds to 3% decline in mortality), the mortality risk of the 80-year-old men in the later cohort will be 12% lower and in the same-age women 15% lower than in the earlier cohort. The walking speed improvements ranged between 0.2 and 0.4 m/s; a 0.1 m/s improvement in walking speed corresponds to substantially better mobility³². In addition, the risk for mobility limitations due to low muscle strength is meaningfully lower in the later-born cohorts. The present results are unique in

that they derive from multiple highly relevant maximal physical performance tests assessed with identical highly standardized measures in two comparable cohorts examined approximately one generation apart. These results provide us with novel information about differences in functional aging in people growing old during different historical periods.

Various explanations can be offered for the current results. The first is that the later cohort had more propitious life course exposures that positively affected their health and functioning. The earlier cohorts were born in 1910-1915, when Finland was largely agricultural, undeveloped and still part of the Russian empire until 1917. Children worked from an early age, experienced the turmoil following the Civil War in 1918, and as young adults, they took part in the Second World War. The later cohort was born in 1938-1942. During the 1940s, many reforms were implemented, including the provision of school meals for all children free of charge and longer obligatory education. This improved the nutritional situation, especially for children from lower income homes, and delayed their entry into the labor market. These societal reforms may underlie our findings of increased height and weight in the latter cohort, which is mostly a result of better nutrition³³. With the rapid development of the country in the 1950s, access to secondary and tertiary education improved and the female disadvantage in education decreased³⁴. This is in line with the doubling of length of education between the earlier and later cohorts. Higher education is associated with better jobs, and better economic conditions and psychosocial resources and with more beneficial health behavior, all of which contribute to better health and functioning. Heavy manual work in earlier life is associated with increased risk for problems in health and functioning in older age³⁵. The regression analyses indicated that positive secular trends in the covariates were important aspects underlying improved muscle strength and walking speed in the later cohort. Longer education in men and increased leisure-time physical activity levels in both sexes were associated with better walking speed whereas increased body size and physical activity level were associated with better muscle strength. However, the association of physical activity with cohort differences in physical performance can be interpreted in two ways: high physical activity may result from better physical performance, or vice versa³⁶.

Many of the birth cohort effects remained unexplained by the variables available in our data. Other potential explanations include improved medical care and better access to health care. In addition, working conditions has improved through legislation protecting employees and improved technological solutions. However, we can probably rule out genetic differences between the cohorts as an explanation: since the resettlement of the Karelian population in Finland during the Second World War, there has been very little immigration. In addition, we do not believe that selective mortality explains the results. Mortality prior to the age 75 or 80 years was lower in the latter cohort than the earlier cohort, making the later cohort less rather than more selected.

In the lung function tests, the results were somewhat inconsistent. The later-born cohort performed better in the FVC test, which measures the total amount of air that can be forcibly exhaled. However, cohort differences in exhaled airflow were small or non-existent. In contrast to an earlier finding among 75-year-olds, we noted a positive change only among the 80-year-olds in the FEV1 test and no improvement in PEF¹⁷. In our study, cohort differences in lung function were partly explained by the greater body size in the later-born cohort. Increased education, potentially indicating better working conditions and health habits also explained better lung function in the younger cohort. Smoking is the main reason for decreased pulmonary function and chronic airway obstruction. In our study, the proportion of ever smokers in the later cohort was lower among men and higher among women, a finding in line with previous reports³⁷. However, it had only a minor impact on the cohort differences. Environmental factors pertaining to pulmonary health have possibly worsened during the past few decades due to urbanization, exposures to emissions of biomass fuels and other causes of environmental pollution. Long-term exposure to ambient air pollutants have been shown to result in impaired lung function and an increasing prevalence of obstructive lung diseases, which may explain why the improvement in FEV1 was smaller than that in FVC and not evident in the 75-year-old cohort³⁸.

The better physical performance in the later-born cohort can be explained by their slower rate-of-change with increasing age, a higher lifetime maximum in physical performance, or a combination of the two. Between the years 1989 and 2017, the remaining life expectancy in Finland has increased by around three years among 75-year-olds and two years among 80-year-olds¹. Having more years to death at these ages, together with the current results, suggests that today's older people are functionally younger than people of the same age one generation earlier. Our findings support the hypothesis of the postponement of disability to older ages³⁹, although our data cannot be used to support or reject the compression of morbidity hypothesis, which continues to be debated⁴⁰. Nevertheless, the results point toward more years spent with higher functional capacity at least among current 75- and 80-year-old adults in Finland. However, it is unclear whether this positive trend applies to younger cohorts. Beller et al.¹² showed an opposing trend in grip strength among younger cohorts, which may stem from changes in health-related lifestyles, such as increased sedentary lifestyle and obesity. In addition, differences in historical and economic developments and cultural factors may also result in mixed trends in physical performance in different countries.

The main strength of this study is the use of standardized maximal performance assessments of multiple functions conducted with identical methods 28 years apart. Muscle strength, walking speed and respiratory functions describe the intrinsic physiological capacity of older adults. Another strength of the study is that we compared men and women of exactly the same ages. Participant recruitment was also comparable in both studies. We found that non-participants did not differ between the studies in terms of self-rated health or reasons to decline participation, a finding that supports the comparability of the cohorts. However, we cannot rule out the possibility that some unmeasured influences underlying the participation rates may have affected the results. In addition, our study is unique in the length of the interval between the studies being almost three decades.

The study has also its limitations. First, the participation rate in the later study was lower than in the earlier, which could mean that the participants in the later study represent a healthier section of the target population than those in the earlier study. Because non-participants did not differ between the studies, we may assume

that the cohorts were comparable. However, it is still possible that because of the smaller participation rate, the later cohort is more selected and potentially healthier group, and we cannot completely rule out the possibility of selection bias explaining partly the results. The measurement equipment for assessing grip strength and knee extension strength was identical in both times, and methodological differences probably do not explain the observed differences between the cohorts. However, we cannot rule out the possibility that for lung function test, some systematic measurement difference may have affected the results. Another possible limitation is that the cohort differences in comorbidity could not be included in the analyses due to changes in the diagnostics, treatment and recording of chronic conditions over the past three decades. The results may be unique to Finland; however, it is likely that they can be generalized to other countries that have undergone similar societal changes during the last 100 years. We do not have data on life-course exposures earlier in the participants' lives, information which would have strengthened our conclusions on the possible reasons for the cohort differences.

To conclude, the present study suggests improved physical performance, especially in walking speed and muscle strength, in the more recent birth cohorts of 75- and 80-year-old Finnish adults. These functional traits underlie mobility, activities of daily living and participation in social life. The results may help to identify potentially unrecognized resources of older adults and encourage their continued engagement in valued activities in later life.

Contributors

K. Koivunen: Conception and design of the study, drafting the manuscript, data preparation, analysis and interpretation. E. Sillanpää: Contribution to the design of the study, data analysis and interpretation, drafting the manuscript. M. Munukka: Data preparation, analysis and interpretation, critical revision for important intellectual content. E. Portegijs: Data collection, critical revision for important intellectual content. T. Rantanen: Conception and design of the study, acquiring the funding for conducting the research, conducting the study and acquiring data, data analysis and interpretation, drafting the manuscript.

Conflicts of interest

T.R. serves on the Journal of Gerontology: Medical Sciences editorial board. Otherwise, the authors declare no conflicts of interest.

Funding

This work was supported by the European Research Council (grant number 693045 to TR) and the Academy of Finland (grant number 310526 to TR). The Evergreen 1 project has been supported by the Academy of Finland, Finnish Social Insurance Institution, Finnish Ministry of Education, Finnish Ministry of Social Affairs and Health, City of Jyväskylä, and the Association of Finnish Lion Clubs and the Scandinavian Red Feather Project. The content of this manuscript does not reflect the official opinion of the European Union. Responsibility for the information and views expressed in the manuscript lies entirely with the authors.

Accepted Manuscript

References

1. Official statistics of finland (OSF): Deaths [e-publication]. Accessed 1.8., 2019.
2. Heikkinen E, Kauppinen M, Rantanen T, et al. Cohort differences in health, functioning and physical activity in the young-old finnish population. *Aging Clin Exp Res*. 2011;23(2):126-134. <https://link-springer-com.ezproxy.jyu.fi/article/10.3275%2F6932>. doi: 10.3275/6932.
3. Beltrán-Sánchez H, Jiménez MP, Subramanian SV. Assessing morbidity compression in two cohorts from the health and retirement study. *Journal of Epidemiology and Community Health*. 2016;70(10):1011-1016. <http://dx.doi.org/10.1136/jech-2015-206722>. doi: 10.1136/jech-2015-206722.
4. Hung WW, Ross JS, Boockvar KS, Siu AL. Recent Trends in Chronic Disease, Impairment and Disability Among Older Adults in the United States. <https://www.ncbi.nlm.nih.gov/pubmed/21851629>. doi: 10.1186/1471-2318-11-47.
5. Christensen K, Thinggaard M, Oksuzyan A, et al. Physical and cognitive functioning of people older than 90 years: A comparison of two danish cohorts born 10 years apart. *Lancet, The*. 2013;382(9903):1507-1513. <https://www.clinicalkey.es/playcontent/1-s2.0-S0140673613607771>. doi: 10.1016/S0140-6736(13)60777-1.

6. Chen Y, Sloan FA. Explaining disability trends in the U.S. elderly and Near- Elderly population. *Health Services Research*. 2015;50(5):1528-1549.
<https://onlinelibrary.wiley.com/doi/abs/10.1111/1475-6773.12284>. doi: 10.1111/1475-6773.12284.
7. Hung WW, Ross JS, Boockvar KS, Siu AL. Recent trends in chronic disease, impairment and disability among older adults in the united states. *BMC geriatrics*. 2011;11(1):47.
<https://www.ncbi.nlm.nih.gov/pubmed/21851629>. doi: 10.1186/1471-2318-11-47.
8. Sarkeala T, Nummi T, Vuorisalmi M, Hervonen A, Jylhä M. Disability trends among nonagenarians in 2001–2007: Vitality 90+ study. *Eur J Ageing*. 2011;8(2):87-94. doi: 10.1007/s10433-011-0188-2.
9. Studenski S, Perera S, Patel K, et al. Gait speed and survival in older adults. *JAMA*. 2011;305(1):50-58. <http://dx.doi.org/10.1001/jama.2010.1923>. doi: 10.1001/jama.2010.1923.
10. Rantanen T, Guralnik JM, Foley D, et al. Midlife hand grip strength as a predictor of old age disability. *JAMA*. 1999;281(6):558-560. <http://dx.doi.org/10.1001/jama.281.6.558>. doi: 10.1001/jama.281.6.558.
11. Strand BH, Bergland A, Jørgensen L, Schirmer H, Emaus N, Cooper R. Do more recent born generations of older adults have stronger grip? A comparison of three cohorts of 66- to 84-year-olds in the tromsø study. *The journals of gerontology. Series A, Biological sciences and medical sciences*. 2019;74(4):528-533. <https://www.ncbi.nlm.nih.gov/pubmed/30312369>. doi: 10.1093/gerona/gly234.
12. Tomkinson GR, Kidokoro T, Dufner T, Noi S, Fitzgerald JS, Mcgrath RP. Temporal trends in handgrip strength for older japanese adults between 1998 and 2017. *Age and ageing*. 2020.
<https://www.ncbi.nlm.nih.gov/pubmed/32037458>. doi: 10.1093/ageing/afaa021.

13. Wranger LS, Elmståhl S, Ekström H. Physical performance in relation to birth cohort: A comparison of 60 year old swedish men and women born twelve years apart. *Archives of Gerontology and Geriatrics*. 2019;82:22-27.
<https://www.sciencedirect.com/science/article/pii/S016749431930010X>. doi: 10.1016/j.archger.2019.01.010.
14. Dodds R, Pakpahan E, Granic A, Davies K, Sayer A. The recent secular trend in grip strength among older adults: Findings from the english longitudinal study of ageing. *Eur Geriatr Med*. 2019;10(3):395-401. doi: 10.1007/s41999-019-00174-4.
15. Beller J, Miething A, Regidor E, Lostao L, Epping J, Geyer S. Trends in grip strength: Age, period, and cohort effects on grip strength in older adults from germany, sweden, and spain. *SSM - Population Health*. 2019;9:100456. <http://dx.doi.org/10.1016/j.ssmph.2019.100456>. doi: 10.1016/j.ssmph.2019.100456.
16. Hörder H, Skoog I, Johansson L, Falk H, Frändin K. Secular trends in frailty: A comparative study of 75-year olds born in 1911-12 and 1930. *Age and Ageing*. 2015;44(5):817-822.
<https://academic-oup-com.ezproxy.jyu.fi/ageing/article/44/5/817/52206>. doi: 10.1093/ageing/afv084.
17. Vincent Wai-Ming Lak, Ingmar Skoog, Xinxin Guo. Secular trends in lung function and its relation to survival in swedish 75 year olds 1976-2006. *Age and Ageing*. 2012;41(6):735-740.
<https://www.ncbi.nlm.nih.gov/pubmed/22843793>. doi: 10.1093/ageing/afs098.
18. Cooper R, Kuh D, Hardy R, Mortality Review Group. Objectively measured physical capability levels and mortality: Systematic review and meta-analysis. *BMJ*. 2010;341(7774):639.
<http://dx.doi.org/10.1136/bmj.c4467>. doi: 10.1136/bmj.c4467.

19. Wagner K, Cameron-Smith D, Wessner B, Franzke B. Biomarkers of aging: From function to molecular biology. *Nutrients*. 2016;8(6):338. <https://www.ncbi.nlm.nih.gov/pubmed/27271660>. doi: 10.3390/nu8060338.
20. Newman AB, Sanders JL, Kizer JR, et al. Trajectories of function and biomarkers with age: The CHS all stars study. *International journal of epidemiology*. 2016;45(4):1135-1145. <https://www.ncbi.nlm.nih.gov/pubmed/27272182>. doi: 10.1093/ije/dyw092.
21. Sillanpaa E, Stenroth L, Bijlsma AY, et al. Associations between muscle strength, spirometric pulmonary function and mobility in healthy older adults. *AGE*. 2014;36(4):1-11. <https://www.narcis.nl/publication/RecordID/oai:research.vu.nl:publications%2Fdf80b4d6-6076-4831-ae5c-dd7f02716c1d>. doi: 10.1007/s11357-014-9667-7.
22. Rantanen T, Era P, Heikkinen E. Maximal isometric strength and mobility among 75-year-old men and women. *Age Ageing*. 1994;23(2):132-137. doi: 10.1093/ageing/23.2.132 [doi].
23. Rantanen T, Era P, Heikkinen E. Physical activity and the changes in maximal isometric strength in men and women from the age of 75 to 80 years. *J Am Geriatr Soc*. 1997;45(12):1439-1445. doi: 10.1111/j.1532-5415.1997.tb03193.x [doi].
24. Heikkinen E. Background, design, and methods of the evergreen project. *Journal of Aging and Physical Activity*. 1998;6(2):106-120.
25. Rantanen T, Saajanaho M, Karavirta L, et al. Active aging – resilience and external support as modifiers of the disablement outcome: AGNES cohort study protocol. *BMC public health*. 2018;18(1):565-21. <https://www.ncbi.nlm.nih.gov/pubmed/29716566>. doi: 10.1186/s12889-018-5487-5.

26. Rantanen T, Era P, Heikkinen E. Physical activity and the changes in maximal isometric strength in men and women from the age of 75 to 80 years. *Journal of the American Geriatrics Society*. 1997;45(12):1439-1445. <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1532-5415.1997.tb03193.x>. doi: 10.1111/j.1532-5415.1997.tb03193.x.
27. Miller MR, Hankinson J, Brusasco V, et al. Standardisation of spirometry. *European Respiratory Journal*. 2005;26(2):319-338. <http://erj.ersjournals.com/cgi/content/abstract/26/2/319>. doi: 10.1183/09031936.05.00034805.
28. Fonseca R, Michaud P, Zheng Y. The effect of education on health: Evidence from national compulsory schooling reforms. *SERIEs*. 2020;11(1):83-103. <https://search.proquest.com/docview/2267589569>. doi: 10.1007/s13209-019-0201-0.
29. Samson MM, Meeuwse IB, Crowe A, Dessens JA, Duursma SA, Verhaar HJ. Relationships between physical performance measures, age, height and body weight in healthy adults. *Age and ageing*. 2000;29(3):235-242. <https://www.ncbi.nlm.nih.gov/pubmed/10855906>. doi: 10.1093/ageing/29.3.235.
30. Talaminos Barroso A, Márquez-Martín E, Roa Romero L, Ortega F. Factors affecting lung function: A review of the literature. *Archivos de bronconeumologia*. 2018;54(6):327-332. <https://search.proquest.com/docview/2010374439>. doi: 10.1016/j.arbres.2018.01.030.
31. Sallinen J, Stenholm S, Rantanen T, Heliövaara M, Sainio P, Koskinen S. Hand- Grip strength cut points to screen older persons at risk for mobility limitation. *Journal of the American Geriatrics Society*. 2010;58(9):1721-1726. <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1532-5415.2010.03035.x>. doi: 10.1111/j.1532-5415.2010.03035.x.

32. Perera S, Mody SH, Woodman RC, Studenski SA. Meaningful change and responsiveness in common physical performance measures in older adults. *J Am Geriatr Soc.* 2006;54(5):743-749. doi: JGS701 [pii].
33. Cole TJ. The secular trend in human physical growth: A biological view. *Economics and Human Biology.* 2003;1(2):161-168.
<https://www.sciencedirect.com/science/article/pii/S1570677X02000333>. doi: 10.1016/S1570-677X(02)00033-3.
34. Breen R, Luijkx R, Müller W, Pollak R. Long-term trends in educational inequality in europe: Class inequalities and gender differences 1. *European Sociological Review.* 2010;26(1):31-48. doi: 10.1093/esr/jcp001.
35. Kulmala J, Ngandu T, Pajala S, et al. Leisure-time and occupational physical activity in early and late adulthood in relation to later life physical functioning. *Journal of physical activity & health.* 2016;13(10):1079-1087. <https://www.ncbi.nlm.nih.gov/pubmed/27254853>. doi: 10.1123/jpah.2015-0664.
36. Rantanen T, Guralnik JM, Sakari-Rantala R, et al. Disability, physical activity, and muscle strength in older women: The women's health and aging study. *Archives of Physical Medicine and Rehabilitation.* 1999;80(2):130-135.
<https://www.sciencedirect.com/science/article/pii/S0003999399901090>. doi: 10.1016/S0003-9993(99)90109-0.
37. Fred C. Pampel. Cigarette use and the narrowing sex differential in mortality. *Population and Development Review.* 2002;28(1):77-104. <https://www.jstor.org/stable/3092758>. doi: 10.1111/j.1728-4457.2002.00077.x.

38. Wang M, Aaron CP, Madrigano J, et al. Association between long-term exposure to ambient air pollution and change in quantitatively assessed emphysema and lung function. *JAMA*. 2019;322(6):546-556. doi: 10.1001/jama.2019.10255 [doi].
39. Manton KG. Changing concepts of morbidity and mortality in the elderly population. *Milbank Mem Fund Q Health Soc*. 1982;60(2):183-244.
40. Chatterji S, Byles J, Cutler D, Seeman T, Verdes E. Health, functioning, and disability in older adults--present status and future implications. *Lancet*. 2015;385(9967):563-575. doi: 10.1016/S0140-6736(14)61462-8 [doi].

Accepted Manuscript

Captions for Tables and Illustrations

Figure 1. Flow Chart for Evergreen and Evergreen II.

Table 1. Descriptive statistics and cohort differences of 75- and 80-year-old men and women born in 1910 and 1914 (Evergreen cohort) and born in 1938-1939 and 1942-1943 (Evergreen II cohort).

Table 2. Cohort differences in physical performance of 75- and 80-year-old men and women born in 1910 and 1914 (Evergreen cohort) and born in 1938-1939 and 1942-1943 (Evergreen II cohort).

Table 3. Linear regression of the association between the birth cohort and physical performance measures in men.

Table 4. Linear regression of the association between the birth cohort and physical performance measures in women.

Accepted Manuscript

Table 1. Descriptive statistics and cohort differences of 75- and 80-year-old men and women born in 1910 and 1914 (Evergreen cohort) and born in 1938-1939 and 1942-1943 (Evergreen II cohort).

	75 years		Cohort	80 years		Cohort
	Evergreen	Evergreen II	difference	Evergreen	Evergreen II	difference
			p-value ^a			p-value ^a
Men, n	104	183		60	132	
Age, mean (SD)	75.3 (0.3)	75.4 (0.4)	0.171	79.6 (0.3)	79.6 (0.4)	0.626
Years of education, mean (SD)	6.2 (3.5)	12.2 (4.4)	<0.001	5.9 (4.1)	11.9 (4.4)	<0.001
Height, cm, mean (SD)	169.5 (6.2)	172.7 (6.0)	<0.001	169.1 (6.5)	172.3 (6.1)	0.001
Weight, kg, mean (SD)	74.1 (10.7)	80.4 (13.0)	<0.001	75.3 (12.8)	80.1 (12.6)	0.017
Self-rated health, n (%)			<0.001			<0.001
Very good / Good	13 (13)	106 (58)		10 (17)	60 (46)	
Average	79 (76)	72 (39)		35 (59)	69 (52)	
Poor / Very poor	12 (12)	5 (3)		14 (24)	3 (2)	

Physical activity, n (%)			<0.001			<0.001
Low	30 (29)	10 (6)		24 (40)	15 (12)	
Moderate	62 (60)	127 (70)		33 (55)	94 (72)	
High	12 (12)	44 (24)		3 (5)	21 (16)	
Smoking, n (%)						
Never	33 (34)	87 (48)	0.027	19 (33)	72 (56)	0.004
Current / Former	64 (66)	95 (52)		38 (67)	56 (44)	
Women, n	191	251		145	160	
Age, mean (SD)	75.3 (0.3)	75.4 (0.4)	0.557	79.6 (0.3)	79.7 (0.4)	0.517
Years of education, mean (SD)	6.1 (3.2)	12.0 (4.1)	<0.001	5.7 (3.1)	11.8 (6.2)	<0.001
Height, cm, mean (SD)	155.8 (5.6)	159.4 (5.1)	<0.001	155.5 (5.4)	158.2 (5.5)	<0.001
Weight, kg, mean (SD)	67.5 (11.6)	71.1 (12.4)	0.002	64.5 (10.2)	69.7 (12.0)	<0.001
Self-rated health, n (%)			<0.001			<0.001
Very good / Good	27 (14)	137 (55)		18 (13)	67 (42)	
Average	139 (73)	108 (43)		93 (65)	85 (53)	
Poor / Very poor	25 (13)	6 (2)		33 (23)	8 (5)	
Physical activity, n (%)			<0.001			<0.001
Low	42 (22)	29 (12)		48 (34)	21 (13)	
Moderate	139 (74)	190 (76)		92 (65)	120 (76)	

High	7 (4)	30 (12)	2 (1)	18 (11)
Smoking, n (%)			0.005	0.051
Never	167 (90)	201 (80)	133 (93)	136 (86)
Current / Former	18 (10)	49 (20)	10 (7)	22 (14)

Note: ^a t-test for continuous variables and chi-square test for categorical variables.

Accepted Manuscript

Table 2. Cohort differences in physical performance of 75- and 80-year-old men and women born in 1910 and 1914 (Evergreen cohort) and born in 1938-1939 and 1942-1943 (Evergreen II cohort).

	75 years				80 years			
	Evergreen		Evergreen II		Evergreen		Evergreen II	
	Mean (SD)	Mean (SD)	% (95% CI)	p-value ^a	Mean (SD)	Mean (SD)	% (95% CI)	p-value ^a
Men								
Walking speed, m/s	1.8 (0.5)	2.0 (0.4)	11 (5, 18)	<0.001	1.5 (0.5)	1.8 (0.4)	20 (11, 33)	<0.001
Grip strength, N	374 (89)	406 (74)	9 (3, 14)	0.001	309 (80)	364 (74)	18 (10, 26)	<0.001
Knee extension strength, N	362 (99)	452 (102)	25 (18, 32)	<0.001	332 (73)	397 (101)	20 (12, 27)	<0.001

FVC, l	2.9 (0.7)	3.4 (0.8)	17 (12, 24)	<0.001	2.6 (0.7)	3.3 (0.6)	27 (15, 30)	<0.001
FEV1, l	2.5 (0.6)	2.6 (0.6)	4 (-0.7, 12)	0.081	2.2 (0.6)	2.5 (0.5)	14 (9, 25)	<0.001
PEF, l/s	7.4 (2.0)	7.4 (2.0)	0 (-7, 6)	0.841	6.9 (2.1)	7.4 (1.7)	7 (1, 17)	0.093
Increased risk for mobility limitation, n (%)	48 (47.5%)	48 (26.5%)		<0.001	39 (70.9%)	68 (51.9%)		0.012
Women								
Walking speed, m/s	1.5 (0.4)	1.7 (0.3)	18 (13, 22)	<0.001	1.2 (0.3)	1.6 (0.3)	33 (22, 34)	<0.001
Grip strength, N	227 (58)	238 (52)	5 (0.2, 9)	0.042	172 (55)	215 (44)	25 (19, 32)	<0.001
Knee extension strength, N	241 (73)	302 (81)	25 (19, 31)	<0.001	188 (63)	277 (82)	47 (38, 56)	<0.001
FVC, l	2.2 (0.5)	2.5 (0.5)	14 (12, 20)	<0.001	1.9 (0.5)	2.3 (0.4)	21 (18, 29)	<0.001
FEV1, l	1.9 (0.4)	1.9 (0.4)	0 (-4, 4)	0.825	1.6 (0.4)	1.8 (0.3)	12 (4, 14)	0.001

PEF, l/s	5.0 (1.2)	4.9 (1.1)	-2 (-6, 3)	0.544	4.8 (1.4)	5.0 (1.3)	4 (-3, 10)	0.318
Increased risk for mobility limitation, n (%)	67 (35.4%)	69 (27.7%)		0.052	104 (75.4%)	70 (44.3%)		<0.001

Note. CI, confidence interval; FVC, forced vital capacity; FEV1, forced expiratory volume in 1 second; PEF, peak expiratory flow. ^a t-test

Table 3. Linear regression of the association between the birth cohort and physical performance measures in men.

	Men 75 years			Men 80 years		
	Birth cohort		Model	Birth cohort		Model
	β (SE)	p-value	Adjusted r^2	β (SE)	p-value	Adjusted r^2
Walking speed, m/s						
Birth cohort	0.211 (0.055)	<0.001	0.046	0.330 (0.075)	<0.001	0.089
+ education	0.029 (0.066)	0.663	0.105	0.226 (0.089)	0.012	0.106
+ height	0.189 (0.057)	0.001	0.051	0.294 (0.077)	<0.001	0.097
+ weight	0.256 (0.056)	<0.001	0.076	0.346 (0.076)	<0.001	0.093
+ PA	0.073 (0.054)	0.176	0.211	0.174 (0.072)	0.016	0.275
+ smoking	0.181 (0.057)	0.002	0.049	0.271 (0.077)	0.001	0.098
Grip strength, N						
Birth cohort	32.2 (9.9)	0.001	0.033	54.9 (12.2)	<0.001	0.094
+ education	19.1 (12.2)	0.118	0.036	51.4 (14.4)	<0.001	0.095
+ height	19.3 (9.9)	0.052	0.108	41.5 (11.8)	0.001	0.193
+ weight	23.8 (10.0)	0.018	0.069	46.4 (12.0)	<0.001	0.147
+ PA	16.7 (10.2)	0.104	0.091	50.0 (12.6)	<0.001	0.109
+ smoking	25.2 (10.1)	0.016	0.025	49.5 (12.7)	<0.001	0.092
Knee extension strength, N						

Birth cohort	89.4 (12.6)	<0.001	0.151	65.0 (15.1)	<0.001	0.087
+ education	85.1 (15.4)	<0.001	0.142	58.4 (17.6)	0.001	0.093
+ height	81.9 (12.9)	<0.001	0.163	58.7 (15.4)	<0.001	0.101
+ weight	83.0 (12.9)	<0.001	0.161	58.3 (15.1)	<0.001	0.114
+ PA	68.7 (12.9)	<0.001	0.209	54.1 (15.2)	<0.001	0.133
+ smoking	85.0 (13.0)	<0.001	0.131	56.8 (15.3)	<0.001	0.100
FVC, l						
Birth cohort	0.521 (0.091)	<0.001	0.103	0.604 (0.103)	<0.001	0.158
+ education	0.378 (0.111)	0.047	0.108	0.584 (0.122)	<0.001	0.157
+ height	0.391 (0.089)	<0.001	0.199	0.451 (0.099)	<0.001	0.279
+ weight	0.562 (0.093)	<0.001	0.117	0.580 (0.105)	<0.001	0.165
+ PA	0.417 (0.096)	<0.001	0.133	0.510 (0.108)	<0.001	0.192
+ smoking	0.485 (0.095)	<0.001	0.097	0.548 (0.105)	<0.001	0.171
FEV1, l						
Birth cohort	0.135 (0.077)	0.081	0.007	0.364 (0.086)	<0.001	0.091
+ education	-0.011 (0.094)	0.905	0.025	0.356 (0.103)	<0.001	0.084
+ height	0.037 (0.076)	0.631	0.093	0.242 (0.084)	0.004	0.205
+ weight	0.161 (0.079)	0.044	0.010	0.334 (0.088)	<0.001	0.093
+ PA	0.048 (0.082)	0.556	0.032	0.282 (0.091)	0.002	0.117
+ smoking	0.097 (0.080)	0.223	0.022	0.282 (0.087)	0.001	0.128
PEF, l/s						

Birth cohort	-0.049 (0.244)	0.841	0.000	0.533 (0.292)	0.069	0.013
+ education	-0.483 (0.296)	0.104	0.011	0.515 (0.346)	0.138	0.009
+ height	-0.307 (0.244)	0.209	0.055	0.265 (0.295)	0.370	0.067
+ weight	-0.028 (0.251)	0.912	0.000	0.390 (0.294)	0.186	0.042
+ PA	-0.437 (0.254)	0.087	0.054	0.276 (0.310)	0.373	0.029
+ smoking	-0.158 (0.252)	0.531	0.004	0.397 (0.102)	0.188	0.014

Note. Each covariate was added in the model one at a time with birth cohort; β , unstandardized beta indicates mean cohort difference (reference group Evergreen cohort); SE, standard error; PA, Physical activity; FVC, forced vital capacity, FEV1, forced expiratory volume in 1 second; PEF, peak expiratory flow.

Accepted Manuscript

Table 4. Linear regression of the association between the birth cohort and physical performance measures in women.

	Women 75 years			Women 80 years		
	Birth cohort		Model	Birth cohort		Model
	β (SE)	p-value	Adjusted r^2	β (SE)	p-value	Adjusted r^2
Walking speed, m/s						
Birth cohort	0.261 (0.033)	<0.001	0.122	0.349 (0.038)	<0.001	0.224
+ education	0.146 (0.041)	<0.001	0.151	0.318 (0.044)	<0.001	0.231
+ height	0.240 (0.035)	<0.001	0.127	0.329 (0.039)	<0.001	0.228
+ weight	0.274 (0.032)	<0.001	0.213	0.398 (0.037)	<0.001	0.287
+ PA	0.205 (0.031)	<0.001	0.275	0.274 (0.036)	<0.001	0.347
+ smoking	0.275 (0.034)	<0.001	0.138	0.354 (0.038)	<0.001	0.224
Grip strength, N						
Birth cohort	10.8 (5.31)	0.042	0.007	43.5 (5.8)	<0.001	0.158
+ education	9.09 (6.76)	0.181	0.003	41.9 (7.3)	<0.001	0.161
+ height	-0.09 (5.40)	0.987	0.084	35.8 (5.7)	<0.001	0.227
+ weight	8.71 (5.33)	0.103	0.022	37.0 (5.8)	<0.001	0.206
+ PA	6.18 (5.37)	0.251	0.036	38.4 (5.9)	<0.001	0.191
+ smoking	11.3 (5.4)	0.038	0.006	43.2 (5.9)	<0.001	0.153
Knee extension strength, N						
Birth cohort	60.4 (7.5)	<0.001	0.127	88.7 (8.6)	<0.001	0.264


+ education	55.0 (9.6)	<0.001	0.122	99.2 (10.1)	<0.001	0.271
+ height	49.7 (7.8)	<0.001	0.159	82.6 (8.8)	<0.001	0.276
+ weight	55.5 (7.4)	<0.001	0.165	83.5 (8.8)	<0.001	0.272
+ PA	54.4 (7.6)	<0.001	0.150	77.1 (8.7)	<0.001	0.308
+ smoking	61.1 (7.6)	<0.001	0.127	87.6 (8.7)	<0.001	0.256
FVC, l						
Birth cohort	0.342 (0.046)	<0.001	0.115	0.439 (0.052)	<0.001	0.195
+ education	0.280 (0.059)	<0.001	0.115	0.340 (0.066)	<0.001	0.201
+ height	0.199 (0.043)	<0.001	0.289	0.356 (0.050)	<0.001	0.307
+ weight	0.373 (0.046)	<0.001	0.113	0.441 (0.054)	<0.001	0.193
+ PA	0.304 (0.046)	<0.001	0.154	0.415 (0.054)	<0.001	0.189
+ smoking	0.341 (0.047)	<0.001	0.111	0.434 (0.053)	<0.001	0.189
FEV1, l						
Birth cohort	0.008 (0.038)	0.825	0.000	0.148 (0.043)	0.001	0.036
+ education	-0.024 (0.048)	0.618	0.000	0.086 (0.055)	0.123	0.044
+ height	-0.086 (0.037)	0.023	0.122	0.088 (0.042)	0.039	0.134
+ weight	0.008 (0.038)	0.835	0.000	0.142 (0.045)	0.002	0.033
+ PA	-0.023 (0.038)	<0.001	0.036	0.133 (0.045)	0.004	0.029
+ smoking	0.019 (0.039)	0.020	0.008	0.148 (0.044)	0.001	0.032
PEF, l/s						
Birth cohort	-0.076 (0.122)	0.535	0.000	0.156 (0.156)	0.318	0.000

+ education	-0.241 (0.156)	0.124	0.002	0.049 (0.201)	0.809	-0.001
+ height	-0.277 (0.125)	0.028	0.052	0.048 (0.160)	0.763	0.022
+ weight	-0.101 (0.123)	0.412	0.001	0.125 (0.162)	0.441	-0.002
+ PA	-0.145 (0.125)	0.246	0.009	0.031 (0.161)	0.849	0.015
+ smoking	-0.050 (0.124)	0.688	0.001	0.163 (0.160)	0.308	-0.003

Note. Each covariate was added in the model one at a time with birth cohort; β , unstandardized beta indicates mean cohort difference (reference group Evergreen cohort); SE, standard error; PA, Physical activity; FVC, forced vital capacity, FEV1, forced expiratory volume in 1 second; PEF, peak expiratory flow

Accepted Manuscript

Figure 1



Accepted Manuscript