

This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Luisto, Rami; Pankka, Pekka

Title: Stoïlow's theorem revisited

Year: 2020

Version: Accepted version (Final draft)

Copyright: © 2019 Elsevier GmbH. All rights reserved.

Rights: CC BY-NC-ND 4.0

Rights url: https://creativecommons.org/licenses/by-nc-nd/4.0/

Please cite the original version:

Luisto, R., & Pankka, P. (2020). Stoïlow's theorem revisited. Expositiones Mathematicae, 38(3), 303-318. https://doi.org/10.1016/j.exmath.2019.04.002

STOÏLOW'S THEOREM REVISITED

RAMI LUISTO AND PEKKA PANKKA

ABSTRACT. Stoïlow's theorem from 1928 states that a continuous, open, and light map between surfaces is a discrete map with a discrete branch set. This result implies that such maps between orientable surfaces are locally modeled by power maps $z \mapsto z^k$ and admit a holomorphic factorization.

The purpose of this expository article is to give a proof of this classical theorem having readers in mind that are interested in continuous, open and discrete maps.

1. INTRODUCTION

Stoïlow's classical theorem in [16] states that a continuous, open and light map between surfaces is a discrete map which has a discrete branch set. In what follows, we call this theorem Stoilow's discreteness theorem.

Recall that a continuous map $f: X \to Y$ between topological spaces is *light* if the pre-image $f^{-1}(y)$ of each point $y \in Y$ is totally disconnected, and *discrete* if $f^{-1}(y)$ is a discrete subset of X. A continuous map is *open* if the image of each open set is an open set. The *branch set* B_f of a continuous map $f: X \to Y$ is the set of points $x \in X$ at which f fails to be a local homeomorphism.

In [16] Stoïlow shows that these maps are locally modeled by power maps $z \mapsto z^k$; see [16, p. 372] for the discussion. This local description indicates a deep connection between continuous, open and light maps and holomorphic maps between surfaces. This connection was found by Stoïlow [17, p. 120] and Whyburn [19, Theorem X.5.1, p. 198] and [20, p. 103]: For a continuous, open and light map $f: S \to S'$ between orientable Riemann surfaces there exists a Riemann surface \tilde{S} and a homeomorphism $h: S \to \tilde{S}$ for which $f \circ h^{-1}: \tilde{S} \to S'$ is a holomorphic map; the Riemann surface \tilde{S} in this statement is naturally the Riemann surface associated to the map f. The first edition of [20], published in 1956, does not give this result a specific name, but already in the second edition from 1964 the result is referred as Stoilow's theorem.

In this expository article we discuss the proof of Stoïlow's discreteness theorem having readers in mind that are interested in discrete and open maps,

Date: April 1, 2019.

²⁰¹⁰ Mathematics Subject Classification. 30-02.

R.L. has been partially supported by a grant of the Finnish Academy of Science and Letters, the Academy of Finland (grant 288501 'Geometry of subRiemannian groups') and by the European Research Council (ERC Starting Grant 713998 GeoMeG 'Geometry of Metric Groups').

P.P. has been partially supported by the Academy of Finland projects #256228 and #297258.

such as quasiregular maps (see e.g. [15]) or Thurston maps (see e.g. [4]). For this reason we separate Stoïlow's theorem into two parts: the discreteness of the map and the discreteness of the branch set.

Theorem 1.1 (Stoïlow, 1928). Let Ω be a domain in \mathbb{C} and $f: \Omega \to \mathbb{C}$ a continuous, open and light map. Then f is a discrete map.

Theorem 1.2 (Stoïlow, 1928). Let Ω be a domain in \mathbb{C} and $f: \Omega \to \mathbb{C}$ a continuous, open and discrete map. Then B_f is a discrete set.

It is interesting to notice that both results stem from path-lifting arguments. Indeed, after path-lifting results are established, standard applications of the Jordan curve theorem yield both Theorems 1.1 and 1.2. For continuous, open and discrete maps, we may use Rickman's path-lifting theorem [14] and for continuous, open and light maps the method of Floyd [7]. Floyd's method suffices for all our purposes and we recall it in Section 3; see [11] for a more detailed discussion on path-lifting methods.

Having Theorems 1.1 and 1.2 at our disposal, it is a straightforward covering-space argument to show that continuous, open and light maps between surfaces are locally modeled by power maps.

Theorem 1.3 (Stoïlow, 1928). Let $f: \Sigma \to \Sigma'$ be a continuous, open and light map between surfaces. For each $x \in \Sigma$, there exists $k \in \mathbb{N}$, a neighborhood U of z, and homeomorphisms $\psi: U \to \mathbb{D}$ and $\phi: fU \to \mathbb{D}$ for which $\psi(x) = 0, \ \phi(f(x)) = 0$, and the diagram



commutes.

Remark 1.4. For maps between oriented surfaces, we may alternatively state that there exist orientation-preserving homeomorphisms $\psi: U \to \mathbb{D}$ and $\phi: fU \to \mathbb{D}$ for which the local model for f is $z \mapsto z^k$, if f is orientationpreserving, and $z \mapsto \overline{z}^k$ if f is orientation-reversing; see the proof of Theorem 1.3 in Section 5.

This local version of Stoïlow's discreteness theorem yields the following global factorization theorem. Indeed, by Theorem 1.3, a continuous, light and open map $f: \Sigma \to S'$ induces a conformal structure on Σ making it a Riemann surface \tilde{S} ; we refer to [4, Lemma A.10] for a short proof. Thus we may consider f as a holomorphic map $f: \tilde{S} \to S'$. If the surface Σ a priori carries a conformal structure, and we consider Σ as a Riemann surface S, we may take the homeomorphism h in the factorization to be the identity homeomorphism $\Sigma \to \Sigma$.

Theorem (Stoïlow's factorization theorem, 1938). Let $f: S \to S'$ be a continuous, open and light map between Riemann surfaces. Then there exists a Riemann surface \tilde{S} and a homeomorphism $h: S \to \tilde{S}$ such that $f \circ h^{-1}: \tilde{S} \to S'$ is a holomorphic map.

Recall that an orientable topological surface carries a conformal structure. Indeed, by a classical theorem of Radó, every 2-manifold can be triangulated (see e.g. [12, Theorem 8.3, p. 60] or [1, Section II.8, p. 105]) and every triangulated orientable surface has a conformal structure (see e.g. [1, II.2.5E, Theorem, p. 127] or [3, Section 2.2, pp. 9–11]). In this way we recover the interpretation that a continuous, open and light map between orientable surfaces is a holomorphic map between Riemann surfaces.

As the authors' interest for these theorems of Stoïlow stems from their role in the theory of quasiregular maps, we finish this introduction with a related remark. In the quasiconformal literature Stoïlow's theorem typically refers to the result that each quasiregular map $S \to S'$ between Riemann surfaces factors into a holomorphic map $S \to S'$ and a quasiconformal homeomorphism $S \to S$. The proof of this result is analytic in its nature and based on the Beltrami equation. We refer to Astala, Iwaniec, and Martin [2, Theorem 5.5.1] for a detailed discussion.

Acknowledgments. The authors would like to thank Mario Bonk for his encouragement to write this expository article and for several helpful remarks. We are also grateful to the anonymous referee for many comments which considerably improved the exposition.

2. Preliminaries

In the complex plane \mathbb{C} we denote the unit ball as \mathbb{D} and the unit circle as \mathbb{S}^1 . For a path $c: [0,1] \to \mathbb{C}$ we set |c| to be the image set of c. An injective path $\beta: [0,1] \to \mathbb{C}$ is called an *arc*. For a set $A \subset \mathbb{C}$ and a radius r > 0 we denote by B(A,r) the set of points whose distance from A is strictly less than r.

A set is called *totally disconnected* if its connected components are points. A totally disconnected closed set in the plane has topological dimension zero. We recall this fact as the following lemma; see [10, Section II.2, p. 14] for a proof.

Lemma 2.1. Let C be a closed and totally disconnected set in \mathbb{C} . Then every point $x \in C$ has a neighborhood basis consisting of neighborhoods U satisfying $C \cap \partial U = \emptyset$.

Let Ω be a planar domain and $f: \Omega \to \mathbb{C}$ be a continuous, open and light map. A domain $U \subset \Omega$ is a normal domain (for f) if U is compactly contained in Ω and $\partial f U = f \partial U$. For each $x \in \Omega$ and r > 0, we denote by U(x, f, r) the component of $f^{-1}B(f(x), r)$ containing the point x. The following lemma shows that domains U(x, f, r) are normal domains of x for all r > 0 small enough. The proof here is essentially identical to the proof in the case when f is discrete; see e.g. [18, Lemma 5.1] where the discreteness is used essentially only to guarantee that Lemma 2.1 holds.

Lemma 2.2. Let Ω be a planar domain, $f: \Omega \to \mathbb{C}$ a continuous, open and light map, and let $x \in \Omega$. Then there exists a radius $r_x > 0$ such that for all $r \leq r_x$ the domain U(x, f, r) is a normal domain containing x and fU(x, f, r) = B(f(x), r). Proof. Let $x \in \Omega$. By Lemma 2.1 there exists a precompact neighborhood V of x for which $\partial V \cap f^{-1}{f(x)} = \emptyset$. Since ∂V is compact and does not contain pre-images of x, we may fix $r_x > 0$ for which $B(f(x), r_x) \cap f \partial V = \emptyset$.

Let $0 < r < r_x$ and set U := U(x, f, r). We show first that U is a precompact domain contained in V. Since

$$f(U \cap \partial V) \subset fU \cap f\partial V \subset B(f(x), r) \cap f\partial V = \emptyset,$$

we have $U \cap \partial V = \emptyset$. Since U is a connected neighborhood of x by definition, we therefore have that $U \subset V$. Thus \overline{U} is compact as a closed subset of the compact set \overline{V} .

We claim next that $\partial f U = f \partial U$. To show the inclusion $\partial f U \subset f \partial U$, we note first that, by precompactness of U and continuity of f, we have that $\overline{fU} = f\overline{U}$. On the other hand, since f is open and U is a domain, both Uand fU are open sets. Thus

$$\partial fU = \overline{fU} \setminus fU = (f\overline{U}) \setminus fU \subset f(\overline{U} \setminus U) = f\partial U.$$

This proves the first inclusion.

We show now the second inclusion $f\partial U \subset \partial fU$. Since f is continuous, we have $f\partial U \subset f\overline{U} \subset \overline{fU}$ and so it suffices to show that $fU \cap f\partial U = \emptyset$. Suppose towards contradiction that $fU \cap f\partial U \neq \emptyset$ and let $z \in fU \cap f\partial U$. Since f is an open map and U is a domain there exists a radius s > 0 such that $B(z, s) \subset fU$.

Fix $w \in (\partial U) \cap f^{-1}\{z\}$. Since f is continuous, $f^{-1}B(x,r)$ is open and thus intersects U as $w \in (f^{-1}B(x,r)) \cap \partial U$. In particular, the *w*-component of $f^{-1}B(x,r)$, denoted W, intersects U. Now, as the open set U is the *x*component of $f^{-1}B(x,r) \supset f^{-1}B(z,s)$, it must in fact contain W, since the components of nested sets are also nested. This implies that $w \notin \partial U$, which is a contradiction. Thus $f\partial U \subset \partial fU$ and we conclude that $f\partial U = \partial fU$. Thus U is a normal domain.

Finally, we wish to show that fU = B(f(x), r). To this end we first note that since f is an open map and U a domain, fU is an open set in B(f(x), r). Next we show that fU must also be closed in B(f(x), r). Suppose towards contradiction that there exists a point $z \in B(f(x), r) \cap \partial fU$. Since $\partial fU =$ $f\partial U$, we may fix a point $w \in \partial U$ with f(w) = z. Now for s < r - d(f(x), z)we see that the w-component V of $f^{-1}B(z, s)$ is a neighborhood of w that intersects U. Since

$$V \subset f^{-1}B(z,s) \subset f^{-1}B(x,r)$$

and both U and V are connected we must have by the definition of U that $V \subset U$. But this is a contradiction since $w \in \partial U$, but V is now a neighborhood of w in U. Thus we conclude that fU is both open and closed in B(f(x), r), and so in particular fU = B(f(x), r).

We show next that, when restricted to a normal domain, a continuous, open and light map is a proper map and furthermore the components of the pre-image of a compact set map surjectively to that compact set. For continuous, open and discrete maps this claim is straightforward to prove, but for continuous, open and light maps the proof is more involved. The proofs of the following two lemmas are a special case of a more general result of Whyburn; see [19, Theorem 7.4, p. 147 and Theorem 7.5, p. 148].

Lemma 2.3. Let U be a normal domain of a continuous, open and light map $f: \Omega \to \mathbb{C}$. Then the restriction $f|_U: U \to fU$ is a proper map.

Proof. Let $K \subset fU$ be a compact set. Suppose towards contradiction that $U \cap f^{-1}K$ is not compact. Then there exists a sequence (x_j) in $U \cap f^{-1}K$ that has no subsequence converging in $U \cap f^{-1}K$. Since \overline{U} is compact, the sequence (x_j) does have a subsequence (y_j) converging to a point $y_0 \in \overline{U}$. Since $(f^{-1}K) \cap U$ is closed in U, we conclude that $y_0 \in \partial U$. Now by the continuity of f and closedness of K, we have that $f(y_0) \in K$. Since $y_0 \in \partial U$ and U is a normal domain, this implies that $K \cap \partial fU \neq \emptyset$, which is a contradiction, since K is a compact subset of the domain fU. We conclude that $f|_U$ is a proper map.

Lemma 2.4. Let U be a normal domain of a continuous, open and light map $f: \Omega \to \mathbb{C}$. Then for any compact connected set $K \subset fU$, and each component E of $(f|_U)^{-1}K$, the restriction $f|_E: E \to K$ is surjective.

Proof. Let $K \subset fU$ be a compact and connected set, and let $\tilde{E} = (f|_U)^{-1}K$. We note that, since f is open, also the restriction $f|_{\tilde{E}} \colon \tilde{E} \to K$ is open.

Suppose towards contradiction that there exists a component $E \subset \tilde{E}$ and a point $p \in K$ for which $p \notin fE$. Since $f|_U \colon U \to fU$ is a proper map, the pre-image $P = U \cap f^{-1}\{p\}$ is a compact set which does not intersect E.

Since \tilde{E} is a compact subset of the plane, it is especially a compact Hausdorff space. Thus the components of \tilde{E} equal the quasicomponents of \tilde{E} , see e.g. [6, Theorem 6.1.23]; recall that the quasicomponent of a point is the intersection of all open and closed sets containing that point. Thus we may choose, for every $x \in P$, a separation $\{A_x, B_x\}$ of the set \tilde{E} , i.e., two disjoint open sets $A_x, B_x \subset U$ covering \tilde{E} for which $E \subset A_x$ and $x \in B_x$. Since P is compact, the cover $\{B_x : x \in P\}$ has a finite subcover $\{B_{x_1}, \ldots, B_{x_n}\}$.

For each j = 1, ..., n, the open set A_{x_j} contains the set E and satisfies $A_{x_j} \cap B_{x_j} = \emptyset$. Thus the intersection

$$A = \bigcap_{j=1}^{n} A_{x_j}$$

is a neighborhood of E satisfying $A \cap P = \emptyset$ and $E \cap \partial A = \emptyset$. Now fA is an open set in fU that contains fE but does not contain p. Since K is a connected set that intersects both fA and its complement $CfA \ni p$, we have $(\partial fA) \cap K \neq \emptyset$.

Finally, on the one hand, by the openness of f we have $\partial f A \subset f \partial A$. On the other hand, due to the definition of the sets A_j and thus of A, $(\partial A) \cap f^{-1}K = \emptyset$. This is a contradiction and so the original claim holds. \Box

3. Path lifting after Floyd and Stoïlow

In this section we discuss path-lifting, which is one of the main tools in the forthcoming proofs. The following theorem is, essentially, due to Stoïlow [16, pp. 354–358] and its idea was generalized by Floyd [7] to the setting of compact metric spaces. We include here a version of Floyd's proof in the planar setting for the sake of completeness of the exposition; note that the proof would go through also in the setting of locally compact metric spaces. **Theorem 3.1.** Let $\Omega \subset \mathbb{C}$ be a planar domain and $f: \Omega \to \mathbb{C}$ a continuous, open and light map. Let U be a normal domain in Ω . Then for any path $\beta: [0,1] \to fU$ and any point $x_0 \in U \cap f^{-1}\{\beta(0)\}$ there exists a path $\alpha: [0,1] \to U$ satisfying $\alpha(0) = x_0$ and $f \circ \alpha = \beta$.

We formulate a few lemmas for the proof of Theorem 3.1. For the statement of these auxiliary results we set $\mathscr{C}(\Omega)$ to be the collection of all nonempty compact subsets of $\Omega \subset \mathbb{C}$. This space is given the topology induced by the *Hausdorff distance* d_H of compact sets, that is, the distance $d_H(A, A')$ of compact sets $A, A' \subset \mathbb{C}$ is

$$d_H(A, A') = \inf \left\{ \varepsilon > 0 \mid A \subset B(A', \varepsilon), A' \subset B(A, \varepsilon) \right\}.$$

We show first that a continuous map $f: \Omega \to \mathbb{C}$ induces a continuous map $f_*: \mathscr{A} \to \mathscr{A}$. For the properties of the Hausdorff distance and convergence, we refer to [5, pp. 70–77] and merely comment on the continuity of the induced map f_* .

Lemma 3.2. For a continuous mapping $f: \Omega \to \mathbb{C}$ the induced map

 $f_* \colon \mathscr{C}(\Omega) \to \mathscr{C}(\mathbb{C}), \quad A \mapsto fA,$

is continuous in the topology induced by the Hausdorff distance.

Proof. Let $A \subset \Omega$ be a compact set, denote $\operatorname{dist}(A, \partial \Omega) =: s_0$ and set $\varepsilon > 0$. Since Ω is locally compact, the continuous map f is locally uniformly continuous and so there exists $\delta \in (0, \frac{s_0}{4})$ such that $d(f(x), f(y)) < \varepsilon$ for all $x, y \in B(A, \frac{s_0}{2})$ with $d(x, y) < \delta$. In particular, $fB(A, \delta) \subset B(fA, \varepsilon)$.

Let $A' \subset \Omega$ be a compact set for which $d_H(A, A') < \delta$. We show that $d_H(fA, fA') < \varepsilon$. Note first that since $fB(A, \delta) \subset B(fA, \varepsilon)$ and $A' \subset B(A, \delta)$, we have $fA' \subset B(fA, \varepsilon)$. On the other hand, for any $y_0 \in fA$ we can fix $x_0 \in A$ such that $f(x_0) = y_0$. Since $d_H(A, A') < \delta$ there now exists a point $z_0 \in A'$ with $d(z_0, x_0) < \delta$, and so by the aforementioned uniform continuity $d(f(z_0), y_0) < \varepsilon$, so in particular $fA \subset B(fA', \varepsilon)$. Thus $d_H(fA, fA') < \varepsilon$ and the induced map f_* is continuous.

The following lemma is contained e.g. in [19, pp. 131, 148], but we include a modern proof for the reader's convenience.

Lemma 3.3. Let $\Omega \subset \mathbb{C}$ be a planar domain and $f: \Omega \to \mathbb{C}$ a continuous, open and light map. Suppose U is a normal domain in Ω . Then for any $\varepsilon > 0$ there exists a constant $\delta > 0$ having the property that, for any continuum $C' \subset fU$ satisfying diam $(C') < \delta$, the components of $U \cap f^{-1}C'$ have diameter strictly less than ε .

Proof. Suppose there exists $\varepsilon_0 > 0$ having the property that, for each $n \in \mathbb{N}$, there exists a continuum $C'_n \subset fU$ having diameter at most 1/n and a component C_n of $U \cap f^{-1}C'_n$ having diameter at least ε_0 . Note that by Lemma 2.4 $f(C_n) = C'_n$.

Since both \overline{U} and \overline{fU} are compact, we may, after passing to subsequences if necessary, assume that the sequences (C'_n) and (C_n) converge in the Hausdorff metric to a point $z \in \overline{fU}$ and to a continuum $C \subset \overline{U}$ with diam $(C) \geq \varepsilon_0$, respectively. Then, by the continuity of f_* , i.e. Lemma 3.2, $fC = \{z\}$. Since f is light, this is a contradiction. Thus the claim is true.



FIGURE 1. An example of an $(f, \beta, 2)$ -prelift of a path.

With these preliminary results we can now turn to the proof of Theorem 3.1. To improve the clarity of the exposition we define some further auxiliary concepts. The crucial idea from Floyd is, in the setting of Theorem 3.1, to define for any subdivison $0 = t_0 < \ldots < t_k = 1$ of the unit interval a continuum $C \subset U$ such that $fC = |\beta|$ and $C \cap f^{-1}(\beta([t_j, t_{j+1}]))$ is a component of $f^{-1}(\beta([t_j, t_{j+1}]))$ for each $j = 0, \ldots, k - 1$.

For the purpose of lifting paths, the dyadic subdivisions turn out to be natural. We formalize Floyd's idea by introducing the concept of a prelift.

Definition 3.4. For a continuous, open and light mapping $f: \Omega \to \mathbb{C}$, a normal domain U of f and a path $\beta: [a,b] \to fU$ we call a continuum $C = C(f,\beta,n) \subset U$ an (f,β,n) -prelift of β in U if

- (1) for each $k = 1, ..., 2^n$, $C \cap f^{-1}(\beta[(k-1)2^{-n}, k2^{-n}])$ is a component of $f^{-1}(\beta[(k-1)2^{-n}, k2^{-n}])$, and
- (2) for each $k = 0, \dots 2^n, C \cap f^{-1}(\beta(k2^{-n})) \neq \emptyset$.

We say that a prelift C starts from $x_0 \in f^{-1}(\beta(a))$ if $x_0 \in C$.

The restriction of a prelift $C(f,\beta,n)$ to an interval $[c,d] \subset [a,b]$ is just a subcontinuum of C that is also an $(f,\beta|_{[c,d]},n)$ -prelift. Note that an (f,β,n) prelift is also a (f,β,m) -prelift for all $m \leq n$.

Finally we denote by w(C) the mesh of a prelift $C = C(f, \beta, n)$, that is

$$w(C) := \max_{k=1,\dots,2^n} \operatorname{diam} \left(C \cap f^{-1}(\beta[(k-1)2^{-n}, k2^{-n}]) \right).$$

Note that as an immediate corollary of Lemma 3.3 and the uniform continuity of β we see that $w(C(f, \beta, n)) \to 0$ as $n \to \infty$.

A posteriori, for planar continuous, open and light mappings the definition of a prelift turns out to be overly verbose; the prelifts for continuous, open and discrete mappings are merely boundedly finite unions of path segments by Rickman's path-lifting theorem in [14].

Lemma 3.5. In the setting of Theorem 3.1, for any $n \in \mathbb{N}$ and any given $x_0 \in U \cap f^{-1}\{\beta(0)\}$, there exists an (f, β, n) -prelift of β in U starting from x_0 .

Proof. Fix $n \in \mathbb{N}$ and $x_0 \in U \cap f^{-1}\{\beta(a)\}$. We set C_0 to be the component of $U \cap f^{-1}(\beta[0, 2^{-n}])$ containing x_0 . By Lemma 2.4 this component C_0 maps surjectively onto $\beta([0, 2^{-n}])$.

Suppose C_{k-1} has been defined for some $k < 2^n$. By Lemma 2.4 C_{k-1} is mapped surjectively onto the set $\beta[(k-1)2^{-n}, k2^{-n}]$, which intersects $\beta[k2^{-n}, (k+1)2^{-n}]$ by the continuity of β . In particular C_{k-1} contains a pre-image of the point $f^{-1}{\beta(k2^{-n})}$, and so we may fix a component C_k of $f^{-1}(\beta[k2^{-n}, (k+1)2^{-n}])$ such that C_k and C_{k-1} intersect. Again by Lemma 2.4 this component C_k maps surjectively onto $\beta([k2^{-n}, (k+1)2^{-n}])$.

We now set $C := \bigcup_{j=0}^{2^n-1} C_j$ and note immediately that C is the required (f, β, n) -prelift of β .

We are now ready to prove Theorem 3.1. The idea of the proof relies on taking a sequence of prelifts given by Lemma 3.5 and then constructing a subsequence of these prelifts that is, in a sense, converging on each subinterval of [0, 1]. Curious readers might be interested in comparing this to the properties of ultralimits where one can, in a compact metric space, choose a converging subsequence for each sequence in a consistent manner via a principal ultrafilter in \mathbb{N} ; see e.g. [5, pp. 77-80].

Proof of Theorem 3.1. Recall that by Lemma 2.3 the restriction of a continuous, open and light mapping to a normal domain is a proper map.

Fix a (f, β, n) -prelift C_n for each $n \in \mathbb{N}$. We recursively construct a nested sequence of subsequences of (C_n) . First note that since (C_n) is a sequence of continua in a precompact domain U, it has a subsequence $(C_j^0)_j$ converging to a continuum $C \subset \overline{U}$ with respect to the Hausdorff metric. Furthermore since $f(C_n) \subset |\beta|$ with $|\beta| \cap \partial fU = \emptyset$, we have $C \subset U$.

Suppose now that the subsequence $(C_m^{j-1})_m$ has already been defined. We define the next subsequence $(C_m^j)_m$ in 2^j steps. First note that a restriction of the prelifts C_m^{j-1} to the interval $[0, 2^{-j}]$ gives rise to a sequence of $(f, \beta|_{[0,2^{-j}]}, j)$ -prelifts. As before we may take a subsequence (S_n^0) of $(C_m^j)_m$ such that restricted sequence converges. Now suppose that a subsequence $(S_n^k)_n$ of $(C_m^j)_m$ has been defined for $k < 2^j - 1$. We set (S_n^{k+1}) to be the subsequence of (S_n^k) such that the restrictions of these prelifts S_n^k to the interval $[k2^{-j}, (k+1)2^{-j}]$ form a converging sequence. Finally we set $(C_m^j)_m = (S_m^{2j})_m$.

From these nested sequences $(C_n^j)_n$, $j \in \mathbb{N}$, we take the diagonal, i.e. we set $P_n = C_n^n$, and note that the sequence (P_n) has the property that for any $j \in \mathbb{N}$ it is the subsequence of $(C_n^j)_n$ after possibly omitting finitely many elements. In particular, for any dyadic interval $\sigma \subset [0, 1]$, the restrictions of P_n to the interval σ form a converging sequence in U with respect to the Hausdorff metric. We show next that the limit of P_n , denoted P, is a path and thus a lift of β under f.

Fix $t_0 \in [0, 1]$. Note that for any $m \in \mathbb{N}$ there are at most two intervals of the form $[k2^{-m}, (k+1)2^{-m}]$ containing t_0 . Call the union of these two intervals σ^m . Since the restrictions of P_n to any of the intervals σ^m converge as $n \to \infty$, we thus see that with respect to the Hausdorff metric,

$$\left(P_n \cap f^{-1}(\beta(\sigma^n))\right) = \bigcap_{k=1}^n \left(P_n \cap f^{-1}(\beta(\sigma^k))\right) \to P \cap f^{-1}(\beta(t_0)).$$

Furthermore, since the mesh of the prelifts P_n tends to zero as $n \to \infty$, we note that also

diam
$$(P_n \cap f^{-1}(\beta(\sigma^n))) \to 0$$
 as $n \to \infty$.

Thus $P \cap f^{-1}(\beta(t_0))$ is a point; call it $\alpha(t_0)$. We next note that since the mesh of the prelifts P_n tends to zero, there exists for any $\epsilon > 0$ an integer $m_{\epsilon} \in \mathbb{N}$ such that $d(\alpha(t_0), \alpha(t)) < \epsilon$ whenever $|t_0 - t| < 2^{-m_{\epsilon}}$. Thus the mapping $t \mapsto \alpha(t)$ is continuous.

The path $\alpha: [0,1] \to U$ is now a path that maps onto β under f, and $\alpha(0) = x_0$ by construction of the sequence (C_n) . Furthermore, since the prelifts were defined via pre-image components of pieces of $|\beta|$, we see that in fact $f \circ \alpha = \beta$ by Lemma 3.2. Thus α is a lift of β in U, and the claim holds true.

We end this section with a proposition on the uniqueness of lifts into simply connected planar domains. Note that this claim clearly fails for maps between more general surfaces, even between spheres, and in higher dimensions.

Proposition 3.6. Let $\Omega \subset \mathbb{C}$ be a simply connected planar domain, $f: \Omega \to \mathbb{C}$ a continuous open map, and let $\beta_1, \beta_2: [0,1] \to \Omega$ be lifts of the same line segment $\alpha: [0,1] \to f\Omega$ for which $\beta_1(0) = \beta_2(0)$ and $\beta_1(1) = \beta_2(1)$. Then $\beta_1 = \beta_2$.

Proposition 3.6 is an almost immediate consequence of the following version of the Jordan curve theorem; for a proof we refer to [12, Section 4] or [13, Theorem 1.10, p.33]. Furthermore we emphasize here that by the boundary of a domain U compactly contained in a planar domain Ω we mean the boundary relative to Ω .

Theorem (The Jordan curve theorem). Let $\Omega \subset \mathbb{C}$ be a simply connected planar domain and let $c: \mathbb{S}^1 \to \Omega$ be an injective continuous map. Then $\Omega \setminus |c|$ consists of two domains, exactly one of which has a compact closure in Ω . Both of these domains have the image of the curve c as their boundary.

Besides the Jordan curve theorem we need also the following result stating that a line segment is not a boundary of a bounded domain in the plane. We formulate the statement of the lemma in slightly more general form for later use.

Lemma 3.7. Let Ω be a precompact planar domain and $\beta : [0,1] \to \Omega$ a line segment. Suppose r > 0 is a radius such that the closed balls $\overline{B}(\beta(0), r)$ and $\overline{B}(\beta(1), r)$ are disjoint and contained in Ω . Then there exists no precompact domain $W \subset \Omega$ such that

$$\partial W \subset |\beta| \cup \overline{B}(\beta(0), r) \cup \overline{B}(\beta(1), r)$$

and

$$(|\beta| \setminus (\overline{B}(\beta(0), r) \cup \overline{B}(\beta(1), r))) \cap \partial W \neq \emptyset.$$

Proof. Towards contradiction, suppose there exists such a precompact domain W. Let

$$z \in \left(|\beta| \setminus (\overline{B}(\beta(0), r) \cup \overline{B}(\beta(1), r)) \right) \cap \partial W$$

and take a radius s > 0 such that $\overline{B}(w, s)$ intersects neither of the closed balls $\overline{B}(\beta(0), r)$ and $\overline{B}(\beta(1), r)$.

Now the line segment β divides B(w, s) into two domains, one of which must be contained in W since $B(w, s) \cap \partial W$ is contained in the line segment $|\beta|$. Denote that domain by V. Let now $\alpha \colon [0, 1) \to \Omega$ be a line segment, perpendicular to β , that starts from w, intersects V and ends at the boundary of Ω . Since α does not intersect either of the balls $\overline{B}(\beta(0), r)$ and $\overline{B}(\beta(1), r)$, we conclude that it intersects the boundary of W at most once at $\alpha(0)$. Thus $\alpha(0, 1) \subset W$ and so W is not precompact as $\alpha(1) \notin W$. \Box

Proof of Proposition 3.6. Suppose $\beta_1 \neq \beta_2$. Then there exists $t_0 \in [0, 1]$ for which $\beta_1(t_0) \neq \beta_2(t_0)$. Let

$$a = \sup\{t \in [0, t_0] \mid \beta_1(t) = \beta_2(t)\} \text{ and } b = \inf\{t \in [t_0, 1] \mid \beta_1(t) = \beta_2(t)\}.$$

Then there exists an injective continuous map $c \colon \mathbb{S}^1 \to \Omega$ for which $|c| = |\beta_1|_{[a,b]}| \cup |\beta_2|_{[a,b]}|$, and so $|f \circ c| = \alpha[a,b]$. By the Jordan curve theorem, one of the components of $\Omega \setminus |c|$, say U, is a precompact subdomain of Ω with $\partial U = |c|$.

Thus fU is a precompact domain in \mathbb{C} and $\partial fU \subset |\alpha|$. This is a contradiction with Lemma 3.7 since α is a line segment. The claim follows. \Box

4. Proof of Theorem 1.1

We use now the path-lifting result, Theorem 3.1, and the Jordan curve theorem to prove Theorem 1.1, that is, to show that a continuous, open and light planar map is discrete. We follow here an idea of Stoïlow that the discreteness of the mapping follows from the finiteness of lifts of line segments. Stoïlow calls the following proposition, together with the existence of lifts, *Théorème Fondamental* ([16, p. 361]).

Proposition 4.1. Let Ω be a planar domain and $f: \Omega \to \mathbb{C}$ a continuous, open and light map. Let $x \in \Omega$ and let r > 0 be so small that U := U(x, f, r)is a normal domain contained in a simply connected neighborhood of x in Ω . Suppose $\beta: [0,1] \to B(f(x),r)$ is a ray $t \mapsto tz_0 + f(x)$, where $z_0 \in \mathbb{S}^1$. Then there exists at most finitely many lifts of β in U.

With this proposition at our disposal, Theorem 1.1 follows immediately.

Proof of Theorem 1.1 assuming Proposition 4.1. Let $x_0 \in \Omega$ be a point and fix a normal domain U_0 of x_0 contained in some simply connected neighborhood of x in Ω . By Theorem 3.1 a ray in fU_0 starting from $f(x_0)$ has a lift starting from each pre-image of $f(x_0)$ in U_0 . By Proposition 4.1, such a ray has only finitely many lifts in U_0 . Thus the set $U_0 \cap f^{-1}{f(x_0)}$ is finite and so f is a discrete map. \Box

For the proof of Proposition 4.1 we require a few auxiliary results. The first one is the following technical lemma about the intersection properties of a sequence of sets which will be applied to the images of a sequence of paths.

STOÏLOW'S THEOREM REVISITED



FIGURE 2. Constructing a "rectangular Jordan curve" in the proof of Proposition 4.1.

The proof requires no geometric properties of paths or planar topology; this result is actually a version of the infinite Ramsey theorem for two colors; see e.g. [8, Theorem 5, p. 16].

Lemma 4.2. Let X be a set and (A_n) be a sequence of subsets of X. Then there exists a subsequence (C_n) of (A_n) for which either

(a) all the sets C_n are mutually disjoint, or

(b) any two of the sets C_n intersect.

Proof. We construct fist a subsequence (B_n) of (A_n) with the property that for any $k \in \mathbb{N}$ either

- (i) $B_j \cap B_k = \emptyset$ for all j > k, or
- (ii) $B_j \cap B_k \neq \emptyset$ for all j > k.

We construct the sequence (B_n) recursively together with a sequence of nested auxiliary subsequences $(B_n^k)_n$ of (A_n) . Set first $B_0 = A_0$. Now either B_0 intersects infinitely many elements of $(A_n)_{n\geq 1}$ or is disjoint from infinitely many elements of $(A_n)_{n\geq 1}$. Choose one such infinite collection and call it (B_n^0) .

Suppose that (B_0, \ldots, B_k) and $((B_n^0), \ldots, (B_n^k))$ have been defined. We set $B_{k+1} = B_0^k$ and note that either B_{k+1} intersects infinitely many elements of $(B_n^k)_{n\geq 1}$ or is disjoint from infinitely many elements of $(B_n^{k+1})_{n\geq 1}$. Choose one such infinite collection and call it (B_n^{k+1}) .

Since for all $k \in \mathbb{N}$ the sequence $(B_n^k)_n$ is a subsequence of $(B_n^{k-1})_n$, we note that the required property holds for (B_n) , i.e. for any $k \in \mathbb{N}$ either (i) or (ii) holds.

To complete the proof we note that there are two types of elements in the sequence (B_n) ; those satisfying condition (i) and those satisfying condition (ii). Thus there must be infinitely many of elements of either type (i) or type (ii). Call one such collection (C_n) . Now either all elements of (C_n) intersect all of the consequtive elements of the sequence or all elements of (C_n) intersect none of the consequtive elements of the sequence. Thus we

see that (C_n) satisfies the condition in the statement of the lemma and so the proof is complete.

Finally, we require the following lemma that allows us to extend pairs of disjoint arcs into a Jordan curve.

Lemma 4.3. Let Ω be a planar domain with $A, B \subset \Omega$ domains such that $\overline{A} \cap \overline{B} = \emptyset$ and $\overline{A}, \overline{B} \subset \Omega$. Suppose there exists two arcs $\alpha, \beta \colon [0,1] \to \Omega$ for which $\alpha(0), \beta(0) \in A, \alpha(1), \beta(1) \in B$ and $|\alpha| \cap |\beta| = \emptyset$. Then there exists a Jordan curve $c \colon \mathbb{S}^1 \to \Omega$ for which $|\alpha| \setminus (A \cup B) \subset |c|, |\beta| \setminus (A \cup B) \subset |c|$ and $|c| \subset A \cup B \cup |\alpha| \cup |\beta|$.

Proof. Planar domains are arc-connected, so there exists arcs $\gamma_A \colon [0,1] \to A$ and $\gamma_B \colon [0,1] \to B$ connecting the pairs of points $(\alpha(0), \beta(0))$ and $(\alpha(1), \beta(1))$, respectively. Fix

$$t_A^1 := \sup\{t \in [0,1] \mid \gamma_A(t) \in |\alpha|\}, \quad t_A^2 := \inf\{t \in [t_A^1, 1] \mid \gamma_A(t) \in |\beta|\}, \\ t_B^1 := \sup\{t \in [0,1] \mid \gamma_B(t) \in |\alpha|\}, \text{ and } t_B^2 := \inf\{t \in [t_B^1, 1] \mid \gamma_B(t) \in |\beta|\}.$$

We note that the restrictions $\gamma_A|_{[t_A^1, t_A^2]}$ and $\gamma_B|_{[t_B^1, t_B^2]}$ are arcs that meet the paths α and β only at their endpoints. Thus we may define the Jordan curve c as a concatenation of these two restrictions together with the restrictions of α and β to suitable subintervals.

Proof of Proposition 4.1. Suppose a ray $\beta \colon [0,1] \to fU$ has infinitely many mutually distinct lifts $\tilde{\beta}_n \colon [0,1] \to U$, $n \in \mathbb{N}$. By passing to a subsequence if necessary, we may assume that $\tilde{\beta}_n(0) \to x_0 \in \overline{U}$ and $\tilde{\beta}_n(1) \to y_0 \in \overline{U}$ as $n \to \infty$. Since U is a normal domain and $\beta(0), \beta(1) \in B(f(x), r)$, we have, in fact, that $x_0, y_0 \in U$.

By Lemma 4.2 we may assume, by passing again to a subsequence if necessary, that for the sequence $(\tilde{\beta}_n)$ either

(a) the images of the lifts β_n are mutually disjoint, or

(b) any two of the images of the lifts $\tilde{\beta}_n$ intersect.

Suppose first that (a) holds. Let s > 0 be a radius for which $V = U(x_0, f, s)$ and $V' = U(y_0, f, s)$ are mutually disjoint normal domains of x_0 and y_0 , respectively. Let $M, N \in \mathbb{N}$ be distinct indices for which

$$\beta_M(0), \beta_N(0) \in V$$
 and $\beta_M(1), \beta_N(1) \in V'$.

Then there exists by Lemma 4.3 a Jordan curve $c \colon \mathbb{S}^1 \to U$ for which

$$|c| \subset V \cup V' \cup |\tilde{\beta}_M| \cup |\tilde{\beta}_N|$$
 and $|c| \not\subset V \cup V'$.

The image $|f \circ c|$ is a continuum contained in the ray $|\beta|$ and two mutually disjoint disks $B(f(x_0), s)$ and $B(f(y_0), s)$ located at the endpoints of β ; see Figure 2. Let $K = |\beta| \cup (\bar{B}(f(x_0), s) \cup \bar{B}(f(y_0), s))$.

By the Jordan curve theorem the curve c bounds a precompact domain W in U, having boundary equal to |c|. Now, on the one hand, the boundary $\partial f W$ of f W is contained in $|c| \subset K$. Also note that since U = U(x, f, r), we in particular have that fU = B(f(x), r) and so fW is a precompact domain in a disk. On the other hand, the closure of fW contains the segment $|\beta| \setminus (B(f(x_0), s) \cup B(f(y_0), s))$. Thus, by connectedness of fW, we have $fW \supset B(f(x), r) \setminus K$. In particular, fW is not precompact in B(f(x), r).



FIGURE 3. Path having infinitely many lifts with a joint starting point in the proof of Proposition 4.1.

By Lemma 3.7 there can be no such domain fW. This is a contradiction, and the proof in case (a) is complete.

Suppose now that (b) holds. By Proposition 3.6 one of the sets

 $\{\tilde{\beta}_j(1) \mid j \in \mathbb{N}\}$ or $\{\tilde{\beta}_j(0) \mid j \in \mathbb{N}\}$

must be infinite. Thus we may assume, by changing the direction of the path β and by passing to a subsequence if necessary, that the endpoints $\tilde{\beta}_i(1)$ are all distinct.

Let $\gamma: [0,1] \to B(f(y_0),r)$ be a ray for which $\gamma(0) = f(y_0)$ and $|\gamma| \cap |\beta| = f(y_0)$; see Figure 3. For each $n \in \mathbb{N}$, let $\tilde{\gamma}_n$ be a lift of γ starting from $\tilde{\beta}_n(1)$. We show next that all the lifts $\tilde{\gamma}_n$ are pair-wise disjoint. Towards contradiction suppose not and take $\tilde{\gamma}_k$ and $\tilde{\gamma}_j$ such that $\tilde{\gamma}_k(s_0) = \tilde{\gamma}_j(s_0)$ for some $s_0 \in [0,1]$. Now by the property (b) the two lifts $\tilde{\beta}_k$ and $\tilde{\beta}_j$ with corresponding indices also intersect; $\tilde{\beta}_k(t_0) = \tilde{\beta}_j(t_0)$ for some $t_0 \in [0,1]$. We may assume t_0 is the first such intersection point, i.e. either $t_0 = 0$ or $\tilde{\beta}_k(t) \neq \tilde{\beta}_j(t)$ for all $t < t_0$. For i = k, j we define the concatenations $\theta_i: [0,1] \to U$ by setting

$$\theta_i(t) = \begin{cases} \tilde{\beta}_i(2t), & t \in [0, 1/2] \\ \tilde{\gamma}_i(2t-1), & t \in [1/2, 1]. \end{cases}$$

Now applying Proposition 3.6 to the restrictions

 $\theta_k|_{[t_0/2,(s_0+1)/2]}$ and $\theta_j|_{[t_0/2,(s_0+1)/2]}$

gives rise to a contradiction and we conclude that all the lifts $\tilde{\gamma}_n$ are pair-wise disjoint.

Now the argument of case (a) applies to the sequence $(\tilde{\gamma}_n)$, which is again a contradiction and the proof is complete.

5. Proofs of Theorems 1.2 and 1.3

Theorem 1.2 is a direct consequence of the following proposition. For the statement, we say that a normal domain U of a continuous, open and discrete map $f: \Omega \to \mathbb{C}$ is a normal neighborhood of x (with respect to f), if $\overline{U} \cap f^{-1}{f(x)} = {x}$. The existence of such normal neighborhoods follows immediately by noting that on the one hand for the discrete map f and any point $x \in \Omega$, we have $d(x, f^{-1}{f(x)} \setminus {x}) > 0$ and on the other hand that for continuous, open and discrete mappings diam $(U(x, f, r)) \to 0$ as $r \to 0$. The latter property in turn follows e.g. from Lemma 3.3, but see also [15, Lemma I.4.9].

Proposition 5.1. Let Ω be a planar domain and let $f: \Omega \to \mathbb{C}$ be a continuous, open and discrete map. Let $x_0 \in \Omega$ and let r > 0 be so small that $U_0 := U(x_0, f, r)$ is a normal neighborhood of x_0 contained in a simply connected domain in Ω . Then $U_0 \cap B_f \subset \{x_0\}$.

Note that since $f: \Omega \to \mathbb{C}$ is a priori both continuous and open, the branch set B_f of f is actually the set of points at which f is not locally injective. Indeed, since a continuous and open bijection is a homeomorphism, we conclude that a continuous and open locally injective map is a local homeomorphism. In particular, if f is locally injective at $x \in \Omega$, then x has a neighborhood $U \subset \Omega$ for which $f|_U: U \to fU$ is a homeomorphism and fU is an open neighborhood of f(x). Thus $x \notin B_f$.

Proof of Proposition 5.1. Suppose there exists $b \in (U_0 \cap B_f) \setminus \{x_0\}$ and let $U \subset U_0$ be a normal neighborhood of b. Since $b \in B_f$, the map f is not locally injective at b. Thus we may fix a point $y_0 \in fU$ for which $\#(U \cap f^{-1}\{y_0\}) \ge 2$. Note that since U is a normal neighborhood of b, $U \cap f^{-1}\{f(b)\} = \{b\}$ and so $y_0 \ne f(b)$; likewise $y_0 \ne f(x_0)$.

Note that fU is a planar disk. Thus we may fix two (piecewise linear) arcs $\alpha: [0,1] \to fU$ and $\beta: [0,1] \to fU_0 \setminus \alpha(0,1]$ satisfying $\alpha(0) = \beta(0) = y_0$, $\alpha(1) = f(b)$, and $\beta(1) = f(x_0)$. Let also $z_1, z_2 \in U \cap f^{-1}\{y_0\}, z_1 \neq z_2$. By Theorem 3.1, there exists, for i = 1, 2, lifts $\tilde{\alpha}_i: [0,1] \to U$ and $\tilde{\beta}_i: [0,1] \to U_0$ of α and β , respectively, satisfying $\tilde{\alpha}_i(0) = \tilde{\beta}_i(0) = z_i$ for i = 1, 2. Since U_0 and U are normal neighborhoods of x_0 and b, respectively, we have $\tilde{\alpha}_i(1) = b$, $\tilde{\beta}_i(1) = x_0$ for i = 1, 2.

For i = 1, 2, let $\tilde{\gamma}_i \colon [0, 1] \to U$ be the path

$$t \mapsto \begin{cases} \tilde{\alpha}_i(2t), & t \in [0, 1/2] \\ \tilde{\beta}_i(2t-1), & t \in [1/2, 1]. \end{cases}$$

Since $\tilde{\gamma}_1(0) = \tilde{\gamma}_2(0)$, $\tilde{\gamma}_1(1) = \tilde{\gamma}_2(1)$, and U is contained in a simply connected domain, we have by Proposition 3.6 that $\tilde{\gamma}_1 = \tilde{\gamma}_2$. This is a contradiction and the claim follows.

We finish with a simple proof of Theorem 1.3 based on Theorem 1.2 and a covering argument. For the basic properties and terminology concerning covering maps and covering neighborhoods we refer to [9].

Proof of Theorem 1.3. Let $z \in \Sigma$. Since the theorem posits the existence of some neighborhood, the claim is local in nature and thus we may assume that Σ and Σ' are planar domains.

Let r > 0 be so small that U(z, f, r) is a normal neighborhood of z contained in some simply connected domain in Σ . Denote

$$U' = U(z, f, r) \setminus \{z\}, B' = B(f(z), r) \setminus \{f(z)\} \text{ and } \mathbb{D}' = \mathbb{D} \setminus \{0\}.$$

Since f is discrete by Theorem 1.1,

$$U' \cap f^{-1}\{y\} \subset \overline{U} \cap f^{-1}\{y\}$$

is finite for any $y \in fU$. Furthermore, by Proposition 5.1 $f|_{U'}: U' \to B'$ is a local homeomorphism and since U is a normal neighborhood, $f|_{U'}$ is also a proper map. Thus as a proper local homeomorphism $f|_{U'}$ is a covering map. Indeed, for any point $y_0 \in B'$ the pre-image $U' \cap f^{-1}\{y_0\}$ is a finite set $\{x_1, \ldots, x_k\}$ and we may fix disjoint open sets U_j such that the restriction $f|_{U_j}: U_j \to fU_j, j = 1, \ldots, k$, is a homeomorphism. The open set $\bigcap_{j=1}^k f(U_j)$ is now a covering neighborhood of y_0 , i.e. the restriction of f to its pre-image components is a homeomorphism onto the set.

As a finite cover of B', the domain U' is a topological punctured disk and we conclude that U is a topological disk. Let $h_1: U \to \mathbb{D}$ and $h_2: B(f(z), r) \to \mathbb{D}$ be homeomorphisms with $h_1(z) = 0$ and $h_2(f(z)) = 0$. Let also

$$g := h_2 \circ f \circ (h_1|_{U'})^{-1} \colon \mathbb{D}' \to \mathbb{D}'.$$

Since $f|_{U'}: U' \to B'$ is a covering map, so is g. Thus the induced map $g_*: \pi_1(\mathbb{D}') \to \pi_1(\mathbb{D}')$ is of the form $m \mapsto km$ for some $k \in \mathbb{Z} \setminus \{0\}$. Note that $k \neq 0$, since the induced homomorphism g_* is injective by the homotopy lifting property of covering maps.

We consider first the case k > 0. Let $\zeta_k : \mathbb{D}' \to \mathbb{D}'$ be the covering map $z \mapsto z^k$, and let $h' : \mathbb{D}' \to \mathbb{D}'$ be the lift of $g : \mathbb{D}' \to \mathbb{D}'$ under ζ_k . Then $g = \zeta_k \circ h'$ and h' is a homeomorphism, since it is an injective covering map. The homeomorphism h' extends to a homeomorphism $h : \mathbb{D} \to \mathbb{D}$ by the continuity of ζ_k and hence $f|_U = h_2 \circ \zeta_k \circ h \circ h_1 =: \phi^{-1} \circ \zeta_k \circ \psi$.

In the case k < 0 we repeat the previous argument by replacing the homeomorphism h_2 by the homeomorphism $k_2: B(f(x), r) \to \mathbb{D}$ defined by $x \mapsto \overline{h(z)}$.

References

- L. V. Ahlfors and L. Sario. *Riemann surfaces*. Princeton Mathematical Series, No. 26. Princeton University Press, Princeton, N.J., 1960.
- [2] K. Astala, T. Iwaniec, and G. Martin. Elliptic partial differential equations and quasiconformal mappings in the plane, volume 48 of Princeton Mathematical Series. Princeton University Press, Princeton, NJ, 2009.
- [3] L. Bers. Riemann surfaces: Lectures by Lipman Bers. New York University, 1958.
- [4] M. Bonk and D. Meyer. Expanding Thurston Maps. Mathematical Surveys and Monographs. American Mathematical Society, 2017.
- [5] M. R. Bridson and A. Haefliger. Metric spaces of non-positive curvature, volume 319 of Grundlehren der Mathematischen Wissenschaften. Springer-Verlag, Berlin, 1999.
- [6] R. Engelking. General topology, volume 6 of Sigma Series in Pure Mathematics. Heldermann Verlag, Berlin, second edition, 1989.
- [7] E. E. Floyd. Some characterizations of interior maps. Ann. of Math. (2), 51:571–575, 1950.
- [8] R. L. Graham, B. L. Rothschild, and J. H. Spencer. *Ramsey theory*. Wiley-Interscience Series in Discrete Mathematics and Optimization. John Wiley & Sons, Inc., New York, second edition, 1990.
- [9] A. Hatcher. Algebraic topology. Cambridge University Press, Cambridge, 2002.
- [10] W. Hurewicz and H. Wallman. Dimension Theory. Princeton Mathematical Series, volume 4. Princeton University Press, Princeton, N. J., 1941.
- [11] R. Luisto. A characterization theorem for BLD-mappings between metric spaces. The Journal of Geometric Analysis, 27:2081–2097, 2017.

- [12] E. E. Moise. Geometric topology in dimensions 2 and 3, volume 47 of Graduate Texts in Mathematics. Springer-Verlag, New York-Heidelberg, 1977.
- [13] C. Pommerenke. Univalent functions. Vandenhoeck & Ruprecht, Göttingen, 1975. Studia Mathematica/Mathematische Lehrbücher, Band XXV.
- [14] S. Rickman. Path lifting for discrete open mappings. Duke Math. J., 40:187–191, 1973.
- [15] S. Rickman. Quasiregular mappings, volume 26 of Ergebnisse der Mathematik und ihrer Grenzgebiete (3). Springer-Verlag, Berlin, 1993.
- [16] S. Stoïlow. Sur les transformations continues et la topologie des fonctions analytiques. Ann. Sci. Éc. Norm. Supér. (3), 45:347–382, 1928.
- [17] S. Stoïlow. Leçons sur les principes topologiques de la théorie des fonctions analytiques. Professées à la Sorbonne et à l'université de Cernauti. X + 148 p. Paris, Gauthier-Villars (Collection de monographies sur la théorie des fonctions), 1938.
- [18] J. Väisälä. Discrete open mappings on manifolds. Ann. Acad. Sci. Fenn. Ser. A I No., 392:10, 1966.
- [19] G. T. Whyburn. Analytic topology. American Mathematical Society Colloquium Publications, Vol. XXVIII. American Mathematical Society, Providence, R.I., 1942.
- [20] G. T. Whyburn. *Topological analysis*. Second, revised edition. Princeton Mathematical Series, No. 23. Princeton University Press, Princeton, N.J., 1964.

DEPARTMENT OF MATHEMATICS AND STATISTICS, P.O. BOX 35, FI-40014 UNIVER-SITY OF JYVÄSKYLÄ, FINLAND AND DEPARTMENT OF MATHEMATICAL ANALYSIS, SOKOLOVSKA 83, PRAHA 8, 186 75, CHARLES UNIVERSITY IN PRAGUE

E-mail address: rami.luisto@gmail.com

DEPARTMENT OF MATHEMATICS AND STATISTICS, P.O. BOX 68 (GUSTAF HÄLLSTRÖMIN KATU 2B), FI-00014 UNIVERSITY OF HELSINKI, FINLAND

E-mail address: pekka.pankka@helsinki.fi