

This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Turunen, Katri Maria; Kokko, Katja; Kekäläinen, Tiia; Alén, Markku; Hänninen, Tuomo; Pynnönen, Katja; Laukkanen, Pia; Tirkkonen, Anna; Törmäkangas, Timo; Sipilä, Sarianna

Title: Associations of neuroticism with falls in older adults : do psychological factors mediate the association?

Year: 2022

Version: Accepted version (Final draft)

Copyright: © 2020 Taylor & Francis

Rights: In Copyright

Rights url: <http://rightsstatements.org/page/InC/1.0/?language=en>

Please cite the original version:

Turunen, K. M., Kokko, K., Kekäläinen, T., Alén, M., Hänninen, T., Pynnönen, K., Laukkanen, P., Tirkkonen, A., Törmäkangas, T., & Sipilä, S. (2022). Associations of neuroticism with falls in older adults : do psychological factors mediate the association?. *Aging and Mental Health*, 26(1), 77-85. <https://doi.org/10.1080/13607863.2020.1841735>

1 Associations of Neuroticism with Falls in Older Adults: Do Psychological Factors Mediate
2 the Association?

3 Authors:

4 Turunen Katri Maria (PhD), Gerontology Research Center and Faculty of Sport and Health
5 Sciences, University of Jyväskylä, Jyväskylä, Finland

6 Kokko Katja (PhD), Gerontology Research Center and Faculty of Sport and Health Sciences,
7 University of Jyväskylä, Jyväskylä, Finland

8 Kekäläinen Tiia (PhD), Gerontology Research Center and Faculty of Sport and Health
9 Sciences, University of Jyväskylä, Jyväskylä, Finland

10 Alen Markku (MD, PhD), Department of Medical Rehabilitation, Oulu University Hospital,
11 Oulu, Finland

12 Hänninen Tuomo (PhD), Department of Neurology, Kuopio University Hospital, Kuopio,
13 Finland

14 Pynnönen Katja (PhD), Gerontology Research Center and Faculty of Sport and Health
15 Sciences, University of Jyväskylä, Jyväskylä, Finland

16 Laukkanen Pia (MD, PhD), Gerontology Research Center and Faculty of Sport and Health
17 Sciences, University of Jyväskylä, Jyväskylä, Finland

18 Tirkkonen Anna (MSc), Gerontology Research Center and Faculty of Sport and Health
19 Sciences, University of Jyväskylä, Jyväskylä, Finland

20 Törmäkangas Timo (PhD), Gerontology Research Center and Faculty of Sport and Health
21 Sciences, University of Jyväskylä, Jyväskylä, Finland

- 22 Sipilä Sarianna (PhD), Gerontology Research Center and Faculty of Sport and Health
23 Sciences, University of Jyväskylä, Jyväskylä, Finland
- 24 Corresponding author: Katri Turunen, katri.m.turunen@jyu.fi

25 Abstract

26 Objectives: Neuroticism predicts falls in older people. In addition, concern about falling and
27 depressive symptoms are associated with fall risk. This study examined whether concern
28 about falling and depressive symptoms mediate the association between neuroticism and
29 falls.

30 Method: Cross-sectional data on 314 community-dwelling people aged 70-85 years were
31 utilized. Neuroticism was assessed with a short modified form of the Eysenck Personality
32 Inventory. Indoor and outdoor falls during the past year were self-reported. Concern about
33 falling was assessed with the Falls Efficacy Scale-International and depressive symptoms
34 with the Geriatric Depression Scale-15. Path modeling was used to examine the associations
35 between variables.

36 Results: Mediating pathways linking neuroticism and falls were found: neuroticism was
37 positively associated with concern about falling, which was subsequently linked to indoor
38 falls (indirect effect $\beta=0.34$, $P=0.002$) and recurrent outdoor falls ($\beta=0.19$, $P=0.045$).
39 Moreover, a pathway from neuroticism to indoor falls through depressive symptoms was also
40 found ($\beta=0.21$, $P=0.054$). In other words, higher neuroticism was associated with higher
41 concern about falling and depressive symptoms, both of which were linked to falls. The
42 associations were independent of age, sex, use of psychotropic, chronic diseases, persistent
43 pain, physical performance, physical activity, and executive functioning that are known risk
44 factors for falls.

45 Discussion: The results indicate that concern about falling and depressive symptoms mediate
46 the association between neuroticism and falling. Longitudinal studies are needed to confirm
47 the causality of the findings and to examine the potential to reduce falls by targeting concern
48 about falling and depressive symptoms among older adults higher in neuroticism.

49 KEYWORDS: Aged, Cross-sectional Studies, Accidental Falls, Risk Factors, Fear,
50 Personality.

51 Introduction

52 More than one-third of community-dwelling older adults fall each year, and approximately
53 ten percent of these falls result in serious injury such as a fracture. Falling continues to
54 present a challenge despite extensive knowledge of the intrinsic physiological risk factors,
55 such as impaired balance and vision, poor muscle strength, medication and extrinsic risk
56 factors such as environmental hazards. Some of the known risk factors contributing to falling
57 are modifiable (Deandrea et al., 2010). However, psychological factors such as personality
58 traits, depressive symptomatology and concern about falling have received less attention. A
59 better understanding of these psychological factors could be informative about the
60 mechanisms underlying falls and thus help in the development of more effective fall-
61 prevention interventions for older people at increased risk for falling.

62 Personality traits, defined as individuals' characteristic ways of behaving, thinking and
63 feeling, are relatively stable across both contexts and time (Caspi, 1998; McCrae & Costa,
64 2008). The personality traits included in the well-established Five-Factor Model (Digman,
65 1990) are neuroticism, extraversion, openness, conscientiousness, and agreeableness. The
66 present article focuses on neuroticism, as this trait is recognized as an important contributory
67 factor to falls in older adults (Canada et al., 2019). Neuroticism can be scored on a
68 continuum, ranging from emotional stability (low neuroticism) to emotional instability (high
69 neuroticism). It consists of multiple facets such as anxiety and feeling worried, tense and
70 fearful (McRae & Costa, 2008) that may contribute to the variance in fall-related
71 psychological concerns. Neuroticism has been linked to several health outcomes that
72 contribute to the higher risk of falling in old age, including frailty (Stephan et al., 2017),
73 muscle weakness (Tolea et al., 2012), physical inactivity (Sutin et al., 2016), cognitive
74 impairment (Terracciano et al., 2017), depressive symptoms (Hakulinen et al., 2015) and
75 anxiety (Kotov et al., 2007).

76 A recent study by Canada and colleagues (2019) found that higher neuroticism increased fall
77 risk among older adults over an 11-year follow-up. Their longitudinal study with a large
78 sample (N = 4 759) of older adults aged from 65 to 99 years confirmed previous cross-
79 sectional findings among older adults living independently in a retirement community
80 (Kloseck et al., 2007). The latter study also showed that higher neuroticism was associated
81 with lower confidence in avoiding a fall when performing daily tasks (Kloseck et al., 2007).
82 Similarly, a link between neuroticism and higher concern about falling has been found among
83 community-dwelling women over 70 (Mann et al., 2006). Concern about falling is an
84 important psychological factor in older people, as about one in three without a falls history
85 and about two in three with a falls history have expressed concern about falling (Kumar et al.,
86 2016). Moreover, concern about falling has been shown to increase fall risk, even in older
87 adults with no physiological risk factors for falls (Delbaere et al., 2010^a). Similarly,
88 depressive symptoms have been linked with an increased risk of falling, independently of the
89 presence of a higher physiological fall risk and poorer executive functioning (Kvelde et al.,
90 2015).

91 Despite the evidence on the associations between neuroticism and falls (Canada et al., 2019),
92 neuroticism and concern about falling (Mann et al., 2006), neuroticism and depressive
93 symptoms (Hakulinen et al., 2015), concern about falling and falls (Delbaere et al., 2010^a)
94 and depressive symptoms and falls (Kvelde et al. 2015), the associations between
95 neuroticism, concern about falling, depressive symptoms and actual falls have not previously
96 been investigated simultaneously. This study explored the associations between neuroticism,
97 concern about falling, depressive symptoms and indoor and outdoor falls. Based on the
98 existing evidence, we hypothesized, first, that neuroticism is associated with falls and,
99 second, that the relationship between neuroticism and falls is mediated by increased concern
100 about falling and depressive symptoms. In other words, we assumed that higher neuroticism

101 is associated with higher concern about falling and depressive symptoms, which in turn are
102 linked to actual falls. Given the multivariate nature of falls and differences between indoor
103 and outdoor fall risk profiles (Kelsey et al., 2010), we investigated indoor and outdoor falls
104 separately, taking into account several related factors.

105 Method

106 Participants

107 This cross-sectional study used baseline data gathered for a randomized controlled trial (the
108 PASSWORD study, Sipilä et al., 2018). Community-dwelling 70- to 85-year-old men and
109 women were randomly selected from the population register. The study recruitment strategy
110 and inclusion and exclusion criteria have been published earlier (Sipilä et al., 2018). Briefly,
111 inclusion criteria were sedentary or at most moderately active (walking < 150 min/wk., no
112 regular resistance training), able to walk 500 m without assistance and a score of ≥ 24 points
113 in a Mini Mental State Examination. Exclusion criteria included a severe chronic condition,
114 medication that could affect study participation, contraindications for physical exercise,
115 excessive use of alcohol, difficulties in communication due to severe hearing or vision
116 problems, and another family member already participating in the study. The flow chart of
117 the study is shown in Figure 1. Finally, 314 participants formed the study population. The
118 PASSWORD study was approved by the Ethical Committee of Central Finland Health Care
119 District (14/12/2016, ref.:11/2016). All participants signed an informed consent before the
120 baseline measurements.

121

122 Measurements

123 *Falls*

124 Information on indoor and outdoor falls during the previous year was collected
125 retrospectively by a structured questionnaire. The questions (two questions answered
126 separately) were: “How many times have you fallen indoors/outdoors during the previous
127 year?” The response options were 1= none, 2= once, 3= two to four times, 4= five to seven
128 times, and 5= eight times or more. For the analyses, participants who reported no falls during
129 the previous year were coded as “non-fallers”; those who reported ≥ 1 falls indoors were
130 coded as “indoor fallers”; and those who reported one fall outdoors, but no indoor falls, were
131 coded as “single outdoor fallers” (Pajala et al., 2008). In addition, those who had ≥ 2 outdoor
132 falls, but no indoor falls, were coded as “recurrent outdoor fallers” (Kelsey et al., 2012). We
133 used the dichotomized variables for indoor falls, single outdoor and recurrent outdoor falls
134 (0=no, 1= yes).

135

136 *Personality traits*

137 The trait of neuroticism in the short form of the Eysenck Personality Inventory modified by
138 Floderus (1974) was used. In the inventory, neuroticism is measured by 10 items (e.g., “Do
139 you often feel listless and tired without any special reason?”). Participants were asked to
140 answer ‘yes’ (= 1) or ‘no’ (= 0) to each item. The total score, ranging from 0 to 10, with
141 higher scores indicating a higher degree of neuroticism, was used in the analysis. Because a
142 missing value may result in excessively low sum scores, missing values were imputed by
143 calculating the probability of a positive response (2.9 % of participants had one missing
144 value). The short version of the Eysenck Personality Inventory is a widely used self-
145 completion tool that has been validated in many populations (Floderus-Myrhed et al., 1980;
146 Rose et al., 1988). Cronbach’s alpha for the neuroticism subscale was 0.72.

147 *Concern about falling*

148 The Falls Efficacy Scale-International (FES-I, Yardley et al., 2005) was used to assess level
149 of concern about falling when carrying out a wide range of activities, such as walking on
150 slippery, uneven or sloping surfaces, cleaning the house, shopping, or going to a social event.
151 The questionnaire contains 16 items, each rated on a four-point scale (1=not at all concerned
152 to 4= very concerned). Total scores, ranging from 16 to 64, were computed, a higher score
153 indicating more concern about falling. The psychometric properties of the FES-I have been
154 reported to be excellent (Yardley et al., 2005). In our population, Cronbach's alpha was 0.90.
155 Widely used cut-points differentiating between low (16–19), moderate (20–27) and high (28–
156 64) concern about falling have been presented by Delbaere and colleagues (2010^b).

157 *Depressive symptoms*

158 Depressive symptoms were interviewed using the 15-item Geriatric Depression Scale (GDS-
159 15, Yesavage & Sheikh, 1986) during the nurse examination. Summary scores, ranging from
160 0 to 11 (theoretical range 0-15), were computed; a score of zero to four is considered to be
161 within the normal range and a score of five or more to indicate depressive symptoms
162 (Almeida & Almeida, 1999). In this study, the GDS-15 was used as a continuous variable and
163 Cronbach's alpha was 0.60.

164 *Covariates*

165 *Executive function*

166 Executive functioning was assessed with the Trail Making Tests parts A and B (Reitan,
167 1958). Part A is a psychomotor speed task requiring the participant to connect as quickly as
168 possible and in ascending order a series of randomly dispersed circles containing numbers.
169 Part B requires the participant to shift attention by connecting as quickly as possible and in
170 alternating sequential order a series of randomly dispersed circles containing numbers and

171 letters. The time taken to complete each task is recorded and the difference in the time taken
172 to accomplish Parts B and A calculated. Smaller time differences indicate better performance.

173 *Physical performance and level of physical activity*

174 Physical performance was measured using the Short Physical Performance Battery (SPPB,
175 Guralnik et al., 1994). The SPPB includes habitual walking speed over four meters, five-time
176 chair rise, and standing balance tests. Summary scores range from 0 to 12, with higher scores
177 indicating better performance. Self-reported level of physical activity over the previous
178 month was assessed with a validated single question comprising 7 categories (Hirvensalo et
179 al., 1998): 0=I do not move more than is necessary in my daily routines/chores, 1=I go for
180 casual walks and engage in light outdoor recreation 1-2 times a week; 2=I go for casual walks
181 and engage in light outdoor recreation several times a week; 3=I engage, 1-2 times a week, in
182 brisk physical activity (e.g. yard work, walking, and cycling) to the point of perspiring and
183 some degree of breathlessness; 4=Several times a week (3-5), I engage in brisk physical
184 activity (e.g. yard work, walking, and cycling) to the point of perspiring and some degree of
185 breathlessness; 5=I do keep-fit exercise several times a week in a way that causes rather
186 strong shortness of breath and sweating during the activity; 6 = I participate in competitive
187 sports and maintain my fitness through regular training. Categories 5 and 6 were combined
188 with category 4 (only one participant reported a level 5 and no participant a level 6 activity).
189 For the background analysis, the variable was recoded into three categories: low (categories 0
190 and 1), medium (categories 2 and 3) and high (category 4).

191 To assess health status, information on chronic diseases, presence of persistent pain and
192 medication was collected by self-reports and from the national integrated patient information
193 system by the study physician. *Number of chronic diseases* was calculated as the sum of
194 conditions from the following list: high blood pressure, heart disease, stroke, lung disease,

195 diabetes, arthritis, arthrosis, chronic back disease, osteoporosis, and cancer. *Use of*
196 *psychotropic drugs* (no/yes) including opiates, benzodiazepines, anticholinergic agents,
197 dopaminergic agents, and antidepressants was documented according to the Anatomical
198 Therapeutic Chemical (ATC) classification. *Persistent musculoskeletal pain* in the lower
199 body was assessed by the question "During the past six months, have you suffered from pain
200 in the low back, hip, knee, ankle or foot daily or almost daily for at least one month?".
201 Participants who reported pain in one or more of these body regions were considered to have
202 persistent pain. In addition, *age*, *sex* (1=male, 2=female) and *education* in years were treated
203 as sociodemographic factors.

204 Statistics

205 Descriptive statistics were computed according to fall status (non-faller, single outdoor
206 faller/recurrent outdoor faller, indoor faller). Differences in continuous variable means
207 between the fall groups were examined using one-way ANOVA. Concern about falling and
208 neuroticism were non-normally distributed due to skewed data. Transformations [100/original
209 variable for concern about falling and $\ln(1+\text{original variable})$ for neuroticism] were used to
210 normalize the distribution. Pairwise group differences were tested with post-hoc Tukey's
211 multiple comparisons tests. A chi-square test was used for categorical variables according to
212 fall status. Associations between variables were calculated by Spearman's rank correlation
213 coefficients.

214 It has been shown that neuroticism is associated with falls (Canada et al., 2019) and concern
215 about falling (Mann et al., 2006) and that a bidirectional relation exists between concern
216 about falling and falls (Delbaere et al., 2010a). It was therefore hypothesized that concern
217 about falling would mediate the relationship between neuroticism and falls. Similarly,
218 neuroticism was assumed to predict depressive symptoms (Hakulinen et al., 2015), which in

219 turn are associated with increased fall risk (Kvelde et al. 2015). These factors were included
220 in the conceptual framework that was tested in this study (Figure 2). Previously identified
221 major confounders affecting the association between neuroticism and falls, neuroticism and
222 concern about falling/depressive symptoms and concern about falling/depressive symptoms
223 and falls were age, sex, use of psychotropic, chronic diseases, persistent pain, physical
224 performance, physical activity, and executive functioning. These factors were thus controlled
225 for in the analyses.

226 Path analysis was conducted to examine the pathways between neuroticism and indoor and
227 outdoor falls. We started from the conceptual framework adjusted for the confounders. We
228 used the maximum likelihood approach with Monte Carlo integration (5 000 nodes) which is
229 able to handle complex path models, including the present dichotomous fall outcome
230 variables, and missing values, assuming that missing data were generated by the missing-at-
231 random mechanism. The exploratory mediator model was used to investigate the pathways
232 between neuroticism and indoor and recurrent outdoor falling with concern about falling and
233 depressive symptoms as possible mediators.

234 In addition, differences in the mediation model parameters between males and females were
235 assessed by a likelihood ratio test based on gender-group modeling. Thus, we compared the
236 mediation parameters by gender (gender-specific model) to the mediation parameters in a
237 model where these parameters were constrained equal across the gender groups (pooled
238 gender model). The latter is a nested model of the former, and a non-significant result
239 indicates that the parameter estimates from the two models are statistically similar. Hence,
240 the pooled-gender model is taken to be a sufficient and more parsimonious description of the
241 associations (see, e.g., Jöreskog et al., 1993). The significance level was set at 0.05 (if the
242 95% confidence interval for a path coefficient did not contain the value zero, the parameter
243 was considered statistically significantly different from zero). We did not correct p-values for

244 multiple testing. Analyses were performed with Mplus Version 5.21 (Muthen & Muthen
245 1998-2009, Los Angeles, CA).

246 Results

247 Descriptive statistics are presented in Table 1. Mean participant age was 74.5 years (SD 3.8),
248 body mass index 28 kg² (5) and 60% were women. Thirty-seven participants (12%) had fallen
249 indoors, and the majority of them (n=33, 89%) had also had at least one outdoor fall. In all,
250 160 participants reported outdoor falls: 103 participants (33%) reported one fall, and 57
251 (18%) recurrent falls outdoors in the previous year. Table 1 presents the data on the 87
252 participants reporting one outdoor fall but no indoor falls, the 40 participants reporting ≥ 2
253 (recurrent) outdoor falls but no indoor falls, and the 37 participants reporting ≥ 1 indoor falls.
254 Those who had fallen either indoors or recurrently outdoors during the previous year had
255 reported higher concern about falling, and indoor fallers had also reported more depressive
256 symptoms than non-fallers and single-outdoor fallers. Recurrent outdoor fallers were also
257 more likely to have persistent musculoskeletal pain than non-fallers and single-outdoor
258 fallers. Compared to those reporting only one fall outdoors, indoor fallers had more
259 depressive symptoms and poorer physical performance. In addition, indoor fallers were more
260 concerned about falling, scored higher on neuroticism, and were more likely to be inactive
261 than those who reported one outdoor fall. However, except for their higher frequency in the
262 low physical activity group, indoor fallers did not differ from recurrent outdoor fallers (Table
263 1). In total, 156 participants (50%) were moderately concerned about falling (FES-I score 20
264 to 27) and 45 (14%) were highly concerned about falling (FES-I score more than 28).

265 The inter-correlations are presented in Table 2. Neuroticism correlated significantly with
266 indoor, single outdoor and recurrent outdoor falls. Neuroticism also correlated with concern
267 about falling and depressive symptoms. In turn, concern about falling correlated with indoor,
268 and recurrent outdoor falls, and depressive symptoms with indoor falls.

269 Figure 3 presents path coefficients for the final model estimating concern about falling and
270 depressive symptoms as mediators of the relationship between neuroticism and indoor falls.
271 In the adjusted model, neuroticism was positively associated with concern about falling. In
272 turn, concern about falling was positively associated with falling indoors. The indirect effect
273 of neuroticism on indoor falls via concern about falling was significant ($\beta = 0.34$, 95% CI
274 0.124, 0.561; $P = 0.002$). In addition, neuroticism was associated with indoor falls through
275 depressive symptoms ($\beta = 0.21$, 95% CI -0.004, 0.423; $P = 0.054$). Thus, the total effect of
276 neuroticism on indoor falls through the two indirect paths was $\beta = 0.55$, 95% CI 0.274, 0.831,
277 $P < 0.001$. This model explained 17% ($R^2=0.169$) of the variance in indoor falls. Likelihood
278 ratio tests indicated no statistically significant differences in the path coefficients across the
279 gender groups ($\chi^2_{df=11} = 15.25$, $P = 0.172$).

280 Neuroticism, concern about falling and depressive symptoms were not associated with having
281 only one outdoor fall. Therefore, the mediation analysis for having one outdoor fall was not
282 performed. However, we tested the hypothesis that the association between neuroticism and
283 recurrent outdoor falls is mediated by concern about falling and depressive symptoms.
284 Neuroticism was significantly associated with concern about falling, which in turn was
285 related to recurrent outdoor falls (Figure 4). The mediation analysis revealed a statistically
286 significant indirect pathway between neuroticism and recurrent outdoor falls through concern
287 about falling ($\beta = 0.19$, 95% CI 0.004, 0.371, $P = 0.045$). Depressive symptoms were not
288 associated with recurrent outdoor falls. This model accounted for 10% ($R^2=0.096$) of the
289 variance in recurrent outdoor falls. We found no statistically significant differences in the
290 mediation parameters in the gender-group analysis (recurrent outdoor falls $\chi^2_{df=10} = 13.22$, $P =$
291 0.212).

292

293 Discussion

294 The main aim of the present study was to examine whether concern about falling and/or
295 depressive symptoms function as mediators between neuroticism and falling. The correlations
296 showed that neuroticism was associated with both indoor and recurrent outdoor falls. The
297 mediation analysis revealed that a higher score on neuroticism was positively associated with
298 higher concern about falling, which, in turn, was associated with indoor falls and recurrent
299 outdoor falls. In addition, neuroticism was associated with higher depressive symptoms,
300 which in turn were linked to indoor falls only. Thus, our findings suggest that concern about
301 falling mediates the association between neuroticism and recurrent outdoor falls and that
302 concern about falling and depressive symptoms mediate the relationship between neuroticism
303 and indoor falls. The associations between neuroticism and falls remained significant when
304 adjusted for traditional risk factors for falls, such as poor physical performance, low level of
305 physical activity, poor executive functioning, persistent pain, chronic diseases, and the use of
306 psychotropic medications. These results emphasize that, to prevent falls among older people,
307 interventions are needed that also take account of psychological factors.

308 Our findings extend those of a previous study reporting a link between neuroticism and
309 higher fall risk (Canada et al., 2019) by examining indoor and outdoor fall outcomes
310 separately and the mediator role of higher concern about falling and depressive symptoms in
311 the association between neuroticism and falls. Indoor and outdoor falls differ in older adults:
312 relatively healthy and physically active older adults more often fall outdoors than indoors,
313 whereas indoor falls are associated with activity restriction, health problems and poorer
314 physical performance (Kelsey et al., 2010; Mänty et al., 2009). In line with the literature, the
315 present participants reporting indoor falls tended to be more inactive, to be more concerned
316 about falling, and to have more depressive symptoms than those reporting outdoor falls.
317 Among these relatively healthy, community-living older adults, those who had fallen indoors

318 also seemed to be prone to recurrent outdoor falls, supporting earlier findings that recurrent
319 falls are indicative of an underlying high-risk state (Deandre et al., 2010). Non-recurrent
320 outdoor falls, in turn, seem to be more coincidental and may be related to contextual factors
321 such as environmental hazards (Nyman et al., 2013).

322 This study supports earlier findings showing that higher concern about falling (Delbaere et
323 al., 2010^a) and higher depressive symptomatology (Kvelde et al. 2015) are associated with
324 increased fall risk, even in older adults without obvious physical risk factors. Among older
325 adults who score higher on neuroticism, increased concern about falling and decreased
326 confidence on their ability to undertake activities of daily living without falling may relate to
327 a persistent and dysfunctional disruption of attention and more cautious behavior (Delbaere et
328 al., 2010^a). Concern about falling may, for example, impact negatively on gait. It can lead to
329 so called “cautious” gait, which in turn results in decreased walking stability and increased
330 risk of falling (Delbaere et al., 2009). A recent study by Tuerk and colleagues (2016) showed
331 that increased concern about falling was associated with decreased gray matter volume in the
332 brain areas important for motor control and hence gait safety. Interestingly, their results
333 suggested that this relationship was explained by generalized anxiety and neuroticism rather
334 than physical risk factors for falls. In addition, persons overly concerned about falling or
335 having depressive feelings may restrict their activities of daily living (Zijlstra et al., 2007),
336 resulting in increased physical inactivity, loss of muscle strength and impaired physical
337 condition, all of which are known common risk factors for falls (Deandrea et al., 2010).
338 Moreover, there is evidence that physical inactivity (Sutin et al., 2016), poor muscle strength
339 (Tolea et al., 2012) and deteriorated physical condition (Stephan et al., 2017) are linked with
340 neuroticism. However, we controlled for these common risk factors for falls and found that
341 increased fall risk was a result not only of physical deterioration but also of the psychological
342 challenges experienced by persons scoring high on emotional instability.

343 Individuals scoring high on neuroticism may have more depressive symptomatology, general
344 worries and anxiety (Kotov et al., 2007), factors that have also typically been associated with
345 fall-related psychological concerns (Payette et al., 2015) and lower self-efficacy (O'Shea et
346 al., 2016). In the present study, having depressive symptoms was strongly associated with
347 neuroticism and also directly with indoor falls despite the exclusion of individuals with
348 clinical depression. In addition, Yasunaga and Yaguchi (2014) found that poor self-efficacy
349 mediated the association between neuroticism and a lower level of physical activity among
350 relatively healthy older adults. Concern about falling among older adults who are emotionally
351 unstable may result from the interplay between multiple factors such as lower self-efficacy,
352 anxiety or worrying, depressive mood, and persistent pain. For example, persistent pain that
353 interferes with activities of daily living considerably reduces self-efficacy and increases older
354 adults' concern about falling (Stubbs et al., 2014). Prolonged lower body pain has also been
355 associated with impaired balance (Lihavainen et al., 2009) and risk for falls (Leveille et al.,
356 2009) in older people. Among the present relatively healthy older adults, persistent
357 musculoskeletal pain was related in particular to recurrent outdoor falls. Sex did not moderate
358 the relationship between neuroticism and falls tested in our mediation models, although
359 women reported higher rates of concern about falling than men, as also found previously
360 (Delbaere et al., 2010). Thus, several factors may lead to concern about falling and hence to
361 less effective coping strategies (Loft et al., 2018). The latter in turn increase vulnerability to
362 falls in older adults with relatively high scores on the personality trait of neuroticism.

363 Whereas balance and muscle strength training have proven to be effective in reducing falls
364 (Sherrington et al., 2019), the effects of training on reducing concern about falling are less
365 clear (Kumar et al., 2016). In addition to exercise, successful interventions may need
366 behavioral components aimed at reducing psychological concerns and depressive mood and
367 enhancing self-efficacy (Ziljstra et al., 2009). Moreover, enhancing self-efficacy may

368 improve coping with persistent pain (Turner et al., 2005). Consequently, multicomponent
369 interventions that better take individuals' characteristics into account may be the most
370 effective way of preventing falls in older adults.

371 This study has its limitations. First, the cross-sectional design does not allow any conclusions
372 to be drawn on causality. On the other hand, it has been argued that the association between
373 fall-related psychological concerns and falls is bidirectional (Friedman et al., 2002). In
374 addition, the personality trait of neuroticism is relatively stable throughout adulthood (Kokko
375 et al., 2015) and old age (Steunenberg et al., 2005). Future research could build upon present
376 findings by examining these associations over time. In addition, our measure of neuroticism
377 (10 items) did not allow examination of the facets of this trait. The use of a broader measure
378 of neuroticism would be needed to understand whether specific facets of the trait may be
379 driving these results. A further, major, limitation is that falls were measured retrospectively,
380 and hence may be affected by recall bias. Moreover, the participants with a physician
381 diagnosis of major depression were excluded from the study. Thus, the data may under- or
382 overestimate falls (Sanders et al., 2009) and under-estimate the effect of depressive
383 symptoms. Future studies should use prospective monitoring with daily falls calendars and
384 separate the injurious falls from falls with less severe consequences. There is evidence that
385 higher levels of anxiety, for example, predict fall-related fractures (Catalano et al. 2018).
386 Thus, the possibility of an enhanced risk of fall-related fractures among older people with
387 higher level of neuroticism, should be considered in future works.

388 A strength is the use of detailed and valid assessments of several well-known risk factors for
389 falls. A large sample of community-dwelling older adults with sedentary or at most moderate
390 levels of physical activity but free from severe chronic conditions participated, allowing us to
391 generalize the results to the healthy older adults. Levels of concern about falling (Tuerk et al.,
392 2016) and fall levels (Lord et al., 1993; Schoene et al., 2014), were representative of those

393 among community-dwelling older adults. In addition, the mean scores for neuroticism were
394 comparable to those previously found in large population-based studies of adults (Navrady et
395 al., 2019).

396 In conclusion, this study provides new evidence on the mechanism through which a higher
397 score on neuroticism, also known as emotional instability, may contribute to increased fall
398 risk. Fall-related psychological concerns such as concern about falling could become an
399 important target for more personalized interventions aimed at reducing the incidence of falls
400 in community-dwelling older adults.

401 Funding: This work was supported by the Academy of Finland (296843) and the Ministry of
402 Education and Culture of Finland (OKM/49/626/2017, OKM/72/626/2018,
403 OKM/92/626/2019).

404

405 Acknowledgments

406 We are grateful to all the PASSWORD participants who contributed their time and
407 information. We also thank the staff members who helped with the PASSWORD data
408 collection.

409 References

- 410 Almeida, O. P., & Almeida, S. A. (1999). Short versions of the geriatric depression scale: a
411 study of their validity for the diagnosis of a major depressive episode according to ICD-10
412 and DSM-IV. *International Journal of Geriatric Psychiatry*, 14, 858-865. doi:
413 10.1002/(sici)1099-1166(199910)14:10<858::aid-gps35>3.0.co;2-8
- 414 Canada, B., Stephan, Y., Sutin, A., & Terracciano, A. (2019). Personality and falls among
415 older adults: evidence from a longitudinal cohort. *The Journals of Gerontology, Series B:
416 Psychological and Social Sciences*, gbz040, doi: 10.1093/geronb/gbz040
- 417 Caspi, A. (1998). Personality development across the life course. In W. Damon & N.
418 Eisenberg (Eds.), *Handbook of child psychology: Social, emotional, and personality
419 development* (pp. 311–388). John Wiley & Sons Inc.
- 420 Catalano, A., Martino, G., Bellone, F., Gaudio, A., Lasco, C., Langher, V., Lasco, A., &
421 Morabito, N. (2018). Anxiety levels predict fracture risk in postmenopausal women assessed
422 for osteoporosis. *Menopause*, 25(10), 1110–1115. doi:10.1097/GME.0000000000001123
- 423 Deandrea, S., Lucenteforte, E., Bravi, F., Foschi, R., La Vecchia, C., & Negri, E. (2010).
424 Risk factors for falls in community-dwelling older people: a systematic review and meta-
425 analysis. *Epidemiology*, 21, 658–668. doi: 10.1097/EDE.0b013e3181e89905
- 426 Delbaere, K.^a, Close, J. C., Brodaty, H., Sachdev, P., & Lord, S. R. (2010). Determinants of
427 disparities between perceived and physiological risk of falling among elderly people: cohort
428 study. *The British Medical Journal*, 18;341:c4165. doi: 10.1136/bmj
- 429 Delbaere, K.^b, Close, J. C., Mikolaizak, A. S., Sachdev, P. S., Brodaty, H., & Lord, S. R.
430 (2010). The Falls Efficacy Scale International (FES-I). A comprehensive longitudinal
431 validation study. *Age and Ageing*, 39, 210-216. doi: 10.1093/ageing/afp225

432 Delbaere, K., Sturnieks, D. L., Crombez, G., & Lord, S. R. (2009). Concern about falls elicits
433 changes in gait parameters in conditions of postural threat in older people. *The Journals of*
434 *Gerontology, Series A, Biological Sciences and Medical Sciences*, 64, 237-242. doi:
435 10.1093/gerona/gln014

436 Digman, J. M. (1990). Personality structure: Emergence of the five-factor model. *Annual*
437 *Review of Psychology*, 41, 417–440. doi:10.1146/annurev.ps.41.020190.002221

438 Floderus, B. (1974). Psycho-social factors in relation to coronary heart disease and associated
439 risk factors. *Nordisk Hygienisk Tidskrift*, Suppl 6, 7–148.

440 Floderus-Myrhed B., Pedersen, N., & Rasmuson, I. (1980). Assessment of heritability for
441 personality, based on a short-form of the Eysenck Personality Inventory: a study of 12,898
442 twin pairs. *Behavior Genetics*, 10, 153-162. doi: 10.1007/bf01066265

443 Friedman, S. M., Munoz, B., West, S. K., Rubin, G. S., & Fried, L. P. (2002). Falls and fear
444 of falling: which comes first? A longitudinal prediction model suggests strategies for primary
445 and secondary prevention. *Journal of the American Geriatrics Society*, 50, 1329-1335.
446 10.1046/j.1532-5415.2002.50352.x

447 Guralnik, J. M., Simonsick, E. M., Ferrucci, L., Glynn, R. J., Berkman, L. F., Blazer, D. G.,
448 Scherr, P.A., & Wallace, R. B. (1994). A short physical performance battery assessing lower
449 extremity function: association with self-reported disability and prediction of mortality and
450 nursing home admission. *Journal of Gerontology*, 49, 85. doi: 10.1093/geronj/49.2.M85

451 Hakulinen, C., Elovainio, M., Pulkki-Råback, L., Virtanen, M., Kivimäki, M., & Jokela, M.
452 (2015). Personality and depressive symptoms: Individual participant meta-analysis of 10
453 cohort studies. *Depression and Anxiety*, 32, 461–470. doi:10.1002/da.22376

454 Hirvensalo, M., Lampinen, P., & Rantanen, T. (1998). Physical exercise in old age: An eight-
455 year follow-up study on involvement, motives, and obstacles among persons age 65–84.
456 *Journal of Aging and Physical Activity*, 6, 157–168. doi: 10.1123/japa.6.2.157

457 Jöreskog, K.G., & Sörbom, D. (1993). LISREL 8: Structural Equation Modeling with the
458 SIMPLIS Command Language. Scientific Software International; Lawrence Erlbaum
459 Associates, Inc.

460 Kelsey, J. L., Berry, S. D., Procter-Gray, E., Quach, L., Nguyen, U. S., Li, W., Kiel, D. P.,
461 Lipsitz, L. A., & Hannan, M. T. (2010). Indoor and outdoor falls in older adults are different:
462 the maintenance of balance, independent living, intellect, and Zest in the Elderly of Boston
463 Study. *Journal of the American Geriatrics Society*, 58, 2135-2141. doi: 10.1111/j.1532-
464 5415.2010.03062.x

465 Kelsey, J. L., Procter-Gray, E., Berry, S.D., Hannan, M.T., Kiel, D.P., Lipsitz, L.A., & Li, W.
466 (2012). Reevaluating the implications of recurrent falls in older adults: location changes the
467 inference. *Journal of the American Geriatrics Society*, 60, 517-24. doi: 10.1111/j.1532-
468 5415.2011.03834.x.

469 Klooseck, M., Hobson, S., Crilly, R., Vandervoort, A., & Ward-Griffin, C. (2007). The
470 influence of personality on falling and engagement in daily activities by community-dwelling
471 older adults. *Physical & Occupational Therapy in Geriatrics*, 26, 1–17.
472 doi:10.1080/J148v26n01_01

473 Kokko, K., Rantanen, J., & Pulkkinen, L. (2015). Associations between mental well-being
474 and personality from a life span perspective. In M. Blatny (Ed.), *Personality and well-being*
475 *across the life-span* (pp. 134–159). Palgrave Macmillan

476 Kotov, R., Watson, D., Robles, J. P., & Schmidt, N. B. (2007). Personality traits and anxiety
477 symptoms: the multilevel trait predictor model. *Behaviour Research and Therapy*, 45, 1485-
478 503. doi:10.1016/j.brat.2006.11.011

479 Kumar, A., Delbaere, K., Zijlstra, G. A., Carpenter, H., Iliffe, S., Masud, T., Skelton, D.,
480 Morris, R., & Kendrick, D. (2016). Exercise for reducing fear of falling in older people
481 living in the community: Cochrane systematic review and meta-analysis. *Age and Ageing*, 45,
482 345-352. doi: 10.1093/ageing/afw036

483 Kvelde, T., Lord, SR., Close, JCT., Reppermund, S., Kochan, NA., Sachdev P, Brodaty, H.,
484 & Delbaere, K. Depressive symptoms increase fall risk in older people, independent of
485 antidepressant use, and reduced executive and physical functioning. *Archives of Gerontology*
486 *and Geriatrics*, 60, 190-195. doi: 10.1016/j.archger.2014.09.003.

487 Leveille, S. G., Jones, R. N., Kiely, D. K., Hausdorff, J. M., Shmerling, R. H., Guralnik, J.
488 M., Kiel, D. P., Lipsitz, L. A., & Bean, J. F. (2009). Chronic musculoskeletal pain and the
489 occurrence of falls in an older population. *JAMA*, 302, 2214-2221. doi:
490 10.1001/jama.2009.1738

491 Lihavainen, K., Sipilä, S., Rantanen, T., Sihvonen, S., Sulkava, R., & Hartikainen, S. (2010).
492 Contribution of musculoskeletal pain to postural balance in community-dwelling people aged
493 75 years and older. *The Journals of Gerontology, Series A, Biological Sciences and Medical*
494 *Sciences*, 65, 990-996. doi: 10.1093/gerona/glq052

495 Loft, C. C., Jones, F. W., & Kneebone, I. I. (2018). Falls self-efficacy and falls incidence in
496 community-dwelling older people: the mediating role of coping. *International*
497 *Psychogeriatrics*, 30, 727-733. doi: 10.1017/S1041610217002319

498 Lord, S. R., Ward, J. A., Williams, P., & Anstey, K. J. (1993). An epidemiological study of
499 falls in older community-dwelling women: the Randwick falls and fractures study. *Australian*
500 *Journal of Public Health*, 17, 240-245. <https://doi.org/10.1111/j.1753-6405.1993.tb00143.x>

501 Mann, R., Birks, Y., Hall, J., Torgerson, D., & Watt, I. (2006). Exploring the relationship
502 between fear of falling and neuroticism: a cross-sectional study in community-dwelling
503 women over 70. *Age and Ageing*, 35, 143–147. doi:10.1093/ageing/afj013

504 McCrae, R. R., & Costa, P. T. (2008). Empirical and theoretical status of the five-factor
505 model of personality traits. In G. J. Boyle, G. Matthews, & D. H. Saklofske (Eds.), *The SAGE*
506 *handbook of personality theory and assessment, Vol. 1. Personality theories and models* (pp.
507 273–294). Sage Publications, Inc.

508 Mänty, M., Heinonen, A., Viljanen, A., Pajala, S., Koskenvuo, M., Kaprio, J., & Rantanen, T.
509 (2009). Outdoor and indoor falls as predictors of mobility limitation in older women. *Age and*
510 *Ageing*, 38, 757-761. doi: 10.1093/ageing/afp178

511 Navrady, L. B., Adams, M. J., Chan, S. W. Y.; Major Depressive Disorder Working Group of
512 the Psychiatric Genomics Consortium, Ritchie, S. J., & McIntosh, A. M. (2018). Genetic risk
513 of major depressive disorder: the moderating and mediating effects of neuroticism and
514 psychological resilience on clinical and self-reported depression. *Psychological Medicine*, 48,
515 1890-1899. doi: 10.1017/S0033291717003415

516 Nyman, S. R., Ballinger, C., Phillips, J. E., & Newton, R. (2013). Characteristics of outdoor
517 falls among older people: a qualitative study. *BMC Geriatrics*, 18, 13, 125. doi:
518 10.1186/1471-2318-13-125

519 O'Shea, D. M., Dotson, V. M., & Fieo, R. A. (2017). Aging perceptions and self-efficacy
520 mediate the association between personality traits and depressive symptoms in older adults.
521 *International Journal of Geriatric Psychiatry*, 32, 1217-1225. doi: 10.1002/gps.4584

522 Pajala, S., Era, P., Koskenvuo, M., Kaprio, J., Törmäkangas, T., & Rantanen, T. (2008).
523 Force platform balance measures as predictors of indoor and outdoor falls in community-
524 dwelling women aged 63-76 years. *The Journals of Gerontology, Series A, Biological*
525 *Sciences and Medical Sciences*, 63, 171-178. doi: 10.1093/gerona/63.2.171.

526 Payette, M. C., Bélanger, C., Léveillé, V., & Grenier, S. (2016). Fall-related psychological
527 concerns and anxiety among community-dwelling older adults: systematic review and meta-
528 Analysis. *PLoS One*, 4,e0152848. doi: 10.1371/journal.pone.0152848

529 Reitan, R. M. (1958). Validity of the trail making test as an Indicator of organic brain
530 damage. *Perceptual and Motor Skills*, 8, 271–276. doi: 10.2466/pms.1958.8.3.271

531 Rose, R. J., Koskenvuo, M., Kaprio, J., Sarna, S., & Langinvainio, H. (1988). Shared genes,
532 shared experiences, and similarity of personality: data from 14,288 adult Finnish co-twins.
533 *Journal of Personality and Social Psychology*, 54, 161-171. doi: 10.1037//0022-
534 3514.54.1.161

535 Sanders, K. M., Hayles, A. L., Kotowicz, M. A., Nicholson, G. C. (2009). Monitoring falls in
536 cohort studies of community-dwelling older women. *Journal of the American Geriatrics*
537 *Society*, 57, 733–734. <http://dx.doi.org/10.1111/j.1532-5415.2009.02205.x>

538 Schoene, D., Smith, S. T., Davies, T. A., Delbaere, K., & Lord, S. R. (2014). A Stroop
539 Stepping Test (SST) using low-cost computer game technology discriminates between older
540 fallers and non-fallers. *Age and Ageing*, 43, 285–289. doi: 10.1093/ageing/aft157

541 Sherrington, C., Fairhall, N., Wallbank, G., Tiedemann, A., Michaleff, Z. A., Howard, K.,
542 Clemson, L., Hopewell, S., & Lamb, S. (2019). Exercise for preventing falls in older people
543 living in the community: an abridged Cochrane systematic Review. *British Journal of Sports*
544 *Medicine*, pii: bjsports-2019-101512. doi: 10.1136/bjsports-2019-101512

545 Sipilä, S., Tirkkonen, A., Hänninen, T., Laukkanen, P., Alen, M., Fielding, R.A, Kivipelto,
546 M., Kokko, K., Kulmala, J., Rantanen, T., Sihvonen, S. E., Sillanpää, E., Stigsdotter-Neely,
547 A., &. Törmäkangas, T. (2018). Promoting safe walking among older people: the effects of a
548 physical and cognitive training intervention vs. physical training alone on mobility and falls
549 among older community-dwelling men and women (the PASSWORD study): design and
550 methods of a randomized controlled trial. *BMC Geriatrics*, 18, 215. doi: 10.1186/s12877-
551 018-0906-0

552 Stephan, Y., Sutin, A. R., Canada, B., & Terracciano, A. (2017). Personality and frailty:
553 Evidence from four samples. *Journal of Research in Personality*, 66, 46–53.
554 doi:10.1016/j.jrp.2016.12.006

555 Steunenberg, B., Twisk, J. W., Beekman, A. T., Deeg, D. J., & Kerkhof, A. J. (2005).
556 Stability and change of neuroticism in aging. *The Journals of Gerontology: Series B*
557 *Psychological and Social Sciences*, 60, P27–P33. doi:10.1093/geronb/60.1.p27

558 Stubbs, B., Eggermont, L. H., Patchay, S., & Schofield, P. A. (2014). Pain interference is
559 associated with psychological concerns related to falls in community-dwelling older adults:
560 multisite observational study. *Physical Therapy*, 94, 1410-1420. doi: 10.2522/ptj.20140093

561 Sutin, A. R., Stephan, Y., Luchetti, M., Artese, A., Oshio, A., & Terracciano, A. (2016). The
562 five-factor model of personality and physical inactivity: A meta-analysis of 16 samples.
563 *Journal of Research in Personality*, 63, 22–28. doi:10.1016/j.jrp.2016.05.001

564 Terracciano, A., Stephan, Y., Luchetti, M., Albanese, E., Sutin, A. R. (2017). Personality
565 traits and risk of cognitive impairment and dementia. *Journal of Psychiatric Research*, 89,
566 22-27. doi: 10.1016/j.jpsychires.2017.01.011

567 Tolea, M. I., Terracciano, A., Simonsick, E. M., Metter, E. J., Costa, P. T. Jr., & Ferrucci, L.
568 (2012). Associations between personality traits, physical activity level, and muscle strength.
569 *Journal of Research in Personality*, 46, 264–270. doi:10.1016/j.jrp.2012.02.002

570 Tuerk, C., Zhang, H., Sachdev, P., Lord, S. R., Brodaty, H., Wen, W., & Delbaere, K. (2016).
571 Regional gray matter volumes are related to concern about falling in older people: A voxel-
572 based morphometric study. *The Journals of Gerontology, Series A, Biological Sciences and*
573 *Medical Sciences*, 71, 138-44. doi: 10.1093/gerona/glu242

574 Turner, J. A., Ersek, M., & Kemp, C. (2005). Self-efficacy for managing pain is associated
575 with disability, depression, and pain coping among retirement community residents with
576 chronic pain. *Journal of Pain*, 6, 471-479. doi: 10.1016/j.jpain.2005.02.011

577 Zijlstra, G. A., van Haastregt, J. C., van Eijk, J. T., van Rossum, E., Stalenhoeef, P. A., &
578 Kempen, G. I. (2007). Prevalence and correlates of fear of falling, and associated avoidance
579 of activity in the general population of community-living older people. *Age and Ageing*, 36,
580 304-309. doi:10.1093/ageing/afm021

581 Zijlstra, G. A., Van Haastregt, J. C., Ambergen, T., Van Rossum, E., Van Eijk, J. T.,
582 Tennstedt, S. L., & Kempen, G. I. (2009). Effects of a multicomponent cognitive behavioral
583 group intervention on fear of falling and activity avoidance in community-dwelling older
584 adults: results of a randomized controlled trial. *Journal of the American Geriatrics Society*,
585 57, 2020–2028. <https://doi.org/10.1111/j.1532-5415.2009.02489.x>

586 Yasunaga, A., & Yaguchi, K. (2014). Personality traits, self-efficacy for exercise, and
587 exercise levels in older Japanese adults. *The Japanese Journal of Health Psychology*, 27, 1–
588 11. https://doi.org/10.11560/jahp.27.1_1

589 Yardley, L., Beyer, N., Hauer, K., Kempen, G., Piot-Ziegler, C., & Todd, C. (2005).
590 Development and initial validation of the falls efficacy scale-international (FES-I). *Age and*
591 *Ageing*, 34, 614–619. doi: 10.1093/ageing/afi196

592 Yesavage, J. A. & Sheikh, J. I. (1986). Geriatric depression scale (GDS) recent evidence and
593 development of a shorter version. *Clinical Gerontologist: The Journal of Aging and Mental*
594 *Health*, 5, 165-173. https://doi.org/10.1300/J018v05n01_09

Table 1. Participant characteristics and the fall status, n=314.

Variables	<u>Non-fallers</u> ¹	<u>Outdoors fallers</u>		<u>Indoor fallers</u> ⁴	1 vs 2	1 vs 3	1 vs 4	2 vs 3	2 vs 4	3 vs 4
	(n=150)	Single ² (n=87)	Recurrent ³ (n=40)	(n=37)	P-value	P-value	P-value	P-value	P-value	P-value
Age, years [mean (SD)]	74.7 (3.6)	73.9 (3.7)	75.0 (4.4)	74.3 (4.1)	.410	.957	.970	.399	.917	.874
Sex [n (%) female]	95 (63)	50 (58)	24 (60)	19 (51)	.372	.699	.181	.788	.530	.455
Level of education [n (%) highest]	33 (22)	21 (24)	6 (15)	6 (16)	.705	.330	.438	.242	.328	.883
Executive function, TMTΔ seconds [mean (SD)]	93.0 (61.1)	76.7 (32.2)	90.2 (47.1)	95.9 (54.9)	.101	.991	.990	.532	.248	.532
Number of chronic diseases, [mean (SD)]	1.5 (1.1)	1.5 (1.2)	1.6 (1.2)	1.7 (1.2)	.983	.632	.939	.961	.938	.928

Depressive symptoms, GDS-15 score [mean (SD)]	1.4 (1.4)	1.6 (1.6)	1.8 (1.7)	2.5 (2.5)	.798	.512	.002	.919	.032	.919
Use of psychotropic [n (%) yes]	21 (14)	11 (13)	7 (18)	6 (16)	.768	.579	.731	.466	.597	.881
Physical performance, SPPB score [mean (SD)]	10.1 (1.6)	10.5 (1.4)	10.2 (1.2)	9.6 (2.0)	.314	.991	.286	.786	.024	.345
Level of physical activity [n (%) yes]										
Low	60 (40)	32 (37)	13 (33)	21 (57)	.624	.386	.065	.639	.040	.032
Medium	76 (51)	42 (48)	20 (50)	10 (27)						
High	14 (9)	13 (15)	7 (17)	6 (16)						
Presence of persistent pain [n (%) yes]	43 (29)	32 (37)	23 (58)	14 (38)	.175	.001	.278	.033	.947	.084
Concern about falling, FES-I score [mean (SD)]	21.6 (4.6)	21.5 (4.1)	24.3 (4.9)	26.5 (8.4)	.970	.010	< .001	.016	< .001	.358

Neuroticism, score [mean (SD)]	2.9 (2.2)	2.7 (2.0)	3.7 (2.1)	4.5 (2.7)	.960	.166	.011	.105	.007	.808
--------------------------------	-----------	-----------	-----------	-----------	------	------	------	------	------	------

Notes: P-values were calculated for continuous variables using one-way ANOVA post hoc comparisons, Tukey's test. Between-group differences in categorical variables were tested using Chi-square test.

Table 2. Inter-correlations of the study variables

Variable	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
1. Single outdoor falls	1	-.29**	-.23**	-.11	-.09	-.01	.02	.12*	-.03	-.02	.12*	.05	.00	-.03	-.10
2. Recurrent outdoor falls		1	.26**	.21**	-.26**	.10	.17**	-.02	.04	.02	.05	.04	-.07	-.05	.03
3. Indoor falls			1	.19**	-.18**	.12*	.02	-.09	.02	.03	.04	-.09	-.04	-.06	-.02
4. Neuroticism				1	.37**	.48**	.11*	-.09	.27*	.11	-.00	-.02	-.05	.07	.11
5. Concern about falling					1	-.27**	-.24**	.32**	-.09	-.15**	-.04	.04	.02	-.26**	-.08
6. Depressive symptoms						1	.13*	-.22**	.23**	.17**	.00	-.15**	.03	.04	.07
7. Persistent pain							1	-.20**	.11	.27**	.05	-.11*	-.09	.08	.06
8. Physical performance								1	-.04	-.20**	-.22**	.19**	.14*	-.28**	-.17**
9. Use of psychotropic									1	-.00	.04	-.03	-.02	.06	.03
10. Chronic diseases										1	.15**	-.14*	-.15**	-.04	.20**
11. Executive function											1	-.05	-.37**	-.13*	.32**

12. Level of physical activity												1	.04	-.03	-.06
13. Level of education													1	.14*	-.02
14. Sex														1	.04
15. Age															1

Notes: *p<.05, **p<.01.




Figure 1. Flow chart of the study




Figure 2. Conceptual framework tested.




Figure 3. Standardized path coefficients (95% confidence intervals) of the mediator model for neuroticism, concern about falling and indoor falls among community-dwelling older people (n=314). Note: the model is controlled for age, sex, use of psychotropic, chronic diseases, persistent pain, physical performance, physical activity, and executive functioning; non-significant control variables were removed from the final mediation model.




Figure 4. Standardized path coefficients (95% confidence intervals) of the mediator model for neuroticism, concern about falling and recurrent outdoor falls among community-dwelling older people (n=314). Note: the model is controlled for age, sex, use of psychotropic, chronic diseases, persistent pain, physical performance, physical activity, and executive functioning; only significant associations of these variables are presented.

