

**This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.**

**Author(s):** Zhu, Qihao; Fettinger, James C.; Vasko, Petra; Power, Philip P.

**Title:** Interactions of a Diplumbyne with Dinuclear Transition Metal Carbonyls to Afford Metalloplumbylenes

**Year:** 2020

**Version:** Accepted version (Final draft)

**Copyright:** © 2020 American Chemical Society

**Rights:** In Copyright

**Rights url:** <http://rightsstatements.org/page/InC/1.0/?language=en>

**Please cite the original version:**

Zhu, Q., Fettinger, J. C., Vasko, P., & Power, P. P. (2020). Interactions of a Diplumbyne with Dinuclear Transition Metal Carbonyls to Afford Metalloplumbylenes. *Organometallics*, 39(24), 4629-4636. <https://doi.org/10.1021/acs.organomet.0c00659>

# Interactions of a Diplumbyne with Dinuclear Transition Metal Carbonyls to Afford Metalloplumbylenes

Qihao Zhu, Petra Vasko, James C. Fettinger, and Philip P. Power\*

## Abstract:

The metathesis reactions of the diplumbyne  $\text{Ar}^{\text{iPr}_6}\text{PbPbAr}^{\text{iPr}_6}$  ( $\text{Ar}^{\text{iPr}_6} = -\text{C}_6\text{H}_3-2,6-(\text{C}_6\text{H}_2-2,4,6-\text{iPr}_3)_2$ ) with the dinuclear metal carbonyls  $\text{Mn}_2(\text{CO})_{10}$ ,  $\text{Fe}_2(\text{CO})_9$ ,  $\text{Co}_2(\text{CO})_8$  under mild conditions afforded the complexes  $\text{Mn}(\text{CO})_5(\text{PbAr}^{\text{iPr}_6})$  (**1**),  $\text{Fe}(\text{CO})_4(\text{PbAr}^{\text{iPr}_6})_2$  (**2**),  $\text{Co}_4(\text{CO})_9(\text{PbAr}^{\text{iPr}_6})_2$  (**3**), respectively. Complexes **1-3** were structurally characterized by single-crystal X-ray diffraction, and by  $^1\text{H}$ ,  $^{13}\text{C}\{^1\text{H}\}$ ,  $^{59}\text{Co}\{^1\text{H}\}$ , and  $^{207}\text{Pb}\{^1\text{H}\}$  NMR, UV-Vis, and IR spectroscopy. They are rare examples of species formed by the direct reaction of a group 14 dimetallyne with transition metal carbonyls. Complexes **1** and **2** feature Mn-Pb or Fe-Pb single bonds whereas in **3** a Co-Pb cluster is formed in which the plumbylidyne moiety bridges either an edge or a face of a  $\text{Co}_4$  carbonyl cluster.


## Introduction

Organometallic complexes containing bonds between transition metals and low oxidation state group 14 element moieties are of interest because of their potential for a diverse reactivity thus arises mainly from the unsaturated character of the group 14 element bound directly to the transition metal.<sup>1</sup> For the heavier group 14 elements, early work by Marks<sup>2</sup> and Lappert<sup>3</sup> reported the synthesis of the first stannylene-transition metal complexes which were characterized spectroscopically<sup>2-3</sup> and structurally.<sup>3</sup> These initial examples have been followed by numerous other  $\text{L}_n\text{MER}_2$  (L = ligand; M = transition metal; E = heavier group 14 element; R = organic & related substituents)


complexes that have incorporated the other heavier group 14 elements and various transition metals.<sup>4-6</sup> Related to these complexes are species of formula  $L_nM-ER$  in which the transition metal is  $\sigma$ -bonded directly to the group 14 element. The first such complex,  $Mes^*GeFe(CO)_2R$  ( $Mes^* = \text{supermesityl}$ , i.e.  $C_6H_2-2,4,6-^tBu_3$ ;  $R = Cp/Cp^*$ ), which was reported by Jutzi and Leue in 1994, was characterized spectroscopically.<sup>7</sup> Later Power and coworkers reported the synthesis of metallostanlylene<sup>8</sup>,  $Cp(CO)_3MSnAr^{Me6}$ , and metalloplumbylene<sup>9</sup>,  $Cp(CO)_3MSnAr^{iPr6}$  ( $M = Cr, Mo, \text{ or } W$ ;  $Ar^{Me6} = -C_6H_3-2,6-(C_6H_2-2,4,6-Me_3)_2$ ;  $Ar^{iPr6} = -C_6H_3-2,6-(C_6H_2-2,4,6-^iPr_3)_2$ ), analogues, and related species<sup>10</sup>. Attempts to generate the corresponding metallogermylenes led to the fortuitous discovery of a metallogermylyne featuring a molybdenum-germanium triple bond<sup>11</sup> via carbon monoxide elimination (Scheme 1). In contrast, the corresponding chemistry with the chromium or tungsten species afforded several examples of the desired singly bonded metallogermylenes.<sup>12</sup> Work by Filippou and coworkers disclosed other routes to the metallogermylynes<sup>13</sup> and metallogermylenes (Scheme 1) and expanded their results to include the corresponding complexes with triple bonds to tin<sup>14</sup> and lead.<sup>15</sup>

---

*Power et. al*<sup>11</sup>



*Filippou et. al*<sup>13</sup>



X = halide;  $Ar^{Me6} = -C_6H_3-2,6-(C_6H_2-2,4,6-Me_3)_2$   
 $Cp^* = C_5Me_5$ ;  $dppe = Ph_2PCH_2CH_2PPh_2$

---

### Scheme 1. Synthetic routes to metallogermylynes.

More recent work<sup>16</sup> has shown that the above species can be synthesized by metathesis

1  
2  
3  
4 of the dimetallynes REER (R = terphenyl; E = Ge,<sup>17</sup> Sn,<sup>18</sup> or Pb<sup>19</sup>) with the single or  
5  
6 triple bonded transition metal species, (CO)<sub>3</sub>CpMo-MoCp(CO)<sub>3</sub> or  
7  
8 (CO)<sub>2</sub>CpMo≡MoCp(CO)<sub>2</sub>.<sup>16</sup> In addition, the distannyne and digermynes were reacted  
9  
10 with the group 6 carbonyls M(CO)<sub>6</sub> to form the cluster species, {Ar<sup>iPr4</sup>EM(CO)<sub>4</sub>}<sub>2</sub> (E =  
11  
12 Ge, Sn; Ar<sup>iPr4</sup> = -C<sub>6</sub>H<sub>3</sub>-2,6-(C<sub>6</sub>H<sub>3</sub>-2,6-<sup>i</sup>Pr<sub>3</sub>)<sub>2</sub>).<sup>20</sup> However, the reactivity of diplumbynes  
13  
14 towards simple transition metal carbonyls remains hardly explored. Herein, we report  
15  
16 the syntheses and characterizations of metalloplumbylenes Mn(CO)<sub>5</sub>(PbAr<sup>iPr6</sup>) (**1**),  
17  
18 Fe(CO)<sub>4</sub>(PbAr<sup>iPr6</sup>)<sub>2</sub> (**2**), Co<sub>4</sub>(CO)<sub>9</sub>(PbAr<sup>iPr6</sup>)<sub>2</sub> (**3**) by reactions of the diplumbyne,  
19  
20 Ar<sup>iPr6</sup>PbPbAr<sup>iPr6</sup>, with the dinuclear metal carbonyls, Mn<sub>2</sub>(CO)<sub>10</sub>, Fe<sub>2</sub>(CO)<sub>9</sub>, Co<sub>2</sub>(CO)<sub>8</sub>.  
21  
22  
23  
24  
25  
26  
27  
28  
29

## 30 **Experimental Section**

31  
32 General Procedures. All operations were carried out by using modified Schlenk  
33  
34 techniques or in a Vacuum Atmospheres OMNI-Lab drybox under an atmosphere of  
35  
36 dry argon or nitrogen. The lead compounds were manipulated with careful exclusion of  
37  
38 light due to the tendency of low-valent lead compounds to decompose or  
39  
40 disproportionate under illumination. Solvents were dried over an alumina column and  
41  
42 degassed prior to use.<sup>21</sup> The compound Ar<sup>iPr6</sup>PbPbAr<sup>iPr6</sup> was prepared according to  
43  
44 literature procedures.<sup>19, 22</sup> Metal carbonyls were used as purchased without further  
45  
46 purification. Na[Mn(CO)<sub>5</sub>] was prepared by a modified literature procedure (see SI).<sup>23-</sup>  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
25 The <sup>1</sup>H, <sup>13</sup>C{<sup>1</sup>H}, <sup>59</sup>Co{<sup>1</sup>H}, and <sup>207</sup>Pb{<sup>1</sup>H} NMR spectra were recorded on a Varian  
The <sup>1</sup>H and <sup>13</sup>C{<sup>1</sup>H} NMR spectra were referenced to the  
residual solvent signals in C<sub>6</sub>D<sub>6</sub>. The <sup>59</sup>Co{<sup>1</sup>H} NMR spectrum was referenced to an

external standard of a saturated solution of  $K_3Co(CN)_6$  ( $\delta$  0.0 ppm) in  $D_2O$ . The  $^{207}Pb\{^1H\}$  NMR spectra were referenced to an external standard of  $PbMe_4$  ( $\delta$  0.0 ppm) in  $CDCl_3$ . UV-Visible spectra were recorded in dilute hexane solutions in 3.5 mL quartz cuvettes using an Olis 17 Modernized Cary 14 UV-Vis/NIR spectrophotometer. Infrared spectra were collected on a Bruker Tensor 27 ATR-FTIR spectrometer.

**$Mn(CO)_5(PbAr^{iPr6})$  (1).** Method A: ca. 50 mL of THF were added to a heavy-walled Teflon tapped Schlenk flask charged with 0.395 g (0.29 mmol)  $Ar^{iPr6}PbPbAr^{iPr6}$  and 0.122 g (0.31 mmol)  $Mn_2(CO)_{10}$ . The flask was then sealed, and the solution was stirred at ambient temperature for 2 days, and then heated to ca. 50 °C for 5 days, during which time the color changed from dark brown to dark green. The solution was cooled to room temperature, and filtered. The solution was then concentrated under reduced pressure to ca. 10 mL, and cooled to a ca. -18 °C for one week to afford the product as teal crystals of  $Mn(CO)_5(PbAr^{iPr6})$  (1). Yield: 0.098 g, 19 %.

Method B: 0.46 g (2.11 mmol)  $Na[Mn(CO)_5]$  in ca. 40 mL  $Et_2O$  solution was added dropwise over 15 minutes to a Schlenk flask charged with 1.472 g (0.96 mmol)  $[Ar^{iPr6}Pb(\mu-Br)]_2$  and ca. 30 mL of  $Et_2O$  at 0 °C. After stirring for 2 days, during which time the color changed from yellow to teal, solvent was removed and the residue was extracted with hexanes, and filtered. The hexanes were removed under reduced pressure and ca. 20 mL of  $Et_2O$  was added. Storage of the solution at room temperature overnight afforded teal crystals of  $Mn(CO)_5(PbAr^{iPr6})$  (1) that were suitable for X-ray diffraction studies. Yield: 1.392 g, 82 %.  $^1H$  NMR (600 MHz,  $C_6D_6$ , 298K, ppm):  $\delta$  = 1.09 (br,

1  
2  
3  
4 12H, p-CH(CH<sub>3</sub>)<sub>2</sub>), 1.19 (d, 12H, o-CH(CH<sub>3</sub>)<sub>2</sub>), <sup>3</sup>J<sub>H,H</sub> = 6.6 Hz, 1.38 (d, 12H, o-  
5  
6 CH(CH<sub>3</sub>)<sub>2</sub>), <sup>3</sup>J<sub>H,H</sub> = 6.6 Hz, 2.77 (sept, 2H, p-CH(CH<sub>3</sub>)<sub>2</sub>), <sup>3</sup>J<sub>H,H</sub> = 7.2 Hz, 3.40 (br, 4H,  
7  
8 o-CH(CH<sub>3</sub>)<sub>2</sub>) <sup>3</sup>J<sub>H,H</sub> = 5.8 Hz; 7.12 (br, 4H, m-Trip); 7.55 (t, 1H, p-C<sub>6</sub>H<sub>3</sub>) <sup>3</sup>J<sub>H,H</sub> = 7.3 Hz,  
9  
10 8.03 (d, 2H, m-C<sub>6</sub>H<sub>3</sub>) <sup>3</sup>J<sub>H,H</sub> = 7.3 Hz. <sup>13</sup>C{<sup>1</sup>H} NMR (151 MHz, C<sub>6</sub>D<sub>6</sub>, 298K, ppm): δ =  
11  
12 23.54 (o-CH(CH<sub>3</sub>)<sub>2</sub>), 23.95 (o-CH(CH<sub>3</sub>)<sub>2</sub>), 27.44 (p-CH(CH<sub>3</sub>)<sub>2</sub>), 30.90 (o-CH(CH<sub>3</sub>)<sub>2</sub>),  
13  
14 34.78 (p-CH(CH<sub>3</sub>)<sub>2</sub>), 122.49 (br, m-Trip), 125.30 (p-C<sub>6</sub>H<sub>3</sub>), 128.30 (m-C<sub>6</sub>H<sub>3</sub>), 133.02  
15  
16 (i-Trip), 140.54 (p-Trip), 146.38 (o-Trip), 149.17 (o-C<sub>6</sub>H<sub>3</sub>), 210.94 (br, CO), 282.93 (br,  
17  
18 CO). <sup>207</sup>Pb{<sup>1</sup>H} NMR (125 MHz, C<sub>6</sub>D<sub>6</sub>, 298K, ppm): δ = 8007.2. ATR-FTIR: ν<sub>CO</sub> (cm<sup>-1</sup>):  
19  
20 1958(s), 2060(m). UV-vis (hexane) λ<sub>max</sub> = nm (ε = M<sup>-1</sup>·cm<sup>-1</sup>): 332 (13700), 671 (1640).  
21  
22 **Fe(CO)<sub>4</sub>(PbAr<sup>iPr6</sup>)<sub>2</sub> (2)**. ca. 50 mL of Et<sub>2</sub>O was added to a Schlenk flask charged with  
23  
24 0.452 g (0.33 mmol) Ar<sup>iPr6</sup>PbPbAr<sup>iPr6</sup> and 0.132 g (0.36 mmol) Fe<sub>2</sub>(CO)<sub>9</sub>. The solution  
25  
26 was stirred for 2 days, during which time the color changed from dark brown to green.  
27  
28 The solution was filtered, concentrated under reduced pressure to ca. 10 mL, and cooled  
29  
30 to ca. -18 °C for one week to afford crystals of Fe(CO)<sub>4</sub>(PbAr<sup>iPr6</sup>)<sub>2</sub> (2) as green needles  
31  
32 that were suitable for X-ray diffraction studies. Yield: 0.295 g, 58 %. <sup>1</sup>H NMR (600  
33  
34 MHz, C<sub>6</sub>D<sub>6</sub>, 298K, ppm): δ = 1.13 (d, 12H, p-CH(CH<sub>3</sub>)<sub>2</sub>), <sup>3</sup>J<sub>H,H</sub> = 5.5 Hz. 1.28 (d, 12H,  
35  
36 o-CH(CH<sub>3</sub>)<sub>2</sub>), <sup>3</sup>J<sub>H,H</sub> = 6.2 Hz, 1.36 (d, 12H, o-CH(CH<sub>3</sub>)<sub>2</sub>), <sup>3</sup>J<sub>H,H</sub> = 7.0 Hz, 2.85 (sept, 2H,  
37  
38 p-CH(CH<sub>3</sub>)<sub>2</sub>), <sup>3</sup>J<sub>H,H</sub> = 6.6 Hz, 3.46 (broad, 4H, o-CH(CH<sub>3</sub>)<sub>2</sub>) <sup>3</sup>J<sub>H,H</sub> = 5.8 Hz; 7.12 (s, 4H,  
39  
40 m-Trip); 7.51 (t, 1H, p-C<sub>6</sub>H<sub>3</sub>) <sup>3</sup>J<sub>H,H</sub> = 7.1 Hz, 8.06 (d, 2H, m-C<sub>6</sub>H<sub>3</sub>) <sup>3</sup>J<sub>H,H</sub> = 7.9 Hz.  
41  
42 <sup>13</sup>C{<sup>1</sup>H} NMR (151 MHz, C<sub>6</sub>D<sub>6</sub>, 298K, ppm): δ = 23.46 (o-CH(CH<sub>3</sub>)<sub>2</sub>), 24.24 (o-  
43  
44 CH(CH<sub>3</sub>)<sub>2</sub>), 27.14 (p-CH(CH<sub>3</sub>)<sub>2</sub>), 30.66 (o-CH(CH<sub>3</sub>)<sub>2</sub>), 34.73 (p-CH(CH<sub>3</sub>)<sub>2</sub>), 120.73  
45  
46 (m-Trip), 124.99 (p-C<sub>6</sub>H<sub>3</sub>), 128.29 (m-C<sub>6</sub>H<sub>3</sub>), 133.60 (i-Trip), 141.21 (p-Trip), 145.95  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60


(o-Trip), 149.00 (o-C<sub>6</sub>H<sub>3</sub>), 295.48 (br, CO). <sup>207</sup>Pb{<sup>1</sup>H} NMR (125 MHz, C<sub>6</sub>D<sub>6</sub>, 298K, ppm): not observed. ATR-FTIR:  $\nu_{\text{CO}}$  (cm<sup>-1</sup>): 1889(m), 1958(m), 2005(m), 2041(w).

UV-vis (hexane)  $\lambda_{\text{max}}$  = nm ( $\epsilon = \text{M}^{-1}\cdot\text{cm}^{-1}$ ): 341 (22700), 505 (3140), 641 (3000).

**Co<sub>4</sub>(CO)<sub>9</sub>(PbAr<sup>iPr6</sup>)<sub>2</sub> (3)**. ca. 50 mL of hexane was added to a Schlenk flask charged with 0.330 g (0.24 mmol) Ar<sup>iPr6</sup>PbPbAr<sup>iPr6</sup> and 0.168 g (0.49 mmol) Co<sub>2</sub>(CO)<sub>8</sub>. After stirring overnight, during which time the color changed from dark brown to dark red, the solution was filtered, concentrated under reduced pressure to ca. 15 mL, and cooled to ca. -30 °C for one week to afford crystals of Co<sub>4</sub>(CO)<sub>9</sub>(PbAr<sup>iPr6</sup>)<sub>2</sub> (**3**) as dark maroon blocks that were suitable for X-ray diffraction studies. Yield: 0.299 g, 67 %. <sup>1</sup>H NMR (600 MHz, C<sub>6</sub>D<sub>6</sub>, 298K, ppm):  $\delta$  = 1.11 (d, 12H, p-CH(CH<sub>3</sub>)<sub>2</sub>), <sup>3</sup>J<sub>H,H</sub> = 6.6 Hz. 1.38 (d, 12H, o-CH(CH<sub>3</sub>)<sub>2</sub>), <sup>3</sup>J<sub>H,H</sub> = 6.9 Hz, 1.56 (d, 12H, o-CH(CH<sub>3</sub>)<sub>2</sub>), <sup>3</sup>J<sub>H,H</sub> = 6.9 Hz, 2.92 (sept, 2H, p-CH(CH<sub>3</sub>)<sub>2</sub>), <sup>3</sup>J<sub>H,H</sub> = 7.1 Hz, 3.16 (broad, 4H, o-CH(CH<sub>3</sub>)<sub>2</sub>) <sup>3</sup>J<sub>H,H</sub> = 7.4 Hz; 7.26 (s, 4H, m-Trip); 7.38 (t, 1H, p-C<sub>6</sub>H<sub>3</sub>) <sup>3</sup>J<sub>H,H</sub> = 5.9 Hz, 7.88 (d, 2H, m-C<sub>6</sub>H<sub>3</sub>) <sup>3</sup>J<sub>H,H</sub> = 7.3 Hz. <sup>13</sup>C{<sup>1</sup>H} NMR (151 MHz, C<sub>6</sub>D<sub>6</sub>, 298K, ppm):  $\delta$  = 23.47 (o-CH(CH<sub>3</sub>)<sub>2</sub>), 24.05 (o-CH(CH<sub>3</sub>)<sub>2</sub>), 26.14 (p-CH(CH<sub>3</sub>)<sub>2</sub>), 31.08 (o-CH(CH<sub>3</sub>)<sub>2</sub>), 34.92 (p-CH(CH<sub>3</sub>)<sub>2</sub>), 123.15 (m-Trip), 128.89 (p-C<sub>6</sub>H<sub>3</sub>), 134.74 (m-C<sub>6</sub>H<sub>3</sub>), 138.52 (i-Trip), 143.57 (p-Trip), 146.42 (o-Trip), 149.99 (o-C<sub>6</sub>H<sub>3</sub>), 213.17 (br, CO). <sup>59</sup>Co NMR (142 MHz, C<sub>6</sub>D<sub>6</sub>, 298K, ppm):  $\delta$  = -173.0 (Co(1), relative intensity = 1), -654.4 (Co(2) and Co(3), relative intensity = 2), -2016.0 (Co(4), relative intensity = 1). <sup>207</sup>Pb{<sup>1</sup>H} NMR (125 MHz, C<sub>6</sub>D<sub>6</sub>, 298K, ppm):  $\delta$  = 9010.4. ATR-FTIR:  $\nu_{\text{CO}}$  (cm<sup>-1</sup>): 1820(m), 1849(m), 1948(m), 1972(s), 2008(m), 2035(m). UV-vis (hexane)  $\lambda_{\text{max}}$  = nm ( $\epsilon = \text{M}^{-1}\cdot\text{cm}^{-1}$ ): 368 (shoulder, 57700).

## Results and Discussion

**Synthesis.** Metathetical exchange between group 14 dimetallynes,  $\text{Ar}^{\text{iPr}_4}\text{MMAr}^{\text{iPr}_4}$  or  $\text{Ar}^{\text{iPr}_6}\text{MMAr}^{\text{iPr}_6}$  ( $\text{M} = \text{Ge}, \text{Sn}, \text{or Pb}$ ) and  $(\text{CO})_3\text{CpMo-MoCp}(\text{CO})_3$  or  $(\text{CO})_2\text{CpMo}\equiv\text{MoCp}(\text{CO})_2$ , was shown earlier to afford the complexes,  $\text{Ar}^{\text{iPr}_4}\text{M}\equiv\text{MoCp}(\text{CO})_2$ ,  $\text{Ar}^{\text{iPr}_6}\text{M}\equiv\text{MoCp}(\text{CO})_2$ , and  $\text{Ar}^{\text{iPr}_4}\text{M-MoCp}(\text{CO})_3$  or  $\text{Ar}^{\text{iPr}_6}\text{M-MoCp}(\text{CO})_3$ .<sup>16</sup> In particular, the relatively weak Pb-Pb bond of the diplumbyne<sup>22</sup> is expected to facilitate these reactions with metal-metal (M-M) bonded transition metal carbonyl dimers, hence we studied the reactions of the nominally M-M bonded dinuclear metal carbonyls  $\text{Mn}_2(\text{CO})_{10}$ ,  $\text{Fe}_2(\text{CO})_9$ ,  $\text{Co}_2(\text{CO})_8$  with the diplumbyne,  $\text{Ar}^{\text{iPr}_6}\text{PbPbAr}^{\text{iPr}_6}$ . Compounds  $\text{Mn}(\text{CO})_5(\text{PbAr}^{\text{iPr}_6})$  (**1**),  $\text{Fe}(\text{CO})_4(\text{PbAr}^{\text{iPr}_6})_2$  (**2**),  $\text{Co}_4(\text{CO})_9(\text{PbAr}^{\text{iPr}_6})_2$  (**3**) were attained by the reaction of the diplumbyne with the dinuclear metal carbonyls either in ethereal solvents ( $\text{Et}_2\text{O}$  or THF) or in hexanes (Scheme 2). For compounds **2** and **3**, the syntheses were carried at ambient temperature, and the products were isolated in moderate to good yields. A similar approach to the



### Scheme 2. Reactions of $\text{Ar}^{\text{iPr}_6}\text{PbPbAr}^{\text{iPr}_6}$ with metal carbonyls.

synthesis of **1** afforded a color change from dark brown to green. However, the major signals in the  $^1\text{H}$  NMR spectrum of the residue after removal of solvent were attributed



1  
2  
3  
4 to the unreacted starting materials. To avoid the propensity for decomposition of the  
5  
6 diplumbyne under irradiation, gentle heating was applied to the reaction solution after  
7  
8 stirring at room temperature for two days. This approach afforded **1** in low (ca. 20%)  
9  
10 yield. In contrast, an alternative approach to the synthesis of **1** via salt metathesis of  
11  
12  $[\text{Ar}^{\text{iPr}_6}\text{Pb}(\mu\text{-Br})_2]$  and  $\text{Na}[\text{Mn}(\text{CO})_5]$  afforded **1** in a much higher (82%) yield.  
13  
14

15  
16 The synthesis of **1** by reaction of  $\text{Ar}^{\text{iPr}_6}\text{PbPbAr}^{\text{iPr}_6}$  and  $\text{Mn}_2(\text{CO})_{10}$  is formally a  
17  
18 metathetical exchange between the  $\text{Pb}\equiv\text{Pb}$  'triple' bond and the Mn-Mn single bond.  
19  
20 However, for steric reasons it seems unlikely that this reaction proceeds via intact  
21  
22 dinuclear metal carbonyls. In the cases of **1-3**, it is proposed that the diplumbyne,  
23  
24  $\text{Ar}^{\text{iPr}_6}\text{PbPbAr}^{\text{iPr}_6}$ , dissociates into  $\text{Ar}^{\text{iPr}_6}\text{Pb}\cdot$  radical fragment<sup>16, 26</sup>, followed by  
25  
26 subsequent reaction with  $\text{Mn}_2(\text{CO})_{10}$ ,  $\text{Fe}_2(\text{CO})_9$ , or  $\text{Co}_2(\text{CO})_8$ .  
27  
28  
29  
30  
31


32  
33 The yields of the reactions between diplumbyne and metal carbonyls are possibly  
34  
35 affected by the transition metal-metal (M-M) bond strength. Recently, Shaik and  
36  
37 coworkers presented computational studies on the M-M bonds in transition metal  
38  
39 complexes using of ab initio valence-bond methods, in which the M-M bonds of the  
40  
41 3d-series (group 3-10) are considered as pure charge shift bonds.<sup>27</sup> The Mn-Mn bond  
42  
43 in  $\text{Mn}_2(\text{CO})_{10}$  is calculated to have a bond dissociation energy of 22.9 kcal/mol with a  
44  
45 charge-shift resonance energy of 24.5 kcal/mol (experimental  $38.0 \pm 5$  kcal/mol)<sup>28</sup>  
46  
47 suggesting that the Mn-Mn bond is the strongest M-M bond among  $\text{Mn}_2(\text{CO})_{10}$ ,<sup>27</sup>  
48  
49  $\text{Fe}_2(\text{CO})_9$  (no Fe-Fe bond),<sup>29</sup> and  $\text{Co}_2(\text{CO})_8$  (9.8 kcal/mol),<sup>27</sup> which may account for the  
50  
51 relatively low yield of **1** in the reaction between  $\text{Ar}^{\text{iPr}_6}\text{PbPbAr}^{\text{iPr}_6}$  and  $\text{Mn}_2(\text{CO})_{10}$ .  
52  
53  
54  
55  
56  
57

58 Both the teal crystals of **1**, and green crystals of **2** are highly air-sensitive, and an  
59  
60

1  
2  
3  
4 immediate color change is observed upon their exposure to air. Dark maroon crystals  
5  
6 of **3** can be briefly handled in air without displaying significant color change.  
7  
8  
9 Hydrocarbon solutions of **1**, **2**, and **3** were prone to decomposition by light over time,  
10  
11 with lead metal deposition and a lightening of the color of the solution.  
12  
13  
14  
15  
16


17 **Structures.** The solid-state molecular structures of compounds **1**, **2**, and **3** were  
18  
19 determined by single crystal X-ray crystallography. Selected bond distances and bond  
20  
21 angles are given in Table 1, and the molecular structures of **1**, **2**, and **3** are illustrated in  
22  
23 Figures 1-3, respectively.  
24  
25  
26

27 Compound **1** crystallized in the I2/a space group. The manganese atom in compound **1**  
28  
29 is six-coordinate with a near-octahedral coordination geometry. The Pb- Mn distance is  
30  
31 2.838(4) Å, which is slightly longer than the Pb-Mn single bonds in  
32  
33 [Pb{Mn(CO)<sub>5</sub>}<sub>3</sub>][AlCl<sub>4</sub>]<sup>30</sup> (2.750(1)-2.785(1) Å), and the Pb-Mn single bond in the  
34  
35 dimetalloborylene complex, (Ph<sub>3</sub>Pb)[Mn(η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)(CO)<sub>2</sub>]<sub>2</sub>B, 2.703(7) Å.<sup>31</sup> The Pb-Mn  
36  
37 bond distance in **1** is slightly shorter than the sum (2.85 Å) of the covalent radii of  
38  
39 Mn(1.39 Å) and Pb(1.46 Å).<sup>32-34</sup> The interligand angles at the Mn atom range from  
40  
41 82.65(8)° to 96.75(11)° which deviate somewhat (<7.5°) from the 90° value expected  
42  
43 for idealized octahedral coordination. The C(1)-Pb(1)-Mn(1) angle at the lead atom is  
44  
45 106.45(5)°.  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



**Figure 1.** Thermal ellipsoid (50%) plot of **1**. H atoms and disordered groups are not shown for clarity, the organic substituents are shown as wire frames.

The molecular structure of **2** showed that it has a six-coordinated iron atom with four carbonyl ligands and two mutually *cis*-PbAr<sup>iPr6</sup> moieties, which afford distorted octahedral coordination at iron. The interligand angles at the iron atom range from 75.0(5)° to 106.5(7)°. The deviation of interligand angles from ideal octahedral geometry at iron is possibly due to the steric hindrance between the two *cis*-PbAr<sup>iPr6</sup> moieties. The two Pb-Fe bond distances are 2.830(3) Å, and 2.804(3) Å, which are slightly longer than the sum (2.78 Å) of the covalent radii of Pb(1.46 Å) and Fe(1.32 Å).<sup>32-34</sup> The Pb-Fe bond lengths in **2** are slightly longer than the Pb-Fe distances, (2.736(2) Å and 2.728(2) Å), observed in [Et<sub>2</sub>PbFe(CO)<sub>4</sub>]<sub>2</sub> by Wrackmeyer and coworkers.<sup>35</sup> The relatively long Fe-Pb and Mn-Pb distances in **1** and **2** are consistent




**Figure 2.** Thermal ellipsoid (50%) plot of **2**. H atoms and disordered groups are not shown for clarity, the organic substituents are shown as wire frames.

with a +2 oxidation state of lead as the  $\sigma$ -bonds are likely to be dominated by the p-orbital character from the Pb(II) atom, assuming that the transition metal carbonyl fragment carries a negative charge.<sup>9</sup> The interligand angles at the lead atom(s) of **1** ( $106.45(5)^\circ$ ) and **2** ( $100.50(3)^\circ$ ,  $102.70(3)^\circ$ ) are slightly narrower than those in the  $\text{Cp}(\text{CO})_3\text{M}-\text{PbAr}^{\text{iPr}_6}$  ( $\text{M}=\text{Cr}, \text{Mo}, \text{W}$ ) complexes ( $108.6(2)^\circ$ - $113.58(9)^\circ$ ),<sup>9</sup> possibly due to the steric differences between carbonyl groups and cyclopentadienyl group. The structures of **1** and **2** can also be rationalized by the 18-electron rule where the  $\text{PbAr}^{\text{iPr}_6}$  moieties are 1-electron ligands bound to the transition metals.<sup>9</sup>

Compound **3** crystallized as dark maroon blocks in the  $\text{P2}_1/\text{c}$  space group. The molecular structure of **3** displays a near-tetrahedral  $\text{Co}_4$  core bonded to a total of nine

1  
2  
3  
4 carbonyls and two  $\text{PbAr}^{\text{iPr}_6}$  moieties. Seven of the carbonyls are terminally bound, with  
5  
6 the remaining two carbonyls and one of the  $\text{PbAr}^{\text{iPr}_6}$  moieties each bridging one of the  
7  
8 three edges linking the apical cobalt atom with the three basal cobalt atoms. The  
9  
10 remaining  $\text{PbAr}^{\text{iPr}_6}$  moiety bridges the face formed from the three basal cobalt atoms.  
11  
12 The apical cobalt atom carries only one terminal carbonyl, while each of the basal cobalt  
13  
14 atoms carry two terminal carbonyls. The structure of **3** therefore differs from that of  
15  
16  $\text{Co}_4(\text{CO})_{12}$ , where the apical atom is connected to three terminal carbonyls. The  
17  
18 structure of compound **3** conforms to Wade's rules which predict a *nido* structure for  
19  
20 the tetracobalt tetrahedron core as it has  $n+2$  ( $n$ =no. of vertices) skeletal electron pairs  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



**Figure 3.** Thermal ellipsoid (50%) plot of **3**. H atoms and disordered groups are not shown for clarity, the organic substituents are shown as wire frames.

for cluster bonding.<sup>36-37</sup> The edge-bridging Pb-Co bonds are 2.5008(7) Å and 2.5150(6) Å, while the average of the face-bridging Pb-Co distances is 2.5750(8) Å, which is

**Table 1.** Selected bond distances (Å) and angles (°) for **1-3**.

	<b>1</b> (M = Mn)	<b>2</b> (M = Fe)	<b>3</b> (M = Co)
Pb-C	2.288(2)	Pb(1)-C(1) 2.313(13) Pb(2)-C(37) 2.307(13)	Pb(1)-C(1) 2.229(2) Pb(2)-C(37) 2.214(2)
Pb-M	2.8376(4)	Pb(1)-Fe(1) 2.800(2) Pb(2)-Fe(1) 2.829(2)	2.5078(4) (average of edge-bridging) 2.5735(4) (average of face-bridging)
M-CO (average)	1.839(3)	1.772(19)	1.775(3) (terminal) 1.932(3) (bridging)
M-Pb-C	106.45(5)	C(1)-Pb(1)-Fe(1) 102.70(3)  C(37)-Pb(2)-Fe(1) 100.50(3)	C(37)-Pb(2)-Co(4) 130.44(7) C(37)-Pb(2)-Co(1) 162.43(7) (edge-bridging)  C(1)-Pb(1)-Co(1) 128.18(6) C(1)-Pb(1)-Co(2) 153.22(6) C(1)-Pb(1)-Co(3) 146.90(6) (face-bridging)

slightly longer than the edge-bridging distance as expected for the higher coordination number. The Co-Co lengths are similar to those of the Co-Co bonds (2.441(14) Å to 2.527(10) Å) in cobalt carbonyl clusters<sup>38</sup> except that the Co(1)-Co(4) distance bridged by the PbAr<sup>iPr</sup><sub>6</sub> moiety which is 2.7472(9) Å, much longer than an average Co-Co bond, 2.499(1) Å.<sup>39-41</sup> The calculated structure of the unsaturated compound Co<sub>4</sub>(CO)<sub>11</sub> reported by King and coworkers<sup>42</sup> was predicted to have a μ<sub>4</sub>-CO group bridging all four cobalt atoms, where the structure of **3** resembles one of the predicted structures with a butterfly array of cobalt atoms. The calculated structure indicated a long Co...Co distance between the two wingtips cobalt atoms of 2.974 Å (B3LYP) or 3.105 Å (BP86), in the Co<sub>4</sub> butterfly, 0.2~0.3 Å longer than Co(1)-Co(4) in the structure of **3**.<sup>42</sup> The structure of **3** displays wide interligand angles at the lead atoms (see Table 1) thus arises from the bridging nature of the PbAr<sup>iPr</sup><sub>6</sub> moieties and the higher coordination number

1  
2  
3  
4 at lead atom. In addition, the sum of the angles at Pb(2) is 359.26(7)°, consistent with  
5  
6 a planar coordination geometry. The Pb-C distances in **1-3** range from 2.214(2) Å to  
7  
8 2.313(13) Å which are within the range of reported divalent aryllead species.<sup>43-45</sup> The  
9  
10 transition metal-carbon distances to the carbonyl groups are comparable to their parent  
11  
12 metal carbonyls.<sup>46</sup>

13  
14  
15  
16 **Spectroscopy.** <sup>1</sup>H and <sup>13</sup>C{<sup>1</sup>H} NMR spectra of **1**, **2**, and **3** indicate the high purity of  
17  
18 the samples. The <sup>1</sup>H NMR spectra of **1** and **2** each displayed a set of signals which  
19  
20 correspond to the ligand, Ar<sup>iPr6</sup>.<sup>45</sup> Peak broadenings were observed in the <sup>1</sup>H NMR  
21  
22 spectrum of **1** possibly due to restricted rotations of the flanking rings, which has been  
23  
24 noted previously.<sup>22</sup> The <sup>1</sup>H NMR spectrum of **3** displayed two sets of peaks attributable  
25  
26 to the ligand which differ by approximately 0.01 ppm due to the two slightly different  
27  
28 environments of the face-bridging and edge-bridging lead atoms.

29  
30  
31  
32  
33  
34  
35 The <sup>13</sup>C{<sup>1</sup>H} NMR spectra of **1**, **2**, and **3** displayed signals similar to those in the  
36  
37 previously reported complexes, (η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)(CO)<sub>3</sub>M-PbAr<sup>iPr6</sup> (M=Cr, Mo, W), including  
38  
39 **3** where there was only one set of signals from the ligand, Ar<sup>iPr6</sup>. The <sup>13</sup>C{<sup>1</sup>H} NMR  
40  
41 spectrum of **1** displayed two peaks attributable to the carbonyl groups at 210.94 ppm,  
42  
43 and 282.93 ppm (Table 2) in an approximate integration ratio of 4:1, which correspond  
44  
45 to the COs cis and trans to the PbAr<sup>iPr6</sup> substituents. There is no indication of restricted  
46  
47 rotation around the Pb(1)-Mn(1) axis in the <sup>13</sup>C{<sup>1</sup>H} NMR spectrum, consistent with a  
48  
49 single bond between Pb and Mn. The <sup>13</sup>C{<sup>1</sup>H} NMR spectra of **2** and **3** each showed a  
50  
51 single broad signal at 295.48 ppm, and 213.17 ppm for the carbonyl groups,  
52  
53 respectively. (see Table 2) The <sup>13</sup>C{<sup>1</sup>H} NMR spectrum of **2** can be understood by  
54  
55  
56  
57  
58  
59  
60


1  
2  
3  
4 having an overall  $C_{2v}$  symmetry for its solid-state structure. It is unclear of missing  
5  
6 signals in the  $^{13}C\{^1H\}$  NMR spectrum of **3**, where it has been previously reported of  
7  
8 missing apical  $^{13}C$  signal in the  $^{13}C\{^1H\}$  NMR spectrum of  $Co_4(CO)_{12}$ .<sup>47</sup>

9  
10  
11 The  $^{207}Pb\{^1H\}$  NMR spectra of **1** and **3** displayed broad signals at 8007.2 ppm, and  
12  
13 9010.4 ppm ( Table 2) which are in fair agreement with  $^{207}Pb$  NMR signals observed  
14  
15 for  $(\eta^5-C_5H_5)(CO)_3M-PbAr^{iPr_6}$  (M=Cr, Mo, W), 9374-9659 ppm,<sup>9</sup> are also the chemical  
16  
17 shift range for two-coordinated lead.<sup>44</sup> Signal broadening in  $^{207}Pb\{^1H\}$  NMR spectrum  
18  
19 of **3** could be caused by coupling to multiple  $^{59}Co$  atoms as  $^{59}Co$  is a 7/2 spin nucleus  
20  
21 and thus quadrupolar.<sup>48</sup> The  $^{59}Co\{^1H\}$  NMR spectrum of **3** displayed three signals at -  
22  
23 173.0 ppm, -654.4 ppm, and -2016.0 ppm, respectively. The intensity ratio of the three  
24  
25 signals is 1:2:1 which corresponds to the three different chemical environments for the  
26  
27 cobalt atoms seen in the crystal structure of **3**. The chemical shifts in the  $^{59}Co$  NMR  
28  
29 spectrum of tetrahedral clusters supported by carbonyl ligands are generally unaffected  
30  
31 by edge-bridging or face-bridging metals.<sup>48</sup> The signal at -173.0 ppm can be assigned  
32  
33 to Co(1), the more intense signal at -654.4 ppm can be assigned to both Co(2) and Co(3)  
34  
35 where Co(1), Co(2) and Co(3) are all bonded to two terminal carbonyl ligands in its  
36  
37 solid state, and the signal at -2016.0 ppm to Co(4), which is bonded to two bridged  
38  
39 carbonyl ligands and one terminal carbonyl ligand, respectively, suggesting that solid-  
40  
41 state structure of **3** is also present in the solution structure. The chemical shifts of the  
42  
43 signals are close to those chemical shifts reported for  $Co_4(CO)_{12}$ , -668 ppm (apical Co),  
44  
45 and -2032 ppm (basal Co) in  $CDCl_3$ .<sup>48</sup>

46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58 The UV-vis spectra of **1** and **2** showed absorptions at 671 nm and 641 nm (Table 2),  
59  
60



1  
2  
3  
4 respectively, which are attributed to the n-p transitions of the lead atoms as depicted in  
5  
6 DFT calculations (Figure 4). These absorptions are close to the reported values for ( $\eta^5$ -  
7  
8  $C_5H_5$ )(CO)<sub>3</sub>M-PbAr<sup>iPr</sup><sub>6</sub> 611-624 nm.<sup>9</sup> However, the UV-vis spectrum of **3** displayed an  
9  
10 absorption at 368 nm that is blue-shifted compared to absorptions of plumbylenes,  
11  
12 possibly due to the large extinction coefficient of cobalt-centered electronic transition  
13  
14 where the absorption is very similar to what was reported for Co<sub>4</sub>(CO)<sub>12</sub> (372 nm).<sup>49</sup>  
15  
16  
17  
18  
19  
20



21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
**Figure 4.** Depiction of calculated HOMO (1a) and LUMO (1b) of **1**, calculated HOMO (2a) and LUMO (2b) of **2**. HOMO-LUMO gap of **1** is calculated at 340 kJ/mol, and 325 kJ/mol for **2**, hydrogens are omitted for clarity and the isovalue is set at 0.05.

Compounds **1-3** were also characterized by IR spectroscopy. The IR spectrum of **1**

1  
2  
3  
4 displayed a similar pattern to that of the corresponding complex,  $(\text{Ph}_3\text{PAu})\text{Mn}(\text{CO})_5$   
5  
6  $(1961(\text{vs}) \text{ cm}^{-1}, 2062(\text{s}) \text{ cm}^{-1})$ .<sup>50</sup> The IR spectrum of **1** showed two vibrational modes  
7  
8 of the carbonyl groups,  $A_1$   $(1958(\text{s}) \text{ cm}^{-1})$ , and  $E$   $(2060(\text{m}) \text{ cm}^{-1})$ , which correspond to  
9  
10 axial and equatorial CO stretching modes, respectively. The 4-band  $(2A_1+B_1+B_2)$   
11  
12 pattern in the IR spectrum of **2** is similar to what was reported for  
13  
14  $(\text{NHC})\text{Al}(\text{Br})[(\text{Fe}(\text{CO})_4)]$ <sup>51</sup> (NHC = N-heterocyclic carbene) and  $\text{cis-Fe}(\text{CO})_4(\text{SnR}_3)_2$ <sup>52</sup>.  
15  
16 CO stretching frequencies of **1** were calculated to be  $1974 \text{ cm}^{-1}$  for axial,  $1975 \text{ cm}^{-1}$  for  
17  
18 equatorial, and  $2055 \text{ cm}^{-1}$  for axial + equatorial. CO stretching frequencies of **2** were  
19  
20 calculated to be  $1893 \text{ cm}^{-1}$ ,  $1921 \text{ cm}^{-1}$ ,  $1967 \text{ cm}^{-1}$ , and  $2008 \text{ cm}^{-1}$ . The IR spectrum of  
21  
22 **3** showed similarities with the calculated (BP86) vibrational frequencies of the  
23  
24 theoretical

25  
26  
27  
28  
29  
30  
31  
32  
33 **Table 2.** Selected spectroscopic data for **1-3**.

	<b>1</b> (M = Mn)	<b>2</b> (M = Fe)	<b>3</b> (M = Co)
UV-vis (nm) <sup>a</sup>	332, 671	341, 505, 641	368
IR( $\tilde{\nu}_{\text{CO}}$ ) $\text{cm}^{-1}$	1958(s), 2060(m)	1889(m), 1958(m), 2005(m), 2041(w)	1820(m), 1849(m), 1948(m), 1972(s), 2008(m), 2035(m).
<sup>13</sup> C NMR (CO) (ppm) <sup>b</sup>	210.94, 282.93	295.48	213.17
<sup>207</sup> Pb NMR (ppm) <sup>c</sup>	8007.2	-	9010.4

34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60 a. UV-vis spectra were collected in hexanes at 298K. b. <sup>13</sup>C{<sup>1</sup>H} NMR spectra were

1  
2  
3  
4 collected in C<sub>6</sub>D<sub>6</sub> at 298K. c. <sup>207</sup>Pb{<sup>1</sup>H} NMR spectra were collected in C<sub>6</sub>D<sub>6</sub> at 298K.  
5  
6 butterfly complex Co<sub>4</sub>(CO)<sub>11</sub>, where the vibrational frequencies have relatively low  
7  
8 wavenumber values of 1820 cm<sup>-1</sup> and 1849 cm<sup>-1</sup>, which can be attributed to the bridging  
9  
10 carbonyl groups, and that 2008(m) cm<sup>-1</sup> and 2035(m) cm<sup>-1</sup> to the terminal carbonyl  
11  
12 groups(2020(s), 2026(s), 2032(s)).<sup>42</sup>  
13  
14  
15

## 16 **Conclusion**

17  
18  
19 In conclusion, the syntheses and characterizations of complexes Mn(CO)<sub>5</sub>(PbAr<sup>iPr6</sup>) (1),  
20  
21 Fe(CO)<sub>4</sub>(PbAr<sup>iPr6</sup>)<sub>2</sub> (2), Co<sub>4</sub>(CO)<sub>9</sub>(PbAr<sup>iPr6</sup>)<sub>2</sub> (3), have been described. They are rare  
22  
23 examples of complexes synthesized by interaction of group 14 dimetallyne and  
24  
25 dinuclear transition metal carbonyls. The reactivities of diplumbyne and dinuclear  
26  
27 metal carbonyls can be attributed to the charge-shift bond character of Pb≡Pb triple  
28  
29 bond and M-M single bond. The syntheses of **1-3** are postulated to proceed via the  
30  
31 reactions of Ar<sup>iPr6</sup>Pb• radical fragment from the dissociation of Ar<sup>iPr6</sup>PbPbAr<sup>iPr6</sup> with  
32  
33 Mn<sub>2</sub>(CO)<sub>10</sub>, Fe<sub>2</sub>(CO)<sub>9</sub>, or Co<sub>2</sub>(CO)<sub>8</sub>. Further studies of these synthesized  
34  
35 metalloplumbylenes with small molecules are currently underway.  
36  
37  
38  
39  
40  
41  
42  
43  
44

## 45 **ASSOCIATED CONTENT**

### 46 Supporting Information

47  
48 The Supporting Information is available free of charge on the ACS Publications website  
49  
50 at <https://pubs.acs.org/doi/10.1021/acs.organomet.xxxxxxx>.  
51  
52

53  
54 Synthetic details for Na[Mn(CO)<sub>5</sub>], Computational details, NMR, IR and electronic  
55  
56 spectra of the described compounds (PDF)  
57  
58  
59  
60

1  
2  
3  
4 Crystal data for CCDC 2018923 (CIF)

5  
6 Crystal data for CCDC 2018925 (CIF)

7  
8  
9 Crystal data for CCDC 2027035 (CIF)

10  
11  
12 Optimized Cartesian coordinates (XYZ)

### 13 14 **Accession Codes**

15  
16  
17 CCDC 2018923, 2018925, 2027035 contain the supplementary crystallographic data  
18  
19 for this paper. These data can be obtained free of charge via  
20  
21 [www.ccdc.cam.ac.uk/data\\_request/cif](http://www.ccdc.cam.ac.uk/data_request/cif), or by emailing [data\\_request@ccdc.cam.ac.uk](mailto:data_request@ccdc.cam.ac.uk),  
22  
23  
24 or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road,  
25  
26  
27 Cambridge CB2 1EZ, UK; fax: +44 1223 336033.

### 28 29 30 **AUTHOR INFORMATION**

#### 31 32 **Corresponding Author**

33 **Philip P. Power** – *Department of Chemistry, University of California, Davis,*  
34 *California 95616, United States;*  
35 *orcid.org/0000-0002-6262-3209; Email: [pppower@ucdavis.edu](mailto:pppower@ucdavis.edu)*

#### 36 37 38 **Authors**

39 **Qihao Zhu** – *Department of Chemistry, University of California, Davis, California*  
40 *95616, United States;*

41 *orcid.org/0000-0002-5566-4491*

42 **Petra Vasko** – *Department of Chemistry, Nanoscience Center, University of Jyväskylä,*  
43 *P.O. Box 35, FI-40014 University of Jyväskylä, Finland;*

44 *orcid.org/0000-0003-4202-6869*

45 **James C. Fettinger** – *Department of Chemistry, University of California, Davis,*  
46 *California 95616, United States;*

47 *orcid.org/0000-0002-6428-4909*

#### 48 49 50 51 52 **Notes**

53 The authors declare no competing financial interest.

### 54 55 56 **ACKNOWLEDGMENTS**

57 We are grateful to the National Science Foundation (NSF) for financial support (Grants  
58 CHE-1565501) and for the dual-source X-ray diffractometer (Grant 0840444). PV  
59  
60

would like to thank the Academy of Finland (project number 314794) and Prof. Heikki M. Tuononen for providing computational resources.

## References

1. Fischer, E. O., On the Way to Carbene and Carbyne Complexes. *Advances in Organometallic Chemistry*, 14; Elsevier: Burlington, 1976; pp 1-32.
2. Marks, T. J., Dialkylgermylene- and -stannylene-pentacarbonylchromium complexes. *J. Am. Chem. Soc.* **1971**, *93*, 7090-7091.
3. Cotton, J. D.; Davison, P. J.; Goldberg, D. E.; Lappert, M. F.; Thomas, K. M., Co-ordination chemistry of heavy-atom group IV donors, and the crystal and molecular structure of  $[(\text{Me}_3\text{Si})_2\text{CH}]_2\text{SnCr}(\text{CO})_5$ . *J. Chem. Soc., Chem. Commun.* **1974**, 893-895.
4. Petz, W., Transition-metal complexes with derivatives of divalent silicon, germanium, tin, and lead as ligands. *Chem. Rev.* **1986**, *86*, 1019-1047.
5. Hashimoto, H.; Tobita, H., Recent advances in the chemistry of transition metal–silicon/germanium triple-bonded complexes. *Coord. Chem. Rev.* **2018**, *355*, 362-379.
6. Álvarez-Rodríguez, L.; Cabeza, J. A.; García-Álvarez, P.; Polo, D., The transition-metal chemistry of amidinatosilylenes, -germylenes and -stannylenes. *Coord. Chem. Rev.* **2015**, *300*, 1-28.
7. Jutzi, P.; Leue, C., (Supermesityl)chlorogermylene (Supermesityl =  $\text{Mes}^* = 2,4,6\text{-tBu}_3\text{C}_6\text{H}_2$ ): Synthesis and Derivatization to (Supermesityl)ferriogermynes. *Organometallics* **1994**, *13*, 2898-2899.
8. Eichler, B. E.; Phillips, A. D.; Haubrich, S. T.; Mork, B. V.; Power, P. P., Synthesis, Structures, and Spectroscopy of the Metallostannylenes  $(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_3\text{M-Sn-C}_6\text{H}_3\text{-2,6-Ar}_2$  (M = Cr, Mo, W; Ar =  $\text{C}_6\text{H}_2\text{-2,4,6-Me}_3$ ,  $\text{C}_6\text{H}_2\text{-2,4,6-Pr}_3$ ). *Organometallics* **2002**, *21*, 5622-5627.
9. Pu, L.; Power, P. P.; Boltz, I.; Herbst-Irmer, R., Synthesis and Characterization of the Metalloplumbylenes  $(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_3\text{M-Pb-C}_6\text{H}_3\text{-2,6-Trip}_2$  (M = Cr, Mo, or W; Trip =  $-\text{C}_6\text{H}_2\text{-2,4,6-i-Pr}_3$ ). *Organometallics* **2000**, *19*, 352-356.
10. Lei, H.; Guo, J.-D.; Fettinger, J. C.; Nagase, S.; Power, P. P., Synthesis, Characterization, and CO Elimination of Ferrio-Substituted Two-Coordinate Germylenes and Stannylenes. *Organometallics* **2011**, *30*, 6316-6322.
11. Simons, R. S.; Power, P. P.,  $(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2\text{MoGeC}_6\text{H}_3\text{-2,6-Mes}_2$ : A Transition-Metal Germylyne Complex. *J. Am. Chem. Soc.* **1996**, *118*, 11966-11967.
12. Pu, L.; Twamley, B.; Haubrich, S. T.; Olmstead, M. M.; Mork, B. V.; Simons, R. S.; Power, P. P., Triple Bonding to Germanium: Characterization of the Transition Metal Germylynes  $(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2\text{M}:\text{Ge-C}_6\text{H}_3\text{-2,6-Mes}_2$  (M = Mo, W; Mes =  $-\text{C}_6\text{H}_2\text{-2,4,6-Me}_3$ ) and  $(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2\text{M}:\text{Ge-C}_6\text{H}_3\text{-2,6-Trip}_2$  (M = Cr, Mo, W; Trip =  $-\text{C}_6\text{H}_2\text{-2,4,6-i-Pr}_3$ ) and the Related Single Bonded Metallogermylenes  $(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_3\text{M-Ge-C}_6\text{H}_3\text{-2,6-Trip}_2$  (M = Cr, W). *J. Am. Chem. Soc.* **2000**, *122*, 650-656.
13. Filippou, A. C.; Philippopoulos, A. I.; Portius, P.; Neumann, D. U., Synthesis and Structure of the Germylyne Complexes  $\text{trans-[X(dppe)}_2\text{W}\equiv\text{Ge}(\eta^1\text{-Cp}^*)]$  (X=Cl, Br, I) and Comparison of the  $\text{W}\equiv\text{E}$  Bonds (E=C, Ge) by Density Functional Calculations. *Angew. Chem. Int. Ed.* **2000**, *39*, 2778-2781.
14. Filippou, A. C.; Portius, P.; Philippopoulos, A. I.; Rohde, H., Triple Bonding to Tin: Synthesis and Characterization of the Stannylyne Complex  $\text{trans-[Cl(PMe}_3)_4\text{W}\equiv\text{Sn-C}_6\text{H}_3\text{-2,6-Mes}_2]$ . *Angew. Chem. Int. Ed.* **2003**, *42*, 445-447.
15. Filippou, A. C.; Weidemann, N.; Schnakenburg, G.; Rohde, H.; Philippopoulos, A. I., Tungsten–

Lead Triple Bonds: Syntheses, Structures, and Coordination Chemistry of the Plumbidyne Complexes  $\text{trans-[X(PMe}_3)_4\text{W}\equiv\text{Pb(2,6-Trip}_2\text{C}_6\text{H}_3)]$ . *Angew. Chem. Int. Ed.* **2004**, *43*, 6512-6516.

16. Queen, J. D.; Phung, A. C.; Caputo, C. A.; Fettinger, J. C.; Power, P. P., Metathetical Exchange between Metal-Metal Triple Bonds. *J. Am. Chem. Soc.* **2020**, *142*, 2233-2237.

17. Stender, M.; Phillips, A. D.; Wright, R. J.; Power, P. P., Synthesis and Characterization of a Digermanium Analogue of an Alkyne. *Angew. Chem. Int. Ed.* **2002**, *41*, 1785-1787.

18. Phillips, A. D.; Wright, R. J.; Olmstead, M. M.; Power, P. P., Synthesis and Characterization of 2,6-Dipp<sub>2</sub>-H<sub>3</sub>C<sub>6</sub>SnC<sub>6</sub>H<sub>3</sub>-2,6-Dipp<sub>2</sub> (Dipp = C<sub>6</sub>H<sub>3</sub>-2,6-Pr<sub>2</sub>): A Tin Analogue of an Alkyne. *J. Am. Chem. Soc.* **2002**, *124*, 5930-5931.

19. Pu, L.; Twamley, B.; Power, P. P., Synthesis and Characterization of 2,6-Trip<sub>2</sub>H<sub>3</sub>C<sub>6</sub>PbPbC<sub>6</sub>H<sub>3</sub>-2,6-Trip<sub>2</sub> (Trip = C<sub>6</sub>H<sub>2</sub>-2,4,6-i-Pr<sub>3</sub>): A Stable Heavier Group 14 Element Analogue of an Alkyne. *J. Am. Chem. Soc.* **2000**, *122*, 3524-3525.

20. McCrea-Hendrick, M. L.; Caputo, C. A.; Linnera, J.; Vasko, P.; Weinstein, C. M.; Fettinger, J. C.; Tuononen, H. M.; Power, P. P., Cleavage of Ge-Ge and Sn-Sn Triple Bonds in Heavy Group 14 Element Alkyne Analogues (EAr<sup>i</sup>Pr<sub>4</sub>)<sub>2</sub> (E = Ge, Sn; Ar<sup>i</sup>Pr<sub>4</sub> = C<sub>6</sub>H<sub>3</sub>-2,6(C<sub>6</sub>H<sub>3</sub>-2,6-Pr<sub>2</sub>)<sub>2</sub>) by Reaction with Group 6 Carbonyls. *Organometallics* **2016**, *35*, 2759-2767.

21. Pangborn, A. B.; Giardello, M. A.; Grubbs, R. H.; Rosen, R. K.; Timmers, F. J., Safe and Convenient Procedure for Solvent Purification. *Organometallics* **1996**, *15*, 1518-1520.

22. Queen, J. D.; Bursch, M.; Seibert, J.; Maurer, L. R.; Ellis, B. D.; Fettinger, J. C.; Grimme, S.; Power, P. P., Isolation and Computational Studies of a Series of Terphenyl Substituted Diplumbynes with Ligand Dependent Lead-Lead Multiple-Bonding Character. *J. Am. Chem. Soc.* **2019**, *141*, 14370-14383.

23. King R. B., Stone F. G. A., Sodium Salts of Carbonyl Hydrides Prepared in Ethereal Media. In *Inorg. Synth.*; McGraw-Hill book: New-York, 1963; Vol. 7, pp 196-201.

24. Reimer, K. J., Shaver, A., Quick, M.H. and Angelici, R.J., Pentacarbonylmanganese Halides. In *Inorg. Synth.*, Reagents for Transition Metal Complex and Organometallic Syntheses; Wiley-Interscience: Hoboken, 2006; pp 154-159.

25. Hicks, J.; Juckel, M.; Paparo, A.; Dange, D.; Jones, C., Multigram Syntheses of Magnesium(I) Compounds Using Alkali Metal Halide Supported Alkali Metals as Dispersible Reducing Agents. *Organometallics* **2018**, *37*, 4810-4813.

26. Lai, T. Y.; Tao, L.; Britt, R. D.; Power, P. P., Reversible Sn-Sn Triple Bond Dissociation in a Distannyne: Support for Charge-Shift Bonding Character. *J. Am. Chem. Soc.* **2019**, *141*, 12527-12530.

27. Joy, J.; Danovich, D.; Kaupp, M.; Shaik, S., Covalent vs Charge-Shift Nature of the Metal-Metal Bond in Transition Metal Complexes: A Unified Understanding. *J. Am. Chem. Soc.* **2020**, *142*, 12277-12287.

28. Goodman, J. L.; Peters, K. S.; Vaida, V., The determination of the manganese-manganese bond strength in Mn<sub>2</sub>(CO)<sub>10</sub> using pulsed time-resolved photoacoustic calorimetry. *Organometallics* **1986**, *5*, 815-816.

29. Pan, S.; Zhao, L.; Dias, H. V. R.; Frenking, G., Bonding in Binuclear Carbonyl Complexes M<sub>2</sub>(CO)<sub>9</sub> (M = Fe, Ru, Os). *Inorg. Chem.* **2018**, *57*, 7780-7791.

30. Wolf, S.; Fenske, D.; Klopffer, W.; Feldmann, C., [Pb{Mn(CO)<sub>5</sub>}<sub>3</sub>][AlCl<sub>4</sub>]: a lead-manganese carbonyl with AlCl<sub>4</sub>-linked PbMn<sub>3</sub> clusters. *Dalton Trans.* **2019**, *48*, 4696-4701.

31. Braunschweig, H.; Damme, A.; Dewhurst, R. D.; Kramer, T.; Östreicher, S.; Radacki, K.; Vargas, A., Ditopic Ambiphilicity of an Anionic Dimetalloborylene Complex. *J. Am. Chem. Soc.* **2013**, *135*, 2313-2320.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
32. Clementi, E.; Raimondi, D. L.; Reinhardt, W. P., Atomic Screening Constants from SCF Functions. II. Atoms with 37 to 86 Electrons. *J. Chem. Phys.* **1967**, *47*, 1300-1307.
33. Pyykkö, P.; Atsumi, M., Molecular Single-Bond Covalent Radii for Elements 1–118. *Chem. Eur. J.* **2009**, *15*, 186-197.
34. Cordero, B.; Gómez, V.; Platero-Prats, A. E.; Revés, M.; Echeverría, J.; Cremades, E.; Barragán, F.; Alvarez, S., Covalent radii revisited. *Dalton Trans.* **2008**, 2832-2838.
35. Herberhold, M.; Tröbs, V.; Milius, W.; Wrackmeyer, B., Carbonyliron-Lead Complexes: Multinuclear Magnetic Resonance Study in Solution and X-Ray Structure Determination of  $[\text{Et}_2\text{PbFe}(\text{CO})_4]_2$ . *Z. Naturforsch. B* **1994**, *49*, 1781.
36. Wade, K., The structural significance of the number of skeletal bonding electron-pairs in carboranes, the higher boranes and borane anions, and various transition-metal carbonyl cluster compounds. *J. Chem. Soc. D* **1971**, 792-793.
37. Mingos, D. M. P., A General Theory for Cluster and Ring Compounds of the Main Group and Transition Elements. *Nat. Phys.* **1972**, *236*, 99-102.
38. Wei, C. H., Structural analyses of tetracobalt dodecacarbonyl and tetrarhodium dodecacarbonyl. Crystallographic treatments of a disordered structure and a twinned composite. *Inorg. Chem.* **1969**, *8*, 2384-2397.
39. Plečnik, C. E.; Liu, S.; Chen, X.; Meyers, E. A.; Shore, S. G., Lanthanide–Transition-Metal Carbonyl Complexes: New  $[\text{Co}_4(\text{CO})_{11}]^{2-}$  Clusters and Lanthanide(II) Isocarbonyl Polymeric Arrays. *J. Am. Chem. Soc.* **2004**, *126*, 204-213.
40. Chini, P., The closed metal carbonyl clusters. *Inorg. Chim. Acta* **1968**, *2*, 31-51.
41. Farrugia, L. J.; Braga, D.; Grepioni, F., A structural redetermination of  $\text{Co}_4(\text{CO})_{12}$ : evidence for dynamic disorder and the pathway of metal atom migration in the crystalline phase. *J. Organomet. Chem.* **1999**, *573*, 60-66.
42. Zhang, X.; Li, Q.-s.; Xie, Y.; King, R. B.; Schaefer, H. F., A Carbonyl Group Bridging Four Metal Atoms in a Homoleptic Carbonylmetal Cluster: The Remarkable Case of  $-\text{Co}_4(\text{CO})_{11}$ . *Eur. J. Inorg. Chem.* **2008**, 2158-2164.
43. Simons, R. S.; Pu, L.; Olmstead, M. M.; Power, P. P., Synthesis and Characterization of the Monomeric Diaryls  $\text{M}\{\text{C}_6\text{H}_3\text{-2,6-Mes}_2\}_2$  (M = Ge, Sn, or Pb; Mes = 2,4,6-Me<sub>3</sub>C<sub>6</sub>H<sub>2</sub>-) and Dimeric Aryl–Metal Chlorides  $[\text{M}(\text{Cl})\{\text{C}_6\text{H}_3\text{-2,6-Mes}_2\}]_2$  (M = Ge or Sn). *Organometallics* **1997**, *16*, 1920-1925.
44. Kano, N.; Tokitoh, N.; Okazaki, R., Synthesis and X-ray Crystal Structure of Bis{2,4,6-tris[bis(trimethylsilyl)methyl]phenyl}dibromoplumbane: The First Monomeric Diorganodihaloplumbane in the Crystalline State. *Organometallics* **1997**, *16*, 2748-2750.
45. McCrea-Hendrick, M.; Bursch, M.; Gullett, K.; Maurer, L.; Fettinger, J.; Grimme, S.; Power, P., Counterintuitive Interligand Angles in the Diaryls  $\text{E}\{\text{C}_6\text{H}_3\text{-2,6-(C}_6\text{H}_2\text{-2,4,6- iPr}_3\text{)}_2\}_2$  (E = Ge, Sn, or Pb) and Related Species: The Role of London Dispersion Forces. *Organometallics* **2018**, *37*.
46. Elschenbroich, C., *Organometallic*. Wiley-VCH: Weinheim, 2006; pp. 356-372.
47. Aime, S.; Gobetto, R.; Osella, D.; Milonej, L.; Hawkes, G. E.; Randall, E. W., Reinvestigation of the solution structure of  $\text{Co}_4(\text{CO})_{12}$  by <sup>13</sup>C and <sup>59</sup>Co NMR. 2. *J. Magn. Reson. (1969)* **1985**, *65*, 308-315.
48. Richert, T.; Elbayed, K.; Raya, J.; Granger, P.; Braunstein, P.; Rosé, J., <sup>59</sup>Co NMR in Tetrahedral Clusters. *Magn. Reson. Chem.* **1996**, *34*, 689-696.
49. Friedel, R. A.; Wender, I.; Shufler, S. L.; Sternberg, H. W., Spectra and Structures of Cobalt Carbonyls. *J. Am. Chem. Soc.* **1955**, *77*, 3951-3958.
50. Coffey, C. E.; Lewis, J.; Nyholm, R. S., 339. Metal–metal bonds. Part I. Compounds of gold(0)

with the carbonyls of manganese, iron, and cobalt. *J. Chem. Soc.* **1964**, 1741-1749.

51. Tan, G.; Szilvási, T.; Inoue, S.; Blom, B.; Driess, M., An Elusive Hydridoaluminum(I) Complex for Facile C–H and C–O Bond Activation of Ethers and Access to Its Isolable Hydridogallium(I) Analogue: Syntheses, Structures, and Theoretical Studies. *J. Am. Chem. Soc.* **2014**, *136*, 9732-9742.

52. Pomeroy, R. K.; Vancea, L.; Calhoun, H. P.; Graham, W. A. G., Stereochemically nonrigid six-coordinate metal carbonyl complexes. 3. The series cis-Fe(CO)<sub>4</sub>(SnR<sub>3</sub>)<sub>2</sub> (R = methyl, ethyl, propyl, butyl, phenyl, chloro) and the x-ray structure of tetracarbonylbis(triphenylstannyl)iron(II). *Inorg. Chem.* **1977**, *16*, 1508-1514.

## For Table of Contents

