

This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Daya Bay collaboration; JUNO collaboration

Title: Optimization of the JUNO liquid scintillator composition using a Daya Bay antineutrino detector

Year: 2021

Version: Accepted version (Final draft)

Copyright: © 2021 Elsevier

Rights: CC BY-NC-ND 4.0

Rights url: https://creativecommons.org/licenses/by-nc-nd/4.0/

Please cite the original version:

Daya Bay collaboration, JUNO collaboration. (2021). Optimization of the JUNO liquid scintillator composition using a Daya Bay antineutrino detector. Nuclear Instruments and Methods in Physics Research Section A: Accelerators Spectrometers Detectors and Associated Equipment, 988, Article 164823. https://doi.org/10.1016/j.nima.2020.164823

Optimization of the JUNO liquid scintillator composition using a Daya Bay antineutrino detector

A. Abusleme, T. Adam, S. Ahmad, S. Aiello, M. Akram, N. Ali, F.P. An, G.P. An, Q. An, G. Andronico, N. Anfimov, V. Antonelli, T. Antoshkina, B. Asavapibhop, J.P.A.M. de André, A. Babic, A.B. Balantekin, W. Baldini, M. Baldoncini, H.R. Band, A. Barresi, E. Baussan, M. Bellato, E. Bernieri, D. Biare, T. Birkenfeld, M. Bishai, S. Blin, D. Blum, S. Blyth, C. Bordereau, A. Brigatti, R. Brugnera, A. Budano, P. Burgbacher, M. Buscemi, S. Bussino, J. Busto, I. Butorov, A. Cabrera, H. Cai, X. Cai, Y.K. Cai, Z.Y. Cai, A. Cammi, A. Campeny, C.Y. Cao, G.F. Cao, J. Cao, R. Caruso, C. Cerna, I. Chakaberia, J.F. Chang, Y. Chang, H.S. Chen, P.A. Chen, P.P. Chen, S.M. Chen, S.J. Chen, X.R. Chen, Y.W. Chen, Y.X. Chen, Y. Chen, Z. Chen, J. Cheng, Y.P. Cheng, Z.K. Cheng, A. Chepurnov, J.J. Cherwinka, F. Chiarello, D. Chiesa, P. Chimenti, M.C. Chu, A. Chukanov, A. Chuvashova, Clementi, B. Clerbaux, S. Conforti Di Lorenzo, D. Corti, S. Costa, F.D. Corso, J.P. Cummings, O. Dalager, C. De La Taille, F.S. Deng, J.W. Deng, Z. Deng, Z.Y. Deng, W. Depnering, M. Diaz, X.F. Ding, Y.Y. Ding, B. Dirgantara, S. Dmitrievsky, M.V. Diwan, T. Dohnal, G. Donchenko, J.M. Dong, D. Dornic, E. Doroshkevich, J. Dove, M. Dracos, F. Druillole, S.X. Du, S. Dusini, M. Dvorak, D.A. Dwyer, T. Enqvist, H. Enzmann, A. Fabbri, L. Fajt, D.H. Fan, L. Fan, C. Fang, J. Fang, A. Fatkina, D. Fedoseev, V. Fekete, L.C. Feng, Q.C. Feng, G. Fiorentini, R. Ford, A. Formozov, A. Fournier, S. Franke, J.P. Gallo, H.N. Gan, F. Gao, A. Garfagnini, A. Göttel, C. Genster, M. Giammarchi, A. Giaz, N. Giudice, F. Giuliani, M. Gonchar, G.H. Gong, H. Gong, O. Gorchakov, Y. Gornushkin, M. Grassi, C. Grewing, M. Gromov, V. Gromov, M.H. Gu, W.Q. Gu, X.F. Gu, Y. Gu, M.Y. Guan, N. Guardone, M. Gul, C. Guo, J.Y. Guo, L. Guo, W.L. Guo, X.H. Guo, Y.H. Guo, Z. Guo, M. Haacke, R.W. Hackenburg, P. Hackspacher, C. Hagner, R. Han, Y. Han, S. Hans, M. He, W. He, K.M. Heeger, T. Heinz, Y.K. Heng, R. Herrera, A. Higuera, D.J. Hong, Y.K. Hor, S.J. Hou, Y.B. Hsiung, B.Z. Hu, H. Hu, J.R. Hu, J. Hu, S.Y. Hu, T. Hu, Z.J. Hu, C.H. Huang, G.H. Huang, H.X. Huang, Q.H. Huang, W.H. Huang, X.T. Huang, Y.B. Huang, P. Huber, J.Q. Hui, L. Huo, W.J. Huo, C. Huss, S. Hussain, A. Insolia, A. Ioannisian, D. Ioannisyan, R. Isocrate, D.E. Jaffe, K.L. Jen, X.L. Ji, X.P. Ji, X.Z. Ji, H.H. Jia, J.J. Jia, S.Y. Jian, D. Jiang, X.S. Jiang, R.Y. Jin, X.P. Jing, R.A. Johnson, C. Jollet, D. Jones, J. Joutsenvaara, S. Jungthawan, L. Kalousis, P. Kampmann, L. Kang, M. Karagounis, N. Kazarian, S.H. Kettell, A. Khan, W. Khan, K. Khosonthongkee, P. Kinz, S. Kohn, D. Korablev, K. Kouzakov, M. Kramer, A. Krasnoperov, S. Krokhaleva, Z. Krumshteyn, A. Kruth, N. Kutovskiy, P. Kuusiniemi, B. Lachacinski, T. Lachenmaier, C. Landini, T.J. Langford, J. Lee, J.H.C. Lee, F. Lefevre, L. Lei, R. Lei,



R. Leitner, J. Leung, C. Li, D.M. Li, F. Li, F. Li, H.T. Li, H.L. Li, J. Li, J.J. Li, J.Q. Li, K.J. Li, M.Z. Li, N. Li, N. Li, Q.J. Li, Q.J. Li, R.H. Li, S.C. Li, S.F. Li, S.J. Li, T. Li, T. Li, W.D. Li, W.G. Li, X.M. Li, X.N. Li, X.L. Li, X.Q. Li, Y. Li, Y.F. Li, Z.B. Li, Z.Y. Li, H. Liang, H. Liang, J.J. Liang, D. Liebau, A. Limphirat, S. Limpijumnong, C.J. Lin, G.L. Lin, S.X. Lin, T. Lin, Y.H. Lin, J.J. Ling, J.M. Link, I. Lippi, L. Littenberg, B.R. Littlejohn, F. Liu, H. Liu, H. Liu, H.B. Liu, H.D. Liu, H.J. Liu, H.T. Liu, J.C. Liu, J.L. Liu, M. Liu, Q. Liu, Q. Liu, R.X. Liu, S.Y. Liu, S.B. Liu, S.L. Liu, X.W. Liu, Y. Liu, A. Lokhov, P. Lombardi, K. Loo, S. Lorenz, C. Lu, C. Lu, H.Q. Lu, J.B. Lu, J.G. Lu, S.X. Lu, X.X. Lu, B. Lubsandorzhiev, S. Lubsandorzhiev, L. Ludhova, K.B. Luk, F.J. Luo, G. Luo, P.W. Luo, S. Luo, W.M. Luo, V. Lyashuk, Q.M. Ma, S. Ma, X.B. Ma, X.Y. Ma, Y.Q. Ma, Y. Malyshkin, F. Mantovani, Y.J. Mao, S.M. Mari, F. Marini, S. Marium, C. Marshall, C. Martellini, G. Martin-Chassard, D.A. Martinez Caicedo, A. Martini, J. Martino, D. Mayilyan, K.T. McDonald, R.D. McKeown, A. Müller, G. Meng, Y. Meng, A. Meregaglia, E. Meroni, D. Meyhöfer, M. Mezzetto, J. Miller, L. Miramonti, S. Monforte, P. Montini, M. Montuschi, N. Morozov, P. Muralidharan, J. Napolitano, M. Nastasi, D.V. Naumov, E. Naumova, I. Nemchenok, A. Nikolaev, F.P. Ning, Z. Ning, H. Nunokawa, L. Oberauer, J.P. Ochoa-Ricoux, A. Olshevskiy, F. Ortica, H.R. Pan, A. Paoloni, J. Park, N. Parkalian, S. Parmeggiano, S. Patton, T. Payupol, V. Pec, D. Pedretti, Y.T. Pei, N. Pelliccia, A.G. Peng, H.P. Peng, J.C. Peng, F. Perrot, P.A. Petitjean, L.F. Pineres Rico, A. Popov, P. Poussot, W. Pratumwan, E. Previtali, C.S.J. Pun, F.Z. Qi, M. Qi, S. Qian, X. Qian, X.H. Qian, H. Qiao, Z.H. Qin, S.K. Qiu, M. Rajput, G. Ranucci, N. Raper, A. Re, H. Rebber, A. Rebii, B. Ren, J. Ren, C.M. Reveco, T. Rezinko, B. Ricci, M. Robens, M. Roche, N. Rodphai, L. Rohwer, A. Romani, R. Rosero, B. Roskovec, C. Roth, X.C. Ruan, X.D. Ruan, S. Rujirawat, A. Rybnikov, A. Sadovsky, P. Saggese, G. Salamanna, A. Sangka, N. Sanguansak, U. Sawangwit, J. Sawatzki, F. Sawy, M. Schever, J. Schuler, C. Schwab, K. Schweizer, D. Selivanov, A. Selyunin, A. Serafini, G. Settanta, M. Settimo, M. Shahzad, G. Shi, J.Y. Shi, Y.J. Shi, V. Shutov, A. Sidorenkov, F. Simkovic, C. Sirignano, J. Siripak, M. Sisti, M. Slupecki, M. Smirnov, O. Smirnov, T. Sogo-Bezerra, J. Songwadhana, B. Soonthornthum, A. Sotnikov, O. Sramek, W. Sreethawong, A. Stahl, L. Stanco, K. Stankevich, D. Stefanik, H. Steiger, H. Steiner, J. Steinmann, M. Stender, V. Strati, A. Studenikin, G.X. Sun, L.T. Sun, J.L. Sun, S.F. Sun, X.L. Sun, Y.J. Sun, Y.Z. Sun, N. Suwonjandee, M. Szelezniak, J. Tang, Q. Tang, Q. Tang, X. Tang, A. Tietzsch, I. Tkachev, T. Tmej, K. Treskov, G. Troni, W. Trzaska, W.-H. Tse, C.E. Tull, C. Tuve, S. van Waasen, J. Vanden Boom, N. Vassilopoulos, V. Vedin, G. Verde, M. Vialkov, B. Viaud, B. Viren, C. Volpe, V. Vorobel, L. Votano, P. Walker, C. Wang, C.H. Wang, E. Wang, G.L. Wang, J. Wang, J. Wang, K.Y. Wang, L. Wang, M.F. Wang, M. Wang, M. Wang, N.Y. Wang, R.G. Wang, S.G. Wang, W. Wang, W. Wang, W.S. Wang, X. Wang, X.Y. Wang, Y. Wang, Y. Wang, Y. Wang, Y.F. Wang, Y.G. Wang,

Y.M. Wang, Y.Q. Wang, Z. Wang, Z. Wang, Z.M. Wang, Z.Y. Wang, A. Watcharangkool, H.Y. Wei, L.H. Wei, W. Wei, Y.D. Wei, L.J. Wen, K. Whisnant, C.G. White, C. Wiebusch, S.C.F. Wong, H.L.H. Wong, B. Wonsak, E. Worcester, C.H. Wu, D.R. Wu, F.L. Wu, Q. Wu, W.J. Wu, Z. Wu, M. Wurm, J. Wurtz, C. Wysotzki, Y.F. Xi, D.M. Xia, Y.G. Xie, Z.Q. Xie, Z.Z. Xing, D.L. Xu, F.R. Xu, H.K. Xu, J.L. Xu, J. Xu, M.H. Xu, T. Xu, Y. Xu, Y. Xu, T. Xue, B.J. Yan, X.B. Yan, Y.P. Yan, A.B. Yang, C.G. Yang, H. Yang, J. Yang, L. Yang, X.Y. Yang, Y.F. Yang, Y.Z. Yang, H.F. Yao, Z. Yasin, J.X. Ye, M. Ye, U. Yegin, M. Yeh, F. Yermia, P.H. Yi, Z.Y. You, B.L. Young, B.X. Yu, C.X. Yu, C.Y. Yu, H.Z. Yu, M. Yu, X.H. Yu, Z.Y. Yu, C.Z. Yuan, Y. Yuan, Z.X. Yuan, Z.Y. Yuan, B.B. Yue, N. Zafar, A. Zambanini, P. Zeng, S. Zeng, T.X. Zeng, Y.D. Zeng, L. Zhan, C. Zhang, F.Y. Zhang, G.Q. Zhang, H.H. Zhang, H.Q. Zhang, J. Zhang, J.B. Zhang, J.W. Zhang, P. Zhang, Q.M. Zhang, T. Zhang, X.M. Zhang, X.T. Zhang, Y. Zhang, Y.H. Zhang, Y.M. Zhang, Y.P. Zhang, Y.X. Zhang, Y.Y. Zhang, Y.Y. Zhang, Z.J. Zhang, Z.P. Zhang, Z.Y. Zhang, Z.Y. Zhang, F.Y. Zhao, J. Zhao, R. Zhao, S.J. Zhao, T.C. Zhao, D.Q. Zheng, H. Zheng, M.S. Zheng, Y.H. Zheng, W.R. Zhong, J. Zhou, L. Zhou, N. Zhou, S. Zhou, X. Zhou, J. Zhu, K.J. Zhu, H.L. Zhuang, L. Zong, J.H. Zou

PII:	S0168-9002(20)31220-1
DOI:	https://doi.org/10.1016/j.nima.2020.164823
Reference:	NIMA 164823
To appear in:	Nuclear Inst. and Methods in Physics Research, A
Received date :	24 July 2020
Revised date :	13 October 2020
Accepted date :	27 October 2020

Please cite this article as: A. Abusleme, T. Adam, S. Ahmad et al., Optimization of the JUNO liquid scintillator composition using a Daya Bay antineutrino detector, *Nuclear Inst. and Methods in Physics Research, A* (2020), doi: https://doi.org/10.1016/j.nima.2020.164823.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Elsevier B.V. All rights reserved.

Optimization of the JUNO liquid scintillator composition using a Daya Bay antineutrino detector

Abstract

To maximize the light yield of the liquid scintillator (LS) for the Jiangmen Underground Neutrino Observatory (JUNO), a 20 t LS sample was produced in a pilot plant at Daya Bay. The optical properties of the new LS in various compositions were studied by replacing the gadolinium-loaded LS in one antineutrino detector. The concentrations of the fluor, PPO, and the wavelength shifter, bis-MSB, were increased in 12 steps from 0.5 g/L and <0.01 mg/L to 4 g/L and 13 mg/L, respectively. The numbers of total detected photoelectrons suggest that, with the optically purified solvent, the bis-MSB concentration does not need to be more than 4 mg/L. To bridge the one order of magnitude in the detector size difference between Daya Bay and JUNO, the Daya Bay data were used to tune the parameters of a newly developed optical model. Then, the model and tuned parameters were used in the JUNO simulation. This enabled to determine the optimal composition for the JUNO LS: purified solvent LAB with 2.5 g/L PPO, and 1 to 4 mg/L bis-MSB.

Keywords: neutrino, liquid scintillator, light yield

Preprint submitted to Nucl. Instr. Meth. A

1 1. Introduction

Liquid scintillator (LS) detectors readout by photomultiplier tubes (PMTs) have supported neutrino physics 2 for several decades, from the discovery of neutrinos in the 1950s [1], to the precise measurement of the neutrino 3 mass squared splitting Δm_{21}^2 at KamLAND [2], the precise measurement of solar neutrinos at Borexino [3], 4 and the observation of the θ_{13} -driven neutrino oscillation at Daya Bay [4]. Given the high light yield, good 5 transparency and relatively low price, this kind of detector is also adopted by the Jiangmen Underground Neutrino 6 Observatory (JUNO) [5, 6], which utilizes 20 kt LS with one of the physics goals of determining the neutrino mass 7 ordering. Since the sensitivity comes from a precise measurement of the fine structure in the oscillated neutrino 8 spectrum, a crucial requirement on the JUNO detector is the excellent energy resolution, ~3% at 1 MeV, corresponding 9 to at least 1100 detected photoelectrons (p.e.) per MeV of deposited energy [5]. The number of detected p.e. per MeV 10 is referred to as ObsLY hereafter. The higher ObsLY is, the better the energy resolution and the physics sensitivity are. 11 12 Thus, one of the keys of the JUNO detector development is to maximize ObsLY.

In recent LS experiments, a widely used solvent is linear alkylbenzene (LAB), with 2,5-diphenyloxazole (PPO) as 13 the fluor and p-bis-(o-methylstyryl)-benzene (bis-MSB) as the wavelength shifter. The ionization of a charged particle 14 excites the LAB molecules. A fraction of the excitation energy is transferred to the PPO. Scintillation photons are 15 generated from the de-excitation of PPO molecules. The total number of photons released by PPO is defined as the 16 initial light yield, which rapidly increases with the PPO concentration before reaching 2 g/L. Above this concentration 17 the increase becomes much less steep. The wavelengths of most initial photons are shifted to longer wavelengths by 18 the absorption and re-emission of bis-MSB. This shift is crucial as the long wavelength avoids spectral self-absorption 19 by the solvent and allows the photons to reach PMTs far away from the energy deposit points. Eventually, ObsLY is 20 a joint effect of the initial light yield, the photon absorption and re-emission during propagation, and the wavelength-21 dependent PMT quantum efficiency (QE). To obtain the maximum ObsLY, these aspects must be simultaneously 22 optimized. 23

There have been many studies that independently measured the optical properties of LS, such as Refs. [7, 8, 9] for the initial light yields, Refs. [10, 11, 12] for the transparency. To completely deal with the competing photon absorption and subsequent re-emission processes of the LS components, a comprehensive optical model was developed and reported in Ref. [13]. However, the parameters used in the model were obtained from bench-top experiments with a typical detector size of a few centimeters. Before usage in JUNO, a spherical LS detector with an inner diameter of

²⁹ 35.4 m, the model and its parameters should be validated based on data collected in a larger detector.

This requirement motivated a dedicated LS experiment at Daya Bay. A LS pilot plant was built by the JUNO collaboration in the underground LS hall of Daya Bay. One Daya Bay antineutrino detector (AD) [14, 15] in the Experimental Hall 1 (EH1-AD1) stopped data taking in January 2017. The 20 t gadolinium-loaded LS (Gd-LS) in the innermost cylindrical vessel with 3 m in diameter and height was replaced with purified LS produced by the pilot plant. PPO and bis-MSB concentrations were 0.5 g/L and less than 0.01 mg/L, and then increased in 12 steps to 4 g/L and 13 mg/L, respectively. The ObsLY in the 13 samples was measured at 0.5% precision level. The above-mentioned

³⁶ optical model was successfully tuned to the data. This allowed to identify the optimal scintillator composition for the ³⁷ final JUNO detector.

The structure of this paper is as follows: Sec. 2 describes the Daya Bay AD and the LS replacement experiment. Section 3 presents results of the ObsLY measurements. Section 4 discusses the determination of the JUNO LS composition using the optical model tuned to Daya Bay data.

41 2. The JUNO LS pilot experiment at Daya Bay

The Daya Bay reactor neutrino experiment started data taking on December 24, 2011. With millions of 42 $\overline{\nu}_e$ interactions detected in eight identically designed ADs in three underground EHs, many physics results have been 43 produced, including the current world-leading measurements of the neutrino mixing angle θ_{13} and the squared mass-44 splitting $|\Delta m_{33}^2|$ [16], precise measurements of the reactor $\bar{\nu}_e$ flux and spectrum [17, 18, 19], and stringent limits on the 45 existence of light sterile neutrinos [20, 21]. Three nested cylindrical volumes in each AD are separated by concentric 46 acrylic vessels (IAV, OAV), as shown in Fig. 1. The innermost volume is filled with 20 t of Gd-LS, serving as the 47 48 primary $\bar{\nu}_e$ target. It is surrounded by ~22 t of non-loaded LS to detect γ -rays escaping from the target volume. The outermost volume is filled with mineral oil to shield the LS from natural radioactivity. A total of 192 8-inch 49 PMTs (Hamamatsu R-5912) are installed on the steel vessel to detect scintillation photons. There are three Automated 50 Calibration Units (ACUs) on the top of each AD to calibrate the energy response along the vertical axes at the detector 51

center (ACU-A), the edge of the Gd-LS volume (ACU-B, removed for the replacement experiment), and the LS
 volume (ACU-C). Details of the detector systems are reported in Refs. [14, 15, 22].



Figure 1: Schematic of a Daya Bay antineutrino detector. The three cylindrical volumes, defined by two acrylic vessels, are filled with Gd-LS, LS, and mineral oil from the innermost to the outermost. Three Automated Calibration Units are installed on top of the detector to calibrate the detector's energy response. Two interfaces are installed on the top of EH1-AD1 for the LS replacement experiment, one through the central overflow tank and the other one using the port of ACU-B.

Figure 2 shows photographs of the LS experiment, including the pilot plant, the LS replacement system and 54 EH1-AD1. The pilot plant built in the LS hall consisted of four primary subsystems for purification in sequence: the 55 Al_2O_3 column, the distillation, the water extraction and the steam stripping. In addition there were several supporting 56 subsystems for the PPO and bis-MSB dissolution, and the generation of purified water and nitrogen. To control the 57 optical qualities, an apparatus with a 1-m long tube was built to measure the light attenuation in the liquids. The 58 20 t LAB produced by the Jinling LAB factory was purified by the Al₂O₃ column for optical transparency and then 59 distillated for radiopurity. About 11.6 kg PPO produced by the Haiso Technology Co., LTD was dissolved and added 60 to the LAB. Thus, the initial LS composition was 0.5 g/L PPO without bis-MSB. Before filling AD1, the mixture went 61 through the water extraction and steam stripping systems for radiopurity. Details of the distillation and steam stripping 62 systems have been reported in Ref. [23]. The attenuation length of the purified LAB was measured to be 25.1 m at 63 the photon wavelength of 430 nm, and the attenuation length of the initial LS was 23.8 m. The uncertainties of these 64 values were about 2 m, primarily from the relatively short light path compared to the attenuation length. 65

A replacement system was set up in EH1 to drain the Gd-LS and refill the IAV with new LS. The system consisted 66 of three pumps, three buffer tanks and stainless steel pipes connecting to the two interfaces on AD1. The central 67 interface A for injecting liquids was installed via the central overflow tank while the draining was performed via an 68 acrylic tube inserted to the bottom of IAV through the interface B. The ACU-B was removed for installing the interface 69 B. In fact, the liquids could be filled or drained from either interfaces. The system could run in a self-circulation or 70 a full-circulation mode. In the former mode, liquids were pumped from the IAV to a 300 L buffer tank and then 71 re-inserted into the IAV. In the latter mode, liquids in the IAV were drained and sent to the facilities in the LS hall for 72 further processing, while newly produced liquids or the re-processed ones were returned to the IAV. 73

To avoid mixing of the old Gd-LS and the new LS, the Gd-LS was replaced first by purified water that was then 74 replaced by the new LS. This replacement method was also required to prevent destructive stress on the acrylic vessels. 75 From February 16 to February 22 2017, the Gd-LS was drained at a rate of about 300 L/h through the central interface 76 while purified water was filled through the side interface. Then, the new LS with 0.5 g/L PPO was filled through the 77 central interface while water was drained through the side one. Limited by the position of the central interface A, 78 about 10 L Gd-LS could not be drained out. This resulted in a residual bis-MSB concentration of less than 0.01 mg/L, 79 confirmed by the light absorption measurement with a UV-vis spectrometer. In addition, since the tube through the 80 81 interface B could not touch the IAV bottom, a layer of water with a thickness of about 1 cm was left. Eventually, the





Figure 2: Photographs of EH1-AD1, the replacement system in EH1, and the pilot plant in the LS hall (Hall 5).

Step	PPO	bis-MSB	Date of calibration	LS temperature	ObsLY (p.e./MeV)
Initial	0.5 g/L	<0.01 mg/L	April 28, 2017	22.6 °C	123.7
1	1.0 g/L	<0.01 mg/L	May 28, 2017	22.6 °C	150.3
2	2.0 g/L	<0.01 mg/L	June 4, 2017	22.5 °C	167.7
3	2.0 g/L	0.1 mg/L	June 9, 2017	22.6 °C	177.2
4	2.0 g/L	1.0 mg/L	June 13, 2017	22.6 °C	183.2
5	2.0 g/L	4.0 mg/L	June 18, 2017	22.6 °C	184.3
6	2.0 g/L	7.0 mg/L	June 23, 2017	22.6 °C	184.8
7	2.5 g/L	7.0 mg/L	June 29, 2017	22.6 °C	189.5
8	3.0 g/L	7.0 mg/L	July 5, 2017	22.6 °C	191.6
9	3.5 g/L	7.0 mg/L	July 11, 2017	22.6 °C	192.6
10	4.0 g/L	7.0 mg/L	July 17, 2017	22.7 °C	192.6
11	4.0 g/L	10.0 mg/L	July 22, 2017	22.7 °C	193.0
12	4.0 g/L	13.0 mg/L	July 27, 2017	22.7 °C	193.3

Table 1: Summary of the LS experiment. Each LS composition change took 4 to 5 days, including a slow addition of PPO or bis-MSB over 1.5 days followed by at least 3 days of self-circulation. The ObsLY was measured to a precision of 0.5% using a ⁶⁰Co calibration source in the detector center.

replacement was successfully finished on March 7. To obtain the radiopurity of LS with 0.5 g/L PPO, the replacement

system was shut down after ten days of self-circulation. The LS radiopurity was measured later in May to wait for the
 decay of the ²²²Rn contamination.

Beginning on May 20, PPO and bis-MSB were added in 12 steps as summarized in Table 1. In each step, the 85 replacement system was working in the full-circulation mode. The LS was pumped out with a 300 L/h speed and sent 86 to a buffer tank of the water extraction system in the LS hall. The PPO or bis-MSB was dissolved and slowly added to 87 the buffer tank in 36 hours. During this time about half of the total LS volume was circulated. Then, the replacement 88 system ran in the self-circulation mode with a 300 L/h rate for about three days to obtain the uniform fluor distribution 89 in the IAV. The fluor concentration was measured every 12 hours with a UV-vis spectrometer. In general, after two 90 days of self-circulation the concentration was stabilized at the target value. In the following the detector response was 91 calibrated by deploying ⁶⁰Co calibration sources along ACU-A and ACU-C. The proceduce generally took about six 92 hours. ObsLY was determined using the data collected with the ⁶⁰Co source deployed in the detector center. 93

94 **3.** The light yield measurements

The scintillation photons are detected by the 192 PMTs of the AD, operating at an average gain of 1×10^7 . All the

PMTs were working without problems over the three-months period. In the Daya Bay readout system, after an initial

fast amplification, the PMT signal is fed to a pulse shaping circuit consisting of a differential CR and four integrating 97 RC circuits (CR-(RC)⁴), and then amplified by a factor of ten. The integrated value, sampled by a 40-MHz 12-bit 98 ADC, is used as an estimate of the PMT charge output [24]. The PMT gains were separately calibrated in the thirteen 99 measurements, using the PMT dark noises captured by the data acquisition system. The CR-(RC)⁴ shaping circuit, 100 combined with the time distribution of detected light, introduces a $\sim 10\%$ nonlinearity in the charge measurement 101 of a single channel, dubbed the electronics nonlinearity. From the initial concentration to the 6th step in Table 1, 102 this electronics nonlinearity was carefully measured and corrected with the help of a full Flash ADC readout system 103 following the method reported in Ref. [25]. From the 7th to 12th steps, the Flash ADC readout system was not working 104 well. Therefore, the measured nonlinearity in the 6th step was used to correct nonlinearities in the 7th to 12th steps. 105 Such corrections were feasible because the time distributions of detected light were found to be stable in the 6th to 12th 106 steps. 107

For each concentration, ObsLY was determined by measuring the scintillation light originating from the two 108 γ -rays of ⁶⁰Co decays, corresponding to a total deposit energy of 2.505 MeV. Figure 3 and Table 1 summarize the 109 measured light yields with the 13 LS compositions. For the first three steps with less than 0.01 mg/L bis-MSB from 110 residual Gd-LS, the light yield increased by more than 40% with PPO concentrations increasing from 0.5 g/L to 2 g/L. 111 Adding 1 mg/L bis-MSB further increased the light yield by 10%. However, no significant increase was found when 112 further raising the bis-MSB concentration. This indicates that for the very transparent LAB, scintillation light emitted 113 by PPO would either be absorbed and re-emitted by bis-MSB, or directly reach the PMTs. Adding more bis-MSB 114 does only shift the fractions of photons from PPO and bis-MSB when reaching the PMTs. In addition, after adding 115

bis-MSB, the increase of PPO to more than 2.5 g/L yielded no obvious effect on ObsLY, suggesting the initial light

yield had reached the plateau for particles with low energy deposit density, such as γ 's and e^{\pm} 's.



Figure 3: Measured light yields ObsLY versus PPO and bis-MSB concentrations. The labels of the horizontal axis are the concentrations of PPO (g/L) plus that of bis-MSB (mg/L). The vertical error bar (0.2%) is statistical only and too small to be visible. The points in each colored box are measured at the same PPO or bis-MSB concentration.

To compare ObsLY of different LS compositions, we have considered the relative uncertainties mainly arising from three sources: the LS temperature variation, the statistical fluctuations in the determination of PMT gains and of the 60 Co peak, and the electronics nonlinearity correction. The LS temperature was monitored using four sensors and was found to be stable within 0.2 °C over the three months, resulting in a less than 0.1% light yield variation based on the measurements in Ref. [26]. The second term is estimated to be at 0.2% level according to the fitted energy peak position of the 60 Co source. The third term is less than 0.5% for each measurement as discussed in Ref. [25]. Combining the three sources, the uncertainty of each light yield measurement is estimated to be 0.5%.

From August 2017 to January 2019, several rounds of radiopurity studies have been carried out. In this period, the LS composition was kept at 4 g/L PPO and 13 mg/L bis-MSB. A stable ObsLY was found within $\pm 0.5\%$ as shown in Fig. 4.



Figure 4: ObsLY of the new LS with 4 g/L PPO and 13 mg/L bis-MSB kept stable within $\pm 0.5\%$ (the shaded band) between August 2017 and January 2019. The uncertainties are dominated by statistics of the 60 Co calibration events.

4. Optimization of the JUNO LS composition

The ObsLY measurements performed at Daya Bay are of great importance to future LS experiments, such as 129 JUNO, for the determination of the LS composition. However, ObsLY of Daya Bay cannot be directly used in JUNO, 130 since ObsLY is affected by a few factors, such as the initial light yield, the self-absorption and re-emission effects 131 during propagation, and the PMT QE spectra. The 20 kt LS of JUNO are contained in a spherical acrylic vessel 132 with an inner diameter of 35.4 m. Scintillation light is detected by about 18,000 20-inch PMTs, including 5,000 133 Hamamatsu R-12860 dynode PMTs, and 13,000 NNVT GDG-6201 PMTs with a microchannel plate (MCP-PMT) 134 instead of a dynode structure. In Daya Bay the new LS was studied in a cylindrical vessel with 3 m in height and 135 diameter, and Hamamatsu R-5912 8-inch PMTs were used. 136

To take these differences into account, a newly developed optical model [13] has been implemented in the 137 simulation of both experiments. The model is designed to thoroughly deal with the competing photon absorption 138 and subsequent re-emission processes of the LS components. It starts with primary scintillation photons emitted by 139 PPO. During the propagation, a photon could either be absorbed, or be scattered, or vanish when reaching optical 140 boundaries such as PMTs. The absorption could happen on any LS component, according to the Beer-Lambert law 141 and the absorption spectrum of each component. A new photon with longer wavelength may be emitted if the original 142 photon is absorbed by PPO or bis-MSB. The re-emission probability is defined as the fluorescence quantum efficiency. 143 In this model, scattering of optical photons happens via the Rayleigh process. Once a photon is scattered, it changes 144 direction and continues propagation. The Rayleigh scattering length of LAB is found to be 27.0±2.3 m at 430 nm [27]. 145 Thus, key optical parameters in the model consist of emission spectra of PPO and bis-MSB, absorption spectra of LAB, 146 PPO and bis-MSB, and wavelength-dependent fluorescence quantum efficiencies of PPO and bis-MSB. 147

The measurements of the key optical parameters are described below and in Ref. [13]. The emission spectra 148 of PPO and bis-MSB were well measured with a Fluorolog Tau-3 spectrometer as shown in Fig. 5. The absorption 149 spectrum of each LS component used in Daya Bay and the pilot plant was measured using a Shimadzu UV2550 UV-150 vis spectrometer and quartz cuvettes with different light paths up to 10 cm. To overcome the large uncertainty due 151 to limited cuvette sizes and the long absorption length at photon wavelength of 430 nm, a 1-m long tube was used 152 to measure the attenuation length. Then, the absorption length were obtained by subtracting the Rayleigh scattering 153 length from the attenuation one. Comparison of absorption spectra among the Daya Bay original liquids and the 154 purified JUNO liquids is shown in Fig. 6. The purification significantly improved the transparencies of LAB and 155 PPO. The general method to measure fluorescence QE was using the combination of a fluorescence spectrometer 156 and a UV-Vis spectrometer. An average QE spectrum from several measurements [10, 11, 12] was adopted in the 157 simulation, shown as the default QE spectra in Fig. 5. Due to intrinsic difficulties of several corrections in this method, 158



the measured efficiencies had relatively large uncertainties, typically 5%.

Figure 5: Emission spectra of PPO and bis-MSB in arbitrary units, and their QE efficiencies. The default PPO QE spectra are from the average of three bench-top measurements, while the tuned ones are from the tuning based on Daya Bay data taken in LS experiment in Table 1.



Figure 6: Comparison of the absorption spectra of LAB and PPO used in Daya Bay and produced by JUNO pilot plant. The purification significantly improved the optical transparency.

The optical model has been implemented in the Geant4 [28] based Daya Bay simulation. The residual 1-cm 160 water at the bottom of IAV has been included in the simulation by assuming a perfect surface between the water and 161 the LS. The bis-MSB of less than 0.01 mg/L from the residual Gd-LS is also included. The simulated light yields 162 with respect to bis-MSB concentrations are compared with the experimental data, as shown in Fig. 7. If the absorption 163 spectra of Daya Bay liquids are used, a much steeper ObsLY increase is found, because LAB, which does not re-emit 164 a new photon after the absorption, strongly competes with bis-MSB in the wavelength range of 350 to 400 nm. More 165 bis-MSB leads to more photons shifting to the wavelength range above 420 nm, in which the liquids are much more 166 transparent. Thus, the ObsLY monotonically increases with the bis-MSB concentration. Once the absorption spectra 167 of JUNO liquids are employed, the importance of bis-MSB is significantly reduced, and most of the photons emitted 168

- ¹⁶⁹ by PPO could reach the Daya Bay LS contained in the outer acrylic vessel before absorption by LAB. However, at
- ¹⁷⁰ bis-MSB concentrations of smaller than 1 mg/L, the bis-MSB plays a less important role in the data compared to the
- simulation. Varying the absorption spectra of each component at wavelengths longer than 420 nm does not reduce
- the discrepancy. Changing the height of the bis-MSB QE spectrum, and shifting the cutoff position of the PPO and
- ¹⁷³ bis-MSB QE spectra have minor impacts on the discrepancy. Eventually, the PPO fluorescence QE is increased by ¹⁷⁴ 5% as shown in Fig. 5. Once a photon is absorbed by PPO, the probability of re-emitting a new photon with longer
- wavelength is closer to 1. In this way, the discrepancy between simulation and data is improved from about 2% to
- ¹⁷⁶ better than 1%.



Figure 7: Comparison of the measured and simulated light yields with 2 g/L PPO and various bis-MSB concentrations. Each group is normalized at 1 mg/L bis-MSB. The JUNO LS data in the Daya Bay detector are drawn as the black dots. If the optical properties of Daya Bay LS are used in the simulation, large discrepancies are found as the black squares. Using the default optical properties of JUNO LS in Fig. 5 and Fig. 6, the simulation results (blue inverted triangles) have a much better agreement with data. After increasing the PPO fluorescence quantum efficiencies, the agreement is improved from 2% to 1% (red triangles) at low bis-MSB concentrations. For better visibility, the triangles and inverted triangles are slightly shifted to the right and the left, respectively.

The first application of the tuned model is to extract the initial light yields with different PPO concentrations. As mentioned before, ObsLY is a joint effect of the initial light yield, the absorption and re-emission, and the PMT response. A set of simulation is performed for the Daya Bay AD with LS compositions with different PPO concentrations and 7 mg/L bis-MSB. The same initial light yield is used in the simulation to solely study the selfabsorption effect of PPO. The simulation results are listed in Table 2. Each 0.5 g/L PPO increase leads to about 0.5% loss of ObsLY in the simulation. Thus, the initial light yields are obtained by dividing the Daya Bay measured ObsLY with the simulated ones, and will be used in the determination of the JUNO LS composition.

PPO concentration	ObsLY		Initial light world
FFO concentration	Measured	Simulated	finnañ figint yield
2.0 g/L	1	1	1
2.5 g/L	1.025	0.994	1.031
3.0 g/L	1.037	0.991	1.046
3.5 g/L	1.042	0.986	1.057
4.0 g/L	1.042	0.982	1.061

Table 2: Relative ObsLY with respect to PPO concentrations in the data and the simulation, normalized at 2 g/L PPO. In the simulation the initial light yield is fixed to solely study the PPO self-absorption effects. The initial light yields are extracted by dividing the measured values with the simulated ones.

The optical model, the measured absorption spectra, the tuned fluorescence quantum efficiencies, and the extracted initial light yields have been employed in the JUNO simulation. The simulated ObsLY with respect to PPO

- and bis-MSB concentrations is shown in Fig. 8. The normalization point is chosen as 2.5 g/L PPO and 2 mg/L bis-186
- MSB, at which the largest light yield is found. Although the ObsLY monotonically increases with PPO concentrations 187
- at Daya Bay, 2.5 g/L PPO is preferred at JUNO due to the non-negligible self-absorption in the larger detector. The 188
- optimal bis-MSB concentration could be in the range of 1 mg/L to 4 mg/L, since the ObsLY difference is less than 1% 189
- in this range. Combining the absorption of each component, and the scattering of LAB, the LS attenuation length with 190 this composition is about 20.4 m at 430 nm. It fulfills the requirement in the JUNO Conceptual Design Report [5].
- 191



Figure 8: The simulated ObsLY with respect to PPO and bis-MSB concentrations in the JUNO detector. The tuned optical parameters, and the initial light yields in Table 2 are used in the simulation. For better visibility, the bis-MSB concentrations are divided to two groups which are shown in the top and bottom panel, respectively

5. Summary 192

A precise measurement of ObsLY in various LS compositions has been performed in a Daya Bay AD, by replacing 193 the Gd-LS with purified LS produced in a pilot plant in the underground LS hall. For γ 's and e^{\pm} 's, ObsLY reaches a 194 plateau for PPO concentrations larger than 2.5 g/L. In addition, if the solvent is optically purified, the ObsLY increase 195 with respect to bis-MSB concentrations is negligible for concentrations larger than 4 mg/L. A novel optical model has 196 been employed to describe the complicated optical process in the liquids. The predicted ObsLY in different bis-MSB 197 concentrations agrees with these measurements within 1%. The initial light yields in various PPO concentrations 198 are extracted by subtracting the PPO self-absorption effect predicted by the optical model. To find the optimal LS 199 composition of JUNO, the model and the tuned optical parameters are used in the JUNO simulation. The results 200 suggest that in the JUNO-scale detector, either a PPO concentration larger than 2.5 g/L, or a bis-MSB concentration 201 bigger than 4 mg/L, would reduce the ObsLY. The JUNO LS composition is optimized to be the purified LAB with 202 2.5 g/L PPO and (1-4) mg/L bis-MSB. The optimization method can also be used in other future LS experiments. 203

6. Acknowledgements 204

We are grateful for the ongoing cooperation from the China General Nuclear Power Group and China Light and 205 Power Company. 206

Daya Bay is supported in part by the Ministry of Science and Technology of China, the U.S. Department of Energy, the Chinese Academy of Sciences, the CAS Center for Excellence in Particle Physics, the National Natural

209 Science Foundation of China, the Guangdong provincial government, the Shenzhen municipal government, the China

210 General Nuclear Power Group, the Research Grants Council of the Hong Kong Special Administrative Region of

²¹¹ China, the MOE in Taiwan, the U.S. National Science Foundation, the Ministry of Education, Youth, and Sports of

the Czech Republic, the Charles University Research Centre UNCE, the Joint Institute of Nuclear Research in Dubna,

Russia, the National Commission of Scientific and Technological Research of Chile, We acknowledge Yellow River

Engineering Consulting Co., Ltd., and China Railway 15th Bureau Group Co., Ltd., for building the underground laboratory.

JUNO is supported by the Chinese Academy of Sciences, the National Key R&D Program of China, the CAS Center for Excellence in Particle Physics, the Joint Large-Scale Scientific Facility Funds of the NSFC and CAS,

Wuyi University, and the Tsung-Dao Lee Institute of Shanghai Jiao Tong University in China, the Institut National de
 Physique Nucléaire et de Physique de Particules (IN2P3) in France, the Istituto Nazionale di Fisica Nucleare (INFN)

²²⁰ in Italy, the Fond de la Recherche Scientifique (F.R.S-FNRS) and FWO under the "Excellence of Science C EOS in

221 Belgium, the Conselho Nacional de Desenvolvimento Científico e Tecnològico in Brazil, the Agencia Nacional de

²²² Investigación y Desarrollo in Chile, the Charles University Research Centre and the Ministry of Education, Youth,

and Sports in Czech Republic, the Deutsche Forschungsgemeinschaft (DFG), the Helmholtz Association, and the

²²⁴ Cluster of Excellence PRISMA+ in Germany, the Joint Institute of Nuclear Research (JINR), Lomonosov Moscow

225 State University, and Russian Foundation for Basic Research (RFBR) in Russia, the MOST and MOE in Taiwan, the

226 Chulalongkorn University and Suranaree University of Technology in Thailand, and the University of California at

- 227 Irvine in USA.
- [1] C. L. Cowan, F. Reines, F. B. Harrison, H. W. Kruse, A. D. McGuire, Detection of the free neutrino: A Confirmation, Science 124 (1956)
 103–104. doi:10.1126/science.124.3212.103.
- [2] S. Abe, et al., Precision Measurement of Neutrino Oscillation Parameters with KamLAND, Phys. Rev. Lett. 100 (2008) 221803. arXiv: 0801.4589, doi:10.1103/PhysRevLett.100.221803.

[3] C. Arpesella, et al., Direct Measurement of the Be-7 Solar Neutrino Flux with 192 Days of Borexino Data, Phys. Rev. Lett. 101 (2008) 091302.
 arXiv:0805.3843, doi:10.1103/PhysRevLett.101.091302.

[4] F. P. An, et al., Observation of electron-antineutrino disappearance at Daya Bay, Phys. Rev. Lett. 108 (2012) 171803. arXiv:1203.1669, doi:10.1103/PhysRevLett.108.171803.

- [5] Z. Djurcic, et al., JUNO Conceptual Design ReportarXiv:1508.07166.
- [6] F. An, et al., Neutrino Physics with JUNO, J. Phys. G43 (3) (2016) 030401. arXiv:1507.05613, doi:10.1088/0954-3899/43/3/
 030401.
- [7] P. Lombardi, F. Ortica, G. Ranucci, A. Romani, Decay time and pulse shape discrimination of liquid scintillators based on novel solvents, Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment 701 (2013) 133 - 144. doi:https://doi.org/10.1016/j.nima.2012.10.061.
- 242 URL http://www.sciencedirect.com/science/article/pii/S0168900212012077
- [8] X.-C. Ye, et al., Preliminary study of light yield dependence on LAB liquid scintillator composition, Chin. Phys. C39 (9) (2015) 096003.
 arXiv:1506.00237, doi:10.1088/1674-1137/39/9/096003.
- [9] J. Cumming, S. Hans, M. Yeh, Improving Light Yield Measurements for Low-Yield Scintillators, Nucl. Instrum. Meth. A 925 (2019) 1–5.
 arXiv:1810.02885, doi:10.1016/j.nima.2019.01.014.
- [10] X.-B. Li, H.-L. Xiao, J. Cao, J. Li, X.-C. Ruan, Y.-K. Heng, Timing properties and pulse shape discrimination of LAB-based liquid scintillator, Chinese Physics C 35 (11) (2011) 1026–1032. doi:10.1088/1674-1137/35/11/009.
- 249 URL https://doi.org/10.1088%2F1674-1137%2F35%2F11%2F009
- [11] X.-F. Ding, L.-J. Wen, X. Zhou, Y.-Y. Ding, X.-C. Ye, L. Zhou, M.-C. Liu, H. Cai, J. Cao, Measurement of the fluorescence quantum yield of
 bis-MSB, Chin. Phys. C 39 (12) (2015) 126001. arXiv:1506.00240, doi:10.1088/1674-1137/39/12/126001.

[12] C. Buck, B. Gramlich, S. Wagner, Light propagation and fluorescence quantum yields in liquid scintillators, JINST 10 (09) (2015) P09007.
 arXiv:1509.02327, doi:10.1088/1748-0221/10/09/P09007.

Y. Zhang, Z. Yu, X. Li, Z. Deng, L. Wen, A complete optical model for liquid-scintillator detectors, Nuclear Instruments and Methods in
 Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment 967 (2020) 163860. doi:https://doi.
 org/10.1016/j.nima.2020.163860.

257 URL http://www.sciencedirect.com/science/article/pii/S0168900220303648

[14] F. P. An, et al., The Detector System of The Daya Bay Reactor Neutrino Experiment, Nucl. Instrum. Meth. A811 (2016) 133–161. arXiv: 1508.03943, doi:10.1016/j.nima.2015.11.144.

[15] H. Band, et al., Assembly and Installation of the Daya Bay Antineutrino Detectors, JINST 8 (2013) T11006. arXiv:1309.1557, doi:
 10.1088/1748-0221/8/11/T11006.

[16] D. Adey, et al., Measurement of the Electron Antineutrino Oscillation with 1958 Days of Operation at Daya Bay, Phys. Rev. Lett. 121 (24)
 (2018) 241805. arXiv:1809.02261, doi:10.1103/PhysRevLett.121.241805.

[17] F. P. An, et al., Improved Measurement of the Reactor Antineutrino Flux and Spectrum at Daya Bay, Chin. Phys. C41 (1) (2017) 013002.
 arXiv:1607.05378, doi:10.1088/1674-1137/41/1/013002.

[18] F. P. An, et al., Evolution of the Reactor Antineutrino Flux and Spectrum at Daya Bay, Phys. Rev. Lett. 118 (25) (2017) 251801. arXiv:

- ²⁶⁷ 1704.01082, doi:10.1103/PhysRevLett.118.251801.
- [19] D. Adey, et al., Extraction of the ²³⁵U and ²³⁹Pu Antineutrino Spectra at Daya Bay, Phys. Rev. Lett. 123 (11) (2019) 111801. arXiv:

- 269 1904.07812, doi:10.1103/PhysRevLett.123.111801.
- [20] F. P. An, et al., Improved Search for a Light Sterile Neutrino with the Full Configuration of the Daya Bay Experiment, Phys. Rev. Lett. 117 (15)
 (2016) 151802. arXiv:1607.01174, doi:10.1103/PhysRevLett.117.151802.
- [21] P. Adamson, et al., Improved Constraints on Sterile Neutrino Mixing from Disappearance Searches in the MINOS, MINOS+, Daya Bay, and
 Bugey-3 ExperimentsarXiv: 2002.00301.
- [22] F. An, et al., The muon system of the Daya Bay Reactor antineutrino experiment, Nucl. Instrum. Meth. A 773 (2015) 8–20. arXiv:1407.
 0275, doi:10.1016/j.nima.2014.09.070.
- [23] P. Lombardi, et al., Distillation and stripping pilot plants for the JUNO neutrino detector: Design, operations and reliability, Nucl. Instrum.
 Meth. A925 (2019) 6–17. arXiv:1902.05288, doi:10.1016/j.nima.2019.01.071.
- [24] Q.-j. Li, Z. Wang, J.-f. Lv, X.-b. Yan, W.-q. Jiang, J.-f. Chang, Front-End electronics system of pmt readout for daya bay reactor Neutrino
 Experiment, in: Proceedings, 2009 IEEE Nuclear Science Symposium and Medical Imaging Conference (NSS/MIC 2009): Orlando, Florida,
 October 25-31, 2009, 2009, pp. 1821–1823. doi:10.1109/NSSMIC.2009.5402197.
- [25] D. Adey, et al., A high precision calibration of the nonlinear energy response at Daya Bay, Nucl. Instrum. Meth. A940 (2019) 230-242.
 arXiv:1902.08241, doi:10.1016/j.nima.2019.06.031.
- [26] D.-M. Xia, B.-X. Yu, X.-B. Li, X.-L. Sun, Y.-Y. Ding, L. Zhou, J. Cao, W. Hu, X.-C. Ye, H.-T. Chen, X.-F. Ding, B. Du, Temperature dependence of the light yield of the LAB-based and mesitylene-based liquid scintillators, Chinese Physics C 38 (11) (2014) 116001. doi: 10.1088/1674-1137/38/11/116001.
- 286 URL https://doi.org/10.1088%2F1674-1137%2F38%2F11%2F116001
- [27] X. Zhou, Q. Liu, M. Wurm, Q. Zhang, Y. Ding, Z. Zhang, Y. Zheng, L. Zhou, J. Cao, Y. Wang, Rayleigh scattering of linear alkylbenzene in
- large liquid scintillator detectors, Rev. Sci. Instrum. 86 (7) (2015) 073310. arXiv:1504.00987, doi:10.1063/1.4927458.
- [28] S. Agostinelli, et al., GEANT4: A Simulation toolkit, Nucl. Instrum. Meth. A506 (2003) 250–303. doi:10.1016/S0168-9002(03)
 01368-8.

Appendix A. Author list

A. Abusleme^{δ,1}, T. Adam^{δ,2}, S. Ahmad^{δ,3}, S. Aiello^{δ,4}, M. Akram^{δ,3}, N. Ali^{δ,3}, F. P. An^{δ,4,5}, G. P. An^{δ,6}, O. An^{δ,7}, G. Andronico^{δ,4} 292 N. Anfimov^{δ 8}, V. Antonelli^{δ 9}, T. Antoshkina^{δ 8}, B. Asavapibhop^{δ ,10}, J. P. A. M. de André^{δ 2}, A. Babic^{δ ,11}, A. B. Balantekin^{μ ,12}, W. Baldini^{δ ,13}, M. Baldoncini^{δ ,13}, H. R. Band^{μ ,14}, A. Barresi^{δ ,15}, E. Baussan^{δ 2}, M. Bellato^{δ ,16}, E. Bernieri^{δ ,17}, D. Biare^{δ 8}, 293 294 T. Birkenfeld^{δ ,18}, M. Bishai^{μ ,19}, S. Blin^{δ ,20}, D. Blum^{δ ,21}, S. Blyth^{δ , μ ,22}, C. Bordereau^{δ ,23,24}, A. Brigatti^{δ ,9}, R. Brugnera^{δ ,25}, 295 A. Budano^{6,17}, P. Burgbacher^{6,21}, M. Buscemi^{6,4}, S. Bussino^{6,17}, J. Busto^{6,26}, I. Butorov^{6,8}, A. Cabrera^{6,20}, H. Cai^{6,27}, X. Cai^{6,6} 296 Y. K. Cai^{δ,6}, Z. Y. Cai^{δ,6}, A. Cammi^{δ,28}, A. Campeny^{δ,1}, C. Y. Cao^{δ,6}, G. F. Cao^{δ,μ,6}, J. Cao^{δ,μ,6}, R. Caruso^{δ,4}, C. Cerna^{δ,23} 297 I. Chakaberia^{δ ,29}, J. F. Chang^{δ , μ ,6}, Y. Chang^{δ , μ ,24}, H. S. Chen^{μ ,6}, P. A. Chen^{δ ,22}, P. P. Chen^{δ ,30}, S. M. Chen^{δ , μ ,31}, S. J. Chen^{δ ,32}, X. R. Chen^{δ ,33}, Y. W. Chen^{δ ,34}, Y. X. Chen^{δ , μ ,35}, Y. Chen^{δ , μ ,36}, Z. Chen^{δ ,6}, J. Cheng^{δ , μ ,6}, Y. P. Cheng^{δ , μ ,37}, Z. K. Cheng^{μ ,36}, 298 299 A. Chepurnov^{6,38}, J. J. Cherwinka^{*μ*,12}, F. Chiarello^{6,16}, D. Chiesa^{6,15}, P. Chimenti^{6,39}, M. C. Chu^{*μ*,40}, A. Chukanov^{6,8}, A. Chuvashova^{6,8} 300 . Clementi^{6,41}, B. Clerbaux^{6,42}, S. Conforti Di Lorenzo^{6,20}, D. Corti^{6,16}, S. Costa^{6,4}, F. D. Corso^{6,16}, J. P. Cummings^{4,43}, O. Dalager^{4,44}, 301 C. De La Taille^{6,20}, F. S. Deng^{4,7}, J. W. Deng^{6,27}, Z. Deng^{6,31}, Z. Y. Deng^{6,6}, W. Depnering^{6,45}, M. Diaz^{6,1}, X. F. Ding^{6,9} 302 Y. Y. Ding^{6,µ,6}, B. Dirgantara^{6,46}, S. Dmitrievsky^{6,8}, M. V. Diwan^{µ,19}, T. Dohnal^{6,µ,47}, G. Donchenko^{6,38}, J. M. Dong^{6,31}, D. Dornic^{6,26}, 303 E. Doroshkevich^{δ,48}, J. Dove^{μ,49}, M. Dracos^{δ,2}, F. Druillole^{δ,23}, S. X. Du^{δ,50}, S. Dusini^{δ,16}, M. Dvorak^{δ,μ,47}, D. A. Dwyet^{μ,51}, 304 T. Enqvist^{6,52}, H. Enzmann^{6,45}, A. Fabbri^{6,17}, L. Fajt^{6,11}, D. H. Fan^{6,53}, L. Fan^{6,6}, C. Fang^{6,54}, J. Fang^{6,6}, A. Fatkina^{6,8}, D. Fedoseev^{6,8} 305 V. Fekete^{δ ,11}, L. C. Feng^{δ ,34}, Q. C. Feng^{δ ,55}, G. Fiorentini^{δ ,13}, R. Ford^{δ ,9}, A. Formozov^{δ ,9}, A. Fournier^{δ ,23}, S. Franke^{δ ,56}, J. P. Gallo^{μ ,57}, H. N. Gan^{δ ,58}, F. Gao^{δ ,18}, A. Garfagnini^{δ ,25}, A. Göttel^{δ ,37,18}, C. Genster^{δ ,37}, M. Giammarchi^{δ ,9}, A. Giaz^{δ ,25}, N. Giudice^{δ ,4}, 306 307 F. Giuliani^{6,59}, M. Gonchar^{6,8}, G. H. Gong^{6,µ,31}, H. Gong^{6,µ,31}, O. Gorchakov^{6,8}, Y. Gornushkin^{6,8}, M. Grassi^{6,25}, C. Grewing^{6,60}, 308 P. Gullandez, M. Goldenalez, G. H. Goldenez, H. Goldenez, O. Goldadovez, I. Goldushinez, M. Glassiew, C. Glewingers, M. Gromov^{6,38}, V. Gromov^{6,8}, M. H. Gu^{6,6}, W. Q. Gu^{μ ,19}, X. F. Gu^{δ ,50}, Y. Gu^{δ ,61}, M. Y. Guan^{δ ,6}, N. Guardone^{δ ,4}, M. Gul^{δ ,3}, C. Guo^{δ ,6}, J. Y. Guo^{$\delta\mu$,36}, L. Guo^{μ ,31}, W. L. Guo^{δ ,66}, X. H. Guo^{$\delta\mu$,62}, Y. H. Guo^{$\delta\mu$,63,37}, Z. Guo^{μ ,31}, M. Haacke^{δ ,1}, R. W. Hackenburg^{μ ,19}, P. Hackspacher^{δ ,45}, C. Hagner^{δ ,64}, R. Han^{δ ,65}, Y. Han^{δ ,20}, S. Hans^{μ ,191}, M. He^{$\delta\mu$,66}, W. He^{δ ,6}, K. M. Heeger^{μ ,14}, T. Heinz^{δ ,21}, Y. K. Heng^{$\delta\mu$,6}, R. Herrera^{δ ,1}, A. Higuera^{μ ,66}, D. J. Hong^{δ ,54}, Y. K. Hor^{μ ,36}, S. J. Hou^{δ ,6}, Y. B. Hsiung^{$\delta\mu$,22}, B. Z. Hu^{$\delta\mu$,22}, H. Hu^{δ ,36}, J. R. Hu^{$\delta\mu$,6}, J. Hu^{δ ,6}, S. Y. Hu^{δ ,67}, T. Hu^{$\delta\mu$,6}, Z. J. Hu^{$\delta\mu$,36}, C. H. Huang^{δ ,36}, G. H. Huang^{$\delta\mu$,6}, H. X. Huang^{$\delta\mu$,67}, Q. H. Huang^{δ ,2}, W. H. Huang^{δ ,29}, X. T. Huang^{$\delta\mu$,29}, Y. B. Huang^{$\delta\mu$,654}, P. Huber^{μ ,68}, J. Q. Hui^{δ ,59}, L. Huo^{$\delta\mu$,54}, Y. L. Ian^{$\delta\mu$,69}</sub> 309 310 311 312 313 314 C. Huss^{6,23}, S. Hussain^{6,3}, A. Insolia^{6,4}, A. Ioannisian^{6,69}, D. Ioannisyan^{6,69}, R. Isocrate^{6,16}, D. E. Jaffe^{4,19}, K. L. Jen^{6,4,34}, X. L. Ji^{6,4,6} 315 X. P. Ji^{µ,19}, X. Z. Ji^{6,36}, H. H. Jia^{6,70}, J. J. Jia^{6,27}, S. Y. Jian^{6,67}, D. Jiang^{6,7}, X. S. Jiang^{6,6}, R. Y. Jin^{6,6}, X. P. Jing^{6,6}, R. A. Johnson^{µ,71}, 316 C. Jollet^{6,23}, D. Jones^{4,72}, J. Joutsenvaara^{6,52}, S. Jungthawan^{6,46}, L. Kalousis^{6,2}, P. Kampmann^{6,37,18}, L. Kang^{6,4,30}, M. Karagounis^{6,60}, 317 N. Kazarian^{δ,69}, S. H. Kettell^{µ,19}, A. Khan^{δ,36}, W. Khan^{δ,63}, K. Khosonthongkee^{δ,46}, P. Kinz^{δ,34}, S. Kohn^{µ,73}, D. Korablev^{δ,8}, K. Kouzakov^{δ,38}, M. Kramer^{µ,51,73}, A. Krasnoperov^{δ,8}, S. Krokhaleva^{δ,48}, Z. Krumshteyn^{δ,8}, A. Kruth^{δ,60}, N. Kutovskiy^{δ,8} 318 319 P. Kuusiniemi^{6,52}, B. Lachacinski^{6,23}, T. Lachenmaier^{6,21}, C. Landini^{6,9}, T. J. Langford^{µ,14}, J. Lee^{µ,51}, J. H. C. Lee^{µ,74}, F. Lefevre^{6,75}. 320 L. Lei $^{\delta,31}$, R. Lei $^{\delta,\mu,30}$, R. Leitner $^{\delta,\mu,47}$, 4]J. Leung $^{\delta,\mu,74}$, C. Li $^{\delta,29}$, D. M. Li $^{\delta,50}$, F. Li $^{\delta,46}$, F. Li $^{\delta,31}$, H. T. Li $^{\delta,36}$, H. L. Li $^{\delta,66}$, F. Li $^{\delta,41}$, H. T. Li $^{\delta,36}$, H. L. Li $^{\delta,66}$, F. Li $^{\delta,41}$, H. T. Li $^{\delta,46}$, H. L. Li $^{\delta,66}$, F. Li $^{\delta,41}$, H. T. Li $^{\delta,46}$, H. L. Li $^{\delta,41}$, H. T. Li $^{\delta,41}$, H. T. Li $^{\delta,41}$, H. T. Li $^{\delta,46}$, H. L. Li $^{\delta,46}$, H. L. Li $^{\delta,41}$, H. T. Li{^{\delta,41}}, H. T. Li $^{\delta,41}$, H. T. Li{^{\delta,41}}, H. 321 J. Li^{δ,6}, J. J. Li^{μ,31}, J. Q. Li^{δ,36}, K. J. Li^{δ,36}, M. Z. Li^{δ,6}, N. Li^{δ,76}, N. Li^{δ,6}, Q. J. Li^{δ,76}, Q. J. Li^{μ,6}, R. H. Li^{δ,6}, S. C. Li^{μ,68} 322 S. F. Li^{δ,μ,30}, S. J. Li^{δ,36}, T. Li^{δ,36}, T. Li^{δ,29}, W. D. Li^{δ,μ,6}, W. G. Li^{δ,6}, X. M. Li^{δ,67}, X. N. Li^{δ,μ,6}, X. L. Li^{δ,67}, X. Q. Li^{μ,70}, Y. Li^{δ,30} 323 Y. F. Li^{6,µ,6}, Z. B. Li^{6,µ,56}, Z. Y. Li^{6,36}, H. Liang^{6,67}, H. Liang^{6,47}, J. J. Liang^{6,54}, D. Liebau^{6,60}, A. Limphirat^{6,46}, S. Limpijumnong^{6,46} 324

¹Now at: Department of Chemistry and Chemical Technology, Bronx Community College, Bronx, New York 10453, USA

C. J. Lin^{μ ,51}, G. L. Lin^{$\delta\mu$,34}, S. X. Lin^{$\delta\mu$,30}, T. Lin^{$\delta\delta$,} Y. H. Lin^{$\delta34$}, J. J. Ling^{$\delta\mu$,36}, J. M. Link^{μ ,68}, I. Lippi^{δ ,16}, L. Littenberg^{μ ,19}, 325 B. R. Littlejohn^{4,57}, F. Liu^{6,35}, H. Liu^{6,36}, H. Liu^{6,61}, H. B. Liu^{6,54}, H. D. Liu^{6,50}, H. J. Liu^{6,77}, H. T. Liu^{6,36}, J. C. Liu^{6,4,6}, J. L. Liu^{6,4,59,78}. 326 M. Liu^{6,77}, Q. Liu^{6,79}, Q. Liu^{6,7}, R. X. Liu^{6,6}, S. Y. Liu^{6,6}, S. B. Liu^{6,7}, S. L. Liu^{6,6}, X. W. Liu^{6,36}, Y. Liu^{6,6}, A. Lokhov^{6,38}. 327 P. Lombardi^{δ ,9}, K. Loo^{δ ,52}, S. Lorenz^{δ ,45}, C. Lu^{μ ,80}, C. Lu^{δ ,58}, H. Q. Lu^{δ ,46}, J. B. Lu^{δ ,81}, J. G. Lu^{δ ,6}, S. X. Lu^{δ ,50}, X. X. Lu^{δ ,6}, 328 B. Lubsandorzhiev^{6,48}, S. Lubsandorzhiev^{6,48}, L. Ludhova^{6,37,18}, K. B. Luk^{4,73,51}, F. J. Luo^{6,6}, G. Luo^{6,36}, P. W. Luo^{6,36}, S. Luo^{6,82}, 329 W. M. Luo^{6,6}, V. Lyashuk^{6,48}, Q. M. Ma^{6,6}, S. Ma^{6,6}, X. B. Ma^{6,µ,35}, X. Y. Ma^{6,µ,6}, Y. Q. Ma^{µ,6}, Y. Malyshkin^{6,17}, F. Mantovani^{6,13}, 330 Y. J. Mao^{6,83}, S. M. Mari^{6,17}, F. Marini^{6,25}, S. Marium^{6,3}, C. Marshall^{4,51}, C. Martellini^{6,17}, G. Martin-Chassard^{6,20}, D. A. Martinez 331 Caicedo^{4,57}, A. Martini^{6,84}, J. Martino^{6,75}, D. Mayilyan^{6,69}, K. T. McDonald^{4,80}, R. D. McKeown^{4,85,86}, A. Müller^{6,21}, G. Meng^{6,16} 332 Y. Meng^{δ,μ,59}, A. Meregaglia^{6,23}, E. Meroni^{δ,9}, D. Meyhöfer^{δ,64}, M. Mezzetto^{δ,16}, J. Miller^{δ,87}, L. Miramonti^{δ,9}, S. Monforte^{δ,4} 333 P. Montini^{6,17}, M. Montuschi^{6,13}, N. Morozov^{6,8}, P. Muralidharan^{6,60}, J. Napolitano^{4,72}, M. Nastasi^{6,15}, D. V. Naumov^{6,4,8}. 334 E. Naumova^{6,4,8}, I. Nemchenok^{6,8}, A. Nikolaev^{6,38}, F. P. Ning^{6,6}, Z. Ning^{6,6}, H. Nunokawa^{6,88}, L. Oberauer^{6,56}, J. P. Ochoa-335 Ricoux^{6,µ,44,1}, A. Olshevskiy^{6,µ,8}, F. Ortica^{6,41}, H. R. Pan^{6,µ,22}, A. Paoloni^{6,84}, J. Park^{µ,68}, N. Parkalian^{6,60}, S. Parmeggiano^{6,9} 336 S. Patton^{4,51}, T. Payupol^{6,10}, V. Pec^{6,47}, D. Pedretti^{6,25}, Y. T. Pei^{6,6}, N. Pelliccia^{6,41}, A. G. Peng^{6,77}, H. P. Peng^{6,7}, J. C. Peng^{4,49} 337 F. Perrot^{6,23}, P. A. Petitjean^{6,42}, L. F. Pineres Rico^{6,39}, A. Popov^{6,38}, P. Poussot^{6,2}, W. Pratumwan^{6,46}, E. Previtali^{6,15}, C. S. J. Pun^{4,74} 338 F. Z. Qi^{δ,μ,6}, M. Qi^{δ,μ,32}, S. Qian^{δ,6}, X. Qian^{μ,19}, X. H. Qian^{δ,6}, H. Qiao^{δ,83}, Z. H. Qin^{δ,6}, S. K. Qiu^{δ,77}, M. Rajput^{δ,3}, G. Ranucci^{δ,9}, 339 N. Raper^{$\delta\mu$,36}, A. Re^{$\delta,9$}, H. Rebber^{$\delta,64$}, A. Rebii^{$\delta,23$}, B. Ren^{$\delta,30$}, J. Ren^{$\delta\mu$,67}, C. M. Reveco^{μ ,44}, T. Rezinko^{δ 8}, B. Ricci^{$\delta,13$}, M. Robens^{$\delta,60$}, M. Roche^{$\delta,23$}, N. Rodphai^{$\delta,10$}, L. Rohwer^{$\delta,64$}, A. Romani^{$\delta,41$}, R. Rosero^{μ ,19}, B. Roskovec^{$\delta\mu$,44}, C. Roth^{$\delta,60$}, X. C. Ruan^{$\delta\mu$,67}, X. D. Ruan^{$\delta,54$}, S. Rujirawat^{$\delta,46$}, A. Rybnikov^{$\delta,8$}, A. Sadovsky^{$\delta,8$}, P. Saggese^{$\delta,9}$ </sup>, G. Salamanna^{$\delta,17$}, A. Sangka^{$\delta,89$}, N. Sanguansak^{$\delta,46$}, 340 341 342 U. Sawangwit^{6,89}, J. Sawatzki^{6,56}, F. Sawy^{6,25}, M. Schever^{6,37,18}, J. Schulet^{6,2}, C. Schwab^{6,2}, K. Schweizer^{6,56}, D. Selivanov^{6,8} 343 A. Selyunin^{6,8}, A. Serafini^{6,13}, G. Settanta^{6,17}, M. Settimo^{6,75}, M. Shahzad^{6,3}, G. Shi^{6,31}, J. Y. Shi^{6,6}, Y. J. Shi^{6,31}, V. Shutov^{6,8} 344 A. Sidorenkov^{6,48}, F. Simkovic^{6,11}, C. Sirignano^{6,25}, J. Siripak^{6,46}, M. Sisti^{6,15}, M. Slupecki^{6,52}, M. Smirnov^{6,36}, O. Smirnov^{6,8} 345 T. Sogo-Bezerra^{6,75}, J. Songwadhana^{6,46}, B. Soonthornthum^{6,89}, A. Sotnikov^{6,8}, O. Sramek^{6,47}, W. Sreethawong^{6,46}, A. Stahl^{6,18}, 346 L. Stanco^{6,16}, K. Stankevich^{6,38}, D. Stefanik^{6,11}, H. Steiger^{6,56}, H. Steiner^{4,73,51}, J. Steinmann^{6,18}, M. Stender^{6,64}, V. Strati^{6,13}, 347 A. Studenikin^{6,38}, G. X. Sun^{6,6}, L. T. Sun^{6,6}, J. L. Sun^{4,90}, S. F. Sun^{6,35}, X. L. Sun^{6,6}, Y. J. Sun^{6,7}, Y. Z. Sun^{6,6}, N. Suwonjandee^{6,10}, 348 M. Szelezniak^{6,2}, J. Tang^{6,36}, Q. Tang^{6,36}, Q. Tang^{6,77}, X. Tang^{6,6}, A. Tietzsch^{6,21}, I. Tkachev^{6,48}, T. Tmej^{µ,47}, K. Treskov^{6,µ,8} 349 G. Troni^{*δ*,1}, W. Trzaska^{*δ*,52}, W.-H. Tse^{*μ*,40}, C. E. Tull^{*μ*,51}, C. Tuve^{*δ*,4}, S. van Waasen^{*δ*,60}, J. Vanden Boom^{*δ*,60}, N. Vassilopoulos^{*δ*,6}, 350 V. Vedin^{6,91}, G. Verde^{6,4}, M. Vialkov^{6,38}, B. Viaud^{6,75}, B. Viren^{µ,19}, C. Volpe^{6,20}, V. Vorobel^{6,µ,47}, L. Votano^{6,84}, P. Walker^{6,1}, 351 C. Wang^{6,30}, C. H. Wang^{6,4,24}, E. Wang^{6,50}, G. L. Wang^{6,55}, J. Wang^{6,7}, J. Wang^{6,4,36}, K. Y. Wang^{6,6}, L. Wang^{6,6}, M. F. Wang^{6,6} 352 M. Wang^{$\delta,77$}, M. Wang^{$\delta,\mu,29$}, N. Y. Wang^{$\mu,62$}, R. G. Wang^{$\delta,\mu,6$}, S. G. Wang^{$\delta,83$}, W. Wang^{$\delta,\mu,36$}, W. Wang^{$\delta,\mu,32$}, W. S. Wang^{$\delta,0,02$} 353 X. Wang^{δ,μ,76}, X. Y. Wang^{δ,36}, Y. Wang^{δ,31}, Y. Wang^{μ,32}, Y. Wang^{δ,53}, Y. F. Wang^{δ,6}, Y. G. Wang^{δ,27}, Y. M. Wang^{δ,32}, Y. Q. Wang^{δ,31} 354 Z. Wang^{6,\mu,31}, Z. Wang^{6,\mu,6}, Z. M. Wang^{6,\mu,6}, Z. Y. Wang^{6,31}, A. Watcharangkool^{6,89}, H. Y. Wei^{µ,19}, L. H. Wei^{6,µ,6}, W. Wei^{6,6} 355 Y. D. Wei^{6,30}, L. J. Wen^{6,4,6}, K. Whisnant^{4,92}, C. G. White^{4,57}, C. Wiebusch^{6,18}, S. C. F. Wong^{6,36}, H. L. H. Wong^{4,73,51}, B. Wonsak^{6,64} 356 E. Worcester^{μ ,19}, C. H. Wu^{δ ,34}, D. R. Wu^{δ , μ ,6}, F. L. Wu^{δ , μ ,32}, Q. Wu^{δ , μ ,29}, W. J. Wu^{δ , μ ,27}, Z. Wu^{δ ,6}, M. Wurm^{δ ,45}, J. Wurtz^{δ ,2} 357 C. Wysotzki^{δ,18}, Y. F. Xi^{δ,58}, D. M. Xia^{δμ,93}, Y. G. Xie^{δ,6}, Z. Q. Xie^{δμ,6}, Z. Z. Xing^{δμ,6}, D. L. Xu^{δ,78,59}, F. R. Xu^{δ,61}, H. K. Xu^{δμ,6} 358 J. L. Xu^{δ,μ,6}, J. Xu^{δ,62}, M. H. Xu^{δ,6}, T. Xu^{μ,31}, Y. Xu^{δ,70}, Y. Xu^{δ,37,18}, T. Xue^{μ,31}, B. J. Yan^{δ,6}, X. B. Yan^{δ,6}, Y. P. Yan^{δ,46}, A. B. Yang^{δ,6} 359 C. G. Yang^{\delta,\mu,6}, H. Yang^{\delta,6}, J. Yang^{\delta,50}, L. Yang^{\delta,4,30}, X. Y. Yang^{\delta,6}, Y. F. Yang^{\delta,42}, Y. Z. Yang^{\mu,31}, H. F. Yao^{\delta,\mu,6}, Z. Yasin^{\delta,3}, J. X. Ye^{\delta,6}, 360 C. O. Falig *, H. Falig *, J. Falig *, E. Falig *, A. H. Yuág *, H. Falig *, 361 362 363 G. Q. Zhang^{6,6}, H. H. Zhang^{6,4,36}, H. Q. Zhang^{6,6}, J. Zhang^{6,6}, J. B. Zhang^{6,55}, J. W. Zhang^{6,4,6}, P. Zhang^{6,6}, Q. M. Zhang^{6,4,63}, 364 T. Zhang^{6,59}, X. M. Zhang^{6,6}, X. T. Zhang^{6,4,6}, Y. Zhang^{6,6}, Y. H. Zhang^{6,6}, Y. M. Zhang^{6,4,36}, Y. P. Zhang^{6,6}, Y. X. Zhang^{4,90} 365 Y. Y. Zhang^{6,6}, Y. Y. Zhang^{6,\mu,59}, Z. J. Zhang^{6,\mu,30}, Z. P. Zhang^{4,7}, Z. Y. Zhang^{6,27}, Z. Y. Zhang^{4,6}, F. Y. Zhao^{6,33}, J. Zhao^{6,4,6}, 366 R. Zhao^{6,36}, S. J. Zhao^{6,50}, T. C. Zhao^{6,6}, D. Q. Zheng^{6,61}, H. Zheng^{6,30}, M. S. Zheng^{6,67}, Y. H. Zheng^{6,79}, W. R. Zhong^{6,61}, J. Zhou^{6,67}, 367 L. Zhou^{*δ,μ,6,*} N. Zhou^{*δ,7,*} S. Zhou^{*δ,6,*} X. Zhou^{*δ,27,*} J. Zhu^{*δ,36,*} K. J. Zhu^{*δ,6,*} H. L. Zhuang^{*δ,μ,6,*} L. Zong^{*δ,31,*} J. H. Zou^{*δ,μ,6,*} 368

000	
370	¹ Pontificia Universidad Católica de Chile, Santiago, Chile
371	² 1PHC, Université de Strasbourg, CNRS/IN2P3, F-67037 Strasbourg, France
372	³ Pakistan Institute of Nuclear Science and Technology, Islamabad, Pakistan
373	⁴ INFN Catania and Dipartimento di Fisica e Astronomia dell Università di Catania, Catania, Italy
374	⁵ East China University of Science and Technology, Shanghai, China
375	⁶ Institute of High Energy Physics, Beijing, China
376	⁷ University of Science and Technology of China, Hefei, China
377	⁸ Joint Institute for Nuclear Research, Dubna, Russia
378	⁹ INFN Sezione di Milano and Dipartimento di Fisica dell Università di Milano, Milano, Italy
379	¹⁰ Department of Physics, Faculty of Science, Chulalongkorn University, Bangkok, Thailand
380	¹¹ Comenius University Bratislava, Faculty of Mathematics, Physics and Informatics, Bratislava, Slovakia
381	¹² University of Wisconsin, Madison, Wisconsin 53706, USA
382	¹³ Department of Physics and Earth Science, University of Ferrara and INFN Sezione di Ferrara, Ferrara, Italy
383	¹⁴ Wright Laboratory and Department of Physics, Yale University, New Haven, Connecticut 06520, USA ¹⁵ INFN Milano Bicocca and University
384	of Milano Bicocca, Milano, Italy
385	¹⁶ INFN Sezione di Padova, Padova, Italy

12

386	¹⁷ University of Roma Tre and INFN Sezione Roma Tre, Roma, Italy
387	¹⁸ III. Physikalisches Institut B, RWTH Aachen University, Aachen, Germany
388	¹⁹ Brookhaven National Laboratory, Upton, New York 11973, USA
389	²⁰ IJCLab, Université Paris-Saclay, CNRS/IN2P3, 91405 Orsay, France
390	²¹ Eberhard Karls Universität Tübingen, Physikalisches Institut, Tübingen, Germany
391	²² Department of Physics, National Taiwan University, Taipei
392	²³ Université de Bordeaux, CNRS, CENBG-IN2P3, F-33170 Gradignan, France
393	²⁵ National United University, Miao-Li
394	²³ Dipartimento di Fisica e Astronomia dell'Universita' di Padova and INFN Sezione di Padova, Padova, Italy
395	²⁰ Centre de Physique des Particules de Marseille, Marseille, France
396	²¹ Wuhan University, Wuhan, China
397	²⁰ INFN Milano Bicocca and Politecnico of Milano, Milano, Italy
398	³⁰ D
399	³⁶ Dongguan University of Technology, Dongguan, China
400	³² Tsinghua University, Beijing, China
401	³³ Lecience of Medical Chiversity, Nanjing, China
402	³⁴ Institute of Modern Physics, Chinese Accaemy of Sciences, Lanzhou, China
403	³⁵ Institute of Physics Wational Children in the State of the State
404	³⁶ Surv Yat Sar University Oniversity, Beijing, China
405	³⁷ Foundation Conversity, Guangzhou, Chuna
406	** Forschungszentrum Jutich Gmort, Nuclear Physics institute IKP-2, Jutich, Germany
407	³⁹ Lomonosov Moscow State University, Moscow, Russia
408	40 Criversidade Estadual de Londrina, Evnarina, Brazil
409	⁴¹ human in the second difference of the sec
410	⁴⁴ INFN Sezione di Perugia and Dipartimento di Chimica, Biologia e Biotecnologie dell'Università di Perugia, Perugia, Italy
411	⁴² Universite Libre de Bruxelles, Brussels, Belgium
412	⁴³ Siena College, Loudonville, New York 12211, USA
413	** Department of Physics and Astronomy, University of California, Irvine, California, USA
414	⁴⁻³ Institute of Physics, Johannes-Gutenberg Universität Mainz, Mainz, Germany
415	⁴⁶ Suranaree University of Technology, Nakhon Ratchasima, Thailand
416	⁴ Charles University, Faculty of Mathematics and Physics, Prague, Czech Republic
417	⁴⁶ Institute for Nuclear Research of the Russian Academy of Sciences, Moscow, Russia
418	* Department of Physics, University of Illinois at Urbana-Champaign, Urbana, Illinois 61801, USA
419	⁵⁰ School of Physics and Microelectronics, Zhengzhou University, Zhengzhou, China
420	⁵¹ Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA
421	⁵² University of Jyvaskyla, Department of Physics, Jyvaskyla, Finland
422	³³ Wuyi University, Jiangmen, China
423	⁵⁴ Guangxi University, Nanning, China
424	⁵⁵ Harbin Institute of Technology, Harbin, China
425	⁵⁶ Technische Universität München, München, Germany
426	⁵⁷ Department of Physics, Illinois Institute of Technology, Chicago, Illinois 60616, USA
427	⁵⁸ Institute of Hydrogeology and Environmental Geology, Chinese Academy of Geological Sciences, Shijiazhuang, China
428	⁵⁹ School of Physics and Astronomy, Shanghai Jiao Tong University, Shanghai, China
429	⁶⁰ Forschungszentrum Jülich GmbH, Central Institute of Engineering, Electronics and Analytics - Electronic Systems(ZEA-2), Jülich, Germany
430	⁶¹ Jinan University, Guangzhou, China
431	⁶² Beijing Normal University, Beijing, China
432	⁶³ Xi'an Jiaotong University, Xi'an, China
433	⁶⁴ Institute of Experimental Physics, University of Hamburg, Hamburg, Germany
434	⁶⁵ Beijing Institute of Spacecraft Environment Engineering, Beijing, China
435	⁶⁶ Department of Physics, University of Houston, Houston, Texas 77204, USA
436	⁶⁷ China Institute of Atomic Energy, Beijing, China
437	68 Center for Neutrino Physics, Virginia Tech, Blacksburg, Virginia 24061, USA
438	⁶⁹ Yerevan Physics Institute, Yerevan, Armenia
439	⁷⁰ Nankai University, Tianjin, China
440	⁷¹ Department of Physics, University of Cincinnati, Cincinnati, Ohio 45221, USA
441	⁷² Department of Physics, College of Science and Technology, Temple University, Philadelphia, Pennsylvania 19122, USA
442	⁷³ Department of Physics, University of California, Berkeley, California 94720, USA
443	⁷⁴ Department of Physics, The University of Hong Kong, Pokfulam, Hong Kong, China
444	⁷⁵ SUBATECH, Université de Nantes, IMT Atlantiaue, CNRS-IN2P3, Nantes, France
445	⁷⁶ College of Electronic Science and Engineering, National University of Defense Technology, Changsha, China
446	⁷⁷ The Radiochemistry and Nuclear Chemistry Group in University of South China. Henevane. China
447	⁷⁸ Tsung-Dao Lee Institute, Shanghai Jiao Tong University, Shanehai, China
448	⁷⁹ University of Chinese Academy of Sciences, Beijing China
449	⁸⁰ Joseph Henry Laboratories, Princeton University Princeton New Jersey 08544 USA
450	⁸¹ Jilin University, Chanochun China
451	⁸² Xiamen University, Xiamen, China
452	⁸³ School of Physics, Peking University Reijing China
453	⁸⁴ Laboratori Nazionali di Frascati dell'INFN. Roma Italy

454	⁸⁵ California Institute of Technology, Pasadena, California 91125, USA
455	⁸⁶ College of William and Mary, Williamsburg, Virginia 23187, USA
456	⁸⁷ Universidad Tecnica Federico Santa Maria, Valparaiso, Chile
457	⁸⁸ Pontificia Universidade Catolica do Rio de Janeiro, Rio, Brazil
458	⁸⁹ National Astronomical Research Institute of Thailand, Chiang Mai, Thailand
459	⁹⁰ China General Nuclear Power Group, Shenzhen, China
460	⁹¹ Institute of Electronics and Computer Science, Riga, Latvia
461	⁹² Iowa State University, Ames, Iowa 50011, USA
462	⁹³ Chongqing University, Chongqing, China
463	^µ Daya Bay collaboration
464	^δ JUNO collaboration

Statement about the "CRediT author statement":

Due to the large number of participants in this experiment, no CRediT taxonomy has been filled in.

Declaration of interests

 \boxtimes The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

□ The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: