

This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Cayol, Claire; Giermek, Anna; Gomez-Chamorro, Andrea; Hytönen, Jukka; Kallio, Eva; Mappes, Tapio; Salo, Jemiina; Voordouw, Maarten Jeroen; Koskela, Esa

Title: *Borrelia afzelii* alters reproductive success in a rodent host

Year: 2018

Version: Accepted version (Final draft)

Copyright: © 2018 the Authors

Rights: In Copyright

Rights url: <http://rightsstatements.org/page/InC/1.0/?language=en>

Please cite the original version:

Cayol, C., Giermek, A., Gomez-Chamorro, A., Hytönen, J., Kallio, E., Mappes, T., Salo, J., Voordouw, M. J., & Koskela, E. (2018). *Borrelia afzelii* alters reproductive success in a rodent host. *Proceedings of the Royal Society B : Biological Sciences*, 285(1884), Article 20181056. <https://doi.org/10.1098/rspb.2018.1056>

1 ***Borrelia afzelii* alters reproductive success in a rodent host**

2
3 Claire Cayol^{1*}, Anna Giermek², Andrea Gomez-Chamorro³, Jukka Hytönen⁴, Eva Riikka
4 Kallio^{1,5}, Tapio Mappes¹, Jemiina Salo⁴, Maarten Jeroen Voordouw³, Esa Koskela¹

5
6 **Affiliations**

7
8 ¹Department of Biological and Environmental Science. P.O. Box 35, University of Jyväskylä, FI-
9 40014, Jyväskylä, Finland; claire.c.cayol@jyu.fi; tapio.mappes@jyu.fi; esa.m.koskela@jyu.fi

10 ²Institute of Environmental Sciences. Gronostajowa 7, 30-387, Jagiellonian University, Cracow,
11 Poland; anna.giermek@doctoral.uj.edu.pl

12 ³Institut de Biologie, Laboratoire d'Ecologie et Evolution des Parasites. Rue Emile-Argand 11,
13 Université de Neuchâtel, CH-2000 Neuchâtel, Switzerland; maarten.voordouw@unine.ch;
14 andrea.gomez@unine.ch

15 ⁴Institute of Biomedicine. Kiinamyllynkatu 13, University of Turku, FI-20520 Turku, Finland;
16 jukhyt@utu.fi; jemiina.neuvonen@utu.fi

17 ⁵Department of Ecology and Genetics, P.O. Box 3000, FI-90014 University of Oulu, Oulu,
18 Finland; Eva.Kallio@oulu.fi

19 *Correspondence to: claire.c.cayol@gmail.com; +358 40 484 3717; Fax: +358 (0)14 617 239

20
21 **Keywords:** *Borrelia afzelii*; fitness; host-pathogen interaction; *Myodes glareolus*; zoonosis;
22 natural host

23 **Abstract**

24 The impact of a pathogen on the fitness and behaviour of its natural host depends upon the host-
25 parasite relationship in a given set of environmental conditions. Here, we experimentally
26 investigated the effects of *Borrelia afzelii*, one of the etiological agents of Lyme disease in
27 humans, on the fitness of its natural rodent host, the bank vole (*Myodes glareolus*), in semi-
28 natural conditions with two contrasting host population densities. Our results show that *B. afzelii*
29 can modify the reproductive success and spacing behaviour of its rodent host, whereas host
30 survival was not affected. Infection impaired the breeding probability of large bank voles.
31 Reproduction was hastened in infected females without alteration of the offspring size at birth.
32 At low density, infected males produced fewer offspring, fertilised fewer females and had lower
33 mobility than uninfected individuals. Meanwhile, the infection did not affect the proportion of
34 offspring produced or the proportion of mating partner in female bank voles. Our study is the
35 first to show that *B. afzelii* infection alters the reproductive success of the natural host. The
36 effects observed can reflect the sickness behaviour due to the infection or they can be a
37 consequence of a manipulation of the host behaviour by the bacteria.

38 **1. Introduction**

39 The impact of pathogens on the physiology, behaviour and fitness of their natural hosts is a key
40 determinant for the co-evolution between the pathogen and the host [1–4]. Identifying the effect
41 of a pathogen on all components of host fitness is also essential for predicting the population
42 dynamics of a host-pathogen association and is fundamental for anticipating zoonotic outbreaks
43 [5–8]. However, the study of the impact of parasites on their natural hosts often focuses on host
44 survival [3,9–11], despite the recognition that host reproduction is an important component of
45 host fitness [12–14]. Indeed, subtle effects of an endemic pathogen on the reproduction of its
46 natural host can influence the population dynamics of the wild host [15,16] and ultimately the
47 population dynamics of the pathogen [17].

48 Numerous studies have shown that pathogen virulence depends on ecological factors such
49 as temperature and nutrition [18–21]. Another important ecological factor is host population
50 density because it generates intra-specific competition for limited resources such as space, food
51 and mating partners [22–24]. High host density is therefore expected to exacerbate pathogen
52 virulence. Fluctuations in population density are typical in many small mammal species such as
53 rodents [25]. However, experimental studies on density-dependent costs of infection in rodents
54 are rare because it is often difficult to manipulate host density in an ecologically relevant way
55 (but see [10,26,27]).

56 Spirochete bacteria belonging to the *Borrelia burgdorferi* sensu lato (s. l.) complex cause
57 Lyme borreliosis in humans, which is the most common vector-borne disease in the northern
58 hemisphere [28,29]. *Borrelia afzelii*, which is transmitted by *Ixodes* ticks and hosted by rodents,
59 is the most common etiological agent of human Lyme borreliosis in Europe [28,30]. While Lyme

60 borreliosis causes serious morbidity in humans [31,32], there is currently no clear evidence that
61 *Borrelia burgdorferi* s. l. reduces the fitness of the rodent or avian reservoir hosts [6,9,33–36].
62 However, most studies that investigate the virulence of *Borrelia burgdorferi* s. l. pathogens were
63 correlational and have focused on host survival and, to date, the potential effects on host
64 reproductive success have been ignored (but see [37,38] for physiological cost and effect on host
65 behaviour, respectively).

66 We conducted a field experiment to test whether *B. afzelii* reduces the survival and
67 reproductive success of its rodent host, the bank vole (*Myodes glareolus*). Rodent populations
68 are often strongly influenced by density-dependent effects [25]. We, therefore, hypothesised that
69 the detrimental effects of *B. afzelii* infection on the fitness of bank voles would be more
70 pronounced at high population density. Here, we show that while *B. afzelii* did not affect the host
71 survival, the infection impaired the reproduction of large bank voles, and unexpectedly, that male
72 bank voles had lower reproductive performances at low population density.

73

74 **2. Methods**

75 **(a) Ethical statement**

76 The Finnish Animal Experiment Board approved the trapping and handling methods used in this
77 study under the authorisations ESAVI/3834/04.10.03/2011, ESAVI/7256/04.10.07/2014 and
78 ESAVI/3457/04.10.07/2015.

79

80 **(b) Experimental design**

81 The schedule of the experimental procedure is shown in Fig. S1, and all methods are detailed in
82 the supplementary material. Male and female bank voles (*Myodes glareolus*) from the lab colony
83 at the University of Jyväskylä were measured and assigned to either the *B. afzelii* infection group
84 (injected with a local strain of *B. afzelii*) or the uninfected control group (injected with PBS). All
85 infected and uninfected voles (total of 136 individuals, 68 females and 68 males) were released
86 in 12 large outdoor vegetated enclosures (each 0.2 ha) that were assigned to “high” density (16
87 individuals per enclosure, 8 females and 8 males, half of each sex infected, 5 enclosures) and
88 “low” density (8 individuals per enclosure, 4 females and 4 males, half of each sex infected, 7
89 enclosures) treatments. In the enclosures, the bank voles could move and reproduce freely for 18
90 days, which is the minimum gestation length in females. During this period, spacing behaviour
91 was monitored using live trapping. At the end of this period, all trapped individuals were taken to
92 the laboratory for measurements and monitoring of parturition. Male reproductive success was
93 determined by paternity analyses conducted on the offspring born in the laboratory.

94

95 **(c) Measurements**

96 Before the enclosure period, individuals were weighed, and the head width was measured with a
97 calliper ruler (Electronic Digital Caliper, Scala). These measurements were taken into account
98 when experimental animals were assigned to treatments and enclosures. An ear tissue sample
99 was taken for paternity analysis. A blood sample was taken for an ELISA targeting *B.*
100 *burgdorferi* s. l.-specific IgG antibodies [39] (supplementary material).

101 After the enclosure period, the body measurements and blood sampling were carried out
102 as described above. Males were processed shortly after the trapping day, gravid females were
103 processed after parturition, and females that were not gravid were processed at the end of the
104 experiment. Pups were measured (body mass and head width) within 24 hours of parturition. All
105 measurements were performed blind regarding the infection treatment and density treatment.

106

107 **(d) Statistical analysis**

108 All statistical analyses were carried out using the statistical software R version 3.1.1. Survival of
109 bank voles in the enclosures and individual breeding probability are binary variables. For
110 survival, individuals were assigned 0 or 1 depending on whether they were trapped at the end of
111 the experiment or not. For the assessment of the breeding probability, individuals were assigned
112 0 or 1 depending on whether their number of produced offspring was zero or at least one.

113 Moreover, two response variables of reproductive success were calculated: (1) ‘relative number
114 of offspring’ is the proportion of offspring produced in an enclosure by a given individual, (2)
115 ‘relative number of partners’ is the proportion of partners with which a given individual
116 produced offspring. Eventually, space trapping data allowed to calculate two different home
117 range variables: home range perimeter (m) and home range surface (m²) (Table S1).

118 In the statistical analyses, the injection (*Borrelia afzelii* vs. PBS) was used to define “infection”
119 treatment (infected vs. uninfected), and the population density in the enclosure defined the
120 “density” treatment (“low” vs. “high”). The explanatory variables of the full models always
121 included the two experimentally manipulated factors, *i.e.* the infection treatment and the
122 population density in the enclosure, sex, body mass before injection (BM) and relevant two- and

123 three-way interactions. Enclosure ID was included as a random effect in all models. Three-way
124 interactions involving vole sex were expected in models assessing bank vole reproductive
125 success because the drivers of reproductive success differ between male and female bank voles
126 [40–43]. When these three-way interactions were significant in the full model (see Table S1),
127 separate analyses were conducted for males and females to ease the interpretation of the
128 interactions. Otherwise, reductions of the full models were carried out starting from the non-
129 significant interactions (see supplementary material).

130 For gravid females, the parturition delay was calculated as the difference in the number of
131 days between the date the first litter was observed and the parturition date for the other pregnant
132 females. This variable was modelled as a function of infection, density, BM, and the interaction
133 infection \times density. Moreover, offspring body mass at birth and head width at birth were
134 modelled as a function of the infection status of the mother and father, density and all their two-
135 and three-way interactions. Offspring sex and litter size were included as covariates. Enclosure
136 ID, mother ID and father ID were included as random effects.

137 To analyse the data, we used generalized linear mixed models (GLMMs) with an error
138 distribution that was either normal (home range perimeter, home range surface, body mass and
139 head width of offspring), binomial (survival, breeding probability and variables describing
140 reproductive success: relative number of offspring and relative number of partners) or negative
141 binomial (female parturition delay).

142

143

144 **3. Results**

145 Out of the 68 female and 68 male bank voles released into the enclosures at the beginning of the
146 experiment, 48 females and 30 males (one of which was found dead in the trap) were recovered,
147 and the remaining 58 individuals were considered as dead. Of these 58 individuals, 56 were
148 never observed during the 14 trapping occasions and 2 were not observed during the six last
149 trapping occasions. As we did not observe any introduction of unmarked wild bank voles in the
150 enclosures, and all trapped animals were found in their original enclosure, we assume that
151 missing animals died, rather than escaped. Of the 78 captured individuals, 39 were from the *B.*
152 *afzelii* infection group (24 females, 15 males), and 39 were from the control group (24 females,
153 15 males, including the individual found dead in the trap). There was no effect of *B. afzelii*
154 infection or population density on the survival of bank voles (GLMM: $p > 0.35$, Table S4), but
155 females survived better than male bank voles (GLMM: $p < 0.01$, Table S4).

156

157 **(a) *B. afzelii* infection reduces the breeding probability of large bank voles**

158 Based on the paternity test, 39 of 68 males reproduced during the study (18 of the 30 males that
159 were trapped and 21 of the 38 males that were not trapped). For the analysis of reproductive
160 success, all males were included, regardless of whether they were trapped or not at the end of the
161 study. Out of the 48 captured females, 45 gave birth in the laboratory. We found that the effect of
162 *B. afzelii* infection on bank vole breeding probability was dependent on body size: among small
163 individuals, there was no difference in the breeding probability between infected and uninfected
164 animals. However, uninfected individuals had significantly higher breeding probability than *B.*
165 *afzelii*-infected individuals among large bank voles (GLMM: body mass \times infection, $p = 0.05$,

166 Table 1 and Fig. 1; Table S4 and Fig S4).

167

168 **(b) *B. afzelii* infection reduces male reproductive success at low density**

169 Reproductive success was further explored as the analysis of the relative number of produced
170 offspring and the relative number of partners. The three-way interaction infection \times density \times sex
171 was significant for the relative number of offspring (GLMM: $p = 0.02$, Table S1) and the relative
172 number of partners (GLMM: $p = 0.03$, Table S1), providing evidence that infection and breeding
173 density affected these components of reproductive success differently in males and females. In
174 male bank voles, the relative number of offspring and partners were associated with *B. afzelii*
175 infection status, but the effect differed between the population density treatments (Table 1, Fig.
176 2, Table S2). In low density, uninfected control males sired a higher relative number of offspring
177 (0.42) and fertilised a higher relative number of females (0.43) than *B. afzelii*-infected males
178 (0.05 offspring sired and 0.10 female fertilised). Conversely, in high density there was no effect
179 of the infection treatment: the relative number of offspring sired by uninfected and infected
180 males were 0.13 and 0.11, and the relative number of females fertilised by uninfected and
181 infected males were 0.18 and 0.19, respectively (GLMM: $p = 0.004$ and $p = 0.02$, Table 1, Fig.
182 2). For female bank voles, the relative number of offspring and partners were not affected by the
183 infection (the proportion of offspring produced by uninfected and infected females was 0.27 and
184 0.26, respectively; GLMM: $p = 0.79$, Table S3). As expected, population density influenced the
185 relative number of offspring produced by a female bank vole (relative number of offspring
186 produced by females from low and high-density enclosures was 0.41 and 0.18, respectively;
187 GLMM: estimate on the logit scale (SE): density = 0.85 (0.25), $p < 0.001$, Table S3).

188

189 **(c) *B. afzelii* infection reduces male home range at low density**

190 We found evidence that male and female bank voles differ in their spacing behaviour as the
191 three-way interaction infection \times density \times sex was significant for home range surface and home
192 range perimeter (GLMM: $p < 0.01$, $p = 0.03$, respectively, Table S1). For the uninfected male
193 bank voles, the home range surface was significantly larger in the low-density enclosures (808
194 m^2) compared to the high-density enclosures (378 m^2) (LMM: $p = 0.003$, Table 1, Fig. 3). In
195 contrast, the home range surface of the *B. afzelii*-infected male bank voles was not significantly
196 different between the low density (360 m^2) and high density (524 m^2) enclosures (Table 1, Fig.
197 3). Female home range surface and perimeter were not affected by the infection or the density
198 treatments (Table S3).

199

200 **(d) Infection caused early reproduction in female bank voles**

201 Of the 48 females captured from the enclosures, 45 were gravid and produced a total of 226
202 pups, with a mean number of 5 pups per female (range: 1–7). *B. afzelii*-infected females
203 reproduced on average 3 days earlier than uninfected control females (GLMM: $p = 0.003$, Fig. 4,
204 Table 1) and this effect was independent of the population density (GLMM: $p = 0.30$, Table 1).
205 The size of the offspring at birth was not affected by the infection treatment of the mother or
206 father or population density (LMM for all variables: $p > 0.05$, Table S6).

207

208

209 **4. Discussion**

210 We examined the hypothesis that *B. afzelii* infection reduces the reproductive success of the
211 rodent host and we tested the density-dependence of this effect. We found that *B. afzelii* infection
212 had density-dependent and statistically differing effects on the relative numbers of partners and
213 offspring of male and female bank voles. In males, infected individuals kept at low population
214 density sired a lower proportion of offspring, fertilised a lower proportion of females and
215 displayed smaller home range surface than uninfected males (Fig. 2, 3). In females, by contrast,
216 *B. afzelii* infection did not affect the relative offspring number, relative number of partners and
217 home range surface, but infected individuals gave birth *ca.* 3 days earlier than uninfected
218 individuals. The offspring size (head width and body mass) was not affected by the mother's
219 infection status (Fig. 4, Table S6). Finally, in both sexes, infection reduced the breeding
220 probability of large individuals but did not affect their survival (Fig. 1, Table S4, Table S5).

221 Previous studies found no evidence that infection with *Borrelia burgdorferi* s. l. reduces
222 the fitness of natural hosts; however, most of them were correlational or focused on another
223 genospecies than *B. afzelii*. For instance, capture-mark-recapture (CMR) studies on wild
224 populations of the white-footed mouse (*Peromyscus leucopus*) or the black-legged kittiwake
225 found no effect of infection with *B. burgdorferi* s. l. on the survival of these hosts [9,34,35].
226 Similarly, we found that infection with *B. afzelii* did not impair survival of the bank vole.
227 Another study on white-footed mice found no effect of *B. burgdorferi* s. s. on the wheel running
228 behaviour over the six weeks following experimental infection [6]. In our study, by contrast, the
229 effect of infection on home range size may result from altered running behaviour. A recent study
230 reports a trend in increased foraging behaviour in white-footed mice treated with an anti-*B.*

231 *burgdorferi* vaccine compared with sham-treated individuals, suggesting similarly to our finding,
232 a wider ranging behaviour in individuals with low or with no infection burden [38]. To our
233 knowledge, our study is the first to address the effect of *B. afzelii* infection on host reproduction
234 experimentally under field conditions. Studying the effects of infections on host reproduction is
235 challenging in wild rodent populations, and reproduction is often a latent variable inferred from
236 observed variables. Our experimental setting allows controlling for several sources of variation
237 and confounding factors (*e.g.* age of the host), and we were able to estimate the reproductive
238 success reliably.

239 The experimental infection was performed by peritoneal injection of the bacteria rather
240 than the natural infection route, which involves *Ixodes* ticks. The infection dose and route were
241 based on the literature [33,36,44–46]. The intraperitoneal route was chosen as it has been shown
242 to give more widely disseminated infection than the subcutaneous route [47]. The use of
243 injection instead of the natural transmission route can be debatable, *e.g.* due to the lack of tick
244 salivary compounds that enhance the infectivity of *Borrelia burgdorferi* s. l. [48,49]. Molecules
245 present in tick saliva promote the infection by manipulating or depressing the immune system
246 (*e.g.* Salps) [50]. The injection of *B. burgdorferi* s. l. with tick salivary gland extract led to higher
247 infection success with higher bacterial dissemination, so-called saliva-assisted transmission
248 [50,51]. The lack of these molecules could lead to misestimation of the effects of the infection on
249 the host. However, the injection allows the experimenter to control for the bacterial dose, and it
250 eliminates the variation linked to the tick vectorial capacity [52], hence ensuring a controlled
251 exposure of the study animals to the bacteria. We acknowledge that needle inoculation mimics
252 only grossly the infection via tick bite. However, we can expect any observed effect to be caused

253 by the *B. afzelii* infection given our controlled experimental conditions.

254 The demonstration of fitness-related costs caused by *B. burgdorferi* s. l. infection is
255 important for understanding the evolution of resistance in natural hosts. Recent field studies on
256 the bank vole suggest that polymorphism at the Toll-like receptor 2 (TLR2) gene, a pathogen
257 recognition receptor of the innate immune system, was associated with variation in susceptibility
258 to *B. afzelii* [53,54]. The prevalence of *B. afzelii* infection in bank voles that were homozygous
259 for the C2 resistance allele was half that of the bank voles that were homozygous for the C1
260 susceptibility allele [53]. A study of the TLR2 polymorphism in bank vole populations across
261 Europe found that the resistance allele against *B. afzelii* (C2) was more common in countries
262 with a high incidence of human Lyme disease [55]. This result led Tschirren (2015) to suggest
263 that *B. afzelii* was driving the evolution of the resistance allele at the TLR2 gene in European
264 bank vole populations. However, without clear evidence of reduced fitness in infected rodents,
265 the mechanism of selection was unclear. Our demonstration that infection with *B. afzelii* reduces
266 male reproductive success supports the hypothesis that this pathogen could be driving selection
267 on the TLR2 gene in bank vole populations.

268 The effect of the infection on the relative number of offspring sired and the relative
269 number of females fertilised by a given male bank vole was density-dependent. In the low-
270 density populations, uninfected control males fertilised more females and fathered more
271 offspring compared to the infected males and males kept in high population density (Fig. 2). This
272 result was counter-intuitive, as we predicted that the negative effects of high population density,
273 such as reduced per capita food availability, more aggressive interactions, and potentially higher
274 stress levels, would exacerbate the cost of *B. afzelii* infection [10,42,56–58]. Three hypotheses

275 can explain this result. First, several studies have shown that the strength of male-male
276 competition can vary with population density in a non-linear fashion [See, for instance, 59–61].
277 For example, males can modify their reproductive strategy in high population density leading to
278 lower rates of aggression and lower reproductive success [59,62]. Second, as estimates of the
279 relative number of partners and the relative number of offspring were based on paternity tests,
280 cryptic female choice (*i.e.* a female choice that occurs in the reproductive tract of the female,
281 leading to fertilisation bias in favour of specific males [63,64]) might have occurred. Thus, a
282 density-dependent female cryptic choice favouring healthy males in low-density populations
283 cannot be excluded. Finally, a spurious effect linked to the length of our experiment, which
284 covers only one reproductive episode, cannot be ruled out [65].

285 In the low-density populations, uninfected control males had larger home range sizes than
286 infected males whereas, in the high-density enclosures, there was no significant difference in the
287 home range size between uninfected and infected male bank voles (Fig. 3). One possible
288 explanation for this density-dependent home range reduction is that at high density, males may
289 reduce their exploratory behaviour to avoid encountering other males and having to engage in
290 aggressive male-male interactions. Moreover, at high density, with eight females available in the
291 enclosure, the chance for a male to encounter a receptive female might be higher than in the low-
292 density enclosure where only four females are available. Indeed, female bank voles are territorial
293 and hyperdispersed [41,66]. Consequently, at low density, male bank voles may need to explore
294 a larger home range to search for receptive females than at high density. As expected, the
295 uninfected males had a larger home range in low population density, whereas the infected males
296 presumably allocated resources to their immune response instead of explorative behaviour. In

297 contrast, female bank voles had a smaller home range size than males, that was not affected by
298 population density, reflecting the territorial behaviour of females especially, during late gestation
299 when the space trapping took place [41,66,67].

300 We found that the cost of infection was more important in large bank voles, which are the
301 most frequently infested with ticks and *B. burgdorferi* s. l. in nature [9,68,69]. Large infected
302 individuals showed reduced reproductive success compared to large healthy individuals. Food
303 resource is generally known to constrain reproduction and food addition has been shown to
304 enhance reproductive success in similar outdoor enclosure setups [42,70,71]. These food
305 constraints might have a more negative effect on the large individuals, which have greater
306 energetic needs [72]. Infected large voles showed altered breeding probability regardless of the
307 population density.

308 Infected females plastically modified their life history and reproduced *ca.* 3 days earlier
309 than uninfected females without alteration of the size of the offspring at birth, *i.e.* without signs
310 of premature birth (Fig. 4, Table 1, Table S3). In nature, reproducing females give birth to 1 or 2
311 litters per reproductive season [73], and most individuals live only one season. The biological
312 importance of giving birth three days earlier is not clear, as concerns population dynamics. At
313 the individual level, early reproduction can be a compensatory strategy if parasites reduce the
314 reproductive success of the adult host later in life via morbidity, mortality or castration [74–76].
315 According to the terminal investment theory, individuals maximise their fitness by allocating
316 resources to immediate reproduction when the prospects for future reproduction are reduced, for
317 example by chronic infection [27,77–79]. It remains to be estimated whether *B. afzelii* impairs
318 reproduction of female bank vole during the late stage of infection.

319 In summary, our study shows, for the first time, that the zoonotic pathogen *B. afzelii* can
320 influence the reproductive success of its rodent host. The effect of the infection on the relative
321 number of offspring and partners differed between male and female bank voles. Although large
322 body size favoured reproduction in uninfected individuals, this size benefit disappeared if the
323 individual was infected with *B. afzelii*. In males, infected individuals kept at low population
324 density displayed smaller home range surface than uninfected males. Lower mobility can be a
325 consequence of sickness behaviour due to the infection. On the other hand, predation risk by
326 small carnivores generally increases with vole mobility [80]. By reducing home range size,
327 infection with *B. afzelii* could lower the predation risk of male bank voles by small carnivores,
328 enhancing at the same time, its own fitness [81]. The hypothesis of manipulation of the rodent
329 host by *B. afzelii* is yet to be explored.

330

331 **Supplementary information is available for this paper.**

332

333 **Data accessibility statement:** The dataset analysed during the current study is available in the
334 JYX repository, <http://urn.fi/URN:NBN:fi:jyu-201806133148> [82].

335

336 **Competing interests:** The authors declare no competing financial interests.

337

338 **Authors' contributions:** C.C., E.K., T.M., E.R.K. conceived the study. The fieldwork and
339 laboratory work were carried out by all authors, with the helpers listed in the

340 Acknowledgements. C.C., E.K., T.M., E.R.K., M.V. analysed the results, with input from the

341 other authors. All authors contributed to the interpretation and critical revision. C.C. led the
342 writing of the paper. All authors gave final approval for publication.

343

344 **Acknowledgements:** We thank Anja Siukkola, Susanne Varjola and Joannes Van Cann and the
345 Konnevesi research station for their assistance in the field. We thank Sami Kyröläinen, Elina
346 Virtanen, Annukka Pietikäinen, Julia Honkasalo, Anouk Sarr and Olivier Rais for their help in
347 the lab. Swanne Gordon gave constructive feedback at an early stage of this manuscript. We
348 thank Maarten Voordouw's and Johanna Mappes's lab meetings for their comments. We thank
349 two anonymous referees for valuable comments on the manuscript.

350

351 **Fundings:** This project was supported by the Kone Foundation, the University of Jyväskylä and
352 the Academy of Finland (Eva R. Kallio 250524, 310104; Esa Koskela 257340 and Tapio Mappes
353 132190, 268670).

354 **References**

- 355 1. Anderson RM, May RM. 1979 Population biology of infectious diseases: Part I. *Nature*
356 **280**, 361–367. (doi:10.1038/280361a0)
- 357 2. Scott ME, Dobson A. 1989 The role of parasites in regulating host abundance. *Parasitol.*
358 *today* **5**, 176–183. (doi:10.1016/0169-4758(89)90140-3)
- 359 3. Cattadori IM, Haydon DT, Hudson PJ. 2005 Parasites and climate synchronize red grouse
360 populations. *Nature* **433**, 737–741. (doi:10.1038/nature03276)
- 361 4. Smith MJ, White A, Sherratt JA, Telfer S, Begon M, Lambin X. 2008 Disease effects on
362 reproduction can cause population cycles in seasonal environments. *J. Anim. Ecol.* **77**,
363 378–389. (doi:10.1111/j.1365-2656.2007.01328.x)
- 364 5. Anderson RM, May RM. 1978 Regulation and stability of host-parasite population
365 interactions. *J. Anim. Ecol.* **47**, 219–247.
- 366 6. Schwanz LE, Voordouw MJ, Brisson D, Ostfeld RS. 2011 *Borrelia burgdorferi* has
367 minimal impact on the Lyme disease reservoir host *Peromyscus leucopus*. *Vector borne*
368 *zoonotic Dis.* **11**, 117–124. (doi:10.1089/vbz.2009.0215)
- 369 7. Feore SM, Bennett M, Chantrey J, Jones T, Baxby D, Begon M. 1997 The effect of
370 cowpox virus infection on fecundity in bank voles and wood mice. *Proc. Biol. Sci.* **264**,
371 1457–1461. (doi:10.1098/rspb.1997.0202)
- 372 8. Karesh WB *et al.* 2012 Ecology of zoonoses: Natural and unnatural histories. *Lancet* **380**,
373 1936–1945. (doi:10.1016/S0140-6736(12)61678-X)

- 374 9. Voordouw MJ, Lachish S, Dolan MC. 2015 The Lyme disease pathogen has no effect on
375 the survival of its rodent reservoir host. *PLoS One* , 1–26.
- 376 10. Kallio ER, Voutilainen L, Vapalahti O, Vaheri A, Henttonen H, Koskela E, Mappes T.
377 2007 Endemic hantavirus infection impairs the winter survival of its rodent host. *Ecology*
378 **88**, 1911–1916. (doi:10.1890/06-1620.1)
- 379 11. Dobson AP, Hudson PJ. 1992 Regulation and stability of a free-living host-parasite
380 system : *Trichostrongylus tenuis* in red grouse . II . Population models. *J. Anim. Ecol.* **61**,
381 487–498.
- 382 12. Ehrlich PR, Roughgarden J, Roughgarden J. 1987 *The Science of Ecology*. University of
383 Missouri-Columbia: Macmillan.
- 384 13. Begon M, Harper JL, Townsend CR. 1996 *Ecology: Individuals, Populations and*
385 *Communities*. London: Blackwell Science.
- 386 14. Cockburn A. 1991 *An Introduction to Evolutionary Ecology*. London: Wiley.
- 387 15. Lafferty KD. 1993 Effects of parasitic castration on growth, reproduction and population
388 dynamics of the marine snail *Cerithidea californica*. *Mar. Ecol. Prog. Ser.* **96**, 229–237.
389 (doi:10.3354/meps096229)
- 390 16. Telfer S, Bennett M, Bown K, Carslake D, Cavanagh R, Hazel S, Jones T, Begon M. 2005
391 Infection with cowpox virus decreases female maturation rates in wild populations of
392 woodland rodents. *Oikos* **109**, 317–322. (doi:10.1111/j.0030-1299.2005.13734.x)
- 393 17. Begon M, Telfer S, Smith MJ, Burthe S, Paterson S, Lambin X. 2009 Seasonal host
394 dynamics drive the timing of recurrent epidemics in a wildlife population. *Proc. R. Soc. B-*

- 395 *Biological Sci.* **276**, 1603–1610. (doi:10.1098/rspb.2008.1732)
- 396 18. Blanford S, Thomas MB, Pugh C, Pell JK. 2003 Temperature checks the Red Queen?
397 Resistance and virulence in a fluctuating environment. *Ecol. Lett.* **6**, 2–5.
398 (doi:10.1046/j.1461-0248.2003.00387.x)
- 399 19. Mitchell SE, Rogers ES, Little TJ, Read AF. 2005 Host-parasite and genotype-by-
400 environment interactions: temperature modifies potential for selection by a sterilizing
401 pathogen. *Evolution (N. Y.)*. **59**, 70–80. (doi:10.1111/j.0014-3820.2005.tb00895.x)
- 402 20. Scholthof KB. 2007 The disease triangle: pathogens, the environment and society. *Nat.*
403 *Rev. Microbiol.* **5**, 152–156. (doi:10.1038/nrmicro1596)
- 404 21. Ilmonen P, Hakkarainen H, Koivunen V, Korpimäki E, Mullie A, Ilrnonen P, Korpimäki
405 E, Shutler D. 1999 Parental effort and blood parasitism in Tengmalm's owl: effects of
406 natural and experimental variation in food abundance. *Oikos* **86**, 79–86.
- 407 22. Begon M, Mortimer M, Thompson DJ. 2009 *Population ecology: a unified study of*
408 *animals and plants*. Oxford: John Wiley & Sons.
- 409 23. Krebs CJ. 1970 Microtus population biology : Behavioral changes associated with the
410 population cycle in *M. ochrogaster* and *M. pennsylvanicus*. *Ecology* **51**, 34–52.
- 411 24. Ostfeld RS, Canham CD, Pugh SR. 1993 Intrinsic density-dependent regulation of vole
412 populations. *Nature* **366**, 259–261. (doi:10.1038/366259a0)
- 413 25. Krebs CJ, Myers JH. 1978 Population cycles in small mammals. *Adv. Ecol. Res.* **8**, 267–
414 399. (doi:http://dx.doi.org/10.1016/S0065-2504(08)60280-9)

- 415 26. Burthe S, Telfer S, Begon M, Bennett M, Smith A, Lambin X. 2008 Cowpox virus
416 infection in natural field vole *Microtus agrestis* populations: significant negative impacts
417 on survival. *J. Anim. Ecol.* **77**, 110–9. (doi:10.1111/j.1365-2656.2007.01302.x)
- 418 27. Kallio ER, Helle H, Koskela E, Mappes T, Vapalahti O. 2015 Age-related effects of
419 chronic hantavirus infection on female host fecundity. *J. Anim. Ecol.* **84**, 1264–1272.
420 (doi:10.1111/1365-2656.12387)
- 421 28. Kurtenbach K, Hanincová K, Tsao JI, Margos G, Fish D, Ogden NH. 2006 Fundamental
422 processes in the evolutionary ecology of Lyme borreliosis. *Nat. Rev. Microbiol.* **4**, 660–
423 669. (doi:10.1038/nrmicro1475)
- 424 29. Rizzoli A, Hauffe HC, Carpi G, Vourc’h GI, Neteler M, Rosà R. 2011 Lyme borreliosis in
425 Europe. *Euro Surveill* **16**.
- 426 30. Hanincová K, Schäfer SM, Etti S, Sewell HS, Taragelová V, Ziak D, Labuda M,
427 Kurtenbach K. 2003 Association of *Borrelia afzelii* with rodents in Europe. *Parasitology*
428 **126**, 11–20. (doi:10.1017/S0031182002002548)
- 429 31. Stanek G, Wormser GP, Gray J, Strle F. 2012 Lyme borreliosis. *Lancet* **379**, 461–473.
430 (doi:10.1016/S0140-6736(11)60103-7)
- 431 32. Wang G, van Dam AP, Schwartz I, Dankert J. 1999 Molecular typing of *Borrelia*
432 *burgdorferi* sensu lato: taxonomic, epidemiological, and clinical implications. *Clin.*
433 *Microbiol. Rev.* **12**, 633–653.
- 434 33. Bey RF, Loken KI, Wu CC, Lin TL. 1995 Experimental infection of the red-backed vole
435 (*Clethrionomys gapperi*) with *Borrelia burgdorferi*. *J. Wildl. Dis.* **31**, 428–431.

- 436 34. Chambert T, Staszewski V, Lobato E, Choquet R, Carrie C, Mccoy KD, Tveraa T,
437 Boulinier T. 2012 Exposure of black-legged kittiwakes to Lyme disease spirochetes:
438 Dynamics of the immune status of adult hosts and effects on their survival. *J. Anim. Ecol.*
439 **81**, 986–995. (doi:10.1111/j.1365-2656.2012.01979.x)
- 440 35. Hofmeister EK, Ellis BA, Glass GE, Childs JE. 1999 Longitudinal study of infection with
441 *Borrelia burgdorferi* in a population of *Peromyscus leucopus* at a lyme disease-enzootic
442 site in Maryland. *Am. J. Trop. Med. Hyg.* **60**, 598–609.
- 443 36. Moody KD, Terwilliger GA, Hansen GM, Barthold SW. 1994 Experimental *Borrelia*
444 *burgdorferi* infection in *Peromyscus leucopus*. *J. Wildl. Dis.* **30**, 155–61.
445 (doi:10.7589/0090-3558-30.2.155)
- 446 37. Norte AC, Costantini D, Araújo PM, Eens M, Ramos JA, Heylen D. 2018 Experimental
447 infection by microparasites affects the oxidative balance in their avian reservoir host the
448 blackbird *Turdus merula*. *Ticks Tick. Borne. Dis.* **9**, 720–729.
449 (doi:10.1016/j.ttbdis.2018.02.009)
- 450 38. Ostfeld RS, Brisson D, Oggenfuss K, Devine J, Levy MZ, Keesing F. 2018 Effects of a
451 zoonotic pathogen, *Borrelia burgdorferi* , on the behavior of a key reservoir host. *Ecol.*
452 *Evol.* , 1–10. (doi:10.1002/ece3.3961)
- 453 39. Salo J, Jaatinen A, Soderstrom M, Viljanen MK, Hytonen J. 2015 Decorin binding
454 proteins of *Borrelia burgdorferi* promote arthritis development and joint specific post-
455 treatment DNA persistence in mice. *PLoS One* **10**, 1–17.
456 (doi:10.1371/journal.pone.0121512)

- 457 40. Sikorski MD, Wójcik AM. 1990 Mating system and reproductive success in a free-living
458 population of the bank vole, *Clethrionomys glareolus*. *Soc. Syst. Popul. Cycles Voles* ,
459 193–202.
- 460 41. Koskela E, Mappes T, Ylonen H. 1997 Territorial behaviour and reproductive success of
461 bank vole *Clethrionomys glareolus* females. *J. Anim. Ecol.* **66**, 341–349.
462 (doi:10.2307/5980)
- 463 42. Koskela E, Jonsson P, Hartikainen T, Mappes T. 1998 Limitation of reproductive success
464 by food availability and litter size in the bank vole, *Clethrionomys glareolus*. *Proc. R. Soc.*
465 *London B Biol. Sci.* **165**, 1129–1134.
- 466 43. Mills S, Grapputo A, Jokinen I, Koskela E, Mappes T, Oksanen TA, Poikonen T. 2009
467 Testosterone-mediated effects on fitness related phenotypic traits and fitness. *Am. Nat.*
468 **173**, 475–487. (doi:10.1086/597222)
- 469 44. McLean RG, Ubico SR, Cooksey LM. 1993 Experimental infection of the eastern
470 chipmunk (*Tamias striatus*) with the Lyme disease spirochete (*Borrelia burgdorferi*). *J*
471 *Wildl Dis* **29**, 527–532.
- 472 45. Baum E, Hue F, Barbour AG. 2012 Experimental infections of the reservoir species
473 *Peromyscus leucopus* with diverse strains of *Borrelia burgdorferi*, a Lyme disease agent.
474 *MBio* **3**. (doi:10.1128/mBio.00434-12)
- 475 46. Johnson RC, Kodner CL, Russell ME. 1986 Vaccination of hamsters against experimental
476 infection with *Borrelia burgdorferi*. *Zentralbl Bakteriol Mikrobiol Hyg A* **263**, 45–48.
- 477 47. Schwan TG, Burgdorfer W, Schrumpf ME, Karstens RH. 1988 The urinary bladder, a

- 478 consistent source of *Borrelia burgdorferi* in experimentally infected white-footed mice
479 (*Peromyscus leucopus*). *J Clin Microbiol* **26**, 893–895.
- 480 48. Gern L, Schaible UE, Simon MM. 1993 Mode of inoculation of the Lyme disease agent
481 *Borrelia burgdorferi* influences infection and immune responses in inbred strains of mice.
482 *J. Infect. Dis.* **167**, 971–975.
- 483 49. Ribeiro JMC. 1995 How ticks make a living. *Parasitol. Today* **11**, 91–93.
484 (doi:10.1016/0169-4758(95)80162-6)
- 485 50. Ramamoorthi N *et al.* 2005 The Lyme disease agent exploits a tick protein to infect the
486 mammalian host. *Nature* (doi:10.1038/nature03812)
- 487 51. Nuttall P, Labuda M, Bowman A. 2008 Saliva-assisted transmission of tick-borne
488 pathogens. In *Ticks: Biology, Disease and Control*, pp. 205–219. Cambridge Univ Press.
489 (doi:10.1017/CBO9780511551802.011)
- 490 52. de la Fuente J *et al.* 2017 Tick-pathogen interactions and vector competence:
491 Identification of molecular drivers for tick-borne diseases. *Front. Cell. Infect. Microbiol.*
492 **7**. (doi:10.3389/fcimb.2017.00114)
- 493 53. Tschirren B, Andersson M, Scherman K, Westerdahl H, Mittl PR, Raberg L. 2013
494 Polymorphisms at the innate immune receptor TLR2 are associated with *Borrelia* infection
495 in a wild rodent population. *Proc Biol Sci* **280**, 20130364. (doi:10.1098/rspb.2013.0364)
- 496 54. Tschirren B, Råberg L, Westerdahl H. 2011 Signatures of selection acting on the innate
497 immunity gene Toll-like receptor 2 (TLR2) during the evolutionary history of rodents. *J.*
498 *Evol. Biol.* **24**, 1232–1240. (doi:10.1111/j.1420-9101.2011.02254.x)

- 499 55. Tschirren B. 2015 *Borrelia burgdorferi* sensu lato infection pressure shapes innate immune
500 gene evolution in natural rodent populations across Europe. *Biol. Lett.* **11**, 20150263.
501 (doi:10.1098/rsbl.2015.0263)
- 502 56. Bartolomucci A. 2007 Social stress, immune functions and disease in rodents. *Front.*
503 *Neuroendocrinol.* **28**, 28–49. (doi:10.1016/j.yfrne.2007.02.001)
- 504 57. Bian JH, Du SY, Wu Y, Cao YF, Nie XH, He H, You ZB. 2015 Maternal effects and
505 population regulation: Maternal density-induced reproduction suppression impairs
506 offspring capacity in response to immediate environment in root voles *Microtus*
507 *oeconomus*. *J. Anim. Ecol.* **84**, 326–336. (doi:10.1111/1365-2656.12307)
- 508 58. Forbes KM, Mappes T, Sironen T, Strandin T, Stuart P, Meri S, Vapalahti O, Henttonen
509 H, Huitu O. 2016 Food limitation constrains host immune responses to nematode
510 infections. *Biol. Lett.* **12**, 20160471.
- 511 59. Jirotkul M. 1999 Population density influences male–male competition in guppies. *Anim.*
512 *Behav.* **58**, 1169–1175. (doi:10.1006/anbe.1999.1248)
- 513 60. Hughes NK, Banks PB. 2016 Olfactory contacts mediate plasticity in male aggression
514 with variable male density. *J. Mammal.* **97**, 444–454. (doi:10.1093/jmammal/gyv188)
- 515 61. Milner RNC, Jennions MD, Backwell PRY. 2012 Keeping up appearances: male fiddler
516 crabs wave faster in a crowd. *Biol. Lett.* **8**, 176–178. (doi:10.1098/rsbl.2011.0926)
- 517 62. Judge PG, De Waal FBM. 1993 Conflict avoidance among rhesus monkeys: Coping with
518 short-term crowding. *Anim. Behav.* **46**, 221–232. (doi:10.1006/anbe.1993.1184)
- 519 63. Kokko H, Brooks R, Jennions MD, Morley J. 2003 The evolution of mate choice and

- 520 mating biases. *Proc. R. Soc. B Biol. Sci.* **270**, 653–664. (doi:10.1098/rspb.2002.2235)
- 521 64. Andersson M, Simmons LW. 2006 Sexual selection and mate choice. *Trends Ecol. Evol.*
522 **21**, 296–302. (doi:10.1016/j.tree.2006.03.015)
- 523 65. Oksanen TA, Koivula M, Koskela E, Mappes T. 2007 The cost of reproduction induced
524 by body size at birth and breeding density. *Evolution (N. Y.)*. **61**, 2822–2831.
525 (doi:10.1111/j.1558-5646.2007.00245.x)
- 526 66. Ostfeld RS. 1985 Limiting resources and territoriality in microtine rodents. *Am. Nat.* **126**,
527 1–15. (doi:10.1086/660279)
- 528 67. Wolff JO. 1993 Why are female small mammals territorial? *Oikos* **68**, 364–370.
529 (doi:10.2307/3544853)
- 530 68. Cayol C, Koskela E, Mappes T, Siukkola A, Kallio ER. 2017 Temporal dynamics of the
531 tick *Ixodes ricinus* in northern Europe: epidemiological implications. *Parasites and*
532 *Vectors* **10**, 166. (doi:10.1186/s13071-017-2112-x)
- 533 69. Bunikis J, Tsao J, Luke CJ, Luna MG, Fish D, Barbour AG. 2004 *Borrelia burgdorferi*
534 infection in a natural population of *Peromyscus leucopus* mice: a longitudinal study in an
535 area where Lyme Borreliosis is highly endemic. *J. Infect. Dis.* **189**, 1515–1523.
536 (doi:10.1086/382594)
- 537 70. Jonsson P, Hartikainen T, Koskela ESA, Mappes T. 2002 Determinants of reproductive
538 success in voles: Space use in relation to food and litter size manipulation. *Evol. Ecol.* **16**,
539 455–467. (doi:10.1023/A:1020854525220)
- 540 71. Desy EA, Batzli GO, Liu J. 1990 Effects of food and predation on behaviour of prairie

- 541 voles: a field experiment. *Oikos* **58**, 159–168. (doi:10.2307/3545423)
- 542 72. White CR, Seymour RS. 2003 Mammalian basal metabolic rate is proportional to body
543 mass^{2/3}. *Proc. Natl. Acad. Sci.* **100**, 4046–4049. (doi:10.1073/pnas.0436428100)
- 544 73. Koivula M, Koskela E, Mappes T, Oksanen T a. 2003 Cost of reproduction in the wild:
545 Manipulation of reproductive effort in the bank vole. *Ecology* **84**, 398–405.
- 546 74. Blackwell AD, Tamayo MA, Beheim B, Trumble BC, Stieglitz J, Hooper PL, Martin M,
547 Kaplan H, Gurven M. 2015 Helminth infection, fecundity, and age of first pregnancy
548 inwomen. *Science (80-.)*. **350**, 6–9.
- 549 75. Minchella DJ, Loverde PT. 1981 A cost of increased early reproductive effort in the snail
550 *Biomphalaria glabrata*. *Am. Nat.* **118**, 876–881. (doi:10.1086/283879)
- 551 76. Weil ZM, Martin LB, Workman JL, Nelson RJ. 2006 Immune challenge retards seasonal
552 reproductive regression in rodents: evidence for terminal investment. *Biol. Lett.* **2**, 393–
553 396. (doi:10.1098/rsbl.2006.0475)
- 554 77. Clutton-Brock TH. 1984 Reproductive effort and terminal investment in iteroparous
555 animals. *Am. Nat.* **123**, 212–229. (doi:10.1086/284198)
- 556 78. Tersago K, Crespín L, Verhagen R, Leirs H. 2012 Impact of Puumala virus infection on
557 maturation and survival in bank voles: a capture-mark-recapture analysis. *J. Wildl. Dis.*
558 **48**, 148–156. (doi:10.7589/0090-3558-48.1.148)
- 559 79. Velando A, Drummond H, Torres R. 2006 Senescent birds redouble reproductive effort
560 when ill: confirmation of the terminal investment hypothesis. *Proc. R. Soc. B Biol. Sci.*
561 **273**, 1443–1448. (doi:10.1098/rspb.2006.3480)

- 562 80. Norrdahl K, Korpimäki E. 1998 Does mobility or sex of voles affect risk of predation by
563 mammalian predators? *Ecology* **79**, 226–232. (doi:10.2307/176877)
- 564 81. Hofmeester TR, Jansen PA, Wijnen HJ, Coipan EC, Fonville M, Prins HHT, Sprong H,
565 van Wieren SE. 2017 Cascading effects of predator activity on tick-borne disease risk.
566 *Proc. R. Soc. B Biol. Sci.* **284**, 20170453. (doi:10.1098/rspb.2017.0453)
- 567 82. Cayol C, Giermek A, Gomez-Chamorro A, Hytönen J, Kallio ER, Mappes T, Salo J,
568 Voordouw MJ, Koskela E. 2018 Data from: *Borrelia afzelii* alters reproductive success in
569 a rodent host. *JYX*. (<http://urn.fi/URN:NBN:fi:jyu-201806133148>).
- 570

571 **Figure captions**

572 Figure 1. The estimated probability of reproduction for a bank vole (\pm 95% CI) depends on their
573 *B. afzelii* infection treatment (uninfected individuals in black, N = 68, infected individuals in
574 grey, N = 68) and their body size (measured as the body mass before injection). In small bank
575 voles, there is no effect of *B. afzelii* infection on breeding probability. In large bank voles, by
576 contrast, uninfected individuals have higher breeding probability than infected individuals. The
577 observed values are shown with open circles.

578

579 Figure 2. The estimated reproductive success of male bank voles depends on the interaction
580 between *B. afzelii* infection (uninfected individuals in black, N = 34, infected individuals in grey,
581 N = 34) and population density. (A) Predicted proportion of females successfully fertilised by a
582 male bank vole (\pm 95% CI) as a function of infection and density. (B) Predicted proportion of
583 offspring sired by a male bank vole (\pm 95% CI) as a function of infection and density (Table S2).
584 The observed values are shown with open circles.

585

586 Figure 3. The estimated home range (in m²) of male bank voles in the enclosures (\pm 95% CI)
587 depends on the interaction between *B. afzelii* infection (uninfected individuals in black, N = 13,
588 infected individuals in grey, N = 14) and population density. At low population density,
589 uninfected males have much larger home ranges than *B. afzelii*-infected males. At high
590 population density, infection with *B. afzelii* does not affect the home range of male bank voles
591 (Table S2). The observed values are shown with open circles.

592

593 Figure 4. Estimated parturition delay in female bank voles (\pm 95% CI) depends on *B. afzelii*
594 infection (uninfected individuals in black, N = 23, infected individuals in grey, N = 22) and
595 population density (Table 1 and S4). The observed values are shown with open circles.

596

597 **Table caption**

598 Table 1: Selected final models for reproductive success and spacing behaviour in bank voles.

599 BM: centred value of body mass before injection; HW: centred value of head width before
600 injection, low: low population density, inf: infected bank voles; σ^2 is the variance attributable to
601 random effect, SD is the standard deviation, SE is the standard error. **Significant effects are**
602 **shown in bold.**