

**This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.**

**Author(s):** Wiberg, R. A. W.; Tyukmaeva, V.; Hoikkala, A.; Ritchie, M. G.; Kankare, M.

**Title:** Cold adaptation drives population genomic divergence in the ecological specialist, *Drosophila montana*

**Year:** 2021

**Version:** Published version

**Copyright:** © 2021 the Authors

**Rights:** CC BY 4.0

**Rights url:** <https://creativecommons.org/licenses/by/4.0/>

**Please cite the original version:**

Wiberg, R. A. W., Tyukmaeva, V., Hoikkala, A., Ritchie, M. G., & Kankare, M. (2021). Cold adaptation drives population genomic divergence in the ecological specialist, *Drosophila montana*. *Molecular Ecology*, 30(15), 3783-3796. <https://doi.org/10.1111/mec.16003>

# Cold adaptation drives population genomic divergence in the ecological specialist, *Drosophila montana*

R. A. W. Wiberg<sup>1</sup>  | V. Tyukmaeva<sup>2</sup>  | A. Hoikkala<sup>2</sup>  | M. G. Ritchie<sup>1</sup>  |  
M. Kankare<sup>2</sup> 

<sup>1</sup>Centre for Biological Diversity, School of Biology, University of St Andrews, St Andrews, UK

<sup>2</sup>Department of Biological and Environmental Science, University of Jyväskylä, Jyväskylä, Finland

## Correspondence

R. A. W. Wiberg, Centre for Biological Diversity, School of Biology, University of St Andrews, St Andrews, KY16 9TH, Scotland, UK.

Email: raw.wiberg@protonmail.com

## Present address

R. A. W. Wiberg, Department of Environmental Sciences, Zoological Institute, University of Basel, Basel, Switzerland  
V. Tyukmaeva, Centre d'Ecologie, Fonctionnelle et Evolutive, CNRS, Montpellier, France

## Funding information

Ella ja Georg Ehrnroothin Säätiö; Academy of Finland, Grant/Award Number: 267244, 268214 and 322980; Natural Environment Research Council, Grant/Award Number: NE/L501852/1 and NE/P000592/1

## Abstract

Detecting signatures of ecological adaptation in comparative genomics is challenging, but analysing population samples with characterised geographic distributions, such as clinal variation, can help identify genes showing covariation with important ecological variation. Here, we analysed patterns of geographic variation in the cold-adapted species *Drosophila montana* across phenotypes, genotypes and environmental conditions and tested for signatures of cold adaptation in population genomic divergence. We first derived the climatic variables associated with the geographic distribution of 24 populations across two continents to trace the scale of environmental variation experienced by the species, and measured variation in the cold tolerance of the flies of six populations from different geographic contexts. We then performed pooled whole genome sequencing of these six populations, and used Bayesian methods to identify SNPs where genetic differentiation is associated with both climatic variables and the population phenotypic measurements, while controlling for effects of demography and population structure. The top candidate SNPs were enriched on the X and fourth chromosomes, and they also lay near genes implicated in other studies of cold tolerance and population divergence in this species and its close relatives. We conclude that ecological adaptation has contributed to the divergence of *D. montana* populations throughout the genome and in particular on the X and fourth chromosomes, which also showed highest interpopulation  $F_{ST}$ . This study demonstrates that ecological selection can drive genomic divergence at different scales, from candidate genes to chromosome-wide effects.

## KEYWORDS

chill coma recovery time, cline populations, cold tolerance, CTmin, *D. montana*, environmental adaptation, genomic divergence

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2021 The Authors. *Molecular Ecology* published by John Wiley & Sons Ltd.

## 1 | INTRODUCTION

The geographic structure of a species is a result of its phylogeographic history, influenced by past and present dispersal, population demography, and selection. Obtaining genome-wide data on genetic polymorphisms across multiple populations of a species is becoming relatively easy, but interpreting the patterns of geographic variation in such data and identifying genes which vary primarily due to selection remains challenging. Often simple "outlier" approaches using genome scans which measure genetic differentiation such as  $F_{ST}$  or Dxy are adopted, but results are difficult to interpret due to confounding effects of selection, drift and population structure, or genomic features such as inversions and other causes of variation in recombination rate (Cruickshank & Hahn, 2014; Noor & Bennett, 2009; Ravinet et al., 2017; Wolf & Ellegren, 2016). If environmental data are available, we can use associations with such factors to help identify loci where genetic differentiation covaries with this environmental variation. Some genome scan methods can incorporate environmental variation and simultaneously fit effects for covariance with environmental factors, while controlling for effects of population demography (Foll & Gaggiotti, 2008; de Villemereuil & Gaggiotti, 2015). This approach has successfully identified genetic variation associated with altitude in humans, among other examples (Foll et al., 2014; Gautier, 2015; de Villemereuil & Gaggiotti, 2015) and has become a useful approach to investigate the ecological adaptations underlying population divergence.

Clinal patterns of variation in phenotypes or gene frequencies have a long history of being used to infer selection along ecotones, and analyses of cline shape can sometimes identify loci under direct selection from others showing clinal variation for other reasons, such as phylogeographic history (Barton & Gale, 1993). Such studies can be very powerful, especially when independent parallel clines are available. For example, Kolaczowski et al., (2011) sampled isofemale lines from extremes of a cline in Australian populations of *D. melanogaster*, and found many genes implicated in clinally varying phenotypes to show highest differentiation. Also, Bergland et al., (2014), and Kapun et al., (2016) sampled North American clines in *D. melanogaster* and *D. simulans* over several years to uncover clinal variation at a genome scale. Bergland et al., (2014) also found consistent fluctuations in allele frequencies for a population sampled over several seasons, indicating a regular response to seasonally varying selection pressures. On the other hand, Machado et al., (2015) concluded that migration and gene flow play a greater role than adaptation in the overall clinality of genomic variants in *D. simulans* than *D. melanogaster*. While the two species share a significant proportion of the genes showing clinal variation, their differences in overwintering ability, migration and population bottlenecks probably act as additional drivers of differences in patterns of variation between them (Machado et al., 2015). Similar studies of clinal variation in phenotypes and allele frequencies have also been carried out in other insects (Paolucci et al., 2016), plants (Bradbury et al., 2013; Chen et al., 2012), mammals (Carneiro et al., 2013; Hoekstra et al., 2004), fish (Vines et al., 2016), and other organisms (Endler, 1973,

1977; Takahashi, 2015). However, more studies are needed that simultaneously compare patterns of variation in allele frequencies and potentially causal environmental variation or ecologically important traits while controlling for population structure, as such studies are necessary to determine if adaptation to climate is directly driving patterns of genetic differentiation.

Here, we investigated geographic variation at both the phenotypic and genetic levels in *Drosophila montana* samples from two continents. This species has spread around the northern hemisphere (Throckmorton, 1969) and both mtDNA and microsatellite data have revealed genetically distinct Finnish and North American populations (Mirol et al., 2007). Moreover, more recent modelling of genome-wide SNP frequencies suggest that the Finnish-North American split happened around 1.75 Mya and that of the North American populations shortly after that (Garlovsky et al., 2020). It is one of the most cold-tolerant *Drosophila* species (Kellermann et al., 2012) and the basic cold tolerance of *D. montana* flies can increase towards the cold seasons through two mechanisms, photoperiodic reproductive diapause (Vesala & Hoikkala, 2011) and cold-acclimation induced by a decrease in day length and/or temperature (Kauranen et al., 2019; Vesala, Salminen, Kostal, et al., 2012; Vesala, Salminen, Laiho, et al., 2012). *D. montana* populations have been found to show clinal variation in the critical day length required for diapause induction (CDL; Lankinen et al., 2013; Tyukmaeva et al., 2011). There is also a correlation between CDL and latitudinally covarying climatic factors such as the mean temperature of the coldest month (Tyukmaeva et al., 2020). In addition, *D. montana* populations from different geographic regions show variation in their courtship cues and mate choice (Klappert et al., 2007; Routtu et al., 2007), which has led to partial reproductive isolation between some distant populations (Jennings et al., 2011, 2014). At the genetic level, differential gene expression studies have identified candidate genes underlying diapause (Kankare et al., 2010, 2016), perception of day length (Parker et al., 2016), and cold acclimation (Parker et al., 2015). Furthermore, a quasi-natural selection experiment for shorter CDL, accompanied by a decrease in cold-tolerance, induced widespread changes in loci with potential roles with these traits (Kauranen et al., 2019). Finally, population genomic analyses have identified several outlier loci when examining differentiation between North American and European populations (Parker et al., 2018). All this makes *D. montana* an interesting example of nascent speciation, potentially influenced by ecological adaptation.

Here, we sought to ask to what extent patterns in the genomic divergence of *D. montana* populations across two continents are correlated with climatic variation and phenotypic responses to cold adaptation. We performed pooled whole-genome sequencing (pool-seq) on six different populations and used Bayesian methods to examine the association between genomic differentiation between populations and environmental variables across both continents. We also phenotyped populations for two different cold tolerance measures; critical thermal minimum (CTmin), and chill coma recovery time (CCRT), and investigated the associations between them and the genetic and climatic data. Ultimately, we

asked if the genomic loci showing an association between genetic and environmental differentiation also showed association with population differentiation in cold tolerance phenotypes, and examined the possible overlap between the set of genes close to candidate SNPs with sets of candidate genes from previous studies of cold adaptation in *D. montana*. If population differentiation is driven by ecological selection then we would predict the extreme cold adaptation of *D. montana* to have left a signature of genomic divergence associated with environmental and phenotypic differentiation across these loci.

## 2 | MATERIALS AND METHODS

### 2.1 | Sample collections and DNA extraction


We collected samples of 49–50 wild-caught flies from six *D. montana* populations from a range of latitudes from 66°N to 38°N in the spring of 2013 or 2014. Four of these populations represented a range of latitudes in North America (N.A.), and two populations were from a range in Finland (Figure 1; Table 1). Samples of wild-caught flies from the six populations were stored in ethanol (the male/female ratio varied across samples; Table 1) and DNA of individual flies was extracted using CTAB solution and phenol-chloroform-isoamylalcohol purifications in 2016. Genomic DNA was extracted from individual flies and quantified using Qubit (Thermo Fisher Scientific), and an equal amount of DNA from each individual (50 ng) was pooled into the final sample. Sequencing was performed at the Finnish Functional Genomics Centre in Turku, Finland ([www.btk.fi/](http://www.btk.fi/)

functional-genomics) on the Illumina HiSeq3000 platform (paired-end reads, read length = 150 bp, estimated coverage ~121x).

### 2.2 | Phenotyping

We measured the critical thermal minimum (CT<sub>min</sub>) and chill coma recovery time (CCRT) of flies from six populations. Fly samples for these tests were collected for five populations (Seward, Terrace, Ashford, Crested Butte, and Korpilahti) from mass-bred populations that have been maintained in the laboratory since 2013–2014. For the Oulanka population, flies were collected from three isofemale strains (established in 2014), because the mass-bred population had been contaminated by another species. The mass-bred populations were originally established from F4 progenies of 20 isofemale strains each (400 flies) and have been maintained in constant temperature ( $19 \pm 1^\circ\text{C}$ ) and light (24 h of light, to prevent the flies from entering diapause) regimes for about 20–25 generations before the experiment. All flies were supplied with fresh malt medium in half-pint bottles every week (Lakovaara, 1969). The newly emerged flies were collected using light CO<sub>2</sub> anaesthesia within 24 h after emergence, separated by sex and placed in malt-vials in the same conditions until sexual maturity (20–21 days old) and were then used in CT<sub>min</sub> and CCRT tests. The same individual flies were first assayed for CT<sub>min</sub> and then for CCRT, the flies were not anaesthetised before these tests.

We assayed a total of 328 females and 302 males for CT<sub>min</sub> and CCRT. These assays were done in batches of between 22 and 30 flies, and split evenly by sex (21 batches in total). Between 32 and 46 (mean 39) flies per population per sex were tested (for the



**FIGURE 1** Maps of all *D. montana* populations in this study. Panels show population locations from (a) Finland and (b) North America showing the locations of all populations sampled. Labelled, blue circles give the locations of populations sampled for phenotyping and sequencing

**TABLE 1** The sources of genomic samples (coordinates and the name of the nearest town), altitude of the sampling site, the year in which sampling was performed, and the number of males and females sampled (M/F) for each pool

Source	Sampling site	Year	M/F
USA, Alaska	Seward 60°9'N; 149°27'W Altitude 35 m	2013	30/20
Canada, British Columbia	Terrace 54°27'N; 128°34'W Altitude 217 m	2014	22/27
USA, Washington	Ashford, 46°45'N; 121°57'W Altitude 573 m	2013	16/34
USA, Colorado	Crested Butte 38°54'N; 106°57'W Altitude 2,900 m	2013	36/13
Finland	Oulanka 66°22'N; 29°20'E Altitude 337 m	2013	25/25
Finland	Korpilahti 62°20'N; 25°34'E Altitude 133 m	2013	27/23

Oulanka isofemale strains, between 15 and 39 flies per strain per sex were used). CTmin tests are based on detecting the temperature (CTmin) at which flies lose neuromuscular function and enter reversible state of chill coma (Andersen et al., 2015). In these tests, the flies were placed into tubes sealed with parafilm and submerged into a 30% glycol-water mixture in Julabo F32-HL chamber. The temperature was decreased at the rate of 0.5°C per min (from 19°C to -6°C) and CTmin was determined by eye, as the temperature at which a fly was unable to stand on its legs. Immediately after the CTmin test, the temperature was set to -6°C and the flies were left in this temperature for 16 h. Vials were then quickly taken out of the glycol-water bath and the flies' CCRT was determined as the time required for the flies to recover from chill coma and stand on their legs. The ambient room temperatures was recorded during trials but in an initial analysis including source population and room temperature there was no statistically significant effect of room temperature on CCRT ( $F_{5,569} = 0.72, p = .4$ ), this variable was therefore left out of further analyses.

To investigate population differences in CTmin and CCRT phenotypes we fit general additive mixed models (GAMMs) in R (v. 3.6.3; R Development Core Team, 2020) using the "mgcv" R package (Wood, 2004). In simple linear models including population, sex, and experimental batch as a fixed effects, experimental batch had an effect on CTmin ( $F_{20,533} = 2.8, p < .01$ ), although compared to population ( $F_{5,533} = 12, p < .01$ ), and sex ( $F_{1,533} = 11.5, p < .01$ ) this effect was not strong. Experimental batch had a stronger effect on CCRT ( $F_{20,533} = 2.83, p < .01$ ), compared to population ( $F_{5,533} = 4.5, p < .01$ ) and sex ( $F_{1,533} = 1.5, p < .22$ ). We therefore included it as a random effect in GAMM analyses. The full GAMMs included altitude, latitude, and sex as fixed effects and experimental batch

as a random effect. We used a cubic regression spline as the basis smoothing function for both altitude and latitude. The raw data for all phenotyping are given in Table S1. The full models are shown in the Supporting Information (Supplementary Material).

## 2.3 | Bioclimatic variables and population geography

We obtained representative climate data from the WorldClim database (Hijmans et al., 2005) for each *D. montana* population sampled for the pool-seq (see above), as well as for 18 additional populations of this species (Table S1; Tyukmaeva et al., 2020). We downloaded the climate data and extracted the values corresponding to population coordinates using the R package "raster" (version 2.5-8; Hijmans et al., 2016). In total this amounts to 55 bioclimatic variables for each population (Table S1). To reduce the number of variables in the data set a principle components analysis (PCA) was performed using the "PCA()" function from the R package "FactoMineR" (version 1.28; Lê et al., 2008). Principle components were kept for further analysis if their eigenvalues were >1. PCA scores for each population were z-transformed using the "scale()" function in base R. Additionally, CTmin and CCRT were summarised to a mean value for each population. In total, this gives four "environmental" variables measured for each population (PC1, PC2, CTmin, and CCRT).

## 2.4 | Mapping, SNP calling and genomic analysis

Quality of raw reads was checked with FASTQC (v. 0.11.5; Andrews, 2015) and reads were trimmed using TRIMMOMATIC (v. 0.32; Bolger et al., 2014; see Supporting Information (Supplementary Material) for full trimming parameters). Trimmed reads were mapped to the *D. montana* reference genome (Parker et al., 2018) using BWA MEM (v. 0.7.7; Li, 2013) with the default options but keeping only alignments with a mapping quality of >20 following best practice guidelines for pool-seq (Schlötterer et al., 2014). Duplicate alignments were removed with SAMTOOLS RMDUP (v 1.3.1; Li et al., 2009) and regions around indels were realigned using PICARD (version 1.118; Broad Institute), GATK (v. 3.2-2; McKenna et al., 2010) and SAMTOOLS. Separate.bam files for each of the sequenced populations were finally merged using BAMTOOLS (v. 2.4.0; Barnett et al., 2011).

Over 80% of reads were properly mapped and retained in all samples. The mean coverage for Seward samples was nearly twice that of the other samples (Figure S1 and Figure S2). To remove the potential for this difference to cause artefacts in downstream analyses, the.bam files for Seward were downsampled to contain 94.1 million reads (the average across the five remaining populations). Median empirical coverage was between ~62 and 88x (Table S2; Figure S3) and much less variable among the populations, allowing common maximum and minimum thresholds to be set based on the aggregate distribution. Allele counts for each population at each genomic position were obtained with samtools mpileup (version 1.3.1; Li et al.,

2009) using options to skip indel calling as well as ignoring reads with a mapping quality <20 and sites with a base quality <15. This was followed by the heuristic SNP calling software PoolSNP using a minimum count of 5 to call an allele, and a minimum coverage of 37 and a maximum coverage <95th percentile of the scaffold-wide coverage distribution to call a SNP (Kapun et al., 2020). Even if all these filters were passed, an allele was not considered if its frequency was <0.001. Finally, we only considered SNPs on scaffolds >10 kb in length. The final set consisted of 2,190,511 biallelic SNPs that could be placed on scaffolds ordered along the chromosomes and were used in downstream analyses.

To test for an association between the four environmental variables and genetic differentiation we used BAYESENV (v. 1.1; Foll & Gaggiotti, 2008; de Villemereuil & Gaggiotti, 2015). BAYESENV fits a model of  $F_{ST}$  to population differentiation for each locus, incorporating environmental differentiation as a predictor (included as the parameter  $g$ ) while fitting two locus-specific effects, one for environmental selection, and the other for other processes (demography, other types of selection). It therefore controls for confounding effects of population structure/relatedness in testing for an association with environmental variables. BAYESENV was run with five pilot runs of 1000 iterations each, followed by a main chain of 4000 iterations of which 2000 were discarded as burnin. Four MCMC chains were run for each analysis to evaluate convergence of the chains to common parameter estimates. Because of the unbalanced number of males (and the resulting variation in ratios of X:Y chromosomes) in the pools, BAYESENV analyses were performed separately on SNPs that could be assigned to the autosomal linkage groups (chromosomes) and the X chromosome. Raw count data were used for the autosomal data. For X linked SNPs, allele count data were scaled to the known number of X chromosomes in the pool using  $n_{eff}$  the effective sample size taking into account the multiple rounds of binomial sampling inherent to a pool-seq design (Feder et al., 2012; Kolaczowski et al., 2011).  $N_{eff}$  scales the allele counts at each SNP downwards based on the known number of chromosomes in the pool (see Table 1).

Chains were assessed for convergence with the "coda" R package (v. 0.19-1; Plummer et al., 2006). Convergence was reached across the four chains for most analyses and parameters (potential scale reduction factors (PSRFs) of ~1 in a Gelman-Rubin diagnostic test; Figures S4–S7), except for analyses of autosomal SNPs and PC2 as the environmental variable which showed mild convergence problems (PSRF = 1.71), although parameter estimates agreed well with the other chains. Thus, this first chain (black line in Figures S4–S7) was discounted for all parameters and only estimates from the remaining three chains were used. The union of significant SNPs (using  $q$ -values for the  $g$  parameter, describing the association between genetic differentiation at a locus and environmental differentiation, to control the FDR at 0.05, i.e., SNPs with  $q$ -values <0.05) across these chains were taken as the final candidate SNPs.

Finally, population genetic statistics ( $\pi$ , and Tajima's  $D$ ) were computed in windows of 10 kb with a step size of 5 kb using methods implemented for pool-seq data (Kapun et al., 2020). Windows


contained a mean of  $488.2 \pm 1.2$  SNPs (mean  $\pm$  SE). These statistics were only computed for scaffolds with a length >50 kb. SNP-wise  $F_{ST}$  was computed for each population with the R package "poolfstat" (v. 1.1.1; Hivert et al., 2018) by first computing all pairwise values, and then deriving population specific  $F_{ST}$  values by averaging across all pairwise values where a population was included. We computed 95% confidence intervals for mean SNP-wise  $F_{ST}$  on each chromosome from the distribution of mean  $F_{ST}$  values across 100 bootstrap samples of SNPs. See the Supplementary Material (Supporting Information) for pseudocode commands of the key pipeline steps.

We performed GO term enrichment analyses with DAVID (v. 6.8; Huang et al., 2009a, 2009b) and GOWINDA (v. 1.12; Kofler & Schlötterer, 2012), which accounts for differences in gene length. For DAVID, since the *D. montana* annotation contains information about orthologs in *D. virilis*, we extracted all genes within 10 kb upstream or downstream of a candidate SNP with an ortholog in *D. virilis* and submitted them to DAVID. For analyses with GOWINDA, because the gene-sets need to be given manually, we obtained gene-sets from FuncAssociate3 (Berriz et al., 2009). As above, we considered SNPs within 10 kb of a gene (--gene-definition updownstream 10,000) and performed 1 million simulations to obtain the empirical  $p$ -value distribution. Regulatory elements, such as enhancers, and transcription factor binding sites, can occur up to 1 Mb up- or downstream from a target gene in other species (Chan et al., 2010; Maston et al., 2006; Pennacchio et al., 2013; Werner et al., 2010) but generally lie within 2 kb of a gene region in *D. melanogaster* (Arnosti, 2003), thus 10 kb represents a compromise.

### 3 | RESULTS

#### 3.1 | Cold tolerance measures, bioclimatic variables and population geography


Across individuals, there was no evidence of an association between minimum critical temperature (CTmin; pooled across sexes) and the chill coma recovery time (CCRT;  $cor = 0.08, p = .07$ ). Neither was there evidence for an association between these traits across populations ( $cor = 0.22, p = .68$ ). CTmin was significantly different between sexes, with females being more cold tolerant ( $t = 3.14, p = .002$ ). While latitude appears to have a nonlinear effect on CTmin (Figure 2a and Figure S8A), the effective degrees of freedom (edf = 1) of the partial effect of latitude suggested a largely linear effect after accounting for altitude ( $F = 40.4, p < .001$ ). Altitude had a more complex effect (edf = 1.262,  $F = 24.0, p < .001$ ), but with a strong altitude outlier in Crested Butte (Table 1), this result should be taken with some caution. Moreover, the overall adjusted R-squared was low at only 0.09. For CCRT, only latitude had a significant effect (edf = 1,  $F = 17.2, p < .001$ ; Figure 2b). Taken together, CTmin was lower for populations at higher latitudes as well as for populations at higher altitudes (Figure 2a and Figure S8A). As one would expect, CTmin showed on average lower values (Figure 2a and Figure S8A), and CCRTs were



**FIGURE 2** Population phenotypes. (a) and (b) show the variation in CTmin and CCRT across populations and latitude, respectively. Solid and dashed lines show the predicted values from the partial effect of latitude from the best model (see Results) for males and females, respectively. In (b) although the points are plotted separately for males and females, the best model only included latitude as a covariate. In both panels the points and error bars give the mean and standard errors, respectively. Figure S8 shows the full data for each population

shorter (Figure 2b and Figure S8B), at higher latitudes, meaning that more northern and higher altitude populations show higher cold tolerance.

We performed a principal component analysis (PCA) of the WorldClim climate data for a total of 24 *D. montana* populations, from which we had collected samples and where climate data were available. We wanted to examine whether the populations chosen for the cold tolerance tests and genome sequencing were representative of the range of environmental variation experienced by the species. These analyses identified four principal components (PCs) that together explained about 98% of the variation (Figure 3a and Figure S9). The first two PCs separated the populations roughly by a measure of distance from the coast (PC1) and then by latitude and altitude (PC2). PC1 explained ~54% of the variation (Figure 3a and b) and loaded heavily on climate and biological variables associated with precipitation and temperature such as “mean temperature of coldest



**FIGURE 3** Principle component analysis (PCA). (a) The distribution of all populations along the two first PC axes. The loadings of variables on each axis can be found in Table S3. Blue circles give the populations that have been pool-sequenced for this study, red squares and triangles give the other Finnish and North American populations, respectively. (b) and (c) give PCs 1 and 2 as a function of latitude and altitude, respectively. The legend for altitude in both (b) and (c) is given in (c)

quarter", "precipitation of wettest month", "annual precipitation" (see Figure S9 and Table S3). Meanwhile, PC2 explained ~23% of the variation (Figure 3a and c) and loaded heavily on biological variables that are associated with latitudinal clinality, e.g., "mean diurnal [temperature] range," and "isothermality" which is the diurnal range divided by the mean "annual [temperature] range". The remaining PCs (PC3 and PC4) explained about 11.5% and 5% of the variation, respectively and did not capture as much of the climatic variation, and were therefore not considered further. Latitude (Spearman's rank correlation:  $\rho = -0.50$ ,  $p = .01$ ) but not altitude ( $\rho = -0.19$ ,  $p = .39$ ) correlated significantly with PC1. However, both altitude ( $\rho = -0.50$ ,  $p = 0.02$ ) and latitude ( $\rho = -0.59$ ,  $p = .003$ ) correlated with PC2 (see also Figure 3c). Importantly, these patterns were fairly robust also when performing PCA using only the six populations for which genomic data were collected. Loadings on PC1 were highly comparable, and although correlations with latitude and altitude did not achieve statistical significance, they were similar in magnitude and direction (latitude vs. PC1:  $\rho = 0.43$ , altitude vs. PC1:  $\rho = -0.14$ ,  $p = .8$ ). For PC2 the correlations were also not statistically significant and differed for latitude both in magnitude and direction ( $\rho = 0.6$ ,  $p = .2$ ), while for altitude, the correlation remained similar and marginally non-significant ( $\rho = -0.82$ ,  $p = .06$ ). Moreover, while the range of the sequenced populations covered ~70% of the range of all populations for PC1, they covered only ~50% of the range for PC2 (Figure 3a). This suggests that environmental variation across the six populations selected for sequencing reflects that experienced by all 24 populations at least for the PC1 axis. Therefore, the relationship between environmental variables and genetic differentiation in the samples selected for pool-seq is likely to reflect true patterns across populations of *D. montana*, though power might be somewhat reduced for PC2. For all subsequent analyses, we used the results of the PCA using all 24 populations, and to examine the association between climate and phenotype, we compared these across the six populations. CTmin was positively correlated with PC1 (Pearson's correlation coefficient ( $\text{cor}$ ) = 0.94,  $p < .01$ ) and had a marginally non-significant association with PC2 ( $\text{cor}$  = 0.75,  $p = .09$ ). However, CCRT showed no relationship with either PC1 ( $\text{cor}$  = 0.09,  $p = .87$ ) or PC2 ( $\text{cor}$  = -0.46,  $p = .36$ ) although the small sample sizes ( $N = 6$  in all cases) make reliable conclusions difficult (see Figure S10).

### 3.2 | Genomic analyses

The number of SNPs with a significant ( $q$ -value  $< 0.05$ ) association between population-based  $F_{ST}$ , the two cold tolerance measures (CCRT and CTmin), and the two PCs of the bioclimatic data varied from 312 (chromosome 3, CCRT) to 2480 (chromosome 4, PC2) across the chromosomes (Figure 4). Using PC1 as an environmental variable with BAYESCENV gave a total of 2976 and 1528 SNPs with a  $q$ -value  $< 0.05$  on the autosomes and on the X chromosome, respectively. Interestingly, the distribution across the chromosomes was not random. Using the distribution of all SNPs to obtain expected counts, there was a significant deviation from expectation ( $\chi^2 = 2906.4$ ,

d.f. = 4,  $p < .001$ ). There were many more SNPs than expected on chromosome 4 (1432 vs. 954) and on the X chromosome (1528 vs. 526; Figure 4). Results were similar for PC2 with 6607 and 1861 SNPs with a  $q$ -value  $< 0.05$  on the autosomes and X chromosomes, respectively. Again, there was a significant deviation from the expected distribution of SNPs across the chromosomes ( $\chi^2 = 1681.9$ , d.f. = 4,  $p < .001$ ) with an overrepresentation on the fourth (2480 vs. 1794) and the X chromosomes (1861 vs. 989; Figure 4).


We also used average CTmin values per population in similar analyses and found a total of 2668 and 1272 SNPs with a  $q$ -value  $< 0.05$  on the autosomes and X chromosomes, respectively. The pattern of significant deviations from expected distributions ( $\chi^2 = 2,526.2$ , d.f. = 4,  $p < .001$ ) was also due to an excess on the fourth (1383 vs. 835) and the X chromosomes (1272 vs. 460; Figure 4). Similar results were found for CCRT with a total of 2240 and 1228 SNPs with a  $q$ -value  $< 0.05$ , respectively. Once again, there was a significant deviation from the expected distribution of SNPs ( $\chi^2 = 2825.6$ , d.f. = 3,  $p < .001$ ) with an excess on the fourth (1252 vs. 735) and the X chromosomes (1228 vs. 405; Figure 4). Manhattan plots of the distribution of SNPs across chromosomes are given in Figures S11–S14.

To more closely examine the loci implicated in the four BayScEnv analyses, we identified genes within 10 kb of, or containing the candidate SNPs (Table S4). Overall, there is quite a large overlap among these genes with ~39% (1102 in total) of them being shared by all the four analyses (Table S4, Figure S15). This is far more than would be expected (mean  $\pm$  SE:  $3.6 \pm 0.06$ ) by randomly resampling (without replacement) a number of genes equivalent to that associated with top SNPs for each BayScEnv analysis from the *D. montana* annotation, then computing the four-way intersection. This large observed overlap therefore probably reflects a genuine overlap among the underlying top SNPs (data not shown). Some (147, ~13%) of these common genes are novel to *D. montana* (i.e., not annotated in *D. virilis* or in other *Drosophila* spp.) and therefore have no annotation, but 955 have an identifiable *D. virilis* ortholog (Table S4). The second largest set are those genes unique to the analysis of PC2 as an environmental covariate (Figure S15). We performed functional enrichment analyses with DAVID keeping all annotation clusters with an enrichment score  $> 1.3$  (corresponding to an average corrected  $p$ -value of .05). This revealed several common categories of genes associated with the climatic variables and population phenotypes (Table S5). For example, terms associated with membrane and transmembrane structures, immunoglobulins, HAD hydrolase and nucleotide binding were enriched in most of the variables (Table S5). Interestingly, there were also several gene ontology categories that were only enriched in one of the variables, such as glycoside and ATPase hydrolase in CCRT and ion channels and transport, as well as metal binding in PC1 (Table S5). GOWINDA analyses revealed significant enrichment of GO terms (after accounting for variation in gene lengths and correcting for multiple testing) only for genes near SNPs associated with variation in CCRT across populations. Interestingly, the term "carbohydrate derivative binding" was identified among the most enriched terms (Table S6), which agrees closely with some terms identified using DAVID for the same variable (Table



S5). Similarly, for PC2, "nucleotide binding" was among the top scoring enriched terms in both GOWINDA and DAVID analyses, though this was not significant after controlling for multiple testing in GOWINDA (Table S5, Table S6).

We then compared loci near candidate SNPs with genes implicated in previous studies of climatic adaptation in *D. montana* including gene expression studies of traits connected to diapause and cold-tolerance (Kankare et al., 2010, 2016; Parker et al., 2015, 2016). Additionally, several candidate genes have been identified near the most significantly differentiated SNPs among *D. montana* populations from Oulanka (Finland), and from North American populations in Colorado and Vancouver (Parker et al., 2018). Finally, quasi-natural selection experiments identified several genes within 10 kb of SNPs responding to selection for a shorter CDL for diapause induction (Kauranen et al., 2019). We tested for an overlap between the total set of genes within 10kb of outlier SNPs from all of the BayScEnv runs ( $N = 2694$ ) and the candidate gene sets identified in earlier studies (see Table S7 for the gene sets and studies used). We computed a bootstrap distribution of overlaps by sampling 2694



**FIGURE 4** The observed counts of candidate SNPs (i.e., SNPs with a  $q$ -value  $< 0.05$ ) across chromosomes for each environmental variable (see Section 2). The total number of SNPs, and the proportion of all SNPs, on each chromosome are given below each set of bars. The expected counts on each chromosome, obtained from the proportions of all SNPs across chromosomes, are shown as points aligned with each bar


Reference	N genes	N overlaps	Mean (95% CI) resampled N overlaps	Empirical $p$ -value
Kankare et al., (2010)	14	8	2.77 (0, 6)	.0004
Parker et al., (2015)	42	13	6.28 (2, 11)	.001
Kankare et al., (2016)	3,929	946	773.14 (732, 814)	<.0001
Parker et al., (2016)	130	27	18.72 (11, 26)	.01
Parker et al., (2018)	2,114	629	414.7 (382, 448)	<.0001
Kauranen et al., (2019)	1,751	402	344.64 (314, 375)	<.0001

Shown are the citations for the original study, the number of genes identified from the original study, the number of overlapping genes with the candidate gene set from this study, and the mean and 95% CI from 10,000 resampled sets of candidate genes.

random genes from the *D. montana* annotation ( $N = 13,683$ ). For each of the gene sets from previous studies this was done 100 times and the distribution compared to the empirical overlap. Results are given in Table 2. In all the cases the empirical overlap was greater than expected by chance with empirical  $p$ -values  $< 0.05$  (ranging from  $< .0001$  to  $.01$ ; Table 2). The only gene that was found in all five of the previous studies used and in the comparison here is called sidestep II (side-II; Table S8). Unfortunately, there is no information available about the biological processes or molecular functions connected to it. Moreover, from 44 other genes that were common to four of our previous studies and this study (Table S8) most (27) have an ortholog in *D. melanogaster*. These genes have molecular functions such as transmembrane signaling or transporter, acetylcholinesterase, ATP binding, protein serine/threonine kinase, carboxylic ester hydrolase, or Rho guanyl-nucleotide exchange factor activity (Thurmond et al., 2019). Many of the genes are also connected to metal ion, nucleic acid or zinc ion binding (Table S8). After identifying information on molecular or biological function and Interpro domains, eventually only five genes remained for which there was no information available (Table S8).

Examination of population genetic parameters identified the Crested Butte population as anomalous. The distribution of Tajima's  $D$  is centred close to zero in most populations, being slightly more negative in North American populations (Figure S16A). However, Crested Butte is an outlier with a greatly reduced genome-wide Tajima's  $D$  (Figure S16A). Similarly, diversity ( $\pi$ ) is also lower in this population than in other populations. There is no overall relationship between latitude and  $\pi$  (Spearman's  $\rho = 0.14$ ,  $N = 6$ ,  $p = .8$ ; Figure S16B) but there is a strong correlation between latitude and Tajima's  $D$  which is influenced by this population (with Crested Butte:  $\rho = 0.88$ ,  $N = 6$ ,  $p = .03$ ; without Crested Butte:  $\rho = 0.8$ ,  $N = 5$ ,  $p = .13$ ). Although Crested Butte occurs at a much higher altitude ( $> 2800$  m) than other populations neither Tajima's  $D$  nor  $\pi$  correlated significantly with altitude (Tajima's  $D$ :  $\rho = -0.6$ ,  $N = 6$ ,  $p = .24$ ,  $\pi$ :  $\rho = -0.6$ ,  $N = 6$ ,  $p = .24$ ). Furthermore,  $F_{ST}$  was similar across all populations and chromosomes with the exception of Crested Butte which remained an outlier with high  $F_{ST}$  (Figure 5). Bootstrapped 95% confidence intervals are presented to give a guide to statistical significance of differences among populations and confirm that Crested Butte has high  $F_{ST}$  (Figure 5). Finally,  $F_{ST}$  was always highest

**TABLE 2** The overlap of the union of genes within 10 kb of SNPs associated with PC1, PC2, CTmin, and CCRT and previous candidate gene sets (see Table S7)



**FIGURE 5** Mean, SD, and confidence intervals of the  $F_{ST}$  values across SNPs on each chromosome and population.  $F_{ST}$  was summarised for each SNP by computing all pairwise values then deriving population specific  $F_{ST}$  values by averaging across all pairwise values containing that population. The 95% confidence intervals (given by the error bars) are derived from 100 bootstrap samples of SNPs on each chromosome

on chromosome 4 and the X chromosome, complementing the results seen in *BAYESENV* analyses (Figure 4 c.f. Figure 5) and as expected if ecological selection is influencing genomic divergence.

## 4 | DISCUSSION

Detecting genomic signatures of climatic adaptation is an important, but challenging, task. Here we use multiple sources of evidence to study ecological adaptation and population divergence in a highly cold tolerant species of *Drosophila*, *D. montana*. This species is characterised by a wide circumpolar distribution extending to high latitudes both in North America and Europe, and to high altitudes in the southern part of its range in the Rocky Mountains of North America. These habitats impose extreme seasonal and climatic selective pressure. We collected bioclimatic data from 24 populations along a latitudinal gradient of about 2900 km in North America, and six populations from a gradient of 720 km in Finland. We characterised population level cold-tolerance for six populations from these ranges using two methods, critical thermal minimum (CTmin) and chill coma recovery time (CCRT) and show that CTmin were lower and CCRT shorter in higher latitude populations, as one would expect. Thus, northern populations are more cold-tolerant. Finally, we performed pool-seq of these six populations to investigate the association between genomic and environmental differentiation.

The two methods examining cold tolerance gave somewhat different results, as CTmin, but not CCRT, differed significantly between sexes, with females having lower CTmin than males. In an earlier study investigating seasonal changes in *D. montana* CCRTs, only one out of six comparisons showed a significant difference between sexes (Vesala, Salminen, Kostal, et al., 2012; Vesala, Salminen, Laiho,

et al., 2012) and Gibert et al., (2001) did not detect sex-specific differences in CCRT in any of 84 *Drosophila* species they studied. However, several studies of *D. melanogaster* have detected shorter CCRT in females than in males, suggesting that females are more cold tolerant than males (Andersen et al., 2015; Bauerfeind et al., 2014; David et al., 1998), perhaps related to their greater body mass (Wilder et al., 2010). Consequently, the extent and adaptive significance of sex-specific differences in CCRT in *Drosophila* remains unclear.

We derived principal components to summarise WorldClim climatic variables using data from all the 24 populations. The first principal component (PC1) separated populations approximately by a measure of “distance inland” and loaded heavily on climate and variables associated with precipitation and temperature. These results follow the geographic distribution of the populations, for example, the population with highest values for PC1 is Ashford, which is on the Pacific coast and receives most rain, but also experiences warm summers and mild winters. Principal component 2 (PC2), loaded heavily on bioclimatic variables associated with latitudinal clinality, which also mapped onto the populations intuitively as those with higher values on PC2 also occurred at higher latitudes. Interestingly, CTmin values were positively correlated with PC1, but not with PC2, while CCRT showed no relationship with either of these components. This suggests that CTmin and CCRT measure at least slightly different biochemical or physiological mechanisms (Findsen et al., 2014; MacMillan et al., 2012), and could hence be correlated with different climatic variables and also show relatively weak correlation to each other (Andersen et al., 2015). Indeed our results found no significant correlation in CTmin and CCRT across populations. It is also interesting to note that high altitude populations have very similar values on PC1 and PC2 to high latitude populations, most probably reflecting the similar climates in these populations. These similarities in climate are probably also underlying the similarities in CTmin between the high altitude population (Crested Butte) and the high latitude populations from Finland.

The Bayesian analysis identified SNPs showing an association of genetic differentiation with climatic and phenotypic variation. The extent to which the loci associated with the phenotypes and adaptations to different climatic conditions are shared indicates that these are closely associated in influencing genome evolution. We found that genes near SNPs showing a significant association between genetic and climatic differentiation overlapped to a large extent with genetic and phenotypic differentiation among populations. The largest intersection set, containing 1102 genes, was the one containing genes near SNPs associated with all the four variables examined (PC1, PC2, CTmin and CCRT). However, PC2 loads heavily on bioclimatic variables relating to latitude, and our analysis using PC2 as a covariate has a large number of private genes (see Figure 4), suggesting that there is also a substantial amount of genetic variation underlying adaptations to latitude unrelated to the phenotypic measures we have quantified. Moreover, this should be viewed with caution because the range of values for PC2 represented by the sequenced samples involve only 50% of the total variation in this axis. Nevertheless, our

study is an excellent example of how strong ecological selection may be detected in genomic studies. In particular, because the Bayesian methods examining both ecological variables and relevant phenotypes gave significant overlap amongst the associated loci, and that these are further associated with more broad genomic differentiation between populations, gives confidence that we are consistently identifying genes associated with ecological selection.

Analyses of the functional annotation of these genes strengthens our conclusions that climate driven adaptation is important. Regions near SNPs associated with climatic variables were enriched for genes previously identified as candidates related to cold tolerance, diapause and responses to changes in day length in *D. montana* (Kankare et al., 2010, 2016; Parker et al., 2015, 2016). Thus, genetic variation across populations of these flies may be largely shaped by differences in ecological and climatic variation. This ecological specialisation may have also contributed to the divergence of *D. montana* from its relatives. Parker et al., (2018) surveyed the rates of molecular evolution in eleven cold tolerant and noncold tolerant species of *Drosophila*. The genes found to be evolving at faster rates in cold-tolerant species were enriched for many of the same functional categories as in our current study, including for example, membrane and transmembrane proteins and immunoglobulins (Parker et al., 2018).

The fact that many of the same annotation terms are enriched in clusters for all environmental variables in DAVID analyses, suggests that similar genes or biochemical pathways are involved in these adaptations. Membrane proteins and lipids are an important determinant of membrane and cuticular permeability at different temperatures, which in turn has an effect on the resistance to desiccation stress in insects (Gibbs, 2002; Stanziano et al., 2015). Importantly, there is evidence for a close link between the desiccation stress response and cold tolerance across species and in *Drosophila* in particular, suggesting an overlap in some of the mechanisms involved (Sinclair et al., 2007). Cold-hardy lines of *D. melanogaster* are known to exhibit elevated lipid metabolism, perhaps in order to allow rapid lipid membrane modification (Williams et al., 2016) in different environmental conditions. Pleckstrin homology (PH) domain was one of the functional clusters found in both the current study and in the species comparison of Parker et al., (2018). This domain is a flexible module of 100–120 amino acids which interacts with a variety of different ligands, composing a protein-protein interaction platform (Scheffzek & Welti, 2012). As changes in the membrane lipid biochemistry form an integral part of the cold tolerance response, genes associated with PH may assist in homeoviscous adaptation namely, alteration of membrane phospholipid composition to maintain fluidity at low temperatures (Sinensky, 1974). Interestingly, the only gene found in all five of our previous studies of cold tolerance in *D. montana* was side-step II (side-II). Unfortunately, there is no information available for the biological processes or molecular functions associated with this gene, but side-II has protein features including immunoglobulin and immunoglobulin-like domain superfamily (Thurmond et al., 2019) and could be involved in immunological processes during cold response of the flies (Parker et al., 2021). Insects are known to produce

a diverse range of antimicrobial peptides and proteins as part of their immune activity against viruses, bacteria, fungi and parasites (Mylonakis et al., 2016) and hence immune responses could be part of a general stress response in cold tolerance (Ferguson et al., 2016; Sinclair et al., 2013). We also find confirmation of some enriched GO terms in GOWINDA analyses that control for variation in gene length, although overlap is not substantial. In summary, our results indicate that some of the same biochemical processes that are targeted by selection on larger evolutionary scales (i.e., across species), are also involved in local adaptation for different populations within a species, providing a rare bridge between the processes of population differentiation and speciation.

At the chromosomal level, we found an overrepresentation of loci associated with ecological selection on chromosomes X (which corresponds to chromosome 2L in *D. melanogaster*) and 4. It is well-known that X chromosomes can generally evolve quickly due to selection on semi-recessive advantageous loci in the hemizygous sex, and smaller effective population size (Charlesworth et al., 1987) and are often most divergent between closely related species (Abbott et al., 2017; Ellegren et al., 2012). However, there is no obvious reason to expect faster divergence in chromosome 4. *D. montana* populations have considerable chromosomal polymorphism (Morales-Hojas et al., 2007; Patterson & Stone, 1952; Stone et al., 1960) and American populations have been classified into several chromosomal forms on the basis of their geographical origin, size and chromosome structure (Throckmorton, 1982). Moreover, both the X chromosome and chromosome 4 of *D. montana* are known to harbour polymorphic inversions (Morales-Hojas et al., 2007; Stone et al., 1960) and such inversions have often been found to vary clinally in other systems (Chen et al., 2012; Kapun et al., 2016; Kolaczowski et al., 2011). Recent genomic data confirms that there are several inversions in both chromosomes X and 4, but we do not yet have any detailed information on their distributions within populations (Poikela et al., unpublished data). The reduced recombination within inversions can independently capture advantageous alleles under selection (Kirkpatrick & Barton, 2006) and divergence is often greater in chromosomes carrying inverted regions and noncolinear regions (Lohse et al., 2015) and such regions may diverge more quickly during speciation with gene flow. Indeed, here we found that the X chromosome and chromosome 4 always have the highest levels of  $F_{ST}$  across all populations, as expected if the ecological selection on loci on these chromosomes influenced overall patterns of genomic diversion, perhaps due to hitchhiking.

Finally, our findings include intriguing result regarding the Crested Butte population which shows reduced genetic variability and a substantial reduction of Tajima's *D* relative to the other populations. This population occurs at a very high altitude (>2800 m) and also shows reproductive incompatibilities with other populations (Jennings et al., 2014). It may have also been bottlenecked during its adaptation to this high altitude, so population expansion or selected sweeps could be prevalent within this population (Garlovsky et al., 2020). Whether ecological specialisation is associated with the spread of incompatibilities is an intriguing possibility.

## 5 | CONCLUSION

Identifying the genetic variation that underlies population divergence and ultimately speciation remains a challenge. Detecting associations between genetic and environmental differentiation at loci across populations has become a popular approach. Here, we apply Bayesian methods to detect such loci across populations of *Drosophila montana*, which is an extremely cold-tolerant *Drosophila* species where we expect strong ecological selection. We identify many genes that are associated with both climate variables and population-level cold-tolerance phenotypes. These genes overlap with candidate genes from other studies of variation in cold-tolerance in *D. montana* and were also overrepresented on chromosomes X and 4 which may be associated with inversion polymorphisms known to be present in these chromosomes. Our study thus provides a clear example of how strong ecological selection can be detected in genome studies, using Bayesian methods to detect local adaptation in combination with studies of ecologically important variables and phenotypes.

### ACKNOWLEDGEMENTS

This work was supported by a combined Natural Environment Research Council and St Andrews 600th Anniversary PhD Studentship grant (NE/L501852/1) to RAWW, The Ella and Georg Ehrnrooth Fellowship to VT, NERC grant (NE/P000592/1) to MGR, and Academy of Finland project 267244 to AH and projects 268214 and 322980 to MK. Bioinformatics and Computational Biology analyses were supported by the University of St Andrews Bioinformatics Unit which is funded by a Wellcome Trust ISSF award (grant 105621/Z/14/Z). We would also like to thank R. Tapanainen and J. Kinnunen for assistance in laboratory work and Oscar E Gaggiotti for advice on BAYESCEMV analyses. We also thank four anonymous reviewers for their valuable comments that improved the manuscript.

### AUTHOR CONTRIBUTIONS

Anneli Hoikkala, Michael G. Ritchie, and Maaria Kankare conceived the study. Venera Tyukmaeva performed the phenotyping experiments. R. Axel W. Wiberg performed all analyses. R. Axel W. Wiberg, Maaria Kankare and Michael G. Ritchie led the writing of the manuscript. All authors contributed to the final version of the manuscript.

### DATA AVAILABILITY STATEMENT

Raw reads have been deposited with NCBI (<https://www.ncbi.nlm.nih.gov>) under the BioProject accession PRJNA588720.

### ORCID

R. A. W. Wiberg  <https://orcid.org/0000-0002-8074-8670>

V. Tyukmaeva  <https://orcid.org/0000-0002-1864-5101>

A. Hoikkala  <https://orcid.org/0000-0001-5407-7992>

M. G. Ritchie  <https://orcid.org/0000-0001-7913-8675>

M. Kankare  <https://orcid.org/0000-0003-1541-9050>

## REFERENCES

- Abbott, J. K., Norden, A. K., & Hansson, B. (2017). Sex chromosome evolution: Historical insights and future perspectives. *Proceedings of the Royal Society B: Biological Sciences*, 284. <https://doi.org/10.1098/rspb.2016.2806>
- Andersen, J. L., Manenti, T., Sorensen, J. G., MacMillan, H. A., Loeschcke, V., & Overgaard, J. (2015). How to assess *Drosophila* cold tolerance: Chill coma temperature and lower lethal temperature are the best predictors of cold distribution limits. *Functional Ecology*, 29, 55–65. <https://doi.org/10.1111/1365-2435.12310>
- Andrews, S. (2015). FASTQC: A quality control tool for high throughput sequence data. Retrieved from <http://www.bioinformatics.bbsrc.ac.uk/projects/fastqc/>
- Arnosti, D. N. (2003). Analysis and function of transcriptional regulatory elements: Insights from *Drosophila*. *Annual Review of Entomology*, 48, 579–602. <https://doi.org/10.1146/annurev.ento.48.091801.112749>
- Barnett, D. W., Garrison, E. K., Quinlan, A. R., Strömberg, M. P., & Marth, G. T. (2011). BAMTOOLS: A C++ API and toolkit for analyzing and managing BAM files. *Bioinformatics*, 27, 1691–1692. <https://doi.org/10.1093/bioinformatics/btr174>
- Barton, N. H., & Gale, K. S. (1993). Genetic analysis of hybrid zones. In R. G. Harrison (Ed.), *Hybrid zones and the evolutionary process* (pp. 13–45). Oxford University Press.
- Bauerfeind, S. S., Kellermann, V., Moghadam, N. N., Loeschcke, V., & Fischer, K. (2014). Temperature and photoperiod affect stress resistance traits in *Drosophila melanogaster*. *Physiological Entomology*, 39, 237–246. <https://doi.org/10.1111/phen.12068>
- Bergland, A. O., Behrman, E. L., O'Brien, K. R., Schmidt, P. S., & Petrov, D. (2014). Genomic evidence of rapid and stable adaptive oscillations over seasonal time scales in *Drosophila*. *PLoS Genetics*, 10, e1004775. <https://doi.org/10.1371/journal.pgen.1004775>
- Berriz, G. F., Beaver, J. E., Cenik, C., Tasan, M., & Roth, F. P. (2009). Next generation software for functional trend analysis. *Bioinformatics*, 25, 3043–3044. <https://doi.org/10.1093/bioinformatics/btp498>
- Bolger, A. M., Lohse, M., & Usadel, B. (2014). TRIMMOMATIC: A flexible trimmer for Illumina sequence data. *Bioinformatics*, 30, 2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>
- Bradbury, D., Smithson, A., & Krauss, S. L. (2013). Signatures of diversifying selection at EST-SSR loci and association with climate in natural *Eucalyptus* populations. *Molecular Ecology*, 22, 5112–5129. <https://doi.org/10.1111/mec.12463>
- Broad Institute. Retrieved from <http://broadinstitute.github.io/picard/>
- Carneiro, M., Baird, S. J. E., Afonso, S., Ramirez, E., Tarroso, P., Teotónio, H., Villafuerte, R., Nachman, M. W., & Ferrand, N. (2013). Steep clines within a highly permeable genome across a hybrid zone between two subspecies of the European rabbit. *Molecular Ecology*, 22, 2511–2525. <https://doi.org/10.1111/mec.12272>
- Chan, Y. F., Marks, M. E., Jones, F. C., Villarreal, G., Shapiro, M. D., Brady, S. D., Southwick, A. M., Absher, D. M., Grimwood, J., Schmutz, J., Myers, R. M., Petrov, D., Jonsson, B., Schluter, D., Bell, M. A., & Kingsley, D. M. (2010). Adaptive evolution of pelvic reduction in sticklebacks by recurrent deletion of a *Pitx1* enhancer. *Science*, 327, 302–305. <https://doi.org/10.1126/science.1182213>
- Charlesworth, B., Coyne, J. A., & Barton, N. H. (1987). The relative rates of evolution of sex chromosomes and autosomes. *The American Naturalist*, 130, 113–146. <https://doi.org/10.1086/284701>
- Chen, J., Källman, T., Ma, X., Gyllenstrand, N., Zaina, G., Morgante, M., & Lascoux, M. (2012). Disentangling the roles of history and local selection in shaping clinal variation of allele frequencies and gene expression in Norway spruce (*Picea abies*). *Genetics*, 191, 865–881. <https://doi.org/10.1534/genetics.112.140749>
- Cruikshank, T. E., & Hahn, M. W. (2014). Reanalysis suggests that genomic islands of speciation are due to reduced diversity, not

- reduced gene flow. *Molecular Ecology*, 23, 3133–3157. <https://doi.org/10.1111/mec.12796>
- David, R., Gibert, P., Pla, E., Petavy, G., Karan, D., & Moreteau, B. (1998). Cold stress tolerance in *Drosophila*: Analysis of chill coma recovery in *D. melanogaster*. *Journal of Thermal Biology*, 23, 291–299. [https://doi.org/10.1016/S0306-4565\(98\)00020-5](https://doi.org/10.1016/S0306-4565(98)00020-5)
- de Villemereuil, P., & Gaggiotti, O. E. (2015). A new FST-based method to uncover local adaptation using environmental variables. *Methods in Ecology and Evolution*, 6, 1248–1258. <https://doi.org/10.1111/2041-210X.12418>
- Ellegren, H., Smeds, L., Burri, R., Olason, P. I., Backström, N., Kawakami, T., Künstner, A., Mäkinen, H., Nadachowska-Brzyska, K., Qvarnström, A., Uebbing, S., & Wolf, J. B. W. (2012). The genomic landscape of species divergence in *Ficedula* flycatchers. *Nature*, 491, 756–760. <https://doi.org/10.1038/nature11584>
- Endler, J. A. (1973). Gene flow and differentiation. *Science*, 179, 243–250. <https://doi.org/10.1126/science.179.4070.243>
- Endler, J. A. (1977). *Geographic variation, speciation, and clines*. Princeton University Press.
- Feder, A. F., Petrov, D. A., & Bergland, A. O. (2012). LDx: Estimation of linkage disequilibrium from high-throughput pooled resequencing data. *PLoS One*, 7, e48588. <https://doi.org/10.1371/journal.pone.0048588>
- Ferguson, L. V., Heinrichs, D. E., & Sinclair, B. J. (2016). Paradoxical acclimation responses in the thermal performance of insect immunity. *Oecologia*, 181, 77–85. <https://doi.org/10.1007/s00442-015-3529-3526>
- Findsen, A., Pedersen, T. H., Petersen, A. G., Nielsen, O. B., & Overgaard, J. (2014). Why do insects enter and recover from chill coma? Low temperature and high extracellular potassium compromise muscle function in *Locusta migratoria*. *The Journal of Experimental Biology*, 217, 1297–1306. <https://doi.org/10.1242/jeb.098442>
- Foll, M., & Gaggiotti, O. E. (2008). A genome-scan method to identify selected loci appropriate for both dominant and codominant markers: A Bayesian perspective. *Genetics*, 180, 977–993. <https://doi.org/10.1534/genetics.108.092221>
- Foll, M., Gaggiotti, O. E., Daub, J. T., Vatsiou, A., & Excoffier, L. (2014). Widespread signals of convergent adaptation to high altitude in Asia and America. *American Journal of Human Genetics*, 95, 394–407. <https://doi.org/10.1016/j.ajhg.2014.09.002>
- Garlovsky, M. D., Yusuf, L. H., Ritchie, M. G., & Snook, R. R. (2020). Within-population sperm competition intensity does not predict asymmetry in conpopulation sperm precedence. *Philosophical Transactions of the Royal Society*, 375, 20200071. <https://doi.org/10.1098/rstb.2020.0071>
- Gautier, M. (2015). Genome-wide scan for adaptive divergence and association with population-specific covariates. *Genetics*, 201, 1555–1579. <https://doi.org/10.1534/genetics.115.181453>
- Gibbs, A. G. (2002). Lipid melting points and cuticular permeability: New insights into an old problem. *Journal of Insect Physiology*, 48, 391–400. [https://doi.org/10.1016/S0022-1910\(02\)00059-8](https://doi.org/10.1016/S0022-1910(02)00059-8)
- Gibert, P., Moreteau, B., Petavy, G., Karan, D., & David, J. (2001). Chill-coma tolerance, a major climatic adaptation among *Drosophila* species. *Evolution*, 55, 1063–1068. [10.1554/0014-3820\(2001\)055\[1063:cctamc\]2.0.co;2](https://doi.org/10.1554/0014-3820(2001)055[1063:cctamc]2.0.co;2)
- Hijmans, R. J., Cameron, S. E., Parra, J. L., Jones, P. G., & Jarvis, A. (2005). Very high resolution interpolated climate surfaces for global land areas. *International Journal of Climatology*, 25, 1965–1978. <https://doi.org/10.1002/joc.1276>
- Hijmans, R. E., van Etten, J., Cheng, J., Mattiuzzi, M., Sumner, M., Greenberg, J. A., & Wueest, R. (2016). RASTER: Geographic Data Analysis and Modeling. <https://CRAN.R-project.org/package=raster>
- Hivert, V., Leblois, R., Petit, E. J., Gautier, M., & Vitalis, R. (2018). Measuring genetic differentiation from pool-seq data. *Genetics*, 210, 315–330. <https://doi.org/10.1534/genetics.118.300900>
- Hoekstra, H., Drumm, K., & Nachman, M. (2004). Ecological genetics of adaptive color polymorphism in pocket mice: Geographic variation in selected and neutral genes. *Evolution*, 58, 1329–1341. <https://doi.org/10.1111/j.0014-3820.2004.tb01711.x>
- Huang, D. W., Sherman, B. T., & Lempicki, R. A. (2009a). Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nature Protocols*, 4, 44–57. <https://doi.org/10.1038/nprot.2008.211>
- Huang, D. W., Sherman, B. T., & Lempicki, R. A. (2009b). Bioinformatics enrichment tools: paths toward the comprehensive functional analysis of large gene lists. *Nucleic Acids Research*, 37, 1–13. <https://doi.org/10.1093/nar/gkn923>
- Jennings, J. H., Mazzi, D., Ritchie, M. G., & Hoikkala, A. (2011). Sexual and postmating reproductive isolation between allopatric *Drosophila montana* populations suggest speciation potential. *BMC Evolutionary Biology*, 11, <https://doi.org/10.1186/1471-2148-11-68>
- Jennings, J. H., Snook, R. R., & Hoikkala, A. (2014). Reproductive isolation among allopatric *Drosophila montana* populations. *Evolution*, 68, 3095–3108. <https://doi.org/10.1111/evo.12535>
- Kankare, M., Parker, D. J., Merisalo, M., Salminen, T. S., & Hoikkala, A. (2016). Transcriptional differences between diapausing and non-diapausing *D. montana* females reared under the same photoperiod and temperature. *PLoS One*, 11, e0161852. <https://doi.org/10.1371/journal.pone.0161852>
- Kankare, M., Salminen, T., Laiho, A., Vesala, L., & Hoikkala, A. (2010). Changes in gene expression linked with adult reproductive diapause in a northern malt fly species: A candidate gene microarray study. *BMC Ecology*, 10(1), 3. <https://doi.org/10.1186/1472-6785-10-3>
- Kapun, M., Barrón, M. G., Staubach, F., Obbard, D. J., Wiberg, R. A. W., Vieira, J., Goubert, C., Rota-Stabelli, O., Kankare, M., Bogaerts-Márquez, M., Haudry, A., Waidele, L., Kozaretska, I., Pasyukova, E. G., Loeschcke, V., Pascual, M., Vieira, C. P., Serga, S., Montchamp-Moreau, C., ... González, J. (2020). Genomic analysis of European *Drosophila melanogaster* populations reveals longitudinal structure, continent-wide selection, and previously unknown DNA viruses. *Molecular Biology and Evolution*, 37, 2661–2678.
- Kapun, M., Fabian, D. K., Goudet, J., & Flatt, T. (2016). Genomic evidence for adaptive inversion clines in *Drosophila melanogaster*. *Molecular Biology and Evolution*, 33, 1317–1336. <https://doi.org/10.1093/molbev/msw016>
- Kauranen, H., Kinnunen, J., Hiillos, A.-L., Lankinen, P., Hopkins, D., Wiberg, R. A. W., Ritchie, M. G., & Hoikkala, A. (2019). Selection for reproduction under short photoperiods changes diapause-associated traits and induces widespread genomic divergence. *Journal of Experimental Biology*, 222, jeb205831. <https://doi.org/10.1242/jeb.205831>
- Kellermann, V., Overgaard, J., Hoffmann, A. A., Flojgaard, C., Svenning, J. C., & Loeschcke, V. (2012). Upper thermal limits of *Drosophila* are linked to species distributions and strongly constrained phylogenetically. *Proceedings of the National Academy of Sciences USA*, 109, 16228–16233. <https://doi.org/10.1073/pnas.1207553109>
- Kirkpatrick, M., & Barton, N. (2006). Chromosome inversions, local adaptation and speciation. *Genetics*, 173, 419–434. <https://doi.org/10.1534/genetics.105.047985>
- Klappert, K., Mazzi, D., Hoikkala, A., & Ritchie, M. G. (2007). Male courtship song and female preference variation between phylogeographically distinct populations of *drosophila montana*. *Evolution*, 61, 1481–1488. <https://doi.org/10.1111/j.1558-5646.2007.00125.x>
- Kofler, R., & Schlötterer, C. (2012). GOWINDA: Unbiased analysis of gene set enrichment for genome-wide association studies. *Bioinformatics*, 28, 2084–2085. <https://doi.org/10.1093/bioinformatics/bts315>
- Kolaczowski, B., Kern, A. D., Holloway, A. K., & Begun, D. J. (2011). Genomic differentiation between temperate and tropical Australian populations of *Drosophila melanogaster*. *Genetics*, 187, 245–260. <https://doi.org/10.1534/genetics.110.123059>

- Lakovaara, S. (1969). Malt as a culture medium for *Drosophila* species. *Drosophila Information Service*, 44, 128.
- Lankinen, P., Tyukmaeva, V. I., & Hoikkala, A. (2013). Northern *Drosophila montana* flies show variation both within and between cline populations in the critical day length evoking reproductive diapause. *Journal of Insect Physiology*, 59, 745–751. <https://doi.org/10.1016/j.jinsphys.2013.05.006>
- Lê, S., Josse, J., & Husson, F. (2008). FactoMineR: An R package for multivariate analysis. *Journal of Statistical Software*, 25, 1–18. <https://doi.org/10.18637/jss.v025.i01>
- Li, H. (2013). Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. arXiv:1303.3997v1 [q-bio.GN] Retrieved from: <https://arxiv.org/abs/1303.3997>
- Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., & Durbin, R., The 1000 Genome Project Data Processing Subgroup. (2009). The Sequence Alignment/Map format and SAMtools. *Bioinformatics*, 25, 2078–2079. <https://doi.org/10.1093/bioinformatics/btp352>
- Lohse, K., Clarke, M., Ritchie, M. G., & Etges, W. J. (2015). Genome-wide tests for introgression between cactophilic *Drosophila* implicate a role of inversions during speciation. *Evolution*, 69, 1178–1190. <https://doi.org/10.1111/evo.12650>
- Machado, H. E., Bergland, A. O., O'Brien, K., Behrman, E. L., Schmidt, P. S., & Petrov, D. (2015). Comparative population genomics of latitudinal variation in *Drosophila simulans* and *Drosophila melanogaster*. *Molecular Ecology*, 25, 723–740. <https://doi.org/10.1111/mec.13446>
- MacMillan, H. A., Williams, C. M., Staples, J. F., & Sinclair, B. J. (2012). Reestablishment of ion homeostasis during chill-coma recovery in the cricket *Gryllus pennsylvanicus*. *Proceedings of the National Academy of Sciences of USA*, 109, 20750–20755. <https://doi.org/10.1073/pnas.1212788109>
- Maston, G. A., Evans, S. K., & Green, M. R. (2006). Transcriptional regulatory elements in the human genome. *Annual Review of Genomics and Human Genetics*, 7, 29–59. <https://doi.org/10.1146/annurev.genom.7.080505.115623>
- McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., Garimella, K., Altshuler, D., Gabriel, S., Daly, M., & DePristo, M. A. (2010). The Genome Analysis Toolkit: A MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Research*, 20, 1297–1303. <https://doi.org/10.1101/gr.107524.110>
- Mirol, P. M., Schäfer, M. A., Orsing, L., Routtu, J., Schlötterer, C., Hoikkala, A., & Butlin, R. K. (2007). Phylogeographic patterns in *Drosophila montana*. *Molecular Ecology*, 16, 1085–1097. <https://doi.org/10.1111/j.1365-294X.2006.03215.x>
- Morales-Hojas, R., Päällysaho, S., Vieira, C. P., Hoikkala, A., & Vieira, J. (2007). Comparative polytene chromosome maps of *D. montana* and *D. virilis*. *Chromosoma*, 116, 21–27. <https://doi.org/10.1007/s00412-006-0075-3>
- Mylonakis, E., Podsiadlowski, L., Muhammed, M., & Vilcinskis, A. (2016). Diversity, evolution and medical applications of insect antimicrobial peptides. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 371, 20150290. <https://doi.org/10.1098/rstb.2015.0290>
- Noor, M. A. F., & Bennett, S. M. (2009). Islands of speciation or mirages in the desert? Examining the role of restricted recombination in maintaining species. *Heredity*, 103, 439–444. <https://doi.org/10.1038/hdy.2009.151>
- Paolucci, S., Salis, L., Vermeulen, C. J., Beukeboom, L. W., & van de Zande, L. (2016). QTL analysis of the photoperiodic response and clinal distribution of period alleles in *Nasonia vitripennis*. *Molecular Ecology*, 25, 4805–4817. <https://doi.org/10.1111/mec.13802>
- Parker, D. J., Envall, T., Ritchie, M. G., & Kankare, M. (2021). Sex-specific responses to cold in a very cold-tolerant, northern *Drosophila* species. *Heredity*, 126(4), 695–705. <https://doi.org/10.1038/s41437-020-00398-2>
- Parker, D. J., Ritchie, M. G., & Kankare, M. (2016). Preparing for Winter: The transcriptomic response associated with different day lengths in *Drosophila montana*. G3: *Genes, Genomes, Genetics*, 6, 1373–1381. <https://doi.org/10.1534/g3.116.027870>
- Parker, D. J., Vesala, L., Ritchie, M. G., Laiho, A., Hoikkala, A., & Kankare, M. (2015). How consistent are the transcriptome changes associated with cold acclimation in two species of the *Drosophila virilis* group? *Heredity*, 115, 13–21. <https://doi.org/10.1038/hdy.2015.6>
- Parker, D. J., Wiberg, R. A. W., Trivedi, U., Tyukmaeva, V. I., Gharbi, K., Butlin, R. K., Hoikkala, A., Kankare, M., & Ritchie, M. G. (2018). Inter- and intra-specific genomic divergence in *Drosophila montana* shows evidence for cold adaptation. *Genome Biology and Evolution*, 10, 2086–2101. <https://doi.org/10.1093/gbe/evy147>
- Patterson, J. T., & Stone, W. S. (1952). *Evolution in the Genus Drosophila*. Macmillan.
- Pennacchio, L. A., Bickmore, W., Dean, A., Nobrega, M. A., & Bejerano, G. (2013). Enhancers: five essential questions. *Nature Reviews Genetics*, 14, 288–295. <https://doi.org/10.1038/nrg3458>
- Plummer, M., Best, N., Cowles, K., & Vines, K. (2006). CODA: Convergence diagnosis and output analysis for MCMC. *R News*, 6, 7–11.
- R Development Core Team. (2020). *R: A language and environment for statistical computing*. Retrieved from <http://www.r-project.org>
- Ravinet, M., Faria, R., Butlin, R. K., Galindo, J., Bierne, N., Rafajlović, M., Noor, M. A. F., Mehlig, B., & Westram, A. M. (2017). Interpreting the genomic landscape of speciation: A road map for finding barriers to gene flow. *Journal of Evolutionary Biology*, 30, 1450–1477. <https://doi.org/10.1111/jeb.13047>
- Routtu, J., Mazzi, D., van der Linde, K., Mirol, P., Butlin, R., & Ritchie, M. G. (2007). The extent of variation in male song, wing and genital characters among allopatric *Drosophila montana* populations. *Journal of Evolutionary Biology*, 20, 1591–1601. <https://doi.org/10.1111/j.1420-9101.2007.01323.x>
- Scheffzek, K., & Welti, S. (2012). Pleckstrin homology (PH) like domains – versatile modules in protein–protein interaction platforms. *FEBS Letters*, 586, 2662–2673. <https://doi.org/10.1016/j.febslet.2012.06.006>
- Schlötterer, C., Tobler, R., Kofler, R., & Nolte, V. (2014). Sequencing pools of individuals – mining genome-wide polymorphism data without big funding. *Nature Reviews Genetics*, 15, 749–763. <https://doi.org/10.1038/nrg3803>
- Sinclair, B. J., Ferguson, L. V., Salehipour-shirazi, G., & MacMillan, H. A. (2013). Cross-tolerance and cross-talk in the cold: Relating low temperatures to desiccation and immune stress in insects. *Integrative and Comparative Biology*, 53, 545–556. <https://doi.org/10.1093/icb/ict004>
- Sinclair, B. J., Nelson, S., Nilson, T. L., Roberts, S. P., & Gibbs, A. G. (2007). The effect of selection for desiccation resistance on cold tolerance of *Drosophila melanogaster*. *Physiological Entomology*, 32, 322–327. <https://doi.org/10.1111/j.1365-3032.2007.00585.x>
- Sinensky, M. (1974). Homeoviscous adaptation—A homeostatic process that regulates the viscosity of membrane lipids in *Escherichia coli*. *Proceedings of the National Academy of Sciences of USA*, 71, 522–525. <https://doi.org/10.1073/pnas.71.2.522>
- Stanziano, J. R., Sové, R. J., Rundle, H. D., & Sinclair, B. J. (2015). Rapid desiccation hardening changes the cuticular hydrocarbon profile of *Drosophila melanogaster*. *Comparative Biochemistry and Physiology*, 180, 38–42. <https://doi.org/10.1016/j.cbpa.2014>
- Stone, W. S., Guest, W. C., & Wilson, F. D. (1960). The evolutionary implications of the cytological polymorphism and phylogeny of the *virilis* group of *Drosophila*. *Proceedings of the National Academy of Sciences of USA*, 46, 350–361. <https://doi.org/10.1073/pnas.46.3.350>
- Takahashi, Y. (2015). Mechanisms and tests for geographic clines in genetic polymorphisms. *Population Ecology*, 57, 355–362. <https://doi.org/10.1007/s10144-014-0474-x>
- Throckmorton, L. H. (1969). *The virilis Species Group. The Genetics and Biology of Drosophila*, Vol. 3 (pp. 227–296).

- Throckmorton, L. H. (1982). The virilis species group. In M. A. Ashburner, H. L. Carson, & J. N. Thompson (Eds.), *The genetics and biology of Drosophila* (Vol. 3b, pp. 227–297). London: Academic Press.
- Thurmond, J., Goodman, J. L., Strelets, V. B., Attrill, H., Gramates, L. S., Marygold, S. J., Matthews, B. B., Millburn, G., Antonazzo, G., Trovisco, V., Kaufman, T. C., Calvi, B. R., Perrimon, N., Gelbart, S. R., Agapite, J., Broll, K., Crosby, L., Santos, G., Emmert, D., ... The FlyBase Consortium. (2019). FlyBase 2.0: The next generation. *Nucleic Acids Research*, 47, 759–765. <https://doi.org/10.1093/nar/gky1003>
- Tyukmaeva, V. I., Lankinen, P., Kinnunen, J., Kauranen, H., & Hoikkala, A. (2020). Latitudinal clines in the timing and temperature-sensitivity of photoperiodic reproductive diapause in *Drosophila montana*. *Ecography*, 43, 1–10. <https://doi.org/10.1111/ecog.04892>
- Tyukmaeva, V. I., Salminen, T. S., Kankare, M., Knott, E., & Hoikkala, A. (2011). Adaptation to a seasonally varying environment: A strong latitudinal cline in reproductive diapause combined with high gene flow in *Drosophila montana*. *Ecology and Evolution*, 1, 160–168. <https://doi.org/10.1002/ece3.14>
- Vesala, L., & Hoikkala, A. (2011). Effects of photoperiodically induced reproductive diapause and cold hardening on the cold tolerance of *Drosophila montana*. *Journal of Insect Physiology*, 57, 46–51. <https://doi.org/10.1016/j.jinsphys.2010.09.007>
- Vesala, L., Salminen, T. S., Kostal, V., Zahradnickova, H., & Hoikkala, A. (2012). Myo-inositol as a main metabolite in overwintering flies: Seasonal metabolomic profiles and cold stress tolerance in a northern drosophilid fly. *Journal of Experimental Biology*, 215, 2891–2897. <https://doi.org/10.1242/jeb.069948>
- Vesala, L., Salminen, T. S., Laiho, A., Hoikkala, A., & Kankare, M. (2012). Cold tolerance and cold-induced modulation of gene expression in two *Drosophila virilis* group species with different distributions. *Insect Molecular Biology*, 21, 107–118. <https://doi.org/10.1111/j.1365-2583.2011.01119.x>
- Vines, T. H., Dalziel, A. C., Albert, A. Y. K., Veen, T., Schulte, P. M., & Schluter, D. (2016). Cline coupling and uncoupling in a stickleback hybrid zone. *Evolution*, 70, 1023–1038. <https://doi.org/10.1111/evo.12917>
- Werner, T., Koshikawa, S., Williams, T. M., & Carroll, S. B. (2010). Generation of a novel wing colour pattern by the Wingless morphogen. *Nature*, 464, 1143–1148. <https://doi.org/10.1038/nature08896>
- Wilder, S. M., Mayntz, D., Toft, S., Rypstra, A. L., Pilati, A., & Vanni, M. J. (2010). Intraspecific variation in prey quality: A comparison of nutrient presence in prey and nutrient extraction by predators. *Oikos*, 119, 350–358. <https://doi.org/10.1111/j.1600-0706.2009.17819.x>
- Williams, C. M., McCue, M. D., Sunny, N. E., Szejner-Sigal, A., Morgan, T. J., Allison, D. B., & Hahn, D. A. (2016). Cold adaptation increases rates of nutrient flow and metabolic plasticity during cold exposure in *Drosophila melanogaster*. *Proceedings of the Royal Society B: Biological Sciences*, 283, 20161317. <https://doi.org/10.1098/rspb.2016.1317>
- Wolf, J. B. W., & Ellegren, H. (2016). Making sense of genomic islands of differentiation in light of speciation. *Nature Reviews Genetics*, 18, 87–100. <https://doi.org/10.1038/nrg.2016.133>
- Wood, S. N. (2004). Stable and efficient multiple smoothing parameter estimation for generalized additive models. *Journal of the American Statistical Association*, 99, 673–686. <https://doi.org/10.1198/016214504000000980>

## SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

**How to cite this article:** Wiberg, R. A. W., Tyukmaeva V., Hoikkala A., Ritchie M. G., & Kankare M. (2021). Cold adaptation drives population genomic divergence in the ecological specialist, *Drosophila montana*. *Molecular Ecology*, 30, 3783–3796. <https://doi.org/10.1111/mec.16003>