

This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Plutenko, Maksym O.; Haukka, Matti; Husak, Alina O.; Golenya, Irina A.; Mulloev, Nurullo U.

Title: A second solvatomorph of poly[[μ 4-N,N'-(1,3,5-oxadiazinane-3,5-diyl)bis(carbamoylmethanoato)]nickel(II)dipotassium] : crystal structure, Hirshfeld surface analysis and semi-empirical geometry optimization

Year: 2021

Version: Published version

Copyright: © Authors, 2021

Rights: CC BY 4.0

Rights url: <https://creativecommons.org/licenses/by/4.0/>

Please cite the original version:

Plutenko, M. O., Haukka, M., Husak, A. O., Golenya, I. A., & Mulloev, N. U. (2021). A second solvatomorph of poly[[μ 4-N,N'-(1,3,5-oxadiazinane-3,5-diyl)bis(carbamoylmethanoato)]nickel(II)dipotassium] : crystal structure, Hirshfeld surface analysis and semi-empirical geometry optimization. *Acta Crystallographica Section E : Crystallographic Communications*, 77(12), Article E77.
<https://doi.org/10.1107/s2056989021011774>

A second solvatomorph of poly[[μ_4 - N,N' -(1,3,5-oxadiazinane-3,5-diyl)bis(carbamoylmethanoato)]-nickel(II)dipotassium]: crystal structure, Hirshfeld surface analysis and semi-empirical geometry optimization

Received 13 October 2021
Accepted 5 November 2021

Edited by B. Therrien, University of Neuchâtel, Switzerland

Keywords: nickel(II) complex; template reaction; pseudomacrocyclic ligand; hydrazide-based ligand; SHAPE analysis; Hirshfeld surface analysis; semi-empirical geometry optimization; crystal structure.

CCDC reference: 2120378

Supporting information: this article has supporting information at journals.iucr.org/e

Maksym O. Plutenko,^{a*} Matti Haukka,^b Alina O. Husak,^{a,c} Irina A. Golenya^a and Nurullo U. Mulloev^d

^aDepartment of Chemistry, National Taras Shevchenko University, Volodymyrska, Street 64, 01601 Kyiv, Ukraine,


^bDepartment of Chemistry, University of Jyväskylä, P.O. Box 35, FI-40014 Jyväskylä, Finland, ^cPBMR Labs Ukraine, Murmanska 1, 02094 Kiev, Ukraine, and ^dThe Faculty of Physics, Tajik National University, Rudaki Avenue 17, 734025 Dushanbe, Tajikistan. *Correspondence e-mail: plutenkom@gmail.com

The title compound, poly[triaquabis[μ_4 - N,N' -(1,3,5-oxadiazinane-3,5-diyl)bis(carbamoylmethanoato)]dinickel(II)tetrapotassium], $[\text{K}_4\text{Ni}_2(\text{C}_7\text{H}_6\text{N}_4\text{O}_7)_2 \cdot (\text{H}_2\text{O})_3]_n$, is a second solvatomorph of poly[[μ_4 - N,N' -(1,3,5-oxadiazinane-3,5-diyl)bis(carbamoylmethanoato)]nickel(II)dipotassium] reported previously [Plutenko *et al.* (2021). *Acta Cryst.* E77, 298–304]. The asymmetric unit of the title compound includes two structurally independent complex anions $[\text{Ni}(\text{C}_7\text{H}_6\text{N}_4\text{O}_7)]^{2-}$, which exhibit an L-shaped geometry and consist of two almost flat fragments perpendicular to one another: the 1,3,5-oxadiazinane fragment and the fragment including other atoms of the anion. The central Ni atom is in a square-planar N_2O_2 coordination arrangement formed by two amide N and two carboxylate O atoms. In the crystal, the title compound forms a layered structure in which layers of negatively charged complex anions and positively charged potassium cations are stacked along the *a*-axis direction. The polymeric framework is stabilized by a system of hydrogen-bonding interactions in which the water molecules act as donors and the carboxylic, amide and water O atoms act as acceptors.


1. Chemical context

In 1976, the products of the metal-templated reaction of hydrazide and aldehyde were separated and structurally described (Clark *et al.*, 1976). It was further shown that such a synthetic strategy makes it possible to obtain complexes with 3d metals in high oxidation states. In particular, there are several works devoted to copper(III) complexes obtained by this method (Oliver & Waters, 1982; Fritsky *et al.*, 1998, 2006). Moreover, the preparation of an unprecedentedly stable iron(IV) clathrochelate complex was reported (Tomyň *et al.*, 2017). Some such compounds are promising redox catalysts, as has been shown by Pap *et al.* (2011) and Shylin *et al.* (2019). Thus, the study of the conditions and peculiarities of hydrazide-aldehyde template interactions, as well as the isolation and characterization of their products, is an important task in modern coordination chemistry.

This work is a continuation of our investigation of the interaction of oxalohydrazidehydroxamic acid with formaldehyde and nickel(II) salts. Here we report the crystal structure of the title compound poly[triaquabis[μ_4 - N,N' -(1,3,5-



oxadiazinane-3,5-diyl)bis(carbamoylmethanoato)]dinickel(II)tetrapotassium] $[(2K_2[Ni(L-2H)] \cdot 3H_2O)_n, \mathbf{2}]$, which is the solvatomorph of the earlier published (Plutenko *et al.*, 2021) complex poly[pentaaquabis[μ_n - N,N' -(1,3,5-oxadiazinane-3,5-diyl)bis(carbamoylmethanoato)]nickel(II)tetrapotassium], $[(2K_2[Ni(L-2H)] \cdot 4.8H_2O)_n, \mathbf{1}]$, $H_2L = N,N'$ -(1,3,5-oxadiazinane-3,5-diyl)bis(aminooxoacetic acid)]. Both compounds can be obtained in a similar fashion as the result of a one-pot template reaction (see Fig. 1).



2. Structural commentary

The title compound, $\mathbf{2}$, $(2K_2[Ni(L-2H)] \cdot 3H_2O)_n$, crystallizes in space group $P2_1/c$, while the previously reported compound $\mathbf{1}$, $(2K_2[Ni(L-2H)] \cdot 4.8H_2O)_n$, crystallizes in $Pbca$. Similarly to $\mathbf{1}$, the asymmetric unit of $\mathbf{2}$ (Fig. 2) includes two structurally independent complex anions $[Ni(L-2H)]^{2-}$ (namely *A* and *B*, which contain Ni1 and Ni1*B*, respectively). In addition, the unit cell of $\mathbf{2}$ also contains four potassium cations and three solvent water molecules.




Figure 1
A plausible mechanism for the formation of the $[Ni(L-2H)]^{2-}$ complex anion.




Figure 2
The asymmetric unit of $\mathbf{2}$ with displacement ellipsoids shown at the 50% probability level.

Similarly to $\mathbf{1}$, the complex anion $[Ni(L-2H)]^{2-}$ has an L-shaped geometry and consists of two almost flat fragments perpendicular to one another: the 1,3,5-oxadiazinane fragment and the fragment including other atoms of the anion. The dihedral angles between the mean planes formed by the non-hydrogen atoms of these fragments are $95.06(8)$ and $94.06(8)^\circ$ for Ni1 and Ni1*B*, respectively. The ligand molecule is coordinated in a tetradentate $\{O_{\text{carboxyl}}, N_{\text{amide}}, N_{\text{amide}}, O_{\text{carboxyl}}\}$ -mode. The central atom of the complex anion exhibits a square-planar coordination arrangement with the N_2O_2 chromophore. The deviation of the Ni^{II} atom from the mean plane defined by the donor atoms is $0.0073(13)$ and $0.0330(12)$ Å for Ni1 and Ni1*B*, respectively.

The Ni—N bond distances are in the range $1.836(3)$ – $1.849(3)$ Å and Ni—O bond lengths are $1.877(2)$ – $1.897(2)$ Å, which is typical for square-planar nickel complexes with similar ligands (Fritsky *et al.*, 1998) and close to the Ni—N and Ni—O bond distances of $\mathbf{1}$. The O—M—O', O—M—N and N—M—N' bond angles have typical values for a square-planar arrangement. The bite angles O1—Ni1—N4, N1—Ni1—O2 and N1—Ni1—N4 deviate from 90° , which is the result of the formation of the five-membered chelate rings. The N—N', N—C and C—O bond lengths of the ligand have typical values for coordinated deprotonated hydrazide and carboxyl groups.

3. Supramolecular features

In the crystal, the nickel(II) complex anions $[Ni(L-2H)]^{2-}$ form layers parallel to the *bc* plane (Fig. 3*a*). Neighbouring complex anion layers are sandwiched by layers of potassium counter-cations (Fig. 4). Thus, negatively charged complex anion layers and positively charged potassium cationic layers are stacked along the *a*-axis direction. It is useful to note that a similar layered structure motif was observed in the crystal of the previously published compound $\mathbf{1}$. However, in the crystal of $\mathbf{1}$ the NiN_2O_2 plane is almost perpendicular to the complex anion layer plane (Fig. 3*b*): the angle between NiN_2O_2 and the *ab* plane is $84.43(4)$ and $85.03(5)^\circ$ for Ni1 and Ni1*B*,




Figure 3
Layers formed by the nickel(II) complex anions $[\text{Ni}(\text{L}-2\text{H})]^{2-}$ in the crystals of (a) compound **2** and (b) compound **1**.

respectively. In contrast, in the crystal of **2** the angle between NiN_2O_2 and the bc plane is $78.30(8)$ and $86.29(7)^\circ$ for Ni1 and Ni1B, respectively.

The demarcation of bonded and non-bonded $\text{K}-X$ interactions ($X = \text{N}$ or O) is still an unclear and debatable problem (Alvarez, 2013). Therefore, the criteria of such demarcation used in this paper need to be detailed. Based on the aforementioned publication (Alvarez, 2013), we propose 3.7 \AA as the maximal distance for $\text{K}-\text{N}$ bonds. Recently, it was shown (Gagné & Hawthorne, 2016) that $\text{K}-\text{O}$ main and maximal bond distances depend on the coordination number of K. The results of this work permits 3.4 , 3.5 and 3.6 \AA to be proposed as the maximal distances for $\text{K}-\text{O}$ bonds in the case of potassium coordination numbers 7, 8 and 9, respectively. In addition, $\text{K} \cdots \text{N}_{\text{amide}}$ interactions were determined as non-bonding because the existence of such bonds would lead to the presence of unstable three-membered $\text{KN}_{\text{amide}}\text{N}_{\text{oxadiazinane}}$ rings with extremely small $\text{N}-\text{K}-\text{N}'$ angles.

The potassium cations are bound to the nickel(II) complex anions through the carboxylic O atoms (K4) the carboxylic and the amide O atoms (K1, K2) or through the amide O and the oxadiazinane N atoms (K3). In addition, the potassium




Figure 4
Crystal packing of the title compound in a stick model, showing the coordination polyhedra of the potassium cations. H atoms are omitted for clarity.

Table 1

Values for continuous shapes measures (CSHM) of the polyhedra centred by the potassium cations.

Shape	CSHM	
	K1	K2
Heptagon ($D7h$)	28.515	29.484
Hexagonal pyramid ($C6v$)	17.225	20.349
Pentagonal bipyramid ($D5h$)	5.142	3.122
Capped octahedron ($C3v$)	7.539	7.840
Capped trigonal prism ($C2v$)	6.374	5.639
Johnson pentagonal bipyramid J13 ($D5h$)	8.789	6.943
Johnson elongated triangular pyramid J7 ($C3v$)	16.352	20.453
	K3	
Enneagon ($D9h$)	32.593	
Octagonal pyramid ($C8v$)	23.087	
Heptagonal bipyramid ($D7h$)	14.962	
Johnson triangular cupola J3 ($C3v$)	12.759	
Capped cube J8 ($C4v$)	9.046	
Spherical-relaxed capped cube ($C4v$)	7.600	
Capped square antiprism J10 ($C4v$)	6.360	
Spherical capped square antiprism ($C4v$)	5.020	
Tricapped trigonal prism J51 ($D3h$)	6.694	
Spherical tricapped trigonal prism ($D3h$)	5.698	
Tridiminished icosahedron J63 ($C3v$)	11.379	
Hula-hoop ($C2v$)	6.577	
Muffin (Cs)	3.691	
	K4	
Octagon ($D8h$)	33.086	
Heptagonal pyramid ($C7v$)	18.988	
Hexagonal bipyramid ($D6h$)	14.426	
Cube (Oh)	10.884	
Square antiprism ($D4d$)	5.463	
Triangular dodecahedron ($D2d$)	5.187	
Johnson gyrobifastigium J26 ($D2d$)	11.775	
Johnson elongated triangular bipyramid J14 ($D3h$)	26.080	
Biaugmented trigonal prism J50 ($C2v$)	6.413	
Biaugmented trigonal prism ($C2v$)	6.587	
Snub diphenoid J84 ($D2d$)	7.862	
Triakis tetrahedron (Td)	11.175	
Elongated trigonal bipyramid ($D3h$)	20.295	

cations have contacts with the O atoms of water molecules, with the amide and the carboxylic O atoms, and with the oxadiazinane O and N atoms of neighbouring complex anions. The K1 and K2 cations exhibit an O_6N coordination, while the K3 and K4 cations exhibit O_8N and O_7N coordinations, respectively.

For an evaluation of the coordination geometry of each potassium cation, *SHAPE 2.1* software (Llunell *et al.*, 2013) was used. A *SHAPE* analysis of the potassium coordination sphere (Table 1, Fig. 5) yields the lowest continuous shape measure (CSHM) value for a distorted pentagonal bipyramid (5.142 for K1 and 3.122 for K2), a distorted muffin (3.691 for K3) and a distorted triangular dodecahedron (5.187 for K4). For K4, comparable CSHM values were obtained for a square antiprism (5.463).

The polyhedra around the neighbouring potassium cations are connected with each other through common vertices (K1 with K3, K1 with K4, K2 with K4), edges (K3 with K4) and faces (K1 with K2, K1 with K3, K2 with K3). The $\text{K}-\text{O}$ bond lengths are in the range $2.628(2)$ – $3.271(3) \text{ \AA}$, $\text{K}-\text{N}$ $2.887(3)$ – $3.025(3) \text{ \AA}$, which is close to those reported for the structures of related carboxylate and amide complexes (Fritsky *et al.*, 1998; Mokhir *et al.*, 2002).




Figure 5
Polyhedral views of the coordination environments for the potassium cations.

The polymeric framework of **2** is stabilized by an extensive system of hydrogen-bonding interactions in which the water molecules act as donors and the carboxylic, the amide and the water O atoms act as acceptors (Table 2). Similarly to **1**, the hydrogen bonds are localized mainly at the potassium cation layers (Fig. 6). Moreover, in comparison to **1**, the unit cell of **2** contains a smaller number of water molecules, which causes a smaller number of hydrogen-bond interactions in the crystal structure.

4. Hirshfeld analysis

The Hirshfeld surface analysis (Spackman & Jayatilaka, 2009) and the associated two-dimensional fingerprint plots (McKinnon *et al.*, 2007) were performed with *CrystalExplorer17* (Turner *et al.*, 2017). The Hirshfeld surfaces of the




Figure 6
Crystal packing of the title compound. C—H hydrogen atoms are omitted for clarity. Hydrogen bonds are indicated by dashed lines.

Table 2
Hydrogen-bond geometry (Å, °).

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
O8—H8O...O9 ⁱ	0.85	2.02	2.869 (4)	173
O8—H8P...O4B ⁱⁱ	0.85	2.01	2.858 (3)	166
O9—H9P...O4 ⁱⁱⁱ	0.86	1.91	2.722 (3)	157
O9—H9O...O6B ^{iv}	0.86	2.07	2.864 (3)	153
O10—H10P...O4 ^v	0.88	2.02	2.887 (3)	168
O10—H10O...O7B ^{vi}	0.87	2.04	2.882 (3)	164

Symmetry codes: (i) $x, -y + \frac{1}{2}, z + \frac{1}{2}$; (ii) $-x, y + \frac{1}{2}, -z + \frac{1}{2}$; (iii) $-x + 1, -y + 1, -z$; (iv) $x, -y + \frac{1}{2}, z - \frac{1}{2}$; (v) $-x + 1, y + \frac{1}{2}, -z + \frac{1}{2}$; (vi) $x, y + 1, z$.

complex anions are colour-mapped with the normalized contact distance (d_{norm}) from red (distances shorter than the sum of the van der Waals radii) through white to blue (distances longer than the sum of the van der Waals radii).

The Hirshfeld surface of the title compound is mapped over d_{norm} , in the colour ranges -0.6388 to 0.9164 a.u. and -0.6768 to 0.7286 a.u. for Ni1 and Ni1B complex anions, respectively (Fig. 7). Similarly to **1**, the complex anions of **2** are connected to the other elements of the crystal packing mainly *via* the amide and the carboxylic O atoms. However, in contrast to **1**, one of the oxadiazinane O atoms of **2** is also involved in intermolecular bond formation.

A fingerprint plot delineated into specific interatomic contacts contains information related to specific intermolecular interactions. The blue colour refers to the frequency of occurrence of the (d_i, d_e) pair with the full fingerprint plot outlined in gray. Fig. 8a and 9a show the two-dimensional fingerprint plots of the sum of the contacts contributing to the Hirshfeld surface represented in normal mode for the Ni1 and Ni1B complex anions, respectively.

The most significant contribution to the Hirshfeld surface is from O...H/H...O contacts (36.9% and 38.7% for the Ni1 and Ni1B complex anions, respectively; Fig. 8b and 9b). In addition, O...K/K...O (20.9% and 18.2% for the Ni1 and Ni1B complex anions; Fig. 8c and 9c) and H...H (10.4% and 13.1% for the Ni1 and Ni1B complex anions, respectively;




Figure 7
The Hirshfeld surfaces of the Ni1 (A) and Ni1B (B) complex anions mapped over d_{norm} .




Figure 8
 (a) Full two-dimensional fingerprint plot of the NiI complex anion and those delineated into (b) O...H/H...O (36.9%) (c) O...K/K...O (20.9%) and (d) H...H (10.4%) contacts.

Fig. 8d and 9d) make very significant contributions to the total Hirshfeld surface. This indicates that there are more K...O contacts and fewer O...H contacts compared to the crystal of **1**.




Figure 9
 (a) Full two-dimensional fingerprint plot of the NiIB complex anion and those delineated into (b) O...H/H...O (38.7%) (c) O...K/K...O (18.2%) and (d) H...H (13.1%) contacts.

Table 3

Comparison of selected geometric data (\AA , $^\circ$; mean values) for the NiI complex anion from calculated and X-ray data.

Geometric parameter	X-ray	PM7	DFTB	GFN2-xTB
Oxadiazinane ring				
C—O	1.434	1.413	1.467	1.410
C—N	1.463	1.489	1.463	1.452
Carboxylate moiety				
C—O	1.287	1.276	1.451	1.260
C=O	1.233	1.224	1.196	1.208
Hydrazide moiety				
C—O	1.249	1.232	1.227	1.216
C—N	1.321	1.357	1.393	1.332
N—N	1.432	1.413	1.413	1.415
C—N _{amide} —Ni—N _{oxadiazine}	175.74	133.89	169.00	162.81
Ni coordination arrangement				
Ni—O	1.892	1.776	1.780	1.871
Ni—N	1.840	1.955	1.974	1.871
O—Ni—N chelate	85.24	93.35	81.32	82.94
O—Ni—N non-chelate	178.29	173.19	162.52	176.77
N—Ni—N	85.53	88.09	90.73	94.40

5. Geometry optimization

The searching of computationally ‘cheap’ but still sufficiently accurate methods of transition-metal complex geometry optimization is an important task of modern computational chemistry. The geometry optimization calculations were carried out with three semi-empirical methods: PM7, DFTB and GFN2-xTB. The PM7 (Stewart, 2013) calculations were performed with *MOPAC2016* software (Stewart, 2016). The DFTB calculations were carried out with the *DFTB+* software package (Hourahine *et al.*, 2020) using the ‘mio-1-1’ (Elstner *et al.*, 1998) and the ‘trans3d-0-1’ (Zheng *et al.*, 2007) Slater–Koster parameterization sets. The GFN2-xTB (Bannwarth *et al.*, 2019) calculations were applied with *xtb 6.4* package (Grimme, 2019). The geometry of the NiI complex anion obtained from the crystal structure was used as the starting geometry for the calculations.

In general, for all described semi-empirical methods, the calculated geometric parameters of the oxadiazinane ring are in reasonable agreement with experimental values (see Table 3). On the other hand, the accuracy of the non-oxadiazinane fragment geometry prediction varies greatly depending on the method. The worst agreement with experiments is from the PM7 method, mainly because of the pyramidalization of the amide nitrogen atom (Table 3). Such non-planarity of the amide fragment is a well-known problem of the PMx methods (Feigel & Strassner, 1993). In contrast, the DFTB method predicts the amide geometric parameters with high accuracy but demonstrates longer than experimental carboxylate C—O bonds and a slight tetragonal distortion of the nickel(II) coordination polyhedra (Table 3). The best results were obtained with the GFN2-xTB method for which the calculated geometric parameters correlate nicely with experimental values (Table 3). The maximal difference between the calculated and the experimental bond lengths concerns the C—O lengths (shorter than the experimental values within 0.024–0.033 Å). A superimposed analysis of the NiI complex anion with its optimized structure gives an




Figure 10
Structural overlay between the experimental (blue) and optimized (orange) structures.

RMSD of 0.131 Å (Fig. 10). Thus, the GFN2-xTB method is a promising geometry prediction method for transition-metal complexes based on hydrazide and carboxylate ligands.

6. Database survey

A search in the Cambridge Structural Database (CSD version 5.39, update of May 2018; Groom *et al.*, 2016) resulted in 11 hits dealing with 3d-metal complexes with macrocyclic or pseudo-macrocyclic ligands formed by template binding of several hydrazide groups by formaldehyde molecules. These complexes contain the following 3d metals: Ni^{II} (Fritsky *et al.*, 1998), Cu^{II} (Clark *et al.*, 1976; Fritsky *et al.*, 2006), Cu^{III} (Oliver & Waters, 1982; Fritsky *et al.*, 1998, Fritsky *et al.*, 2006) and Fe^{IV} (Tomyn *et al.*, 2017). Thus, such macrocyclic and pseudo-macrocyclic ligand systems exhibit a tendency to stabilize the high oxidation states of 3d metals.

7. Synthesis and crystallization

A solution of Ni(ClO₄)₂·6H₂O (0.091 g, 0.25 mmol) in 5 ml of water was added to a warm solution of oxalohydrazide-hydroxamic acid (0.06 g, 0.5 mmol) in 5 ml of water. The resulting light-green mixture was stirred with heating (320–330 K) for 20 min and then 1 ml of 4M KOH solution was added. As a result, the colour of the solution changed to pink. After 5 min of stirring, 0.03 g of the paraformaldehyde (1 mmol) was added and stirring with heating (323–333 K) was continued for 30 min. The resulting orange solution was left for crystallization by slow evaporation in air. After one week, orange crystals of **2** suitable for X-ray diffraction studies were obtained. The crystals were filtered off, washed with diethyl ether and dried in the air. Yield 0.044 g (42%). Elemental analysis for C₁₄H₁₈N₈O₁₇K₄Ni₂ (mol. mass 844.12), calculated, %: C 19.92; H 2.15; N 13.27; Found, %: C 19.69; H 2.16; N 13.11. UV–vis (H₂O), λ_{max} (ε, mol⁻¹ dm³ cm⁻¹): 520 nm (1380). IR (KBr, cm⁻¹): 3420 *br* ν(O–H) stretch, 2981, 2910, 2860 ν(C–H) stretch, 1643 (*vs*) ν(C=O) amide I, 1590 ν_{as}(COO⁻), 1435 ν_s(COO⁻).

Table 4
Experimental details.

Crystal data	
Chemical formula	[K ₄ Ni ₂ (C ₇ H ₆ N ₄ O ₇) ₂ (H ₂ O) ₃]
<i>M</i> _r	844.18
Crystal system, space group	Monoclinic, <i>P</i> ₂ ₁ / <i>c</i>
Temperature (K)	100
<i>a</i> , <i>b</i> , <i>c</i> (Å)	20.3825 (5), 7.7039 (3), 17.3078 (6)
β (°)	98.240 (2)
<i>V</i> (Å ³)	2689.69 (16)
<i>Z</i>	4
Radiation type	Mo <i>K</i> α
μ (mm ⁻¹)	2.12
Crystal size (mm)	0.15 × 0.09 × 0.08
Data collection	
Diffractometer	Bruker Kappa APEXII CCD
Absorption correction	Multi-scan (<i>SADABS</i> ; Sheldrick, 2008)
<i>T</i> _{min} , <i>T</i> _{max}	0.746, 0.842
No. of measured, independent and observed [<i>I</i> > 2σ(<i>I</i>)] reflections	25068, 6148, 5118
<i>R</i> _{int}	0.043
(sin θ/λ) _{max} (Å ⁻¹)	0.650
Refinement	
<i>R</i> [<i>F</i> ² > 2σ(<i>F</i> ²)], <i>wR</i> (<i>F</i> ²), <i>S</i>	0.041, 0.082, 1.14
No. of reflections	6148
No. of parameters	406
H-atom treatment	H-atom parameters constrained
Δρ _{max} , Δρ _{min} (e Å ⁻³)	0.63, -0.45

Computer programs: *COLLECT* (Bruker, 2008), *DENZO/SCALEPACK* (Otwinowski & Minor, 1997), *SUPERFLIP* (Palatinus & Chapuis, 2007), *SHELXL2018/1* (Sheldrick, 2015), *DIAMOND* (Brandenburg, 2009) and *SHELXL97* (Sheldrick, 2008).

8. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 4. H atoms were positioned geometrically (O–H = 0.85–0.88, C–H = 0.99 Å) and refined as riding with *U*_{iso}(H) = 1.2 *U*_{eq}(O, C).

Funding information

This work was supported by the Ministry of Education and Science of Ukraine: Grant of the Ministry of Education and Science of Ukraine for perspective development of a scientific direction 'Mathematical sciences and natural sciences' at Taras Shevchenko National University of Kyiv.

References

- Alvarez, S. (2013). *Dalton Trans.* **42**, 8617–8636.
 Bannwarth, C., Ehlert, S. & Grimme, S. (2019). *J. Chem. Theory Comput.* **15**, 1652–1671.
 Brandenburg, K. (2009). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
 Bruker (2008). *COLLECT*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Clark, G. R., Skelton, B. W. & Waters, T. N. (1976). *J. Chem. Soc. Dalton Trans.* pp. 1528–1536.
 Elstner, M., Porezag, D., Jungnickel, G., Elsner, J., Haugk, M., Frauenheim, Th., Suhai, S. & Seifert, G. (1998). *Phys. Rev. B*, **58**, 7260–7268.
 Feigel, M. & Strassner, T. (1993). *J. Mol. Struct. Theochem*, **283**, 33–48.

- Fritsky, I. O., Kozłowski, H., Kandal, O. M., Haukka, M., Świątek-Kozłowska, J., Gumienna-Kontecka, E. & Meyer, F. (2006). *Chem. Commun.* pp. 4125–4127.
- Fritsky, I. O., Kozłowski, H., Sadler, P. J., Yefetova, O. P., Świątek-Kozłowska, J., Kalibabchuk, V. A. & Głowiak, T. (1998). *J. Chem. Soc. Dalton Trans.* pp. 3269–3274.
- Gagné, O. C. & Hawthorne, F. C. (2016). *Acta Cryst.* **B72**, 602–625.
- Grimme, S. (2019). *xtb 6.4*. Mulliken Center for Theoretical Chemistry, University of Bonn, Bonn, Germany.
- Groom, C. R., Bruno, I. J., Lightfoot, M. P. & Ward, S. C. (2016). *Acta Cryst.* **B72**, 171–179.
- Hourahine, B., Aradi, B., Blum, V., Bonafé, F., Buccheri, A., Camacho, C., Cevallos, C., Deshayé, M. Y., Dumitrică, T., Dominguez, A., Ehlert, S., Elstner, M., van der Heide, T., Hermann, J., Irle, S., Kranz, J. J., Köhler, C., Kowalczyk, T., Kubař, T., Lee, I. S., Lutsker, V., Maurer, R. J., Min, S. K., Mitchell, I., Negre, C., Niehaus, T. A., Niklasson, A. M. N., Page, A. J., Pecchia, A., Penazzi, G., Persson, M. P., Řezáč, J., Sánchez, C. G., Sternberg, M., Stöhr, M., Stuckenberg, F., Tkatchenko, A., Yu, V. W. & Frauenheim, T. (2020). *J. Chem. Phys.* **152**, 124101.
- Llunell, M., Casanova, D., Cirera, J., Alemany, P. & Alvarez, S. (2013). *SHAPE*. Barcelona, Spain.
- 9 McKinnon, J. J., Jayatilaka, D. & Spackman, M. A. (2007). *Chem. Commun.* pp. 3814–3816.
- Mokhir, A. A., Gumienna-Kontecka, E., Świątek-Kozłowska, J. J., Petkova, E. G., Fritsky, I. O., Jerzykiewicz, L., Kapshuk, A. A. & Sliva, T. Yu. (2002). *Inorg. Chim. Acta*, **329**, 113–121.
- Oliver, K. J. & Waters, T. N. (1982). *J. Chem. Soc. Chem. Commun.* pp. 1111–1112.
- Otwinowski, Z. & Minor, W. (1997). *Methods Enzymol.* **276**, 307–326.
- Palatinus, L. & Chapuis, G. (2007). *J. Appl. Cryst.* **40**, 786–790.
- Pap, J. S., Szywriel, Ł., Rowińska-Żyrek, M., Nikitin, K., Fritsky, I. O. & Kozłowski, H. J. (2011). *J. Mol. Catal. A Chem.* **334**, 77–82.
- Plutenko, M. O., Haukka, M., Husak, A. O., Iskenderov, T. S. & Mulloev, N. U. (2021). *Acta Cryst.* **E77**, 298–304.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Sheldrick, G. M. (2015). *Acta Cryst.* **C71**, 3–8.
- Shylin, S. I., Pavliuk, M. V., D'Amario, L., Mamedov, F., Sá, J., Berggren, G. & Fritsky, I. O. (2019). *Chem. Commun.* **55**, 3335–3338.
- Spackman, M. A. & Jayatilaka, D. (2009). *CrystEngComm*, **11**, 19–32.
- Stewart, J. J. P. (2013). *J. Mol. Model.* **19**, 1–32.
- Stewart, J. J. P. (2016). *MOPAC2016*. Stewart Computational Chemistry, Colorado Springs, CO, USA.
- Tomyn, S., Shylin, S. I., Bykov, D., Ksenofontov, V., Gumienna-Kontecka, E., Bon, V. & Fritsky, I. O. (2017). *Nat. Commun.* **8**, 14099.
- Turner, M. J., McKinnon, J. J., Wolff, S. K., Grimwood, D. J., Spackman, P. R., Jayatilaka, D. & Spackman, M. A. (2017). *CrystalExplorer17*. University of Western Australia.
- Zheng, G., Witek, H. A., Bobadova-Parvanova, P., Irle, S., Musaev, D. G., Prabhakar, R., Morokuma, K., Lundberg, M., Elstner, M., Köhler, C. & Frauenheim, T. (2007). *J. Chem. Theory Comput.* **3**, 1349–1367.

supporting information

Acta Cryst. (2021). E77 [https://doi.org/10.1107/S2056989021011774]

A second solvatomorph of poly[[μ_4 -N,N'-(1,3,5-oxadiazinane-3,5-diyl)bis-(carbamoylmethanoato)]nickel(II)dipotassium]: crystal structure, Hirshfeld surface analysis and semi-empirical geometry optimization

Maksym O. Plutenko, Matti Haukka, Alina O. Husak, Irina A. Golenya and Nurullo U. Mulloev

Computing details

Data collection: *COLLECT* (Bruker, 2008); cell refinement: *DENZO/SCALEPACK* (Otwinowski & Minor, 1997); data reduction: *DENZO/SCALEPACK* (Otwinowski & Minor, 1997); program(s) used to solve structure: SUPERFLIP (Palatinus & Chapuis, 2007); program(s) used to refine structure: *SHELXL2018/1* (Sheldrick, 2015); molecular graphics: *DIAMOND* (Brandenburg, 2009); software used to prepare material for publication: *SHELXL97* (Sheldrick, 2008).

Poly[triaquabis[μ_4 -N,N'-(1,3,5-oxadiazinane-3,5-diyl)bis(carbamoylmethanoato)]dinickel(II)tetrapotassium]

Crystal data

[K₄Ni₂(C₇H₆N₄O₇)₂(H₂O)₃]
M_r = 844.18
 Monoclinic, *P2₁/c*
a = 20.3825 (5) Å
b = 7.7039 (3) Å
c = 17.3078 (6) Å
 β = 98.240 (2)°
V = 2689.69 (16) Å³
Z = 4

F(000) = 1704
D_x = 2.085 Mg m⁻³
 Mo *K*α radiation, λ = 0.71073 Å
 Cell parameters from 12179 reflections
 θ = 1.0–30.0°
 μ = 2.12 mm⁻¹
T = 100 K
 Orange, block
 0.15 × 0.09 × 0.08 mm

Data collection

Bruker Kappa APEXII CCD
 diffractometer
 Radiation source: fine-focus sealed tube
 Horizontally mounted graphite crystal
 monochromator
 Detector resolution: 16 pixels mm⁻¹
 φ scans and ω scans with κ offset
 Absorption correction: multi-scan
 (SADABS; Sheldrick, 2008)

T_{min} = 0.746, *T_{max}* = 0.842
 25068 measured reflections
 6148 independent reflections
 5118 reflections with *I* > 2σ(*I*)
R_{int} = 0.043
 θ_{\max} = 27.5°, θ_{\min} = 2.5°
h = -26→26
k = -10→10
l = -22→22

Refinement

Refinement on *F*²
 Least-squares matrix: full
R[*F*² > 2σ(*F*²)] = 0.041
wR(*F*²) = 0.082
S = 1.14

6148 reflections
 406 parameters
 0 restraints
 Hydrogen site location: mixed
 H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + 8.0539P]$$

where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$

$$\Delta\rho_{\max} = 0.63 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.45 \text{ e } \text{\AA}^{-3}$$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Ni1	0.47434 (2)	0.39867 (5)	0.07544 (2)	0.01087 (9)
K1	0.17619 (4)	0.53183 (9)	0.11705 (4)	0.01892 (16)
K2	0.19572 (3)	0.36084 (9)	-0.07023 (4)	0.01670 (15)
K3	0.27474 (3)	0.87045 (9)	0.01255 (4)	0.01761 (15)
K4	0.37676 (4)	0.05611 (9)	-0.14504 (4)	0.02025 (16)
O1	0.41934 (10)	0.2347 (3)	0.01663 (13)	0.0155 (5)
O2	0.55123 (10)	0.2576 (3)	0.08187 (12)	0.0138 (4)
O3	0.66101 (11)	0.2786 (3)	0.11647 (14)	0.0178 (5)
O4	0.63860 (11)	0.6017 (3)	0.18521 (13)	0.0175 (5)
O5	0.43257 (10)	0.6088 (3)	0.23723 (12)	0.0144 (4)
O6	0.28491 (10)	0.5095 (3)	0.04127 (13)	0.0144 (5)
O7	0.31296 (10)	0.1927 (3)	-0.03135 (13)	0.0145 (4)
O8	0.23728 (13)	0.4224 (4)	0.26454 (15)	0.0296 (6)
H8O	0.234956	0.311841	0.266821	0.044*
H8P	0.220280	0.460928	0.303719	0.044*
O9	0.22781 (12)	0.4456 (3)	-0.21414 (15)	0.0260 (6)
H9P	0.270092	0.459040	-0.210682	0.039*
H9O	0.209842	0.522979	-0.245902	0.039*
O10	0.32908 (13)	1.1134 (4)	0.14694 (15)	0.0306 (6)
H10O	0.287037	1.137097	0.135669	0.046*
H10P	0.338827	1.126557	0.197749	0.046*
N1	0.52942 (12)	0.5586 (3)	0.13120 (14)	0.0113 (5)
N2	0.51068 (13)	0.7242 (3)	0.15852 (15)	0.0124 (5)
N3	0.39268 (13)	0.6979 (3)	0.10530 (15)	0.0131 (5)
N4	0.39871 (13)	0.5297 (3)	0.07165 (15)	0.0122 (5)
C1	0.60452 (15)	0.3371 (4)	0.11225 (18)	0.0137 (6)
C2	0.59253 (16)	0.5163 (4)	0.14705 (18)	0.0146 (6)
C3	0.45459 (15)	0.7945 (4)	0.10429 (19)	0.0140 (6)
H3A	0.447429	0.917052	0.118117	0.017*
H3B	0.466161	0.792495	0.050662	0.017*
C4	0.49161 (15)	0.7092 (4)	0.23624 (18)	0.0140 (6)
H4A	0.528350	0.654533	0.271547	0.017*
H4B	0.484584	0.826836	0.256513	0.017*
C5	0.37883 (16)	0.6825 (4)	0.18499 (18)	0.0155 (6)
H5A	0.368739	0.799144	0.204239	0.019*
H5B	0.338951	0.609318	0.185349	0.019*

C6	0.34231 (15)	0.4539 (4)	0.04236 (17)	0.0117 (6)
C7	0.35790 (15)	0.2774 (4)	0.00678 (17)	0.0122 (6)
Ni1B	0.01223 (2)	0.08795 (5)	0.09162 (2)	0.01243 (9)
O1B	0.08679 (11)	0.2257 (3)	0.07813 (13)	0.0166 (5)
O2B	-0.04769 (11)	0.2677 (3)	0.05952 (13)	0.0156 (5)
O3B	-0.15683 (11)	0.3148 (3)	0.03336 (14)	0.0184 (5)
O4B	-0.17542 (11)	-0.0118 (3)	0.10065 (13)	0.0164 (5)
O5B	0.00334 (11)	-0.1490 (3)	0.25474 (13)	0.0170 (5)
O6B	0.18685 (11)	-0.1330 (3)	0.14427 (14)	0.0185 (5)
O7B	0.19555 (11)	0.1959 (3)	0.07804 (14)	0.0200 (5)
N1B	-0.06184 (13)	-0.0419 (3)	0.10402 (15)	0.0133 (5)
N2B	-0.06258 (13)	-0.2169 (3)	0.13205 (15)	0.0134 (5)
N3B	0.05905 (13)	-0.2534 (3)	0.14984 (15)	0.0135 (5)
N4B	0.07292 (13)	-0.0845 (3)	0.12181 (15)	0.0140 (5)
C1B	-0.10848 (16)	0.2241 (4)	0.05852 (18)	0.0144 (6)
C2B	-0.11930 (16)	0.0406 (4)	0.09058 (17)	0.0137 (6)
C3B	-0.00599 (15)	-0.3127 (4)	0.10940 (19)	0.0146 (6)
H3B1	-0.006462	-0.300402	0.052383	0.017*
H3B2	-0.011318	-0.437434	0.120686	0.017*
C4B	-0.05815 (16)	-0.2189 (4)	0.21681 (18)	0.0152 (6)
H4B1	-0.095254	-0.150425	0.232299	0.018*
H4B2	-0.062785	-0.339827	0.234578	0.018*
C5B	0.05751 (16)	-0.2475 (4)	0.23366 (19)	0.0158 (6)
H5B1	0.054706	-0.367518	0.253325	0.019*
H5B2	0.099552	-0.196407	0.259598	0.019*
C6B	0.13592 (16)	-0.0447 (4)	0.12181 (18)	0.0154 (6)
C7B	0.14124 (15)	0.1389 (4)	0.08983 (18)	0.0143 (6)

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ni1	0.0117 (2)	0.00866 (18)	0.01215 (19)	0.00042 (14)	0.00137 (14)	-0.00210 (15)
K1	0.0225 (4)	0.0128 (3)	0.0229 (4)	0.0021 (3)	0.0082 (3)	0.0018 (3)
K2	0.0181 (4)	0.0124 (3)	0.0182 (3)	0.0002 (3)	-0.0022 (3)	-0.0009 (3)
K3	0.0162 (3)	0.0130 (3)	0.0229 (4)	-0.0001 (3)	0.0006 (3)	0.0043 (3)
K4	0.0253 (4)	0.0118 (3)	0.0265 (4)	-0.0019 (3)	0.0134 (3)	-0.0026 (3)
O1	0.0140 (11)	0.0129 (11)	0.0195 (12)	-0.0001 (9)	0.0021 (9)	-0.0028 (9)
O2	0.0149 (11)	0.0116 (10)	0.0149 (11)	0.0018 (8)	0.0020 (8)	-0.0011 (9)
O3	0.0124 (11)	0.0158 (11)	0.0253 (13)	0.0022 (9)	0.0030 (9)	-0.0007 (10)
O4	0.0151 (11)	0.0154 (11)	0.0207 (12)	-0.0003 (9)	-0.0016 (9)	-0.0013 (9)
O5	0.0159 (11)	0.0146 (11)	0.0135 (11)	-0.0009 (9)	0.0044 (8)	0.0014 (9)
O6	0.0135 (11)	0.0127 (11)	0.0168 (11)	0.0013 (9)	0.0016 (9)	-0.0016 (9)
O7	0.0144 (11)	0.0138 (11)	0.0148 (11)	-0.0021 (9)	0.0005 (8)	-0.0015 (9)
O8	0.0368 (16)	0.0286 (14)	0.0253 (14)	0.0059 (12)	0.0103 (11)	0.0025 (11)
O9	0.0159 (13)	0.0345 (15)	0.0268 (14)	-0.0018 (11)	0.0006 (10)	0.0045 (11)
O10	0.0259 (14)	0.0390 (16)	0.0269 (14)	0.0052 (12)	0.0038 (11)	0.0047 (12)
N1	0.0137 (13)	0.0095 (12)	0.0107 (12)	0.0008 (10)	0.0022 (10)	-0.0013 (10)
N2	0.0151 (13)	0.0104 (12)	0.0118 (12)	0.0006 (10)	0.0018 (10)	-0.0017 (10)

N3	0.0177 (14)	0.0090 (12)	0.0125 (13)	0.0001 (10)	0.0021 (10)	-0.0040 (10)
N4	0.0176 (14)	0.0075 (12)	0.0116 (12)	0.0018 (10)	0.0024 (10)	-0.0028 (10)
C1	0.0176 (16)	0.0137 (15)	0.0105 (14)	-0.0010 (12)	0.0042 (12)	0.0031 (12)
C2	0.0179 (17)	0.0124 (15)	0.0141 (15)	-0.0020 (12)	0.0039 (12)	0.0019 (12)
C3	0.0158 (16)	0.0091 (14)	0.0164 (15)	0.0002 (12)	0.0002 (12)	-0.0002 (12)
C4	0.0174 (16)	0.0141 (15)	0.0106 (14)	0.0009 (12)	0.0022 (12)	-0.0019 (12)
C5	0.0177 (16)	0.0152 (15)	0.0139 (15)	-0.0005 (12)	0.0036 (12)	-0.0020 (12)
C6	0.0149 (15)	0.0110 (14)	0.0092 (14)	-0.0001 (12)	0.0023 (11)	0.0008 (11)
C7	0.0165 (16)	0.0098 (14)	0.0105 (14)	-0.0003 (11)	0.0026 (11)	0.0020 (11)
Ni1B	0.0128 (2)	0.00920 (19)	0.0149 (2)	-0.00029 (15)	0.00073 (15)	0.00233 (15)
O1B	0.0159 (12)	0.0116 (11)	0.0224 (12)	-0.0001 (9)	0.0028 (9)	0.0032 (9)
O2B	0.0159 (12)	0.0122 (11)	0.0179 (11)	-0.0007 (9)	-0.0001 (9)	0.0019 (9)
O3B	0.0165 (12)	0.0134 (11)	0.0246 (13)	0.0028 (9)	0.0003 (9)	0.0034 (10)
O4B	0.0149 (12)	0.0140 (11)	0.0203 (12)	0.0005 (9)	0.0021 (9)	0.0004 (9)
O5B	0.0202 (12)	0.0165 (12)	0.0138 (11)	0.0000 (9)	0.0007 (9)	-0.0009 (9)
O6B	0.0139 (11)	0.0157 (11)	0.0251 (12)	0.0009 (9)	0.0004 (9)	-0.0003 (10)
O7B	0.0164 (12)	0.0168 (12)	0.0277 (13)	-0.0025 (9)	0.0067 (10)	0.0025 (10)
N1B	0.0168 (14)	0.0075 (12)	0.0152 (13)	-0.0003 (10)	0.0009 (10)	0.0000 (10)
N2B	0.0158 (13)	0.0095 (12)	0.0149 (13)	0.0000 (10)	0.0021 (10)	0.0021 (10)
N3B	0.0154 (13)	0.0094 (12)	0.0154 (13)	-0.0013 (10)	0.0014 (10)	0.0048 (10)
N4B	0.0146 (13)	0.0101 (12)	0.0170 (13)	-0.0006 (10)	0.0018 (10)	0.0021 (10)
C1B	0.0197 (17)	0.0128 (15)	0.0109 (14)	-0.0019 (12)	0.0023 (12)	-0.0021 (12)
C2B	0.0188 (17)	0.0114 (14)	0.0104 (14)	-0.0001 (12)	0.0004 (12)	-0.0018 (11)
C3B	0.0133 (15)	0.0122 (15)	0.0176 (16)	-0.0013 (12)	0.0002 (12)	-0.0008 (12)
C4B	0.0157 (16)	0.0129 (15)	0.0165 (15)	-0.0008 (12)	0.0009 (12)	0.0037 (12)
C5B	0.0177 (17)	0.0127 (15)	0.0168 (16)	0.0014 (12)	0.0014 (12)	0.0041 (12)
C6B	0.0170 (16)	0.0154 (16)	0.0137 (15)	0.0001 (12)	0.0017 (12)	-0.0035 (12)
C7B	0.0163 (16)	0.0144 (15)	0.0122 (15)	-0.0005 (12)	0.0017 (12)	-0.0007 (12)

Geometric parameters (Å, °)

Ni1—N4	1.836 (3)	O5—C4	1.433 (4)
Ni1—N1	1.844 (3)	O5—C5	1.434 (4)
Ni1—O1	1.887 (2)	O6—C6	1.244 (4)
Ni1—O2	1.897 (2)	O7—C7	1.236 (4)
K1—O6B ⁱ	2.628 (2)	O8—H8O	0.8546
K1—O7B	2.717 (2)	O8—H8P	0.8575
K1—O6	2.737 (2)	O9—H9P	0.8615
K1—O8	2.805 (3)	O9—H9O	0.8567
K1—O3B ⁱⁱ	2.834 (2)	O10—H10O	0.8704
K1—O1B	2.998 (2)	O10—H10P	0.8792
K1—N3B ⁱ	3.025 (3)	N1—C2	1.317 (4)
K1—C7B	3.130 (3)	N1—N2	1.431 (3)
K1—C6B ⁱ	3.368 (3)	N2—C4	1.457 (4)
K1—K2	3.5736 (10)	N2—C3	1.474 (4)
K1—K3	3.8927 (10)	N3—N4	1.433 (3)
K2—O6	2.708 (2)	N3—C5	1.452 (4)
K2—O7	2.717 (2)	N3—C3	1.467 (4)

K2—O3B ⁱⁱ	2.725 (2)	N4—C6	1.324 (4)
K2—O9	2.743 (3)	C1—C2	1.540 (4)
K2—O4B ⁱⁱⁱ	2.760 (2)	C3—H3A	0.9900
K2—O7B	2.864 (2)	C3—H3B	0.9900
K2—N2B ⁱⁱⁱ	2.984 (3)	C4—H4A	0.9900
K2—C6	3.401 (3)	C4—H4B	0.9900
K2—C7	3.443 (3)	C5—H5A	0.9900
K2—C2B ⁱⁱⁱ	3.458 (3)	C5—H5B	0.9900
K2—K3 ^{iv}	4.2728 (10)	C6—C7	1.544 (4)
K3—O7 ⁱ	2.742 (2)	Ni1B—N4B	1.839 (3)
K3—O3B ⁱⁱ	2.810 (2)	Ni1B—N1B	1.849 (3)
K3—O4B ⁱⁱ	2.826 (2)	Ni1B—O2B	1.877 (2)
K3—O6	2.827 (2)	Ni1B—O1B	1.895 (2)
K3—O3 ^v	2.976 (2)	O1B—C7B	1.287 (4)
K3—N3	3.003 (3)	O2B—C1B	1.281 (4)
K3—O10	3.066 (3)	O3B—C1B	1.235 (4)
K3—O6B ⁱ	3.095 (2)	O4B—C2B	1.249 (4)
K3—O7B ⁱ	3.271 (2)	O5B—C5B	1.430 (4)
K3—C2B ⁱⁱ	3.475 (3)	O5B—C4B	1.434 (4)
K3—C6	3.501 (3)	O6B—C6B	1.255 (4)
K3—C1B ⁱⁱ	3.512 (3)	O7B—C7B	1.235 (4)
K3—H10O	2.9446	N1B—C2B	1.323 (4)
K4—O7	2.721 (2)	N1B—N2B	1.434 (3)
K4—O4 ^v	2.733 (2)	N2B—C4B	1.457 (4)
K4—O3 ^{vi}	2.756 (2)	N2B—C3B	1.469 (4)
K4—O5 ^{vii}	2.779 (2)	N3B—N4B	1.431 (3)
K4—N2 ^v	2.887 (3)	N3B—C5B	1.456 (4)
K4—O2 ^{vi}	2.955 (2)	N3B—C3B	1.480 (4)
K4—O8 ^{vii}	3.047 (3)	N4B—C6B	1.320 (4)
K4—C1 ^{vi}	3.095 (3)	C1B—C2B	1.546 (4)
K4—O1	3.127 (2)	C3B—H3B1	0.9900
K4—C7	3.201 (3)	C3B—H3B2	0.9900
K4—C2 ^v	3.354 (3)	C4B—H4B1	0.9900
K4—C5 ^{vii}	3.474 (3)	C4B—H4B2	0.9900
O1—C7	1.282 (4)	C5B—H5B1	0.9900
O2—C1	1.291 (4)	C5B—H5B2	0.9900
O3—C1	1.229 (4)	C6B—C7B	1.528 (4)
O4—C2	1.254 (4)		
N4—Ni1—N1	95.53 (11)	N2 ^v —K4—O1	72.10 (7)
N4—Ni1—O1	85.30 (10)	O2 ^{vi} —K4—O1	88.24 (6)
N1—Ni1—O1	178.66 (11)	O8 ^{vii} —K4—O1	123.68 (7)
N4—Ni1—O2	177.92 (11)	C1 ^{vi} —K4—O1	104.79 (7)
N1—Ni1—O2	85.18 (10)	O7—K4—C7	22.25 (7)
O1—Ni1—O2	94.02 (9)	O4 ^v —K4—C7	70.87 (7)
O6B ⁱ —K1—O7B	165.68 (7)	O3 ^{vi} —K4—C7	106.35 (7)
O6B ⁱ —K1—O6	95.50 (7)	O5 ^{vii} —K4—C7	162.77 (8)
O7B—K1—O6	70.41 (7)	N2 ^v —K4—C7	86.82 (8)

O6B ⁱ —K1—O8	96.75 (8)	O2 ^{vi} —K4—C7	104.10 (7)
O7B—K1—O8	83.03 (8)	O8 ^{vii} —K4—C7	100.36 (8)
O6—K1—O8	97.64 (7)	C1 ^{vi} —K4—C7	113.38 (8)
O6B ⁱ —K1—O3B ⁱⁱ	75.65 (7)	O1—K4—C7	23.34 (7)
O7B—K1—O3B ⁱⁱ	100.07 (7)	O7—K4—C2 ^v	74.75 (7)
O6—K1—O3B ⁱⁱ	66.54 (7)	O4 ^v —K4—C2 ^v	20.73 (7)
O8—K1—O3B ⁱⁱ	161.16 (8)	O3 ^{vi} —K4—C2 ^v	167.98 (8)
O6B ⁱ —K1—O1B	147.71 (7)	O5 ^{vii} —K4—C2 ^v	110.16 (7)
O7B—K1—O1B	45.58 (6)	N2 ^v —K4—C2 ^v	43.27 (7)
O6—K1—O1B	110.52 (7)	O2 ^{vi} —K4—C2 ^v	136.36 (7)
O8—K1—O1B	98.30 (7)	O8 ^{vii} —K4—C2 ^v	95.92 (8)
O3B ⁱⁱ —K1—O1B	96.98 (7)	C1 ^{vi} —K4—C2 ^v	160.86 (8)
O6B ⁱ —K1—N3B ⁱ	58.41 (7)	O1—K4—C2 ^v	63.19 (7)
O7B—K1—N3B ⁱ	135.57 (7)	C7—K4—C2 ^v	61.97 (8)
O6—K1—N3B ⁱ	147.07 (7)	O7—K4—C5 ^{vii}	151.88 (7)
O8—K1—N3B ⁱ	104.66 (8)	O4 ^v —K4—C5 ^{vii}	108.05 (7)
O3B ⁱⁱ —K1—N3B ⁱ	86.25 (7)	O3 ^{vi} —K4—C5 ^{vii}	72.21 (7)
O1B—K1—N3B ⁱ	90.10 (7)	O5 ^{vii} —K4—C5 ^{vii}	23.29 (7)
O6B ⁱ —K1—C7B	171.25 (8)	N2 ^v —K4—C5 ^{vii}	97.79 (7)
O7B—K1—C7B	23.02 (7)	O2 ^{vi} —K4—C5 ^{vii}	79.33 (7)
O6—K1—C7B	92.84 (8)	O8 ^{vii} —K4—C5 ^{vii}	73.58 (7)
O8—K1—C7B	84.73 (8)	C1 ^{vi} —K4—C5 ^{vii}	67.75 (8)
O3B ⁱⁱ —K1—C7B	105.44 (8)	O1—K4—C5 ^{vii}	162.73 (7)
O1B—K1—C7B	24.11 (7)	C7—K4—C5 ^{vii}	173.84 (8)
N3B ⁱ —K1—C7B	112.86 (8)	C2 ^v —K4—C5 ^{vii}	119.08 (8)
O6B ⁱ —K1—C6B ⁱ	19.63 (7)	C7—O1—Ni1	113.14 (19)
O7B—K1—C6B ⁱ	166.68 (8)	C7—O1—K4	81.58 (17)
O6—K1—C6B ⁱ	106.83 (7)	Ni1—O1—K4	147.60 (10)
O8—K1—C6B ⁱ	110.29 (8)	C1—O2—Ni1	113.09 (19)
O3B ⁱⁱ —K1—C6B ⁱ	67.44 (7)	C1—O2—K4 ^{vi}	83.85 (17)
O1B—K1—C6B ⁱ	128.70 (7)	Ni1—O2—K4 ^{vi}	148.49 (10)
N3B ⁱ —K1—C6B ⁱ	42.60 (7)	C1—O3—K4 ^{vi}	94.03 (19)
C7B—K1—C6B ⁱ	152.81 (8)	C1—O3—K3 ^v	127.3 (2)
O6B ⁱ —K1—K2	120.51 (6)	K4 ^{vi} —O3—K3 ^v	86.44 (6)
O7B—K1—K2	52.02 (5)	C2—O4—K4 ^v	108.80 (19)
O6—K1—K2	48.63 (5)	C4—O5—C5	110.2 (2)
O8—K1—K2	128.47 (6)	C4—O5—K4 ^{viii}	133.50 (17)
O3B ⁱⁱ —K1—K2	48.67 (5)	C5—O5—K4 ^{viii}	106.70 (16)
O1B—K1—K2	69.47 (5)	C6—O6—K2	113.52 (19)
N3B ⁱ —K1—K2	124.43 (6)	C6—O6—K1	147.0 (2)
C7B—K1—K2	63.86 (6)	K2—O6—K1	82.02 (6)
C6B ⁱ —K1—K2	116.01 (6)	C6—O6—K3	112.75 (18)
O6B ⁱ —K1—K3	52.41 (5)	K2—O6—K3	105.41 (7)
O7B—K1—K3	114.82 (5)	K1—O6—K3	88.76 (6)
O6—K1—K3	46.56 (5)	C7—O7—K2	116.01 (19)
O8—K1—K3	115.66 (6)	C7—O7—K4	101.30 (18)
O3B ⁱⁱ —K1—K3	46.14 (5)	K2—O7—K4	119.99 (8)
O1B—K1—K3	139.50 (5)	C7—O7—K3 ^{iv}	123.27 (19)

N3B ⁱ —K1—K3	101.21 (5)	K2—O7—K3 ^{iv}	103.04 (7)
C7B—K1—K3	134.40 (6)	K4—O7—K3 ^{iv}	91.98 (7)
C6B ⁱ —K1—K3	60.40 (6)	K1—O8—K4 ^{viii}	135.32 (10)
K2—K1—K3	72.16 (2)	K1—O8—H8O	108.7
O6—K2—O7	63.15 (6)	K4 ^{viii} —O8—H8O	94.8
O6—K2—O3B ⁱⁱ	68.50 (7)	K1—O8—H8P	116.0
O7—K2—O3B ⁱⁱ	130.88 (7)	K4 ^{viii} —O8—H8P	91.6
O6—K2—O9	108.86 (7)	H8O—O8—H8P	106.1
O7—K2—O9	91.30 (7)	K2—O9—H9P	109.4
O3B ⁱⁱ —K2—O9	96.29 (8)	K2—O9—H9O	127.8
O6—K2—O4B ⁱⁱⁱ	127.95 (7)	H9P—O9—H9O	107.0
O7—K2—O4B ⁱⁱⁱ	71.65 (7)	K3—O10—H10O	73.8
O3B ⁱⁱ —K2—O4B ⁱⁱⁱ	153.83 (7)	K3—O10—H10P	146.5
O9—K2—O4B ⁱⁱⁱ	96.17 (8)	H10O—O10—H10P	105.9
O6—K2—O7B	68.65 (7)	C2—N1—N2	116.8 (3)
O7—K2—O7B	71.37 (7)	C2—N1—Ni1	116.5 (2)
O3B ⁱⁱ —K2—O7B	99.16 (7)	N2—N1—Ni1	126.66 (19)
O9—K2—O7B	161.85 (8)	N1—N2—C4	110.7 (2)
O4B ⁱⁱⁱ —K2—O7B	73.70 (7)	N1—N2—C3	109.7 (2)
O6—K2—N2B ⁱⁱⁱ	153.59 (7)	C4—N2—C3	109.4 (2)
O7—K2—N2B ⁱⁱⁱ	129.35 (7)	N1—N2—K4 ^v	104.06 (16)
O3B ⁱⁱ —K2—N2B ⁱⁱⁱ	98.31 (7)	C4—N2—K4 ^v	116.19 (18)
O9—K2—N2B ⁱⁱⁱ	94.92 (7)	C3—N2—K4 ^v	106.62 (17)
O4B ⁱⁱⁱ —K2—N2B ⁱⁱⁱ	57.71 (7)	N4—N3—C5	110.6 (2)
O7B—K2—N2B ⁱⁱⁱ	92.22 (7)	N4—N3—C3	109.3 (2)
O6—K2—C6	19.59 (7)	C5—N3—C3	109.7 (2)
O7—K2—C6	44.71 (7)	N4—N3—K3	106.96 (16)
O3B ⁱⁱ —K2—C6	86.20 (7)	C5—N3—K3	107.10 (18)
O9—K2—C6	99.55 (7)	C3—N3—K3	113.13 (18)
O4B ⁱⁱⁱ —K2—C6	114.21 (7)	C6—N4—N3	115.7 (2)
O7B—K2—C6	72.17 (7)	C6—N4—Ni1	116.7 (2)
N2B ⁱⁱⁱ —K2—C6	164.29 (7)	N3—N4—Ni1	127.1 (2)
O6—K2—C7	45.08 (7)	O3—C1—O2	125.2 (3)
O7—K2—C7	18.82 (7)	O3—C1—C2	120.4 (3)
O3B ⁱⁱ —K2—C7	112.08 (7)	O2—C1—C2	114.4 (3)
O9—K2—C7	93.17 (7)	O3—C1—K4 ^{vi}	62.64 (17)
O4B ⁱⁱⁱ —K2—C7	90.08 (7)	O2—C1—K4 ^{vi}	71.65 (16)
O7B—K2—C7	72.22 (7)	C2—C1—K4 ^{vi}	145.90 (19)
N2B ⁱⁱⁱ —K2—C7	147.43 (7)	O4—C2—N1	127.9 (3)
C6—K2—C7	26.07 (7)	O4—C2—C1	121.8 (3)
O6—K2—C2B ⁱⁱⁱ	134.22 (7)	N1—C2—C1	110.2 (3)
O7—K2—C2B ⁱⁱⁱ	88.06 (7)	O4—C2—K4 ^v	50.47 (16)
O3B ⁱⁱ —K2—C2B ⁱⁱⁱ	134.59 (7)	N1—C2—K4 ^v	86.15 (18)
O9—K2—C2B ⁱⁱⁱ	106.44 (8)	C1—C2—K4 ^v	146.74 (19)
O4B ⁱⁱⁱ —K2—C2B ⁱⁱⁱ	19.29 (7)	N3—C3—N2	113.3 (2)
O7B—K2—C2B ⁱⁱⁱ	68.67 (7)	N3—C3—H3A	108.9
N2B ⁱⁱⁱ —K2—C2B ⁱⁱⁱ	42.10 (7)	N2—C3—H3A	108.9
C6—K2—C2B ⁱⁱⁱ	126.35 (7)	N3—C3—H3B	108.9

C7—K2—C2B ⁱⁱⁱ	105.43 (7)	N2—C3—H3B	108.9
O6—K2—K1	49.34 (5)	H3A—C3—H3B	107.7
O7—K2—K1	99.18 (5)	O5—C4—N2	112.9 (2)
O3B ⁱⁱ —K2—K1	51.36 (5)	O5—C4—H4A	109.0
O9—K2—K1	143.89 (6)	N2—C4—H4A	109.0
O4B ⁱⁱⁱ —K2—K1	119.94 (5)	O5—C4—H4B	109.0
O7B—K2—K1	48.41 (5)	N2—C4—H4B	109.0
N2B ⁱⁱⁱ —K2—K1	104.38 (6)	H4A—C4—H4B	107.8
C6—K2—K1	66.77 (5)	O5—C5—N3	113.3 (2)
C7—K2—K1	86.75 (5)	O5—C5—K4 ^{viii}	50.01 (13)
C2B ⁱⁱⁱ —K2—K1	108.36 (5)	N3—C5—K4 ^{viii}	151.1 (2)
O6—K2—K3 ^{iv}	87.29 (5)	O5—C5—H5A	108.9
O7—K2—K3 ^{iv}	38.69 (5)	N3—C5—H5A	108.9
O3B ⁱⁱ —K2—K3 ^{iv}	147.21 (6)	K4 ^{viii} —C5—H5A	99.6
O9—K2—K3 ^{iv}	112.80 (6)	O5—C5—H5B	108.9
O4B ⁱⁱⁱ —K2—K3 ^{iv}	40.67 (5)	N3—C5—H5B	108.9
O7B—K2—K3 ^{iv}	49.92 (5)	K4 ^{viii} —C5—H5B	65.4
N2B ⁱⁱⁱ —K2—K3 ^{iv}	94.10 (5)	H5A—C5—H5B	107.7
C6—K2—K3 ^{iv}	74.71 (5)	O6—C6—N4	128.1 (3)
C7—K2—K3 ^{iv}	53.87 (5)	O6—C6—C7	123.0 (3)
C2B ⁱⁱⁱ —K2—K3 ^{iv}	52.13 (5)	N4—C6—C7	108.9 (3)
K1—K2—K3 ^{iv}	96.17 (2)	O6—C6—K2	46.89 (15)
O7 ⁱ —K3—O3B ⁱⁱ	130.32 (7)	N4—C6—K2	162.9 (2)
O7 ⁱ —K3—O4B ⁱⁱ	70.29 (6)	C7—C6—K2	78.47 (16)
O3B ⁱⁱ —K3—O4B ⁱⁱ	60.07 (7)	O6—C6—K3	48.13 (15)
O7 ⁱ —K3—O6	157.32 (7)	N4—C6—K3	87.33 (18)
O3B ⁱⁱ —K3—O6	65.69 (6)	C7—C6—K3	147.92 (19)
O4B ⁱⁱ —K3—O6	121.88 (7)	K2—C6—K3	79.28 (7)
O7 ⁱ —K3—O3 ^v	88.18 (7)	O7—C7—O1	124.8 (3)
O3B ⁱⁱ —K3—O3 ^v	92.43 (7)	O7—C7—C6	119.9 (3)
O4B ⁱⁱ —K3—O3 ^v	88.69 (7)	O1—C7—C6	115.2 (3)
O6—K3—O3 ^v	73.94 (6)	O7—C7—K4	56.45 (16)
O7 ⁱ —K3—N3	108.15 (7)	O1—C7—K4	75.08 (16)
O3B ⁱⁱ —K3—N3	120.86 (7)	C6—C7—K4	148.43 (19)
O4B ⁱⁱ —K3—N3	168.61 (7)	O7—C7—K2	45.16 (15)
O6—K3—N3	55.78 (7)	O1—C7—K2	164.7 (2)
O3 ^v —K3—N3	79.96 (7)	C6—C7—K2	75.46 (16)
O7 ⁱ —K3—O10	64.62 (7)	K4—C7—K2	90.18 (8)
O3B ⁱⁱ —K3—O10	136.54 (7)	N4B—Ni1B—N1B	95.93 (11)
O4B ⁱⁱ —K3—O10	115.82 (7)	N4B—Ni1B—O2B	178.25 (11)
O6—K3—O10	117.23 (7)	N1B—Ni1B—O2B	85.75 (10)
O3 ^v —K3—O10	130.90 (7)	N4B—Ni1B—O1B	85.46 (10)
N3—K3—O10	71.96 (7)	N1B—Ni1B—O1B	178.61 (11)
O7 ⁱ —K3—O6B ⁱ	115.43 (7)	O2B—Ni1B—O1B	92.86 (9)
O3B ⁱⁱ —K3—O6B ⁱ	69.03 (7)	C7B—O1B—Ni1B	112.2 (2)
O4B ⁱⁱ —K3—O6B ⁱ	94.66 (6)	C7B—O1B—K1	83.73 (17)
O6—K3—O6B ⁱ	84.13 (6)	Ni1B—O1B—K1	152.27 (11)
O3 ^v —K3—O6B ⁱ	155.90 (7)	C1B—O2B—Ni1B	113.4 (2)

N3—K3—O6B ⁱ	96.08 (7)	C1B—O3B—K2 ⁱⁱ	132.9 (2)
O10—K3—O6B ⁱ	68.32 (7)	C1B—O3B—K3 ⁱⁱ	114.8 (2)
O7 ⁱ —K3—O7B ⁱ	64.91 (6)	K2 ⁱⁱ —O3B—K3 ⁱⁱ	105.44 (8)
O3B ⁱⁱ —K3—O7B ⁱ	92.46 (6)	C1B—O3B—K1 ⁱⁱ	124.0 (2)
O4B ⁱⁱ —K3—O7B ⁱ	66.72 (6)	K2 ⁱⁱ —O3B—K1 ⁱⁱ	79.97 (6)
O6—K3—O7B ⁱ	136.08 (6)	K3 ⁱⁱ —O3B—K1 ⁱⁱ	87.20 (7)
O3 ^v —K3—O7B ⁱ	147.96 (7)	C2B—O4B—K2 ⁱⁱⁱ	113.82 (19)
N3—K3—O7B ⁱ	123.44 (7)	C2B—O4B—K3 ⁱⁱ	110.98 (19)
O10—K3—O7B ⁱ	53.97 (6)	K2 ⁱⁱⁱ —O4B—K3 ⁱⁱ	99.81 (7)
O6B ⁱ —K3—O7B ⁱ	52.04 (6)	C5B—O5B—C4B	109.9 (2)
O7 ⁱ —K3—C2B ⁱⁱ	87.34 (7)	C6B—O6B—K1 ^{iv}	115.7 (2)
O3B ⁱⁱ —K3—C2B ⁱⁱ	43.53 (7)	C6B—O6B—K3 ^{iv}	107.9 (2)
O4B ⁱⁱ —K3—C2B ⁱⁱ	19.60 (7)	K1 ^{iv} —O6B—K3 ^{iv}	85.30 (7)
O6—K3—C2B ⁱⁱ	108.83 (7)	C7B—O7B—K1	97.60 (19)
O3 ^v —K3—C2B ⁱⁱ	99.07 (7)	C7B—O7B—K2	115.0 (2)
N3—K3—C2B ⁱⁱ	164.39 (7)	K1—O7B—K2	79.58 (6)
O10—K3—C2B ⁱⁱ	118.22 (7)	C7B—O7B—K3 ^{iv}	106.4 (2)
O6B ⁱ —K3—C2B ⁱⁱ	78.38 (7)	K1—O7B—K3 ^{iv}	155.85 (9)
O7B ⁱ —K3—C2B ⁱⁱ	64.41 (7)	K2—O7B—K3 ^{iv}	88.02 (6)
O7 ⁱ —K3—C6	138.54 (7)	C2B—N1B—N2B	117.3 (3)
O3B ⁱⁱ —K3—C6	83.02 (7)	C2B—N1B—Ni1B	115.9 (2)
O4B ⁱⁱ —K3—C6	133.08 (7)	N2B—N1B—Ni1B	126.6 (2)
O6—K3—C6	19.12 (6)	N1B—N2B—C4B	110.4 (2)
O3 ^v —K3—C6	63.31 (7)	N1B—N2B—C3B	109.6 (2)
N3—K3—C6	41.19 (7)	C4B—N2B—C3B	109.0 (2)
O10—K3—C6	110.86 (7)	N1B—N2B—K2 ⁱⁱⁱ	106.33 (17)
O6B ⁱ —K3—C6	98.09 (7)	C4B—N2B—K2 ⁱⁱⁱ	106.21 (18)
O7B ⁱ —K3—C6	148.73 (7)	C3B—N2B—K2 ⁱⁱⁱ	115.23 (18)
C2B ⁱⁱ —K3—C6	124.54 (7)	N4B—N3B—C5B	110.0 (2)
O7 ⁱ —K3—C1B ⁱⁱ	112.89 (7)	N4B—N3B—C3B	109.1 (2)
O3B ⁱⁱ —K3—C1B ⁱⁱ	18.63 (7)	C5B—N3B—C3B	109.6 (2)
O4B ⁱⁱ —K3—C1B ⁱⁱ	43.75 (7)	N4B—N3B—K1 ^{iv}	103.54 (16)
O6—K3—C1B ⁱⁱ	84.28 (7)	C5B—N3B—K1 ^{iv}	109.54 (18)
O3 ^v —K3—C1B ⁱⁱ	99.41 (7)	C3B—N3B—K1 ^{iv}	114.86 (18)
N3—K3—C1B ⁱⁱ	138.92 (7)	C6B—N4B—N3B	116.8 (3)
O10—K3—C1B ⁱⁱ	128.16 (7)	C6B—N4B—Ni1B	116.5 (2)
O6B ⁱ —K3—C1B ⁱⁱ	67.97 (7)	N3B—N4B—Ni1B	126.6 (2)
O7B ⁱ —K3—C1B ⁱⁱ	77.36 (7)	O3B—C1B—O2B	125.3 (3)
C2B ⁱⁱ —K3—C1B ⁱⁱ	25.56 (7)	O3B—C1B—C2B	119.7 (3)
C6—K3—C1B ⁱⁱ	101.59 (7)	O2B—C1B—C2B	115.0 (3)
O7 ⁱ —K3—H100	64.6	O3B—C1B—K3 ⁱⁱ	46.60 (16)
O3B ⁱⁱ —K3—H100	122.7	O2B—C1B—K3 ⁱⁱ	160.3 (2)
O4B ⁱⁱ —K3—H100	102.2	C2B—C1B—K3 ⁱⁱ	75.88 (17)
O6—K3—H100	124.2	O4B—C2B—N1B	128.7 (3)
O3 ^v —K3—H100	144.2	O4B—C2B—C1B	121.8 (3)
N3—K3—H100	86.8	N1B—C2B—C1B	109.5 (3)
O10—K3—H100	16.5	O4B—C2B—K2 ⁱⁱⁱ	46.89 (15)
O6B ⁱ —K3—H100	58.1	N1B—C2B—K2 ⁱⁱⁱ	87.87 (18)

O7B ⁱ —K3—H100	37.8	C1B—C2B—K2 ⁱⁱⁱ	150.1 (2)
C2B ⁱⁱ —K3—H100	102.2	O4B—C2B—K3 ⁱⁱ	49.41 (15)
C6—K3—H100	122.8	N1B—C2B—K3 ⁱⁱ	154.7 (2)
C1B ⁱⁱ —K3—H100	112.1	C1B—C2B—K3 ⁱⁱ	78.56 (17)
O7—K4—O4 ^v	76.22 (7)	K2 ⁱⁱⁱ —C2B—K3 ⁱⁱ	76.10 (7)
O7—K4—O3 ^{vi}	93.28 (7)	N2B—C3B—N3B	113.8 (3)
O4 ^v —K4—O3 ^{vi}	157.43 (7)	N2B—C3B—H3B1	108.8
O7—K4—O5 ^{vii}	174.39 (7)	N3B—C3B—H3B1	108.8
O4 ^v —K4—O5 ^{vii}	107.36 (7)	N2B—C3B—H3B2	108.8
O3 ^{vi} —K4—O5 ^{vii}	81.84 (7)	N3B—C3B—H3B2	108.8
O7—K4—N2 ^v	107.43 (7)	H3B1—C3B—H3B2	107.7
O4 ^v —K4—N2 ^v	58.50 (7)	O5B—C4B—N2B	112.4 (3)
O3 ^{vi} —K4—N2 ^v	143.96 (7)	O5B—C4B—H4B1	109.1
O5 ^{vii} —K4—N2 ^v	78.18 (7)	N2B—C4B—H4B1	109.1
O7—K4—O2 ^{vi}	108.20 (7)	O5B—C4B—H4B2	109.1
O4 ^v —K4—O2 ^{vi}	156.33 (7)	N2B—C4B—H4B2	109.1
O3 ^{vi} —K4—O2 ^{vi}	45.97 (6)	H4B1—C4B—H4B2	107.9
O5 ^{vii} —K4—O2 ^{vi}	70.34 (6)	O5B—C5B—N3B	113.5 (2)
N2 ^v —K4—O2 ^{vi}	98.69 (7)	O5B—C5B—H5B1	108.9
O7—K4—O8 ^{vii}	80.97 (7)	N3B—C5B—H5B1	108.9
O4 ^v —K4—O8 ^{vii}	75.67 (7)	O5B—C5B—H5B2	108.9
O3 ^{vi} —K4—O8 ^{vii}	83.04 (7)	N3B—C5B—H5B2	108.9
O5 ^{vii} —K4—O8 ^{vii}	95.62 (7)	H5B1—C5B—H5B2	107.7
N2 ^v —K4—O8 ^{vii}	128.35 (8)	O6B—C6B—N4B	129.5 (3)
O2 ^{vi} —K4—O8 ^{vii}	127.72 (7)	O6B—C6B—C7B	121.0 (3)
O7—K4—C1 ^{vi}	107.76 (8)	N4B—C6B—C7B	109.5 (3)
O4 ^v —K4—C1 ^{vi}	175.75 (8)	O6B—C6B—K1 ^{iv}	44.70 (15)
O3 ^{vi} —K4—C1 ^{vi}	23.33 (7)	N4B—C6B—K1 ^{iv}	90.89 (19)
O5 ^{vii} —K4—C1 ^{vi}	68.56 (7)	C7B—C6B—K1 ^{iv}	149.0 (2)
N2 ^v —K4—C1 ^{vi}	120.66 (8)	O7B—C7B—O1B	124.2 (3)
O2 ^{vi} —K4—C1 ^{vi}	24.50 (7)	O7B—C7B—C6B	120.1 (3)
O8 ^{vii} —K4—C1 ^{vi}	103.22 (8)	O1B—C7B—C6B	115.7 (3)
O7—K4—O1	44.19 (6)	O7B—C7B—K1	59.38 (17)
O4 ^v —K4—O1	79.06 (6)	O1B—C7B—K1	72.16 (17)
O3 ^{vi} —K4—O1	107.45 (7)	C6B—C7B—K1	150.1 (2)
O5 ^{vii} —K4—O1	140.07 (6)		
N4—Ni1—O1—C7	-5.2 (2)	K3—C6—C7—O1	121.4 (3)
O2—Ni1—O1—C7	172.8 (2)	O6—C6—C7—K4	80.9 (5)
N4—Ni1—O1—K4	106.9 (2)	N4—C6—C7—K4	-98.5 (4)
O2—Ni1—O1—K4	-75.08 (19)	K2—C6—C7—K4	65.4 (3)
N1—Ni1—O2—C1	-6.6 (2)	K3—C6—C7—K4	18.5 (7)
O1—Ni1—O2—C1	172.3 (2)	O6—C6—C7—K2	15.5 (3)
N1—Ni1—O2—K4 ^{vi}	111.6 (2)	N4—C6—C7—K2	-163.9 (2)
O1—Ni1—O2—K4 ^{vi}	-69.5 (2)	K3—C6—C7—K2	-46.9 (3)
N4—Ni1—N1—C2	-179.1 (2)	N4B—Ni1B—O1B—C7B	4.4 (2)
O2—Ni1—N1—C2	2.9 (2)	O2B—Ni1B—O1B—C7B	-175.1 (2)
N4—Ni1—N1—N2	0.5 (2)	N4B—Ni1B—O1B—K1	-117.2 (2)

O2—Ni1—N1—N2	-177.6 (2)	O2B—Ni1B—O1B—K1	63.2 (2)
C2—N1—N2—C4	-90.7 (3)	N1B—Ni1B—O2B—C1B	-1.5 (2)
Ni1—N1—N2—C4	89.7 (3)	O1B—Ni1B—O2B—C1B	178.5 (2)
C2—N1—N2—C3	148.5 (3)	N4B—Ni1B—N1B—C2B	177.3 (2)
Ni1—N1—N2—C3	-31.1 (3)	O2B—Ni1B—N1B—C2B	-3.2 (2)
C2—N1—N2—K4 ^v	34.8 (3)	N4B—Ni1B—N1B—N2B	2.3 (3)
Ni1—N1—N2—K4 ^v	-144.78 (16)	O2B—Ni1B—N1B—N2B	-178.2 (2)
C5—N3—N4—C6	83.1 (3)	C2B—N1B—N2B—C4B	-86.4 (3)
C3—N3—N4—C6	-156.0 (3)	Ni1B—N1B—N2B—C4B	88.5 (3)
K3—N3—N4—C6	-33.3 (3)	C2B—N1B—N2B—C3B	153.5 (3)
C5—N3—N4—Ni1	-88.9 (3)	Ni1B—N1B—N2B—C3B	-31.5 (3)
C3—N3—N4—Ni1	32.0 (3)	C2B—N1B—N2B—K2 ⁱⁱⁱ	28.4 (3)
K3—N3—N4—Ni1	154.80 (16)	Ni1B—N1B—N2B—K2 ⁱⁱⁱ	-156.67 (16)
N1—Ni1—N4—C6	-172.8 (2)	C5B—N3B—N4B—C6B	89.9 (3)
O1—Ni1—N4—C6	8.2 (2)	C3B—N3B—N4B—C6B	-149.9 (3)
N1—Ni1—N4—N3	-1.0 (2)	K1 ^{iv} —N3B—N4B—C6B	-27.1 (3)
O1—Ni1—N4—N3	-179.9 (2)	C5B—N3B—N4B—Ni1B	-85.9 (3)
K4 ^{vi} —O3—C1—O2	-36.9 (3)	C3B—N3B—N4B—Ni1B	34.3 (3)
K3 ^v —O3—C1—O2	51.7 (4)	K1 ^{iv} —N3B—N4B—Ni1B	157.07 (16)
K4 ^{vi} —O3—C1—C2	141.0 (2)	N1B—Ni1B—N4B—C6B	-179.7 (2)
K3 ^v —O3—C1—C2	-130.4 (2)	O1B—Ni1B—N4B—C6B	0.3 (2)
K3 ^v —O3—C1—K4 ^v	88.59 (18)	N1B—Ni1B—N4B—N3B	-3.9 (3)
Ni1—O2—C1—O3	-173.4 (2)	O1B—Ni1B—N4B—N3B	176.1 (2)
K4 ^{vi} —O2—C1—O3	34.2 (3)	K2 ⁱⁱ —O3B—C1B—O2B	-57.3 (4)
Ni1—O2—C1—C2	8.5 (3)	K3 ⁱⁱ —O3B—C1B—O2B	156.7 (2)
K4 ^{vi} —O2—C1—C2	-143.9 (2)	K1 ⁱⁱ —O3B—C1B—O2B	52.5 (4)
Ni1—O2—C1—K4 ^v	152.39 (15)	K2 ⁱⁱ —O3B—C1B—C2B	124.0 (3)
K4 ^v —O4—C2—N1	-41.1 (4)	K3 ⁱⁱ —O3B—C1B—C2B	-22.0 (3)
K4 ^v —O4—C2—C1	139.8 (2)	K1 ⁱⁱ —O3B—C1B—C2B	-126.2 (2)
N2—N1—C2—O4	2.0 (5)	K2 ⁱⁱ —O3B—C1B—K3 ⁱⁱ	146.0 (3)
Ni1—N1—C2—O4	-178.4 (3)	K1 ⁱⁱ —O3B—C1B—K3 ⁱⁱ	-104.2 (2)
N2—N1—C2—C1	-178.8 (2)	Ni1B—O2B—C1B—O3B	-173.6 (3)
Ni1—N1—C2—C1	0.8 (3)	Ni1B—O2B—C1B—C2B	5.2 (3)
N2—N1—C2—K4 ^v	-28.5 (2)	Ni1B—O2B—C1B—K3 ⁱⁱ	-115.0 (6)
Ni1—N1—C2—K4 ^v	151.09 (15)	K2 ⁱⁱⁱ —O4B—C2B—N1B	-35.3 (4)
O3—C1—C2—O4	-5.2 (5)	K3 ⁱⁱ —O4B—C2B—N1B	-147.0 (3)
O2—C1—C2—O4	173.0 (3)	K2 ⁱⁱⁱ —O4B—C2B—C1B	144.7 (2)
K4 ^{vi} —C1—C2—O4	79.9 (4)	K3 ⁱⁱ —O4B—C2B—C1B	33.1 (3)
O3—C1—C2—N1	175.5 (3)	K3 ⁱⁱ —O4B—C2B—K2 ⁱⁱⁱ	-111.64 (18)
O2—C1—C2—N1	-6.3 (4)	K2 ⁱⁱⁱ —O4B—C2B—K3 ⁱⁱ	111.64 (18)
K4 ^{vi} —C1—C2—N1	-99.3 (4)	N2B—N1B—C2B—O4B	1.8 (5)
O3—C1—C2—K4 ^v	60.0 (5)	Ni1B—N1B—C2B—O4B	-173.7 (3)
O2—C1—C2—K4 ^v	-121.8 (3)	N2B—N1B—C2B—C1B	-178.3 (2)
K4 ^{vi} —C1—C2—K4 ^v	145.1 (2)	Ni1B—N1B—C2B—C1B	6.2 (3)
N4—N3—C3—N2	-69.2 (3)	N2B—N1B—C2B—K2 ⁱⁱⁱ	-23.2 (2)
C5—N3—C3—N2	52.2 (3)	Ni1B—N1B—C2B—K2 ⁱⁱⁱ	161.31 (15)
K3—N3—C3—N2	171.75 (18)	N2B—N1B—C2B—K3 ⁱⁱ	-73.4 (6)
N1—N2—C3—N3	69.0 (3)	Ni1B—N1B—C2B—K3 ⁱⁱ	111.1 (5)

C4—N2—C3—N3	-52.6 (3)	O3B—C1B—C2B—O4B	-8.7 (4)
K4 ^v —N2—C3—N3	-178.98 (19)	O2B—C1B—C2B—O4B	172.4 (3)
C5—O5—C4—N2	-57.2 (3)	K3 ⁱⁱ —C1B—C2B—O4B	-25.0 (3)
K4 ^{viii} —O5—C4—N2	162.24 (18)	O3B—C1B—C2B—N1B	171.3 (3)
N1—N2—C4—O5	-66.0 (3)	O2B—C1B—C2B—N1B	-7.5 (4)
C3—N2—C4—O5	54.9 (3)	K3 ⁱⁱ —C1B—C2B—N1B	155.0 (2)
K4 ^v —N2—C4—O5	175.62 (17)	O3B—C1B—C2B—K2 ⁱⁱⁱ	48.9 (5)
C4—O5—C5—N3	56.9 (3)	O2B—C1B—C2B—K2 ⁱⁱⁱ	-129.9 (3)
K4 ^{viii} —O5—C5—N3	-151.9 (2)	K3 ⁱⁱ —C1B—C2B—K2 ⁱⁱⁱ	32.6 (4)
C4—O5—C5—K4 ^{viii}	-151.2 (2)	O3B—C1B—C2B—K3 ⁱⁱ	16.3 (3)
N4—N3—C5—O5	66.4 (3)	O2B—C1B—C2B—K3 ⁱⁱ	-162.6 (3)
C3—N3—C5—O5	-54.2 (3)	N1B—N2B—C3B—N3B	68.4 (3)
K3—N3—C5—O5	-177.35 (19)	C4B—N2B—C3B—N3B	-52.5 (3)
N4—N3—C5—K4 ^{viii}	18.2 (5)	K2 ⁱⁱⁱ —N2B—C3B—N3B	-171.76 (18)
C3—N3—C5—K4 ^{viii}	-102.4 (4)	N4B—N3B—C3B—N2B	-69.9 (3)
K3—N3—C5—K4 ^{viii}	134.4 (3)	C5B—N3B—C3B—N2B	50.6 (3)
K2—O6—C6—N4	158.3 (3)	K1 ^{iv} —N3B—C3B—N2B	174.42 (18)
K1—O6—C6—N4	-88.6 (5)	C5B—O5B—C4B—N2B	-59.4 (3)
K3—O6—C6—N4	38.5 (4)	N1B—N2B—C4B—O5B	-63.8 (3)
K2—O6—C6—C7	-21.0 (3)	C3B—N2B—C4B—O5B	56.6 (3)
K1—O6—C6—C7	92.1 (4)	K2 ⁱⁱⁱ —N2B—C4B—O5B	-178.71 (19)
K3—O6—C6—C7	-140.8 (2)	C4B—O5B—C5B—N3B	57.7 (3)
K1—O6—C6—K2	113.1 (4)	N4B—N3B—C5B—O5B	67.1 (3)
K3—O6—C6—K2	-119.79 (19)	C3B—N3B—C5B—O5B	-52.9 (3)
K2—O6—C6—K3	119.79 (19)	K1 ^{iv} —N3B—C5B—O5B	-179.72 (19)
K1—O6—C6—K3	-127.1 (4)	K1 ^{iv} —O6B—C6B—N4B	36.6 (4)
N3—N4—C6—O6	-0.9 (5)	K3 ^{iv} —O6B—C6B—N4B	130.0 (3)
Ni1—N4—C6—O6	171.9 (2)	K1 ^{iv} —O6B—C6B—C7B	-144.6 (2)
N3—N4—C6—C7	178.5 (2)	K3 ^{iv} —O6B—C6B—C7B	-51.1 (3)
Ni1—N4—C6—C7	-8.7 (3)	K3 ^{iv} —O6B—C6B—K1 ^{iv}	93.42 (17)
N3—N4—C6—K2	65.3 (8)	N3B—N4B—C6B—O6B	-1.3 (5)
Ni1—N4—C6—K2	-121.9 (6)	Ni1B—N4B—C6B—O6B	174.9 (3)
N3—N4—C6—K3	26.8 (2)	N3B—N4B—C6B—C7B	179.7 (2)
Ni1—N4—C6—K3	-160.42 (15)	Ni1B—N4B—C6B—C7B	-4.0 (3)
K2—O7—C7—O1	-164.9 (2)	N3B—N4B—C6B—K1 ^{iv}	23.5 (2)
K4—O7—C7—O1	-33.2 (3)	Ni1B—N4B—C6B—K1 ^{iv}	-160.30 (15)
K3 ^{iv} —O7—C7—O1	66.7 (4)	K1—O7B—C7B—O1B	33.6 (3)
K2—O7—C7—C6	11.3 (3)	K2—O7B—C7B—O1B	-48.4 (4)
K4—O7—C7—C6	142.9 (2)	K3 ^{iv} —O7B—C7B—O1B	-144.0 (3)
K3 ^{iv} —O7—C7—C6	-117.1 (2)	K1—O7B—C7B—C6B	-145.2 (2)
K2—O7—C7—K4	-131.67 (18)	K2—O7B—C7B—C6B	132.8 (2)
K3 ^{iv} —O7—C7—K4	99.98 (18)	K3 ^{iv} —O7B—C7B—C6B	37.2 (3)
K4—O7—C7—K2	131.67 (18)	K2—O7B—C7B—K1	-82.00 (13)
K3 ^{iv} —O7—C7—K2	-128.3 (2)	K3 ^{iv} —O7B—C7B—K1	-177.60 (16)
Ni1—O1—C7—O7	178.1 (2)	Ni1B—O1B—C7B—O7B	173.5 (3)
K4—O1—C7—O7	28.2 (3)	K1—O1B—C7B—O7B	-30.0 (3)
Ni1—O1—C7—C6	1.8 (3)	Ni1B—O1B—C7B—C6B	-7.7 (3)
K4—O1—C7—C6	-148.1 (2)	K1—O1B—C7B—C6B	148.9 (2)

Ni1—O1—C7—K4	149.87 (15)	Ni1B—O1B—C7B—K1	-156.51 (16)
Ni1—O1—C7—K2	133.7 (7)	O6B—C6B—C7B—O7B	7.6 (5)
K4—O1—C7—K2	-16.1 (8)	N4B—C6B—C7B—O7B	-173.3 (3)
O6—C6—C7—O7	7.3 (4)	K1 ^{iv} —C6B—C7B—O7B	-44.8 (5)
N4—C6—C7—O7	-172.2 (3)	O6B—C6B—C7B—O1B	-171.3 (3)
K2—C6—C7—O7	-8.2 (2)	N4B—C6B—C7B—O1B	7.8 (4)
K3—C6—C7—O7	-55.2 (5)	K1 ^{iv} —C6B—C7B—O1B	136.3 (3)
O6—C6—C7—O1	-176.2 (3)	O6B—C6B—C7B—K1	-72.3 (5)
N4—C6—C7—O1	4.4 (4)	N4B—C6B—C7B—K1	106.7 (4)
K2—C6—C7—O1	168.3 (2)	K1 ^{iv} —C6B—C7B—K1	-124.7 (4)

Symmetry codes: (i) $x, y+1, z$; (ii) $-x, -y+1, -z$; (iii) $-x, -y, -z$; (iv) $x, y-1, z$; (v) $-x+1, -y+1, -z$; (vi) $-x+1, -y, -z$; (vii) $x, -y+1/2, z-1/2$; (viii) $x, -y+1/2, z+1/2$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O8—H8O \cdots O9 ^{viii}	0.85	2.02	2.869 (4)	173
O8—H8P \cdots O4B ^{ix}	0.85	2.01	2.858 (3)	166
O9—H9P \cdots O4 ^v	0.86	1.91	2.722 (3)	157
O9—H9O \cdots O6B ^{vii}	0.86	2.07	2.864 (3)	153
O10—H10P \cdots O4 ^x	0.88	2.02	2.887 (3)	168
O10—H10O \cdots O7B ⁱ	0.87	2.04	2.882 (3)	164

Symmetry codes: (i) $x, y+1, z$; (v) $-x+1, -y+1, -z$; (vii) $x, -y+1/2, z-1/2$; (viii) $x, -y+1/2, z+1/2$; (ix) $-x, y+1/2, -z+1/2$; (x) $-x+1, y+1/2, -z+1/2$.