

This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Dumitru, Adrian; Mäntysaari, Heikki; Paatelainen, Risto; Kaushik, Roy, Salazar, Farid; Schenke, Björn

Title: Azimuthal correlations in diffractive scattering at the Electron-Ion Collider

Year: 2022

Version: Published version

Copyright: © 2022 the Authors

Rights: CC BY 4.0

Rights url: <https://creativecommons.org/licenses/by/4.0/>

Please cite the original version:

Dumitru, A., Mäntysaari, H., Paatelainen, R., Kaushik, R., & Schenke, B. (2022). Azimuthal correlations in diffractive scattering at the Electron-Ion Collider. In DIS 2021 : Proceedings of the 28th International Workshop on Deep-Inelastic Scattering and Related Subjects (Article 102). Stichting SciPost. SciPost Physics Proceedings.
<https://doi.org/10.21468/SciPostPhysProc.8.102>

Azimuthal correlations in diffractive scattering at the Electron-Ion Collider

Adrian Dumitru^{1,2}, Heikki Mäntysaari^{3,4*}, Risto Paatelainen⁴,
Kaushik Roy^{5,6}, Farid Salazar^{6,7,8} and Björn Schenke⁷

1 Department of Natural Sciences, Baruch College, CUNY,
17 Lexington Avenue, New York, NY 10010, USA

2 The Graduate School and University Center, The City University of New York,
365 Fifth Avenue, New York, NY 10016, USA

3 Department of Physics, University of Jyväskylä,
P.O. Box 35, 40014 University of Jyväskylä, Finland

4 Helsinki Institute of Physics, P.O. Box 64, 00014 University of Helsinki, Finland

5 Max-Planck-Institut für Physik, Föhringer Ring 6, 80805 München, Germany

6 Department of Physics and Astronomy, Stony Brook University,
Stony Brook, NY 11794, USA

7 Physics Department, Brookhaven National Laboratory, Bldg. 510A, Upton, NY 11973, USA

8 Center for Frontiers in Nuclear Science (CFNS), Stony Brook University,
Stony Brook, NY 11794, USA

★ heikki.mantysaari@jyu.fi



*Proceedings for the XXVIII International Workshop
on Deep-Inelastic Scattering and Related Subjects,
Stony Brook University, New York, USA, 12-16 April 2021*
doi:[10.21468/SciPostPhysProc.8](https://doi.org/10.21468/SciPostPhysProc.8)

Abstract

We calculate azimuthal correlations between the exclusively produced vector meson and the scattered electron in Deep Inelastic Scattering processes at the future Electron-Ion Collider (EIC). We identify “kinematical” and “intrinsic” contributions to these correlations, and show that the correlations are sensitive to the non-trivial correlations in the gluon distribution of the target. Realistic predictions at the EIC kinematics are provided using two different approaches to describe the dipole-proton interaction at relatively small x .



Copyright A. Dumitru *et al.*

This work is licensed under the Creative Commons

[Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).

Published by the SciPost Foundation.

Received 24-05-2021

Accepted 23-03-2022

Published 13-07-2022

doi:[10.21468/SciPostPhysProc.8.102](https://doi.org/10.21468/SciPostPhysProc.8.102)



Check for
updates

1 Introduction

Exclusive vector particle production at high energy is a powerful probe of the small- x structure of protons and nuclei. This process has two particular advantages. First, as there is no

net color charge transfer to the target, the cross section scales approximately as the gluon density squared [1]. Additionally, as the momentum transfer is Fourier conjugate to the impact parameter, it becomes possible to study the spatial distribution of small- x gluons (and their event-by-event fluctuations) [2–4].

To perform more detailed imaging, it is useful to look at more differential measurements. In Ref. [5] presented in this Talk, exclusive J/ψ and photon production cross sections have been calculated differentially in both squared momentum transfer $|t|$ and the azimuthal angle $\phi_{k\Delta}$ between the outgoing electron and the produced vector particle in the frame where the incoming photon has no transverse momentum.

2 Azimuthal correlations in exclusive cross section

Cross section for exclusive vector particle production in electron-hadron collisions was calculated in Ref. [5]. The cross section can be written as

$$\begin{aligned} \frac{d\sigma^{eA \rightarrow eAV}}{dx_{\mathbb{P}} dQ^2 d|t| d\phi_{k\Delta}} &= \frac{\alpha_{em}}{32\pi^3 Q^2 x_{\mathbb{P}}} \sum_{\lambda'=0,\pm 1} \left\{ (1-y) \left| \langle \widetilde{\mathcal{M}}_{0,\lambda'}^{\gamma^*A,V} \rangle_Y \right|^2 \right. \\ &\quad \left. + \frac{1}{4} [1 + (1-y)^2] \sum_{\lambda=\pm 1} \left| \langle \widetilde{\mathcal{M}}_{\lambda,\lambda'}^{\gamma^*A,V} \rangle_Y \right|^2 \right. \\ &\quad \left. - \frac{\sqrt{2}}{2} (2-y) \sqrt{1-y} \sum_{\lambda=\pm 1} \text{Re} \left(\langle \widetilde{\mathcal{M}}_{0,\lambda'}^{\gamma^*A,V} \rangle_Y \langle \widetilde{\mathcal{M}}_{\lambda,\lambda'}^{\gamma^*A,V} \rangle_Y^* \right) \cos(\phi_{k\Delta}) \right. \\ &\quad \left. + (1-y) \text{Re} \left(\langle \widetilde{\mathcal{M}}_{+1,\lambda'}^{\gamma^*A,V} \rangle_Y \langle \widetilde{\mathcal{M}}_{-1,\lambda'}^{\gamma^*A,V} \rangle_Y^* \right) \cos(2\phi_{k\Delta}) \right\}. \quad (1) \end{aligned}$$

Here $\mathcal{M}_{\lambda,\lambda'}^{\gamma^*A,V}$ corresponds to the scattering amplitude for the $\gamma^* + A \rightarrow V + A$ scattering, where λ and λ' refer to the polarizations of the γ^* and V , respectively, $V = J/\psi$ or $V = \gamma$ and y is the inelasticity variable. Explicit expression for these amplitudes are calculated in Ref. [5] and the results are in agreement with the previous calculation performed in the same Color Glass Condensate (CGC) framework [6].

An important feature in Eq. (1) is that the interference between amplitudes with different polarization states results in non-zero $\cos(\phi_{k\Delta})$ and $\cos(2\phi_{k\Delta})$ modulations. These are in turn related to the dependence on orientation between the impact parameter \mathbf{b}_\perp and dipole size \mathbf{r}_\perp in the dipole-target scattering amplitude $\mathcal{D}(\mathbf{r}_\perp, \mathbf{b}_\perp)$. For example, the amplitude $\widetilde{\mathcal{M}}_{\pm 1, \mp 1}^{\gamma^*A, \gamma^*}$ reads

$$\begin{aligned} \widetilde{\mathcal{M}}_{\pm 1, \mp 1}^{\gamma^*A, \gamma^*} &= -\frac{8N_c e^2 q_f^2}{(2\pi)^2} \int_{\mathbf{b}_\perp} e^{-i\Delta_\perp \cdot \mathbf{b}_\perp} \int_{\mathbf{r}_\perp} e^{\pm 2i\phi_{r\Delta}} \mathcal{D}(\mathbf{r}_\perp, \mathbf{b}_\perp) \\ &\quad \times \int_z e^{-i\delta_\perp \cdot \mathbf{r}_\perp z \bar{z}} \varepsilon_f K_1(\varepsilon_f r_\perp) \varepsilon'_f K_1(\varepsilon'_f r_\perp), \quad (2) \end{aligned}$$

where $\varepsilon_f^2 = Q^2 z \bar{z} + m_f^2$ with $\bar{z} = 1 - z$. As $\phi_{r\Delta}$ is the azimuthal angle between the dipole orientation and the momentum transfer vectors, this amplitude is directly sensitive to the $\cos(2\phi_{rb})$ modulations in the dipole amplitude, where ϕ_{rb} is the angle between the impact parameter and the dipole orientation. This contribution to the azimuthal modulations in Eq. (1) we refer to as the intrinsic contribution. There is also an additional purely kinematical contribution from the non-zero off-forward phase factor $\delta_\perp = \frac{z-\bar{z}}{2} \Delta$.




Figure 1: The modulation coefficient v_n in coherent J/ψ production at $Q^2 = 2 \text{ GeV}^2$ from Ref. [5]. Left: dependence on the momentum fraction x_p . Right: v_2 calculated using the dipole amplitude from the impact parameter dependent MV model (standard setup) and from the LCPT calculation of Ref [12].

3 Numerical setup

The main results in Ref. [5] are obtained using a CGC setup, where the Wilson lines describing the target structure at initial $x = 0.01$ are obtained from the McLerran-Venugopalan model [7, 8] with impact parameter dependence. The non-uniform geometry is included by using a local Gaussian color charge correlator $g^2 \langle \rho(\mathbf{x}_\perp), \rho(\mathbf{y}_\perp) \rangle \sim g^4 \mu^2$, where the density $g^4 \mu^2$ depends on the impact parameter $\mathbf{b}_\perp = (\mathbf{x}_\perp + \mathbf{y}_\perp)/2$, and for the protons a Gaussian distribution $\sim e^{-b_\perp^2/(2B)}$ is assumed. To calculate cross sections at $x < 0.01$, this initial condition is coupled to a JIMWLK evolution equation [9].

The MV model with impact parameter dependence results in a dipole-proton scattering amplitude where the scattering probability is highest when the dipole is oriented parallel to the impact parameter (see [5, 10, 11]). For comparison, in this Contribution we also present preliminary results using a recent dipole-target scattering amplitude obtained as a result of an explicit light cone perturbation theory calculation in Refs. [12, 13] (“LCPT dipole”). In that approach, one takes a non-perturbative wave function constrained by low-energy data to describe the proton structure in the valence quark region, and then calculates perturbatively a gluon emission which allows extension of the calculation towards moderately small x . This calculation results in non-trivial correlations that are absent in the MV model, and for example the dipole-target scattering amplitude is largest when the dipole is perpendicular to the impact parameter.

4 Numerical results

In this Contribution we focus on exclusive J/ψ production in $e + p$ scattering, the results for DVCS are discussed in detail in Ref. [5] as well as $e+A$ collisions where the angular modulations are found to be very small. We show results for the modulation coefficients v_n defined as

$$v_n = \frac{\int_0^{2\pi} d\phi_{k\Delta} e^{ni\phi_{k\Delta}} d\sigma^{e+p \rightarrow e+p+V} / dt d\phi_{k\Delta} dQ^2 dx_p}{\int_0^{2\pi} d\phi_{k\Delta} d\sigma^{e+p \rightarrow e+p+V} / dt d\phi_{k\Delta} dQ^2 dx_p}. \quad (3)$$

The v_1 and v_2 modulations for the exclusive J/ψ production at two different $x_{\mathbb{P}}$ values accessible at the EIC are shown in Fig. 1a using the impact parameter dependent MV model initial condition coupled to the JIMWLK evolution. In case of heavy meson production, the v_1 modulation is found to be extremely small, as it is dominated by the off-forward phase factor whose contribution is small when the dipole size $|r_{\perp}^2| \sim 1/(Q^2 + M_{J/\psi}^2)$ is small. The v_2 modulation is significant, as it receives also a contribution from the $\cos(2\phi_{rb})$ modulation present in the dipole-proton scattering amplitude. The modulations are suppressed at smaller $x_{\mathbb{P}}$, and this change is driven by the JIMWLK evolution. The proton grows towards small $x_{\mathbb{P}}$, which results in smaller density gradients and weaker angular dependence in the dipole scattering amplitude.

Next we compare the v_2 modulation we obtain using the dipole amplitude from the MV model, and from the LCPT calculation discussed briefly in Sec. 3. We calculate v_2 at $x_{\mathbb{P}} = 0.01$, which is not too small and the LCPT calculation, which has not been combined with a small- x evolution equation, can be expected to be applicable. The results are shown in Fig. 1b. As discussed above, the angular dependence in the ‘‘LCPT dipole’’ differs from that of the MV model, which results in v_2 being negative (and the magnitude being smaller) than what is obtained when the (impact parameter dependent) MV model is used. For comparison, we also calculate v_2 using the LCPT dipole where we have artificially removed any dependence on the dipole orientation. In this case we find a small and positive v_2 , where the positive contribution is due to the non-zero off-forward phase $e^{-i\delta_{\perp} \cdot r_{\perp}}$ e.g. in Eq. (2).

5 Conclusion

We have shown in Ref. [5] how azimuthal electron-vector meson correlations can be used to access the angular dependence of the dipole-target scattering amplitude. This, in turn, can also be related to the gluon transversity GPD and to the gluon Wigner distribution [6, 10, 14]. From the Color Glass Condensate based setup, where an impact parameter dependent MV model is coupled to the JIMWLK small- x evolution equation, we predict significant ($\sim 5\%$) azimuthal modulations for exclusive J/ψ production in the electron-proton collisions at the EIC. Much larger modulations are predicted for DVCS, but the Bethe-Heitler contribution may render this measurements more difficult to interpret.

In this Contribution we have studied the model dependence of the predicted azimuthal modulations, by using a dipole amplitude recently obtained as a result of explicit light cone perturbation theory calculations [12] (also presented in this Conference). The dependence on the dipole orientation is very different in these calculations that approach the problem from the dilute valence quark region, compare to the MV model which consider the opposite (dense) limit. We find very different predictions for the v_2 modulations in exclusive J/ψ production, even with opposite signs when the dipole amplitude from the MV model or from the LCPT calculation is used. In the future, it will be interesting to study how the transition from a dilute (described by a LCPT calculation combined with a small- x evolution) to the dense region, where the MV model can be taken to be a more realistic description, takes place.

Funding information This work was supported by the Academy of Finland, projects 314764 (H.M) and 1322507 (R.P). H.M. is supported under the European Union’s Horizon 2020 research and innovation programme STRONG-2020 project (grant agreement no. 824093), and R.P. by the European Research Council grant agreement no. 725369. A.D. thanks the US Department of Energy, Office of Nuclear Physics, for support via Grant DE-SC0002307. F.S. and B.P.S. are supported under DOE Contract No. DE-SC0012704. F.S. and K.R. were also supported by the joint Brookhaven National Laboratory-Stony Brook University Center for

Frontiers in Nuclear Science (CFNS). The content of this article does not reflect the official opinion of the European Union and responsibility for the information and views expressed therein lies entirely with the authors. Computing resources from CSC – IT Center for Science in Espoo, Finland and from the Finnish Grid and Cloud Infrastructure (persistent identifier [urn:nbn:fi:research-infras-2016072533](https://nbn-resolving.org/urn:nbn:fi:research-infras-2016072533)) were used.

References

- [1] M. G. Ryskin, *Diffractional J/ψ electroproduction in LLA QCD*, Z. Phys. C **57**, 89 (1993), doi:[10.1007/BF01555742](https://doi.org/10.1007/BF01555742).
- [2] H. Mäntysaari and B. Schenke, *Evidence of Strong Proton Shape Fluctuations from Incoherent Diffraction*, Phys. Rev. Lett. **117**, 052301 (2016), doi:[10.1103/PhysRevLett.117.052301](https://doi.org/10.1103/PhysRevLett.117.052301).
- [3] H. Mäntysaari and B. Schenke, *Revealing proton shape fluctuations with incoherent diffraction at high energy*, Phys. Rev. D **94**, 034042 (2016), doi:[10.1103/PhysRevD.94.034042](https://doi.org/10.1103/PhysRevD.94.034042).
- [4] H. Mäntysaari, *Review of proton and nuclear shape fluctuations at high energy*, Rep. Prog. Phys. **83**, 082201 (2020), doi:[10.1088/1361-6633/aba347](https://doi.org/10.1088/1361-6633/aba347).
- [5] H. Mäntysaari, K. Roy, F. Salazar and B. Schenke, *Gluon imaging using azimuthal correlations in diffractive scattering at the Electron-Ion Collider*, Phys. Rev. D **103**, 094026 (2021), doi:[10.1103/PhysRevD.103.094026](https://doi.org/10.1103/PhysRevD.103.094026).
- [6] Y. Hatta, B.-W. Xiao and F. Yuan, *Gluon tomography from deeply virtual Compton scattering at small x* , Phys. Rev. D **95**, 114026 (2017), doi:[10.1103/PhysRevD.95.114026](https://doi.org/10.1103/PhysRevD.95.114026).
- [7] L. McLerran and R. Venugopalan, *Computing quark and gluon distribution functions for very large nuclei*, Phys. Rev. D **49**, 2233 (1994), doi:[10.1103/PhysRevD.49.2233](https://doi.org/10.1103/PhysRevD.49.2233).
- [8] L. McLerran and R. Venugopalan, *Gluon distribution functions for very large nuclei at small transverse momentum*, Phys. Rev. D **49**, 3352 (1994), doi:[10.1103/PhysRevD.49.3352](https://doi.org/10.1103/PhysRevD.49.3352).
- [9] A. H. Mueller, *A simple derivation of the JIMWLK equation*, Phys. Lett. B **523**, 243 (2001), doi:[10.1016/S0370-2693\(01\)01343-0](https://doi.org/10.1016/S0370-2693(01)01343-0).
- [10] H. Mäntysaari, N. Mueller, F. Salazar and B. Schenke, *Multigluon Correlations and Evidence of Saturation from Dijet Measurements at an Electron-Ion Collider*, Phys. Rev. Lett. **124**, 112301 (2020), doi:[10.1103/PhysRevLett.124.112301](https://doi.org/10.1103/PhysRevLett.124.112301).
- [11] E. Iancu and A. H. Rezaeian, *Elliptic flow from color-dipole orientation in pp and pA collisions*, Phys. Rev. D **95**, 094003 (2017), doi:[10.1103/PhysRevD.95.094003](https://doi.org/10.1103/PhysRevD.95.094003).
- [12] A. Dumitru, H. Mäntysaari and R. Paatelainen, *Color charge correlations in the proton at NLO: Beyond geometry based intuition*, Phys. Lett. B **820**, 136560 (2021), doi:[10.1016/j.physletb.2021.136560](https://doi.org/10.1016/j.physletb.2021.136560).
- [13] A. Dumitru and R. Paatelainen, *Sub-femtometer scale color charge fluctuations in a proton made of three quarks and a gluon*, Phys. Rev. D **103**, 034026 (2021), doi:[10.1103/PhysRevD.103.034026](https://doi.org/10.1103/PhysRevD.103.034026).
- [14] Y. Hatta, B.-W. Xiao and F. Yuan, *Probing the Small- x Gluon Tomography in Correlated Hard Diffractive Dijet Production in Deep Inelastic Scattering*, Phys. Rev. Lett. **116**, 202301 (2016), doi:[10.1103/PhysRevLett.116.202301](https://doi.org/10.1103/PhysRevLett.116.202301).