

This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Krupchyk, Katya; Liimatainen, Tony; Salo, Mikko

Title: Linearized Calderón problem and exponentially accurate quasimodes for analytic manifolds

Year: 2022

Version: Published version

Copyright: © 2022 the Authors

Rights: CC BY 4.0

Rights url: <https://creativecommons.org/licenses/by/4.0/>

Please cite the original version:

Krupchyk, K., Liimatainen, T., & Salo, M. (2022). Linearized Calderón problem and exponentially accurate quasimodes for analytic manifolds. *Advances in Mathematics*, 403, Article 108362. <https://doi.org/10.1016/j.aim.2022.108362>




ELSEVIER

Contents lists available at ScienceDirect

Advances in Mathematics

www.elsevier.com/locate/aim



Linearized Calderón problem and exponentially accurate quasimodes for analytic manifolds



Katya Krupchyk^a, Tony Liimatainen^{b,*}, Mikko Salo^c

^a Department of Mathematics, University of California, Irvine, CA 92697-3875, USA

^b Department of Mathematics and Statistics, University of Helsinki, Helsinki, PL 68, FI-00014, Finland

^c Department of Mathematics and Statistics, University of Jyväskylä, Jyväskylä, FI-40014, Finland

ARTICLE INFO

Article history:

Received 2 October 2020

Received in revised form 7 March 2022

Accepted 7 March 2022

Available online 5 April 2022

Communicated by C. Fefferman

MSC:

primary 35R30

secondary 35A18, 35A20, 35J25

Keywords:

Inverse problems

Riemannian manifold

Conformally transversally

anisotropic

Gaussian quasimodes

WKB construction

Wave front set

ABSTRACT

In this article we study the linearized anisotropic Calderón problem on a compact Riemannian manifold with boundary. This problem amounts to showing that products of pairs of harmonic functions of the manifold form a complete set. We assume that the manifold is transversally anisotropic and that the transversal manifold is real analytic and satisfies a geometric condition related to the geometry of pairs of intersecting geodesics. In this case, we solve the linearized anisotropic Calderón problem. The geometric condition does not involve the injectivity of the geodesic X-ray transform. Crucial ingredients in the proof of our result are the construction of Gaussian beam quasimodes on the transversal manifold, with exponentially small errors, as well as the FBI transform characterization of the analytic wave front set.

© 2022 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

* Corresponding author.

E-mail addresses: katya.krupchyk@uci.edu (K. Krupchyk), tony.liimatainen@helsinki.fi (T. Liimatainen), mikko.j.salo@jyu.fi (M. Salo).

1. Introduction and statement of results

The inverse conductivity problem posed by Calderón [5] asks to determine the electrical conductivity of a medium from voltage and current measurements on its boundary. This problem is the mathematical model of Electrical Impedance (or Resistivity) Tomography, an imaging method with applications in seismic and medical imaging. It is also one of the most fundamental models of inverse boundary value problems for elliptic partial differential equations. For these reasons both the theoretical and applied aspects of the Calderón problem have been under intense study. We refer to the survey [53] for more information and references.

In this article we are interested in the case where the electrical conductivity of the medium is *anisotropic*, i.e. depends on direction. This can be modeled by a matrix conductivity coefficient, or in geometric terms by having a resistivity coefficient given by a Riemannian metric g on a compact manifold M with smooth boundary. There are many variants of this problem. One of them is the (geometric) Calderón problem for a Schrödinger equation: given a known compact Riemannian manifold (M, g) with smooth boundary and an unknown potential $q \in C^\infty(M)$, determine q from the knowledge of the Cauchy data on ∂M of solutions of the Schrödinger equation

$$(-\Delta_g + q)u = 0 \text{ in } M.$$

Here $-\Delta_g$ is the Laplace-Beltrami operator. This geometric Calderón problem is solved in [21] when $\dim(M) = 2$. The problem is open in general when $\dim(M) \geq 3$ with only partial results available. In particular, the unique determination of q was obtained in [52] in the Euclidean setting, in [27] for hyperbolic manifolds, and in [40], [29] in the real analytic setting. Going beyond these settings, the geometric Calderón problem was only solved in the case when (M, g) is CTA (conformally transversally anisotropic, see Definition 1.1 below) and under the assumption that the geodesic X-ray transform on the transversal manifold is injective [10,12].

The linearized version (at $q = 0$) of the above problem is also of interest, since methods for the linearized problem often give insight to the original problem. In our case, the linearized problem reduces to the following simple question asking whether products of pairs of harmonic functions form a complete set in $L^1(M)$:

Question 1. Let (M, g) be a compact oriented Riemannian manifold with smooth boundary. If $f \in L^\infty(M)$ satisfies

$$\int_M f u_1 u_2 dV_g = 0$$

for all $u_j \in L^2(M)$ with $\Delta_g u_j = 0$ in M , $j = 1, 2$, is it true that $f \equiv 0$?

The methods of [21,10,12] give a positive answer to Question 1 when $\dim(M) = 2$, or when $\dim(M) \geq 3$ and (M, g) is CTA with the transversal manifold having injective geodesic X-ray transform. There have been recent attempts to improve these results when $\dim(M) \geq 3$. In [20], it is proved that Question 1 has a positive answer when (M, g) is a complex Kähler manifold with sufficiently many holomorphic functions. The article [13] establishes a recovery of singularities result: if (M, g) is transversally anisotropic and the transversal manifold satisfies a certain geometric condition, one can recover transversal singularities of f . See also [11], [50] for the linearized Calderón problem with partial data in the Euclidean setting.

The recent related works [14,15,31,32,34,41,47] regarded inverse problems for nonlinear elliptic equations on CTA manifolds as well as some complex manifolds, equipped with a Kähler metric. The linearized problem in Question 1 plays an important role in these results. Moreover, it was proven in [34] that on a general transversally anisotropic manifold, the products of four (instead of pairs of) harmonic functions form a complete set in $L^1(M)$, and this result was used crucially to solve the corresponding inverse problems for semilinear Schrödinger equations. We also mention the related works [22,36], which solve inverse problems for nonlinear wave equations by showing that sets of products of four waves are dense.

In this article we extend the result of [13] and show that if the transversal manifold is additionally real-analytic, Question 1 has a positive answer (i.e. one can recover $f \in L^\infty(M)$ completely, not just some of its singularities).

Let us proceed to state our results. To that end, let us first recall the following definitions, see [10], [12].

Definition 1.1. Let (M, g) be a smooth compact oriented Riemannian manifold of dimension $n \geq 3$ with smooth boundary ∂M .

- (i) (M, g) is called transversally anisotropic if $(M, g) \subset\subset (T, g)$ where $T = \mathbb{R} \times M_0^{\text{int}}$, $g = e \oplus g_0$, (\mathbb{R}, e) is the Euclidean real line, and (M_0, g_0) is a smooth compact $(n-1)$ -dimensional manifold with smooth boundary, called the transversal manifold.
- (ii) (M, g) is called conformally transversally anisotropic (CTA) if (M, cg) is transversally anisotropic, for some positive function $c \in C^\infty(M)$.

Here and in what follows $M_0^{\text{int}} = M_0 \setminus \partial M_0$ stands for the interior of M_0 . By choosing local coordinates x' for M_0 and denoting by x_1 the coordinate on \mathbb{R} , the metric $g = e \oplus g_0$ has the form

$$dx_1^2 + g_{\alpha\beta}(x')dx'^\alpha dx'^\beta,$$

where the Einstein summation convention is used for $\alpha, \beta = 1, \dots, n-1$.

In this work we require that all manifolds are oriented. We mention the recent work [4], which studies the Calderón problem on non-oriented Riemannian surfaces.

Let (M, g) be transversally anisotropic of dimension $n \geq 3$ with a transversal manifold (M_0, g_0) . Next we need some definitions related to the transversal manifold (M_0, g_0) . Following [12], we say that a geodesic $\gamma : [-T_1, T_2] \rightarrow M_0$, $0 < T_1, T_2 < \infty$, is nontangential if $\gamma(-T_1), \gamma(T_2) \in \partial M_0$, $\gamma(t) \in M_0^{\text{int}}$ for all $-T_1 < t < T_2$, and $\dot{\gamma}(-T_1), \dot{\gamma}(T_2)$ are nontangential vectors on ∂M_0 . Following [13], we have the following definition.

Definition 1.2. We say that $(x'_0, \xi'_0) \in S^*M_0^{\text{int}}$ is generated by an admissible pair of geodesics, if there are two nontangential unit speed geodesics

$$\gamma_1 : [-T_1, T_2] \rightarrow M_0, \quad \gamma_2 : [-S_1, S_2] \rightarrow M_0,$$

$0 < T_1, T_2, S_1, S_2 < \infty$, such that

- (i) $\gamma_1(0) = \gamma_2(0) = x'_0$,
- (ii) $\dot{\gamma}_1(0) + \dot{\gamma}_2(0) = t_0 \xi'_0$, for some $0 < t_0 < 2$, where ξ'_0 is understood as an element of $T_{x_0}M_0^{\text{int}}$ by the Riemannian duality,
- (iii) γ_1, γ_2 do not have self-intersections at the point x'_0 , and x'_0 is the only point of their intersections, i.e.

$$\begin{aligned} \gamma_1(t) = x'_0 &\Leftrightarrow t = 0, & \gamma_2(s) = x'_0 &\Leftrightarrow s = 0, \\ \gamma_1(t) = \gamma_2(s) &\Rightarrow \gamma_1(t) = \gamma_2(s) = x'_0. \end{aligned}$$

Let $f \in L^\infty(M)$ and let us extend $f \in L^\infty(M)$ by zero to $(\mathbb{R} \times M_0) \setminus M$. Writing $x = (x_1, x')$ where $x_1 \in \mathbb{R}$, and x' are local coordinates M_0 , we let

$$\widehat{f}(\lambda, x') = \int_{-\infty}^{\infty} e^{-i\lambda x_1} f(x_1, x') dx_1, \quad \lambda \in \mathbb{R}, \quad (1.1)$$

be the *partial Fourier transform* of f with respect to x_1 . We have for each $\lambda \in \mathbb{R}$ that $\widehat{f}(\lambda, \cdot) \in L^\infty(M_0) \cap \mathcal{E}'(M_0^{\text{int}})$.

When X is a real analytic open manifold and $u \in \mathcal{D}'(X)$, we let $WF_a(u) \subset T^*X \setminus \{0\}$ stand for the analytic wave front set of u , see [49, Definition 6.1], [25, Sections 8.5, 9.3]. The set $WF_a(u) \subset T^*X \setminus \{0\}$ is closed conic and we have

$$\pi(WF_a(u)) = \text{singsupp}_a(u),$$

where $\pi : T^*X \rightarrow X$, $(x, \xi) \mapsto x$, is the natural projection and $\text{singsupp}_a(u)$ is the analytic singular support of u , i.e. the smallest closed set such that u is real analytic in the complement [25, Theorem 8.4.5]. In particular, $WF_a(u) = \emptyset$ if and only if u is real analytic on X .

We have the following analytic microlocal result, which is an analog of Theorem 1.1 in [13], established in the C^∞ -case.

Theorem 1.3. *Let (M, g) be a transversally anisotropic manifold of dimension $n \geq 3$ with transversal manifold (M_0, g_0) , and assume that M_0^{int} and $g_0|_{M_0^{\text{int}}}$ are real analytic. Assume furthermore that $f \in L^\infty(M)$ satisfies*

$$\int_M f u_1 u_2 dV_g = 0, \tag{1.2}$$

for all $u_j \in L^2(M)$ with $-\Delta_g u_j = 0$ in M^{int} . Let $(x'_0, \xi'_0) \in S^*M_0^{\text{int}}$ be generated by an admissible pair of geodesics. Then for any $\lambda \in \mathbb{R}$, one has

$$(x'_0, \xi'_0) \notin WF_a(\widehat{f}(\lambda, \cdot)) \subset T^*M_0^{\text{int}} \setminus \{0\}.$$

Here $\widehat{f}(\lambda, \cdot)$ refers to the partial Fourier transformation given by (1.1).

Theorem 1.3 implies the following global result, which gives a positive answer to Question 1 under suitable geometric assumptions.

Theorem 1.4. *Let (M, g) be a transversally anisotropic manifold of dimension $n \geq 3$ and assume that the transversal manifold (M_0, g_0) is connected, M_0^{int} as well as g_0 in M_0^{int} are real analytic. Assume that every point $(x'_0, \xi'_0) \in S^*M_0^{\text{int}}$ is generated by an admissible pair of geodesics. Moreover, assume that $f \in L^\infty(M)$ satisfies (1.2) for all $u_j \in L^2(M)$ with $-\Delta_g u_j = 0$ in M^{int} . Then $f = 0$ in M .*

Remark 1.5. Note that while (M_0^{int}, g_0) is real analytic, Theorem 1.4 does not follow from the existing results in the real analytic setting, as it corresponds to deforming the zero potential by an L^∞ perturbation.

Remark 1.6. In Theorems 1.3 and 1.4 while M^{int} is real analytic, the boundary ∂M need not be real analytic.

As the following example shows, there exist transversally anisotropic manifolds (M, g) with a transversal manifold (M_0, g_0) satisfying the geometric conditions of Theorem 1.4 and with a non-invertible geodesic X-ray transform. Therefore, the geometric Calderón problem is still open on such manifolds while our Theorem 1.4 gives a positive solution to the corresponding linearized problem.

Example 1.7. Let $M_0 = \mathbb{S}^1 \times [0, a]$, $a > 0$, be a cylinder with its usual flat metric g_0 . The geodesics on M_0 are straight lines, circular cross sections, and helices that wind around the cylinder. The geodesic X-ray transform is not invertible, since the kernel contains functions of the form $f(e^{it}, s) = h(s)$ where $h \in C^\infty((0, a))$ integrates to zero over $[0, a]$. However, it is shown in Appendix A that every point $(x'_0, \xi'_0) \in S^*M_0^{\text{int}}$ is generated by an admissible pair of geodesics.

It is established in [13, Lemma 3.1] that if (M_0, g_0) satisfies the strict Stefanov–Uhlmann regularity condition at $(x'_0, \xi'_0) \in S^*M_0^{\text{int}}$, which we now proceed to recall, then (x'_0, ξ'_0) is generated by an admissible pair of geodesics.

Definition 1.8. The transversal manifold (M_0, g_0) satisfies the strict Stefanov–Uhlmann regularity condition at $(x'_0, \xi'_0) \in S^*M_0^{\text{int}}$ if there exists $\eta' \in S^*_{x'_0}M_0^{\text{int}}$ such that $g_0(\xi'_0, \eta') = 0$ and such that the following holds: let $\gamma_{x'_0, \eta'} : [-T_1, T_2] \rightarrow M_0$, $0 < T_1, T_2 < \infty$, be the geodesic with $\gamma_{x'_0, \eta'}(0) = x'_0$, $\dot{\gamma}_{x'_0, \eta'} = \eta'$. We have

- (i) $\gamma_{x'_0, \eta'}$ is nontangential,
- (ii) $\gamma_{x'_0, \eta'}$ contains no points conjugate to x'_0 ,
- (iii) $\gamma_{x'_0, \eta'}$ does not self-intersect for any time $t \in [-T_1, T_2]$.

Hence, if a transversally anisotropic manifold (M, g) is such that the transversal manifold (M_0, g_0) satisfies the strict Stefanov–Uhlmann regularity condition at every point of $S^*M_0^{\text{int}}$ with M_0^{int} and $g_0|_{M_0^{\text{int}}}$ real analytic, and (M_0, g_0) is connected, then Theorem 1.4 holds.

As the following examples demonstrate, there are transversally anisotropic manifolds (M, g) with a transversal manifold (M_0, g_0) satisfying the geometric condition of Theorem 1.4, and with an invertible geodesic X-ray transform. Thus, for such manifolds (M, g) , Theorem 1.4 also follows from [10], [12].

Example 1.9. Let (M_0, g_0) be a *simple* manifold, i.e. a compact simply connected manifold with strictly convex boundary so that no geodesic has conjugate points. Then (M_0, g_0) satisfies the strict Stefanov–Uhlmann regularity condition at any point of $S^*M_0^{\text{int}}$ and thus also the geometric condition in Theorem 1.4. Note that in this case (M, g) is admissible in the sense of [10], and Theorem 1.4 would also follow from [10].

Example 1.10. Let $\mathbb{S}^3 \subset \mathbb{R}^4$ be the unit sphere and let μ be a geodesic arc from the north pole to the south pole of the sphere. Let M_0 be the closure of a neighborhood of μ . It is established in [13] that the manifold M_0 satisfies the strict Stefanov–Uhlmann regularity condition at each point of $S^*M_0^{\text{int}}$. Notice also that the manifold M_0 contains conjugate points, so that it is not simple. However, the geodesic X-ray transform on (M_0, g_0) is injective by [51], and Theorem 1.4 would therefore also follow from [12].

Remark 1.11. We would like to remark that the strict Stefanov–Uhlmann condition is not satisfied for (M_0, g_0) of Example 1.7 since for any $(x'_0, \xi'_0) \in S^*M_0^{\text{int}}$ with ξ'_0 pointing in the direction of the $[0, a]$ factor, the orthogonal geodesics never reach ∂M_0 .

The proof of Theorem 1.3 depends crucially on the construction of Gaussian beam quasimodes along nontangential geodesics on M_0 , with exponentially small errors, as stated in the following result. Before stating the result, let us recall from [49, Chapter 1]

the notion of a classical analytic symbol. Let $V \subset \mathbb{C}^n$ be an open set. We say that $a(x; h) = \sum_{k=0}^{\infty} h^k a_k(x)$ is a (formal) classical analytic symbol in V if $a_k \in \text{Hol}(V)$, $k = 0, 1, 2, \dots$, and for every $\tilde{V} \subset\subset V$, there exists $C = C_{\tilde{V}} > 0$ such that

$$|a_k(x)| \leq C^{k+1} k^k, \quad x \in \tilde{V}, \tag{1.3}$$

$k = 0, 1, 2, \dots$. The classical analytic symbol $a(x; h)$ is said to be elliptic if $a_0 \neq 0$.

We have the following essentially well known result, see [48] and [49], and see also the work by Babich [1] for a sketch of the proof. Notice that here our quasimode construction is performed along the entire geodesic segment contrary to the standard constructions in a neighborhood of a point, see [9]. We use the notation

$$\text{neigh}(p, X)$$

for an open neighborhood of a point $p \in X$, and similarly for neighborhoods of general subsets of other topological spaces in place of p and X above.

Theorem 1.12. *Let (X, g) be a compact Riemannian manifold of dimension $n \geq 2$ with smooth boundary, contained in a real analytic open manifold (\hat{X}, g) of the same dimension with g real analytic in \hat{X} . Let $\gamma : [-T_1, T_2] \rightarrow X$, $0 < T_1, T_2 < \infty$, be a unit speed non-tangential geodesic in X , and let $\lambda \in \mathbb{R}$. For any neighborhood of $\gamma([-T_1, T_2])$, there is a family of C^∞ functions $v(x; h)$ on X , $0 < h \leq 1$, supported in the neighborhood, and $C > 0$ such that*

$$\|(-h^2 \Delta_g - (hs)^2)v\|_{L^2(X)} = \mathcal{O}(e^{-\frac{1}{Ch}}), \quad \|v\|_{L^2(X)} \asymp 1, \tag{1.4}$$

as $h \rightarrow 0$. Here $s = \frac{1}{h} + i\lambda$. The local structure of the family $v(x; h)$ is as follows: let $p \in \gamma([-T_1, T_2])$ and let $t_1 < \dots < t_{N_p}$ be the times in $(-T_1, T_2)$ when $\gamma(t_l) = p$, $l = 1, \dots, N_p$. In a sufficiently small neighborhood V of a point $p \in \gamma([-T_1, T_2])$, we have

$$v|_V = v^{(1)} + \dots + v^{(N_p)},$$

where each $v^{(l)}$ has the form

$$v^{(l)}(x; h) = h^{-\frac{(n-1)}{4}} e^{is\varphi^{(l)}(x)} a^{(l)}(x; h).$$

Here $\varphi = \varphi^{(l)}$ is real analytic in V satisfying for t near t_l ,

$$\varphi(\gamma(t)) = t, \quad \nabla\varphi(\gamma(t)) = \dot{\gamma}(t), \quad \text{Im}(\nabla^2\varphi(\gamma(t))) \geq 0, \quad \text{Im}(\nabla^2\varphi)|_{\dot{\gamma}(t)^\perp} > 0, \tag{1.5}$$

and $a^{(l)}$ is an elliptic classical analytic symbol in a complex neighborhood of p .

We have chosen to give a fairly complete proof of Theorem 1.12 since we are not aware of a detailed treatment in the literature and since we need to have fairly precise information concerning the quasimodes for our applications. We refer to [33] for a related complex Riccati equation, and to [6] for a geometric interpretation of it.

Let us briefly mention how the exponentially small error is achieved in Theorem 1.12. The proof of the theorem is by using the ansatz $v(x; h) = e^{is\varphi(x)}a(x; h)$, which, as usual, leads to solving the eikonal equation for the phase function $\varphi(x)$ and a transport equation for the amplitude $a(x; h)$. We first find an exact analytic solution for the eikonal equation near a geodesic segment of γ . Consequently, the transport equation for the amplitude $a(x; h) = \sum_{k=0}^N h^k a_k(x)$ has analytic coefficients and we find $a(x; h)$ as a classical analytic symbol. This involves adapting the nested neighborhood method of [49]. The error for v being a true eigenfunction then is

$$(-h^2 \Delta_g - (hs)^2)e^{is\varphi} \left(\sum_{j=0}^N h^j a_j \right) = h^{N+2} T_2(a_N), \tag{1.6}$$

where T_2 is a second order operator with analytic coefficients. Cauchy estimates and (1.3) then yield that the error term (1.6) is bounded by $h^{N+2} C^{N+1} N^N$. Letting the order N of the expansions of a depend on h as $N = N(h) = \lceil \frac{1}{h\epsilon C} \rceil$ gives the exponentially small error in the theorem.

The above was based on finding first an exact analytic solution to the eikonal equation $|d\varphi|_g = 1$ near a geodesic segment of γ . To find such a solution, we view the eikonal equation as the Hamilton-Jacobi equation,

$$p(x, \varphi'_x(x)) = 0, \tag{1.7}$$

where $p(x, \xi) = |\xi|_{g(x)}^2 - 1$ is holomorphically continued to a complex domain. When solving the Hamilton-Jacobi equation (1.7) we proceed by a geometric argument of constructing a complex Lagrangian manifold,

$$\Lambda \subset p^{-1}(0),$$

in a complex neighborhood of a segment of the graph of $\hat{\gamma} \subset T^*X$, see [48]. The solution φ is then obtained as a generating function of the Lagrangian Λ , which parametrizes Λ as

$$\Lambda = \{(x, \varphi'_x(x))\}.$$

Extending the argument to a neighborhood of the geodesic segment of γ requires some extra work involving positive Lagrangians.

Let us proceed to explain the main ideas in the proof of Theorem 1.3. Let $\alpha_0 = (x'_0, \xi'_0) \in S^*M_0^{\text{int}}$ be generated by an admissible pair of geodesics $\gamma_1(\alpha_0)$ and $\gamma_2(\alpha_0)$ on

M_0 . We first show that there exists a neighborhood of α_0 in $S^*M_0^{\text{int}}$ such that every point α in the neighborhood is generated by an admissible pair of geodesics $\gamma_1(\alpha)$ and $\gamma_2(\alpha)$ on M_0 . Next we construct two real analytic families of Gaussian beams quasimodes $v_1(\alpha)$ and $v_2(\alpha)$ on M_0 , associated to $\gamma_1(\alpha)$ and $\gamma_2(\alpha)$, respectively, with exponentially small errors. The fact that (M, g) is transversally anisotropic provides us with the limiting Carleman weight $\phi(x) = x_1$ for the Laplacian, and using the technique of Carleman estimates, we convert the families of Gaussian beams $v_1(\alpha)$ and $v_2(\alpha)$ into two families of harmonic functions on M with exponentially small remainder terms. Testing the orthogonality relation (1.2) with the constructed families of harmonic functions leads to the exponential decay of the FBI transform of $\widehat{f}(\lambda, \cdot)$ in a neighborhood of α_0 . Using the FBI characterization of the analytic wave front set, see [49], we conclude the proof. Note that we need to work with families of Gaussian beams to fill out the entire neighborhood of α_0 .

Remark 1.13. Similarly to [13], Theorem 1.3 and Theorem 1.4 are established for transversally anisotropic manifolds rather than CTA manifolds. The reason for this is that the standard reduction

$$c^{\frac{n+2}{4}} \circ (-\Delta_{cg}) \circ c^{-\frac{(n-2)}{4}} = -\Delta_g + q, \quad g = e \oplus g_0,$$

leads to the potential

$$q = -c^{\frac{n+2}{4}} \Delta_g (c^{-\frac{(n-2)}{4}}),$$

see [12], and therefore, to construct harmonic functions with exponentially small remainder terms on a CTA manifold, one has to construct Gaussian beam quasimodes for the conjugated Schrödinger operator,

$$e^{sx_1} (-h^2 \Delta_g + h^2 q) e^{-sx_1},$$

with exponentially small errors. If c is independent of x_1 and real-analytic then so is q , and this construction could be done as in Theorem 1.12. Notice also that for this reason, one can also include a general real analytic potential which is independent of x_1 in the results of Theorem 1.12. However, if c depends on x_1 then the corresponding sequence of transport equations becomes of $\bar{\partial}$ -type, see e.g. [14], [30], which complicates the analysis of Theorem 1.12 further and is therefore not developed here.

The proof of Theorem 1.12 given in Section 2 below shows also that the following result holds.

Corollary 1.14. *Let us make the same assumptions as in Theorem 1.12 and let q be real analytic on X . Then there is a family of C^∞ functions $v_q(x; h)$ on X such that*

$$\|(-h^2(\Delta_g + q) - (hs)^2)v_q\|_{L^2(X)} = \mathcal{O}(e^{-\frac{1}{ch}}), \quad \|v_q\|_{L^2(X)} \asymp 1,$$

as $h \rightarrow 0$. Here $v_q(x; h)$ satisfies the same properties as does $v(x; h)$ in Theorem 1.12.

Let us mention that Gaussian beam quasimode constructions have a long tradition in microlocal analysis, see [2], [44], [45], with applications in the analysis of eigenfunctions, see [54], and inverse problems, see [43] and the references given there.

Finally, let us point out certain related results on a standard geometric version of the Calderón problem, which asks to determine a metric g up to natural gauges (a boundary-fixing diffeomorphism, and also a conformal factor when $\dim(M) = 2$) from the knowledge of Cauchy data on ∂M of solutions of the equation $-\Delta_g u = 0$ in M . This problem was solved in [38] when $\dim(M) = 2$, but for $\dim(M) \geq 3$ it is only known under additional conditions such as the manifold being real-analytic, see [40,38,37], or Einstein [19]. Alternative proofs are given in [3,35]. Interesting counterexamples in the case of measurements on disjoint sets or low regularity coefficients are given in [7,8]. If one allows degenerate coefficients, there are other counterexamples [37,17]. Counterexamples with degenerate coefficients form the basis of invisibility cloaking, see e.g. [53].

The paper is organized as follows. Section 2 is devoted to the construction of exponentially accurate Gaussian beam quasimodes and the proof of Theorem 1.12. Section 3 contains some results concerning properties of geodesics needed in the proof of Theorem 1.3. Section 4 extends Theorem 1.12 to produce analytic families of exponentially accurate Gaussian beam quasimodes. The construction of families of harmonic functions based on Gaussian beam quasimodes is presented in Section 5. Section 6 contains some facts about analytic wave front sets and the proofs of Theorem 1.3 and Theorem 1.4. The admissibility property of geodesics in Example 1.7 is verified in Appendix A.

Acknowledgments

The research of K.K. is partially supported by the National Science Foundation (DMS 1815922 and DMS 2109199). T.L. and M.S. were supported by the Academy of Finland (Finnish Centre of Excellence in Inverse Modelling and Imaging, grant numbers 312121 and 309963), and M.S. was also supported by the European Research Council under Horizon 2020 (ERC CoG 770924). This material is based upon work supported by the National Science Foundation under Grant No. 1440140, while K.K. and M.S. were in residence at MSRI in Berkeley, California, during the semester on Microlocal Analysis in 2019. K.K. is very grateful to Johannes Sjöstrand for a very helpful discussion and for pointing out the reference [48]. The authors would like to thank the anonymous referee for a careful reading and helpful suggestions that have improved the presentation.

2. Exponentially accurate quasimodes. Proof of Theorem 1.12

Let (X, g) be a compact Riemannian manifold of dimension $n \geq 2$ with smooth boundary, contained in a larger real analytic open manifold (\widehat{X}, g) of the same dimension

with g real analytic in \widehat{X} . We extend γ as a unit speed geodesic in \widehat{X} . Let $\varepsilon > 0$ be such that $\gamma(t) \in \widehat{X} \setminus X$ and $\gamma(t)$ has no self-intersection for $t \in [-T_1 - 2\varepsilon, -T_1] \cup (T_2, T_2 + 2\varepsilon]$. This choice of ε is possible since γ is non-tangential. First it follows from [28, Lemma 7.2] that $\gamma|_{[-T_1 - \varepsilon, T_2 + \varepsilon]}$ self-intersects only at finitely many times t_j with

$$-T_1 < t_1 < \dots < t_N < T_2.$$

We also set $t_0 := -T_1 - \varepsilon$ and $t_{N+1} := T_2 + \varepsilon$. An application of [12, Lemma 3.5] shows that there exists an open cover $\{(U_j, \kappa_j)\}_{j=0}^{N+1}$ of $\gamma([-T_1 - \varepsilon, T_2 + \varepsilon])$ consisting of coordinate neighborhoods U_j and real analytic diffeomorphisms κ_j having the following properties:

- (i) $\kappa_j(U_j) = I_j \times B$, where I_j are open intervals and $B = B(0, \delta')$ is an open ball in \mathbb{R}^{n-1} . Here $\delta' > 0$ can be taken arbitrarily small and the same for each U_j ,
- (ii) $\kappa_j(\gamma(t)) = (t, 0)$ for each $t \in I_j$,
- (iii) t_j only belongs to I_j and $\overline{I_j} \cap \overline{I_k} = \emptyset$ unless $|j - k| \leq 1$,
- (iv) $\kappa_j = \kappa_k$ on $\kappa_j^{-1}((I_j \cap I_k) \times B)$.

The corresponding local coordinates $\kappa_j(x) = (t, y) \in U_j$ are called the Fermi coordinates. Here we note that Lemma 3.5 in [12] is established in the C^∞ case, and the real analyticity of the Fermi diffeomorphisms κ_j is obtained by inspection of the proof of Lemma 3.5 in [12], in view of the analyticity of \widehat{X} . As observed in the proof of [12, Lemma 3.5], in the case when γ does not self-intersect, there are Fermi coordinates on a single coordinate neighborhood of $\gamma|_{[-T_1 - \varepsilon, T_2 + \varepsilon]}$ so that (i) and (ii) are satisfied. These coordinates are given by inverting the map

$$(t, y) \mapsto \exp_{\gamma(t)}\left(\sum_{k=1}^{n-1} y^k e_k(t)\right) \in \widehat{X}.$$

Here $e_k(t)$ are the parallel transportations of the last $n - 1$ vectors of an orthonormal frame $\{\dot{\gamma}|_{t=-T_1}, e_1, \dots, e_{n-1}\} \subset T_{\gamma(-T_1)}M$ and \exp is the exponential map of (\widehat{X}, g) .

Our goal is to construct exponentially accurate Gaussian beam quasimodes near $\gamma([-T_1 - \varepsilon, T_2 + \varepsilon])$. We shall start by carrying out the quasimode construction in a fixed coordinate neighborhood $U = U_j$ which we can identify with the set $I \times B$, where $I \subset \mathbb{R}$ is an open interval and $B = B(0, \delta')$ is an open ball in \mathbb{R}^{n-1} with $\delta' > 0$. Without loss of generality, we assume that $0 \in I$. The geodesic γ in the open set U is given by $\Gamma = \{x = (t, y) \in I \times B : y = 0\}$.

Let us consider the following Gaussian beam ansatz,

$$v(t, y; h) = e^{is\varphi(t,y)} a(t, y; h), \quad s = \frac{1}{h} + i\lambda, \quad \lambda \in \mathbb{R}, \tag{2.1}$$

where the phase φ is complex valued with $\text{Im } \varphi(t, y) \geq 0$ and a is an amplitude. We shall proceed to construct the quasimode v so that the phase φ satisfies (1.5) and the amplitude a is an elliptic classical analytic symbol.

2.1. Construction of the phase function φ

We shall proceed using the classical arguments, solving the Hamilton-Jacobi equation in the complex domain and making crucial use of the geometry of positive complex Lagrangians, see [48]. Let us remark here that while we only need the good properties of the phase in the real domain, specifically along the geodesic γ , since the phase function takes complex values, the Hamilton-Jacobi equation holds naturally for the holomorphic extensions in the complex domain. From the geometric point of view, the complex Lagrangian manifold naturally associated to the phase function φ is not confined to the real domain but is a submanifold of the complexified phase space.

First we have

$$e^{-is\varphi}(-h^2\Delta_g - (hs)^2)e^{is\varphi}a = -h^2\Delta_g a - ih(1 + i\lambda h)[2\langle d\varphi, da \rangle_g + (\Delta_g \varphi)a] + (1 + i\lambda h)^2[\langle d\varphi, d\varphi \rangle_g - 1]a. \tag{2.2}$$

In the usual Gaussian beam construction in the C^∞ -setting, one solves the eikonal equation to a large, and sometimes infinite, order along the geodesic, see [43], [2], [44], [45]. Working in the present real analytic setting, it will be natural to solve the eikonal equation

$$\langle d\varphi, d\varphi \rangle_g - 1 = p(x, \varphi'_x(x)) = 0 \tag{2.3}$$

in a full neighborhood of the geodesic. Here

$$p(x, \xi) = |\xi|_g^2 - 1 = G(x)\xi \cdot \xi - 1 \tag{2.4}$$

is the semiclassical principal symbol of the operator $P = -h^2\Delta_g - (hs)^2$, where $G(x) = (g^{jk}(x))$. Since the metric g is real analytic, $p(x, \xi)$ extends to a holomorphic function in an open set of the form $\tilde{U} \times \mathbb{C}^n$, where

$$\tilde{U} \subset \mathbb{C}^n$$

is a complex neighborhood of U .

Let $(x(t), \xi(t)) = \exp(\frac{t}{2}H_p)(0, \xi_0)$ be the integral curve of the Hamiltonian H_p in T^*X , which corresponds to the unit speed geodesic γ , so that

$$\pi_x \left(\exp \left(\frac{t}{2} H_p \right) (0, \xi_0) \right) = \gamma(t), \quad t \in I \subset \mathbb{R},$$

where $\pi_x(x, \xi) = x$, and $\xi_0 = \dot{\gamma}(0)$. Here $\dot{\gamma}(0)$ is viewed as a cotangent vector using the Riemannian duality. Since $(0, \xi_0) \in p^{-1}(0) \cap (U \times \mathbb{R}^n)$, we therefore have $(x(t), \xi(t)) \in p^{-1}(0) \cap (U \times \mathbb{R}^n)$ for all $t \in I$. We have explicitly the Hamilton's equations

$$\begin{cases} \dot{x}(t) = \frac{1}{2}\partial_\xi p(x(t), \xi(t)), \\ \dot{\xi}(t) = -\frac{1}{2}\partial_x p(x(t), \xi(t)), \\ x(0) = 0, \\ \xi(0) = \xi_0. \end{cases} \tag{2.5}$$

Recalling that $x = (t, y) \in U$ and writing $\xi = (\tau, \eta) \in T_x^*X$ for the dual variable, we see from (2.5) that

$$\partial_\tau p(x(t), \xi(t)) \neq 0 \quad \text{for all } t \in I, \tag{2.6}$$

since the t component of $\dot{x}(t)$ is identically 1 in the (t, y) coordinates.

We look for a real analytic solution φ of (2.3) in U such that

$$\text{Im } \varphi(t, y) \geq 0, \quad \text{Im } \varphi(t, 0) = 0, \quad \text{Im } \varphi''_{yy}(t, 0) > 0, \quad t \in I, \tag{2.7}$$

and therefore,

$$\text{Im } \varphi(t, y) \sim |y|^2 = \text{dist}((t, y), \Gamma)^2, \quad (t, y) \in U.$$

Here and throughout the paper the notation $f_1 \sim f_2$ for positive functions f_1 and f_2 means that there exists a fixed $\epsilon > 0$ with $\epsilon f_1 < f_2 < \epsilon^{-1} f_1$. We will find the required real analytic solution of (2.3) as the restriction to $U \subset \mathbb{R}^n$ of a holomorphic function φ in $\tilde{U} \subset \mathbb{C}^n$, solving the following Cauchy problem for the Hamilton-Jacobi equation in the complex domain,

$$\begin{cases} p(x, \varphi'_x(x)) = 0, & x = (t, y) \in \tilde{U}, \\ \varphi(0, y) = \psi(y), \\ \varphi'_x(0) = \xi_0. \end{cases} \tag{2.8}$$

Here we take ψ to be a holomorphic function near $0 \in \mathbb{C}^{n-1}$ such that

$$\text{Im } \psi''_{yy}(0) > 0, \tag{2.9}$$

$\psi(0)$ is real, and so that the compatibility condition $\psi'_y(0) = \eta_0$ in (2.8) holds, with $\xi_0 = (\tau_0, \eta_0) \in \mathbb{R} \times \mathbb{R}^{n-1}$. Note that p, φ and ψ are holomorphic in their variables. For a holomorphic function $f(z_1, \dots, z_N)$ in an open set $V \subset \mathbb{C}^N$ we write $f'_z(z) = (\partial_{z_1} f(z), \dots, \partial_{z_N} f(z))$ for the complex gradient, $f''_{zz}(z) = (\partial_{z_j z_k} f(z))_{j,k=1}^N$ for the complex Hessian, etc. If $z_j = x_j + iy_j$, holomorphicity implies that $\partial_z^\alpha f(z) = \partial_x^\alpha f(z)$




Fig. 1. A Lagrangian submanifold $\Lambda \subset T^*X^{\mathbb{C}}$ satisfying $p(\Lambda) = 0$ is the union of the red integral curves of H_p in $T^*X^{\mathbb{C}}$ passing through $\Lambda' \subset T^*X^{\mathbb{C}}$, which is represented by the black arrows. Here $T^*X^{\mathbb{C}}$ is the cotangent bundle of the complexification of X , which is locally $\mathbb{C}_x^n \times \mathbb{C}_\xi^n$. (For interpretation of the colors in the figure(s), the reader is referred to the web version of this article.)

for any multi-index α . This shows that a holomorphic solution φ of (2.8) in $\tilde{U} \subset \mathbb{C}^n$ indeed yields a real analytic solution of (2.3) in U .

Remark. Let us note that [18, Theorem 5.5] gives the standard Hamilton-Jacobi theory locally near a point in the smooth case, and the extension of this theory to the holomorphic case is discussed in the remark following Theorem 1.8.2 in [23]. However here we need to construct the phase φ enjoying the good properties along the entire geodesic segment, and therefore, we shall give a detailed discussion of the construction below. The condition (2.9) will be crucial for this purpose.

Step 1. Solving near a point. In order to solve (2.8), we start by following the proof of [18, Theorem 5.5], see also [48]. The setting of our proof is illustrated in Fig. 1. To this end, we observe first that in view of (2.6), by the implicit function theorem applied to $p(0, \cdot, \cdot, \cdot)$, in a complex neighborhood of $(0, 0, \tau_0, \eta_0)$ we have $p(0, y, \tau, \eta) = 0$ if and only if $\tau = \lambda(y, \eta)$ where λ is a holomorphic function near $(0, \eta_0) \in \mathbb{C}^{2(n-1)}$ such that $\lambda(0, \eta_0) = \tau_0$.

Let us define

$$\Lambda' := \{(0, y, \tau, \eta) : \eta = \psi'_y(y), \tau = \lambda(y, \eta), y \in \text{neigh}(0, \mathbb{C}^{n-1})\} \subset \mathbb{C}^{2n}.$$

We have that Λ' is a complex manifold of complex dimension $n - 1$ such that

$$\Lambda' \subset p^{-1}(0),$$

which is isotropic in the sense that the restriction of σ to $T\Lambda' \times T\Lambda'$ vanishes:

$$\sigma|_{\Lambda'} = 0. \tag{2.10}$$

Here $\sigma = \sum_{j=1}^n d\xi_j \wedge dx^j$ is the complex symplectic form on (the tangent space of) $\mathbb{C}^{2n} = \mathbb{C}_x^n \times \mathbb{C}_\xi^n$. Indeed, if we fix $(x, \xi) \in \Lambda'$, any vector tangent to Λ' with base point at (x, ξ) is of the form $(0, V^y, V^\tau, V^\eta)$ with $V^\eta = \psi''_{yy} V^y$ and $V^y \in \mathbb{C}^{n-1}$. Applying σ to two such vectors gives $V_1^y \cdot \psi''_{yy} V_2^y - V_2^y \cdot \psi''_{yy} V_1^y = 0$, showing (2.10).

Note also that

$$\Lambda' \cap \mathbb{R}^{2n} = \{(0, 0, \tau_0, \eta_0)\}. \tag{2.11}$$

Indeed, $(0, 0, \tau_0, \eta_0) \in \Lambda' \cap \mathbb{R}^{2n}$ as $(\tau_0, \eta_0) \in \mathbb{R}^n$ and $\psi'_y(0) = \eta_0$. To see the opposite inclusion, let $(0, y, \lambda(y, \eta), \eta = \psi'(y)) \in \Lambda' \cap \mathbb{R}^{2n}$ and Taylor expand $\psi'(y)$ at $y = 0$,

$$\eta = \psi'(y) = \eta_0 + \psi''(0)y + \mathcal{O}(|y|^2), \quad y \in \mathbb{R}^{n-1}.$$

We have $\text{Im } \eta = \text{Im } \psi''(0)y + \mathcal{O}(|y|^2)$, and therefore, in view of (2.9), $\text{Im } \eta = 0$ implies that $y = 0$. This shows (2.11).

Let H_p be the complex Hamilton vector field of p , and let us consider the H_p flowout of Λ' :

$$\Lambda = \left\{ \exp\left(\frac{t}{2}H_p\right)(\rho) : \rho \in \Lambda', t \in \text{neigh}(I, \mathbb{C}) \right\} \subset \mathbb{C}^{2n}.$$

Here if $\mu = \sum_{j=1}^N a_j(z)\partial_{z_j}$ is a holomorphic vector field on an open set $V \subset \mathbb{C}^N$ in the sense that $a_j \in \text{Hol}(V)$, $j = 1, \dots, n$, we can define the flow $\exp(t\mu)(\rho)$, $\rho \in V$, locally for $t \in \text{neigh}(0, \mathbb{C})$, by solving the system of ODE,

$$\begin{cases} \dot{z}_j(t) = a_j(z(t)), & 1 \leq j \leq n, \\ z(0) = \rho, \end{cases}$$

see [16, Section 1] and the references given there.

Then $\Lambda' \subset \Lambda$, and since the flow of H_p preserves p , we have

$$\Lambda \subset p^{-1}(0),$$

and Λ is a \mathbb{C} -Lagrangian submanifold of \mathbb{C}^{2n} , see [18, Proposition 5.4] for a proof in the real case. The proof in the present holomorphic setting is similar. Let us also recall from [18, page 60] that the holomorphic Hamilton vector field H_p is tangent to Λ at each point of Λ . This is because Λ is a Lagrangian contained in $p^{-1}(0)$.

The differential of $\pi_x|_\Lambda$ is bijective at $(0, 0, \tau_0, \eta_0)$ since the differential of π_x is injective and since any Lagrangian submanifold has dimension $\dim(X)$. (The differential of $\pi_x|_\Lambda$ is injective since the differential of the exponential map $TX \rightarrow X$ is injective.) Consequently, there is a function $\varphi \in \text{Hol}(\text{neigh}(0, \mathbb{C}^n))$ such that

$$\Lambda = \Lambda_\varphi := \{(x, \varphi'_x(x)) : x \in \text{neigh}(0, \mathbb{C}^n)\}, \tag{2.12}$$

see [42, Section 5.6, Exercise 4], and also [18, Theorem 5.3] for the real version of this result. We have $\varphi'_x(0) = \xi_0$ and modifying φ by a constant we get $\varphi(0, y) = \psi(y)$, and such a solution is unique.

Step 2. Solving near γ . Let us denote the tangent space of Λ at $(0, 0, \tau_0, \eta_0)$ by Λ_0 and write

$$\Lambda_0 := T_{(0,0,\tau_0,\eta_0)}\Lambda = \{(\delta_x, \delta_\xi) \in \mathbb{C}^n \times \mathbb{C}^n : \delta_\xi = \varphi''_{xx}(0)\delta_x\}, \tag{2.13}$$

where in the second equality we used (2.12).

We claim that Λ_0 is a positive Lagrangian plane in the sense that

$$\frac{1}{i}\sigma(\rho, \bar{\rho}) \geq 0, \quad \rho \in \Lambda_0.$$

To this end, letting $M_0 = \varphi''_{xx}(0)$ and using (2.13), we write $\rho = (\delta_x, M_0\delta_x) \in \Lambda_0$. Then using that M_0 is symmetric, we get

$$\begin{aligned} \frac{1}{i}\sigma(\rho, \bar{\rho}) &= \frac{1}{i}(M_0\delta_x \cdot \bar{\delta}_x - \overline{M_0\delta_x} \cdot \delta_x) = 2\text{Im}(M_0\delta_x \cdot \bar{\delta}_x) \\ &= 2\text{Im}(M_0)\text{Re}\delta_x \cdot \text{Re}\delta_x + 2\text{Im}(M_0)\text{Im}\delta_x \cdot \text{Im}\delta_x, \end{aligned} \tag{2.14}$$

and therefore, it suffices to prove that

$$\text{Im } M_0 \geq 0. \tag{2.15}$$

In doing so, using (2.8), we write

$$M_0 = \begin{pmatrix} \varphi''_{tt}(0, 0) & \varphi''_{ty}(0, 0) \\ \varphi''_{yt}(0, 0) & \psi''_{yy}(0) \end{pmatrix}. \tag{2.16}$$

Using that H_p is tangent to Λ_φ , we see that $\exp(\frac{t}{2}H_p)(0, \xi_0) = (x(t), \varphi'_x(x(t)))$ is real for $t \in \text{neigh}(0, \mathbb{R})$, so that $\varphi'_t(t, 0), \varphi'_y(t, 0)$ are real. Hence,

$$\text{Im } M_0 = \begin{pmatrix} 0 & 0 \\ 0 & \text{Im } \psi''_{yy}(0) \end{pmatrix}, \tag{2.17}$$

and therefore, by the condition $\text{Im } \psi''_{yy}(0) > 0$ we imposed on ψ in (2.9), (2.15) follows.

For future reference, let us remark that

$$\Lambda_0 \cap \mathbb{R}^{2n} = \mathbb{R}H_p(0, \xi_0), \tag{2.18}$$

where $\mathbb{R}H_p(0, \xi_0) = \{sH_p(0, \xi_0) : s \in \mathbb{R}\}$. Indeed, we have $H_p(0, \xi_0) \in \Lambda_0 \cap \mathbb{R}^{2n}$ since the H_p vector field is tangent to Λ . On the other hand, if $(\delta_x, M_0\delta_x) \in \Lambda_0 \cap \mathbb{R}^{2n}$, it follows from (2.17) that

$$\delta_x = (\delta_t, 0) = \delta_t \dot{x}(0) = \delta_t p'_\xi(0, \xi_0),$$

where $\delta_t \in \mathbb{R}$. Here in the second equality we used that $(t, 0)$ corresponds to the geodesic in Fermi coordinates. We get $(\delta_x, M_0 \delta_x) = \delta_t (p'_\xi(0, \xi_0), M_0 p'_\xi(0, \xi_0)) = \delta_t H_p((0, \xi_0))$, which shows (2.18). Here in the last equality we used $H_p(0, \xi_0) \in \Lambda_0$.

Let

$$\kappa(t) := \exp\left(\frac{t}{2} H_p\right) : \Lambda \rightarrow \Lambda, \quad t \in I \subset \mathbb{R},$$

and therefore, the differential satisfies

$$d\kappa(t)(0, \xi_0) : \Lambda_0 \rightarrow T_{\kappa(t)(0, \xi_0)} \Lambda.$$

As the canonical transformation $\kappa(t)$ is real for each $t \in I$, $d\kappa(t)(0, \xi_0)$ preserves positivity, see [42, Section 5.6, Exercise 8], and therefore,

$$\Lambda_t := T_{\kappa(t)(0, \xi_0)} \Lambda \subset \mathbb{C}^{2n}$$

is a positive Lagrangian plane, for all $t \in I$.

We claim that Λ_t is transversal to the fiber $F = \{(0, \eta) : \eta \in \mathbb{C}^n\} \subset \mathbb{C}^{2n}$, for all $t \in I$, i.e. $\Lambda_t + F = \mathbb{C}^{2n}$. As $\dim \Lambda_t = n$, we have to show that $\Lambda_t \cap F = \{0\}$. Indeed, let $(0, \eta) \in \Lambda_t \cap F$. Then (2.13) implies that

$$\begin{pmatrix} 0 \\ \eta \end{pmatrix} = d\kappa(t)(0, \xi_0) \begin{pmatrix} \delta_x \\ M_0 \delta_x \end{pmatrix}, \tag{2.19}$$

for some $\delta_x \in \mathbb{C}^n$. We have

$$\begin{aligned} 0 &= \frac{1}{i} \sigma \left(\begin{pmatrix} 0 \\ \eta \end{pmatrix}, \overline{\begin{pmatrix} 0 \\ \eta \end{pmatrix}} \right) = \frac{1}{i} \sigma \left(d\kappa(t)(0, \xi_0) \begin{pmatrix} \delta_x \\ M_0 \delta_x \end{pmatrix}, d\kappa(t)(0, \xi_0) \overline{\begin{pmatrix} \delta_x \\ M_0 \delta_x \end{pmatrix}} \right) \\ &= \frac{1}{i} \sigma \left(\begin{pmatrix} \delta_x \\ M_0 \delta_x \end{pmatrix}, \overline{\begin{pmatrix} \delta_x \\ M_0 \delta_x \end{pmatrix}} \right) = 2 \operatorname{Im} (M_0 \delta_x \cdot \overline{\delta_x}). \end{aligned}$$

As $\operatorname{Im} M_0 \geq 0$, we get $(\operatorname{Im} M_0) \delta_x = 0$, and therefore, (2.17) implies that $\delta_x = \alpha p'_\xi(0, \xi_0)$ for some $\alpha \in \mathbb{C}$. Thus, by (2.19) we obtain that

$$\begin{pmatrix} 0 \\ \eta \end{pmatrix} = d\kappa(t)(0, \xi_0) (\alpha H_p(0, \xi_0)) = \alpha H_p(x(t), \xi(t)) = \alpha \begin{pmatrix} \dot{x}(t) \\ \dot{\xi}(t) \end{pmatrix}.$$

Since $\dot{x}(t) \neq 0$, we get $\alpha = 0$. Hence,

$$\eta = 0,$$

which establishes the claim.

As Λ_t is transversal to the fiber for all $t \in I$, by inspection of the proof of Theorem 5.5 in [18], we conclude that there exists $\varphi \in \text{Hol}(\text{neigh}(I \times B, \mathbb{C}^n))$ such that $\Lambda = \Lambda_\varphi$ and φ solves (2.8). The function φ is a continuation of the one appearing in (2.12). Notice that it is precisely thanks to the fact that the tangent plane Λ_t does not contain any non-zero vector of the form $(0, \eta)$ for all $t \in I$ that the proof of Theorem 5.5 in [18] applies near each point in $I \times \{0\}$, see also [26, Section 24.2].

Step 3. Properties of the solution. Next we shall check that the property (2.7), that is

$$\text{Im } \varphi(t, y) \geq 0, \quad \text{Im } \varphi(t, 0) = 0, \quad \text{Im } \varphi''_{yy}(t, 0) > 0, \quad t \in I,$$

holds for φ . First, $\varphi'_x(x(t)) = \xi(t)$ is real for $t \in I$. Writing

$$\frac{d}{dt} \varphi(x(t)) = \varphi'_x(x(t)) \cdot \dot{x}(t) = \xi(t) \cdot \frac{1}{2} p'_\xi(x(t), \xi(t)),$$

we have

$$\varphi(t, 0) = \psi(0) + \frac{1}{2} \int_0^t \xi(s) \cdot p'_\xi(x(s), \xi(s)) ds = \psi(0) + t, \tag{2.20}$$

as $\xi \cdot p'_\xi(x, \xi) = 2(p(x, \xi) + 1)$. Thus, using that $\psi(0)$ is real, we see that $\text{Im } \varphi(t, 0) = 0$ for $t \in I$. Furthermore, if $\psi(0) = 0$, we get $\varphi(t, 0) = t$.

Let $M(t) = \varphi''_{xx}(x(t))$. Then $M(t)$ is an $n \times n$ complex symmetric matrix depending real analytically on t , such that

$$\text{Im } M(t) \geq 0, \tag{2.21}$$

in view of the positivity of Λ_t . We claim that

$$\text{Im } M(t)|_W > 0, \tag{2.22}$$

where $W \subset \mathbb{R}^n$ is an algebraic supplement to $\mathbb{R}\dot{x}(t)$ so that $\mathbb{R}\dot{x}(t) \oplus W = \mathbb{R}^n$. To that end, let us observe first that

$$\Lambda_t \cap \mathbb{R}^{2n} = d\kappa(t)(0, \xi_0)(\Lambda_0 \cap \mathbb{R}^{2n}) = d\kappa(t)(0, \xi_0)(\mathbb{R}H_p(0, \xi_0)) = \mathbb{R}H_p(x(t), \xi(t)).$$

Here we have used (2.18) in the second equality. Let $v \in W$ be such that

$$\text{Im } M(t)v \cdot v = 0.$$

Hence, by (2.21), we get

$$\text{Im } M(t)v = 0.$$

Thus, $(v, M(t)v) \in \Lambda_t \cap \mathbb{R}^{2n} = \mathbb{R}H_p(x(t), \xi(t))$, and therefore, v is proportional to $p'_\xi(x(t), \xi(t)) = \dot{x}(t)$. This gives that $v = 0$, since $v \in W$. Hence, (2.22) follows, and we get $\text{Im } \varphi''_{yy}(t, 0) > 0$ for all $t \in I$.

Finally, we get $\text{Im } \varphi(t, y) \geq 0$ for all $(t, y) \in U$ by Taylor’s formula and by using that $\varphi'_x(x(t)) = \xi(t)$ is real. We have therefore constructed a real analytic solution φ of (2.3) such that (2.7) holds.

2.2. Construction of the amplitude

We shall follow [49, Theorem 9.3], where the construction of the amplitude as a classical analytic symbol is carried out in a neighborhood of a point, extending the construction to a full neighborhood of a geodesic segment.

We look for the amplitude a in the form of a formal power series in h ,

$$a(x; h) = \sum_{k=0}^{\infty} h^k a_k(x). \tag{2.23}$$

From (2.2), we see that we want to solve the following equation formally in powers of h ,

$$e^{-is\varphi}(-h^2\Delta_g - (hs)^2)e^{is\varphi}a = [-hiL_0 - ih\Delta_g\varphi + h^2(-\Delta_g + \lambda L_0 + \lambda\Delta_g\varphi)]a = 0, \tag{2.24}$$

in a fixed complex domain \tilde{U} , containing Γ . Here

$$L_0 = 2\langle d\varphi, d\cdot \rangle_g = 2G(x)\varphi'_x \cdot \partial_x = p'_\xi(x, \varphi'_x(x)) \cdot \partial_x, \tag{2.25}$$

where p is given in (2.4). The transport equation (2.24) can be written in the following form,

$$(hL_0 + hf(x) + h^2Q(x, D_x))a = 0, \tag{2.26}$$

where $f(x) = \Delta_g\varphi$ is a holomorphic function on \tilde{U} and $Q(x, D_x) = i(-\Delta_g + \lambda L_0 + \lambda\Delta_g\varphi)$ is a holomorphic differential operator of order 2. To solve (2.26), we remark first that the holomorphic vector field L_0 is transversal to each complex hypersurface $H_{t_0} = \{(t, y) \in \text{neigh}(I, \mathbb{C}) \times \text{neigh}(0, \mathbb{C}^{n-1}) : t = t_0 \in I\}$ at $(t_0, 0)$. Indeed,

$$p'_\xi(x(t), \varphi'_x(x(t))) \cdot \partial_x = p'_\tau(x(t), \varphi'_x(x(t)))\partial_t + p'_\eta(x(t), \varphi'_x(x(t))) \cdot \partial_y,$$

where $p'_\tau(x(t), \varphi'_x(x(t))) \neq 0$ for all $t \in I$ since $\partial_\tau p(x(t), \xi(t)) \neq 0$ for all $t \in I$ as noted in (2.6). Thus, substituting (2.23) into (2.26), we get a sequence of transport equations which can all be solved uniquely in a suitable complex domain containing Γ , provided that $a|_{H_{t_0}}$ is prescribed, for some $t_0 \in I$. However, the difficulty here is that we would like our solution $a(x; h)$ to be a classical analytic symbol, and following [49, Section 9], we shall establish this fact making use of the method of “nested neighborhoods” introduced

in [49]. Contrary to [49, Theorem 9.3], where the family of “nested neighborhoods” is considered near a point, here we shall work in such neighborhoods near a piece of the geodesic.

For simplicity, let us take $t_0 = 0$. We look for solution to (2.26) by using convenient coordinates. The coordinates we will use are the usual flowout coordinates (see e.g. [39]), which we show to exist for L_0 on a neighborhood of a given interval.

Lemma 2.1. *Let $J \subset\subset I$ be an open interval. There exist local holomorphic coordinates $(s, z) \in \text{neigh}(\overline{J}, \mathbb{C}) \times \text{neigh}(0, \mathbb{C}^{n-1})$ such that the hyperplane H_0 is given by the equation $s = 0$ and $L_0 = \frac{\partial}{\partial s}$.*

Proof. We continue to work in the Fermi coordinates $x = (t, y)$ and recall from [28] that

$$G(t, y) = (g^{jk}(t, y)) = 1 + \mathcal{O}(|y|^2). \tag{2.27}$$

Now (2.3), (2.4), and (2.27) imply that

$$(\varphi'_t)^2(t, 0) + (\varphi'_y)^2(t, 0) = 1, \tag{2.28}$$

and therefore, it follows from (2.20) and (2.28) that $\varphi'_y(t, 0) = 0$. Hence, Taylor expanding $\varphi(t, y)$ at $y = 0$, we get

$$\varphi(t, y) = \psi(0) + t + \mathcal{O}(|y|^2). \tag{2.29}$$

It follows from (2.25), (2.27), and (2.29) that

$$L_0 = 2(1 + \mathcal{O}(|y|^2)) \begin{pmatrix} 1 + \mathcal{O}(|y|^2) \\ \mathcal{O}(|y|) \end{pmatrix} \cdot \begin{pmatrix} \partial_t \\ \partial_y \end{pmatrix} = 2(1 + \mathcal{O}(|y|^2))\partial_t + \mathcal{O}(|y|) \cdot \partial_y. \tag{2.30}$$

Consider the initial value problem for the flow $\exp(sL_0)(0, z)$,

$$\begin{cases} \partial_s(t, y)(s, z) = L_0((t, y)(s, z)), \\ (t, y)(0, z) = (0, z), \end{cases} \tag{2.31}$$

where $(s, z) \in \text{neigh}(I, \mathbb{C}) \times \text{neigh}(0, \mathbb{C}^{n-1})$. In particular, $y(s, z)|_{z=0} = 0$ and therefore, $y(s, z) = \mathcal{O}(|z|)$. Differentiating the first equation in (2.31) in z_j and using (2.30), we get

$$\begin{cases} \partial_s(\partial_{z_j} t(s, z)) = \mathcal{O}(y(s, z)\partial_{z_j} y) = \mathcal{O}(|z|), \\ \partial_{z_j} t(0, z) = 0. \end{cases} \tag{2.32}$$

Hence,

$$\partial_{z_j} t(s, z) = \mathcal{O}(|z|). \tag{2.33}$$

Consider the holomorphic map

$$F : \text{neigh}(I, \mathbb{C}) \times \text{neigh}(0, \mathbb{C}^{n-1}) \ni (s, z) \mapsto (t, y)(s, z).$$

In view of (2.33), the differential $DF(s, 0)$ is given by

$$DF(s, 0) = \begin{pmatrix} t'_s(s, 0) & 0 & \dots & 0 \\ y'_{1s}(s, 0) & y'_{1z_1}(s, 0) & \dots & y'_{1z_{n-1}}(s, 0) \\ \vdots & \vdots & \dots & \vdots \\ y'_{n-1s}(s, 0) & y'_{n-1z_1}(s, 0) & \dots & y'_{n-1z_{n-1}}(s, 0) \end{pmatrix}, \tag{2.34}$$

where $t'_s(s, 0) = 2(1 + \mathcal{O}(|y(s, 0)|^2)) = 2$. By Liouville’s formula, see [24, Theorem 1.2.5], we know that the last $n - 1$ columns in (2.34) are linearly independent, and therefore, $\det(DF(s, 0)) \neq 0$ for all $s \in \text{neigh}(I, \mathbb{C})$. Furthermore, $F|_{I \times \{0\}}$ is injective as $F(s, 0) = (t(s, 0), 0) = (2s, 0)$. An application of a holomorphic version of [28, Lemma 7.3] allows us to conclude that F is a holomorphic diffeomorphism in $\text{neigh}(\bar{J}, \mathbb{C}) \times \text{neigh}(0, \mathbb{C}^{n-1})$ where $J \subset\subset I$ is an open interval.

Now writing $x = (t, y)$, in view of (2.31), we see that

$$\frac{\partial}{\partial s} u(x(s, y)) = u'_x(x(s, y)) \cdot \dot{x}(s, y) = (L_0 u)(x(s, y)).$$

Finally, it follows from (2.30) and (2.31) that

$$\begin{cases} \partial_s t(s, z) = 2(1 + \mathcal{O}(|z|^2)), \\ t(0, z) = 0, \end{cases}$$

and therefore, $t(s, z) = 2s + \mathcal{O}(|z|^2)s$. Hence, $t = 0$ is equivalent to the fact that $s = 0$, showing that the hyperplane H_0 is given by the equation $s = 0$. \square

Passing to the new holomorphic coordinates provided by Lemma 2.1, and renaming them as $x = (t, y)$, we are led from (2.26) to consider the following initial value problem,

$$\begin{cases} (h \frac{\partial}{\partial t} + hf(x) + h^2 Q(x, D_x))a = 0, \\ a|_{t=0} = w(y; h), \end{cases} \tag{2.35}$$

where $w(y; h)$ is a classical analytic symbol near $0 \in \mathbb{C}^{n-1}$. We would like to find a classical analytic symbol a solving (2.35). Here f is a holomorphic function, and Q is a holomorphic differential operator of order 2. To that end, it suffices to solve the following problem,

$$\begin{cases} (h \frac{\partial}{\partial t} + hf(x) + h^2 Q(x, D_x))a = hv, \\ a|_{t=0} = 0, \end{cases} \tag{2.36}$$

where $v(x; h)$ is a classical analytic symbol in $\text{neigh}(\overline{J}, \mathbb{C}) \times \text{neigh}(0, \mathbb{C}^{n-1})$. This is because a solution a to (2.36) with $v = -(\frac{\partial}{\partial t} + f(x) + hQ(x, D_x))v_0$ and $v_0|_{t=0} = w$, implies that $a + v_0$ solves (2.35). Using that

$$\partial_t + f(t, y) = e^{-F(t,y)} \circ \partial_t \circ e^{F(t,y)},$$

where $F'_t(t, y) = f(t, y)$, we may assume that $f(x) = 0$.

We shall first carry out the analysis of (2.36) under the assumption that the interval J is symmetric about the origin and after a rescaling we may assume that $\overline{J} = [-1, 1]$. Let $\Omega \subset \mathbb{C}^n$ be open such that $[-1, 1]_t \times \{0\}_y \subset \Omega$ and Ω is in the domain of definition of various symbols. Then let $0 < \varepsilon < 1, r > 0$ be small but fixed so that if we set

$$\Omega_0 = \left\{ (t, y) \in \mathbb{C}^n : \frac{|y|}{\varepsilon} + \frac{|\text{Im } t|}{\varepsilon} + |\text{Re } t| < 1 + r \right\}$$

then $\overline{\Omega_0} \subset \Omega$. Consider the family of open sets,

$$\Omega_s = \left\{ (t, y) \in \mathbb{C}^n : \frac{|y|}{\varepsilon} + \frac{|\text{Im } t|}{\varepsilon} + |\text{Re } t| < 1 + r - s \right\},$$

with $0 \leq s < r$. Note that Ω_s is a family of “nested neighborhoods” of $[-1, 1] \times 0$ in the sense of [49, Theorem 9.3], so that we have

- (i) if $s_1 > s_2$ then $\Omega_{s_1} \subset \Omega_{s_2}$,
- (ii) there exists $\delta > 0$ such that for all $s_1 > s_2$ and all $x \in \Omega_{s_1}$ we have the inclusion $B_{\mathbb{C}^n}(x, \delta(s_1 - s_2)) \subset \Omega_{s_2}$.

Given $\mu > 0$, we say that $a \in \mathcal{A}_\mu$, if $a(x; h) = \sum_{k=0}^\infty a_k(x)h^k$, a is holomorphic in Ω , such that for all $s \in (0, r)$,

$$\sup_{\Omega_s} |a_k| \leq \frac{f(a, k)}{s^k} k^k, \tag{2.37}$$

where $f(a, k)$ is the best constant for which (2.37) holds, and

$$\sum_{k=0}^\infty f(a, k) \mu^k := \|a\|_\mu < \infty. \tag{2.38}$$

Now if $a \in \mathcal{A}_\mu$ for some $\mu > 0$ then $f(a, k) \leq C^{k+1}$, $k = 0, 1, 2, \dots$, and therefore, a is a classical analytic symbol on Ω_0 . Let

$$(\partial_t^{-1}a)(t, y) = \int_0^t a(\tau, y) d\tau. \tag{2.39}$$

We shall need the following result, see [49, Theorem 9.3] and [46, Lemma 5.5].

Lemma 2.2. *Let $a \in \mathcal{A}_\mu$ be of the form*

$$a = \sum_{k=2}^{\infty} h^k a_k,$$

and let $b = (h\partial_t)^{-1}a$. Then

$$\|b\|_\mu \leq \mathcal{O}\left(\frac{1}{\mu}\right)\|a\|_\mu. \tag{2.40}$$

Proof. We have

$$b = \sum_{k=2}^{\infty} h^{k-1} \partial_t^{-1} a_k = \sum_{k=1}^{\infty} h^k b_k,$$

where $b_k = \partial_t^{-1} a_{k+1}$. Let us estimate $\sup_{\Omega_s} |b_k|$. To that end, we write

$$b_k(x) = t \int_0^1 a_{k+1}(\sigma t, y) d\sigma$$

We claim that for $0 \leq \sigma \leq 1$, if $x = (t, y) \in \Omega_s$ then

$$(\sigma t, y) \in \Omega_{s+(1-\sigma)|t|}. \tag{2.41}$$

Indeed, using that $0 < \varepsilon < 1$, we get

$$\begin{aligned} \frac{|y|}{\varepsilon} + \frac{\sigma|\text{Im } t|}{\varepsilon} + \sigma|\text{Re } t| &< 1 + r - s - \frac{(1-\sigma)}{\varepsilon}|\text{Im } t| - (1-\sigma)|\text{Re } t| \\ &< 1 + r - (s + (1-\sigma)|t|), \end{aligned}$$

showing $(\sigma t, y) \in \Omega_{s+(1-\sigma)|t|}$ as claimed. It follows from (2.37), (2.41) that for $x \in \Omega_s$, we have

$$\begin{aligned} |b_k(x)| &\leq |t| f(a, k+1) (k+1)^{k+1} \int_0^1 \frac{d\sigma}{(s + (1-\sigma)|t|)^{k+1}} \\ &= f(a, k+1) (k+1)^{k+1} |t| \int_0^1 \frac{d\sigma}{(s + \sigma|t|)^{k+1}} \\ &= f(a, k+1) (k+1)^{k+1} \int_0^{|t|} \frac{d\sigma}{(s + \sigma)^{k+1}} \end{aligned}$$

$$\begin{aligned} &\leq f(a, k + 1)(k + 1)^{k+1} \int_0^\infty \frac{d\sigma}{(s + \sigma)^{k+1}} = f(a, k + 1)(k + 1)^{k+1} \int_s^\infty \frac{d\sigma}{\sigma^{k+1}} \\ &= f(a, k + 1) \frac{(k + 1)^{k+1}}{ks^k}. \end{aligned}$$

Here we have used that $k \geq 1$. Thus, for any $0 < s < r$, we get

$$\sup_{\Omega_s} |b_k| \leq f(a, k + 1) \frac{(1 + 1/k)k^k(1 + 1/k)^k}{s^k} \leq \frac{2ef(a, k + 1)}{s^k} k^k,$$

and therefore by the definition of $f(b, k)$, see (2.37), we have

$$f(b, k) \leq 2ef(a, k + 1), \quad k = 1, 2, \dots$$

Using (2.38), we obtain that

$$\|b\|_\mu = \sum_{k=1}^\infty f(b, k)\mu^k \leq \sum_{k=1}^\infty 2ef(a, k + 1)\mu^k = \frac{2e}{\mu} \|a\|_\mu,$$

establishing (2.40). \square

Now applying to $(h\partial_t)^{-1}$ to (2.36), we get

$$a + (h\partial_t)^{-1}h^2Q(x, D_x)a = \partial_t^{-1}v. \tag{2.42}$$

Here $\partial_t^{-1}v$ is a classical analytic symbol in Ω_0 . To proceed, we need the following result.

Lemma 2.3. *Let $a \in \mathcal{A}_\mu$. Then $(h\partial_t)^{-1}h^2Q(x, D_x)a \in \mathcal{A}_\mu$ with*

$$\|(h\partial_t)^{-1}h^2Q(x, D_x)a\|_\mu \leq \mathcal{O}(\mu) \|a\|_\mu \tag{2.43}$$

Proof. Writing $a(x) = \sum_{k=0}^\infty a_k(x)h^k$, we get

$$h^2Q(x, D_x)a = \sum_{k=2}^\infty h^kQ(x, D_x)a_{k-2}.$$

For $s_1 > s_2$, in view of the property (ii) of the “nested neighborhoods” Ω_s , and (2.37), we obtain for $k = 2, 3, \dots$ that

$$\sup_{\Omega_{s_1}} |Q(x, D_x)a_{k-2}| \leq \frac{C}{(s_1 - s_2)^2} \sup_{\Omega_{s_2}} |a_{k-2}| \leq \frac{C}{(s_1 - s_2)^2} \frac{f(a, k - 2)}{s_2^{k-2}} (k - 2)^{k-2}. \tag{2.44}$$

The Cauchy estimate was used here in the first inequality. Taking $0 < s_2 = \frac{k-2}{k} s_1 < s_1$ for $k = 3, 4, \dots$, we get from (2.44) that

$$\sup_{\Omega_{s_1}} |Q(x, D_x)a_{k-2}| \leq \frac{Cf(a, k-2)}{s_1^k} k^k,$$

and therefore, in view of (2.37),

$$f(Q(x, D_x)a_{k-2}, k) \leq Cf(a, k-2).$$

Thus, by the definition of $\|\cdot\|_\mu$, see (2.38), we obtain that

$$\|h^2Q(x, D_x)a\|_\mu \leq \sum_{k=2}^\infty \mu^k Cf(a, k-2) \leq \mathcal{O}(\mu^2)\|a\|_\mu. \tag{2.45}$$

Lemma 2.2 together with (2.45) implies that

$$\|(h\partial_t)^{-1}h^2Q(x, D_x)a\|_\mu \leq \mathcal{O}(1/\mu)\|h^2Q(x, D_x)a\|_\mu \leq \mathcal{O}(\mu)\|a\|_\mu,$$

establishing (2.43). \square

It follows from Lemma 2.3 that

$$\|\tilde{Q}^j w\|_\mu \leq \mathcal{O}(\mu^j)\|w\|_\mu, \quad \tilde{Q} := (h\partial_t)^{-1}h^2Q(x, D_x),$$

for $w \in \mathcal{A}_\mu$, and therefore, by Neumann series argument, we have that the equation (2.42),

$$a + (h\partial_t)^{-1}h^2Q(x, D_x)a = \partial_t^{-1}v,$$

has a unique solution a with $\|a\|_\mu < \infty$ for $\mu > 0$ small enough. Thus, a is a classical analytic symbol in Ω_0 .

We shall next proceed to solve (2.35) when the interval J is not necessarily symmetric with respect to the origin, $\bar{J} = [a, b]$ where $a < 0 < b$. Without loss of generality, we may assume that $0 < a + b$. Let $N \geq 0$ be the largest integer such that $(N + 1)|a| < b$. We first solve (2.35) with the initial condition prescribed at $t = 0$ in the symmetric interval $[a, |a|]$ and obtain a unique classical analytic symbol $a^{(0)}$ in a complex neighborhood of $[a, |a|] \times \{0\}$. Next we solve the initial value problem,

$$\begin{cases} (h\partial_t + h^2Q(x, D_x))a^{(1)} = 0, \\ a^{(1)}|_{t=|a|} = a^{(0)}|_{t=|a|}, \end{cases} \tag{2.46}$$

in a complex neighborhood of $[0, 2|a|] \times \{0\}$. Continuing this process and working the symmetric intervals of the form $[(j - 1)|a|, (j + 1)|a|]$, $j = 2, \dots, N$, we construct a classical analytic symbol in a complex neighborhood of $[a, (N + 1)|a|] \times \{0\}$ solving (2.35). Finally solving (2.46) with the initial condition prescribed at $t = (N + 1)|a|$ in a

complex neighborhood of $[2(N + 1)|a| - b, b] \times \{0\}$, we obtain a classical analytic symbol in a complex neighborhood of $[a, b] \times \{0\}$ solving (2.35).

Furthermore, demanding that $a|_{t=0}$ should be an elliptic classical analytic symbol near 0 in \mathbb{C}^{n-1} , we conclude that the classical analytic symbol $a(x; h)$ is elliptic in the sense that $a_0 \neq 0$. This completes the construction of the amplitude as an elliptic classical analytic symbol.

It follows from (2.24) that for all $N \geq 1$,

$$e^{-is\varphi}(-h^2\Delta_g - (hs)^2)e^{is\varphi} \left(\sum_{j=0}^N h^j a_j \right) = h^{N+2}(-\Delta_g a_N + \lambda(L_0 + \Delta_g \varphi)a_N)$$

in a complex neighborhood \tilde{U} of Γ . Using (1.3) and Cauchy’s estimates, we obtain after an arbitrarily small decrease of \tilde{U} that

$$\left| e^{-is\varphi}(-h^2\Delta_g - (hs)^2)e^{is\varphi} \left(\sum_{k=0}^N h^k a_k \right) \right| \leq h^{N+2} C^{N+1} N^N.$$

Choosing $N = N(h) = \lceil \frac{1}{heC} \rceil$, we obtain that

$$\left| e^{-is\varphi}(-h^2\Delta_g - (hs)^2)e^{is\varphi} \left(\sum_{j=0}^{N(h)} h^k a_k \right) \right| \leq \mathcal{O}(1)e^{-\frac{1}{C_1 h}}, \quad C_1 > 0,$$

for all $0 < h \leq 1$. Note that we also have

$$\left| \sum_{j=0}^{N(h)} h^k a_k \right| \leq C \sum_{k=0}^{N(h)} e^{-k} \leq C \frac{e}{e-1}.$$

In the estimate above we used $k \leq N(h) = \lceil 1/(heC) \rceil$ and $|a_k| \leq C^{k+1} k^k$, which holds since a is classical analytic symbol.

Let $\chi \in C_0^\infty(\mathbb{R}^{n-1})$ be such that $0 \leq \chi \leq 1$, $\chi = 1$ for $|y| \leq 1/4$ and $\chi = 0$ for $|y| \geq 1/2$. In view of (2.1) we set

$$v(t, y; h) = h^{-\frac{(n-1)}{4}} e^{is\varphi(t,y)} a(t, y; h), \quad a(t, y; h) = \left(\sum_{k=0}^{N(h)} h^k a_k \right) \chi\left(\frac{y}{\delta'}\right). \tag{2.47}$$

Here $\delta' > 0$ is chosen sufficiently small so that $\chi(y/\delta')$ is zero outside the set where we have constructed the functions φ and a_k . Since $\text{Im} \varphi(t, y) \geq \frac{|y|^2}{C}$, $C > 0$, the cutoff function χ does not destroy the exponential smallness of the error, and we see that v satisfies after an arbitrarily small decrease of the real domain U ,

$$\|v\|_{L^2(U)} \asymp 1, \quad \|(-h^2\Delta - (hs)^2)v\|_{L^2(U)} = \mathcal{O}(e^{-\frac{1}{C_1 h}}), \quad C > 0, \tag{2.48}$$

as $h \rightarrow 0$. Here for the first bound, we use the fact that $a_0 \neq 0$.

2.3. Gluing the local quasimodes together

Let us now return to the open Fermi cover $(U_j)_{j=0}^{N+1}$ of $\gamma([-T_1 - \varepsilon, T_2 + \varepsilon])$, replacing it if necessary by a slightly smaller relatively compact subcover. We have constructed φ_0 real analytic in U_0 solving the Cauchy problem (2.8) in a complex neighborhood of U_0 so that $\varphi_0(t, 0) = t$ for all $t \in I_0$ and $\text{Im } \varphi_0''(t, 0) > 0$ for all $t \in I_0$. In order to construct $\varphi = \varphi_1$ real analytic in U_1 , we pick $t_0 \in I_0 \cap I_1$ and solve in a complex neighborhood of U_1 ,

$$\begin{cases} p(x, \varphi'_x(x)) = 0, \\ \varphi|_{t=t_0} = \varphi_0(t_0, y), \\ \varphi'_x(t_0) = \xi(t_0). \end{cases}$$

We get φ_1 such that $\varphi_1 = \varphi_0$ near $(t_0, 0) \in U_0 \cap U_1$, and thus, by unique continuation, $\varphi_1 = \varphi_0$ in $U_0 \cap U_1$, assuming as we may that $U_0 \cap U_1$ is connected. Note that in view of (2.20), we have $\varphi_1(t, 0) = t$ for all $t \in I_1$. Continuing in this way, we obtain φ_j real analytic in U_j , $0 \leq j \leq N + 1$, such that $\varphi_j = \varphi_{j+1}$ in $U_j \cap U_{j+1}$ and (2.7) holds for every φ_j .

Next let $a^{(0)}(t, y; h)$ be an elliptic classical analytic symbol in a complex neighborhood of U_0 obtained by solving (2.26). To get $a^{(1)}(t, y; h)$, we solve the sequence of transport equations (2.26) with $\varphi = \varphi_1$ and with $a^{(1)}|_{t=t_0} = a^{(0)}|_{t=t_0}$. Thus, by uniqueness and analytic continuation, $a^{(1)} = a^{(0)}$ in $U_0 \cap U_1$. Continuing in the same way, we get v_0, v_1, \dots, v_{N+1} such that

$$v_j = v_{j+1} \quad \text{in } U_j \cap U_{j+1}. \tag{2.49}$$

Let $\chi_j = \chi_j(t) \in C_0^\infty(I_j)$ be such that $\sum_{j=0}^{N+1} \chi_j = 1$ near $[-T_1 - \varepsilon, T_2 + \varepsilon]$, and define our quasimode v globally by

$$v = \sum_{j=0}^{N+1} \chi_j v_j.$$

Let $p_1, \dots, p_R \in X^{\text{int}}$ be the distinct points where the geodesic γ self-intersects, and let $-T_1 < t_1 < \dots < t_N < T_2$ be the times of self-intersections. Let V_1, \dots, V_R be small neighborhoods in X around p_j , $j = 1, \dots, R$. Then choosing δ' in (2.47) small enough we obtain an open cover of a neighborhood of $\gamma[-T_1, T_2]$ in \widehat{X} ,

$$\text{supp } (v(\cdot; h)) \cap X \subset (\cup_{j=1}^R V_j) \cup (\cup_{k=1}^S W_k), \tag{2.50}$$

where in each V_j , the quasimode is a finite sum,

$$v(\cdot; h)|_{V_j} = \sum_{l:\gamma(t_l)=p_j} v_l(\cdot; h), \tag{2.51}$$

and in each W_k (where there are no self-intersecting points), in view of (2.49), there is some $l(k)$ so that the quasimode is given by

$$v(\cdot; h)|_{W_k} = v_{l(k)}(\cdot; h). \tag{2.52}$$

Finally, the bounds (1.4) follows from the bounds (2.48), and the representations (2.51) and (2.52) of v . This completes the proof of Theorem 1.12.

3. Some preliminary results about geodesics

Let (X, g) be a compact Riemannian manifold of dimension $n \geq 2$ with smooth boundary, contained in an open real analytic manifold (\widehat{X}, g) of the same dimension with g real analytic in \widehat{X} . First we have the following analog of [13, Lemma 2.1], established in this work in the smooth case.

Lemma 3.1. *Let $\alpha_0 = (x_0, \xi_0) \in S^*X^{\text{int}}$ and let $\zeta_1, \zeta_2 \in S^*_{x_0}X^{\text{int}}$ be such that*

$$\zeta_1 + \zeta_2 = t_0\xi_0 \tag{3.1}$$

for some $0 < t_0 < 2$. Then there exists a neighborhood U of α_0 in S^*X^{int} and a real analytic map

$$I : U \rightarrow S^*X^{\text{int}} \times S^*X^{\text{int}}, \quad (x, \xi) \mapsto (x, \omega_1(x, \xi)) \times (x, \omega_2(x, \xi))$$

such that

$$I(x_0, \xi_0) = (x_0, \zeta_1) \times (x_0, \zeta_2), \tag{3.2}$$

and

$$\omega_1(x, \xi) + \omega_2(x, \xi) = t_0\xi, \quad (x, \xi) \in U. \tag{3.3}$$

Proof. We follow the proof of [13, Lemma 2.1] with minor changes in the real analytic setting, and the argument is presented here only for the convenience of the reader.

Let x_1, \dots, x_n be real analytic local coordinates on X^{int} centered at x_0 such that $G(0) = 1$. Here $G = G(x) = (g^{jk})$ is the co-metric tensor. It follows from (3.1) upon taking the scalar product with ζ_1, ζ_2 , and ξ_0 , that $\zeta_1 \cdot \xi_0 = \zeta_2 \cdot \xi_0$ and $t_0 = 2\zeta_1 \cdot \xi_0$. Similarly, if (3.3) holds, then $t_0 = 2G(x)\omega_1(x, \xi) \cdot \xi$, and we therefore should have

$$G(x)\omega_1(x, \xi) \cdot \xi = \zeta_1 \cdot \xi_0. \tag{3.4}$$

Furthermore, if (3.3) is valid, then

$$\omega_2(x, \xi) = 2(\zeta_1 \cdot \xi_0)\xi - \omega_1(x, \xi).$$

Note that this implies that $|\omega_2(x, \xi)|_{G(x)} = 1$, provided that (3.4) holds, and therefore, we only need to determine $\omega_1(x, \xi) \in S^*X^{\text{int}}$ depending analytically on (x, ξ) such that $\omega_1(x_0, \xi_0) = \zeta_1$ and (3.4) is valid.

To that end, let us set

$$\zeta(x) = \frac{\zeta_1}{\sqrt{G(x)\zeta_1 \cdot \zeta_1}}.$$

Then $\zeta(x)$ is real analytic in x and $\zeta(0) = \zeta_1$. Let

$$\tilde{\omega}_1(x, \xi) = \zeta(x) + \alpha(x, \xi)\xi,$$

for $(x, \xi) \in \text{neigh}((x_0, \xi_0), S^*\mathbb{R}^n)$, with some $\alpha = \alpha(x, \xi)$ to be chosen. We have

$$G(x)\tilde{\omega}_1(x, \xi) \cdot \tilde{\omega}_1(x, \xi) = 1 + \alpha^2 + 2\alpha G(x)\zeta(x) \cdot \xi.$$

We set

$$\omega_1(x, \xi) = \frac{\tilde{\omega}_1(x, \xi)}{\sqrt{G(x)\tilde{\omega}_1(x, \xi) \cdot \tilde{\omega}_1(x, \xi)}} = \frac{\zeta(x) + \alpha\xi}{\sqrt{1 + \alpha^2 + 2\alpha G(x)\zeta(x) \cdot \xi}}. \tag{3.5}$$

We would like to find α so that $\alpha(0, \xi_0) = 0$ and that (3.4) holds. The former requirement guarantees that $\omega_1(0, \xi_0) = \zeta_1$, and the latter requirement implies that we should have

$$F(x, \xi, \alpha) = 0,$$

where

$$F(x, \xi, \alpha) := (1 - (\zeta_1 \cdot \xi_0)^2)\alpha^2 + 2G(x)\zeta(x) \cdot \xi(1 - (\zeta_1 \cdot \xi_0)^2)\alpha + (G(x)\zeta(x) \cdot \xi)^2 - (\zeta_1 \cdot \xi_0)^2.$$

Note that F is real analytic in $(x, \xi, \alpha) \in \text{neigh}((0, \xi_0), S^*\mathbb{R}^n) \times \mathbb{R}$, $F(0, \xi_0, 0) = 0$, and $F'_\alpha(0, \xi_0, 0) = 2(\zeta_1 \cdot \xi_0)(1 - (\zeta_1 \cdot \xi_0)^2) \neq 0$, as $0 < \zeta_1 \cdot \xi_0 = \frac{t_0}{2} < 1$. Thus, by the implicit function theorem, there is a neighborhood U of $(0, \xi_0)$ and a unique real analytic function $\alpha(x, \xi)$ in U such that $\alpha(0, \xi_0) = 0$ and $F(x, \xi, \alpha) = 0$ if and only if $\alpha = \alpha(x, \xi)$. Hence, $\omega_1(x, \xi)$ given in (3.5) satisfies the conditions of the proposition. This completes the proof. \square

We shall also need the following result.

Lemma 3.2. *Let $\alpha_0 = (x_0, \xi_0) \in S^*X^{int}$. Assume that α_0 is generated by an admissible pair of geodesics $\gamma_1(\alpha_0) : [-T_1(\alpha_0), T_2(\alpha_0)] \rightarrow X$ and $\gamma_2(\alpha_0) : [-S_1(\alpha_0), S_2(\alpha_0)] \rightarrow X$, where $0 < T_1(\alpha_0), T_2(\alpha_0), S_1(\alpha_0), S_2(\alpha_0) < \infty$. Then there exists a neighborhood V of α_0 in S^*X^{int} such that every point $\alpha = (\alpha_x, \alpha_\xi) \in V$ is generated by an admissible pair of geodesics $\gamma_1(\alpha) : [-T_1(\alpha), T_2(\alpha)] \rightarrow X$ and $\gamma_2(\alpha) : [-S_1(\alpha), S_2(\alpha)] \rightarrow X$, which depend real analytically on α .*

Proof. First we have

$$\zeta_1 + \zeta_2 = t_0 \xi_0,$$

for some $0 < t_0 < 2$, where $\zeta_j = \dot{\gamma}_j(\alpha_0)$ is viewed as a cotangent vector, using the Riemannian duality. By Lemma 3.1, there exists a neighborhood U of α_0 in S^*X^{int} and a real analytic map

$$I : U \rightarrow S^*X^{int} \times S^*X^{int}, \quad \alpha = (\alpha_x, \alpha_\xi) \mapsto (\alpha_x, \omega_1(\alpha)) \times (\alpha_x, \omega_2(\alpha))$$

such that

$$I(x_0, \xi_0) = (x_0, \zeta_1) \times (x_0, \zeta_2),$$

and

$$\omega_1(\alpha) + \omega_2(\alpha) = t_0 \alpha_\xi. \tag{3.6}$$

The corresponding unit speed geodesics $\gamma_1(\alpha) : [-T_1(\alpha), T_2(\alpha)] \rightarrow X$ and $\gamma_2(\alpha) : [-S_1(\alpha), S_2(\alpha)] \rightarrow X$, $T_1(\alpha), T_2(\alpha), S_1(\alpha), S_2(\alpha) > 0$, such that $\gamma_j(\alpha)(0) = \alpha_x$, $\dot{\gamma}_j(\alpha)(0) = \omega_j(\alpha)$, are non-tangential being small perturbations of the non-tangential geodesics $\gamma_j(\alpha_0)$, $j = 1, 2$. Hence, the functions $T_j(\alpha)$ and $S_j(\alpha)$ depend continuously on $\alpha \in U$ and in particular, they are bounded after an arbitrarily small decrease of U . Note also that (3.6) implies that $\gamma_1(\alpha)$ and $\gamma_2(\alpha)$ are two distinct geodesics and that are not reverses of each other.

We claim that there is a neighborhood $\tilde{U} \subset U$ of α_0 such that for all $\alpha \in \tilde{U}$, we have

$$\gamma_1(\alpha)(t) = \gamma_1(\alpha)(0) \iff t = 0. \tag{3.7}$$

Indeed, otherwise there exists a sequence $\alpha_k \rightarrow \alpha_0$ as $k \rightarrow \infty$ and $0 \neq t_k \in [-T_1(\alpha_k), T_2(\alpha_k)]$ such that

$$\gamma_1(\alpha_k)(t_k) = \gamma_1(\alpha_k)(0), \tag{3.8}$$

for all k . Assuming, as we may, that $t_k \rightarrow t_0$, we get from (3.8) that $\gamma_1(\alpha_0)(t_0) = \gamma_1(\alpha_0)(0)$. Since the geodesic $\gamma_1(\alpha_0)$ does not self-intersect at $x_0 = \gamma_1(\alpha_0)(0)$, we conclude that $t_0 = 0$. Since $\gamma_1(\alpha_k)(t_k) \rightarrow \gamma_1(\alpha_0)(0) = x_0 \in X^{int}$ as $k \rightarrow \infty$, for all k

sufficiently large, we see that $\gamma_1(\alpha_k)(t_k) \in X^{\text{int}}$. As X is compact, it has a positive injectivity radius $\text{Inj}(X) > 0$. Here we have extended X to a closed manifold to speak about the injectivity radius and the boundary will not cause any problems as $\gamma_1(\alpha_k)(t_k) \in X^{\text{int}}$, for k sufficiently large. Now (3.8) implies that $|t_k| \geq \text{Inj}(X)$ for all k sufficiently large, which is a contradiction as $t_k \rightarrow 0$. Thus, the claim (3.7) follows. The same is true for the family of geodesics $\gamma_2(\alpha)$ for α in a possibly smaller neighborhood of α_0 .

Finally, we claim there is a neighborhood $V \subset \tilde{U}$ of α_0 such that for all $\alpha \in V$, we have

$$\gamma_1(\alpha)(t) = \gamma_2(\alpha)(s) \implies t = s = 0. \tag{3.9}$$

Indeed, otherwise, there exists $\alpha_k \rightarrow \alpha_0$ as $k \rightarrow \infty$, and $t_k \in [-T_1(\alpha_k), T_2(\alpha_k)]$, and $s_k \in [-S_1(\alpha_k), S_2(\alpha_k)]$ such that

$$\gamma_1(\alpha_k)(t_k) = \gamma_2(\alpha_k)(s_k), \tag{3.10}$$

$t_k \neq 0$, and $s_k \neq 0$, for all k . Assuming as we may that $t_k \rightarrow t_0$ and $s_k \rightarrow s_0$ and passing to the limit in (3.10), we obtain that

$$\gamma_1(\alpha_0)(t_0) = \gamma_2(\alpha_0)(s_0),$$

and therefore, as $\gamma_1(\alpha_0)$ and $\gamma_2(\alpha_0)$ are admissible, we get $t_0 = s_0 = 0$. Thus, we get

$$\begin{aligned} \gamma_1(\alpha_k)(t_k) &= \gamma_2(\alpha_k)(s_k) \rightarrow x_0, \\ (\alpha_k)_x &= \gamma_1(\alpha_k)(0) = \gamma_2(\alpha_k)(0) \rightarrow x_0, \end{aligned}$$

as $k \rightarrow \infty$. Note that for k sufficiently large, all the points $\gamma_1(\alpha_k)(t_k)$, $\gamma_1(\alpha_k)(0)$, $\gamma_2(\alpha_k)(s_k)$, $\gamma_2(\alpha_k)(0)$ are in the interior of X . Therefore, $|t_k| \geq \text{Inj}(X)$ and $|s_k| \geq \text{Inj}(X)$ for k sufficiently large, as otherwise, the geodesics $\gamma_1(\alpha_k)$ and $\gamma_2(\alpha_k)$ would intersect at a geodesic ball centered at $(\alpha_k)_x$. This contradicts the fact that $t_k \rightarrow 0$ and $s_k \rightarrow 0$ as $k \rightarrow \infty$, showing the claim. Hence, the pair of geodesics $\gamma_1(\alpha)$, $\gamma_2(\alpha)$ is admissible, for all $\alpha \in V$. \square

4. Analytic families of exponentially accurate Gaussian beam quasimodes

When proving Theorem 1.3 below, we shall need the following consequence of Theorem 1.12.

Corollary 4.1. *Let (X, g) be a compact Riemannian manifold of dimension $n \geq 2$ with smooth boundary, contained in an open real analytic manifold (\tilde{X}, g) of the same dimension with g real analytic in \tilde{X} . Let $\alpha_0 = (x_0, \xi_0) \in S^*X^{\text{int}}$ and let $\gamma_0 : [-T_1, T_2] \rightarrow X$, $0 < T_1, T_2 < \infty$, be a unit speed nontangential geodesic such that $\gamma_0(0) = x_0$, and γ_0 does*

not have self-intersections at x_0 . Let $\gamma(\alpha) : [-T_1(\alpha), T_2(\alpha)] \rightarrow X$, $0 < T_1(\alpha), T_2(\alpha) < \infty$, $\alpha = (\alpha_x, \alpha_\xi) \in \text{neigh}(\alpha_0, S^*X^{\text{int}})$, be a real analytic family of unit speed nontangential geodesics such that $\gamma(\alpha_0) = \gamma_0$, and $\gamma(\alpha)(0) = \alpha_x$. Let $\lambda \in \mathbb{R}$. Then there is a real analytic family of C^∞ functions $v(x, \alpha; h)$ on X , $\alpha \in \text{neigh}(\alpha_0, S^*X^{\text{int}})$, $0 < h \leq 1$, and $C > 0$ such that $\text{supp}(v(\cdot, \alpha; h))$ is confined to a small neighborhood of $\gamma(\alpha)([-T_1(\alpha), T_2(\alpha)])$ for each α , and

$$\|(-h^2\Delta_g - (hs)^2)v(\cdot, \alpha; h)\|_{L^2(X)} = \mathcal{O}(e^{-\frac{1}{Ch}}), \quad \|v(\cdot, \alpha; h)\|_{L^2(X)} \asymp 1, \tag{4.1}$$

as $h \rightarrow 0$, uniformly in α . Here $s = \frac{1}{h} + i\lambda$. The local structure of the family $v(\cdot, \alpha; h)$ in a neighborhood of α_x is as follows:

$$v(x, \alpha; h) = h^{-\frac{(n-1)}{4}} e^{is\varphi(x, \alpha)} a(x, \alpha; h),$$

where $\varphi(x, \alpha)$ is real analytic in (x, α) for $\alpha \in \text{neigh}(\alpha_0, S^*X^{\text{int}})$ and $|x - \alpha_x| < \frac{1}{c}$, $c > 0$, and $a(x, \alpha; h)$ is an elliptic classical analytic symbol near (x_0, α_0) . Furthermore, for t close to 0 and $\alpha \in \text{neigh}(\alpha_0, S^*X^{\text{int}})$, we have

$$\begin{aligned} \varphi(\gamma(\alpha)(t), \alpha) &= t, & \nabla\varphi(\gamma(\alpha)(t), \alpha) &= \dot{\gamma}(\alpha)(t), \\ \text{Im}(\nabla^2\varphi(\gamma(\alpha)(t))) &\geq 0, & \text{Im}(\nabla^2\varphi)|_{\dot{\gamma}(\alpha)(t)^\perp} &> 0. \end{aligned}$$

Proof. The functions $T_1(\alpha)$ and $T_2(\alpha)$ depend continuously on α in a small neighborhood of α_0 , and shrinking the neighborhood further we may assume that $T_1(\alpha), T_2(\alpha)$ are bounded. Let $\varepsilon > 0$ be such that $\gamma(\alpha)(t) \in \widehat{X} \setminus X$ and $\gamma(\alpha)(t)$ has no self-intersection for $t \in [-T_1(\alpha) - 2\varepsilon, -T_1(\alpha)] \cup [T_2(\alpha), T_2(\alpha) + 2\varepsilon]$ for all $\alpha \in \text{neigh}(\alpha_0, S^*X^{\text{int}})$. This choice of ε is possible since $\gamma(\alpha)$ are non-tangential and depend smoothly on α . It follows from [28, Lemma 7.2] that $\gamma(\alpha)|_{[-T_1(\alpha) - \varepsilon, T_2(\alpha) + \varepsilon]}$ self-intersects only at finitely many times $t_j(\alpha)$, $1 \leq j \leq N(\alpha)$, with

$$-T_1(\alpha) < t_1(\alpha) < \dots < t_{N(\alpha)}(\alpha) < T_2(\alpha).$$

First we claim that there is N_0 such that $N(\alpha) \leq N_0 < \infty$ for all α in a small neighborhood of α_0 . This follows by inspection of the arguments in the proof of [28, Lemma 7.2]. Indeed, as explained in [28, Lemma 7.2], if $\gamma(\alpha)(t) = \gamma(\alpha)(s)$ for some $t \neq s$ then $\dot{\gamma}(\alpha)(t) \neq \pm\dot{\gamma}(\alpha)(s)$. Furthermore, if r is smaller than the injectivity radius of some closed manifold containing a fixed neighborhood of $X \cup \gamma(\alpha)([-T_1(\alpha) - 2\varepsilon, T_2(\alpha) + 2\varepsilon])$ for $\alpha \in \text{neigh}(\alpha_0, S^*X^{\text{int}})$, then any two distinct geodesic segments of length $\leq r$ can intersect in at most one point. Partitioning $[-T_1(\alpha) - 2\varepsilon, T_2(\alpha) + 2\varepsilon]$ in disjoint intervals $\{J_k\}_{k=1}^{K(\alpha)}$ of length $\leq r$, we get an injective map

$$\begin{aligned} \{(t, s) \in [-T_1(\alpha) - 2\varepsilon, T_2(\alpha) + 2\varepsilon]^2 : s < t \text{ and } \gamma(\alpha)(t) = \gamma(\alpha)(s)\} \\ \longrightarrow \{(k, l) \in \{1, \dots, K(\alpha)\}^2\}, \\ (t, s) \mapsto (k, l) \text{ such that } t \in J_k, s \in J_l. \end{aligned} \tag{4.2}$$

Since $T_1(\alpha)$ and $T_2(\alpha)$ are bounded for α in a small neighborhood of α_0 , we may assume that $K(\alpha)$ is bounded. Consequently, the cardinality of the set $\{(k, l) \in \{1, \dots, K(\alpha)\}^2\}$ is bounded uniformly in α . The claim follows.

We also set $t_0(\alpha) := -T_1(\alpha) - \varepsilon$ and $t_{N+1}(\alpha) := T_2(\alpha) + \varepsilon$. An inspection of the proof of [12, Lemma 3.5] allows us to conclude that there exists an open cover $\{(U_j(\alpha), \kappa_j(\alpha))\}_{j=0}^{N(\alpha)+1}$ of $\gamma(\alpha)([-T_1(\alpha) - \varepsilon, T_2(\alpha) + \varepsilon])$ consisting of coordinate neighborhoods $U_j(\alpha)$ and real analytic diffeomorphisms $\kappa_j(\alpha)$, depending real analytically on α , such that the following properties hold,

- (i) $\kappa_j(\alpha)(U_j(\alpha)) = I_j \times B$, where I_j are fixed open intervals and $B = B(0, \delta')$ is an open ball in \mathbb{R}^{n-1} . Here $\delta' > 0$ can be taken arbitrarily small and the same for each $U_j(\alpha)$, uniformly for α close to α_0 ,
- (ii) $\kappa_j(\alpha)(\gamma(\alpha)(t)) = (t, 0)$ for each $t \in I_j$,
- (iii) t_j only belongs to I_j and $\overline{I_j} \cap \overline{I_k} = \emptyset$ unless $|j - k| \leq 1$,
- (iv) $\kappa_j(\alpha) = \kappa_k(\alpha)$ on $\kappa_j^{-1}((I_j \cap I_k) \times B)$.

In particular, the open sets $U_j(\alpha)$ are bounded uniformly in α and contain a fixed open set.

Following the proof of Theorem 1.12, and making use of the fact that the geodesics $\gamma(\alpha)$ do not have self-intersections at α_x , for α close to α_0 , we obtain the statement of Corollary 4.1. \square

Remark 4.2. Let us also note that in general the number of self-intersecting times $N(\alpha)$ need not depend continuously on α . To this end, assume that the dimension of the manifold X is > 2 and that the geodesic γ_0 in X has a self-intersection at some point $x_1 \in \gamma_0((-T_1, T_2))$ so that $x_1 = \gamma_0(t_1) = \gamma_0(t_2)$, $t_1 < t_2$. Then one can show that by means of a small perturbation, that one can unwind the loop in the direction orthogonal to the plane spanned by the velocity vectors $\dot{\gamma}(t_1)$ and $\dot{\gamma}(t_2)$.

5. Construction of families of harmonic functions based on Gaussian beam quasimodes

Let (M, g) be a transversally anisotropic manifold of dimension $n \geq 3$ with transversal manifold (M_0, g_0) , and assume that M_0^{int} and $g_0|_{M_0^{\text{int}}}$ are real analytic.

First assume, as we may, that (M, g) is embedded in a compact smooth manifold (N, g) without boundary of the same dimension, and let U be open in N such that $M \subset U$. Our starting point is the following Carleman estimate for $-h^2\Delta$, established in [10].

Proposition 5.1. *Let ϕ be a limiting Carleman weight for $-h^2\Delta$ on U . Then for all $0 < h \ll 1$, we have*

$$h\|u\|_{L^2(N)} \leq C\|e^{\frac{\phi}{h}}(-h^2\Delta)e^{-\frac{\phi}{h}}u\|_{L^2(N)}, \quad C > 0, \tag{5.1}$$

for all $u \in C_0^\infty(M^{\text{int}})$.

Using a standard argument, see [10], we convert the Carleman estimate (5.1) into the following solvability result.

Proposition 5.2. *Let ϕ be a limiting Carleman weight for $-h^2\Delta$ on U . If $h > 0$ is small enough, then for any $v \in L^2(M)$, there is a solution $u \in L^2(M)$ of the equation*

$$e^{\frac{\phi}{h}}(-h^2\Delta)e^{-\frac{\phi}{h}}u = v \quad \text{in } M^{\text{int}},$$

which satisfies

$$\|u\|_{L^2(M)} \leq \frac{C}{h}\|v\|_{L^2(M)}.$$

Now as $M \subset\subset \mathbb{R} \times M_0^{\text{int}}$, there is a compact Riemannian manifold \widetilde{M}_0 of dimension $n-1$ with smooth boundary such that $M \subset\subset \mathbb{R} \times \widetilde{M}_0 \subset\subset \mathbb{R} \times M_0^{\text{int}}$. Note that $(M_0^{\text{int}}, g_0|_{M_0^{\text{int}}})$ is an open real analytic manifold with real analytic metric, and we can use Corollary 4.1 to construct a real analytic family of Gaussian beam quasimodes along nontangential geodesics on \widetilde{M}_0 .

Let us write $x = (x_1, x')$ for local coordinates in $\mathbb{R} \times \widetilde{M}_0$. Let

$$s = \frac{1}{h} + i\lambda, \quad \lambda \in \mathbb{R}, \quad \lambda \text{ fixed.}$$

Note by [10, Lemma 2.9] that $\phi(x) = \pm x_1$ is a limiting Carleman weight for $-h^2\Delta$ on U . We are interested in finding harmonic functions,

$$-\Delta_g u = 0 \quad \text{in } M^{\text{int}}, \tag{5.2}$$

having the form

$$u = u(x, \alpha; h) = e^{-sx_1}(v(x', \alpha; h) + r(x, \alpha; h)),$$

where $v = v(x', \alpha; h)$ is the Gaussian beam quasimode constructed in Corollary 4.1 on the transversal manifold \widetilde{M}_0 , associated to a nontangential unit speed geodesic $\gamma(\alpha)$ on \widetilde{M}_0 depending analytically on $\alpha \in \text{neigh}(\alpha_0, S^*\widetilde{M}_0^{\text{int}})$, and r is a remainder term. Thus, u is a solution of (5.2) provided that r solves

$$e^{\frac{x_1}{h}}(-h^2\Delta_g)e^{-\frac{x_1}{h}}(e^{-i\lambda x_1}r) = -e^{-i\lambda x_1}(-h^2\Delta_{g_0} - (hs)^2)v(x', \alpha; h). \tag{5.3}$$

Proposition 5.2 and Corollary 4.1 imply that there is $r = r(\cdot; \alpha; h) \in L^2(M)$ solving (5.3) such that

$$\|r\|_{L^2(M)} = \mathcal{O}(e^{-\frac{1}{Ch}}), \quad C > 0,$$

as $h \rightarrow 0$, uniformly in $\alpha \in \text{neigh}(\alpha_0, S^*\widetilde{M}_0^{\text{int}})$.

To summarize, we have the following result.

Proposition 5.3. *Let $s = \frac{1}{h} + i\lambda$ with $\lambda \in \mathbb{R}$ being fixed. For all $h > 0$ small enough, there are families of harmonic functions $u_1, u_2 \in L^2(M)$, i.e. $-\Delta_g u_j = 0$ in M^{int} , having the form*

$$\begin{aligned} u_1(x, \alpha; h) &= e^{-sx_1}(v(x', \alpha; h) + r_1(x, \alpha; h)), \\ u_2(x, \alpha; h) &= e^{sx_1}(v(x', \alpha; h) + r_2(x, \alpha; h)), \end{aligned}$$

where $v = v(\cdot, \alpha; h)$ is the family of Gaussian beam quasimodes constructed in Corollary 4.1 on \widetilde{M}_0 , and $r \in L^2(M)$ is such that $\|r_j\|_{L^2(M)} = \mathcal{O}(e^{-\frac{1}{Ch}})$, $C > 0$, as $h \rightarrow 0$, uniformly in $\alpha \in \text{neigh}(\alpha_0, S^*\widetilde{M}_0^{\text{int}})$, $j = 1, 2$.

6. Proofs of Theorem 1.3 and Theorem 1.4

6.1. Some facts about analytic wave front sets

We shall rely on the following characterization of the analytic wave front set, which we recall from [49, Definition 6.1] for the convenience of the reader. In our applications, we have $m = n - 1$.

Definition 6.1. Let $\alpha_0 = (x_0, \xi_0) \in T^*\mathbb{R}^m \setminus \{0\}$, and let $\varphi(x, \alpha)$, $x \in \mathbb{R}^m$, $\alpha = (\alpha_x, \alpha_\xi) \in T^*\mathbb{R}^m \setminus \{0\}$, be analytic defined in a neighborhood of (x_0, α_0) such that

$$\varphi(x, \alpha)|_{x=\alpha_x} = 0, \quad \varphi'_x(x, \alpha)|_{x=\alpha_x} = \alpha_\xi, \tag{6.1}$$

and

$$\text{Im } \varphi(x, \alpha) \geq C_0|x - \alpha_x|^2, \quad x, \alpha \text{ real}, \tag{6.2}$$

for some $C_0 > 0$. Let $a(x, \alpha; h)$ be an elliptic classical analytic symbol defined in a neighborhood of (x_0, α_0) , and let $u \in \mathcal{D}'(X)$, where $X \subset \mathbb{R}^m$ is an open set containing x_0 . We have $\alpha_0 \notin WF_a(u)$ if and only if there is a real neighborhood U of α_0 and $C > 0$ such that

$$\sup_{\alpha \in U} |Tu(\alpha; h)| \leq Ce^{-\frac{1}{Ch}}, \tag{6.3}$$

for $0 < h \leq 1$, where

$$Tu(\alpha; h) = \int e^{\frac{i\varphi(x, \alpha)}{h}} a(x, \alpha; h) \chi(x) \overline{u(x)} dx,$$

and $\chi \in C_0^\infty(X)$ is supported in a small neighborhood of x_0 and $\chi = 1$ near x_0 .

Remark 6.2. It is established in [49, Proposition 6.2] that the condition (6.3) is independent of the choice of χ , a , and φ .

Remark 6.3. Assume that φ , a , and u satisfy the same conditions as in Definition 6.1, and let $\psi \in C_0^\infty(\mathbb{R}^n)$ be supported in a small neighborhood of 0 and $\psi = 1$ near 0. We have $\alpha_0 \notin WF_a(u)$ if and only if there is a real neighborhood \tilde{U} of α_0 and $\tilde{C} > 0$ such that

$$\sup_{\alpha \in \tilde{U}} |\tilde{T}u(\alpha; h)| \leq \tilde{C}e^{-\frac{1}{Ch}}, \tag{6.4}$$

for $0 < h \leq 1$, where

$$\tilde{T}u(\alpha; h) = \int e^{\frac{i\varphi(x,\alpha)}{h}} a(x, \alpha; h) \psi(x - \alpha_x) \overline{u(x)} dx.$$

The condition (6.4) is independent of the choice of ψ .

Remark 6.4. The condition (6.1) in Definition 6.1 and Remark 6.3 can be replaced by the following

$$\varphi(x, \alpha)|_{x=\alpha_x} = f(\alpha) \text{ real valued, } \quad \varphi'_x(x, \alpha)|_{x=\alpha_x} = t_0\alpha_\xi, \tag{6.5}$$

for some fixed $t_0 > 0$. Indeed, we apply Definition 6.1 and Remark 6.3 with $\varphi(x, \alpha)$ replaced by $\frac{1}{t_0}(\varphi(x, \alpha) - f(\alpha))$ and with h replaced by h/t_0 .

Remark 6.5. Since the wave front set $WF_a(u)$ is conic, we may restrict the attention in Definition 6.1 to $\xi_0 \in \mathbb{R}^m$ such that $|\xi_0| = 1$.

6.2. Proof of Theorem 1.3

Let $\alpha_0 = (x'_0, \xi'_0) \in S^*M_0^{\text{int}}$ be generated by an admissible pair of geodesics $\gamma_1(\alpha_0) : [-T_1(\alpha_0), T_2(\alpha_0)] \rightarrow M_0$ and $\gamma_2(\alpha_0) : [-S_1(\alpha_0), S_2(\alpha_0)] \rightarrow M_0$. As $M \subset\subset \mathbb{R} \times M_0^{\text{int}}$, there is a compact Riemannian manifold \tilde{M}_0 of dimension $n - 1$ with smooth boundary such that $M \subset\subset \mathbb{R} \times \tilde{M}_0 \subset\subset \mathbb{R} \times M_0^{\text{int}}$, and $x'_0 \in \tilde{M}_0^{\text{int}}$. Furthermore, we can choose \tilde{M}_0 so that the geodesics $\gamma_1(\alpha_0)$ and $\gamma_2(\alpha_0)$ are nontangential on \tilde{M}_0 , and hence, $\gamma_1(\alpha_0)$ and $\gamma_2(\alpha_0)$ are admissible on \tilde{M}_0 .

Then by Lemma 3.2, there exists a neighborhood V of α_0 in $S^*\tilde{M}_0^{\text{int}}$ such that every point $\alpha = (\alpha_{x'}, \alpha_{\xi'}) \in V$ is generated by an admissible pair of geodesics $\gamma_1(\alpha) : [-T_1(\alpha), T_2(\alpha)] \rightarrow \tilde{M}_0$ and $\gamma_2(\alpha) : [-S_1(\alpha), S_2(\alpha)] \rightarrow \tilde{M}_0$, which depend real-analytically on α . Thus, for all $\alpha \in V$, we have

$$\gamma_1(\alpha)(0) = \gamma_2(\alpha)(0) = \alpha_{x'}, \tag{6.6}$$

$$\dot{\gamma}_1(\alpha)(0) + \dot{\gamma}_2(\alpha)(0) = t_0\alpha_{\xi'}, \tag{6.7}$$

for some $0 < t_0 < 2$ fixed, $\gamma_1(\alpha)$, $\gamma_2(\alpha)$ do not have self-intersections at $\alpha_{x'}$, and $\alpha_{x'}$ is the only point where $\gamma_1(\alpha)$ and $\gamma_2(\alpha)$ intersect, for all $\alpha \in V$.

Let $s_1 = \frac{1}{h} + i\lambda$ and $s_2 = \frac{1}{h}$, where $\lambda \in \mathbb{R}$. Applying Corollary 4.1, we get $v_j(x', \alpha; h)$, $j = 1, 2$, Gaussian beam quasimodes on \widetilde{M}_0 , associated to $\widetilde{\gamma}_j(\alpha)$ on \widetilde{M}_0 , depending real analytically on $\alpha \in V$ such that

$$\|v_j(\cdot, \alpha; h)\|_{L^2(M)} \asymp 1, \quad \|(-h^2\Delta_{g_0} - (hs)^2)v_j(\cdot, \alpha; h)\|_{L^2(M)} = \mathcal{O}(e^{-\frac{1}{C\hbar}}), \quad (6.8)$$

as $h \rightarrow 0$, for some $C > 0$, uniformly in $\alpha \in V$.

An application of Proposition 5.3 gives harmonic functions on M having the form

$$\begin{aligned} u_1(x, \alpha; h) &= e^{-s_1x_1}(v_1(x', \alpha; h) + r_1(x, \alpha; h)), \\ u_2(x, \alpha; h) &= e^{s_2x_1}(v_2(x', \alpha; h) + r_2(x, \alpha; h)), \end{aligned} \quad (6.9)$$

where

$$\|r_j\|_{L^2(M)} = \mathcal{O}(e^{-\frac{1}{C\hbar}}), \quad C > 0, \quad (6.10)$$

as $h \rightarrow 0$, uniformly in $\alpha \in V$.

Substituting the harmonic functions u_1 and u_2 given by (6.9) into (1.2), we get

$$\int_M f e^{-i\lambda x_1} (v_1(x', \alpha; h) + r_1)(v_2(x', \alpha; h) + r_2) dV_g = 0. \quad (6.11)$$

Using (6.10) and (6.8), we see that

$$\int_M f e^{-i\lambda x_1} v_1(x', \alpha; h)v_2(x', \alpha; h) dV_g = \mathcal{O}(e^{-\frac{1}{C\hbar}}), \quad C > 0, \quad (6.12)$$

uniformly in $\alpha \in V$. Let us extend $f \in L^\infty(M)$ by zero to $(\mathbb{R} \times M_0) \setminus M$ and set

$$\widehat{f}(\lambda, x') = \int_{-\infty}^{\infty} e^{-i\lambda x_1} f(x_1, x') dx_1$$

for the Fourier transform with respect to x_1 . Using the fact that $dV_g = dx_1 dV_{g_0}$, we obtain from (6.12) that

$$\int_{\widetilde{M}_0} \widehat{f}(\lambda, x') v_1(x', \alpha; h)v_2(x', \alpha; h) dV_{g_0} = \mathcal{O}(e^{-\frac{1}{C\hbar}}), \quad C > 0, \quad (6.13)$$

uniformly in $\alpha \in V$. Recalling that the geodesics $\gamma_1(\alpha)$ and $\gamma_2(\alpha)$ intersect at $\alpha_{x'}$ only and that

$$\text{supp } (v_j(\cdot, \alpha; h)) \subset \text{small neigh}(\gamma_j(\alpha)), \quad j = 1, 2,$$

we conclude from (6.13) that

$$\int_{\text{neigh}(\alpha_{x'}, M_0)} \widehat{f}(\lambda, x') v_1(x', \alpha; h) v_2(x', \alpha; h) \sqrt{g_0(x')} dx' = \mathcal{O}(e^{-\frac{1}{Ch}}), \tag{6.14}$$

uniformly in $\alpha \in V$. Recalling that the geodesics $\gamma_1(\alpha)$, $\gamma_2(\alpha)$ do not have self-intersections at $\alpha_{x'}$, by Corollary 4.1, we have in a small neighborhood of $\alpha_{x'}$,

$$\begin{aligned} v_1(x', \alpha; h) &= h^{-\frac{(n-2)}{4}} e^{is_1\varphi_1(x', \alpha)} a_1(x', \alpha; h), \\ v_2(x', \alpha; h) &= h^{-\frac{(n-2)}{4}} e^{is_2\varphi_2(x', \alpha)} a_2(x', \alpha; h). \end{aligned} \tag{6.15}$$

Here $\varphi_j(x', \alpha)$ are real analytic in (x', α) in a region of the form $\alpha \in V$ and $|x' - \alpha_{x'}| < 1/c$, which is an open neighborhood of (x'_0, α_0) . Furthermore, $a_j(x', \alpha; h)$ are elliptic classical analytic symbols in a neighborhood of (x'_0, α_0) , $j = 1, 2$. It follows that the neighborhood of $\alpha_{x'}$ occurring as the domain of integration in (6.14) can be taken to be fixed and independent of α . We also have for the geodesic parameters t and s near 0 that

$$\begin{aligned} (\varphi_1)'_{x'}(\gamma_1(\alpha)(t), \alpha) &= \dot{\gamma}_1(\alpha)(t), \quad \text{Im}((\varphi_1)''_{x'x'}(\gamma_1(\alpha)(t), \alpha)) \geq 0, \\ \text{Im}((\varphi_1)''_{x'x'}(\gamma_1(\alpha)(t), \alpha)|_{[\dot{\gamma}_1(\alpha)(t)]^\perp}) &> 0, \end{aligned} \tag{6.16}$$

and

$$\begin{aligned} (\varphi_2)'_{x'}(\gamma_2(\alpha)(s), \alpha) &= \dot{\gamma}_2(\alpha)(s), \quad \text{Im}((\varphi_2)''_{x'x'}(\gamma_2(\alpha)(s), \alpha)) \geq 0, \\ \text{Im}((\varphi_2)''_{x'x'}(\gamma_2(\alpha)(s), \alpha)|_{[\dot{\gamma}_2(\alpha)(s)]^\perp}) &> 0. \end{aligned} \tag{6.17}$$

Now substituting (6.15) into (6.14), we see that

$$\int_{\text{neigh}(\alpha_{x'}, M_0)} e^{\frac{i\varphi(x', \alpha)}{h}} \widehat{f}(\lambda, x') a(x', \alpha; h) dx' = \mathcal{O}(e^{-\frac{1}{Ch}}), \quad h \rightarrow 0, \tag{6.18}$$

uniformly in $\alpha \in V$. Here

$$\varphi(x', \alpha) = \varphi_1(x', \alpha) + \varphi_2(x', \alpha) \tag{6.19}$$

is analytic in a neighborhood of (x'_0, α_0) , and

$$a(x', \alpha; h) = e^{-\lambda\varphi_1(x', \alpha)} a_1(x', \alpha; h) a_2(x', \alpha; h) \sqrt{g_0(x')}$$

is an elliptic classical analytic symbol in a neighborhood of (x'_0, α_0) , since the product of two classical analytic symbols is a classical analytic symbol.

We now claim that the phase function $\varphi(x', \alpha)$ in (6.19) satisfies the conditions (6.5) and (6.2). First, in view of (6.6) and (1.5), we have

$$\varphi(x', \alpha)|_{x'=\alpha_{x'}} = \varphi_1(\gamma_1(\alpha)(0), \alpha) + \varphi_2(\gamma_2(\alpha)(0), \alpha) = 0. \tag{6.20}$$

Using (6.16), (6.17), and (6.7), we get

$$\begin{aligned} \varphi'_{x'}(x', \alpha)|_{x'=\alpha_{x'}} &= (\varphi_1)'_{x'}(\gamma_1(\alpha)(0), \alpha) + (\varphi_2)'_{x'}(\gamma_2(\alpha)(0), \alpha) \\ &= \dot{\gamma}_1(\alpha)(0) + \dot{\gamma}_2(\alpha)(0) = t_0\alpha_{\xi'}. \end{aligned} \tag{6.21}$$

It follows from (6.20) and (6.21) that the condition (6.5) holds. Let us now check the condition (6.2). To this end, Taylor expanding $\varphi(x', \alpha)$ at $x' = \alpha_{x'}$, we get

$$\begin{aligned} \varphi(x', \alpha) &= \varphi(\alpha_{x'}, \alpha) + \varphi'_{x'}(\alpha_{x'}, \alpha) \cdot (x' - \alpha_{x'}) \\ &\quad + \frac{1}{2}\varphi''_{x'x'}(\alpha_{x'}, \alpha)(x' - \alpha_{x'}) \cdot (x' - \alpha_{x'}) + \mathcal{O}(|x' - \alpha_{x'}|^3), \end{aligned}$$

and therefore, in view of (6.20) and (6.21), when x' and α are real, we see that

$$\text{Im } \varphi(x', \alpha) = \frac{1}{2}\text{Im } \varphi''_{x'x'}(\alpha_{x'}, \alpha)(x' - \alpha_{x'}) \cdot (x' - \alpha_{x'}) + \mathcal{O}(|x' - \alpha_{x'}|^3).$$

Hence, the condition (6.2) is equivalent the following condition,

$$\text{Im } \varphi''_{x'x'}(\alpha_{x'}, \alpha) > 0. \tag{6.22}$$

Using (6.16), (6.17), and the fact that the vectors $\dot{\gamma}_1(\alpha)(0)$ and $\dot{\gamma}_2(\alpha)(0)$ are not parallel, we have

$$\text{Im } \varphi''_{x'x'}(\alpha_{x'}, \alpha) = \text{Im } (\varphi_1)''_{x'x'}(\gamma_1(\alpha)(0), \alpha) + \text{Im } (\varphi_2)''_{x'x'}(\gamma_2(\alpha)(0), \alpha) > 0,$$

showing (6.22). Thus, by Remarks 6.3 and 6.4, in view of (6.18), we get $\alpha_0 \notin WF_a(\widehat{f}(\lambda, \cdot)) = WF_a(\widehat{f}(-\lambda, \cdot))$ for all $\lambda \in \mathbb{R}$. Noting that if (1.2) holds for f , it also holds for \overline{f} and $\lambda \in \mathbb{R}$ is arbitrary, we get $\alpha_0 \notin WF_a(\widehat{f}(\lambda, \cdot))$ for all $\lambda \in \mathbb{R}$. This completes the proof of Theorem 1.3.

6.3. Proof of Theorem 1.4

Now since every point $(x'_0, \xi'_0) \in S^*M_0^{\text{int}}$ is generated by an admissible pair of geodesics, by Theorem 1.3, we get $\widehat{f}(\lambda, \cdot)$ is real-analytic in M_0^{int} for all $\lambda \in \mathbb{R}$. The fact that $\widehat{f}(\lambda, \cdot)$ has a compact support in M_0^{int} and that M_0 is connected implies that $\widehat{f}(\lambda, \cdot) = 0$ for all $\lambda \in \mathbb{R}$, and therefore, $f = 0$. This completes the proof of Theorem 1.4.

Appendix A. Discussion related to Example 1.7

Let $M_0 = \mathbb{S}^1 \times [0, a]$, with $a > 0$, be a cylinder with its usual flat metric g_0 . The purpose of this Appendix is to show that every point $(x_0, \xi_0) \in S^*M_0^{\text{int}}$ is generated by an admissible pair of geodesics.

We have

$$T^*(\mathbb{S}^1 \times (0, a)) \simeq T^*\mathbb{S}^1 \times T^*(0, a) \simeq (\mathbb{S}^1 \times \mathbb{R}) \times ((0, a) \times \mathbb{R}) \simeq (\mathbb{S}^1 \times (0, a)) \times \mathbb{R}^2,$$

and therefore, we may identify $S^*_{x_0}M_0^{\text{int}}$ with the unit circle \mathbb{S}^1 in $\mathbb{R}^2 \simeq \mathbb{C}$.

Given $\mathbb{S}^1 \ni \xi_0 = (\xi_{01}, \xi_{02}) \simeq \xi_{01} + i\xi_{02}$, we set

$$\xi_1 = e^{i\alpha}(\xi_{01} + i\xi_{02}) \in \mathbb{S}^1, \quad \xi_2 = e^{-i\alpha}(\xi_{01} + i\xi_{02}) \in \mathbb{S}^1, \tag{A.1}$$

with $\alpha \in (0, 2\pi)$ to be chosen. The geodesics γ_1 and γ_2 on M_0 such that $\gamma_j(0) = x_0$ and $\dot{\gamma}_j(0) = \xi_j$, $j = 1, 2$, are given by

$$\begin{aligned} \gamma_1(t) &= (x_{01} + \xi_{11}t, x_{02} + \xi_{12}t) \in \mathbb{R}/2\pi\mathbb{Z} \times [0, a], \\ \gamma_2(s) &= (x_{01} + \xi_{21}s, x_{02} + \xi_{22}s) \in \mathbb{R}/2\pi\mathbb{Z} \times [0, a]. \end{aligned}$$

The geodesics γ_1 and γ_2 are nontangential provided that

$$\begin{aligned} \xi_{12} &= \text{Im}(e^{i\alpha}(\xi_{01} + i\xi_{02})) = \xi_{02} \cos \alpha + \xi_{01} \sin \alpha \neq 0, \\ \xi_{22} &= \text{Im}(e^{-i\alpha}(\xi_{01} + i\xi_{02})) = \xi_{02} \cos \alpha - \xi_{01} \sin \alpha \neq 0. \end{aligned} \tag{A.2}$$

Note that if γ_1 and γ_2 are nontangential then they do not have self-intersections.

We have in view of (A.1),

$$\xi_1 + \xi_2 = (2 \cos \alpha)\xi_0, \tag{A.3}$$

and therefore, the property (ii) of Definition 1.2 follows with $t_0 = 2 \cos \alpha$, provided that

$$0 < \cos \alpha < 1. \tag{A.4}$$

Note that γ_1 and γ_2 intersect each other if there exist t and s such that

$$\xi_{11}t - \xi_{21}s \in 2\pi\mathbb{Z}, \quad \xi_{12}t = \xi_{22}s. \tag{A.5}$$

Now if we choose α so that

$$|\xi_{11}t - \xi_{21}s| < 2\pi, \quad \xi_{12}t = \xi_{22}s, \tag{A.6}$$

then (A.5) implies that $\xi_1 t = \xi_2 s$, and therefore, $|t| = |s|$. In view of (A.3) and (A.4), we get $t = s = 0$, and hence, x_0 is the only point of intersections of γ_1 and γ_2 .

To achieve (A.6), assuming that (A.2) holds, we estimate

$$\begin{aligned} |\xi_{11}t - \xi_{21}s| &= \frac{|t|}{|\xi_{22}|} |\xi_{22}\xi_{11} - \xi_{21}\xi_{12}| = \frac{|t|}{|\xi_{22}|} |\operatorname{Im}(\bar{\xi}_1 \xi_2)| = \frac{|t|}{|\xi_{22}|} |\sin(2\alpha)| \\ &\leq \frac{a}{|\xi_{12}||\xi_{22}|} |\sin(2\alpha)| = \frac{a}{|\xi_{02}^2 - \sin^2 \alpha|} |\sin(2\alpha)|, \end{aligned}$$

where we use that $0 \leq x_{02} + \xi_{12}t \leq a$, $0 \leq x_{02} \leq a$, and (A.2). Thus, to prove the result, we have to choose $\alpha \in (0, 2\pi)$ so that (A.2), (A.4), and

$$\frac{a}{|\xi_{02}^2 - \sin^2 \alpha|} |\sin(2\alpha)| < 2\pi \quad (\text{A.7})$$

hold. In doing so let us first consider the case when $\xi_{02} \neq 0$. In this case choosing $\alpha > 0$ small enough, depending on a and ξ_{02} , we see that (A.2), (A.4), and (A.7) hold. When $\xi_{02} = 0$, we choose $\alpha = \frac{\pi}{2} - \beta$ with $\beta > 0$ small enough, depending on a . Then (A.7) becomes

$$\frac{a}{|\cos^2 \beta|} |\sin(2\beta)| < 2\pi,$$

which together with (A.2), (A.4) hold for such small β . This completes the proof that every point of $S^*M_0^{\text{int}}$ is generated by an admissible pair of geodesics.

References

- [1] V. Babich, Construction of Gaussian beams with exponentially small residual (Russian), *Zap. Nauč. Semin. POMI* 203 (1992), *Mat. Vopr. Teor. Rasprostr.* Voln. 22, 17–20, 173–174, translation in *J. Math. Sci.* 79 (1996), no. 4, 1169–1171.
- [2] V. Babich, V. Buldyrev, *Short-Wavelength Diffraction Theory. Asymptotic Methods*, Springer Series on Wave Phenomena, vol. 4, Springer-Verlag, Berlin, 1991.
- [3] M. Belishev, Geometrization of rings as a method for solving inverse problems, in: *Sobolev Spaces in Mathematics. III*, *Int. Math. Ser. (N.Y.)*, vol. 10, Springer, New York, 2009, pp. 5–24.
- [4] M. Belishev, D. Korikov, On the EIT problem for nonorientable surfaces, *J. Inverse Ill-Posed Probl.* 29 (3) (2021) 339–349.
- [5] A.P. Calderón, On an inverse boundary value problem, in: *Seminar on Numerical Analysis and Its Applications to Continuum Physics*, Rio de Janeiro, 1980, *Soc. Brasil. Mat.*, Rio de Janeiro, 1980, pp. 65–73, Reprinted in *Comput. Appl. Math.* 25 (2016), no. 2–3, 133–138.
- [6] M. Dahl, A geometric interpretation of the complex tensor Riccati equation for Gaussian beams, *J. Nonlinear Math. Phys.* 14 (1) (2007) 95–111.
- [7] T. Daudé, N. Kamran, F. Nicoleau, On the hidden mechanism behind non-uniqueness for the anisotropic Calderón problem with data on disjoint sets, *Ann. Henri Poincaré* 20 (3) (2019) 859–887.
- [8] T. Daudé, N. Kamran, F. Nicoleau, On nonuniqueness for the anisotropic Calderón problem with partial data, *Forum Math. Sigma* 8 (2020), 17 pp.
- [9] N. Dencker, J. Sjöstrand, M. Zworski, Pseudospectra of semiclassical (pseudo-) differential operators, *Commun. Pure Appl. Math.* 57 (3) (2004) 384–415.
- [10] D. Dos Santos Ferreira, C. Kenig, M. Salo, G. Uhlmann, Limiting Carleman weights and anisotropic inverse problems, *Invent. Math.* 178 (1) (2009) 119–171.
- [11] D. Dos Santos Ferreira, C. Kenig, J. Sjöstrand, G. Uhlmann, On the linearized local Calderón problem, *Math. Res. Lett.* 16 (6) (2009) 955–970.
- [12] D. Dos Santos Ferreira, Y. Kurylev, M. Lassas, M. Salo, The Calderón problem in transversally anisotropic geometries, *J. Eur. Math. Soc.* 18 (11) (2016) 2579–2626.

- [13] D. Dos Santos Ferreira, Y. Kurylev, M. Lassas, T. Liimatainen, M. Salo, The linearized Calderón problem in transversally anisotropic geometries, *Int. Math. Res. Not.* (2022), <https://doi.org/10.1093/imrn/rny234>.
- [14] A. Feizmohammadi, L. Oksanen, An inverse problem for a semi-linear elliptic equation in Riemannian geometries, *J. Differ. Equ.* 269 (6) (2020) 4683–4719.
- [15] A. Feizmohammadi, T. Liimatainen, Y.-H. Lin, An inverse problem for a semilinear elliptic equation on conformally transversally anisotropic manifolds, preprint, <https://arxiv.org/abs/2112.08305>, 2021.
- [16] F. Forstneric, Actions of $(\mathbb{R}, +)$ and $(\mathbb{C}, +)$ on complex manifolds, *Math. Z.* 223 (1) (1996) 123–153.
- [17] A. Greenleaf, M. Lassas, G. Uhlmann, On nonuniqueness for Calderón’s inverse problem, *Math. Res. Lett.* 10 (5–6) (2003) 685–693.
- [18] A. Grigis, J. Sjöstrand, *Microlocal Analysis for Differential Operators. An Introduction*, London Mathematical Society Lecture Note Series, vol. 196, Cambridge University Press, Cambridge, 1994.
- [19] C. Guillarmou, A. Sá Barreto, Inverse problems for Einstein manifolds, *Inverse Probl. Imaging* 3 (2009) 1–15.
- [20] C. Guillarmou, M. Salo, L. Tzou, The linearized Calderón problem on complex manifolds, *Acta Math. Sin. Engl. Ser.* 35 (6) (2019) 1043–1056.
- [21] C. Guillarmou, L. Tzou, Calderón inverse problem with partial data on Riemann surfaces, *Duke Math. J.* 158 (1) (2011) 83–120.
- [22] P. Hintz, G. Uhlmann, J. Zhai, An inverse boundary value problem for a semilinear wave equation on Lorentzian manifolds, *Int. Math. Res. Not.* (2021) rnab088, <https://doi.org/10.1093/imrn/rnab088>.
- [23] L. Hörmander, *Linear Partial Differential Operators*, Springer-Verlag, New York, Berlin-Göttingen-Heidelberg, 1963.
- [24] L. Hörmander, *Lectures on Nonlinear Hyperbolic Differential Equations*, Mathematics & Applications, vol. 26, Springer-Verlag, Berlin, 1997.
- [25] L. Hörmander, *The Analysis of Linear Partial Differential Operators, I*, Springer-Verlag, Berlin, 2003.
- [26] L. Hörmander, *The Analysis of Linear Partial Differential Operators, III*, Springer-Verlag, Berlin, 2003.
- [27] H. Isozaki, Inverse spectral problems on hyperbolic manifolds and their applications to inverse boundary value problems in Euclidean space, *Am. J. Math.* 126 (6) (2004) 1261–1313.
- [28] C. Kenig, M. Salo, The Calderón problem with partial data on manifolds and applications, *Anal. PDE* 6 (8) (2013) 2003–2048.
- [29] R. Kohn, M. Vogelius, Determining conductivity by boundary measurements, *Commun. Pure Appl. Math.* 37 (3) (1984) 289–298.
- [30] K. Krupchyk, G. Uhlmann, Inverse problems for magnetic Schrödinger operators in transversally anisotropic geometries, *Commun. Math. Phys.* 361 (2) (2018) 525–582.
- [31] K. Krupchyk, G. Uhlmann, Inverse problems for nonlinear magnetic Schrödinger equations on conformally transversally anisotropic manifolds, *Anal. PDE* (2022), in press.
- [32] K. Krupchyk, G. Uhlmann, L. Yan, A remark on inverse problems for nonlinear magnetic Schrödinger equations on complex manifolds, *Proc. Am. Math. Soc.* (2022), <https://doi.org/10.1090/proc/16060>.
- [33] A. Katchalov, M. Lassas, *Gaussian Beams and Inverse Boundary Spectral Problems. New Analytic and Geometric Methods in Inverse Problems*, Springer, Berlin, 2004, pp. 127–163.
- [34] M. Lassas, T. Liimatainen, Y.H. Lin, M. Salo, Inverse problems for elliptic equations with power type nonlinearities, *J. Math. Pures Appl.* (9) 145 (2021) 44–82.
- [35] M. Lassas, T. Liimatainen, M. Salo, The Poisson embedding approach to the Calderón problem, *Math. Ann.* 377 (1–2) (2020) 19–67.
- [36] M. Lassas, T. Liimatainen, L. Potenciano-Machado, T. Tyni, Stability estimates for inverse problems for semi-linear wave equations on Lorentzian manifolds, arXiv:2106.12257, 2021.
- [37] M. Lassas, M. Taylor, G. Uhlmann, The Dirichlet-to-Neumann map for complete Riemannian manifolds with boundary, *Commun. Anal. Geom.* 11 (2003) 207–222.
- [38] M. Lassas, G. Uhlmann, On determining a Riemannian manifold from the Dirichlet-to-Neumann map, *Ann. Sci. Éc. Norm. Supér.* (4) 34 (5) (2001) 771–787.
- [39] J.M. Lee, *Introduction to Smooth Manifolds*, Springer, New York, NY, 2013.
- [40] J.M. Lee, G. Uhlmann, Determining anisotropic real analytic conductivities by boundary measurements, *Commun. Pure Appl. Math.* 42 (8) (1989) 1097–1112.
- [41] Y. Ma, L. Tzou, Semilinear Calderón problem on Stein manifolds with Kähler metric, *Bull. Aust. Math. Soc.* 103 (1) (2021) 132–144.

- [42] A. Martinez, *An Introduction to Semiclassical and Microlocal Analysis*, Universitext, Springer-Verlag, New York, 2002.
- [43] L. Oksanen, M. Salo, P. Stefanov, G. Uhlmann, Inverse problems for real principal type operators, preprint, <https://arxiv.org/abs/2001.07599>, 2020.
- [44] J. Ralston, Approximate eigenfunctions of the Laplacian, *J. Differ. Geom.* 12 (1) (1977) 87–100.
- [45] J. Ralston, Gaussian beams and the propagation of singularities, in: *Studies in Partial Differential Equations*, in: *MAA Stud. Math.*, vol. 23, Math. Assoc. America, Washington, DC, 1982, pp. 206–248.
- [46] L. Robbiano, C. Zuily, Analytic theory for the quadratic scattering wave front set and application to the Schrödinger equation, *Astérisque* 283 (2002), vi+128 pp.
- [47] M. Salo, L. Tzou, Inverse problems for semilinear elliptic PDE with measurements at a single point, preprint, <https://arxiv.org/abs/2202.05290>, 2022.
- [48] J. Sjöstrand, Applications of Fourier distributions with complex phase functions, in: *Lecture Notes in Math.*, vol. 459, Springer, Berlin, 1975, pp. 256–282.
- [49] J. Sjöstrand, Singularités Analytiques Microlocales, (French) [Microlocal Analytic Singularities], *Astérisque*, vol. 95, Soc. Math. France, Paris, 1982, pp. 1–166.
- [50] J. Sjöstrand, G. Uhlmann, Local analytic regularity in the linearized Calderón problem, *Anal. PDE* 9 (3) (2016) 515–544.
- [51] P. Stefanov, G. Uhlmann, Integral geometry of tensor fields on a class of non-simple Riemannian manifolds, *Am. J. Math.* 130 (2008) 239–268.
- [52] J. Sylvester, G. Uhlmann, A global uniqueness theorem for an inverse boundary value problem, *Ann. Math. (2)* 125 (1) (1987) 153–169.
- [53] G. Uhlmann, Electrical impedance tomography and Calderón’s problem, *Inverse Probl.* 25 (12) (2009) 123011, 39 pp.
- [54] S. Zelditch, Local and global analysis of eigenfunctions on Riemannian manifolds, in: *Handbook of Geometric Analysis*. No. 1, in: *Adv. Lect. Math.*, vol. 7, Int. Press, Somerville, MA, 2008, pp. 545–658.