

This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Antunes Lima, Rodrigo; Cunha Soares, Fernanda; van Poppel, Mireille; Savinainen, Saija; Mäntyselkä, Aino; Haapala, Eero A.; Lakka, Timo

Title: Determinants of Cognitive Performance in Children and Adolescents : A Populational Longitudinal Study

Year: 2022

Version: Published version

Copyright: © 2022 by the authors

Rights: CC BY 4.0

Rights url: <https://creativecommons.org/licenses/by/4.0/>

Please cite the original version:

Antunes Lima, R., Cunha Soares, F., van Poppel, M., Savinainen, S., Mäntyselkä, A., Haapala, E. A., & Lakka, T. (2022). Determinants of Cognitive Performance in Children and Adolescents : A Populational Longitudinal Study. *International Journal of Environmental Research and Public Health*, 19(15), Article 8955. <https://doi.org/10.3390/ijerph19158955>



Article

Determinants of Cognitive Performance in Children and Adolescents: A Populational Longitudinal Study

Rodrigo Antunes Lima ^{1,*}, Fernanda Cunha Soares ², Mireille van Poppel ³, Saija Savinainen ^{4,5},
Aino Mäntyselkä ⁵, Eero A. Haapala ^{4,6,†} and Timo Lakka ^{4,7,8,†}

- ¹ Research, Innovation and Teaching Unit, Parc Sanitari Sant Joan de Déu, CIBERSAM, 08830 Sant Boi de Llobregat, Spain
- ² Division of Orthodontics and Pediatric Dentistry, Department of Dental Medicine, Karolinska Institutet, 171 77 Stockholm, Sweden; fercsoares@gmail.com
- ³ Institute of Sport Science, University of Graz, 8010 Graz, Austria; mireille.van-poppel@uni-graz.at
- ⁴ Institute of Biomedicine, School of Medicine, University of Eastern Finland, 70211 Kuopio, Finland; saija.savinainen@uef.fi (S.S.); eero.a.haapala@jyu.fi (E.A.H.); timo.lakka@uef.fi (T.L.)
- ⁵ Department of Pediatrics, Kuopio University Hospital, 70211 Kuopio, Finland; aino.mantyselka@kuh.fi
- ⁶ Faculty of Sport and Health Sciences, University of Jyväskylä, 40014 Jyväskylä, Finland
- ⁷ Department of Clinical Physiology and Nuclear Medicine, School of Medicine, Kuopio University Hospital, University of Eastern Finland, 70211 Kuopio, Finland
- ⁸ Kuopio Research Institute of Exercise Medicine, 70100 Kuopio, Finland
- * Correspondence: rodrigoantlima@gmail.com
- † These authors contributed equally to this work.



Citation: Lima, R.A.; Soares, F.C.; van Poppel, M.; Savinainen, S.; Mäntyselkä, A.; Haapala, E.A.; Lakka, T. Determinants of Cognitive Performance in Children and Adolescents: A Populational Longitudinal Study. *Int. J. Environ. Res. Public Health* **2022**, *19*, 8955. <https://doi.org/10.3390/ijerph19158955>

Academic Editor: Paul B. Tchounwou

Received: 17 June 2022

Accepted: 20 July 2022

Published: 23 July 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: We evaluated the determinants of cognitive performance in children and adolescents. This is a longitudinal study, secondary analysis of the Physical Activity and Nutrition in Children (PANIC) study. We assessed 502 children (51.6% girls) at middle childhood (range: 6.6 to 9.0 years), at late childhood, 437 children (51.0% girls, range: 8.8 to 11.2 years), and in 277 adolescents (54.5% girls, range: 15.0 to 17.4 years). Raven's progressive matrices tests estimated the participants' cognitive performance (outcome variable) at all time points. In total, we evaluated 29 factors from various dimensions (prenatal, neonatal, child fitness, lifestyle and anthropometrics). None of the neonatal and anthropometric parameters were associated with cognitive performance. Preeclampsia (prenatal) and listening to music, writing, arts and craft and watching TV (lifestyle) were negatively associated with cognitive performance. Shuttle run and box and block tests (fitness), and playing music, reading and time at the computer (lifestyle) were positive determinants of cognitive performance in children and adolescents. Fitness and lifestyle factors during childhood and adolescence diminished the importance of prenatal factors on cognitive performance and lifestyle factors were especially relevant in regard to cognitive performance. Reading was positively associated with cognitive performance, regardless of age and time dedicated, and should be promoted.

Keywords: children; adolescents; cognition; development; behaviour; lifestyle; health

1. Introduction

Cognitive performance encompasses the ability of processing information, intelligence and reasoning, along with language and memory development [1]. Cognitive development is crucial for adequate self-perception in relation to the social environment and essential for interpersonal connection and interaction [2]. Therefore, cognitive development is pivotal for cultivating our learning and adaptative skills [3]. Ultimately, enhanced cognitive development in youth is related to higher success at school, better job opportunities and higher income, resulting in better quality of life [4,5].

To date, longitudinal studies have reported various factors being associated with cognitive performance in children [6,7] and adolescents [8,9]. In summary, cognitive performance in children and adolescents has been associated with prenatal and neonatal

factors as well as factors related to their childhood and adolescence periods, such as lifestyle, anthropometrics and fitness factors [6–9]. However, there are a number of limitations in the state of the art.

First, only a few studies have long-term follow-ups monitoring children until adolescence [6–14]. Second, although prenatal and neonatal factors have been related to later cognitive performance in early childhood [10,11], it is not well defined whether and how those factors continue to impact cognitive performance later in life [12–14]. Third, it is uncertain whether lifestyle and physical development in childhood overcome or diminish the importance of prenatal and neonatal factors in relation to cognitive performance during childhood and adolescence.

The PANIC study (more information about the study in the Methods Section) followed children (aged six to nine years) until adolescence (aged 15 to 17 years) and contains a dataset with extensive information on factors of the child’s and adolescent’s developmental period [15]. Thus, allowing for a comprehensive analysis of the determinants of cognitive performance in children and adolescents. We appraised all the variables collected in the PANIC study and selected the ones that were associated with cognitive performance in previous studies [6–14]. Taking advantage of the PANIC study dataset, we evaluated prenatal, neonatal, childhood (at two distinct points) and adolescence factors in relation to cognitive performance during childhood and adolescence.

2. Materials and Methods

This is a secondary longitudinal analysis of the PANIC study which is a physical activity and dietary intervention and follow-up study in a population sample of children from eastern Finland. The PANIC study has been described in detail [15]. The study protocol was approved by the Research Ethics Committee of the Hospital District of Northern Savo (Statement 69/2006). The parents or caregivers of the children gave their written informed consent and the children provided their assent to participate. The PANIC study was carried out in accordance with the principles of the Declaration of Helsinki as revised in 2008.

At middle childhood, we assessed 502 children (51.6% girls) of an average age of 7.6 years (range: 6.6 to 9.0 years), at late childhood 437 children (51.0% girls) of 9.8 years of age (range: 8.8 to 11.2 years), and in adolescence 277 participants (54.5% girls) of 15.8 years of age (range: 15.0 to 17.4 years). Tables 1 and 2 present the descriptive characteristics of the participants.

Table 1. Prenatal and neonatal descriptive characteristics of the participants.

	<i>n</i>	Mean	SD
Prenatal			
Maternal weight gain (kg)	1089	5.2	14.2
BMI (Kg/m ²) (1st Trimester)	1203	23.1	4.5
BMI (Kg/m ²) (3rd Trimester)	1104	28.3	4.4
Neonatal			
	<i>n</i>	Mean	SD
Birth height (cm)	1506	50.0	2.1
Birth weight (g)	1515	3535.1	527.6
Independent walking (months)	912	12.0	1.9
Apgar score 1-min (points)	1277	8.7	0.9
Apgar score 5-min (points)	1277	9.0	0.7
Prenatal			
	<i>n</i>		%
Preeclampsia—No	474		96.3
Preeclampsia—Yes	18		3.7
Gestational diabetes mellitus—No	463		92.2
Gestational diabetes mellitus—Yes	39		7.8
Neonatal			
	<i>n</i>		%
Small for gestational age—No	371		90.7
Small for gestational age—Yes	38		9.3

Table 2. Descriptive data on child's fitness, anthropometrics and lifestyle at middle childhood, late childhood and adolescence.

Variables	Middle Childhood			Late Childhood			Adolescence																																																																																																																																																																													
	<i>n</i>	Mean	SD	<i>n</i>	Mean	SD	<i>n</i>	Mean	SD																																																																																																																																																																											
Child Fitness																																																																																																																																																																																				
Sit and reach (cm)	500	−3.2	8.0	422	−6.0	9.7	249	2.8	12.8																																																																																																																																																																											
Handgrip strength right (kg)	504	47.8	9.4	435	62.0	13.5	248	106.2	24.7																																																																																																																																																																											
Handgrip strength left (kg)	504	46.9	9.3	435	61.1	13.6	250	105.6	25.4																																																																																																																																																																											
Sit up (repetitions)	469	10.6	4.8	398	16.9	4.7	246	21.8	5.8																																																																																																																																																																											
Standing long jump (cm)	461	125.8	16.5	393	145.3	20.5	242	191.7	31.6																																																																																																																																																																											
Shuttle run (sec)	456	24.1	2.3	390	22.1	1.7	243	20.6	2.0																																																																																																																																																																											
Box and block (score)	499	101.7	13.4	433	118.1	12.1	244	139.9	17.2																																																																																																																																																																											
Child Anthropometrics																																																																																																																																																																																				
BMI (kg/m ²)	504	16.1	2.1	437	17.3	2.7	276	21.0	3.6																																																																																																																																																																											
BMI SDS	504	−0.2	1.1	437	−0.1	1.1	276	−0.1	1.0																																																																																																																																																																											
Body fat % (excluding the head)	493	21.5	9.1	417	24.6	9.8	265	23.4	10.6																																																																																																																																																																											
Lean mass % (excluding the head)	493	76.2	8.8	417	72.4	9.7	265	72.9	10.4																																																																																																																																																																											
Child Lifestyle																																																																																																																																																																																				
<table border="1"> <thead> <tr> <th colspan="2"></th> <th colspan="2">Middle childhood</th> <th colspan="2">Late childhood</th> <th colspan="2">Adolescence</th> </tr> <tr> <th colspan="2"></th> <th><i>n</i></th> <th>%</th> <th><i>n</i></th> <th>%</th> <th><i>n</i></th> <th>%</th> </tr> </thead> <tbody> <tr> <td rowspan="3">Listening to music</td> <td>0 min/day</td> <td>287</td> <td>57.3</td> <td>209</td> <td>28.5</td> <td>79</td> <td>28.5</td> </tr> <tr> <td>1–30 min/day</td> <td>109</td> <td>21.8</td> <td>92</td> <td>21.5</td> <td>50</td> <td>18.1</td> </tr> <tr> <td>≥30 min/day</td> <td>105</td> <td>21.0</td> <td>128</td> <td>29.8</td> <td>148</td> <td>53.4</td> </tr> <tr> <td rowspan="2">Playing music</td> <td>Yes</td> <td>423</td> <td>84.4</td> <td>327</td> <td>76.2</td> <td>228</td> <td>82.3</td> </tr> <tr> <td>No</td> <td>78</td> <td>15.6</td> <td>102</td> <td>23.8</td> <td>49</td> <td>17.7</td> </tr> <tr> <td rowspan="3">Reading</td> <td>0 min/day</td> <td>131</td> <td>26.2</td> <td>95</td> <td>22.1</td> <td>79</td> <td>28.5</td> </tr> <tr> <td>1–30 min/day</td> <td>174</td> <td>34.7</td> <td>128</td> <td>29.8</td> <td>79</td> <td>28.5</td> </tr> <tr> <td>≥30 min/day</td> <td>196</td> <td>39.1</td> <td>206</td> <td>48.0</td> <td>119</td> <td>42.9</td> </tr> <tr> <td rowspan="3">Writing</td> <td>0 min/day</td> <td>336</td> <td>67.1</td> <td>322</td> <td>75.1</td> <td>103</td> <td>37.2</td> </tr> <tr> <td>1–14 min/day</td> <td>67</td> <td>13.4</td> <td>50</td> <td>11.7</td> <td>32</td> <td>11.6</td> </tr> <tr> <td>≥15 min/day</td> <td>98</td> <td>19.6</td> <td>57</td> <td>13.3</td> <td>142</td> <td>51.3</td> </tr> <tr> <td rowspan="2">Drawing</td> <td>0 min/day</td> <td>125</td> <td>25.0</td> <td>193</td> <td>45.0</td> <td>242</td> <td>87.4</td> </tr> <tr> <td>>0 min/day</td> <td>376</td> <td>75.1</td> <td>236</td> <td>55.0</td> <td>35</td> <td>12.6</td> </tr> <tr> <td rowspan="2">Arts and crafts</td> <td>0 min/day</td> <td>268</td> <td>53.5</td> <td>304</td> <td>70.9</td> <td>244</td> <td>88.1</td> </tr> <tr> <td>>0 min/day</td> <td>233</td> <td>46.5</td> <td>125</td> <td>29.1</td> <td>33</td> <td>11.9</td> </tr> <tr> <td rowspan="3">Watching tv</td> <td>≤30 min/day</td> <td>56</td> <td>11.2</td> <td>42</td> <td>9.8</td> <td>86</td> <td>31.1</td> </tr> <tr> <td>31–90 min/day</td> <td>353</td> <td>70.5</td> <td>280</td> <td>65.3</td> <td>132</td> <td>47.6</td> </tr> <tr> <td>>90 min/day</td> <td>92</td> <td>18.4</td> <td>107</td> <td>24.9</td> <td>59</td> <td>21.3</td> </tr> <tr> <td rowspan="3">Time on the computer</td> <td>0 min/day</td> <td>124</td> <td>24.8</td> <td>64</td> <td>14.9</td> <td>70</td> <td>25.3</td> </tr> <tr> <td>1–60 min/day</td> <td>320</td> <td>63.9</td> <td>275</td> <td>64.1</td> <td>96</td> <td>34.7</td> </tr> <tr> <td>>60 min/day</td> <td>57</td> <td>11.4</td> <td>90</td> <td>21.0</td> <td>111</td> <td>40.1</td> </tr> </tbody> </table>												Middle childhood		Late childhood		Adolescence				<i>n</i>	%	<i>n</i>	%	<i>n</i>	%	Listening to music	0 min/day	287	57.3	209	28.5	79	28.5	1–30 min/day	109	21.8	92	21.5	50	18.1	≥30 min/day	105	21.0	128	29.8	148	53.4	Playing music	Yes	423	84.4	327	76.2	228	82.3	No	78	15.6	102	23.8	49	17.7	Reading	0 min/day	131	26.2	95	22.1	79	28.5	1–30 min/day	174	34.7	128	29.8	79	28.5	≥30 min/day	196	39.1	206	48.0	119	42.9	Writing	0 min/day	336	67.1	322	75.1	103	37.2	1–14 min/day	67	13.4	50	11.7	32	11.6	≥15 min/day	98	19.6	57	13.3	142	51.3	Drawing	0 min/day	125	25.0	193	45.0	242	87.4	>0 min/day	376	75.1	236	55.0	35	12.6	Arts and crafts	0 min/day	268	53.5	304	70.9	244	88.1	>0 min/day	233	46.5	125	29.1	33	11.9	Watching tv	≤30 min/day	56	11.2	42	9.8	86	31.1	31–90 min/day	353	70.5	280	65.3	132	47.6	>90 min/day	92	18.4	107	24.9	59	21.3	Time on the computer	0 min/day	124	24.8	64	14.9	70	25.3	1–60 min/day	320	63.9	275	64.1	96	34.7	>60 min/day	57	11.4	90	21.0	111	40.1
		Middle childhood		Late childhood		Adolescence																																																																																																																																																																														
		<i>n</i>	%	<i>n</i>	%	<i>n</i>	%																																																																																																																																																																													
Listening to music	0 min/day	287	57.3	209	28.5	79	28.5																																																																																																																																																																													
	1–30 min/day	109	21.8	92	21.5	50	18.1																																																																																																																																																																													
	≥30 min/day	105	21.0	128	29.8	148	53.4																																																																																																																																																																													
Playing music	Yes	423	84.4	327	76.2	228	82.3																																																																																																																																																																													
	No	78	15.6	102	23.8	49	17.7																																																																																																																																																																													
Reading	0 min/day	131	26.2	95	22.1	79	28.5																																																																																																																																																																													
	1–30 min/day	174	34.7	128	29.8	79	28.5																																																																																																																																																																													
	≥30 min/day	196	39.1	206	48.0	119	42.9																																																																																																																																																																													
Writing	0 min/day	336	67.1	322	75.1	103	37.2																																																																																																																																																																													
	1–14 min/day	67	13.4	50	11.7	32	11.6																																																																																																																																																																													
	≥15 min/day	98	19.6	57	13.3	142	51.3																																																																																																																																																																													
Drawing	0 min/day	125	25.0	193	45.0	242	87.4																																																																																																																																																																													
	>0 min/day	376	75.1	236	55.0	35	12.6																																																																																																																																																																													
Arts and crafts	0 min/day	268	53.5	304	70.9	244	88.1																																																																																																																																																																													
	>0 min/day	233	46.5	125	29.1	33	11.9																																																																																																																																																																													
Watching tv	≤30 min/day	56	11.2	42	9.8	86	31.1																																																																																																																																																																													
	31–90 min/day	353	70.5	280	65.3	132	47.6																																																																																																																																																																													
	>90 min/day	92	18.4	107	24.9	59	21.3																																																																																																																																																																													
Time on the computer	0 min/day	124	24.8	64	14.9	70	25.3																																																																																																																																																																													
	1–60 min/day	320	63.9	275	64.1	96	34.7																																																																																																																																																																													
	>60 min/day	57	11.4	90	21.0	111	40.1																																																																																																																																																																													

On average, children were born at 39.8 weeks of gestation (± 1.8 weeks). Mothers were 30 years of age on average (range 16 to 44 years), 42.8% did not have any previous births and 31.1% one previous birth; the family income was lower than EUR 30,000 per year for 21.4% of the families and between EUR 30,000 and EUR 60,000 per year for 41.8% of the families.

Children not followed at adolescence presented higher body fat percentage compared to their peers at baseline. A higher proportion of participants born from mothers with preeclampsia were followed at adolescence compared to their peers from mothers who did not have preeclampsia during pregnancy. There was no difference in any other relevant variable because of loss to follow-up.

2.1. Patient and Public Involvement

Patients or the public were not involved in the design, or conduct, or reporting or dissemination plans of our research.

2.2. Outcome

Cognitive performance was estimated twice in childhood (middle childhood at 6.6 to 9.0 years of age, late childhood at 8.8 to 11.2 years of age) by the Raven's coloured progressive matrices and by Raven's progressive matrices in adolescence (15.0 to 17.4 years of age) [16]. Raven's matrices assess abstract reasoning and fluid intelligence; coloured progressive matrices include 36 and progressive matrices 60 different items [16]. All of the questions on the Raven's tests consist of visual geometric design with a missing piece. The participant could choose from six to eight choices to pick from to fill in the missing piece. The test becomes increasingly complex, requiring ever-greater cognitive capacity to encode and analyse information for participants as the test progresses [16].

2.3. Factors

2.3.1. Socioeconomic Background

Socioeconomic background was assessed by the annual household income (family income) and the level of education of the parents. The family income was asked by a structured questionnaire, at middle childhood, late childhood and adolescence, from both parents and coded into three categories (\leq EUR 30,000, EUR 30,001–EUR 60,000 and $>$ EUR 60,000). The level of education was asked by a structured questionnaire from both parents and coded into three categories (vocational school or less, vocational high school, university) based on the highest completed or ongoing degree. If the parents reported different categories, the higher category was used in the analyses.

2.3.2. Prenatal and Neonatal Exposures

Data were collected on maternal age at child's birth, gestational age at birth, maternal weight gain, preeclampsia, maternal body mass index (BMI at 1st and 3rd trimesters during pregnancy), gestational diabetes, singleton pregnancy, number of previous births (parity), smoking status during pregnancy (no smoking, smoked but quit during the first trimester, smoked after the first trimester) and also the participants' birth height, birth weight, independent walking (time in which the child started walking independently) and the one- and five-minute Apgar scores assigned retrospectively from the birth register provided by the National Institute for Health and Welfare. The one- and five-minute Apgar scores evaluate newborns in five criteria: activity (tone), pulse, grimace, appearance and respiration. For each criterion, newborns can receive a score from 0 to 2 [17].

Preeclampsia was defined as hypertension and proteinuria occurring after 20 weeks of gestation. Hypertension was defined as systolic blood pressure \geq 140 mmHg or diastolic blood pressure \geq 90 mm Hg after 20 weeks of gestation. Proteinuria was defined as the urinary excretion of \geq 0.3 g protein in a 24 h specimen, or 0.3 g/L or two \geq 1+ readings on dipstick in a random urine determination with no evidence of urinary tract infection.

2.3.3. Pubertal Status

Because only a few children had entered clinical puberty at middle childhood, current height as a percentage of predicted adult height was used as a measure of pubertal status. The boys' predicted adult height was calculated as follows: the mean of the height of Finnish men (178.6 cm) + [the standard deviation of the height of Finnish men (6.0 cm) \times the deviation of the child's predicted adult height from the average of the predicted adult height of Finnish children]. The girl's predicted adult height was calculated as follows: the mean of the height of Finnish women (165.3 cm) + [the standard deviation of the height of Finnish women (5.4 cm) \times the deviation of the child's predicted adult height from the average of the predicted adult height of Finnish children]. The deviation in the child's predicted adult height from the average of the predicted adult height of Finnish children

was calculated according to the national guidelines as follows: (the arithmetic mean of the father's and mother's height-171)/10 [18].

2.3.4. Child Fitness Exposures

Sit and Reach

Lower back and hamstring muscle flexibility were assessed by the sit-and-reach test [19]. The children were asked to sit down with their heels 25 cm apart at the zero line. A measuring stick was placed to -38 cm from the zero line. The children were asked to reach slowly forward as far as possible while keeping the hands parallel and to repeat the same task three times. The test score was the longest distance in cm reached with the fingertips from the starting line of -38 cm, with a smaller distance reached indicating poorer lower back and hamstring flexibility.

Handgrip

Handgrip strength was assessed by the Martin vigorimeter (Martin, Tuttlingen, Germany). The children were asked to keep their elbow close to the body, their arm flexed at 90° and to press a rubber bulb maximally three times each with their right and left hand. The mean of the best trial of each hand was used in the analyses and was expressed in kilopascals [19].

Sit-Up

The sit-up test was used to assess abdominal muscle strength and endurance [19]. The children were asked to lie down with knees flexed at 90°, feet on the ground and arms behind the neck. The children were told to perform as many sit-ups as possible in 30 s with their elbows touching their knees as the assistant held their feet on the floor. The test score was the number of technically correct sit-ups completed in 30 s.

Standing Long Jump

Lower limb explosive strength was assessed by the standing long jump test [19]. The children were asked to place their feet next to each other, jump as far as possible and land on both feet. The test score was the best result of three attempts in cm.

Shuttle Run

Speed and agility were assessed by the 50 m shuttle run test [20]. Children were asked to run five meters from a starting line to another line as fast as possible, to turn on the line, to run back to the starting line and to repeat until five repetitions were completed. The test score was the running time in seconds, with a longer time indicating a poorer performance.

Box and Block

Manual dexterity and upper-limb movement speed were assessed by the box-and-block test [21]. The children were asked to pick up 150 small wooden cubes (2.5 cm/side) one by one with the dominant hand from one side of a wooden box (53.7 cm × 25.4 cm × 8 cm), to move as many cubes as possible to the other side of the box over 60 s and to repeat the same task with the non-dominant hand. The test score was the number of cubes moved to the other side of the box, with smaller number of cubes moved indicating poorer manual dexterity.

2.3.5. Lifestyle Exposures

The following exposures were assessed at middle and late childhood and adolescence. The time the participants' spent listening to music, playing music, reading, writing, drawing, doing arts and crafts, watching TV and on the computer were inquired by the PANIC Physical Activity Questionnaire administered by the parents with the children [22]. Time spent on each lifestyle factor was queried separately for weekdays and weekends (in

minutes per day). The amount of total time for each factor was calculated by adding the times spent in each factor weighted by the number of weekdays and weekend days. We categorised the time in each factor to better represent the time participants spent in each lifestyle factor.

2.3.6. Body Composition and Anthropometrics

Body lean mass and body fat percentage were measured after emptying the bladder, in a supine position and light clothing and after removing all metal objects, by a Lunar[®] dual-energy X-ray absorptiometry (DXA) device (Lunar Prodigy Advance; GE Medical Systems, Madison, WI, USA) [23]. We excluded the head in the estimation of the participants' fat percentage. Body weight was measured twice after overnight fasting, after emptying the bladder and standing in light underwear using a calibrated InBody[®] 720 bioelectrical impedance device (Biospace, Seoul, Korea) to an accuracy of 0.1 kg. The mean of these two values was used in the analyses. Stature was measured three times in the Frankfurt plane without shoes using a wall-mounted stadiometer to accuracy of 0.1 cm. BMI was calculated as body mass (kg) divided by stature (m) squared. The BMI-standard deviation score (BMI-SDS) was calculated based on Finnish references values [24]. The prevalence of underweight, normal weight, overweight and obesity was defined using the national reference values provided by Saari and colleagues [24].

2.4. Statistical Analysis

We used STATA version 16 for Windows (StataCorp LP, College Station, TX, USA) and R Studio version 2021.9.0.351 (PBC, Boston, MA, USA) for analysis. Means, standard deviations and absolute and relative frequencies described the variables of interest. Chi-square (categorical variables) or two-way ANOVA (numerical variables) was conducted to evaluate possible differences in the outcome or factors comparing participants with complete data with adolescents not monitored at the second follow-up (dropout analysis). We used hierarchical multilevel linear regressions to evaluate the factors longitudinally associated with cognitive performance. Our hierarchical analysis was conducted in the following steps.

2.4.1. First Step: Defining the Adjustments

In addition to child's age, sex, pubertal status and intervention group, defined a priori to be included as adjustments, we evaluated which maternal demographic variables (maternal age, family income, gestational age at birth and parity) would be used as adjustments. We ran multilevel linear regressions between each of the abovementioned variables and cognitive performance adjusted for child's age, sex, pubertal status and intervention group. Maternal age, family income, parity and gestational age at birth were associated with cognitive performance and used as adjustments.

2.4.2. Second Step: Evaluate the Factors Associated with Cognitive Performance

We individually evaluated each of the factors in relation to cognitive performance. All the analyses were adjusted for child's age, sex, pubertal status, intervention group, maternal age, family income, parity and gestational age at birth. Since we were interested in estimating the longitudinal association between the factors and cognitive performance for each of the life phases (middle childhood, late childhood and adolescence), we included an interaction factor between the factor and time (factor * time).

2.4.3. Third Step: Evaluate the Factors, amongst Domains, Associated with Cognitive Performance

All the factors associated ($p \leq 0.05$) with cognitive performance in at least one life phase in the second step were included in this third phase, which was conducted in the following order:

1. All the factors with $p \leq 0.05$ in the prenatal domain plus the variables in the neonatal domain.
2. After selecting all the remaining factors from the prenatal and neonatal domains, we inserted the factors from the child fitness domain into the model.
3. After selecting all the remaining factors from the prenatal, neonatal and child fitness domains, we inserted the factors from the child lifestyle domain in the model.

In this third step, we used the forward stepwise procedure to include and select the factors in the model. We started with the factor with the lowest p value. It is important to highlight that we maintained the interaction term between each factor and time in the third step and that we accounted for the cluster structure of data at school level in all multilevel analyses. Preliminary analyses did not observe collinearity between factors in the models. Of note, the associations between factors and cognitive performance were presented as the predicted score in the Raven's test. Further, to account for the multiple testing, the p -values shown are the sharpened False Discovery Rate (FDR) q -values [25].

3. Results

Figure 1 presents the cognitive performance of the participants during childhood and adolescence. Children at middle childhood scored on the cognitive performance test, on average, 24.0 points (range 4.0 to 35.0 points), and children at late childhood scored 29.1 points on average (range 13.0 to 36.0 points). Adolescents' cognitive performance was 48.0 points on average (range 11.0 to 60.0 points).




Figure 1. Participants' cognitive performance during middle childhood, late childhood and adolescence.

3.1. Prenatal Factors in Relation to Cognitive Performance

From all the prenatal factors tested, only preeclampsia was longitudinally associated with cognitive performance. Particularly, adolescents whose mothers had preeclampsia presented a lower cognitive performance compared to their peers whose mothers did not have preeclampsia (Preeclampsia (No): 50.15, 95% CI 45.44 to 54.86; preeclampsia (Yes): 45.85, 95% CI 39.85 to 51.84; $p = 0.018$; Table 3). However, preeclampsia was not significantly associated with cognitive performance after considering fitness factors (Table 3, Prenatal + Neonatal + Child Fitness model).

3.2. Neonatal Factors in Relation to Cognitive Performance

None of the neonatal factors was longitudinally associated with cognitive performance (Table 3).

3.3. Child's Fitness Factors in Relation to Cognitive Performance

From all the child's fitness factors tested, only the shuttle run and box-and-block tests were longitudinally associated with cognitive performance. One additional second in the shuttle run test was associated with lower cognitive performance at middle childhood (-0.248 points; $p = 0.038$). Each additional score in the box and block test was associated with higher cognitive performance at middle childhood (0.067 points; $p = 0.001$) and adolescence (0.058 points; $p = 0.010$). However, the shuttle run and box and block tests were no longer associated with cognitive performance in the model including Prenatal + Neonatal + Child Fitness factors (Table 3).

3.4. Child's Lifestyle Factors in Relation to Cognitive Performance

A wide range of lifestyle factors was associated with cognitive performance. Specifically, adolescents who listened to music presented lower cognitive performance compared to their peers who did not listen to music. Adolescents who played music showed higher cognitive performance compared to adolescents who did not play music. Reading was longitudinally associated with higher cognitive performance in all developmental phases (middle and late childhood and adolescence). Writing outside the school period was longitudinally associated with lower cognitive performance in middle and late childhood. Adolescents who did arts and crafts showed lower cognitive performance compared to their peers who did not perform arts and crafts. Adolescents who watched TV for more than 30 min/day presented lower cognitive performance compared to their peers who did not watch TV. Children at middle childhood who spent 1 to 60 min/day on the computer exhibited higher cognitive performance compared to their peers who did not dedicate any time to the computer (Table 3, factors individually + adjustments).

A few lifestyle factors remained associated with cognitive performance in the fully adjusted model (Prenatal + Neonatal + Child Fitness + Child Lifestyle + adjustments model). Particularly, adolescents who listened to music presented lower cognitive performance compared to their peers who did not listen to music. Adolescents who played music exhibited higher cognitive performance compared to their peers who did not play music. Reading was associated with higher cognitive performance in all developmental periods. Writing outside the school period was negatively associated with cognitive performance in all developmental periods (middle and late childhood and adolescence). Adolescents who did arts and crafts showed lower cognitive performance compared to their peers who did not perform arts and crafts. Limited time on the computer (1–60 min/day) was associated with higher cognitive performance at middle childhood and adolescence compared to their peers without time on the computer (Table 3, Prenatal + Neonatal + Child Fitness + Child Lifestyle model).

Table 3. Cont.

Factors	Predicted Cognitive Performance, Scores in the Raven's Test											
	Factors Individually			Prenatal + Neonatal			Prenatal + Neonatal + Child Fitness			Prenatal + Neonatal+ Child Fitness + Child Lifestyle		
	Raven's Score	(95% CI)	p	Raven's Score	(95% CI)	p	Raven's Score	(95% CI)	p	Raven's Score	(95% CI)	p
≥0 min/day (ref. (0 min/day))—late childhood	29.30	(28.33 to 30.28)	0.387									
0 min/day—adolescence	49.47	(44.67 to 54.28)										
≥0 min/day (ref. (0 min/day))—adolescence	51.33	(46.22 to 56.43)	0.096									
Arts craft												
0 min/day—middle childhood	22.36	(20.02 to 24.70)							22.38	(20.16 to 24.60)		
≥0 min/day (ref. (0 min/day))—middle childhood	23.39	(21.06 to 25.71)	0.055						23.39	(21.18 to 25.59)		0.059
0 min/day—late childhood	28.96	(28.07 to 29.85)							28.54	(27.67 to 29.42)		
≥0 min/day (ref. (0 min/day))—late childhood	29.10	(27.94 to 30.25)	0.827						28.97	(27.86 to 30.07)		0.296
0 min/day—adolescence	50.83	(46.04 to 55.63)							52.29	(47.68 to 56.89)		
≥0 min/day (ref. (0 min/day))—adolescence	47.80	(42.80 to 52.79)	0.005						48.53	(43.76 to 53.30)		0.041
Watching tv												
<30 min/day—middle childhood	21.94	(19.17 to 24.72)							21.79	(19.15 to 24.44)		
31–90 min/day (ref. (<30 min/day))—middle childhood	23.46	(21.16 to 25.75)	0.097						22.99	(20.81 to 25.18)		0.133
>90 min/day (ref. (<30 min/day))—middle childhood	22.73	(20.26 to 25.20)	0.453						22.50	(20.15 to 24.85)		0.296
<30 min/day—late childhood	30.34	(28.43 to 32.25)							30.04	(28.21 to 31.87)		
31–90 min/day (ref. (<30 min/day))—late childhood	28.68	(27.77 to 29.58)	0.090						28.24	(27.39 to 29.09)		0.059
>90 min/day (ref. (<30 min/day))—late childhood	29.53	(28.34 to 30.73)	0.447						29.13	(27.28 to 30.28)		0.246
<30 min/day—adolescence	51.06	(46.23 to 55.89)							52.35	(47.75 to 56.95)		
31–90 min/day (ref. (<30 min/day))—adolescence	49.08	(44.25 to 53.92)	0.013						50.74	(46.12 to 55.36)		0.050
>90 min/day (ref. (<30 min/day))—adolescence	49.00	(44.09 to 53.91)	0.035						51.0	(46.27 to 55.73)		0.133
Time on the computer												
0 min/day—middle childhood	22.14	(19.68 to 24.60)							21.83	(19.49 to 24.17)		
1–60 min/day (ref. (0 min/day))—middle childhood	23.68	(21.38 to 25.97)	0.015						23.14	(20.95 to 25.32)		0.044
>60 min/day (ref. (0 min/day))—middle childhood	22.87	(20.20 to 25.55)	0.448						22.41	(19.85 to 24.97)		0.307
0 min/day (ref. (0 min/day))—late childhood	28.99	(27.43 to 30.56)							28.85	(27.36 to 30.34)		
1–60 min/day (ref. (0 min/day))—late childhood	29.30	(28.41 to 30.20)	0.705						28.80	(27.93 to 29.67)		0.458
>60 min/day (ref. (0 min/day))—late childhood	28.33	(27.02 to 29.63)	0.487						28.19	(26.94 to 29.45)		0.296
0 min/day—adolescence	49.46	(44.51 to 54.31)							49.73	(45.07 to 54.38)		
1–60 min/day (ref. (0 min/day))—adolescence	50.56	(45.72 to 55.40)	0.234						51.85	(47.21 to 56.50)		0.042
>60 min/day (ref. (0 min/day))—adolescence	48.97	(44.09 to 53.84)	0.596						49.95	(45.27 to 54.63)		0.434
Child Anthropometrics												
BMI (kg/m ²)—middle childhood	−0.017	(−0.269 to 0.235)	0.893									
BMI (kg/m ²)—late childhood	0.075	(−0.136 to 0.285)	0.488									
BMI (kg/m ²)—adolescence	0.800	(−0.150 to 0.310)	0.495									
BMI SDS—middle childhood	0.066	(−0.415 to 0.548)	0.787									
BMI SDS—late childhood	0.111	(−0.413 to 0.634)	0.678									
BMI SDS—adolescence	0.184	(−0.551 to 0.919)	0.624									
Body fat % (excluding the head)—middle childhood	0.009	(−0.050 to 0.068)	0.758									
Body fat % (excluding the head)—late childhood	0.019	(−0.041 to 0.079)	0.543									
Body fat % (excluding the head)—adolescence	0.044	(−0.026 to 0.115)	0.221									
Lean mass % (excluding the head)—middle childhood	−0.009	(−0.070 to 0.053)	0.777									
Lean mass % (excluding the head)—late childhood	−0.018	(−0.079 to 0.043)	0.565									
Lean mass % (excluding the head)—adolescence	−0.043	(−0.115 to 0.029)	0.238									

Legend: Results are presented in the predicted values of the Raven's score. All analyses were adjusted for child's age, sex, pubertal status, intervention group, and maternal age, family income, parity and gestational age at birth. *p*-values are the sharpened False Discovery Rate *q*-values.

4. Discussion

Several factors were longitudinally associated with cognitive performance in children and adolescents. None of the neonatal and anthropometric parameters was associated with cognitive performance. Preeclampsia (prenatal), listening to music, writing, arts and crafts and watching TV (lifestyle) were negatively associated with cognitive performance. On the other hand, the shuttle run and box and block tests (fitness), and playing music, reading and time on the computer (lifestyle) were positive determinants of cognitive performance in children and adolescents. Fitness and lifestyle factors overcame the importance of prenatal factors on cognitive performance and lifestyle factors were especially relevant in regard to cognitive performance. Of note, reading was the most important positive determinant of cognitive performance during childhood and adolescence.

Our findings highlight the importance of all life periods in relation to cognitive development, from conception to behaviours during adolescence. However, there are some key elements to better understand the importance of each lifetime period in relation to cognitive development. Behaviours incorporated later in life may overcome or diminish the importance of being exposed to previous factors negatively related to cognitive performance.

For example, preeclampsia was negatively associated with cognitive performance in adolescents, as demonstrated previously [26,27]. However, preeclampsia was not associated with cognitive performance after considering the participants' fitness factors, particularly the shuttle run and the box and block tests. Those are novel and relevant findings because it might be possible to diminish the importance of deleterious factors that the children might have been exposed to during pregnancy or early childhood in relation to cognitive development.

It is complex to interpret the importance of lifestyle factors in relation to cognitive performance. Some lifestyle factors seem to stimulate, whereas others were negatively associated with cognitive performance. Our findings suggest that lifestyle factors play a major role on cognitive performance, particularly during adolescence.

It seems that limited exposure to listening to music, writing, arts and crafts, and watching TV are positively associated with cognitive performance, since dedicating time to those factors was related to worse cognitive performance. Further, listening to music, writing and arts and crafts remained associated with cognitive performance when considering the other lifestyle factors. It is possible that adolescents who spent additional time outside school writing were the ones exhibiting lower cognitive and academic performance at school. Listening to music and arts and crafts during adolescence might be depriving time for reading, playing music and limited computer time (1–60 min/day), which were positively related to cognitive performance. Importantly, future investigations should assess these findings in depth.

It is possible that a wide range of stimuli is beneficial for cognitive performance during childhood and adolescence [28–30]. Particularly, playing music during adolescence, 1 to 60 min/day on the computers and reading during childhood and adolescence were related to higher cognitive performance. Further, the more children read, the better their cognitive performance. The importance of reading and reading performance is well established in regard to cognitive function and development in children and adolescents, and recent evidence shows that the relationship might be reciprocal [31]. In summary, our findings support the premise that additional time reading is linked to better cognitive performance during childhood and adolescence.

Limitations

This study carries limitations that should be considered in the interpretation of the findings. We ran several models, including on the number of factors. Thus, some models might be over fitted. Nevertheless, we also present each factor in relation to cognitive performance adjusted for confounders, similarly to previous studies. We ran various models, but *p*-values were adjusted for multiple testing to diminish the risk of type I errors. This is not a randomised controlled trial; hence, we cannot infer on the causal relationship between

factors. We observed differences in body fat percentage and proportion of preeclampsia due to loss to follow-up. Moreover, there might be other factors that might impact cognitive development that we did not account for in our study and should be further investigated, such as genetic [32,33] and nutritional factors [34,35].

5. Conclusions

Cognitive performance throughout childhood and adolescence was associated with factors from conception to adolescence. Although some prenatal factors were negatively associated with cognitive performance in children, it seems that better fitness and lifestyle factors during childhood can negate these negative relationships. Spending time on a mix of activities seems beneficial and more cognitive challenging lifestyles seem to be more relevant in later childhood and adolescence. Reading is positively associated with cognitive performance, regardless of age, and should be promoted. Practitioners, future interventions and public health policies should be aware of the probable changes in the role of the various lifestyle factors on cognitive performance in different lifetime phases.

Author Contributions: Conceptualization, R.A.L., F.C.S., M.v.P. and E.A.H.; Data curation, A.M. and T.L.; formal analysis, R.A.L. and F.C.S.; funding acquisition, E.A.H. and T.L.; investigation, T.L.; methodology, S.S., A.M., E.A.H. and T.L.; project administration, S.S.; resources, M.v.P. and S.S.; supervision, F.C.S. and M.v.P.; writing—original draft, R.A.L.; writing—review and editing, F.C.S., M.v.P., S.S., A.M., E.A.H. and T.L. All authors have read and agreed to the published version of the manuscript.

Funding: The PANIC Study was financially supported by the Ministry of Education and Culture of Finland [award/grant number: N/A], Ministry of Social Affairs and Health of Finland [award/grant number: N/A], Research Committee of the Kuopio University Hospital Catchment Area (State Research Funding [award/grant number: N/A]), Finnish Innovation Fund Sitra [award/grant number: N/A], Social Insurance Institution of Finland [award/grant number: N/A], Finnish Cultural Foundation [award/grant number: N/A], Foundation for Paediatric Research [award/grant number: N/A], Diabetes Research Foundation in Finland [award/grant number: N/A], Finnish Foundation for Cardiovascular Research [award/grant number: N/A], Juho Vainio Foundation [award/grant number: N/A], Paavo Nurmi Foundation [award/grant number: N/A], Yrjö Jahnsson Foundation [award/grant number: N/A], and the city of Kuopio [award/grant number: N/A]. Moreover, the PhD students and postdoctoral researchers of The PANIC Study have been supported by Program for Clinical Research and Program for Health Sciences of Doctoral School of University of Eastern Finland [award/grant number: N/A], Finnish Doctoral Programs in Public Health [award/grant number: N/A], Päivikki and Sakari Sohlberg Foundation [award/grant number: N/A], Paulo Foundation [award/grant number: N/A], Jalmari and Rauha Ahokas Foundation [award/grant number: N/A], Aarne and Aili Turunen Foundation [award/grant number: N/A], Finnish Medical Foundation [award/grant number: N/A], Aino Eerola and Orion Trusts of Finnish Medical Foundation [award/grant number: N/A], Finnish Medical Society Duodecim [award/grant number: N/A], the Foundation of Kuopio University Hospital for Scientific Research [award/grant number: N/A], Jenny and Antti Wihuri Foundation [award/grant number: N/A], Otto A. Malm Foundation [award/grant number: N/A], Emil Aaltonen Foundation [award/grant number: N/A], Helena Vuorenmies Foundation [award/grant number: N/A], Orion Research Foundation sr [award/grant number: N/A], Kuopio Naturalists' Society [award/grant number: N/A], Olvi Foundation [award/grant number: N/A] and the city of Kuopio [award/grant number: N/A].

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: The parents or caregivers of the children gave their written informed consent and the children provided their assent to participate. The Research Ethics Committee of the Hospital District of Northern Savo approved the study protocol in 2006 (Statement 69/2006).

Data Availability Statement: The data is not publicly available due to ethical reasons. However, Timo A. Lakka can provide further information on the PANIC study and the PANIC data on a reasonable request.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Donnelly, J.E.; Hillman, C.H.; Castelli, D.; Etnier, J.L.; Lee, S.; Tomporowski, P.; Lambourne, K.; Szabo-Reed, A.N. Physical Activity, Fitness, Cognitive Function, and Academic Achievement in Children: A Systematic Review. *Med. Sci. Sport. Exerc.* **2016**, *48*, 1197–1222. [[CrossRef](#)]
- Children National Research Council (US) Panel to Review the Status of Basic Research on School-Age. *Development During Middle Childhood: The Years From Six to Twelve*; Collins, W.A., Ed.; National Academies Press: Washington, DC, USA, 1984; ISBN 0-309-03478-7.
- Koh, K. Maternal Breastfeeding and Children's Cognitive Development. *Soc. Sci. Med.* **2017**, *187*, 101–108. [[CrossRef](#)]
- Hirano, K.; Imbens, G.W.; Ridder, G. Efficient Estimation of Average Treatment Effects Using the Estimated Propensity Score. *Econometrica* **2003**, *71*, 1161–1189. [[CrossRef](#)]
- Currie, J.; Almond, D. Human Capital Development before Age Five. In *Handbook of Labor Economics*; Elsevier: Amsterdam, The Netherlands, 2011; Volume 4, pp. 1315–1486, ISBN 1573-4463.
- Ford, N.D.; Stein, A.D. Risk Factors Affecting Child Cognitive Development: A Summary of Nutrition, Environment, and Maternal-Child Interaction Indicators for Sub-Saharan Africa. *J. Dev. Orig. Health Dis.* **2016**, *7*, 197–217. [[CrossRef](#)]
- De Greeff, J.W.; Bosker, R.J.; Oosterlaan, J.; Visscher, C.; Hartman, E. Effects of Physical Activity on Executive Functions, Attention and Academic Performance in Preadolescent Children: A Meta-Analysis. *J. Sci. Med. Sport.* **2018**, *21*, 501–507. [[CrossRef](#)]
- Barch, D.M.; Albaugh, M.D.; Avenevoli, S.; Chang, L.; Clark, D.B.; Glantz, M.D.; Hudziak, J.J.; Jernigan, T.L.; Tapert, S.F.; Yurgelun-Todd, D.; et al. Demographic, Physical and Mental Health Assessments in the Adolescent Brain and Cognitive Development Study: Rationale and Description. *Dev. Cogn. Neurosci.* **2018**, *32*, 55–66. [[CrossRef](#)]
- Esteban-Cornejo, I.; Ma Tejero-Gonzalez, C.; Sallis, J.F.; Veiga, O.L. Physical Activity and Cognition in Adolescents: A Systematic Review. *J. Sci. Med. Sport* **2015**, *18*, 534–539. [[CrossRef](#)]
- Sacchi, C.; Marino, C.; Nosarti, C.; Vieno, A.; Visentin, S.; Simonelli, A. Association of Intrauterine Growth Restriction and Small for Gestational Age Status With Childhood Cognitive Outcomes A Systematic Review and Meta-Analysis. *JAMA Pediatr.* **2020**, *174*, 772–781. [[CrossRef](#)]
- Duncan, A.F.; Matthews, M.A. Neurodevelopmental Outcomes in Early Childhood. *Clin. Perinatol.* **2018**, *45*, 377–392. [[CrossRef](#)]
- Gale, C.R.; O'Callaghan, F.J.; Godfrey, K.M.; Law, C.M.; Martyn, C.N. Critical Periods of Brain Growth and Cognitive Function in Children. *Brain* **2004**, *127*, 321–329. [[CrossRef](#)]
- Oddy, W.H.; Kendall, G.E.; Blair, E.; De Klerk, N.H.; Stanley, F.J.; Landau, L.I.; Silburn, S.; Zubrick, S. Breast Feeding and Cognitive Development in Childhood: A Prospective Birth Cohort Study. *Paediatr. Perinat. Epidemiol.* **2003**, *17*, 81–90. [[CrossRef](#)]
- Zhang, H.; Lee, Z.X.; White, T.; Qiu, A. Parental and Social Factors in Relation to Child Psychopathology, Behavior, and Cognitive Function. *Transl. Psychiatry* **2020**, *10*, 80. [[CrossRef](#)]
- Leppänen, M.H.; Haapala, E.A.; Veijalainen, A.; Seppälä, S.; Oliveira, R.S.; Lintu, N.; Laitinen, T.; Tarvainen, M.P.; Lakka, T.A. Associations of Cardiometabolic Risk Factors with Heart Rate Variability in 6- to 8-year-old Children: The PANIC Study. *Pediatr. Diabetes* **2020**, *21*, 251–258. [[CrossRef](#)]
- Raven, J.; Raven, J.; Court, J. *Coloured Progressive Matrices. Manual for Raven's Progressive Matrices and Vocabulary Scales*; Oxford Psychologist Press Ltd.: London, UK, 1998.
- Stark, A.R.; Adamkin, D.H.; Batton, D.G.; Bell, E.F.; Bhutani, V.K.; Denson, S.E.; Engle, W.A.; Martin, G.I.; Blackmon, L.R.; Barrington, K.J.; et al. The Apgar Score. *Pediatrics* **2006**, *117*, 1444–1447.
- Haapala, E.A.; Poikkeus, A.-M.; Tompuri, T.; Kukkonen-Harjula, K.; Leppänen, P.H.T.; Lindi, V.; Lakka, T.A. Associations of Motor and Cardiovascular Performance with Academic Skills in Children. *Med. Sci. Sports Exerc.* **2014**, *46*, 1016–1024. [[CrossRef](#)]
- European Council. *Eurofit: Handbook for the Eurofit Tests of Physical Fitness*, 1st ed.; Council of Europe: Rome, Italy, 1988.
- Léger, L.A.; Mercier, D.; Gadoury, C.; Lambert, J. The Multistage 20 Metre Shuttle Run Test for Aerobic Fitness. *J. Sports Sci.* **1988**, *6*, 93–101. [[CrossRef](#)]
- Jongbloed-Pereboom, M.; Nijhuis-van der Sanden, M.W.G.; Steenbergen, B. Norm Scores of the Box and Block Test for Children Ages 3–10 Years. *Am. J. Occup. Ther.* **2013**, *67*, 312–318. [[CrossRef](#)]
- Vaisto, J.; Eloranta, A.-M.; Viitasalo, A.; Tompuri, T.; Lintu, N.; Karjalainen, P.; Lampinen, E.-K.; Agren, J.; Laaksonen, D.E.; Lakka, H.-M.; et al. Physical Activity and Sedentary Behaviour in Relation to Cardiometabolic Risk in Children: Cross-Sectional Findings from the Physical Activity and Nutrition in Children (PANIC) Study. *Int. J. Behav. Nutr. Phys. Act.* **2014**, *11*, 55. [[CrossRef](#)]
- Tompuri, T.T.; Lakka, T.A.; Hakulinen, M.; Lindi, V.; Laaksonen, D.E.; Kilpeläinen, T.O.; Jääskeläinen, J.; Lakka, H.-M.; Laitinen, T. Assessment of Body Composition by Dual-Energy X-Ray Absorptiometry, Bioimpedance Analysis and Anthropometrics in Children: The Physical Activity and Nutrition in Children Study. *Clin. Physiol. Funct. Imaging* **2015**, *35*, 21–33. [[CrossRef](#)]
- Saari, A.; Sankilampi, U.; Hannila, M.-L.; Kiviniemi, V.; Kesseli, K.; Dunkel, L. New Finnish Growth References for Children and Adolescents Aged 0 to 20 Years: Length/Height-for-Age, Weight-for-Length/Height, and Body Mass Index-for-Age. *Ann. Med.* **2011**, *43*, 235–248. [[CrossRef](#)]
- Verhoeven, K.J.F.; Simonsen, K.L.; McIntyre, L.M. Implementing False Discovery Rate Control: Increasing Your Power. *Oikos* **2005**, *108*, 643–647. [[CrossRef](#)]
- Sverrisson, F.A.; Bateman, B.T.; Aspelund, T.; Skulason, S.; Zoega, H. Preeclampsia and Academic Performance in Children: A Nationwide Study from Iceland. *PLoS ONE* **2018**, *13*, e0207884. [[CrossRef](#)] [[PubMed](#)]

27. Gumusoglu, S.B.; Chilukuri, A.S.S.; Santillan, D.A.; Santillan, M.K.; Stevens, H.E. Neurodevelopmental Outcomes of Prenatal Preeclampsia Exposure. *Trends Neurosci.* **2020**, *43*, 253. [[CrossRef](#)] [[PubMed](#)]
28. Holmes, R.M.; Gardner, B.; Kohm, K.; Bant, C.; Ciminello, A.; Moedt, K.; Romeo, L. The Relationship between Young Children's Language Abilities, Creativity, Play, and Storytelling. *Early Child Dev. Care* **2019**, *189*, 244–254. [[CrossRef](#)]
29. Zhang, Y.; Chen, S.; Li, S.; Zhao, Q.; Zhou, Z.; Huang, F.; Wang, F. The Use of Internet Language Enhances Creative Performance. *J. Gen. Psychol.* **2021**, *148*, 26–44. [[CrossRef](#)]
30. Beverley Lambert, E. Children's Drawing and Painting from a Cognitive Perspective: A Longitudinal Study. *Early Years* **2005**, *25*, 249–269. [[CrossRef](#)]
31. Peng, P.; Kievit, R.A. The Development of Academic Achievement and Cognitive Abilities: A Bidirectional Perspective. *Child Dev. Perspect.* **2020**, *14*, 15–20. [[CrossRef](#)]
32. Le Guen, Y.; Amalric, M.; Pinel, P.; Pallier, C.; Frouin, V. Shared Genetic Aetiology between Cognitive Performance and Brain Activations in Language and Math Tasks. *Sci. Rep.* **2018**, *8*, 17624. [[CrossRef](#)]
33. Tucker-Drob, E.M.; Briley, D.A.; Harden, K.P. Genetic and Environmental Influences on Cognition Across Development and Context. *Curr. Dir. Psychol. Sci.* **2013**, *22*, 349–355. [[CrossRef](#)]
34. Anjos, T.; Altmäe, S.; Emmett, P.; Tiemeier, H.; Closa-Monasterolo, R.; Luque, V.; Wiseman, S.; Pérez-García, M.; Lattka, E.; Demmelmaier, H.; et al. Nutrition and Neurodevelopment in Children: Focus on NUTRIMENTHE Project. *Eur. J. Nutr.* **2013**, *52*, 1825–1842. [[CrossRef](#)]
35. Pinkerton, R.; Oriá, R.B.; Lima, A.A.M.; Rogawski, E.T.; Oriá, M.O.B.; Patrick, P.D.; Moore, S.R.; Wiseman, B.L.; Niehaus, M.D.; Guerrant, R.L. Early Childhood Diarrhea Predicts Cognitive Delays in Later Childhood Independently of Malnutrition. *Am. J. Trop. Med. Hyg.* **2016**, *95*, 1004–1010. [[CrossRef](#)] [[PubMed](#)]