

This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Ikonen, Toni; Romney, Matthew

Title: Quasiconformal geometry and removable sets for conformal mappings

Year: 2022

Version: Accepted version (Final draft)

Copyright: © 2022, The Hebrew University of Jerusalem

Rights: In Copyright

Rights url: <http://rightsstatements.org/page/InC/1.0/?language=en>

Please cite the original version:

Ikonen, T., & Romney, M. (2022). Quasiconformal geometry and removable sets for conformal mappings. *Journal d'Analyse Mathématique*, 148(1), 119-185. <https://doi.org/10.1007/s11854-022-0224-5>

QUASICONFORMAL GEOMETRY AND REMOVABLE SETS FOR CONFORMAL MAPPINGS

TONI IKONEN AND MATTHEW ROMNEY

ABSTRACT. We study metric spaces defined via a conformal weight, or more generally a measurable Finsler structure, on a domain $\Omega \subset \mathbb{R}^2$ that vanishes on a compact set $E \subset \Omega$ and satisfies mild assumptions. Our main question is to determine when such a space is quasiconformally equivalent to a planar domain. We give a characterization in terms of the notion of planar sets that are removable for conformal mappings. We also study the question of when a quasiconformal mapping can be factored as a 1-quasiconformal mapping precomposed with a bi-Lipschitz map.

CONTENTS

1. Introduction	1
2. Preliminaries	8
3. Constructing a metric from a norm field	15
4. Removable implies reciprocal	23
5. Reciprocal implies removable	29
6. Linear Cantor sets: two examples	32
7. Factoring quasiconformal mappings	34
References	46

1. INTRODUCTION

1.1. **Overview.** Let (X, d_X) and (Y, d_Y) be metric spaces with locally finite Hausdorff 2-measure. A homeomorphism $f: X \rightarrow Y$ is *K-quasiconformal* if there exists $K \geq 1$ such that

$$(1) \quad K^{-1} \operatorname{mod} \Gamma \leq \operatorname{mod} f\Gamma \leq K \operatorname{mod} \Gamma$$

for all path families Γ in X , where $\operatorname{mod} \Gamma$ denotes the conformal modulus of Γ . The map f is *quasiconformal* if it is K -quasiconformal for some $K \geq 1$. This definition is generally referred to as the *geometric definition* of quasiconformal mappings, and it is one of several possible generalizations of Euclidean quasiconformal maps to the setting of metric spaces. The definition of modulus, as well as other terms used in this introduction, is reviewed in Section 2.

The *quasiconformal uniformization problem* asks one to determine which metric spaces can be mapped onto a domain in the Euclidean plane or the 2-sphere by a mapping that is quasiconformal, according to one of the several definitions. This problem is based on the classical uniformization theorem, which states that every simply connected Riemannian 2-manifold is *conformally* equivalent to either the

2010 *Mathematics Subject Classification.* Primary 30L10. Secondary 30C35, 52A38, 53B40.

Key words and phrases. quasiconformal mappings, conformal modulus, Finsler metric.

Both authors were supported by the Academy of Finland, project number 308659. The first author was also supported by the Vilho, Yrjö and Kalle Väisälä Foundation. The second author was also supported by the Deutsche Forschungsgemeinschaft, grant SPP 2026.

Euclidean plane, the 2-sphere, or the hyperbolic plane. Outside the 2-dimensional Riemannian setting, conformality is a very strong property, and it is natural to require only quasiconformality. Motivation comes from connections to neighboring fields such as complex dynamics [BM17] and geometric group theory [Bon06].

In the following, let (X, d) be a metric space homeomorphic to a 2-dimensional manifold and having locally finite Hausdorff 2-measure. Such a space is referred to in this paper as a *metric surface*. By *quasiconformal surface*, we mean a metric surface (X, d) that is quasiconformally equivalent to a smooth Riemannian 2-manifold.

The uniformization problem for metric surfaces has been studied recently using various axiomatic approaches. Rajala has proved that a metric surface X homeomorphic to \mathbb{R}^2 is a quasiconformal surface if and only if it satisfies a condition called *reciprocity* (Definition 2.7 below) [Raj17]. Roughly speaking, this condition says that X does not have too many more rectifiable paths, as quantified by conformal modulus, than Euclidean space. In this case, as shown in [Rom19], there exists a quasiconformal map $f: X \rightarrow \Omega \subset \mathbb{R}^2$ that satisfies the modulus inequality

$$\frac{2}{\pi} \operatorname{mod} \Gamma \leq \operatorname{mod} f\Gamma \leq \frac{4}{\pi} \operatorname{mod} \Gamma$$

for all path families Γ in X . This inequality is sharp, as can be shown by considering the plane equipped with either the $\|\cdot\|_1$ - or $\|\cdot\|_\infty$ -norm. These results are extended to arbitrary metric surfaces in [Iko19]. A different approach was taken in a series of papers of Lytchak and Wenger [LW17], [LW18], [LW20] based on the assumption that the space satisfies a *quadratic isoperimetric inequality*.

The goal of the present paper is to understand the uniformization results described above in the context of concrete constructions of metric surfaces. We study a general scheme for constructing surfaces based on specifying a measurable Finsler structure on a planar domain that vanishes on some subset of the plane. The natural problem is to decide when this construction yields a quasiconformal surface.

We provide an answer by linking the uniformization problem for metric surfaces to a separate topic in complex analysis: removable sets for classes of holomorphic functions. There are several notions of removability; see [You15] for a recent survey. For us, the relevant definition is the following. A compact set $E \subset \mathbb{R}^2$ is *removable for conformal mappings* if every conformal embedding $f: \mathbb{R}^2 \setminus E \rightarrow \widehat{\mathbb{R}}^2$ extends to a conformal mapping $\tilde{f}: \widehat{\mathbb{R}}^2 \rightarrow \widehat{\mathbb{R}}^2$, that is, to a Möbius transformation. Here, $\widehat{\mathbb{R}}^2$ denotes the extended plane, which can be identified with \mathbb{S}^2 via stereographic projection. There seems to be no standard terminology for sets satisfying this condition. This is referred to as *S-removability* in the survey [You15], while the terms *set of absolute area zero* and *negligible set for extremal distance* are also used. Note that this is different than the notion of *conformal removability*, which requires that every *homeomorphism* of $\widehat{\mathbb{R}}^2$ that is conformal on the set $\widehat{\mathbb{R}}^2 \setminus E$ be a Möbius transformation.

This connection to removable sets is natural in hindsight but does not appear to have been made before. On the other hand, removable sets are inherently connected to a different type of uniformization problem, namely of multiply connected planar domains onto some canonical class of domain, typically slit domains or circle domains. We recall that whether an arbitrary planar domain can be mapped conformally onto a circle domain is the well-known *Koebe Kreisnormierungsproblem* [HS93]. We hope the present paper will add a new perspective on these various topics.

1.2. Motivating examples. A basic observation, made in Example 2.1 in [Raj17], is that not every metric surface is a quasiconformal surface. A simple example is the following. Define a length pseudometric d_ω on \mathbb{R}^2 via the conformal weight

$\omega = \chi_{\mathbb{R}^2 \setminus \mathbb{D}}$. More precisely, we define the ω -length of an absolutely continuous path γ to be $\ell_\omega(\gamma) = \int_\gamma \omega ds$, and let $d_\omega(x, y) = \inf \ell_\omega(\gamma)$, the infimum taken over all absolutely continuous paths γ connecting x and y . If we let X be the quotient space of \mathbb{R}^2 formed by collapsing the unit disk to a single point, then d_ω induces a metric on X , denoted by \tilde{d}_ω , that is locally Euclidean outside the origin. The space (X, \tilde{d}_ω) , while being homeomorphic to \mathbb{R}^2 , is not quasiconformally equivalent to a planar domain. This is because the family of paths in X that intersect the collapsed point has positive modulus, while the modulus of the family of paths intersecting a single point in the Euclidean plane is zero. This example is included as Example 11.3 in [LW18].

A second example, and the one that comprises Example 2.1 in [Raj17], is a continuous conformal weight ω that vanishes on a Cantor set E of positive area. In this case, d_ω is a metric on \mathbb{R}^2 , and the identity map $(\mathbb{R}^2, \|\cdot\|_2) \rightarrow (\mathbb{R}^2, d_\omega)$ is a homeomorphism. Nevertheless, the vanishing of the weight increases the conformal modulus of path families in (\mathbb{R}^2, d_ω) in a way incompatible with admitting a quasiconformal parametrization by \mathbb{R}^2 .

At the other extreme, it is not hard to show that if the analogous construction is carried out for a set E with Hausdorff dimension smaller than one, then the resulting space is quasiconformally equivalent to the plane. Indeed, the set E is then negligible for length and so has no effect on modulus. What happens in the intermediate situation—when the Hausdorff dimension satisfies $1 \leq \dim_{\mathcal{H}} E < 2$ or when $\mathcal{H}^2(E) = 0$ —is not *a priori* clear and is one of the motivations of our work.

Similar constructions appear in a number of related contexts. One of these is the notion of *strong A_∞ -weight* introduced by David and Semmes in [DS90]. Such a weight determines a metric on \mathbb{R}^2 that is Ahlfors 2-regular and quasisymmetrically equivalent to the plane. Conversely, the Jacobian of a quasisymmetric mapping from \mathbb{R}^2 to an Ahlfors 2-regular metric space induces a strong A_∞ -weight on \mathbb{R}^2 . We do not define this term here but refer the reader to [Sem96, Def. 1.5]. Such weights appear naturally when trying to recognize metric spaces that are bi-Lipschitz embeddable in some Euclidean space. See [DS90, Sem93, Sem96, Laa02, Bis07] for various contributions to this topic. A separate set of papers [BKR98, BHR01] studies metrics on the unit disk defined by conformal weights satisfying a Harnack-type inequality and an area growth condition, and shows that a number of results of classical complex analysis have natural analogues in this setting. All of the metric surfaces constructed in these two sets of papers are quasiconformally equivalent to a planar domain.

In the above examples, when a space fails to be a quasiconformal surface, this is due to the space “collapsing” on the set E where the weight vanishes. In fact, it may be the case that this is essentially the only way that a metric surface can fail to admit a quasiconformal parametrization. This is made precise by the following question of Rajala and Wenger.

Question 1.1. Let (X, d) be a metric space homeomorphic to \mathbb{R}^2 with locally finite Hausdorff 2-measure. Is there in general a domain $\Omega \subset \mathbb{R}^2$ and a surjective continuous monotone mapping $f: \Omega \rightarrow X$ such that f is in the metric Sobolev space $N_{\text{loc}}^{1,2}(\Omega, X)$ and satisfies the one-sided dilatation condition

$$g_f^2(x) \leq K J_f(x)$$

for some constant $K \geq 1$ and almost every $x \in \Omega$?

Here, g_f is the minimal weak upper gradient of f and J_f is the Jacobian of f ; see Section 2.2. We say that $f: \Omega \rightarrow X$ is *monotone* if the preimage of every point $x \in X$ is a connected and compact subset of Ω .

1.3. Setting and main results. We continue with a description of our setting and main results. Let Ω be a planar domain and $E \subset \Omega$ be a compact set that does not separate Ω . We consider a measurable seminorm field $N: \Omega \times \mathbb{R}^2 \rightarrow [0, \infty)$ that vanishes exactly on the set E and satisfies certain mild assumptions, namely lower semicontinuity, local boundedness, and having locally bounded distortion. The seminorm at the point $x \in \Omega$ is denoted throughout this paper by N_x . We think of N as a Finsler structure on \mathbb{R}^2 , determining a Finsler metric on \mathbb{R}^2 , although requiring no regularity beyond the previous assumptions.

For conciseness, and since N_x is a norm for all $x \in \Omega \setminus E$, we use the term *norm field* and not *seminorm field* throughout this paper when referring to N . A norm field N satisfying the above hypotheses is said to be *admissible* (Definition 3.1 below). We define the N -length of an absolutely continuous path $\gamma: I \rightarrow \Omega$ by

$$(2) \quad \ell_N(\gamma) = \int_I N \circ D\gamma(t) dt.$$

In interpreting (2), note that the base point of N is understood to be $\gamma(t)$ even though this is omitted from the notation. One then obtains a pseudometric d_N on Ω by setting $d_N(x, y) = \inf \ell_N(\gamma)$, the infimum taken over all absolutely continuous paths γ from x to y contained in Ω . Let \mathcal{E}_N denote the collection of equivalence classes of points in \mathbb{R}^2 , declaring x to be equivalent to y if $d_N(x, y) = 0$. Then d_N determines a metric on the quotient space $\mathbb{R}^2/\mathcal{E}_N$ denoted by \tilde{d}_N . In Section 3, we describe this construction in more detail.

We make the following definition.

Definition 1.2. The admissible norm field N is *reciprocal* if the corresponding space $(\Omega/\mathcal{E}_N, \tilde{d}_N)$ is reciprocal (Definition 2.7).

The natural problem is to characterize as best as possible those norm fields N that are reciprocal. Our first result is the following.

Theorem 1.3. Let $\Omega \subset \mathbb{R}^2$ be a domain and $E \subset \Omega$ a compact set. If E is removable for conformal mappings, then every admissible norm field $N: \Omega \times \mathbb{R}^2 \rightarrow [0, \infty)$ that vanishes exactly on E is reciprocal.

Recall that our definition of admissibility includes the statement that N is locally bounded. It turns out that this assumption can be relaxed. In Proposition 4.5, we show that Theorem 1.3 still holds provided there exists some $p > 2$ such that the maximal stretching $L(N)$ is in $L^p_{\text{loc}}(\Omega)$. This generalization follows fairly readily from Theorem 1.3 by an approximation argument.

Next, we consider whether some converse to Theorem 1.3 holds. Observe first that the strongest possible converse to Theorem 1.3 is false: a reciprocal norm field N may vanish on a set E that is not removable for conformal mappings. As a simple example, take $E \subset \mathbb{R}^2$ to be a snowflake arc and let $N = \chi_{\mathbb{R}^2 \setminus E} \cdot \|\cdot\|_2$. Since $\mathcal{H}^1_{\|\cdot\|_2}(|\gamma| \cap E) = 0$ for every absolutely continuous path γ , we see that d_N actually coincides with the Euclidean metric. However, it is a basic fact that any set that is removable for conformal mappings is totally disconnected.

On the other hand, if one requires that the norm field N decays fast enough near E and N is reciprocal, then examples of the type just described are not possible. To illustrate this, consider two admissible norm fields N_1 and N_2 that satisfy $N_1 \leq N_2$. Every path that has finite N_2 -length also has finite N_1 -length, while the opposite may fail to be true for a large family of paths. In this sense, the space generated by the smaller norm field N_1 has more rectifiable paths and the reciprocity condition is harder to satisfy. This leads to the following partial converse to Theorem 1.4.

Theorem 1.4. Let $\Omega \subset \mathbb{R}^2$ be a domain and $E \subset \Omega$ a compact set for which $\Omega \setminus E$ is connected, and let $N_p(x) = \min \{1, d_{\|\cdot\|_2}(x, E)^p\} \|\cdot\|_2$. If N_p is reciprocal for some $p > \max \{\dim_{\mathcal{H}} E - 1, 0\}$, then the set E is removable for conformal mappings.

Our method of proof actually yields a stronger conclusion. The relevant property of the norm field N_p , verified in Lemma 5.1 below, is that the quotient map π_N maps E onto a set of zero 1-dimensional Hausdorff measure with respect to the metric \tilde{d}_N . Thus, for any reciprocal norm field N with this property, the corresponding set E on which N vanishes is removable for conformal mappings. For example, one can show that, if E is contained in a continuum F satisfying $\mathcal{H}_{\|\cdot\|_2}^1(F) < \infty$, this condition on the quotient map π_N is satisfied for all admissible norm fields N vanishing on E . For such compact sets, the strongest converse to Theorem 1.3 holds. That is, if any admissible norm field N vanishing exactly on E is reciprocal, then E is removable for conformal mappings.

The lower bound for p in Theorem 1.4 is sharp. Consider an arc $E \subset \mathbb{R}^2$ that is bi-Lipschitz equivalent to $([0, 1], |\cdot|^{1/d})$ for some $d \in (1, 2)$. Then E is a snowflake arc of Hausdorff dimension d . It follows from [Sem96, Theorem 6.3] that the square of the weight $\omega_{d-1}(x) = \min \{1, d_{\|\cdot\|_2}(x, E)^{d-1}\}$ is a strong A_∞ -weight, as defined in [Sem96, Definition 1.5], and hence the norm field N_{d-1} is reciprocal. However, the arc E is not removable for conformal mappings.

Theorems 1.3 and 1.4 show that reciprocal norm fields are almost characterized by whether the set on which they vanish is removable for conformal mappings. We now mention a few facts about removable sets for conformal mappings that are known, many of them coming from a classic paper of Ahlfors–Beurling [AB50]. First, every compact set of positive Hausdorff 2-measure is not removable. Second, every compact set of zero Hausdorff 1-measure is removable. More intriguingly, for Cantor sets $E \subset \mathbb{R} \times \{0\}$ of positive Hausdorff 1-measure, both outcomes are possible. In [AB50], Ahlfors and Beurling construct Cantor sets in $\mathbb{R} \times \{0\}$ of positive \mathcal{H}^1 -measure that are removable for conformal maps, as well as such Cantor sets that are not removable. A similar example in the related context of circle domain uniformization can be found as Theorem 11.1 of an early version of a paper of Schramm [Sch95]. Next, by Theorem 10 in [AB50] and Proposition 3.3 in [KKR19], removable sets for conformal mappings are *metrically removable*: for every $\varepsilon > 0$, each pair of points $x, y \in \mathbb{R}^2$ can be connected by a curve disjoint from $E \setminus \{x, y\}$ that has length at most $\|x - y\|_2 + \varepsilon$. See [HH08] and [KKR19] for more on the topic of metric removability. Removable sets for conformal mappings are also examples of the *quasiextremal distance exceptional sets* considered in [GM85] and the related literature. Finally, an equivalent definition can be given by replacing the word “conformal” with “quasiconformal” in the definition [You15, Prop. 4.7]. Thus the property of removability is invariant under quasiconformal mappings of the complementary domain.

This should be compared with the notion of *removable sets for bounded analytic functions*. The problem of characterizing such sets is known as *Painlevé’s problem* and has received considerable attention, with a satisfactory resolution obtained by Tolsa in [Tol03]. We note here that this is a stronger notion of removability: every set that is removable for bounded analytic functions is removable for conformal mappings. See Proposition 4.3 of [You15] for a proof. For example, a removable set for bounded analytic functions must have Hausdorff dimension at most 1. Moreover, according to David’s resolution of Vitushkin’s conjecture [Dav98], a compact set E with finite Hausdorff 1-measure is removable for bounded analytic functions if and only if it is purely 1-unrectifiable.

Finally, we remark that the notion of uniformly disconnected sets provides a further class of examples to which these results apply. In [Sem96], Semmes studies

metrics of the form d_{N_p} , where N_p is as in Theorem 1.4, with the additional assumption that the set E is *uniformly disconnected*, meaning that there exists $\varepsilon > 0$ with the property that, for any two distinct points $x, y \in E$, there is no sequence of points $x = x_0, x_1, \dots, x_m = y$ in E satisfying $\|x_{j-1} - x_j\|_2 < \varepsilon \|x - y\|_2$ for all $j \in \{1, \dots, m\}$. He proves that for such an E and every $p > 0$, the square of the weight $\omega_p(x) = \min\{1, d_{\|\cdot\|_2}(x, E)^p\}$ is a strong A_∞ -weight and hence the norm field N_p in Theorem 1.4 is reciprocal. Therefore Theorem 1.4 implies that uniformly disconnected Cantor sets are removable for conformal mappings. This removability can alternatively be deduced in many ways from the existing literature. Note in particular that a uniformly disconnected set E can have Hausdorff dimension arbitrarily close to 2.

1.4. Factorization of quasiconformal mappings. This section is motivated by the following factorization problem. Consider a quasiconformal surface (X, d) and corresponding isothermal parametrization $f: \Omega \rightarrow X$, where Ω is a smooth Riemannian surface. Following [Iko19], a quasiconformal homeomorphism $f: \Omega \rightarrow X$ is *isothermal* if it is distortion-minimizing at almost every point in a suitable sense. Roughly speaking, the *pointwise distortion* of f at x is the aspect ratio of the image of a small ball centered at x . The existence of an isothermal parametrization for every quasiconformal surface is established in [Iko19, Corollary 6.3]. See Section 2.5 for the precise definition of distortion and Section 7.1 for the definition of isothermal map. We ask: can one find a metric surface $(\widehat{X}, \widehat{d})$ such that f factors as $f = \widehat{f} \circ P$, where $\widehat{f}: \widehat{X} \rightarrow X$ is 1-quasiconformal and $P: \Omega \rightarrow \widehat{X}$ is bi-Lipschitz? In other words, can one find a ‘‘conformal representative’’ for the space X within the class of bi-Lipschitz surfaces?

If the metric is defined by a continuous reciprocal norm field of bounded distortion, then such a factorization can always be found. Recall that for every domain $\Omega \subset \mathbb{R}^2$ there exists a smooth Riemannian norm field $G = \omega \|\cdot\|_2$ on Ω such that (Ω, d_G) is complete and has Gaussian curvature 0 or -1 . We have the following result.

Proposition 1.5. Let $\Omega \subset \mathbb{R}^2$ be a domain and N a reciprocal norm field with distortion H . If \widehat{N} is continuous outside the set $E = \{x \in \Omega : N_x = 0\}$, then there exists a distance \widehat{d} on Ω such that:

- (i) The identity map $P: (\Omega, d_G) \rightarrow (\Omega, \widehat{d})$ satisfies
- $$(3) \quad d_G(x, y) \leq \widehat{d}(P(x), P(y)) \leq H d_G(x, y)$$
- for all $x, y \in \Omega$.

- (ii) The identity map $\widehat{\iota}: (\Omega, \widehat{d}) \rightarrow (\Omega, d_N)$ is 1-quasiconformal.

If the identity map $\iota: \Omega \rightarrow (\Omega, d_N)$ is isothermal, then it has distortion at most $\sqrt{2}$ [Iko19, Cor. 4.7], and so (3) holds with $H = \sqrt{2}$. The example of the ℓ^∞ -norm on \mathbb{R}^2 shows that the value $H = \sqrt{2}$ in (3) is sharp for the case of general isothermal maps. Since every quasiconformal surface has an isothermal parametrization, this raises the question of finding conditions on N that guarantee that the conclusion of Proposition 1.5 holds with $H = \sqrt{2}$. In turn, this question is related to the regularity of the Beltrami coefficient derived from distance ellipse field corresponding to N and does not appear to have a straightforward answer. We briefly address this issue in Section 7.3.

In general, the conclusion of Proposition 1.5 may fail if N_x is discontinuous outside of E . In the final part of the paper, we present a lengthy construction giving a negative answer to the above factorization question in general. In fact, we obtain the stronger conclusion that no quasiconformal map \widehat{f} in such a factorization can have distortion smaller than that of f .

Theorem 1.6. There is a metric d on \mathbb{R}^2 such that the identity map $\iota: (\mathbb{R}^2, \|\cdot\|_2) \rightarrow (\mathbb{R}^2, d)$ is an isothermal quasiconformal homeomorphism, but ι does not factor as $\iota = \widehat{\iota} \circ P$, where $(\widehat{X}, \widehat{d})$ is a metric surface, $\widehat{\iota}: (\widehat{X}, \widehat{d}) \rightarrow (\mathbb{R}^2, d)$ is quasiconformal with distortion $H(\widehat{\iota}) < \sqrt{2}$ and $P: (\mathbb{R}^2, \|\cdot\|_2) \rightarrow (\widehat{X}, \widehat{d})$ is bi-Lipschitz.

The identity map ι in our construction has distortion $H(\iota) = \sqrt{2}$, so the inequality $H(\widehat{\iota}) < \sqrt{2}$ is sharp.

The metric d in Theorem 1.6 is defined via a lower semicontinuous norm field of the form

$$N_x = \begin{cases} c_x \|\cdot\|_1 & \text{if } x \in F \\ c_x \|\cdot\|_\infty & \text{if } x \notin F \end{cases}$$

for some measurable set $F \subset \mathbb{R}^2$ and measurable function $x \mapsto c_x$, where $0 \leq c_x \leq 1$ and c_x vanishes at a single point. Note that this fits exactly into the construction scheme of this paper, and therefore (\mathbb{R}^2, d) is a quasiconformal surface.

One might initially expect that the metric \widehat{d} on \mathbb{R}^2 defined by

$$\widehat{N}_x = \begin{cases} \|\cdot\|_1 & \text{if } x \in F \\ \sqrt{2} \|\cdot\|_\infty & \text{if } x \notin F \end{cases}$$

with $\widehat{\iota}$ and P the identity map on \mathbb{R}^2 , or some variation on this, gives a factorization satisfying the properties given in Theorem 1.6. Observe that $\|\cdot\|_2 \leq \widehat{N} \leq \sqrt{2} \|\cdot\|_2$ everywhere, so the map P in this situation is bi-Lipschitz. However, the map $\widehat{\iota}$ may fail to be 1-quasiconformal. The reason for this is that the norm field \widehat{N} corresponding to F is typically not lower semicontinuous, in which case the metric tangents of P need not coincide with \widehat{N}_x almost everywhere. Indeed, we prove Theorem 1.6 by specifying explicitly a set F and coefficients c_x for which this failure of 1-quasiconformality occurs for the norm field \widehat{N} defined above, and in fact for any conformal rescaling of \widehat{N} bi-Lipschitz equivalent to the Euclidean norm field.

We now describe our construction in somewhat more detail. The basic idea is to construct a sequence of nested Cantor sets K_i as the intersection of a collection of squares in the plane. This is done so that the odd-indexed Cantor sets are formed from squares in the standard (i.e., non-rotated) alignment, while the even-indexed Cantor sets are formed from squares aligned diagonally. Next, the norm field on $K_i \setminus K_{i+1}$ for odd values of i is defined to be the supremum norm $\|\cdot\|_\infty$, scaled by a constant c_i satisfying $c_i \rightarrow 0$ as $i \rightarrow \infty$, while the norm field for even values of i is defined to be the $\|\cdot\|_1$ -norm, also scaled by a constant c'_i satisfying $c'_i \rightarrow 0$ as $i \rightarrow \infty$. A consequence of the distortion inequality for $\widehat{\iota}$ is that the metric tangents of P and ι cannot differ by more than a fixed amount, up to rescaling. With a suitable choice of constants c_i, c'_i , the alternating arrangement of the Cantor sets K_i then forces the metric tangents of P to be arbitrarily small at some points.

Lytchak–Wenger [LW18] and Creutz–Soultanis [CS19] study similar types of factorizations for *minimal disks* or *solutions to Plateau’s problem* with metric space target, though without trying to optimize the properties of P in the way that we have proposed. Here, we simply remark that the map ι in our example is also an energy-minimizing map (for the Reshetnyak energy) in the sense of these papers on each closed disk. We refer the reader to the above papers for definitions of these terms.

1.5. Outline. Our paper is organized as follows. Section 2 gives an overview of basic results and notation related to metric Sobolev spaces, quasiconformal mappings, and removable sets. In Section 3, we give a detailed overview of the construction of metric spaces from a prescribed norm field under suitable assumptions. In Section 4,

we prove the first of the main results, Theorem 1.3, stating that an admissible norm field is reciprocal if it vanishes exactly on a set that is removable for conformal mappings. In Section 5, we prove the partial converse, Theorem 1.4. Section 6 gives a pair of examples of spaces constructed from conformal weights that each vanish on a linear Cantor set of positive length, one of which is reciprocal and one of which is not. Thus this can be viewed as the borderline case. Finally, Section 7 gives the proof of Proposition 1.5 as well as the construction for Theorem 1.6.

Acknowledgments. We are thankful to Alexander Lytchak, Kai Rajala, and Atte Lohvansuu for discussions about this project and feedback on a draft of this paper. We also thank Dimitrios Ntalampekos and Malik Younsi for discussions related to removable sets for conformal mappings, and in particular for Malik Younsi bringing the paper [AB50] to our attention. Finally, we thank Jarmo Jääskeläinen for discussions related to the Beltrami equation.

2. PRELIMINARIES

2.1. Notation. In this paper, we frequently consider several metrics in close proximity to one another. For this reason, we will consistently use subscripts to denote the metric being referred to. Let (X, d) be a metric space. The open ball centered at a point $x \in X$ of radius $r > 0$ with respect to the metric d is denoted by $B_d(x, r)$.

The Euclidean metric is denoted by $\|\cdot\|_2$. Thus, for example, we write $B_{\|\cdot\|_2}(x, r)$ for an open ball with respect to this metric, and $ds_{\|\cdot\|_2}$ for the Euclidean length element.

We recall the definition of Hausdorff measure. Let (X, d) be a metric space. For all $p \geq 0$, the p -dimensional Hausdorff measure, or Hausdorff p -measure, is defined by

$$\mathcal{H}_X^p(A) = \sup_{\delta > 0} \inf \left\{ \frac{\alpha(p)}{2^p} \sum_{i=1}^{\infty} (\text{diam } A_i)^p : A \subset \bigcup_{i=1}^{\infty} A_i, \text{diam } A_i < \delta \right\}$$

for all sets $A \subset X$, where $\alpha(p) = \pi^{\frac{p}{2}} (\Gamma(\frac{p}{2} + 1))^{-1}$. The constant $\alpha(p)$ is chosen in such a way that $\mathcal{H}_{\mathbb{R}^n}^n$ coincides with the Lebesgue measure \mathcal{L}^n for all positive integers.

If the space X is understood but not the metric d , then we use the notation \mathcal{H}_d^p instead of \mathcal{H}_X^p . The Hausdorff dimension of a set $E \subset X$ is the infimal value of p for which $\mathcal{H}_X^p(E) = 0$ and is denoted by $\dim_{\mathcal{H}_d} E$. For the basics of Hausdorff measure, see for example [AT04, Chapter 2].

Unless otherwise noted, in this paper a metric surface (X, d) is always equipped with the Hausdorff 2-measure generated by the metric d . For example, the phrase *almost every* refers to the Hausdorff 2-measure. Similarly, an interval in \mathbb{R} is equipped with the Lebesgue measure \mathcal{L}^1 .

A *path* is a continuous function from an interval into a metric space. A path in X will typically be denoted by γ . The image of γ is denoted by $|\gamma|$. The *length* of the path $\gamma: [a, b] \rightarrow X$ is defined as

$$\ell_d(\gamma) = \sup \sum_{j=1}^n d(\gamma(t_{j-1}), \gamma(t_j)),$$

the supremum taken over all finite sequences $a = t_0 \leq t_1 \leq \dots \leq t_n = b$. A path is *rectifiable* if it has finite length.

The *metric speed* of a path $\gamma: [a, b] \rightarrow X$ at the point $t \in [a, b]$ is defined as

$$v_\gamma(t) = \lim_{h \rightarrow 0} \frac{d(\gamma(t+h), \gamma(t))}{h}$$

whenever this limit exists. If γ is rectifiable, its metric speed exists at \mathcal{L}^1 -almost every $t \in [a, b]$; see Theorem 2.1 of [Dud07].

A rectifiable path $\gamma: [a, b] \rightarrow X$ is *absolutely continuous* if for all $a \leq s \leq t \leq b$,

$$d(\gamma(t), \gamma(s)) \leq \int_s^t v_\gamma(u) d\mathcal{L}^1(u)$$

with $v_\gamma \in L^1([a, b])$ and \mathcal{L}^1 the Lebesgue measure on the real line. Equivalently, γ is absolutely continuous if it maps sets of \mathcal{L}^1 -measure zero to sets of \mathcal{H}_X^1 -measure zero in its image; see Section 3 of [Dud07].

A path $\tilde{\gamma}: [c, d] \rightarrow X$ is a *reparametrization* of γ if there exists a map $\psi: [a, b] \rightarrow [c, d]$ that is surjective, non-decreasing, and continuous such that $\gamma = \tilde{\gamma} \circ \psi$. If ψ is absolutely continuous, we say that $\tilde{\gamma}$ is an *absolutely continuous reparametrization* of γ . Note that this is different from $\tilde{\gamma}$ itself being an absolutely continuous path.

Every rectifiable path γ has a reparametrization $\tilde{\gamma}: [0, \ell_d(\gamma)] \rightarrow X$ such that the metric speed of $\tilde{\gamma}$ equals one \mathcal{L}^1 -almost everywhere. In this case, we write $\gamma_s = \tilde{\gamma}$, and refer to γ_s as the *unit speed parametrization* of γ . See Chapter 5 of [HKST15] for details.

If γ is rectifiable, the unit speed parametrization γ_s is 1-Lipschitz and hence absolutely continuous [HKST15, Proposition 5.1.8].

Let γ be a rectifiable path. Then the *path integral* of a Borel function $\rho: X \rightarrow [0, \infty]$ over γ is

$$(4) \quad \int_\gamma \rho ds = \int_0^{\ell_d(\gamma)} \rho \circ \gamma_s d\mathcal{L}^1,$$

where \mathcal{L}^1 is the Lebesgue measure on the real line.

If γ is absolutely continuous and $\tilde{\gamma}$ is an absolutely continuous reparametrization of γ , the chain rule for metric speeds [Dud07, Theorem 3.16 and Remark 3.4] states that

$$v_\gamma = (v_{\tilde{\gamma}} \circ \psi) \psi' \in L^1([c, d]),$$

where the right-hand side is understood to be zero whenever the derivative $\psi' = 0$ (even if $v_{\tilde{\gamma}} \circ \psi$ is not defined or is infinite at such a point).

Moreover, for absolutely continuous γ , the unit speed parametrization γ_s is an absolutely continuous reparametrization of γ . Therefore (4) can be restated for absolutely continuous $\gamma: [a, b] \rightarrow X$ as follows:

$$\int_\gamma \rho ds = \int_a^b (\rho \circ \gamma) v_\gamma d\mathcal{L}^1.$$

Given a Borel set $A \subset X$, the length of a path $\gamma: [a, b] \rightarrow X$ in A is defined as $\int_X \#(A \cap \gamma^{-1}(x)) d\mathcal{H}_X^1(x)$, where $\#(A \cap \gamma^{-1}(x))$ is the multiplicity of γ in A . This formula makes sense for paths that are not necessarily rectifiable; see Theorem 2.10.13 [Fed69]. If γ is rectifiable, the number coincides with the path integral of χ_A over γ .

2.2. Metric Sobolev spaces. In this section, we give an overview of the theory of Sobolev spaces in the metric space setting. We refer the reader to the book [HKST15] for a comprehensive introduction to this topic. Throughout this section, assume that (X, d_X) and (Y, d_Y) are each a metric surface: a metric space homeomorphic to a 2-dimensional manifold with locally finite Hausdorff 2-measure.

The conformal modulus provides a basic way of measuring the size of a family of paths. It is a conformal invariant in the Euclidean case, which accounts for both its nomenclature and its usefulness. Let Γ be a family of paths in X . A Borel function $\rho: X \rightarrow [0, \infty]$ is *admissible* for the path family Γ if the path integral $\int_\gamma \rho ds \geq 1$

for all locally rectifiable paths $\gamma \in \Gamma$. The *conformal modulus*, or simply *modulus*, of Γ is

$$\text{mod } \Gamma = \inf \int_X \rho^2 d\mathcal{H}_X^2,$$

where the infimum is taken over all admissible functions ρ .

If ρ is admissible for a path family $\Gamma' \subset \Gamma$ such that $\Gamma \setminus \Gamma'$ has modulus zero, then ρ is said to be *weakly admissible* for Γ . If a property holds for every path $\gamma \in \Gamma$ except in a subfamily of modulus zero, then this property is said to hold *on almost every path*. If $\text{mod } \Gamma < \infty$, then there exists a weakly admissible Borel function $\rho \in L^2(X)$ such that

$$\text{mod } \Gamma = \int_X \rho^2 d\mathcal{H}_X^2.$$

Such a ρ is called a *minimizer* of Γ . Such a minimizer is unique \mathcal{H}_X^2 -almost everywhere.

Let $f: (X, d_X) \rightarrow (Y, d_Y)$ be a mapping between metric surfaces X and Y . A function $g: X \rightarrow [0, \infty]$ is an *upper gradient* of f if

$$d_Y(f(x), f(y)) \leq \int_\gamma g ds$$

for every rectifiable path $\gamma: [0, 1] \rightarrow X$ connecting x to y . The function g is a *weak upper gradient* of f if the same holds for almost every rectifiable path.

The weak upper gradient $g \in L_{\text{loc}}^2(X)$ is *minimal* if it satisfies $g \leq \tilde{g}$ almost everywhere for all weak upper gradients $\tilde{g} \in L_{\text{loc}}^2(X)$ of f . If f has a weak upper gradient $g \in L_{\text{loc}}^2(X)$, then f has a minimal weak upper gradient, which we denote by g_f . The existence of g_f follows from the fact that the weak upper gradients of f form a lattice. This also implies that g_f is unique up to measure zero; see Section 6 of [HKST15] and Section 3 of [Wil12] for details. In general, g_f is only a weak upper gradient.

Proposition 6.3.3 of [HKST15] and countable subadditivity of modulus (see also Lemmas 3.2 and 3.3 of [Wil12]) establish that a Borel function $\rho: X \rightarrow [0, \infty]$ belonging to $L_{\text{loc}}^2(X)$ is a weak upper gradient of f if and only if for almost every absolutely continuous path $\gamma: [a, b] \rightarrow X$, the composition $f \circ \gamma$ is an absolutely continuous path for which the metric speeds $v_{f \circ \gamma}$ and v_γ satisfy

$$(5) \quad v_{f \circ \gamma} \leq (\rho \circ \gamma) v_\gamma$$

\mathcal{L}^1 -almost everywhere on $[a, b]$. Since $\rho \in L_{\text{loc}}^2(X)$ the right-hand side of (5) is integrable on its domain for almost every γ .

Let Z be a metric space such that $\mathcal{H}_{d_Z}^2(Z) < \infty$. Choose a point $y \in Y$, and let $d_y = d_Y(\cdot, y)$. The space $L^2(Z, Y)$ is defined as the set of measurable mappings $f: Z \rightarrow Y$ such that $d_y \circ f$ is in $L^2(Z)$. One can check that this definition is independent of the choice of y .

We define $L_{\text{loc}}^2(X, Y)$ to consist of those measurable mappings $f: X \rightarrow Y$ for which, for all $x \in X$, there is an open set $U \subset X$ containing x such that $f|_U$ is in $L^2(U, Y)$.

The metric Sobolev space $N_{\text{loc}}^{1,2}(X, Y)$ consists of those mappings $f: X \rightarrow Y$ in $L_{\text{loc}}^2(X, Y)$ that have a minimal weak upper gradient $g_f \in L_{\text{loc}}^2(X)$.

For open $U \subset X$ with $\mathcal{H}_X^2(U) < \infty$, we say that $f \in N^{1,2}(U, Y)$ if $f|_U \in N_{\text{loc}}^{1,2}(U, Y)$ in such a way that $g_f|_U \in L^2(U)$ and for some $y \in Y$, $f_y(x) = d_y \circ f|_U \in L^2(U)$.

Next we define the Jacobian of f for continuous $f: X \rightarrow Y$. The *pullback measure* $f^*\mathcal{H}_Y^2$ is defined for Borel sets $A \subset X$ by the formula

$$f^*\mathcal{H}_Y^2(A) = \int_Y \#(A \cap f^{-1}(y)) d\mathcal{H}_Y^2,$$

where $\#(A \cap f^{-1}(y))$ is the multiplicity function of f relative to A . The measure $f^*\mathcal{H}_Y^2$ can be defined equivalently using a suitable Carathéodory construction; see [Fed69, 2.10.10]. In fact, $f^*\mathcal{H}_Y^2$ is a Borel regular outer measure.

If the pullback measure $f^*\mathcal{H}_Y^2$ is locally finite, we say that the *Jacobian* of f is the Radon–Nikodym derivative of $f^*\mathcal{H}_Y^2$ with respect to \mathcal{H}_X^2 . The Jacobian is denoted by J_f . The local finiteness of $f^*\mathcal{H}_Y^2$ and \mathcal{H}_X^2 imply that J_f is locally integrable. See Sections 3.1–3.2 in Volume I of [Bog07] for details on the Radon–Nikodym derivative of a measure.

2.3. Seminorms. We introduce the terminology and notation we use for seminorms. Recall that a *seminorm* S on \mathbb{R}^2 is a function $S: \mathbb{R}^2 \rightarrow [0, \infty)$ satisfying the following conditions for all $v, w \in \mathbb{R}^2$ and $\lambda \in \mathbb{R}$:

- (i) (absolute homogeneity) $S(\lambda v) = |\lambda|S(v)$ whenever $\lambda \in \mathbb{R}$ and $v \in \mathbb{R}^2$;
- (ii) (triangle inequality) $S(v + w) \leq S(v) + S(w)$.

The seminorm S is a *norm* if it has the additional property that $S(v) = 0$ only if $v = 0$. The *maximal stretching* of S is

$$(6) \quad L(S) = \sup \{S(v) : \|v\|_2 \leq 1\}.$$

The *minimal stretching* of S is

$$(7) \quad \omega(S) = \inf \{S(v) : \|v\|_2 \geq 1\}.$$

The *Jacobian* of the seminorm S is

$$J_2(S) = \frac{\pi}{\mathcal{L}^2(\{v : S(v) \leq 1\})}.$$

Observe that $J_2(S) = 0$ in the case that N is only a seminorm. The *distortion* of S is

$$(8) \quad H(S) = \frac{L(S)}{\omega(S)}$$

if $\omega(S) > 0$ and $H(S) = \infty$ otherwise. The latter case occurs if S is a non-zero seminorm that is not a norm. The *outer dilatation* and *inner dilatation* of S are defined by, respectively,

$$K_O(S) = \frac{L(S)^2}{J_2(S)}, \quad K_I(S) = \frac{J_2(S)}{\omega(S)^2}$$

if $J_2(S) > 0$, and $K_O(S) = K_I(S) = \infty$ otherwise. The *maximal dilatation* of S is $K(S) = \max\{K_O(S), K_I(S)\}$. Observe that $K_O(S) \geq 1$ and $K_I(S) \geq 1$.

The seminorm S induces a pseudometric d_S on \mathbb{R}^2 by the formula $d_S(x, y) = S(x - y)$. The identity map $\iota_S: (\mathbb{R}^2, \|\cdot\|_2) \rightarrow (\mathbb{R}^2, d_S)$ has the constant function $L(S)$ as its minimal weak upper gradient and $J_2(S)$ as its Jacobian. Its inverse ι_S^{-1} has the constant function $\omega(S)^{-1}$ as its minimal weak upper gradient.

The following lemma gives a relationship between the maximal dilatation and distortion.

Lemma 2.1. The distortion $H(S)$ and maximal dilatation $K(S)$ of S satisfy $H(S) \leq K(S) \leq H(S)^2$.

Proof. If $\omega(S) = 0$, then $H(S) = K(S) = \infty$. Otherwise, $H(S)$ and $K(S)$ are both finite. Observe the relationship $H(S)^2 = K_O(S)K_I(S) \leq K(S)^2$. On the other hand, the relationships $K_O(S) \geq 1$ and $K_I(S) \geq 1$ imply respectively that $H(S)^2 \geq K_I(S)$ and $H(S)^2 \geq K_O(S)$. We conclude that $H(S)^2 \geq K(S)$. \square

2.4. Metric derivatives of Lipschitz mappings. Throughout this section, we let Ω denote a domain in \mathbb{R}^2 and (X, d) denote a metric space. We refer to Section 2.3 for basic terminology about seminorms.

Definition 2.2. Let $f: (\Omega, \|\cdot\|_2) \rightarrow (X, d)$ be a Lipschitz map. For all $x \in \Omega$ and $v \in \mathbb{R}^2$, the *metric derivative of f at x in the direction v* is

$$(9) \quad N_{f,x}(v) = \limsup_{t \rightarrow 0^+} \frac{d(f(x), f(x + tv))}{t}.$$

A result by Ivanov [Iva08] states the following. Similar results are proved in [Kir94, DCP90, DCP91, DCP95].

Theorem 2.3. Let $f: (\Omega, \|\cdot\|_2) \rightarrow (X, d)$ be a Lipschitz map. There exists a Borel set $N_0 \subset \Omega$ of zero Lebesgue measure such that, for all $x \in \Omega \setminus N_0$ and all $v \in \mathbb{R}^2$, the limit superior in (9) is an actual limit, and $v \mapsto N_{f,x}(v)$ is a seminorm for every $x \in \mathbb{R}^2 \setminus N_0$.

As a consequence of Theorem 2.3, the metric derivative of a Lipschitz map defines a seminorm field on Ω .

Proposition 2.4. Let $f: \Omega \rightarrow X$ be a Lipschitz function and N_f its metric derivative. The maximal stretching $x \mapsto L(N_f(x))$ is a minimal weak upper gradient of f , and f satisfies the change of variables formula

$$(10) \quad \int_{\Omega} \rho(z) J_2(N_{f,z}) d\mathcal{L}^2(z) = \int_X \int_{f^{-1}(x)} \rho(y) d\mathcal{H}^0(y) d\mathcal{H}_d^2(x)$$

for all Borel functions $\rho: \Omega \rightarrow [0, \infty]$.

Proof. Theorem 2.3 implies that the metric derivative, as defined in Definition 2.2, coincides with the metric derivative of Kirchheim [Kir94] \mathcal{L}^2 -almost everywhere in Ω . Kirchheim proves the change of variables formula (10) as Corollary 8 in [Kir94]. That $L(N_f)$ is a minimal weak upper gradient of f is proved in Section 4 of [LW17]. \square

The metric differential can be used to compute the metric speed of an absolutely continuous path.

Lemma 2.5. If $\gamma: [a, b] \rightarrow \Omega$ is an absolutely continuous path, then for almost every $t \in [a, b]$, the metric speed $v_{f \circ \gamma}(t)$ of $f \circ \gamma$ exists and satisfies

$$v_{f \circ \gamma}(t) = N_f \circ D\gamma(t),$$

where $D\gamma(t)$ is the derivative of γ at t .

Proof. Ivanov proves in Proposition 2.7 of [Iva08] that $\ell_d(f \circ \gamma) = \ell_{N_f}(\gamma)$ for every Lipschitz path $\gamma: [a, b] \rightarrow \mathbb{R}^2$. Since every absolutely continuous path has a Lipschitz parametrization, the same result holds for absolutely continuous paths $\gamma: [a, b] \rightarrow \mathbb{R}^2$. The lemma now follows from the Lebesgue differentiation theorem. \square

2.5. Quasiconformal mappings. Recall the geometric definition of quasiconformal mapping given in (1). A result of Williams is that this geometric definition is equivalent to an analytic definition based on metric Sobolev spaces. We state the two-dimensional case of this result, or rather a generalization to the case of continuous monotone maps.

Theorem 2.6 (cf. [Wil12]). Let X and Y be metric surfaces with locally finite Hausdorff 2-measure. Let $f: X \rightarrow Y$ be continuous and monotone and suppose that the pullback measure $f^*\mathcal{H}_Y^2$ is locally finite. The following are equivalent for the same constant $K \geq 1$:

- (i) $\text{mod } \Gamma \leq K \text{ mod } f\Gamma$ for all path families Γ in X .
- (ii) $f \in N_{\text{loc}}^{1,2}(X, Y)$ and satisfies

$$g_f^2(x) \leq K J_f(x)$$

for \mathcal{H}_X^2 -almost every $x \in X$.

Theorem 2.6 can be established using the original proof in [Wil12] with slight modifications which deal with the multiplicity of f . This is omitted here. A similar result can be found as Proposition 3.5 of [LW20].

The *outer dilatation* of f is the smallest constant $K \geq 1$ for which the modulus inequality $\text{mod } \Gamma \leq K \text{ mod } f\Gamma$ holds for all Γ in X . The *inner dilatation* of f is the smallest constant $K \geq 1$ for which $\text{mod } f\Gamma \leq K \text{ mod } \Gamma$ holds for all Γ in X . These are denoted respectively by $K_O(f)$ and $K_I(f)$. Thus a quasiconformal map is a homeomorphism with finite outer and inner dilatation. The *pointwise distortion* of f at $x \in X$ is

$$(11) \quad H_f(x) = g_f(x)g_{f^{-1}}(f(x)).$$

We interpret (11) as $H_f(x) = 1$ whenever $g_f(x) = 0$ or $g_{f^{-1}}(f(x)) = 0$. A consequence of Proposition 3.4 and Corollary 3.8 in [Iko19] is that $H_f(x)$ is independent of the representatives of g_f and $g_{f^{-1}}$ \mathcal{H}_X^2 -almost everywhere. Moreover, Corollary 3.12 of [Iko19] implies that $H_f(x) \leq \sqrt{K_O(f)K_I(f)}$ for \mathcal{H}_X^2 -almost every $x \in X$. The smallest constant $H \geq 1$ for which $H_f(x) \leq H$ for \mathcal{H}_X^2 -almost every $x \in X$ is called the *distortion* of f and denoted by $H(f)$.

Consider now a quasiconformal map $f: \Omega \subset \mathbb{R}^2 \rightarrow X$ that is also Lipschitz. Then the equalities $g_f(x) = L(N_{f,x})$ and $g_{f^{-1}} \circ f(x) = (\omega(N_{f,x}))^{-1}$ hold for \mathcal{L}^2 -almost every $x \in \Omega$ [Iko19, Proposition 4.8]. Consequently, we have the equality $H_f(x) = H(N_{f,x})$ for \mathcal{L}^2 -almost every $x \in \Omega$.

In general, a quasiconformal map $f: \Omega \subset \mathbb{R}^2 \rightarrow X$ must satisfy *Lusin's Condition* (N^{-1}): for every Borel set $E \subset \Omega$ of positive Lebesgue measure, $f(E)$ has positive Hausdorff 2-measure. This is essentially proved in Remark 8.3 or Section 17 of [Raj17]. On the other hand, f need not satisfy *Lusin's Condition* (N): for every Borel set $E \subset \Omega$ of zero Lebesgue measure, $f(E)$ has zero Hausdorff 2-measure. An example of this is given as Proposition 17.1 of [Raj17]. Generalizations of these results are considered in Section 3 of [Iko19].

A uniformization theorem for quasiconformal mappings was proved by Rajala based on the notion of *reciprocity* [Raj17]. Let X be a metric surface. For a set $G \subset X$ and disjoint sets $F_1, F_2 \subset G$, let $\Gamma(F_1, F_2; G)$ denote the family of paths whose images are contained in G that start from F_1 and end in F_2 . A *quadrilateral* is a set Q homeomorphic to $[0, 1]^2$ with boundary consisting of four nonoverlapping boundary arcs, labelled $\xi_1, \xi_2, \xi_3, \xi_4$ in cyclic order.

Definition 2.7. A metric surface X is *reciprocal* if there exists a constant $\kappa \geq 1$ such that

$$(12) \quad \text{mod } \Gamma(\xi_1, \xi_3; Q) \text{ mod } \Gamma(\xi_2, \xi_4; Q) \leq \kappa$$

for every quadrilateral $Q \subset X$, and

$$(13) \quad \lim_{r \rightarrow 0^+} \text{mod } \Gamma(\overline{B}(x, r), X \setminus B(x, R); \overline{B}(x, R)) = 0$$

for all $x \in X$ and $R > 0$ such that $X \setminus B(x, R) \neq \emptyset$.

Note that, for all metric surfaces, the product in (12) is bounded from below by a universal constant $\tilde{\kappa} > 0$ [RR19]. We say that a reciprocal surface is κ -reciprocal if (12) and the corresponding lower bound hold for the constant κ .

Theorem 1.4 in [Raj17] states that a metric surface X homeomorphic to \mathbb{R}^2 is reciprocal if and only if there exists a quasiconformal homeomorphism onto the disk or the Euclidean plane. This result is extended to arbitrary metric surfaces in [Iko19]. More precisely, Theorem 1.3 in [Iko19] states that a metric surface X is locally reciprocal (that is, every point in X has a neighborhood that is reciprocal) if and only if X is quasiconformally equivalent to a smooth Riemannian 2-manifold. In particular, a metric surface that is locally reciprocal is also globally reciprocal.

2.6. Removable sets for conformal mappings. We collect some background on removable sets for conformal mappings. Recall from the introduction that the compact set $E \subset \mathbb{R}^2$ is *removable for conformal mappings* if every conformal embedding $f: \mathbb{R}^2 \setminus E \rightarrow \widehat{\mathbb{R}}^2$ extends to a conformal mapping $F: \widehat{\mathbb{R}}^2 \rightarrow \widehat{\mathbb{R}}^2$. Thus f is the restriction of a Möbius transformation.

This notion exists under several names, including *sets of absolute area zero* and *negligible sets for extremal distance*. This nomenclature reflects the following characterization.

Proposition 2.8. Let $E \subset \mathbb{R}^2$ be compact. The following are equivalent.

- (i) E is removable for conformal mappings.
- (ii) E has absolute area zero: for every conformal embedding $f: \mathbb{R}^2 \setminus E \rightarrow \widehat{\mathbb{R}}^2$, the complementary set $\widehat{\mathbb{R}}^2 \setminus f(\mathbb{R}^2 \setminus E)$ has Lebesgue measure zero.
- (iii) E is negligible for modulus: for every domain $\Omega \subset \mathbb{R}^2$ and pair of disjoint compact sets $F, G \subset \Omega \setminus E$, $\text{mod } \Gamma(F, G; \Omega) = \text{mod } \Gamma(F, G; \Omega \setminus E)$.
- (iv) Any quasiconformal embedding $f: \mathbb{R}^2 \setminus E \rightarrow \widehat{\mathbb{R}}^2$ has an extension to a quasiconformal mapping $F: \widehat{\mathbb{R}}^2 \rightarrow \widehat{\mathbb{R}}^2$.
- (v) For any open set $U \subset \mathbb{R}^2$, every quasiconformal mapping on $U \setminus E$ extends quasiconformally to the whole open set U .

The equivalence of (i), (ii) and (iii) is proved in [AB50]. The equivalence of (i) and (iv) is a consequence of the measurable Riemann mapping theorem. See Proposition 4.7 in [You15] for a proof. The equivalence of (i) and (v) can also be found in [You15] as Proposition 4.6. We see from (iv) and (v) that removability for conformal mappings is a local property and a quasiconformal invariant. If E contains a nontrivial connected component E_0 , then there is a non-Möbius conformal map $f: \mathbb{R}^2 \setminus E_0 \rightarrow \mathbb{R}^2$ such that $\mathbb{R}^2 \setminus f(\mathbb{R}^2 \setminus E_0)$ is the closed unit disk. Thus Property (ii) implies that a removable set for conformal mappings is totally disconnected.

Property (iii) in Proposition 2.8 indicates the connection between quasiconformal uniformization and removable sets. Observe that for each triple F, G , and Ω , $\Gamma(F, G; \Omega \setminus E)$ is a subset of $\Gamma(F, G; \Omega)$ and thus satisfies $\text{mod } \Gamma(F, G; \Omega \setminus E) \leq \text{mod } \Gamma(F, G; \Omega)$. In contrast, the metric space constructions in our paper collapse a domain at the set E and hence *increase* the modulus of a path family, up to a factor related to the dilatation bound of the norm field. Thus Theorem 1.3 and Theorem 1.4 can be summarized roughly by saying that *removing the set E does not decrease the modulus of any path family if and only if collapsing the plane at E does not increase the modulus of any path family*.

3. CONSTRUCTING A METRIC FROM A NORM FIELD

In this section, we give a description of the metric spaces considered in this paper and develop their basic properties. These spaces are constructed from measurable Finsler structures satisfying additional assumptions. The precise definition is given in Section 3.1.

There is a vast literature on Riemannian and Finsler geometry, typically requiring smoothness or at least continuity of the metrics. The idea of constructing metrics from Finsler structures with less regularity has been considered by various previous authors, and so the material in this section is more-or-less standard. In Section 3.2, we include a brief comparison with the existing literature.

We consider here seminorm fields N such that either N_x is a norm or $N_x = 0$ for all $x \in \Omega$. Recall from the introduction that, slightly abusing terminology, we use the term *norm field* to refer to an object of this type. Since a vector $v \in \mathbb{R}^2$ often comes with an implicit basepoint x , we will sometimes write $N(v)$ in place of $N_x(v)$, such as in the expression $N \circ D\gamma$.

3.1. Definition of the metric. Let $\Omega \subset \mathbb{R}^2$ be a domain.

Definition 3.1. A norm field $N: \Omega \times \mathbb{R}^2 \rightarrow [0, \infty)$ is *admissible* if it satisfies the following:

- (i) (lower semicontinuous) For all vectors $v \in \mathbb{R}^2$ and points $x \in \Omega$, we have $N_x(v) \leq \liminf_{y \rightarrow x} N_y(v)$.
- (ii) (locally bounded) For all $x \in \Omega$, there is a neighborhood U of x and $M > 0$ such that $L(N_y) \leq M$ for all $y \in U$.
- (iii) (locally bounded distortion) For all $x \in \Omega$, there is a neighborhood U of x and $H > 0$ such that $L(N_y) \leq H\omega(N_y)$ for all $y \in U$.
- (iv) (nonseparating) The set $E = \{x \in \Omega : N_x = 0\}$ is compact and $\Omega \setminus E$ is connected.

An immediate consequence of having locally bounded distortion is that $N_x(v) = 0$ for some $v \in \mathbb{R}^2 \setminus \{0\}$ if and only if N_x is identically zero.

We use the norm field N to measure the length of an absolutely continuous path $\gamma: [a, b] \rightarrow \Omega$ in the following way. We define the N -length of γ to be

$$\ell_N(\gamma) = \int_a^b N \circ D\gamma(t) dt,$$

where $D\gamma: [a, b] \rightarrow \Omega \times \mathbb{R}^2$ is a Borel representative of the differential of γ .

Definition 3.2. Let N be an admissible seminorm field and $x, y \in \Omega$. The N -distance between x and y is defined as

$$d_N(x, y) = \inf \ell_N(\gamma),$$

where the infimum is taken over absolutely continuous paths γ joining x to y in Ω .

The function d_N is locally finite and satisfies the triangle inequality, but it may happen that $d_N(x, y) = 0$ for distinct points $x, y \in \Omega$. Thus, in general, d_N is only a pseudodistance. Let \mathcal{E}_N be the partition of Ω into equivalence classes of points, where $x, y \in \Omega$ belong to the same equivalence class if $d_N(x, y) = 0$. This yields the quotient space Ω/\mathcal{E}_N and the natural quotient map $\pi_N: \Omega \rightarrow \Omega/\mathcal{E}_N$. The space Ω/\mathcal{E}_N comes equipped with the metric that is the pushforward of d_N under π_N , which we denote by \tilde{d}_N .

A consequence of the local boundedness of N is that the quotient map π_N is locally Lipschitz. In particular, the results described in Section 2.4 apply to the map π_N .

3.2. Remarks on definition of admissible norm fields. We offer a few remarks about Definition 3.1 and give a comparison to the previous literature.

The lower semicontinuity assumption guarantees that the metric tangents of Ω/\mathcal{E}_N coincide with N almost everywhere. This implies, for example, that two conformally equivalent norm fields generate metrics that are 1-quasiconformally equivalent. In general, the metric tangents are not so well-behaved. For example, let $F \subset [0, 1]$ be a Cantor set of positive linear measure, and let $E = F \times F \subset \mathbb{R}^2$. The norm field N defined by

$$N_x = \begin{cases} 2\|\cdot\|_\infty & \text{if } x \in E \\ \|\cdot\|_1 & \text{if } x \notin E \end{cases}$$

generates the same metric as the norm field $\|\cdot\|_1$, despite the fact that they differ on a positive measure set. Indeed, the inequality $\|x - y\|_1 \leq d_N(x, y)$ is immediate for all $x, y \in \mathbb{R}^2$, since $\|\cdot\|_1 \leq 2\|\cdot\|_\infty$. On the other hand, for all $x, y \in \mathbb{R}^2 \setminus E$, there is an ℓ_1 -geodesic from x to y lying in $\mathbb{R}^2 \setminus E$. Thus $d_N(x, y) \leq \|x - y\|_1$ for such x, y . Since E has empty interior, we obtain the inequality $d_N(x, y) \leq \|x - y\|_1$ for all $x, y \in \mathbb{R}^2$. The lower semicontinuity assumption allows us to avoid this type of behaviour; see Lemma 3.6 below.

The fact that $E = \{x \in \Omega : N_x = 0\}$ is non-separating guarantees that the quotient space is homeomorphic to Ω (Corollary 3.9). For example, if E is the Euclidean unit circle and $N = \chi_{\mathbb{R}^2 \setminus E} \|\cdot\|_2$, the resulting quotient space is not a 2-manifold.

Now we discuss some of the related literature on non-smooth Finsler metrics. Perhaps the first investigations into this topic were carried out by Busemann–Mayer in [BM41]. Beginning in the 1940s, the Russian school led by Alexandrov developed a theory of *surfaces of bounded curvature*, also now known as *Alexandrov surfaces*, as a generalization of two-dimensional Riemannian geometry. See [AZ67] and [Res93] for an overview.

Finsler metrics on Lipschitz manifolds were systematically studied by De Cecco–Palmieri in the series of papers [DCP88, DCP90, DCP91, DCP95]. Note that they take a different approach to defining the distance d_N from a norm field N . The idea is to make the distance more robust by making the definition insensitive to changes in N on a set of measure zero. In particular, the norm field N need only be defined on a full measure subset. This is achieved as follows. For a set $F \subset \mathbb{R}^2$ of measure zero, let Γ_F be the family of absolutely continuous paths that intersect F in a set of length zero. Then one defines the metric $d_{N,F}$ as in Definition 3.2 but restricting to paths in Γ_F . Next one defines $D_N(x, y) = \sup d_{N,F}(x, y)$, the supremum taken over all measure zero sets F . This is called the *intrinsic distance* in [DCP95, GPP06] and *essential metric* in [AHPCS18] and further investigated in [CS19]. Observe that if N is continuous, then the essential metric coincides with the metric considered in this paper. However, we do not take this approach, since the norm fields we have in mind typically vanish on a set of measure zero, and we prefer the additional flexibility of only requiring N to be lower semicontinuous.

3.3. Properties of length. In the remainder of this section, we establish properties of admissible norm fields and their corresponding metric. Our first lemma states that the property of lower semicontinuity of N in each direction v can be promoted to lower semicontinuity at a point in all directions uniformly.

Lemma 3.3. Let N be an admissible norm field and $x \in \Omega$. For every $\varepsilon > 0$, there exists $r > 0$ such that

$$N_y(v) \geq (1 - \varepsilon)N_x(v).$$

for all $y \in B(x, r)$ and $v \in \mathbb{R}^2$.

Proof. If N_x is the zero seminorm, then the conclusion follows immediately. Thus we may assume that N_x is a norm. By the positive homogeneity of N , we need only consider vectors $v \in \mathbb{S}^1$. Let $\varepsilon > 0$ and let $\delta = \varepsilon\omega(N_x)$, so that $N_x(v) - \delta \geq (1 - \varepsilon)N_x(v)$ for all $v \in \mathbb{R}^2$. Thus it suffices to show that there exists a radius $r > 0$ such that

$$N_y(v) \geq N_x(v) - \delta$$

for all $y \in B(x, r)$ and $v \in \mathbb{S}^1$.

Assume to the contrary that no such r exists. Then there exist sequences $(y_n) \subset \Omega$ and $(v_n) \subset \mathbb{S}^1$ for which

$$(14) \quad N_{y_n}(v_n) < N_x(v_n) - \delta$$

for all $n \in \mathbb{N}$. By passing to a subsequence, we have that v_n converges to some vector $v \in \mathbb{S}^1$.

Let $M > 0$ be such that $L(N_y) \leq M$ for all y in a neighborhood of x . Then for every sufficiently large $n \in \mathbb{N}$,

$$N_x(v_n) - M \|v - v_n\|_2 \leq N_x(v)$$

and

$$N_{y_n}(v) \leq N_{y_n}(v_n) + M \|v - v_n\|_2.$$

Moreover, the lower semicontinuity of N implies that

$$N_x(v) - \frac{\delta}{2} \leq N_{y_n}(v)$$

for all sufficiently large $n \in \mathbb{N}$. Combining these inequalities yields

$$N_x(v_n) - \left(2M \|v - v_n\|_2 + \frac{\delta}{2}\right) \leq N_{y_n}(v_n).$$

Let n be sufficiently large so that $\|v - v_n\|_2 < \delta(4M)^{-1}$. Then the preceding inequality contradicts (14), and the result follows. \square

The next lemma shows that the metric d_N is locally well-behaved outside of the set E .

Lemma 3.4. Let N be an admissible norm field. For all $x \in \Omega \setminus E$, there exists $r > 0$ such that $\overline{B}(x, r) \subset \Omega \setminus E$ and the quotient map π_N is bi-Lipschitz in the neighborhood $B(x, r)$.

Proof. We let $\omega(x) = \omega(N_x)$ denote the minimal stretching of N . Lemma 3.3 implies that ω is lower semicontinuous. Also $\omega(x) = 0$ if and only if N_x is a seminorm.

Let $x \in \Omega \setminus E$. Let $R > 0$ be such that the closed ball $\overline{B}(x, R)$ is contained in $\Omega \setminus E$ and satisfies $\omega(z) \geq \omega(x)/2$ for all $z \in B(x, R)$. Such an $R > 0$ exists by the lower semicontinuity of the map $z \mapsto \omega(z)$. Moreover, the local boundedness of N implies that the maximal stretching $L(N_z)$ is bounded from above by M for all $z \in B(x, R)$. We conclude that

$$\frac{\omega(x)}{2} \|v\|_2 \leq N_z(v) \leq M \|v\|_2$$

for all $z \in B(x, R)$ and all $v \in \mathbb{R}^2$.

Let $r = R/2$. We claim that

$$\frac{\omega(x) \|y - z\|_2}{4} \leq d_N(y, z) \leq M \|y - z\|_2$$

for all $y, z \in B(x, r)$. Clearly, the line segment from y to z has N -length at most $M \|y - z\|_2$. For the lower bound, consider an arbitrary absolutely continuous

path γ from y to z . If $|\gamma| \subset B(x, R)$, then we have the lower bound $\ell_N(\gamma) \geq \omega(x) \|y - z\|_2 / 2$. If $|\gamma|$ is not contained in $B(x, R)$, then its length is at least

$$\omega(x)(R - r) = \frac{\omega(x)R}{2} \geq \frac{\omega(x) \|y - z\|_2}{4}.$$

Since our path is arbitrary, we obtain $d_N(y, z) \geq \omega(x) \|y - z\|_2 / 4$. We conclude that d_N is bi-Lipschitz equivalent to the Euclidean distance on $B(x, r)$. \square

Lemma 3.5. For \mathcal{L}^2 -almost every $x \in \Omega$, the metric derivative N_{π_N} of π_N at x satisfies

$$N_{\pi_N, x} \leq N_x.$$

Moreover, for every $x \in \Omega$,

$$N_x \leq N_{\pi_N, x}.$$

In particular, the metric derivative N_{π_N} equals N \mathcal{L}^2 -almost everywhere in $x \in \Omega$.

Proof. First, we show that the upper bound $N_{\pi_N, x} \leq N_x$ holds \mathcal{L}^2 -almost everywhere in Ω . Consider a fixed $v \in \mathbb{R}^2 \setminus \{0\}$. The local boundedness of N implies that the function $x \mapsto N_x(v)$ is locally integrable. Consider a rectangle $R \subset \Omega$ with one side parallel to v . There is a family of parallel line segments $\gamma_t: [0, h_0] \rightarrow R$, $\gamma_t(s) = x_t + vs$, that foliate R . Observe that for all t and s , $D\gamma_t(s) = v$. The definition of d_N implies that

$$N_{\pi_N, \gamma_t(s)}(v) \leq \limsup_{h \rightarrow 0^+} \frac{1}{h} \int_{[s, s+h]} N_{\gamma_t(a)}(v) d\mathcal{L}^1(a).$$

According to Lebesgue's differentiation theorem, the lim sup on the right-hand side equals $N_{\gamma_t(s)}(v)$ for \mathcal{L}^1 -almost every $s \in [0, h_0]$. Fubini's theorem implies that

$$N_{\pi_N, x}(v) \leq N_x(v)$$

holds \mathcal{L}^2 -almost everywhere in R . Since R is arbitrary, the same conclusion holds for almost every point in Ω . The first inequality follows.

Next, we show that the inequality $N_x \leq N_{\pi_N, x}$ holds for all $x \in \Omega$. In the case that $x \in E$, the conclusion is immediate since then $N_x = 0$. We consider now the case that $x \in \Omega \setminus E$. Let $v \in \mathbb{R}^2 \setminus \{0\}$ and let $\varepsilon > 0$.

Let $r > 0$ be such that the conclusions of Lemma 3.3 and Lemma 3.4 hold for the point x and the given value of ε . In particular, Lemma 3.4 implies that there exists $\alpha > 0$ such that

$$\alpha^{-1} d_N(y, z) \leq \|y - z\|_2 \leq \alpha d_N(y, z)$$

for all $y, z \in B(x, r)$. Moreover, the local boundedness of N implies that there exists $M > 0$ such that the maximal stretching $L(N_y) \leq M$ for all $y \in B(x, r)$. Let

$$t_0 = \frac{1}{\|v\|_2} \frac{r}{2\alpha} \min \left\{ \frac{1}{\alpha}, \frac{1}{\varepsilon M} \right\}.$$

For all $t \in (0, t_0)$, consider an absolutely continuous path $\gamma_t: [0, 1] \rightarrow \Omega$ joining x to $x + tv$ that satisfies

$$(15) \quad \int_0^1 N \circ D\gamma_t d\mathcal{L}^1 \leq d_N(x, x + tv) + \varepsilon t N_x(v).$$

The right-hand side of (15) is bounded above by $\alpha t \|v\|_2 + \varepsilon M t \|v\|_2 < r/\alpha$. In particular, this implies that

$$(16) \quad |\gamma_t| \subset B_{\|\cdot\|}(x, r).$$

Next, observe that

$$(17) \quad t N_x(v) = N_x(tv) \leq M \|tv\|_2 \leq \alpha M d_N(x, x + tv).$$

Applying now the conclusion of Lemma 3.3 along γ_t , which is allowed due to (16), we have

$$(18) \quad (1 - \varepsilon)N_x(D\gamma_t(s)) \leq N \circ D\gamma_t(s),$$

for almost every $s \in [0, 1]$. Note that the norm field N on the left-hand side has a fixed basepoint.

Since straight line segments are geodesics with respect to the norm N_x , by integrating both sides of (18) and applying (15) and (17), we obtain

$$(19) \quad (1 - \varepsilon)N_x(tv) \leq (1 + \varepsilon\alpha M)d_N(x, x + tv).$$

We divide both sides of (19) by t and let $t \rightarrow 0$. We have

$$(1 - \varepsilon)N_x(v) \leq (1 + \varepsilon\alpha M) \liminf_{t \rightarrow 0} \frac{d_N(x, x + tv)}{t}.$$

The lim inf on the right-hand side is bounded from above by the metric derivative $N_{\pi_N, x}(v)$. The result follows by letting $\varepsilon \rightarrow 0$. \square

Lemma 3.6. For every Borel function $\rho: \Omega/\mathcal{E}_N \rightarrow [0, \infty]$, we have the change of variables formula

$$\int_{\Omega} \rho \circ \pi_N J_2(N) d\mathcal{L}^2 = \int_{\Omega/\mathcal{E}_N} \rho d\mathcal{H}_{d_N}^2.$$

Proof. We proved in Lemma 3.5 the fact that the metric derivative of π_N equals N \mathcal{L}^2 -almost everywhere. Then the change of variables formula Proposition 2.4 implies that $\mathcal{H}_{d_N}^2(\pi_N(E)) = 0$. The fact that π_N is injective in the complement of E implies that the multiplicity term from Proposition 2.4 can be omitted. \square

Lemma 3.7. For every absolutely continuous path γ in Ω , $\ell_N(\gamma) = \ell_{d_N}(\pi_N \circ \gamma)$. In particular, the equality

$$(20) \quad v_{\pi_N \circ \gamma} = N \circ D\gamma$$

holds almost everywhere in the domain of γ .

Proof. An immediate consequence of the definitions is that $\ell_{\tilde{d}_N}(\pi_N \circ \gamma) \leq \ell_N(\gamma)$ for every absolutely continuous γ in Ω . For the other direction, let L denote the N_{π_N} -length of γ :

$$L = \int_I N_{\pi_N} \circ D\gamma(t) d\mathcal{L}^1(t).$$

Since $N(x) \leq N_{\pi_N}(x)$ for all $x \in \Omega$ by Lemma 3.5, we see that $\ell_N(\gamma) \leq L$. By Lemma 2.5, the equality $L = \ell_{\tilde{d}_N}(\pi_N \circ \gamma)$ holds for all absolutely continuous γ . The equality $\ell_N(\gamma) = \ell_{\tilde{d}_N}(\pi_N \circ \gamma)$ now follows. The metric speed identity (20) follows from the Lebesgue differentiation theorem. \square

As a consequence of the previous lemma, whenever $\gamma: I \rightarrow \Omega$ is an absolutely continuous path, we have the integral formula

$$\int_{\pi_N \circ \gamma} \rho ds_N = \int_I (\rho \circ \pi_N)(N \circ D\gamma) d\mathcal{L}^1$$

for all Borel measurable functions $\rho: \Omega/\mathcal{E}_N \rightarrow [0, \infty]$.

3.4. The quotient map.

Proposition 3.8. The quotient map $\pi_N: \Omega \rightarrow \Omega/\mathcal{E}_N$ is locally Lipschitz, locally bi-Lipschitz in the complement of E , and its restriction to $\Omega \setminus E$ is injective.

Moreover, the map π_N is closed and, for all $x \in \pi_N(E)$, the preimage $\pi_N^{-1}(x)$ is a connected and compact subset of E .

Proof. We already proved in Lemma 3.4 that π_N is locally bi-Lipschitz outside of E . Moreover, since N is locally bounded, π_N is locally Lipschitz at all points in Ω .

Next, let $x \in \Omega \setminus E$ and $U \subset \Omega \setminus E$ be a neighborhood of x such that $\pi_N|_U$ is bi-Lipschitz. The bi-Lipschitz property implies that $d_N(x, y) > 0$ for all $y \in U$. Next, let $r > 0$ be small enough so that $\overline{B}_{\|\cdot\|}(x, r) \subset U$, and let $c = \inf\{d_N(x, y) : y \in S_{\|\cdot\|}(x, r)\} > 0$. If $y \in \Omega \setminus U$, then any path from x to y must intersect $S_{\|\cdot\|}(x, r)$, which gives $d_N(x, y) \geq c > 0$. We conclude that π_N is injective in the complement of E .

Next, we prove that $\pi_N^{-1}(\tilde{x})$ is a connected compact subset of E for all $\tilde{x} \in \pi_N(E)$. Let $x \in \pi_N^{-1}(\tilde{x})$ and let K be the component of E containing x .

Let γ be a closed Jordan path in $\Omega \setminus E$ that separates K and the boundary of $\partial\Omega$. See [Why64, Section III.3] for the existence of such a path γ . Let U be the complementary component of $|\gamma|$ containing K .

Let $c = \inf\{d_N(x, z) : z \in \Omega \setminus \overline{U}\}$. The image $|\gamma|$ has a small neighborhood V compactly contained in $\Omega \setminus E$. Every path joining the point x to $\Omega \setminus \overline{U}$ must pass through \overline{V} . The lower semicontinuity of N implies that $N \geq \alpha \|\cdot\|_2$ in \overline{V} for some $\alpha > 0$ and hence that $c > 0$.

Let $y \in \pi_N^{-1}(\tilde{x})$. Let (γ_n) be a sequence of Lipschitz paths joining x to y satisfying

$$\ell_N(\gamma_n) \leq 2^{-n}c$$

for all $n \in \mathbb{N}$. Observe that the image of each path γ_n is contained in \overline{U} . Moreover, for every $z_n \in |\gamma_n|$, we have that $d_N(x, z_n) \leq 2^{-n}c$. This implies that a subsequence of the sets $(|\gamma_n|)$ converges with respect to the Hausdorff distance to a connected subset of $\pi_N^{-1}(\tilde{x}) \cap \overline{U}$. This is a consequence of general properties of Hausdorff convergence in metric spaces; see Proposition 4.4.14 and Theorems 4.4.15 and 4.4.17 in [AT04]. Since y is arbitrary, we conclude that $\pi_N^{-1}(\tilde{x})$ is connected.

Next, we check that $\pi_N^{-1}(\tilde{x}) \subset K$. Observe that, for a given point $z \in \Omega \setminus K$, the Jordan path γ above can be chosen so that K and z are contained in different complementary components. An argument similar to that above shows that $d_N(K, z) > 0$.

The final step is to show that π_N is closed. Let $F \subset \Omega$ be a closed set and let \tilde{x} be a limit point of $\pi_N(F)$. Since $\pi_N^{-1}(\tilde{x})$ is a singleton or contained in a component of E , there is a Jordan domain $U \subset \Omega$ such that ∂U is contained in $\Omega \setminus E$ and separates $\pi_N^{-1}(\tilde{x})$ and $\partial\Omega$. Arguing as in the first part of the proof, we deduce that there is a constant $c > 0$ such that $d_N(\pi_N^{-1}(\tilde{x}), z) \geq c$ for all $z \in \Omega \setminus U$. This implies that \tilde{x} is a limit point of $\pi_N(F \cap \overline{U})$. Let (\tilde{x}_j) be a sequence in $\pi_N(F \cap \overline{U})$ with limit \tilde{x} . Let (y_j) be a sequence in $F \cap \overline{U}$ such that $\pi_N(y_j) = \tilde{x}_j$. The compactness of \overline{U} implies that there is a subsequence (y_{j_k}) that converges to a point $y \in F$. Since $\pi_N|_{\overline{U}}$ is Lipschitz, it follows that (x_{j_k}) converges to $\pi_N(y)$, and moreover that $\tilde{x} = \pi_N(y)$. We conclude that $\tilde{x} \in \pi_N(F)$, and hence that $\pi_N(F)$ is closed. \square

Corollary 3.9. The space Ω/\mathcal{E}_N is homeomorphic to Ω .

Proof. By Proposition 3.8, π_N is a closed and monotone map. Thus each element of the decomposition \mathcal{E}_N is a planar continuum. Since the components of E are non-separating, so are the elements of \mathcal{E}_N . It follows now from the classical theorem of Moore that Ω/\mathcal{E}_N is homeomorphic to Ω . See, for instance, Theorem 25.1 in [Dav86]. \square

Next we study the analytic properties of π_N . A consequence of Proposition 2.4 and Lemma 3.5 is that $x \mapsto L(N_x)$ is a minimal weak upper gradient of π_N . The following lemma identifies the minimal weak upper gradient of the inverse of π_N .

Lemma 3.10. *If $U \subset \Omega$ is an open set such that $\pi_N|_U$ is injective and its inverse h is an element of $N_{\text{loc}}^{1,2}(\pi_N(U), \mathbb{R}^2)$, the function*

$$g = \left(\frac{1}{\omega(N)} \chi_{U \setminus E} \right) \circ h$$

is a minimal weak upper gradient of h .

We use the convention $\frac{1}{0} \cdot 0 = 0$ in Lemma 3.10.

Proof. We show that the function g as in the claim is a weak upper gradient of h . First, the change of variables formula Lemma 3.6 implies that $\mathcal{H}_{d_N}^2(\pi_N(E)) = 0$.

Therefore the paths that have positive \tilde{d}_N -length on $\pi_N(E)$ have zero modulus. Moreover, since h is an element of $N_{\text{loc}}^{1,2}(\pi_N(U), \mathbb{R}^2)$, h maps almost every absolutely continuous path in $\pi_N(U)$ to an absolutely continuous path in U . Thus it suffices to check the upper gradient inequality for a path $\tilde{\gamma}: [0, 1] \rightarrow \pi_N(U)$ that intersects $\pi_N(E)$ in a set of \tilde{d}_N -length zero and along which h is absolutely continuous.

Let $\tilde{\gamma}: [0, 1] \rightarrow \pi_N(U)$ be such a path, and let $x = \tilde{\gamma}(0)$ and $y = \tilde{\gamma}(1)$. Let $\gamma = h \circ \tilde{\gamma}$. Note that the absolute continuity of h along $\tilde{\gamma}$ implies that γ intersects E in a set of Euclidean length zero. Therefore, by reparametrizing, we can assume that the set $J = \gamma^{-1}(\Omega \setminus E)$ has full length in $[0, 1]$.

By Lemma 2.5, the metric speed identity $v_{\tilde{\gamma}} = N \circ D\gamma$ holds \mathcal{L}^1 -almost everywhere for γ . Also for almost every $t \in [0, 1] \setminus \gamma^{-1}(E)$,

$$(21) \quad v_{\gamma}(t) = \|D\gamma(t)\|_2 \leq \frac{1}{\omega(N_{\gamma}(t))} N \circ D\gamma(t) = \frac{1}{\omega(N_{\gamma}(t))} v_{\tilde{\gamma}}(t),$$

where $\omega(N_{\gamma}(t))$ is the minimal stretching of N at $\gamma(t)$. Since $\gamma^{-1}(E)$ has zero measure, we conclude from (21) that for almost every $t \in [0, 1]$,

$$(22) \quad v_{\gamma}(t) \leq \left(\frac{1}{\omega(N)} \chi_{\Omega \setminus E} \right) \circ \gamma(t) v_{\tilde{\gamma}}(t).$$

The right-hand side in (22) equals $g \circ (\pi_N \circ \gamma(t)) v_{\tilde{\gamma}}(t)$. Therefore, integrating both sides of (22) implies that

$$\|h(x) - h(y)\|_2 \leq \int_{\tilde{\gamma}} g ds_{\tilde{d}_N}.$$

The local L^2 -integrability of g follows from the fact that N has locally bounded distortion (Lemma 2.1) and the change of variables formula (Lemma 3.6).

We are left to check that g is a minimal weak upper gradient. Let $\rho \in L_{\text{loc}}^2(\pi_N(U))$ be a weak upper gradient of h . We want to show that $g(x) \leq \rho(x)$ for $\mathcal{H}_{d_N}^2$ -almost every $x \in \pi_N(U)$. The set $\pi_N(E)$ is negligible, so it is sufficient to check this in the complement of $\pi_N(E)$. As h is locally bi-Lipschitz in the complement of $\pi_N(E)$, it suffices to check that

$$(23) \quad g \circ \pi_N(x) = \sup_{v \in \mathbb{S}^1} \frac{1}{N_x(v)} \leq \rho \circ \pi_N(x)$$

\mathcal{L}^2 -almost every $x \in U \setminus E$.

Consider a square $R \subset U \setminus E$ and the accompanying foliation given by

$$\gamma_t(s) = x_0 + sv + tw,$$

where $v, w \in \mathbb{R}^2$ are orthogonal vectors and $s, t \in [-1, 1]$.

The metric speed characterization $v_{\pi_N \circ \gamma} = N \circ D\gamma$ implies that

$$v_{\gamma_t}(s) = \|v\|_2 \leq \rho \circ (\pi_N \circ \gamma_t(s)) N_{\gamma_t(s)}(v)$$

holds almost everywhere along the domain of γ_t for almost every t . Fubini's theorem implies that

$$\|v\|_2 \leq \rho \circ \pi_N(x) N_x(v)$$

holds \mathcal{L}^2 -almost everywhere on R . Equivalently,

$$1 \leq \rho \circ \pi_N(x) N_x \left(\frac{v}{\|v\|_2} \right)$$

\mathcal{L}^2 -almost everywhere on R . We can cover $U \setminus E$ by squares whose sides are parallel to v and w , so we deduce that

$$(24) \quad \frac{1}{N_x \left(\frac{v}{\|v\|_2} \right)} \leq \rho \circ \pi_N(x)$$

\mathcal{L}^2 -almost everywhere on $U \setminus E$.

Let D be a countable dense subset of \mathbb{S}^1 . We have shown that, for \mathcal{L}^2 -almost every $x \in U \setminus E$, (24) holds for every $v \in D$. Consequently, (23) holds for \mathcal{L}^2 -almost every $x \in U \setminus E$. \square

3.5. Local quasiconformality. Let U be a subdomain of Ω such that $\overline{U} \subset \Omega$ is compact. Since the norm field N has locally bounded distortion, there exists $K(U) < \infty$ such that

$$L(N)^2 \leq K(U) J_2(N)$$

for the maximal stretching $L(N)$ and the Jacobian $J_2(N)$. Recall that $L(N)$ is a weak upper gradient of π_N , $J_2(N)$ is the Jacobian of π_N , and that the pullback measure $\pi_N^* \mathcal{H}_{d_N}^2$ is locally finite. Thus Theorem 2.6 implies the following.

Proposition 3.11. For every path family Γ in U , we have that

$$\text{mod } \Gamma \leq K(U) \text{mod } \pi_N \Gamma.$$

If $(\Omega/\mathcal{E}_N, \tilde{d}_N)$ is reciprocal, then it admits some quasiconformal parametrization from a domain in Euclidean space. We show here that the map π_N itself is a quasiconformal parametrization, at least locally.

Proposition 3.12. The metric surface $(\Omega/\mathcal{E}_N, \tilde{d}_N)$ is reciprocal if and only if π_N is a homeomorphism that is locally quasiconformal.

Here, a map $\psi: X \rightarrow Y$ is *locally quasiconformal* if every point $x \in X$ has a neighborhood U such that the restriction of ψ to U is K -quasiconformal for some $K \geq 1$, where the K is allowed to depend on x .

Proof. If π_N is a locally quasiconformal homeomorphism, every point in $Y = (\Omega/\mathcal{E}_N, \tilde{d}_N)$ has a neighborhood that is reciprocal. By Corollary 1.4 of [Iko19], this implies that Y is reciprocal.

Conversely, suppose that Y is reciprocal. It suffices to fix an arbitrary Jordan domain $Q \subset \Omega$ with $\partial Q \cap E = \emptyset$ and check that $\pi_N|_{\text{int } Q}$ is quasiconformal.

The reciprocity of Y implies the existence of a homeomorphism $f: \pi_N(Q) \rightarrow \overline{\mathbb{D}}$ that is $\frac{\pi}{2}$ -quasiconformal in $\pi_N(Q)$. Let $h = f \circ \pi_N|_{\text{int } Q}$. The claim follows if we can check that h is quasiconformal.

The map h satisfies the assumptions of Theorem 2.6 and the property (i). Therefore the quasiconformality of h follows from its injectivity; see Section 3.1 of [AIM09]. We check that h is injective.

Let $y \in \mathbb{D}$ and $C = h^{-1}(y)$. The set C is connected and compact. The modulus of paths joining y to the boundary \mathbb{S}^1 is zero by the 2-Loewner property of $\overline{\mathbb{D}}$

[Hei01, Example 8.24], hence (i) from Theorem 2.6 implies that the modulus of paths joining C to ∂Q is zero. This happens only when C is a singleton by the 2-Loewner property of $\overline{\mathbb{D}}$. Therefore h is injective. \square

Remark 3.13. Proposition 3.12 gives two simple criteria for $(\Omega/\mathcal{E}_N, \tilde{d}_N)$ to fail to be reciprocal. First, if $\mathcal{L}^2(E) > 0$, then π_N is not locally quasiconformal since Lusin's Condition (N^{-1}) is violated. Second, if π_N is not injective, then $(\Omega/\mathcal{E}_N, \tilde{d}_N)$ is not reciprocal.

4. REMOVABLE IMPLIES RECIPROCAL

The objective of this section is to prove Theorem 1.3. An outline of the proof is as follows. First, we give a pair of reductions, Lemma 4.1 and Lemma 4.2, showing that it suffices to consider only the case of admissible norm fields of the form $N = \omega \|\cdot\|_2$ defined on all of \mathbb{R}^2 , for some bounded function $\omega: \mathbb{R}^2 \rightarrow [0, \infty)$. Next, Proposition 4.3 gives a criterion for the mapping π_N in our situation to be quasiconformal: it suffices to show that π_N preserves the modulus of the path families $\Gamma(\xi_1, \xi_3; R)$ and $\Gamma(\xi_2, \xi_4; R)$ for a single rectangle R containing E with boundary edges $\xi_1, \xi_2, \xi_3, \xi_4$.

We complete the proof by verifying the modulus condition of Proposition 4.3. This part is an application of the classical theorem of uniformization onto slit domains. This argument is based on the proof of Theorem 9 in [AB50]. In Section 4.3, we extend Theorem 1.3 by relaxing the assumption that $L(N) \in L_{\text{loc}}^\infty(\Omega)$ to the assumption that $L(N) \in L_{\text{loc}}^p(\Omega)$ for some $p \in (2, \infty)$.

Lemma 4.1. An admissible norm field N on Ω is reciprocal if and only if the norm field $\widehat{N} = \omega(N) \|\cdot\|_2$ induced by the minimal stretching $\omega(N)$ is reciprocal.

Proof. As we see from Proposition 3.12, it suffices to show that the metrics generated by N and \widehat{N} are locally quasiconformally equivalent. Observe first that it follows directly from the definition that $\widehat{N} \leq N$. Since N has locally bounded distortion, every point has a neighborhood U such that $N_x \leq H\widehat{N}_x$ for some $H > 0$ and every $x \in U$. These facts imply that the corresponding distances are locally bi-Lipschitz equivalent. \square

For the following lemma, fix a subdomain $\Omega' \subset \Omega$ that contains E and is compactly contained in Ω . Let $K = \overline{\Omega'}$. Since $N = \omega \|\cdot\|_2$ is locally bounded, there exists $\alpha > 0$ such that $\omega < \alpha$ everywhere on K . We define

$$\widehat{N} = (\omega \chi_K + \alpha \chi_{\mathbb{R}^2 \setminus K}) \|\cdot\|_2.$$

The choice of α implies that \widehat{N} is admissible on \mathbb{R}^2 vanishing exactly on E . Also, \widehat{N} coincides with N in Ω' .

Lemma 4.2. The norm field $N = \omega \|\cdot\|_2$ is reciprocal in Ω if and only if the extension \widehat{N} is reciprocal in \mathbb{R}^2 . Moreover, in either one of these cases the quotient maps $\pi_{\widehat{N}}$ and π_N are 1-quasiconformal homeomorphisms.

Proof. First of all, since N and \widehat{N} are equal in Ω' , there exists a homeomorphism

$$f: \pi_N(\Omega') \rightarrow \pi_{\widehat{N}}(\Omega')$$

for which $\pi_{\widehat{N}} = \pi_N \circ f$ on Ω' . In fact, the map f is a local isometry and hence 1-quasiconformal.

Since the restrictions of π_N and $\pi_{\widehat{N}}$ to the complement of E are locally bi-Lipschitz, we deduce that they are locally quasiconformal if and only if their restrictions to Ω' are locally quasiconformal. These two conditions are equivalent for

the maps since f is quasiconformal. We conclude from Proposition 3.12 that N is reciprocal if and only if \widehat{N} is reciprocal.

We are left to check that if π_N is locally quasiconformal, then it is actually 1-quasiconformal. Combining Theorem 2.6 with the local quasiconformality of π_N , we conclude that $h = \pi_N^{-1}$ has the Sobolev regularity required for Lemma 3.10. Therefore

$$\rho = \left(\frac{1}{\omega} \chi_{\Omega \setminus E} \right) \circ h$$

is a minimal weak upper gradient of h . The change of variables formula Lemma 3.6 and Theorem 2.6 imply that the outer dilatation of h is bounded from above by one. The outer dilatation bound for π_N follows from Proposition 3.11. We conclude that π_N is 1-quasiconformal. The 1-quasiconformality of $\pi_{\widehat{N}}$ is argued in a similar manner. \square

4.1. A criterion for quasiconformality. We prove Proposition 4.3 in this section. We consider an admissible norm field $N = \omega \|\cdot\|_2$ defined on a domain $\Omega \subset \mathbb{R}^2$ vanishing exactly on a non-separating compact set $E \subset \Omega$.

We consider a quadrilateral $Q \subset \Omega$ whose boundary ∂Q does not intersect the set E . Let $(\xi_1, \xi_2, \xi_3, \xi_4)$ be a decomposition of ∂Q into four arcs labelled in counterclockwise order.

Since ∂Q does not intersect E , the restriction of π_N to ∂Q is a homeomorphism (Proposition 3.8). As a consequence of Corollary 3.9, the image $\pi_N Q$ is a Jordan domain with boundary $\pi_N \partial Q$ and the arcs $(\pi_N \xi_1, \pi_N \xi_2, \pi_N \xi_3, \pi_N \xi_4)$ decompose $\pi_N(\partial Q)$.

We fix some notation for the following proof. Let

$$\begin{aligned} \Gamma_1 &= \Gamma(\xi_1, \xi_3; Q) & \text{and} & & \widetilde{\Gamma}_1 &= \Gamma(\pi_N \xi_1, \pi_N \xi_3; \pi_N Q); \\ \Gamma_2 &= \Gamma(\xi_2, \xi_4; Q) & \text{and} & & \widetilde{\Gamma}_2 &= \Gamma(\pi_N \xi_1, \pi_N \xi_3; \pi_N Q). \end{aligned}$$

We defined $\Gamma(F_1, F_2; G)$ in Section 2.5. Notice that $\pi_N \Gamma_1 \subset \widetilde{\Gamma}_1$ and $\pi_N \Gamma_2 \subset \widetilde{\Gamma}_2$.

Proposition 4.3. Let $N = \omega \|\cdot\|_2$ be admissible. If $\text{mod } \Gamma_1 = \text{mod } \widetilde{\Gamma}_1$ and $\text{mod } \Gamma_2 = \text{mod } \widetilde{\Gamma}_2$, then the restriction of π_N to Q is a homeomorphism and 1-quasiconformal.

Proof. Proposition 3.11 and the special form of N imply that the outer dilatation of π_N is one so we only need to check that $\pi_N|_Q$ is injective and that its inverse has its outer dilatation bounded from above by one.

It was proved in [RR19] that there exists a continuous function

$$\widetilde{u}_1: \pi_N Q \rightarrow [0, 1]$$

in the Sobolev space $N^{1,2}(\pi_N Q)$ whose minimal weak upper gradient $\widetilde{\rho}_1$ is a minimizer for $\text{mod } \widetilde{\Gamma}_1$. The function \widetilde{u}_1 satisfies the boundary conditions $\widetilde{u}_1(\pi_N \xi_1) = 0$ and $\widetilde{u}_1(\pi_N \xi_3) = 1$.

Consider $u_1 = \widetilde{u}_1 \circ \pi_N$. Since $N = \omega \|\cdot\|_2$ and π_N has bounded outer dilatation, it is readily verified that $\rho_1 = (\widetilde{\rho}_1 \circ \pi_N) \omega \in L^2(Q)$ is a weak upper gradient of u_1 with L^2 -norm $\text{mod } \widetilde{\Gamma}_1 = \text{mod } \Gamma_1$. Therefore $u_1 \in N^{1,2}(Q)$.

A consequence of Weyl's lemma [AIM09, A.6.10] and continuity of u_1 is that u_1 is harmonic in the interior of Q ; it minimizes the Dirichlet energy among continuous Sobolev maps $u: Q \rightarrow [0, 1]$ with boundary values $u(\xi_1) = 0$ and $u(\xi_3) = 1$.

We repeat the above construction for the path families Γ_2 and $\widetilde{\Gamma}_2$. Let u_2 and \widetilde{u}_2 denote the corresponding functions, where $u_2(\xi_2) = 0$ and $u_2(\xi_4) = 1$.

Let M denote the modulus of Γ_1 . A consequence of Riemann mapping theorem is that $M u_2$ is a harmonic conjugate of u_1 and the restriction of $f = (u_1, M u_2)$

to the interior of Q is conformal. The map extends as a homeomorphism to the boundary ∂Q .

Let $\tilde{f} = (\tilde{u}_1, M\tilde{u}_2)$. Then $f = \tilde{f} \circ \pi_N$ by construction. This identity and bijectivity of f imply that the restriction of π_N to Q is a homeomorphism.

Since the restriction of π_N to Q is a homeomorphism and ∂Q does not intersect E , we find a Jordan neighborhood $U \supset Q$ for which $\pi_N|_U$ is a homeomorphism and that $U \cap E = Q \cap E$ (Proposition 3.8). Let h denote the inverse of $\pi_N|_U$.

We claim that $h \in N_{\text{loc}}^{1,2}(\pi_N(U), U)$. Since π_N is locally bi-Lipschitz in the complement of E and $E \cap U \subset \text{int}(Q)$, it suffices to verify that $h|_V$ is an element of $N_{\text{loc}}^{1,2}(V, U)$, where $V = \pi_N(\text{int}(Q))$. This regularity follows readily since the restriction of f to the interior of Q is locally bi-Lipschitz, \tilde{f} is an element of $N^{1,2}(\pi_N Q, [0, 1] \times [0, M])$, and $h = f^{-1} \circ \tilde{f}$ in V . Now the outer dilatation bound $K_O(h) \leq 1$ follows from Lemma 3.10 and the change of variables formula of π_N . \square

Remark 4.4. Proposition 4.3 is related to a question posed by Rajala in [Raj17]. Rajala asks whether the reciprocal upper bound (12) implies that points have zero modulus in the sense of (13). This is the case whenever $N = \omega \|\cdot\|_2$ is admissible and satisfies the sharp modulus upper bound of one in (12). The proof relies on the existence and Sobolev regularity of energy minimizers of quadrilaterals on metric spaces studied on [Raj17, RR19]. The key observation is that if π_N preserves the modulus of opposite sides of ∂Q , then the quotient map π_N pulls back these minimizers to harmonic minimizers on the Euclidean space.

4.2. Proof of Theorem 1.3. Let $E \subset \mathbb{R}^2$ be removable for conformal mappings. We want to prove that for any domain $\Omega \supset E$ and admissible norm field $N: \Omega \times \mathbb{R}^2 \rightarrow [0, \infty)$ vanishing exactly on E , the quotient space $(\Omega/\mathcal{E}_N, \tilde{d}_N)$ is reciprocal.

As shown in Lemma 4.1 and Lemma 4.2, we only need to consider the case where $\Omega = \mathbb{R}^2$ and $N = \omega \|\cdot\|_2$.

Let $R = [a, b] \times [c, d]$ be a rectangle whose interior contains E . Let $\xi_1 = \{a\} \times [c, d]$, $\xi_2 = [a, b] \times \{c\}$, $\xi_3 = \{b\} \times [c, d]$, and $\xi_4 = [a, b] \times \{d\}$. Let $\Gamma_1 = \Gamma(\xi_1, \xi_3; R)$ and $\Gamma_2 = \Gamma(\xi_2, \xi_4; R)$.

Let $\tilde{\Gamma}_1$ denote the family of paths joining $\pi_N \xi_1$ to $\pi_N \xi_3$ in $\pi_N R$ and $\tilde{\Gamma}_2$ the family of paths joining $\pi_N \xi_2$ to $\pi_N \xi_4$ in $\pi_N R$. We claim that $\text{mod } \tilde{\Gamma}_1 = \text{mod } \Gamma_1$ and $\text{mod } \tilde{\Gamma}_2 = \text{mod } \Gamma_2$. Proposition 4.3 then implies that π_N is 1-quasiconformal in the interior of R . Since R is an arbitrary rectangle containing E , it then follows that π_N is globally 1-quasiconformal.

Observe that the inequalities $\text{mod } \tilde{\Gamma}_1 \geq \text{mod } \Gamma_1$ and $\text{mod } \tilde{\Gamma}_2 \geq \text{mod } \Gamma_2$ hold in general by Proposition 3.11. Thus we only need to verify the opposite inequalities.

A standard fact is that there is a sequence of finitely connected domains $\Omega_k \subset \mathbb{R}^2 \setminus E$ such that $\bar{\Omega}_k \subset \Omega_{k+1}$ for all $k \in \mathbb{N}$, each component of $\partial \Omega_k$ is a closed analytic Jordan path, and $\bigcup_{k=1}^{\infty} \Omega_k = \mathbb{R}^2 \setminus E$. We assume without loss of generality that $\partial R \subset \Omega_1$.

For each $n \in \mathbb{N}$, there exists a conformal embedding $\varphi_n: \Omega_n \rightarrow \mathbb{R}^2$ normalized as

$$\varphi_n(z) = z + \frac{a_{1,n}}{z} + \dots$$

near ∞ such that the real part of $a_{1,n}$ is the smallest among all conformal embeddings $\psi: \Omega_n \rightarrow \mathbb{R}^2$ of the form

$$(25) \quad \psi(z) = z + \frac{a}{z} + \dots$$

See for example Section V.2 of [Gol69].

For each $n \in \mathbb{N}$, the minimizer φ_n is unique and its image is a domain $U_n \subset \mathbb{R}^2$ whose complement consists of finitely many arc segments parallel to the vertical axis.

Fix n and consider $k \geq n$. By the minimality of the real part of $a_{1,n}$, we have that $0 \geq \operatorname{Re}(a_{1,k}) \geq \operatorname{Re}(a_{1,n})$. Hence the mappings $\varphi_k|_{\Omega_n} : \Omega_n \rightarrow \mathbb{R}^2$ form a normal family. See for example the proof of Theorem 1 of [Gol69, Section V.2] for details. A diagonal argument then implies that (φ_n) is a normal family. Thus every subsequence of (φ_n) has a further subsequence converging uniformly on compact sets to a conformal map $f : \mathbb{R}^2 \setminus E \rightarrow \mathbb{R}^2$ satisfying the normalization (25) around ∞ . By the removability of E , the map extends to a Möbius transformation, and thus (25) implies that $f(z) = z$ for all $z \in \mathbb{R}^2$. Hence the sequence (φ_k) itself must converge to the identity map uniformly on compact sets of $\mathbb{R}^2 \setminus E$.

Let Q_n denote the quadrilateral bounded by the Jordan curve $\varphi_n(\partial R)$. The quadrilateral Q_n converges to R with respect to Hausdorff distance as $n \rightarrow \infty$. Let π_1 and π_2 denote projection onto the x -axis and y -axis, respectively, and let $a_n = \sup \pi_1(\varphi_n(\xi_1))$, $b_n = \inf \pi_1(\varphi_n(\xi_3))$, $c_n = \inf \pi_2(\varphi_n(\xi_2))$ and $d_n = \sup \pi_2(\varphi_n(\xi_4))$.

Let $R_n = [a_n, b_n] \times [c_n, d_n]$ and $\widehat{E}_n = \mathbb{R}^2 \setminus \varphi_n(\Omega_n)$. Observe that \widehat{E}_n consists of finitely many vertical slits S_1, \dots, S_m .

There exists n_0 such that for all $n \geq n_0$, the slits \widehat{E}_n are contained in the interior of R_n , and that $0 < b_n - a_n$ and $0 < d_n - c_n$. Fix such an n . We claim that

$$(26) \quad \operatorname{mod} \widetilde{\Gamma}_1 \leq \frac{d_n - c_n}{b_n - a_n}.$$

Consider the function $\rho_n : \mathbb{R}^2 / \mathcal{E}_N \rightarrow [0, \infty]$ defined as zero in the complement of $\pi_N(\Omega_n)$, and otherwise by

$$\rho_n = \left(\frac{\chi_{R_n \setminus \widehat{E}_n}}{b_n - a_n} \circ \varphi_n \frac{J_{\varphi_n}^{-1/2}}{\omega} \right) \circ (\pi_N|_{\Omega_n})^{-1}.$$

We claim that ρ_n is admissible for $\widetilde{\Gamma}_1$. Let $\gamma \in \widetilde{\Gamma}_1$ be locally rectifiable with respect to \widetilde{d}_N .

We consider the restriction of γ to the set $I = \gamma^{-1}(\pi_N \varphi_n^{-1}(R_n \setminus \widehat{E}_n))$. We have

$$\int_{\gamma} \rho_n ds_{\widetilde{d}_N} \geq \int_I \rho_n \circ \gamma v_{\gamma} d\mathcal{L}^1.$$

Here $\theta = \varphi_n \circ (\pi_N|_{\Omega_{n+1}})^{-1} \circ \gamma|_I$ is well-defined and

$$\int_I \rho_n \circ \gamma v_{\gamma} d\mathcal{L}^1 = \int_I \frac{\chi_{R_n \setminus \widehat{E}_n}}{b_n - a_n} \circ \theta v_{\theta} d\mathcal{L}^1.$$

Since \widehat{E}_n consists of finitely many vertical slits, we conclude using the area formula for paths and the projection onto the x -axis that

$$\int_I \frac{\chi_{R_n \setminus \widehat{E}_n}}{b_n - a_n} \circ \theta v_{\theta} d\mathcal{L}^1 \geq \frac{1}{b_n - a_n} \mathcal{L}^1(|\pi_1 \circ \theta|) \geq 1.$$

Therefore

$$\int_{\gamma} \rho_n ds_{\widetilde{d}_N} \geq 1,$$

and we conclude that ρ_n is admissible. The change of variables formulas for π_N and φ_n yield that

$$\int \rho_n^2 d\mathcal{H}_{\widetilde{d}_N}^2 = \frac{d_n - c_n}{b_n - a_n}.$$

This verifies (26). Finally, observe that $d_n - c_n \rightarrow d - c$ and $b_n - a_n \rightarrow b - a$ as $n \rightarrow \infty$. This shows that

$$\text{mod } \tilde{\Gamma}_1 \leq \frac{d - c}{b - a} = \text{mod } \Gamma_1.$$

A similar argument, using conformal mappings onto horizontal slit domains, shows that $\text{mod } \tilde{\Gamma}_2 \leq \text{mod } \Gamma_2$. This completes the proof.

4.3. An extension of Theorem 1.3 to integrable norm fields. In this section we extend Theorem 1.3 to the case of lower semicontinuous norm fields N with locally bounded distortion such that $L(N) \in L^p_{\text{loc}}(\Omega)$ for some $p \in (2, \infty)$. We assume that N vanishes exactly on a compact set $E \subset \Omega$ that is removable for conformal mappings.

For this section, we allow the possibility for N_x to be infinite at some points $x \in \Omega$. To say this more precisely, in the definition of seminorm in Section 2.3, we consider a seminorm to be a function $S: \mathbb{R}^2 \rightarrow [0, \infty]$ satisfying the same assumptions listed there, following the convention that $0 \cdot \infty = 0$. An admissible norm field is now a function $N: \Omega \times \mathbb{R}^2 \rightarrow [0, \infty]$ satisfying the conditions of Definition 3.1, except that local boundedness of N is now replaced by the assumption that $L(N) \in L^p_{\text{loc}}(\Omega)$. Observe that the local boundedness of the distortion then implies that if $N_x(v) = \infty$ for some $v \in \mathbb{R}^2 \setminus \{0\}$, then N_x must have the form

$$N_x(v) = \begin{cases} \infty & \text{if } v \neq 0 \\ 0 & \text{if } v = 0 \end{cases}.$$

In particular, $\omega(N_x) = L(N_x) = \infty$. Note also that the minimal stretching $\omega(N)$ is lower semicontinuous, and that Lemma 3.3 remains true for $x \in \Omega$ with $\omega(N_x) < \infty$.

We define the pseudodistance d_N exactly as in Definition 3.2. It is no longer immediately clear that $d_N(x, y) < \infty$ for any given pair of points $x, y \in \Omega$, but this can be shown from the local L^p -integrability of $L(N_x)$, since $p > 2$. See [LW18, Thm. 1.1] and [CS19, Prop. 5.12] for proofs of the analogous fact in related settings.

As before, we identify $x, y \in \Omega$ if $d_N(x, y) = 0$ and let X denote the corresponding quotient space. Let $\pi: \Omega \rightarrow X$ denote the associated quotient map. The quotient distance d_X on X is defined as follows: for every $x, y \in X$, we set $d_X(x, y) = d_N(\pi^{-1}(x), \pi^{-1}(y))$, observing that this is independent of the choice of element in $\pi^{-1}(x)$ and $\pi^{-1}(y)$ and hence well-defined.

Next, we establish that π is continuous and an element of $N^{1,p}_{\text{loc}}(\Omega, X)$. To this end, for every $x \in \Omega$, $f_x(z) = d_N(x, z)$ is measurable as a consequence of [HKST15, Theorem 9.3.1]. Since $L(N)$ is an upper gradient of π , it is also an upper gradient of f_x . Then Morrey's embedding theorem [HKST15, Theorem 9.2.14] implies that f_x is locally Hölder continuous with Hölder exponent and Hölder constant independent of x . This implies that π is locally Hölder continuous. Thus π is continuous with upper gradient $L(N) \in L^p_{\text{loc}}(\Omega)$. We are now ready for the main result of this section. We recall that N is assumed to vanish on a compact set E removable for conformal mappings.

Proposition 4.5. The metric space X has locally finite Hausdorff 2-measure, and the quotient map π is a locally quasiconformal homeomorphism. In particular, X is a quasiconformal surface.

Proof. We first prove that π is a homeomorphism. To this end, let $\omega(z) = \omega(N_z)$ and $\hat{N}_z = \omega(z) \|\cdot\|_2$ for every $z \in \Omega$. For each $k \in \mathbb{N}$, we define the function $\omega_k: \Omega \rightarrow [0, \infty)$ by

$$\omega_k(z) = \min \{ \omega(z), k \}.$$

Each function ω_k is bounded and lower semicontinuous in Ω , and $\omega_k(z) = 0$ if and only if $\omega(z) = 0$. For every $z \in \Omega$, the sequence $(\omega_k(z))_{k=1}^\infty$ is non-decreasing and converges to $\omega(z)$.

Let $N_k = \omega_k \|\cdot\|_2$ and $d_k = d_{N_k}$. Since N_k is bounded and lower semicontinuous, Theorem 1.3 implies that $\pi_k: \Omega \rightarrow (\Omega, d_k)$ defined by $\pi_k(z) = z$ is a 1-quasiconformal homeomorphism. Since $N_k \leq \omega(N) \|\cdot\|_2 \leq N$ everywhere, we see that

$$d_k(\pi_k(x), \pi_k(y)) \leq d_X(\pi(x), \pi(y))$$

for all $x, y \in \Omega$. Since π_k is a homeomorphism, we see that π is injective. Now the map $\psi_k: X \rightarrow (\Omega, d_k)$ defined by $\psi_k = \pi_k \circ \pi^{-1}$ is 1-Lipschitz, hence $\pi^{-1} = \pi_k^{-1} \circ \psi_k$ is continuous. Therefore π is a homeomorphism.

Recall that N has locally bounded distortion. From this and the fact that, for every $x \in X$, $\pi^{-1}(\overline{B}_X(x, r))$ is compact for sufficiently small $r > 0$, we see that the induced distances d_N and $d_{\widehat{N}}$ are locally bi-Lipschitz equivalent. We assume from this point onwards, without loss of generality, that $N = \widehat{N} = \omega \|\cdot\|_2$.

Let Γ_0 denote the family of paths along which $\omega = L(N)$ fails to be integrable. Since $\omega \in L^p_{\text{loc}}(\Omega) \subset L^2_{\text{loc}}(\Omega)$, the family Γ_0 has zero modulus. Recall from Lemma 3.7 that, for any absolutely continuous path θ in Ω ,

$$\ell_{d_X}(\pi \circ \theta) \geq \lim_{k \rightarrow \infty} \ell_{d_k}(\psi_k \circ \theta) = \lim_{k \rightarrow \infty} \ell_{N_k}(\theta) = \ell_N(\theta),$$

where the latter equality follows from monotone convergence. If $\theta \notin \Gamma_0$, we have $\ell_N(\theta) < \infty$ and the definition of d_X implies $\ell_{d_X}(\pi \circ \theta) \leq \ell_N(\theta)$. So $\ell_{d_X}(\pi \circ \theta) = \ell_N(\theta)$. Since the equality holds for all absolutely continuous paths outside the negligible family Γ_0 , we conclude from Sections 3.3 and 3.4 of [LW18] that $L(N) = \omega$ is a minimal weak upper gradient of $\pi \in N^{1,2}_{\text{loc}}(\Omega, X)$ and the Jacobian of π equals $J_2(N) = \omega^2$ \mathcal{L}^2 -almost everywhere. Since we also have that $\pi \in N^{1,p}_{\text{loc}}(\Omega, X)$ for $p > 2$, it satisfies Lusin's Condition (N) [Vod00, Theorem 7.1]. Therefore, for each compact set $K \subset \Omega$,

$$\mathcal{H}^2_X(\pi(K)) = \int_K \omega^2 d\mathcal{L}_2 < \infty.$$

We conclude that X has locally finite Hausdorff 2-measure. An application of Theorem 2.6 yields that $K_O(\pi) = 1$.

The proof is complete after we verify $K_O(\pi^{-1}) = 1$. Since π_k is 1-quasiconformal for every k , it suffices to verify $K_O(\psi_k) = 1$ for some k . To this end, we fix $k \in \mathbb{N}$ and recall that ψ_k is 1-Lipschitz.

Since π_k is a quasiconformal homeomorphism, it satisfies Lusin's Condition (N^{-1}). This implies that the map π^{-1} satisfies Lusin's Condition (N). As a consequence, the Jacobian of ψ_k coincides with ρ_k^2 for $\rho_k = ((\omega_k/\omega)\chi_{\Omega \setminus E}) \circ \pi^{-1}$.

Since ψ_k is Lipschitz, we have $\psi_k \in N^{1,2}_{\text{loc}}(X, (\Omega, d_k))$. We claim that any minimal weak upper gradient of ψ_k coincides with ρ_k almost everywhere in X . If we verify this, then $K_O(\psi_k) = 1$ follows from Theorem 2.6.

Consider an absolutely continuous path $\gamma: [0, 1] \rightarrow X$ with $|\gamma| \subset X \setminus \pi(E)$. Then $\psi_k \circ \gamma$ is absolutely continuous, and since d_k and $\|\cdot\|_2$ are locally bi-Lipschitz equivalent in a neighborhood of the image of $\theta = \pi_k^{-1} \circ \psi_k \circ \gamma$, the path θ is absolutely continuous with respect to $\|\cdot\|_2$. Then by monotone convergence and Lemma 3.7,

$$\ell_N(\theta) = \lim_{n \rightarrow \infty} \ell_{N_n}(\theta) = \lim_{n \rightarrow \infty} \ell_{d_n}(\psi_n \circ \gamma).$$

Since every ψ_n is 1-Lipschitz,

$$\lim_{n \rightarrow \infty} \ell_{d_n}(\psi_n \circ \gamma) \leq \ell_{d_X}(\gamma).$$

Therefore $\ell_N(\theta) \leq \ell_{d_X}(\gamma) < \infty$, and, by the construction of d_X , $\ell_{d_X}(\gamma) \leq \ell_N(\theta)$.

Since $\ell_N(\theta) = \ell_{d_X}(\gamma)$ holds for every subpath of γ , we see that $v_\gamma = N \circ D\theta = \omega \circ \theta v_\theta$ and $v_{\psi_k \circ \gamma} = \omega_k \circ \theta v_\theta$ almost everywhere in the domain of γ . We conclude from this that

$$(27) \quad v_{\psi_k \circ \gamma} = \rho_k \circ \gamma v_\gamma$$

almost everywhere with respect to the length measure of γ . If $\tilde{\Gamma}_0$ denotes the family of absolutely continuous paths in X that have positive length on the set $\pi(E)$, then $\mathcal{H}_X^2(\pi(E)) = 0$ implies $\text{mod } \tilde{\Gamma}_0 = 0$. The equality (27) remains valid for every absolutely continuous path $\gamma \notin \tilde{\Gamma}_0$, which implies that ρ_k is a weak upper gradient of ψ_k . The minimality of ρ_k is immediate from (27). So any minimal weak upper gradient of ψ_k coincides with ρ_k \mathcal{H}_X^2 -almost everywhere. \square

Remark 4.6. The norm field $N = \omega \|\cdot\|_2$ defined by the weight $\omega(x) = \|x\|_2^{-1} (1 - \log \|x\|_2)^{-1} \in L^2(\mathbb{D})$ induces a complete hyperbolic metric on the punctured unit disk. In particular, the origin is at infinite distance from any other point. Consequently, the assumption $p > 2$ in Proposition 4.5 cannot be relaxed.

5. RECIPROCAL IMPLIES REMOVABLE

This section is dedicated to a proof of Theorem 1.4. Recall that we consider a compact set $E \subset \Omega$ for which $\Omega \setminus E$ is connected, together with the norm field N defined by $N_x = \min\{1, d_{\|\cdot\|_2}(E, x)^p\} \|\cdot\|_2$ for some $p > \max\{\dim_{\mathcal{H}} E - 1, 0\}$. The norm field N induces a decomposition \mathcal{E}_N of Ω , a metric \tilde{d}_N on Ω/\mathcal{E}_N , and a quotient map $\pi: \Omega \rightarrow (\Omega/\mathcal{E}_N, \tilde{d}_N)$, as described in Section 3.

5.1. Decay of the norm field near E . The following lemma states that if N decays to zero sufficiently fast near E , then each component of E collapses to a point under the quotient map π_N .

Lemma 5.1. Let $N_x = \min\{1, d_{\|\cdot\|_2}(x, E)^p\} \|\cdot\|_2$. For all $p > \max\{\dim_{\mathcal{H}} E - 1, 0\}$, $\mathcal{H}_{\tilde{d}_N}^1(\pi_N(E)) = 0$. Consequently, the preimage of every $x \in \pi_N(E)$ is a connected component of E .

Proof. Let $p > \dim_{\mathcal{H}} E - 1$ and let $\varepsilon > 0$. By the definition of Hausdorff dimension, there exists $\delta > 0$ and a countable collection of sets $\mathcal{A} = \{A_j\}$ such that $E \subset \bigcup_j A_j$, $\text{diam}_{\|\cdot\|_2} A_j \leq \delta$ for all j , and

$$\sum_j (\text{diam}_{\|\cdot\|_2} A_j)^{p+1} < \varepsilon.$$

Without loss of generality, we may assume that $A_j \cap E \neq \emptyset$ for all j . Let $d_j = \text{diam}_{\|\cdot\|_2} A_j$. Thus $A_j \subset \overline{B}_{\|\cdot\|_2}(y, d_j)$ for some $y \in E$. By integrating N over the straight-line path from y to a point $z \in A_j$, it follows that

$$d_N(y, z) \leq \int_0^{d_j} t^p dt = \frac{d_j^{p+1}}{p+1}.$$

Thus $\text{diam}_{d_N} A_j \leq 2(p+1)^{-1} d_j^{p+1} < 2(p+1)^{-1} \delta^{p+1}$, and we get $\sum_j \text{diam}_{d_N} A_j < 2(p+1)^{-1} \varepsilon$. This is sufficient to show that $\mathcal{H}_{\tilde{d}_N}^1(\pi_N(E)) = 0$.

Next, let $x \in \pi_N(E)$. Proposition 3.8 implies that $\pi_N^{-1}(x)$ is a subset of a connected component F of E . Since $\pi_N(F)$ is connected and compact subset of $\pi_N(E)$, we have that

$$\text{diam } \pi_N(F) \leq \mathcal{H}_{\tilde{d}_N}^1(\pi_N(F)) \leq \mathcal{H}_{\tilde{d}_N}^1(\pi_N(E)) = 0.$$

Hence $\pi_N(F) = x$ and we must have $F = \pi_N^{-1}(x)$. \square

5.2. Proof of Theorem 1.4. We first observe that if $(\Omega/\mathcal{E}_N, \tilde{d}_N)$ is reciprocal, then the space formed by taking the same set E and the same definition for N , but applied to all points $x \in \mathbb{R}^2$, is also reciprocal. Thus the choice of domain Ω is not relevant for the proof, and we assume for the remainder of the section that $\Omega = \mathbb{R}^2$.

We prove the contrapositive: if E is not removable for conformal mappings, then $(\mathbb{R}^2/\mathcal{E}_N, \tilde{d}_N)$ is not reciprocal.

Let $E \subset \mathbb{R}^2$ be a set that is not removable for conformal mappings. As a consequence of Proposition 2.8, there is a compact set $\hat{E} \subset \mathbb{R}^2$ of positive measure and a conformal map $f: \mathbb{R}^2 \setminus \hat{E} \rightarrow \mathbb{R}^2 \setminus E$. Let $\hat{N} = \chi_{\mathbb{R}^2 \setminus \hat{E}} \|\cdot\|_2$ and let $\hat{\pi}: \mathbb{R}^2 \rightarrow (\mathbb{R}^2/\mathcal{E}_{\hat{N}}, \tilde{d}_{\hat{N}})$ be the associated quotient map. Observe that \hat{N} is an admissible norm field vanishing on the set \hat{E} .

The following lemma states that f extends to a mapping of the respective quotient spaces. For brevity, let $\hat{Y} = \mathbb{R}^2/\mathcal{E}_{\hat{N}}$ and $Y = \mathbb{R}^2/\mathcal{E}_N$.

Lemma 5.2. The map $f: \mathbb{R}^2 \setminus \hat{E} \rightarrow \mathbb{R}^2 \setminus E$ induces a continuous monotone map $\hat{f}: \hat{Y} \rightarrow Y$. That is, there is a monotone map $\hat{f}: \hat{Y} \rightarrow Y$ satisfying $\hat{f} \circ \hat{\pi}(x) = \pi_N \circ f(x)$ for all $x \in \mathbb{R}^2 \setminus \hat{E}$.

Proof. Let $y \in \hat{Y}$, and let \hat{F} denote its preimage under $\hat{\pi}$. If $\hat{F} = \{x\}$ for some point $x \notin \hat{E}$, then we set $\hat{f}(y) = \pi_N \circ f(x)$.

Otherwise, \hat{F} is a subset of some component \hat{A} of \hat{E} . For all $m \in \mathbb{N}$, let $\hat{\gamma}_m$ be a Jordan path with image contained in $B_{\|\cdot\|_2}(\hat{A}, 1/m) \setminus \hat{E}$ that separates \hat{A} and infinity. The curve $|\hat{\gamma}_m|$ is the boundary of a closed region \hat{A}_m containing \hat{A} . We assume without loss of generality that $|\hat{\gamma}_{m+1}| \subset \hat{A}_m$ for all m .

By assumption, $\gamma_m = f \circ \hat{\gamma}_m$ is a Jordan loop whose image bounds a compactly contained domain A_m . Let $A = \bigcap_m \bar{A}_m$. It is immediate that A is nonempty and compact. The intersection is also connected; see for example Section 28 of [Wil70]. This implies that A is a connected component of E . Therefore $\pi_N(A)$ is a point by Lemma 5.1. We define $\hat{f}(y) = \pi_N(A)$.

We now check that \hat{f} is continuous. Let $y \in \hat{Y}$ and let (y_n) be a sequence in \hat{Y} converging to y . Let $\hat{F}_n = \hat{\pi}^{-1}(y_n)$. In the case that $\hat{F} = \{x\}$ for some $x \notin \hat{E}$, the continuity is obvious. Otherwise, we proceed as follows. For each fixed $m \in \mathbb{N}$, $F_n \subset \hat{A}_m$ for sufficiently large n . This implies that $\hat{f}(y_n) \subset \pi_N(A_m)$. Therefore the accumulation points of $\hat{f}(y_n)$ are in the intersection of $\pi_N(A_m)$. Since the intersection equals $\pi_N(A)$, the sequence $\hat{f}(y_n)$ converges to $\pi_N(A) = \hat{f}(y)$. The continuity follows.

By construction, the preimage of a point in \mathbb{R}^2/\mathcal{E} under $\hat{\pi} \circ \hat{f}$ is either a single-point set or a component of \hat{E} . We conclude that \hat{f} is monotone. \square

Let $R = [a, b] \times [c, d] \subset \mathbb{R}^2$ be a rectangle whose interior contains \hat{E} . Let Γ_1 denote the family of paths $\gamma_t: [a, b] \rightarrow \mathbb{R}^2$, where $t \in [c, d]$, defined by $\gamma_t(s) = (s, t)$. Thus Γ_1 is a foliation of R by horizontal paths. Let Γ_2 denote the corresponding foliation of R by vertical paths.

Next, let Q be the Jordan domain bounded by $f(\partial R)$, and let $\xi_1, \xi_2, \xi_3, \xi_4$ denote, respectively, the image of the left, bottom, right, and top side of R . Let $\tilde{\Gamma}_1$ denote the family paths joining $\pi_N \xi_1$ to $\pi_N \xi_3$ in $\pi_N Q$ and $\tilde{\Gamma}_2$ the family of paths joining $\pi_N \xi_2$ to $\pi_N \xi_4$.

By Lemma 4.2, it suffices to show that Y is not 1-reciprocal. Thus the proof is complete after we verify the inequalities

$$(28) \quad 1 < \text{mod } \hat{\pi}\Gamma_1 \text{ mod } \hat{\pi}\Gamma_2$$

and

$$(29) \quad \text{mod } \widehat{\pi}\Gamma_1 \text{ mod } \widehat{\pi}\Gamma_2 \leq \text{mod } \widetilde{\Gamma}_1 \text{ mod } \widetilde{\Gamma}_2.$$

Define the function $P: \mathbb{R}^2 \rightarrow [0, \infty]$ by

$$P(x) = \begin{cases} L(N_{f(x)}) \|D_x f\| & \text{if } x \notin \widehat{E} \\ 0 & \text{if } x \in \widehat{E}. \end{cases}$$

Since N is a weighted Euclidean norm and f is conformal in the complement of \widehat{E} , it follows that $N \circ D_x f(v) = P(x) \|v\|_2$ for all $v \in \mathbb{R}^2$ and all $x \in \mathbb{R}^2 \setminus \widehat{E}$.

We consider the function $\widehat{P}: \mathbb{R}^2/\widehat{\mathcal{E}} \rightarrow [0, \infty]$ defined by taking $\widehat{P}(x) = P(\widehat{\pi}^{-1}(x))$. Observe that \widehat{P} is well-defined since $\widehat{\pi}$ is injective outside of \widehat{E} . Loosely speaking, \widehat{P} is a weak upper gradient of \widehat{f} .

Let $\rho: \widehat{f}(\widehat{R}) \rightarrow [0, \infty]$ be an admissible function for $\widetilde{\Gamma}_1$, and let $\widehat{\rho} = (\rho \circ \widehat{f})\widehat{P}$. We first observe that

$$(30) \quad \int_{\widehat{R}} \widehat{\rho}^2 d\mathcal{H}_d^2 = \int_{\widehat{f}(\widehat{R})} \rho^2 d\mathcal{H}_{d_N}^2.$$

Indeed, the integrals are left unchanged by the removal of $\pi_N(E)$ and $\widehat{\pi}(\widehat{E})$ from both sides. With this reduction, the identity (30) follows from the Jacobian identities $J_f \equiv \|Df\|^2$, $J_{\widehat{\pi}} = \chi_{\mathbb{R}^2 \setminus E}$, and $J_{\pi_N} = L^2(N)$.

Next, we claim that $\widehat{\rho}$ is weakly admissible for $\widehat{\pi}\Gamma_1$. Let $\widehat{\gamma}_t$ denote the image under $\widehat{\pi}$ of the horizontal path γ_t in the quotient space $\mathbb{R}^2/\widehat{\mathcal{E}}$. Lemma 3.7 implies that

$$(31) \quad v_{\widehat{f} \circ \widehat{\gamma}_t}(s) = N \circ Df \circ D\gamma_t(s) = \widehat{P} \circ \widehat{\gamma}_t(s) v_{\widehat{\gamma}_t}(s)$$

for \mathcal{L}^1 -almost every $s \in [a, b] \setminus \gamma_t^{-1}(\widehat{E})$ and that the total variation of $\widehat{\gamma}_t$ in $\widehat{\pi}\widehat{E}$ is zero. Similarly, since $\mathcal{H}_{d_N}^1(\pi_N(E)) = 0$ by Lemma 5.1, the area formula [Fed69, Theorem 2.10.13] for paths implies that the total variation of $\widehat{f} \circ \widehat{\gamma}_t$ in $\pi_N(E)$ is zero. We conclude that $\widehat{f} \circ \widehat{\gamma}_t$ is absolutely continuous as long as the right-hand side of (31) is integrable.

Observe that (30) holds with the characteristic function $\chi_{\widehat{f}(\widehat{R})}$ in place of ρ and \widehat{P} in place of $\widehat{\rho}$. Then an application of Fubini's theorem implies that the function in the right-hand side of (31) is integrable for \mathcal{L}^1 -almost every t . For such t , we conclude from (31) that

$$1 \leq \int_{\widehat{f} \circ \widehat{\gamma}_t} \rho ds = \int_{\widehat{\gamma}_t} \widehat{\rho} ds.$$

Therefore $\widehat{\rho}$ is weakly admissible for $\widehat{\pi}\Gamma_1$, and the equality (30) implies that

$$\text{mod } \widehat{\pi}\Gamma_1 \leq \text{mod } \widetilde{\Gamma}_1.$$

A similar argument applied to the path family $\widehat{\pi}\Gamma_2$ gives $\text{mod } \widehat{\pi}\Gamma_2 \leq \text{mod } \widetilde{\Gamma}_2$. The inequality (29) now follows.

To conclude the proof, we prove (28). Let ρ be admissible for $\widehat{\pi}\Gamma_1$. Then for all $t \in [c, d]$, we have $1 \leq \int_c^d \rho \circ \widehat{\pi} \chi_{\mathbb{R}^2 \setminus \widehat{E}}(s, t) dt$. Applying Fubini's theorem and Hölder's inequality gives

$$d - c \leq \int_R \rho \circ \widehat{\pi} \chi_{\mathbb{R}^2 \setminus \widehat{E}} d\mathcal{L}^2 \leq \left(\int_R \rho \circ \widehat{\pi}^2 \chi_{\mathbb{R}^2 \setminus \widehat{E}} d\mathcal{L}^2 \right)^{1/2} \mathcal{L}^2(R \setminus \widehat{E})^{1/2}.$$

After rearranging and taking the infimum over admissible ρ , we find that

$$(d - c)^2 / \mathcal{L}^2(R \setminus \widehat{E}) \leq \text{mod } \widehat{\pi}\Gamma_1.$$

The analogous argument gives $(b-a)^2/\mathcal{L}^2(R \setminus \widehat{E}) \leq \text{mod } \widehat{\pi}\Gamma_2$. Thus

$$1 < \frac{(b-a)^2(d-c)^2}{\mathcal{L}^2(R \setminus \widehat{E})^2} \leq \text{mod } \widehat{\pi}\Gamma_1 \text{ mod } \widehat{\pi}\Gamma_2.$$

6. LINEAR CANTOR SETS: TWO EXAMPLES

We call a Cantor set $E \subset \mathbb{R} \times \{0\}$ a *linear Cantor set*. As remarked in Section 1.3, a norm field vanishing on a linear Cantor set E of positive length may or may not be reciprocal. For completeness, we include here two explicit examples to illustrate both of these cases. Recall from the discussion following the statement of Theorem 1.4 that a compact set $E \subset [0, 1] \times \{0\}$ is removable for conformal mappings if there exists an admissible norm field N vanishing on E that is reciprocal. Such an E is necessarily a linear Cantor set by Proposition 3.12. Conversely, if there exists an admissible norm field vanishing on a linear Cantor set E that is not reciprocal, then E is not removable for conformal mappings. Versions of these examples are already present in [AB50, Section 7]. A closely related construction, and the one that we directly based Example 6.1 on, is found in Section 11 of an early version of the paper [Sch95].

Example 6.1. We construct a lower semicontinuous weight $\omega: \mathbb{R}^2 \rightarrow [0, \infty]$ that vanishes on a Cantor set $E \subset [0, 1] \times \{0\}$ of positive length such that the space (\mathbb{R}^2, d_ω) is not reciprocal. The idea is to make E sufficiently large so that the modulus of the path family joining $(0, 0)$ to $(0, 1)$ in (\mathbb{R}^2, d_ω) is positive.

Identify $[0, 1]$ with the set $[0, 1] \times \{0\} \subset \mathbb{R}^2$. Let $a_1 = 1/2$, and now define inductively sequences $(a_j), (b_j)$ by the rules $b_j = a_j / \exp(4^j)$ and $a_{j+1} = (a_j - b_j)/2$. Let I_1 be a closed interval centered at $t_1 = 1/2$ of length $2b_1$. Define next intervals I_j inductively as follows. Assume that we have a collection of disjoint intervals I_1, \dots, I_{j-1} . From the complement of $(0, 1) \setminus \bigcup_{k=1}^{j-1} I_k$, choose an open interval J of largest length. Let t_j be the midpoint of J , and let I_j be the closed interval centered at t_j of length $2b_j$. We record the observation that $d_{\|\cdot\|_2}(t_j, \{0, 1\}) = \min\{t_j, 1 - t_j\} \geq a_j$. Let $E = [0, 1] \setminus \bigcup_j I_j$, and let $\omega = \chi_{\mathbb{R}^2 \setminus E}$.

Consider now an interval I_j . Assume in the first case that $t_j \leq 1/2$. For all $t \in (0, t_j - b_j)$, let $\gamma_{j,t}$ be the path that connects t to $2t_j - t$ along the upper semicircle of the circle centered at t_j with radius $t_j - t$. Let Γ_j be the family of all such paths $\gamma_{j,t}$. Observe that Γ_j is a full-modulus subfamily of the family of paths in the upper half-plane H that separate the sets $\overline{B}((t_j, 0), b_j)$ and $H \setminus B((t_j, 0), t_j)$.

Since the metric speed of $\gamma \in \Gamma_j$ with respect to Euclidean distance and with respect to d_ω coincide almost everywhere along γ , the modulus of Γ_j with respect to the metric d_ω equals the Euclidean modulus: $\text{mod}_{d_\omega} \Gamma_j = \log(t_j/b_j)/\pi$. See for example [Hei01, Lemma 7.18]. Next, we consider the case that $t_j > 1/2$. For all $t \in (t_j + b_j, 1)$, let $\gamma_{j,t}$ be the path connecting t to $2t_j - t$ along the upper semicircle of the circle centered at t_j with radius $t - t_j$. In this case, we have $\text{mod}_{d_\omega} \Gamma_j = \log((1 - t_j)/b_j)/\pi$.

We claim that the metric d_ω violates reciprocity condition (13). Let $F_1 = \{(0, 0)\}$ and $F_2 = \{(1, 0)\}$ and let $\Gamma = \Gamma(F_1, F_2; \mathbb{R}^2)$. Recall the notation $\Gamma(F_1, F_2; G)$ defined in Section 2.5.

Observe that Γ is a subfamily of $\Gamma(F_1, \mathbb{R}^2 \setminus \mathbb{D}; \mathbb{R}^2)$, which is majorized by the annular path families $\Gamma(B_{\|\cdot\|_2}(0, \varepsilon), \mathbb{R}^2 \setminus \mathbb{D}; \mathbb{R}^2)$ for all $\varepsilon > 0$. In particular,

$$\text{mod}_{d_\omega} \Gamma(B_{\|\cdot\|_2}(0, \varepsilon), \mathbb{R}^2 \setminus \mathbb{D}; \mathbb{R}^2) \geq \text{mod}_{d_\omega} \Gamma$$

for all $\varepsilon > 0$. Thus it is sufficient to show that $\text{mod}_{d_\omega} \Gamma > 0$.

Let ρ be an admissible function for Γ for the metric d_ω . For each $j \in \mathbb{N}$, let $m_j = \inf\{\int_\gamma \rho ds_\omega : \gamma \in \Gamma_j\}$. If $m_j > 0$, this implies that ρ/m_j is admissible for the

path family Γ_j , and thus that

$$(32) \quad \int_{A_j} \frac{\rho^2}{m_j^2} d\mathcal{H}_\omega^2 \geq \text{mod } \Gamma_j = \frac{\log(\min\{t_j, 1-t_j\}/b_j)}{\pi} \geq \frac{\log(a_j/b_j)}{\pi}.$$

Let γ_j be a path in Γ_j such that $\int_{\gamma_j} \rho ds_\omega \leq \max\{2m_j, 2^{-j-1}\}$. We can define a path $\gamma: [0, 1] \rightarrow \mathbb{R}^2$ by

$$\gamma(t) = (t, \sup_{j \in \mathbb{N}} \pi_2(|\gamma_j| \cap (\{t\} \times \mathbb{R}))).$$

Here, π_2 denotes projection onto the vertical axis. We see that the path γ is d_ω -rectifiable as follows. For every $t \in [0, 1]$ except on a countable set, there is $\varepsilon > 0$ such that $\gamma((t-\varepsilon, t+\varepsilon))$ is a subpath of a single path γ_j . At such points t , γ is locally rectifiable. Otherwise, there is $\varepsilon > 0$ such that the image of $\gamma((t-\varepsilon, t+\varepsilon))$ is contained in the union of the image of two paths γ_j, γ_k . Again, it follows that γ is locally rectifiable at these points. We conclude that γ itself is rectifiable. Observe that

$$1 \leq \int_\gamma \rho ds \leq \sum_{j=1}^{\infty} \int_{\gamma_j} \rho ds \leq \sum_{j=1}^{\infty} \max\{2m_j, 2^{-j-1}\}.$$

This implies that the relationship $m_j \geq 1/(2 \cdot 2^j)$ must hold for some $j \in \mathbb{N}$. This together with (32) gives

$$\begin{aligned} \frac{1}{2 \cdot 2^j} \leq m_j &\leq \left(\frac{\pi}{\log(a_j/b_j)} \right)^{1/2} \left(\int_{\mathbb{R}^2} \rho^2 d\mathcal{H}_{\|\cdot\|_2}^2 \right)^{1/2} \\ &= \left(\frac{\pi}{4^j} \right)^{1/2} \left(\int_{\mathbb{R}^2} \rho^2 d\mathcal{H}_{\|\cdot\|_2}^2 \right)^{1/2}. \end{aligned}$$

This yields the lower bound

$$\frac{1}{4\pi} \leq \int_{\mathbb{R}^n} \rho^2 d\mathcal{H}^2.$$

We conclude that (\mathbb{R}^2, d_ω) is not reciprocal.

Example 6.2. We construct a lower semicontinuous weight $\omega: \mathbb{R}^2 \rightarrow [0, \infty]$ that vanishes on a Cantor set $E \subset [0, 1] \times \{0\}$ of positive length such that the space (\mathbb{R}^2, d_ω) is reciprocal.

As before, identify $[0, 1]$ with the set $[0, 1] \times \{0\} \subset \mathbb{R}^2$. Consider the quadrilateral $Q = [0, 1] \times [-1, 1]$. Let Γ be the family of paths in Q connecting the left and right edges of Q .

Fix for the time being a value $t \in (0, 1/2)$. Let $I = [t, 1-t] \subset (0, 1)$ and let $\omega_1 = \chi_{\mathbb{R}^2 \setminus I}$, noting that ω_1 vanishes on the set I . Let \mathcal{E}_1 denote the decomposition of \mathbb{R}^2 corresponding to I . The weight ω_1 determines a metric \tilde{d}_{ω_1} on $\mathbb{R}^2/\mathcal{E}_1$ that is not reciprocal. Let π_{ω_1} denote the associated quotient map. Note that the metric \tilde{d}_{ω_1} , like all other metrics in this example, agrees with the Euclidean metric locally outside of $\pi_{\omega_1}(I)$. Thus the Hausdorff 2-measure relative to the metric \tilde{d}_{ω_1} coincides with Lebesgue measure.

Let $\tilde{\rho}$ be an admissible function for $\pi_{\omega_1}\Gamma$ with respect to the metric \tilde{d}_{ω_1} satisfying $\int \tilde{\rho}^2 d\mathcal{H}_{\tilde{d}_{\omega_1}}^2 \leq 2 \text{ mod } \pi_{\omega_1}\Gamma$. Since the function

$$\tilde{g} = \frac{\chi_{[0,t] \times [-1,1]} + \chi_{(1-t,1] \times [-1,1]}}{2t}$$

is admissible for $\pi_{\omega_1}\Gamma$, it follows that

$$(33) \quad \int \tilde{\rho}^2 d\mathcal{H}_{\tilde{d}_{\omega_1}}^2 \leq 2 \int \tilde{g}^2 d\mathcal{H}_{\tilde{d}_{\omega_1}}^2 = \frac{2}{t}.$$

Let $\rho = \chi_Q + \tilde{\rho} \circ \pi_{\omega_1}$.

For all $n \in \mathbb{N}$, $i \in \{1, \dots, n\}$, let φ_i^n denote the similarity mapping of \mathbb{R}^2 taking I to the interval $[(i-1)/n, i/n]$. Explicitly, $\varphi_i^n(x) = x/n + ((i-1)/n, 0)$. Let $I_n = \bigcup_{i=1}^n \varphi_i^n(I)$ and let \mathcal{E}_n denote the corresponding decomposition of \mathbb{R}^2 . Let $\omega_n = \chi_{\mathbb{R}^2 \setminus I_n}$, and \tilde{d}_{ω_n} the resulting metric on $\mathbb{R}^2/\mathcal{E}_n$.

Let $\rho_i^n = \rho \circ (\varphi_i^n)^{-1}$. Define now the function $\rho_n: Q \rightarrow [0, \infty]$ by

$$\rho_n(x) = \begin{cases} \rho_i^n(x) & \text{if } x \in \varphi_i^n((0, 1) \times [-1, 1]) \text{ for some } i \in \{1, \dots, n\} \\ 1 & \text{otherwise} \end{cases}.$$

For all $x \in \pi_{\omega_n}(Q)$, we define $\tilde{\rho}_n(x) = \rho_n(\pi_{\omega_n}^{-1}(x))$. We claim that $\tilde{\rho}_n$ is admissible for $\pi_{\omega_n}\Gamma$ with respect to the metric \tilde{d}_{ω_n} .

Let $Q_i^n = [(i-1)/n, i/n] \times [-1, 1]$, and let γ_i^n be a subpath of γ that traverses Q_i^n horizontally. It suffices to show that $\int_{\gamma_i^n} \rho_n \omega_n ds_{\|\cdot\|_2} \geq 1/n$. If γ_i^n does not intersect I_n , then this is clear since $\rho_i^n \geq 1$ on $Q_i^n \setminus I_n$. If γ_i^n is contained in $\varphi_i^n(Q)$, then this is also immediate by the admissibility of $\tilde{\rho}$. Finally, if γ_i^n intersects both I_n and $Q_i^n \setminus \varphi_i^n(Q)$, then γ_i^n must traverse a vertical distance of $1/n$, and again the conclusion follows. We conclude that ρ_n is admissible for Γ .

Next, we have the upper bound

$$(34) \quad \int_Q \rho_n^2 d\mathcal{L}^2 \leq \int_Q 1 d\mathcal{L}^2 + \sum_{i=1}^n \int_{Q_i^n} (\rho_i^n)^2 d\mathcal{L}^2 \leq 2 + \frac{\|\rho\|_{L^2(Q)}^2}{n}.$$

Observe that $2 = \text{mod}_{\|\cdot\|_2} \Gamma$. Thus, by taking n to be sufficiently large, the modulus of Γ with respect to \tilde{d}_{ω_n} becomes arbitrarily close to the Euclidean modulus.

We can now define the Cantor set E as follows. For a given $t \in (0, 1/2)$ and $n \in \mathbb{N}$, let $I(t)$, $I_n(t)$ and $\omega_n(t)$ denote respectively the sets I and I_n and the weight ω_n constructed above. For all $j \in \mathbb{N}$, let $t_j = 2^{-j-2}$, observing that $\mathcal{L}^1(I(t_j)) = 1 - 2t_j$. Let $\tilde{\omega}_j = \omega_{n_j}(t_j)$. By choosing n_j sufficiently large, we can guarantee that

$$\mathcal{L}^1(I_{n_j}(t_j) \cap I_{n_{j-1}}(t_{j-1})) \geq (1 - 4t_j)\mathcal{L}^1(I_{n_{j-1}}(t_{j-1}))$$

and that $\text{mod}_{\tilde{\omega}_j} \Gamma \leq 2 + 1/j$ by applying (33) and (34). Inductively choosing n_j in this manner, we have

$$\mathcal{L}^1\left(\bigcap_{i=1}^j I_{n_i}(t_i)\right) \geq \prod_{i=1}^j (1 - 4t_i) = \prod_{i=1}^j (1 - 2^{-i}).$$

Let $E = \bigcap_{j=1}^{\infty} I_{n_j}(t_j)$ and let $\omega = \chi_{\mathbb{R}^2 \setminus E}$. Then $\mathcal{L}^1(E) = \prod_{j=1}^{\infty} (1 - 2^{-j}) > 0$. Moreover, $\omega \geq \tilde{\omega}_j$ for all $j \in \mathbb{N}$. This fact, combined with Theorem 2.6, yields that $2 \leq \text{mod}_{\pi_{\omega}} \Gamma \leq \text{mod}_{\pi_{\tilde{\omega}_j}} \Gamma$ for all $j \in \mathbb{N}$. We conclude that $\text{mod}_{\pi_{\omega}} \Gamma = 2 = \text{mod}_{\|\cdot\|_2} \Gamma$.

Let Γ^* denote the family of paths connecting the bottom and top edges of Q . It is clear that the function $\rho^* = 1/2\chi_Q$ is admissible for Γ^* with respect to the metric \tilde{d}_{ω} . Thus $\text{mod}_{\tilde{d}_{\omega}} \Gamma^* = 1/2 = \text{mod}_{\|\cdot\|_2} \Gamma^*$. By Proposition 4.3, this suffices to show that \tilde{d}_{ω} is reciprocal.

7. FACTORING QUASICONFORMAL MAPPINGS

The goal of this section is to prove Proposition 1.5 and Theorem 1.6. To prepare for this, we first give in Section 7.1 an overview of isothermal quasiconformal mappings. See [Iko19] for a more complete treatment. Section 7.2 gives the proof of Proposition 1.5. This is followed by a discussion in Section 7.3 of the problem of optimizing the distortion constant in Proposition 1.5. Finally, in Section 7.4, we prove Theorem 1.6.

7.1. Isothermal Parametrizations. Let X be a quasiconformal surface. By Theorem 6.2 in [Iko19], there exists a complete Riemannian surface Y of constant curvature and a quasiconformal map

$$\psi: Y \rightarrow X$$

with minimal pointwise distortion at almost every point: for every other Riemannian surface Z and quasiconformal map $\varphi: Z \rightarrow X$, the inequality

$$(35) \quad (g_\psi(g_{\psi^{-1}} \circ \psi)) \circ (\psi^{-1} \circ \varphi) \leq g_\varphi(g_{\varphi^{-1}} \circ \varphi)$$

holds \mathcal{H}_Z^2 -almost everywhere on Z . Recall that, for example, g_ψ and $g_{\psi^{-1}}$ refer to minimal weak upper gradients of ψ and ψ^{-1} , respectively. In this case, we say that (Y, ψ) is an *isothermal parametrization* of X . By Corollary 4.7 of [Iko19], any isothermal parametrization ψ is quasiconformal with outer dilatation $K_O(\psi)$ at most $4/\pi$ and inner dilatation $K_I(\psi)$ at most $\pi/2$. Also, the pointwise distortion of ψ is bounded from above by $\sqrt{2}$ \mathcal{H}_Y^2 -almost everywhere.

We elaborate on the meaning of (35) in the case when $X = (\mathbb{R}^2, d_N)$ for some norm N . Then we can take $Y = \mathbb{R}^2$ and ψ to be a linear map

$$\psi: \mathbb{R}^2 \rightarrow (\mathbb{R}^2, d_N)$$

such that $g_\psi = L(N \circ \psi)$ and $g_{\psi^{-1}} = \omega(N \circ \psi)^{-1}$. Recall that L and ω denote, respectively, the maximal and minimal stretching, defined in (6) and (7).

The inequality (35) implies that, for all other linear maps $\varphi: \mathbb{R}^2 \rightarrow (\mathbb{R}^2, N)$, we have

$$(36) \quad \frac{L(N \circ \psi)}{\omega(N \circ \psi)} \leq \frac{L(N \circ \varphi)}{\omega(N \circ \varphi)}.$$

In terms of the *distortion* of a norm defined in (8), the inequality (36) implies that $N \circ \psi$ has the smallest possible distortion among such pairs ψ and φ . This can be phrased in terms of the *Banach–Mazur distance* in convex geometry; see [Rom19] and [Iko19, Section 4].

An isothermal parametrization of a quasiconformal surface is essentially unique. This is also part of the content of Theorem 6.2 of [Iko19], partially quoted here.

Theorem 7.1 ([Iko19]). Let $\psi: Y \rightarrow X$ be an isothermal parametrization of X , and $\varphi: Z \rightarrow X$ a quasiconformal map from a Riemannian surface Z onto X . Then φ is isothermal if and only if $\psi^{-1} \circ \varphi$ is a conformal diffeomorphism.

Let N be an admissible reciprocal norm field on \mathbb{R}^2 that vanishes on the compact set $E \subset \mathbb{R}^2$. The following lemma is a consequence of Theorem 4.16 of [Iko19].

Lemma 7.2. The identity map $\iota: \mathbb{R}^2 \rightarrow (\mathbb{R}^2, d_N)$ is isothermal if and only if

$$(37) \quad \frac{L(N_x)}{\omega(N_x)} \leq \frac{L(N_x \circ \varphi)}{\omega(N_x \circ \varphi)}$$

for all $\varphi \in \text{GL}_2$, for \mathcal{L}^2 -almost every $x \in \mathbb{R}^2$.

Observe that (37) is satisfied by the norm $N_x = \|\cdot\|_\infty$, and more generally by any norm N_x whose unit ball is a square. Thus Lemma 7.2 has the following corollary.

Corollary 7.3. Suppose that N is reciprocal and that the unit ball of N_x is a square for \mathcal{L}^2 -almost every $x \in \mathbb{R}^2$. Then the identity map $\iota: \mathbb{R}^2 \rightarrow (\mathbb{R}^2, d_N)$ is isothermal.

7.2. Proof of Proposition 1.5. Recall that we are assuming that N is a reciprocal norm field such that $\pi_N: \Omega \rightarrow (\Omega, d_N)$ is isothermal, and that N is continuous outside the set $E = \{x \in \Omega : N_x = 0\}$.

Let G be a complete Riemannian norm field on Ω of constant Gaussian curvature -1 or 0 , which exists by the classical uniformization theorem. The field is of the form $G = \omega \|\cdot\|_2$ for some smooth ω . Consider the norm field

$$M = \chi_{\Omega \setminus E} \frac{\omega}{\omega(N)} N + \chi_E G.$$

The function $1/\omega(N)$ is continuous in $\Omega \setminus E$ due to the continuity of N outside E . Then the distortion bound on N implies that M is a lower semicontinuous norm field satisfying $G \leq M \leq HG$ everywhere.

Let $\hat{d} = d_M$ denote the distance induced by M . Then

$$d_G \leq \hat{d} \leq Hd_G,$$

so the identity map $P = \pi_M: (\Omega, d_G) \rightarrow (\Omega, \hat{d})$ satisfies (3) and in particular is H -bi-Lipschitz. Lemma 3.5 states that its metric differential coincides with M \mathcal{L}^2 -almost everywhere.

The proof is complete after we show that $\hat{\iota} = \pi_N \circ P^{-1}$ is 1-quasiconformal. Recall that the metric derivatives of π_N and P coincide with N and M , respectively. Then, as a consequence of Corollary 5.15 and Proposition 5.12 of [Iko19], the 1-quasiconformality is equivalent to proving that for \mathcal{L}^2 -almost every $x \in \Omega$, the distortion of the identity map from (\mathbb{R}^2, M_x) to (\mathbb{R}^2, N_x) equals one \mathcal{L}^2 -almost everywhere.

Observe that, by the change of variables formula Lemma 3.6 and the Lusin's Condition (N^{-1}) of π_N , the set E has zero \mathcal{L}^2 -measure, so we only need to check the pointwise distortion in the complement of E . Here the claim is immediate, since $M_z = \omega(z)N_z/\omega(N_z)$ for every $z \in \Omega \setminus E$. We conclude that $\hat{\iota}$ is 1-quasiconformal.

7.3. Remarks on optimal distortion. We discuss the question of when the optimal constant $H = \sqrt{2}$ in (3) in Proposition 1.5 can be achieved. We recall that any planar quasiconformal mapping $f: \Omega \rightarrow \hat{\Omega}$ is a solution of the *Beltrami equation* $f_{\bar{z}} = \mu f_z$, where $\mu: \Omega \rightarrow \mathbb{C}$ is a measurable function satisfying $\|\mu\|_\infty < 1$. Conversely, the *measurable Riemann mapping theorem* provides a homeomorphic solution to the Beltrami equation for any such μ . The function μ is called the *Beltrami coefficient*. Geometrically, the choice of a Beltrami coefficient corresponds to the choice of a measurable ellipse field on Ω modulo rescaling of the ellipses. See Chapter 5 of [AIM09] for an in-depth overview of the topic.

Given a norm field \hat{N} on a domain $\hat{\Omega} \subset \mathbb{R}^2$, one obtains an ellipse field on $\hat{\Omega}$ by associating to each norm \hat{N}_x its distance ellipse, that is, the unique ellipse $\mathcal{E} \subset B_{\hat{N}_x}(0, 1)$ having minimal $\lambda \geq 1$ such that $B_{\hat{N}_x}(0, 1) \subset \lambda \mathcal{E}$. This in turn gives a Beltrami coefficient $\mu_{\hat{N}}$ corresponding to \hat{N} . We refer the reader to [Iko19, Sec. 4] for more details.

This choice of ellipse field also determines an underlying Riemannian structure on the metric space $(\hat{\Omega}, d_{\hat{N}})$. A consequence of the classical slit domain uniformization theorem [AS60, Chapter III, Section 4] and [Iko19, Corollary 6.3] is the existence of a domain $\Omega \subset \mathbb{R}^2$ and a locally quasiconformal map $\psi: \Omega \rightarrow \hat{\Omega}$ such that $\hat{f} = \pi_{\hat{N}} \circ \psi$ is isothermal. Consider the distance $d(x, y) = d_{\hat{N}}(\hat{f}(x), \hat{f}(y))$ on Ω and the norm field $N = \hat{N} \circ D\psi$. Then the identity map $\iota: \Omega \rightarrow (\Omega, d)$ is isothermal and the metric differential of ι exists and equals N \mathcal{L}^2 -almost everywhere. If the norm field N obtained in this manner is continuous and non-zero outside $E =$

$\psi^{-1}(\{x \in \widehat{\Omega} : \widehat{N}_x = 0\})$, then Proposition 1.5 now holds with constant $H = \sqrt{2}$ for the space (Ω, d) and hence the original space $(\widehat{\Omega}, d_{\widehat{N}})$ as well.

The question of when the norm field N is continuous, in turn, depends upon the regularity of the map ψ . In fact, if ψ is \mathcal{C}^1 -continuous in Ω , then N is continuous and non-zero outside E . Since the map ψ arises as a solution to the Beltrami equation, this leads to the question of regularity of solutions to the Beltrami equation. Indeed, if we consider a domain U compactly contained in $\widehat{\Omega}$, the restriction of ψ^{-1} to U solves the Beltrami equation induced by $\mu_{\widehat{N}}|_U$. The \mathcal{C}^1 -continuity of ψ in U is known to hold, for example, when $\mu_{\widehat{N}}|_U$ is \mathcal{C}^1 -continuous, locally Hölder continuous [AIM09, Theorem 15.0.7] or in $W_{\text{loc}}^{1,p}(U)$ for a large enough $p > 1$ depending on the L^∞ -norm of $\mu_{\widehat{N}}|_U$ [BCO19, Proposition 4].

Solutions of the Beltrami equation for $\mu_{\widehat{N}}$, even when \widehat{N} is a continuous Riemannian norm field, need not always be \mathcal{C}^1 -continuous. In the following, we use complex notation $z = z_1 + iz_2$ to denote the point $(z_1, z_2) \in \mathbb{R}^2$ and $\bar{z} = z_1 - iz_2$ to denote the complex conjugate of z . See Section 2.4 of [AIM09] for a brief overview of complex notation. The following example is based on Section 15.1 of [AIM09]. Let

$$\mu(z) = \frac{z}{\bar{z}(1 + \log \|z\|_2^2)}$$

and consider the continuous Riemannian norm field \widehat{N} on $\widehat{\Omega} = B_{\|\cdot\|_2}(0, e^{-1/2})$ defined by $\widehat{N}_z(v) = \|v + \mu(z)\bar{v}\|_2$. Then $\mu(z) = \mu_{\widehat{N}_z}$, where $\mu_{\widehat{N}_z}$ is the Beltrami coefficient corresponding to the \widehat{N} as described earlier in this remark. Even though \widehat{N} is continuous, every solution for the Beltrami equation for $\mu_{\widehat{N}_z} = \mu(z)$ has a discontinuous derivative at the origin. This is seen by considering the particular solution $g(z) = -z \log \|z\|_2^2$ and noticing that the differential Dg is discontinuous at the origin. It is enough to check this property for g since, by the Stoilow factorization theorem [AIM09, Theorem 5.5.1], every other quasiconformal solution is of the form $\Psi \circ g$ for some conformal diffeomorphism Ψ .

For more general norm fields, we have the additional complexity that $\mu_{\widehat{N}}$ can be smooth even though \widehat{N} is not. For example, consider the continuous norm field \widehat{N} defined by $\widehat{N}_z(v) = \|e^{i\|z\|_2} v\|_\infty$. Since the supremum norm is not \mathcal{C}^1 -continuous in $\mathbb{R}^2 \setminus \{0\}$ we see that \widehat{N} is not \mathcal{C}^1 -continuous, for example by considering the basepoint $z = \pi/4$ and the vector $v = 1$, even though $\mu_{\widehat{N}} = 0$. The identity $\mu_{\widehat{N}} = 0$ follows from Corollary 7.3.

7.4. Proof of Theorem 1.6. In this section, we present the construction used to prove Theorem 1.6, namely of a quasiconformal surface whose isothermal parametrization cannot be factored as a bi-Lipschitz mapping postcomposed with a quasiconformal mapping of smaller distortion. We begin by introducing the notation and parameters involved in Section 7.4.1. We develop various properties of this construction in the following subsections, culminating with the proof of Theorem 1.6 in Section 7.4.6.

7.4.1. Notation. Let us introduce the notation used in our construction. Our first task is to construct a sequence of nested Cantor sets, denoted by K_1, K_2, \dots and satisfying $K_1 \supset K_2 \supset \dots$. There are two intermediate steps used in our construction of the sets K_i . First, we define sets E_i^j for all $i, j \in \mathbb{N}$, $j \geq i$, to serve as base collections of squares from which the Cantor sets are extracted. The sets E_i^j are each the union of a collection of congruent closed squares $Q_i^j(k, l)$ that covers almost all of $[0, 1]^2$. The main feature of our construction is that the squares $Q_i^j(k, l)$ have




FIGURE 1. A square $Q_i^j(k, l) \subset E_i^j$ for i odd and the intersection $Q_i^j(k, l) \cap E_{i+1}^j$, shaded gray. The large outer square is $I_i^j(k) \times J_i^j(l)$.

the standard non-rotated alignment for odd values of i , while the square $Q_i^j(k, l)$ are aligned diagonally for even values of i .

In the second intermediate step, we define inductively

$$F_i^j = E_i^j \cap F_{i-1}^j \cap F_i^{j-1} \cap [0, 1]^2$$

for all $i, j \in \mathbb{N}$, $j \geq i$, with the convention that $F_0^j = F_i^{i-1} = [0, 1]^2$ for all i, j . By taking $F_i = \bigcap_j F_i^j$, we obtain a collection of nested Cantor sets. However, for our construction to work, we need the further property that the intersection of the Cantor sets is small. For this reason, we later define K_i to be a subset of F_i with the property that $\text{diam } K_i \rightarrow 0$ as $i \rightarrow \infty$.


We take a moment to fix some additional notation. In the following, let $I = J = [0, 1]$ and let $Q = I \times J = [0, 1]^2$. We identify I with the set $[0, 1] \times \{0\}$ and J with the set $\{0\} \times [0, 1]$. Let π_1 denote the standard projection map from \mathbb{R}^2 onto the first coordinate axis, and let π_2 denote the standard projection map from \mathbb{R}^2 onto the second coordinate axis.

As mentioned above, the even-numbered Cantor sets are formed from squares that are rotated by $\pi/4$ from the standard alignment. Let Q^* denote the square with vertices $(1/2, -1/2)$, $(3/2, 1/2)$, $(1/2, 3/2)$, and $(-1/2, 1/2)$. Let $I^* = J^* = [0, \sqrt{2}]$. We also identify I^* with the set $[0, \sqrt{2}] \times \{0\}$ and J^* with the set $\{0\} \times [0, \sqrt{2}]$. Let $\varphi: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be the orientation-preserving isometry that maps $[0, \sqrt{2}]^2$ onto Q^* and satisfies $\varphi(0, 0) = (1/2, -1/2)$. Explicitly,

$$\varphi(x, y) = (1/2, -1/2) + \frac{1}{\sqrt{2}}(x - y, x + y).$$

Thus $\varphi(I^* \times J^*) = Q^*$. Next, let π_1^* denote the projection map from Q^* onto $\varphi(I^*)$, and let π_2^* denote the projection map from Q^* onto $\varphi(J^*)$. Explicitly, $\pi_1^*(x, y) = \varphi(\pi_1(\varphi^{-1}(x, y)), 0)$ and $\pi_2^*(x, y) = \varphi(0, \pi_2(\varphi^{-1}(x, y)))$.

The definition of the sets E_i^j involves three sets of parameters: $\varepsilon_i^j > 0$, $N_i^j \in \mathbb{N}$, and $a_i^j \in \mathbb{N}$. A short explanation of these parameters is the following. The first parameter ε_i^j gives an upper bound on the proportion of area lost when passing from one step of the construction to the next. The second parameter N_i^j gives the number of subdivisions of the initial interval I or I^* that are made when forming


 FIGURE 2. The sets E_i^j as ordered by \preceq .

the squares that comprise E_i^j . The final parameter a_i^j corresponds to the side length of these squares. The precise relation is that the side length of a square in E_i^j is $(1 - 2(a_i^j)^{-1})/N_i^j$ for i odd and $\sqrt{2}(1 - 2(a_i^j)^{-1})/N_i^j$ for i even.

7.4.2. *Constructing the sets E_i^j .* For two pairs of indices (i, j) and (i', j') , we say that $(i, j) \preceq (i', j')$ if $j < j'$ or if $j = j'$ and $i \leq i'$. The relation \preceq gives an ordering on the set of indices (i, j) . We consider the sets E_i^j as being traversed in this order. We also write $(i, j) \prec (i', j')$ if $j < j'$ or if $j = j'$ and $i < i'$. Recall that here and throughout this proof we consider only those indices $i, j \in \mathbb{N}$ for which $j \geq i$. This ordering is illustrated in Figure 2.

We first choose the parameters $\varepsilon_i^j > 0$ so that they satisfy $\prod_{i,j} (1 - \varepsilon_i^j) \geq 1/2$. The factors in the product are traversed according to the ordering on $\{(i, j)\}$ just defined.

The sets E_i^j are defined for all $i, j \in \mathbb{N}$ satisfying $j \geq i$ in the following way. Assume for the moment that we have made suitable choices of $N_i^j, a_i^j \in \mathbb{N}$. In the case that i is odd, we divide I into N_i^j equal subintervals $I_i^j(k) = [(k-1)/N_i^j, k/N_i^j]$ and J into N_i^j equal subintervals $J_i^j(l) = [(l-1)/N_i^j, l/N_i^j]$. If i is even, we divide I^* into N_i^j equal subintervals $I_i^j(k) = [\sqrt{2}(k-1)/N_i^j, \sqrt{2}k/N_i^j]$ and J^* into N_i^j equal subintervals $J_i^j(l) = [\sqrt{2}(l-1)/N_i^j, \sqrt{2}l/N_i^j]$. This yields a collection of squares $I_i^j(k) \times J_i^j(l)$, where $k, l \in \{1, \dots, N_i^j\}$. If i is odd, let $Q_i^j(k, l)$ be the square of side length $(1 - 2(a_i^j)^{-1})/N_i^j$ with the same center and alignment as $I_i^j(k) \times J_i^j(l)$. If i is even, define $Q_i^j(k, l)$ to be the square of side length $\sqrt{2}(1 - 2(a_i^j)^{-1})/N_i^j$ with the same center and alignment as $\varphi(I_i^j(k) \times J_i^j(l))$. In this case, the square $Q_i^j(k, l)$ is contained in Q^* and is aligned diagonally. Let

$$E_i^j = \bigcup_{k,l} Q_i^j(k, l).$$

The square $Q_i^j(k, l)$, for i odd, is given explicitly by

$$\left[\left(k - 1 + \frac{1}{a_i^j} \right) \frac{1}{N_i^j}, \left(k - \frac{1}{a_i^j} \right) \frac{1}{N_i^j} \right] \times \left[\left(l - 1 + \frac{1}{a_i^j} \right) \frac{1}{N_i^j}, \left(l - \frac{1}{a_i^j} \right) \frac{1}{N_i^j} \right].$$

Let $v_i^j(k, l, 1), v_i^j(k, l, 2), v_i^j(k, l, 3), v_i^j(k, l, 4)$ denote the four vertices of $Q_i^j(k, l)$, traversed counterclockwise from the bottom left. Let $w_i^j(k, l)$ denote the center point of $Q_i^j(k, l)$.

Similarly, the square $Q_i^j(k, l)$, for i even, is the image under φ of the square

$$(38) \quad \left[\left(k - 1 + \frac{1}{a_i^j} \right) \frac{\sqrt{2}}{N_i^j}, \left(k - \frac{1}{a_i^j} \right) \frac{\sqrt{2}}{N_i^j} \right] \times \left[\left(l - 1 + \frac{1}{a_i^j} \right) \frac{\sqrt{2}}{N_i^j}, \left(l - \frac{1}{a_i^j} \right) \frac{\sqrt{2}}{N_i^j} \right].$$

Let $v_i^j(k, l, 1), v_i^j(k, l, 2), v_i^j(k, l, 3), v_i^j(k, l, 4)$ denote the four vertices of $Q_i^j(k, l)$, where $v_i^j(k, l, 1)$ is the image under φ of the bottom left vertex of the square in

(38) and the rest are labelled in counterclockwise order. Let $w_i^j(k, l)$ denote the center point of $Q_i^j(k, l)$.

The values of N_i^j and a_i^j are chosen inductively using the ordering \preceq . Let $N_1^1 = 2$ and choose $a_1^1 \in \mathbb{N}$ so that $\mathcal{L}^2(E_1^1) \geq 1 - \varepsilon_1^1$. For the inductive step, assume that we have chosen $N_{i'}^{j'}$ and $a_{i'}^{j'}$ for some pair (i', j') , and that

$$\mathcal{L}^2 \left(\bigcap_{(i'', j'') \preceq (i', j')} E_{i''}^{j''} \right) \geq \prod_{(i'', j'') \preceq (i', j')} (1 - \varepsilon_{i''}^{j''}).$$

Let (i, j) denote the pair immediately succeeding (i', j') .

Define now $N_i^j = 2a_{i'}^{j'} N_{i'}^{j'}$. We then choose a_i^j so that

$$\mathcal{L}^2 \left(\bigcap_{(i'', j'') \preceq (i, j)} E_{i''}^{j''} \right) \geq \prod_{(i'', j'') \preceq (i, j)} (1 - \varepsilon_{i''}^{j''}).$$

This can be done because E_i^j can be made to have arbitrarily large area in, respectively, Q or Q^* by making a_i^j sufficiently large.

We make the following observation. Fix (i, j) and consider a square $Q_i^j(k, l)$. For all (i', j') such that $(i, j) \prec (i', j')$ and $m, n \in \{1, \dots, N_{i'}^{j'}\}$, the square $I_{i'}^{j'}(m) \times J_{i'}^{j'}(n)$, if i' is odd, or $\varphi(I_{i'}^{j'}(m) \times J_{i'}^{j'}(n))$, if i' is even, is either entirely contained in $Q_i^j(k, l)$, has interior disjoint from $Q_i^j(k, l)$, or intersects $Q_i^j(k, l)$ in a triangle whose vertices are three of the vertices of $I_{i'}^{j'}(m) \times J_{i'}^{j'}(n)$.

We also observe a uniformity to how the squares are distributed. For each i, j, k, l , we divide the square $Q_i^j(k, l)$ into four triangles whose vertices are two adjacent vertices of $Q_i^j(k, l)$ and the midpoint of $Q_i^j(k, l)$. Denote these by $T_i^j(k, l, 1)$, $T_i^j(k, l, 2)$, $T_i^j(k, l, 3)$, $T_i^j(k, l, 4)$, where $T_i^j(k, l, m)$ contains the edge $[v_i^j(k, l, m), v_i^j(k, l, m+1)]$, taking $v_i^j(k, l, 5) = v_i^j(k, l, 1)$.

Lemma 7.4. Let $i, j, i', j' \in \mathbb{N}$, where $(i, j) \prec (i', j')$. For all $k, l \in \{1, \dots, N_i^j\}$ and $m \in \{1, \dots, 4\}$ satisfying $T_i^j(k, l, m) \subset Q$, the sets $T_i^j(k, l, m) \cap E_{i'}^{j'}$ are all congruent.

Proof. This proof depends on the property that $2a_i^j N_i^j$ divides $N_{i'}^{j'}$. As a result, squares at different levels of the construction intersect nicely. We consider the case when i is odd.

First, suppose that i' is also odd. For each $m \in \{1, \dots, 4\}$, consider the edge $e_i^j(k, l, m)$ as defined above. We have $\pi_1(v_i^j(k, l, 1)) = \pi_1(v_i^j(k, l, 4)) = k_1(k, l)/N_{i'}^{j'}$ and $\pi_1(v_i^j(k, l, 3)) = \pi_1(v_i^j(k, l, 2)) = k_3(k, l)/N_{i'}^{j'}$, where

$$k_1(k, l) = \frac{(a_i^j k - a_i^j + 1)N_{i'}^{j'}}{a_i^j N_i^j} \quad \text{and} \quad k_3(k, l) = \frac{(a_i^j k - 1)N_{i'}^{j'}}{a_i^j N_i^j}.$$

Similarly, we have $\pi_2(v_i^j(k, l, 2)) = \pi_2(v_i^j(k, l, 1)) = k_2(k, l)/N_{i'}^{j'}$ and $\pi_4(v_i^j(k, l, 3)) = \pi_4(v_i^j(k, l, 4)) = k_4(k, l)/N_{i'}^{j'}$, where

$$k_2(k, l) = \frac{(a_i^j l - a_i^j + 1)N_{i'}^{j'}}{a_i^j N_i^j} \quad \text{and} \quad k_4(k, l) = \frac{(a_i^j l - 1)N_{i'}^{j'}}{a_i^j N_i^j}.$$

Observe that $k_i(k, l) \in \mathbb{N}$ for all $i \in \{1, \dots, 4\}$. We have then

$$Q_i^j(k, l) = [k_1(k, l)/N_{i'}^{j'}, k_3(k, l)/N_{i'}^{j'}] \times [k_2(k, l)/N_{i'}^{j'}, k_4(k, l)/N_{i'}^{j'}].$$

We conclude from this that the intersection $Q_i^j(k, l) \cap E_{i'}^{j'}$ is precisely the union of the squares

$$\{Q_{i'}^{j'}(k', l') : k_2(k, l) + 1 \leq k' \leq k_4(k, l), k_1(k, l) + 1 \leq l' \leq k_3(k, l)\}.$$

We also observe that

$$|k_3(k, l) - k_1(k, l)| = |k_4(k, l) - k_2(k, l)| = \frac{(a_i^j - 2)N_{i'}^{j'}}{a_i^j N_i^j}.$$

Thus the sets $Q_i^j(k, l) \cap E_{i'}^{j'}$ are congruent for all $k, l \in \{1, \dots, N_i^j\}$. Moreover, notice that each set $Q_i^j(k, l) \cap E_{i'}^{j'}$ is invariant under rotations by $\pi/4$ about the center point $w_i^j(k, l)$. We conclude from this that the sets $T_i^j(k, l, m) \cap E_{i'}^{j'}$ are all congruent.

Next, suppose that i' is even. Consider now a triangle $T_i^j(k, l, m)$. The two shorter edges of $T_i^j(k, l, m)$ are the edges of a rectangle $R_i^j(k, l, m)$ of side length $\sqrt{2}(a_i^j - 2)/(a_i^j N_i^j)$. To keep the exposition more manageable, we write out the argument only for $T_i^j(k, l, 1)$. We compute

$$\begin{aligned} \varphi^{-1}(v_i^j(k, l, 1)) &= \left(\frac{(a_i^j k + a_i^j l - 2a_i^j + 2)\sqrt{2}}{2a_i^j N_i^j}, -\frac{(-a_i^j k + a_i^j l + a_i^j N_i^j)\sqrt{2}}{2a_i^j N_i^j} \right) \\ \varphi^{-1}(v_i^j(k, l, 2)) &= \left(\frac{(a_i^j k + a_i^j l - a_i^j)\sqrt{2}}{2a_i^j N_i^j}, \frac{(-a_i^j k + a_i^j l - a_i^j + 2 + a_i^j N_i^j)\sqrt{2}}{2a_i^j N_i^j} \right). \end{aligned}$$

Comparing this with (38) and using the property that $2a_i^j N_i^j$ divides $N_{i'}^{j'}$, we have

$$\varphi^{-1}(R_i^j(k, l, m)) = \left[\frac{k_1(k, l)\sqrt{2}}{N_{i'}^{j'}}, \frac{k_3(k, l)\sqrt{2}}{N_{i'}^{j'}} \right] \times \left[\frac{k_2(k, l)\sqrt{2}}{N_{i'}^{j'}}, \frac{k_4(k, l)\sqrt{2}}{N_{i'}^{j'}} \right]$$

for some $k_1(k, l), \dots, k_4(k, l) \in \mathbb{N}$ satisfying

$$|k_3(k, l) - k_1(k, l)| = |k_4(k, l) - k_2(k, l)| = \frac{(a_i^j - 2)N_{i'}^{j'}}{2a_i^j N_i^j}.$$

The intersection $R_i^j(k, l, 1) \cap E_{i'}^{j'}$ is the union of the squares

$$\{Q_{i'}^{j'}(k', l') : k_2(k, l) + 1 \leq k' \leq k_4(k, l), k_1(k, l) + 1 \leq l' \leq k_3(k, l)\}.$$

Thus the sets $R_i^j(k, l, 1) \cap E_{i'}^{j'}$ are all congruent, and by symmetry it follows that the sets $T_i^j(k, l, 1) \cap E_{i'}^{j'}$ are all congruent as well.

The case when i is even is similar and its proof is omitted. \square

7.4.3. Constructing the Cantor sets. For all i, j , let $F_0^j = Q$ and $F_i^{i-1} = Q$. Define now

$$F_i^j = E_i^j \cap F_{i-1}^j \cap F_i^{j-1}$$

for all $j \geq i$. Observe that $\bigcap_{(i'', j'') \leq (i, j)} E_{i''}^{j''} \subset F_i^j$, so we have $\mathcal{L}^2(F_i^j) \geq 1/2$ for all i, j . Next, let $F_i = \bigcap_{j \geq i} F_i^j$.

Let $K_0 = \mathbb{R}^2$. For each $i \geq 1$, pick inductively a square $Q_i = Q_i^i(j_i, k_i)$ with the property that $Q_i \subset Q_{i-1}$. Let

$$K_i = F_i \cap Q_i.$$

From Lemma 7.4, it follows that $\mathcal{L}^2(K_i) = \mathcal{L}^2(F_i)/(N_i^i)^2$. Moreover, we have that $\text{diam } K_i \rightarrow 0$ as $i \rightarrow \infty$, and in particular that $\bigcap_i K_i$ is a single point set.

7.4.4. *Dense networks of paths.* The following portion of the argument relates to having a “dense network of paths” at every stage.

We define the following subset of K_i . If i is even, let

$$H_i = K_i \cap \pi_2^{-1}(I \setminus \pi_2(K_{i+1})).$$

If i is odd, let

$$H_i = K_i \cap (\pi_2^*)^{-1}(I^* \setminus \pi_2^*(K_{i+1})).$$

For example, in Figure 1 representing the case where i is odd, a point $x \in K_i \cap Q_i^j(k, l)$ belongs to H_i if the line $t \mapsto x + (t, t)$ does not intersect any of the gray boxes.

Lemma 7.5. For every point $x \in K_{i+1}$ and $r > 0$, the set $H_i \cap B_{\|\cdot\|_2}(x, r)$ has positive \mathcal{L}^2 -measure.

Proof. In the first case, we assume that i is even, and hence that $i + 1$ is odd. Let $x \in K_{i+1}$ and $r > 0$. Consider a square $Q_{i+1}^j(k, l)$ containing x for some j sufficiently large so that $Q_{i+1}^j(k, l) \subset B(x, r/3)$ and such that $Q_{i+1}^j(k, l) \subset Q_{i+1}$. Pick a horizontal edge S of $Q_{i+1}^j(k, l)$ whose interior is contained in $\text{int}(Q_{i+1})$.

Consider now the set E_i^{j+1} . Subdivide S into $(a_{i+1}^j - 2)N_i^{j+1}/(a_{i+1}^j N_{i+1}^j)$ congruent subintervals. Each subinterval is the diagonal of a square $\varphi(I_i^{j+1}(k') \times J_i^{j+1}(l'))$, with corresponding square $Q_i^{j+1}(k', l') \subset E_i^{j+1}$. From such a square $Q_i^{j+1}(k', l')$, we may extract a triangle $T_i^{j+1}(k', l', m')$, as defined prior to the statement of Lemma 7.4, whose interior does not intersect E_{i+1}^j . Observe further that $T_i^{j+1}(k', l', m') \subset Q_i$, so that

$$T_i^{j+1}(k', l', m') \cap F_i = T_i^{j+1}(k', l', m') \cap K_i.$$

As a consequence of Lemma 7.4, we have that

$$\mathcal{L}^2(T_i^{j+1}(k', l', m') \cap F_i) = \frac{\mathcal{L}^2(F_i)}{4(N_i^{j+1})^2}.$$

Moreover, $T_i^{j+1}(k', l', m')$ lies in the neighborhood of $Q_{i+1}^j(k, l)$ of radius

$$1/N_i^{j+1} \leq \text{diam } Q_{i+1}^j(k, l) < 2r/3,$$

so $T_i^{j+1}(k', l', m') \subset B_{\|\cdot\|_2}(x, r)$. Also, we have that

$$\text{int}(T_i^{j+1}(k', l', m')) \cap F_i \subset H_i.$$

This verifies the claim.

The case that i is even and $i + 1$ is odd is similar, and we omit the details. \square

Lemma 7.6. Let $x \in \pi_1(K_i)$ be a Lebesgue density point of $\pi_1(K_i)$, where $i \in \mathbb{N}$ is odd. Let $\delta > 0$, and let $t_0 > 0$ be such that

$$\frac{\mathcal{L}^1(\pi_1(K_i) \cap (x, x + t))}{t} \geq 1 - \delta$$

for all $t \in (0, t_0)$. Then for all $y \in \pi_1(K_i)$ satisfying $|y - x| < 2\delta t$,

$$\frac{\mathcal{L}^1(\pi_1(K_i) \cap (y, y + t))}{t} \geq 1 - 2\delta.$$

The same result holds with π_2 instead of π_1 . If i is even, the corresponding result holds for π_1^* and π_2^* , identifying I^* and J^* with the interval $[0, \sqrt{2}]$.

Proof. The first claim follows from the relationship

$$\mathcal{L}^1(\pi_1(K_i) \cap (y, y + t)) \geq \mathcal{L}^1(\pi_1(K_i) \cap (x, x + t)) - |x - y|.$$

The other claims follow from a similar inequality. \square

7.4.5. *Defining the metric on \mathbb{R}^2 .* Define a norm field N on \mathbb{R}^2 by the formula

$$N_x = \begin{cases} 2^{-i/2} \|\cdot\|_1 & \text{if } x \in K_i \setminus K_{i+1}, i \text{ even,} \\ 2^{-(i-1)/2} \|\cdot\|_\infty & \text{if } x \in K_i \setminus K_{i+1}, i \text{ odd.} \end{cases}$$

The norm field N is admissible in the sense of Definition 3.1, in particular being lower semicontinuous, and induces a metric d on \mathbb{R}^2 as described in Section 3. Observe that N vanishes at a single point. Theorem 1.3 and Corollary 7.3 imply the following.

Proposition 7.7. The identity map $\iota: (\mathbb{R}^2, \|\cdot\|_2) \rightarrow (\mathbb{R}^2, d)$ is an isothermal quasiconformal mapping.

7.4.6. *Proof of Theorem 1.6.* We suppose to the contrary that there is a metric space $(\widehat{X}, \widehat{d})$ such that a factorization $\iota = \widehat{\iota} \circ P$ as in the statement of Theorem 1.6 exists, that is, that P is bi-Lipschitz and that $\widehat{\iota}$ has distortion $H(\widehat{\iota}) < \sqrt{2}$. Since ι is Lipschitz, it follows that $\widehat{\iota}$ is also Lipschitz.

By considering the metric $\widehat{d}(P(x), P(y))$ on \mathbb{R}^2 , we assume without loss of generality that $\widehat{X} = \mathbb{R}^2$ and that $\widehat{\iota}$ and P are each the identity map on \mathbb{R}^2 . Let \widehat{N} denote the metric derivative of the map $P: (\mathbb{R}^2, \|\cdot\|_2) \rightarrow (\mathbb{R}^2, \widehat{d})$ as defined in Definition 2.2.

By assumption, the identity map $\widehat{\iota}: (\mathbb{R}^2, \widehat{d}) \rightarrow (\mathbb{R}^2, d)$ is quasiconformal with $H(\widehat{\iota}) < \sqrt{2}$. Moreover, since P is Lipschitz, there exists $C > 0$ such that $\widehat{N}_x \leq C \|\cdot\|_1$ for every $x \in \mathbb{R}^2$. Let $v = (1, 0)$ and $w = (1/\sqrt{2}, 1/\sqrt{2})$, and let $a = H(\widehat{\iota})/\sqrt{2} < 1$. It suffices to show that, for all $i \geq 0$ and almost every $x \in K_i \setminus K_{i+1}$,

$$(39) \quad \begin{aligned} \widehat{N}_x(w) &\leq Ca^i & \text{if } i \text{ is even,} \\ \widehat{N}_x(v) &\leq Ca^i & \text{if } i \text{ is odd.} \end{aligned}$$

This provides a contradiction. Indeed, given that P is bi-Lipschitz, $\widehat{N}_x(w)$ and $\widehat{N}_x(v)$ are bounded from below for all $x \in \mathbb{R}^2$ by some constant $C' > 0$.

Observe that, when i is even, $N_x(w) = \sqrt{2}N_x(v)$ for all $x \in K_i \setminus K_{i+1}$. Similarly, when i is odd, $N_x(v) = \sqrt{2}N_x(w)$. It follows from Proposition 5.12 of [Iko19] that the pointwise distortion of $\widehat{\iota}$ coincides with the distortion of the identity map from $(\mathbb{R}^2, \widehat{N}_x)$ to (\mathbb{R}^2, N_x) for almost every $x \in \mathbb{R}^2$. As a consequence, for almost every $x \in K_i \setminus K_{i+1}$,

$$(40) \quad \begin{aligned} \widehat{N}_x(v) &\leq a\widehat{N}_x(w) & \text{if } i \text{ is even,} \\ \widehat{N}_x(w) &\leq a\widehat{N}_x(v) & \text{if } i \text{ is odd.} \end{aligned}$$

We verify (39) by induction on i . The claim is immediate for $i = 0$, recalling that $K_0 = \mathbb{R}^2$. For the inductive step, fix $i \geq 1$ and assume that (39) holds for almost every $x \in \mathbb{R}^2 \setminus K_i$. We show that (39) holds for almost every $x \in K_i \setminus K_{i+1}$. Let N_{i-1} denote the set of points in $K_{i-1} \setminus K_i$ for which (39) or (40) fails. We split into two cases based upon whether i is odd or even. The idea is the same in each, but the bookkeeping requires separate statements.

Case 1. Assume that i is odd. By the inductive hypothesis, we have $\widehat{N}_x(v) \leq a\widehat{N}_x(w) \leq Ca^i$ for every $x \in (K_{i-1} \setminus K_i) \setminus N_{i-1}$, where N_{i-1} has \mathcal{L}^2 -measure zero. We claim that $\widehat{N}_x(v) \leq Ca^i$ for almost every $x \in K_i \setminus K_{i+1}$. Assume to the contrary that there exists a set $G \subset K_i \setminus K_{i+1}$ of positive measure and a constant $b > 0$ such that $\widehat{N}_x(v) \geq (C+b)a^i$ for all $x \in G$.

For all $t \in [0, 1]$, let $\gamma_t: I \rightarrow \mathbb{R}^2$ be the path defined by $\gamma_t(s) = (s, t)$. According to Lemma 2.5, for every path γ_t and every subinterval $I' \subset I$,

$$\ell_{\widehat{d}}(\gamma_t|_{I'}) = \int_{I'} \widehat{N}_{\gamma_t(s)}(v) d\mathcal{L}^1.$$

Consider now the interval $[s_0, s_0 + h]$ for some $s_0 \in (0, 1)$ and $h \in (0, 1 - s_0)$. Differentiating, we have for \mathcal{L}^2 -almost every $(s_0, t) \in G$ that

$$\lim_{h \rightarrow 0} \frac{\widehat{d}(\gamma_t(s_0), \gamma_t(s_0 + h))}{h} = \lim_{h \rightarrow 0} \frac{\ell_{\widehat{d}}(\gamma_t|_{[s_0, s_0+h]})}{h} \geq (C + b)a^i.$$

In particular, for almost every $x \in G$, there exists $r_0 > 0$ such that

$$(41) \quad \widehat{d}(x, x + rv) \geq (C + b/2)a^i r$$

for all $r \in (0, r_0)$.

On the other hand, consider now a point $x \in G$ such that $\pi_1^*(x)$ is a Lebesgue density point of $\pi_1^*(K_{i-1})$ and $\pi_2^*(x)$ is a Lebesgue density point of $\pi_2^*(K_{i-1})$. Note that by Fubini's theorem, \mathcal{L}^2 -almost every point in G has this property. Let $\delta > 0$ and let $t_0 = t_0(\delta)$ be such that the hypothesis in Lemma 7.6 is satisfied for both the point $\pi_1^*(x)$ and the point $\pi_2^*(x)$.

For all $\varepsilon > 0$, let $H(x, \varepsilon)$ be the set comprising those points $y \in H_{i-1} \cap \overline{B}_{\|\cdot\|_2}(x, \varepsilon)$ for which

$$\mathcal{H}_{\|\cdot\|_2}^1(N_{i-1} \cap \pi_2^{-1}(\pi_2(y))) = 0.$$

Recall that the set H_{i-1} is defined in Section 7.4.4. By Lemma 7.5, the set $H_{i-1} \cap \overline{B}_{\|\cdot\|_2}(x, \varepsilon)$ has positive \mathcal{L}^2 -measure. Since N_{i-1} has \mathcal{L}^2 -measure zero, an application of Fubini's theorem shows that $H(x, \varepsilon)$ is a full measure subset of $H_{i-1} \cap \overline{B}_{\|\cdot\|_2}(x, \varepsilon)$. Let $r \in (0, t_0)$ and $\varepsilon \in (0, 2\delta r)$.

Consider a point $y \in H(x, \varepsilon)$. Let $\gamma_y: [0, r] \rightarrow \mathbb{R}^2$ be the path defined by $\gamma_y(s) = y + sv$. Lemma 2.5 implies that

$$\ell_{\widehat{d}}(\gamma_y) = \int_{[0, r]} \widehat{N}_{\gamma_y(s)}(v) d\mathcal{L}^1(s),$$

and the definition of $H(x, \varepsilon)$ implies that

$$(42) \quad \widehat{N}_z(v) \leq Ca^i$$

for $\mathcal{H}_{\|\cdot\|_2}^1$ -almost every $z \in K_{i-1} \cap |\gamma_y|$.

Next, we estimate the $\mathcal{H}_{\|\cdot\|_2}^1$ -measure of $K_{i-1} \cap |\gamma_y|$. To this end, observe that the path $\gamma_y^1: [0, r] \rightarrow \mathbb{R}$, $\gamma_y^1(s) = y + sv/\sqrt{2}$, intersects K_{i-1} in a set congruent to $\pi_1^*(K_{i-1}) \cap \pi_1^*(|\gamma_y^1|)$. Similarly, the path $\gamma_y^2: [0, r] \rightarrow \mathbb{R}$, $\gamma_y^2(s) = y + s\bar{w}/\sqrt{2}$, where $\bar{w} = (1/\sqrt{2}, -1/\sqrt{2})$, intersects K_{i-1} in a set congruent to $\pi_2^*(K_{i-1}) \cap \pi_2^*(|\gamma_y^2|)$. Since $|\pi_m^*(y) - \pi_m^*(x)| < 2\delta r$, Lemma 7.6 gives, for $m \in \{1, 2\}$,

$$(43) \quad \frac{\mathcal{H}_{\|\cdot\|_2}^1(K_{i-1} \cap |\gamma_y^m|)}{r/\sqrt{2}} \geq 1 - 2\delta.$$

We combine this with the following observation: for any measurable sets $E_1, E_2 \subset [0, r]$ satisfying $|E_j| \geq (1 - \varepsilon_j)r$ for some $\varepsilon_j \in (0, 1)$, $j \in \{1, 2\}$, the diagonal path $\gamma: [0, r] \rightarrow [0, r]^2$ defined by $\gamma(s) = (s, s)$ intersects $E_1 \times E_2$ in a set of length at least $\sqrt{2}(1 - \varepsilon_1 - \varepsilon_2)r$. Since K_{i-1} is constructed as a product set relative to which γ_y is a diagonal path, we conclude from (43) that

$$(44) \quad \frac{\mathcal{H}_{\|\cdot\|_2}^1(K_{i-1} \cap |\gamma_y|)}{r} \geq 1 - 4\delta.$$

Using (42) and the fact that $\widehat{N}_z(v) \leq C$ for all $z \in \mathbb{R}^2$, the inequality (44) gives

$$\widehat{d}(y, y + rv) \leq (1 - 4\delta)Ca^i r + 4\delta Cr.$$

Next, by making the initial choice of δ sufficiently small, we have

$$\widehat{d}(y, y + rv) \leq (1 + \delta)Ca^i r.$$

From this and the relationship $\widehat{d} \leq Cd_{\|\cdot\|_1} \leq \sqrt{2}Cd_{\|\cdot\|_2}$, it follows that

$$\widehat{d}(x, x + rv) \leq 2\sqrt{2}C\varepsilon + (1 + \delta)Ca^i r.$$

Since $\varepsilon \in (0, 2\delta r)$ is arbitrary, we obtain

$$\widehat{d}(x, x + rv) \leq (1 + \delta)Ca^i r.$$

Since this estimate holds for \mathcal{L}^2 -almost every $x \in G$, this contradicts our earlier statement (41) when δ is sufficiently small. We conclude that $\widehat{N}_x(v) \leq Ca^i$ for almost every $x \in K_i \setminus K_{i+1}$.

Case 2. We now consider the case that i is even. The idea is the same as in the first case, but now everything is rotated by $\pi/4$. By the inductive hypothesis, we have that $\widehat{N}_x(w) \leq a\widehat{N}_x(v) \leq Ca^i$ for every $x \in (K_{i-1} \setminus K_i) \setminus N_{i-1}$. We claim that $\widehat{N}_x(w) \leq Ca^i$ for almost every $x \in K_i \setminus K_{i+1}$. Assume to the contrary that there exists a set $G \subset K_i \setminus K_{i+1}$ of positive measure and a constant $b > 0$ such that $\widehat{N}_x(w) \geq (C + b)a^i$ for all $x \in G$.

For all $t \in J^*$, let $\gamma_t: I^* \rightarrow \mathbb{R}^2$ be the path defined by $\gamma_t(s) = \varphi(s, t)$. Consider as before the interval $[s_0, s_0 + h]$ for some $s_0 \in (0, \sqrt{2})$ and $h \in (0, \sqrt{2} - s_0)$. Differentiating, we have that

$$\lim_{h \rightarrow 0} \frac{\ell_{\widehat{d}}(\gamma_t|_{[s_0, s_0+h]})}{h} \geq (C + b)a^i.$$

In particular, for \mathcal{L}^2 -almost every $x \in G$, there exists $r_0 > 0$ such that

$$(45) \quad \widehat{d}(x, x + rw) \geq (C + b/2)a^i r$$

for all $r \in (0, r_0)$.

On the other hand, consider a point $x \in G$ such that $\pi_1(x)$ is a Lebesgue density point of $\pi_1(K_{i-1})$ and $\pi_2(x)$ is a Lebesgue density point of $\pi_2(K_{i-1})$. Let $\delta > 0$ and let $t_0 = t_0(\delta)$ be the corresponding value in Lemma 7.6. For all $\varepsilon > 0$, define the set $H(x, \varepsilon)$ as the set of points $y \in H_{i-1} \cap \overline{B}_{\|\cdot\|_2}(x, \varepsilon)$ for which

$$\mathcal{H}_{\|\cdot\|_2}^1(N_{i-1} \cap (\pi_2^*)^{-1}(\pi_2^*(y))) = 0.$$

As before, $H(x, \varepsilon)$ is a full measure subset of $H_{i-1} \cap \overline{B}_{\|\cdot\|_2}(x, \varepsilon)$. Let $r \in (0, t_0)$ and $\varepsilon \in (0, 2\delta r)$.

For all $y \in H(x, \varepsilon)$, define the path $\gamma_y: [0, r] \rightarrow \mathbb{R}^2$ by $\gamma_y(s) = y + sw$. Recall from Lemma 2.5 that

$$\ell_{\widehat{d}}(\gamma_y) = \int_{[0, r]} \widehat{N}_{\gamma_y(s)}(w) d\mathcal{L}^1.$$

Moreover, $\widehat{N}_z(w) \leq Ca^i$ for $\mathcal{H}_{\|\cdot\|_2}^1$ -almost every $z \in K_{i-1} \cap |\gamma_y|$. Arguing as in the first case, we obtain the inequality

$$\widehat{d}(y, y + rw) \leq (1 - 4\delta)Ca^i r + 2\delta Cr.$$

Next, by taking δ sufficiently small, we then have $\widehat{d}(y, y + rw) \leq (1 + \delta)Ca^i r$. As before, since $\varepsilon \in (0, 2\delta r)$ is arbitrary,

$$\widehat{d}(x, x + rw) \leq (1 + \delta)Ca^i r,$$

which contradicts (45) for sufficiently small $\delta > 0$. We conclude that $\widehat{N}_x(w) \leq Ca^i$ for almost every $x \in K_i \setminus K_{i+1}$.

REFERENCES

- [AB50] Lars Ahlfors and Arne Beurling. Conformal invariants and function-theoretic null-sets. *Acta Math.*, 83:101–129, 1950.
- [AHPCS18] Nathan Albin, Jared Hoppis, Pietro Poggi-Corradini, and Nageswari Shanmugalingam. Infinity modulus and the essential metric. *J. Math. Anal. Appl.*, 467(1):570–584, 2018.
- [AIM09] Kari Astala, Tadeusz Iwaniec, and Gaven Martin. *Elliptic partial differential equations and quasiconformal mappings in the plane*, volume 48 of *Princeton Mathematical Series*. Princeton University Press, Princeton, NJ, 2009.
- [AS60] Lars V. Ahlfors and Leo Sario. *Riemann surfaces*. Princeton Mathematical Series, No. 26. Princeton University Press, Princeton, N.J., 1960.
- [AT04] Luigi Ambrosio and Paolo Tilli. *Topics on analysis in metric spaces*, volume 25 of *Oxford Lecture Series in Mathematics and its Applications*. Oxford University Press, Oxford, 2004.
- [AZ67] A. D. Aleksandrov and V. A. Zalgaller. *Intrinsic geometry of surfaces*. Translated from the Russian by J. M. Danskin. Translations of Mathematical Monographs, Vol. 15. American Mathematical Society, Providence, R.I., 1967.
- [BCO19] A. L. Baisón, A. Clop, and J. Orobitg. Distributional solutions of the Beltrami equation. *J. Math. Anal. Appl.*, 470(2):1081–1094, 2019.
- [BHR01] Mario Bonk, Juha Heinonen, and Steffen Rohde. Doubling conformal densities. *J. Reine Angew. Math.*, 541:117–141, 2001.
- [Bis07] Christopher J. Bishop. An A_1 weight not comparable with any quasiconformal Jacobian. In *In the tradition of Ahlfors-Bers. IV*, volume 432 of *Contemp. Math.*, pages 7–18. Amer. Math. Soc., Providence, RI, 2007.
- [BKR98] Mario Bonk, Pekka Koskela, and Steffen Rohde. Conformal metrics on the unit ball in Euclidean space. *Proc. London Math. Soc. (3)*, 77(3):635–664, 1998.
- [BM41] Herbert Busemann and Walther Mayer. On the foundations of calculus of variations. *Trans. Amer. Math. Soc.*, 49:173–198, 1941.
- [BM17] Mario Bonk and Daniel Meyer. *Expanding Thurston maps*, volume 225 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2017.
- [Bog07] V. I. Bogachev. *Measure theory. Vol. I, II*. Springer-Verlag, Berlin, 2007.
- [Bon06] Mario Bonk. Quasiconformal geometry of fractals. In *International Congress of Mathematicians. Vol. II*, pages 1349–1373. Eur. Math. Soc., Zürich, 2006.
- [CS19] Paul Creutz and Eleftherios Soultanis. Maximal metric surfaces and the Sobolev-to-Lipschitz property. *arXiv e-prints*, page arXiv:1909.10385, September 2019.
- [Dav86] Robert J. Daverman. *Decompositions of manifolds*, volume 124 of *Pure and Applied Mathematics*. Academic Press, Inc., Orlando, FL, 1986.
- [Dav98] Guy David. Unrectifiable 1-sets have vanishing analytic capacity. *Rev. Mat. Iberoamericana*, 14(2):369–479, 1998.
- [DCP88] Giuseppe De Cecco and Giuliana Palmieri. Intrinsic distance on a Lipschitz Riemannian manifold. *Rend. Sem. Mat. Univ. Politec. Torino*, 46(2):157–170 (1990), 1988.
- [DCP90] Giuseppe De Cecco and Giuliana Palmieri. Length of curves on Lip manifolds. *Atti Accad. Naz. Lincei Cl. Sci. Fis. Mat. Natur. Rend. Lincei (9) Mat. Appl.*, 1(3):215–221, 1990.
- [DCP91] Giuseppe De Cecco and Giuliana Palmieri. Integral distance on a Lipschitz Riemannian manifold. *Math. Z.*, 207(2):223–243, 1991.
- [DCP95] Giuseppe De Cecco and Giuliana Palmieri. LIP manifolds: from metric to Finslerian structure. *Math. Z.*, 218(2):223–237, 1995.
- [DS90] Guy David and Stephen Semmes. Strong A_∞ weights, Sobolev inequalities and quasiconformal mappings. In *Analysis and partial differential equations*, volume 122 of *Lecture Notes in Pure and Appl. Math.*, pages 101–111. Dekker, New York, 1990.
- [Dud07] Jakub Duda. Absolutely continuous functions with values in a metric space. *Real Anal. Exchange*, 32(2):569–581, 2007.
- [Fed69] Herbert Federer. *Geometric measure theory*. Die Grundlehren der mathematischen Wissenschaften, Band 153. Springer-Verlag New York Inc., New York, 1969.
- [GM85] F. W. Gehring and O. Martio. Quasiextremal distance domains and extension of quasiconformal mappings. *J. Analyse Math.*, 45:181–206, 1985.
- [Gol69] G. M. Goluzin. *Geometric theory of functions of a complex variable*. Translations of Mathematical Monographs, Vol. 26. American Mathematical Society, Providence, R.I., 1969.

- [GPP06] Adriana Garroni, Marcello Ponsiglione, and Francesca Prinari. From 1-homogeneous supremal functionals to difference quotients: relaxation and Γ -convergence. *Calc. Var. Partial Differential Equations*, 27(4):397–420, 2006.
- [Hei01] Juha Heinonen. *Lectures on analysis on metric spaces*. Universitext. Springer-Verlag, New York, 2001.
- [HH08] Hrant Hakobyan and David A. Herron. Euclidean quasiconvexity. *Ann. Acad. Sci. Fenn. Math.*, 33(1):205–230, 2008.
- [HKST15] Juha Heinonen, Pekka Koskela, Nageswari Shanmugalingam, and Jeremy T. Tyson. *Sobolev spaces on metric measure spaces*, volume 27 of *New Mathematical Monographs*. Cambridge University Press, Cambridge, 2015. An approach based on upper gradients.
- [HS93] Zheng-Xu He and Oded Schramm. Fixed points, Koebe uniformization and circle packings. *Ann. of Math. (2)*, 137(2):369–406, 1993.
- [Iko19] Toni Ikonen. Uniformization Of Metric Surfaces Using Isothermal Coordinates. *arXiv e-prints*, page arXiv:1909.09113, September 2019.
- [Iva08] S. V. Ivanov. Volumes and areas of Lipschitz metrics. *Algebra i Analiz*, 20(3):74–111, 2008.
- [Kir94] Bernd Kirchheim. Rectifiable metric spaces: local structure and regularity of the Hausdorff measure. *Proc. Amer. Math. Soc.*, 121(1):113–123, 1994.
- [KKR19] Sergei Kalmykov, Leonid V. Kovalev, and Tapio Rajala. Removable sets for intrinsic metric and for holomorphic functions. *J. Anal. Math.*, 139(2):751–772, 2019.
- [Laa02] Tomi J. Laakso. Plane with A_∞ -weighted metric not bi-Lipschitz embeddable to \mathbb{R}^N . *Bull. London Math. Soc.*, 34(6):667–676, 2002.
- [LW17] Alexander Lytchak and Stefan Wenger. Area minimizing discs in metric spaces. *Arch. Ration. Mech. Anal.*, 223(3):1123–1182, 2017.
- [LW18] Alexander Lytchak and Stefan Wenger. Intrinsic structure of minimal discs in metric spaces. *Geom. Topol.*, 22(1):591–644, 2018.
- [LW20] Alexander Lytchak and Stefan Wenger. Canonical parameterizations of metric disks. *Duke Math. J.*, 2020. Advance publication.
- [Raj17] Kai Rajala. Uniformization of two-dimensional metric surfaces. *Invent. Math.*, 207(3):1301–1375, 2017.
- [Res93] Yu. G. Reshetnyak. Two-dimensional manifolds of bounded curvature. In *Geometry, IV*, volume 70 of *Encyclopaedia Math. Sci.*, pages 3–163, 245–250. Springer, Berlin, 1993.
- [Rom19] Matthew Romney. Quasiconformal parametrization of metric surfaces with small dilatation. *Indiana Univ. Math. J.*, 68(3):1003–1011, 2019.
- [RR19] Kai Rajala and Matthew Romney. Reciprocal lower bound on modulus of curve families in metric surfaces. *Ann. Acad. Sci. Fenn. Math.*, 44(2):681–692, 2019.
- [Sch95] Oded Schramm. Transboundary extremal length. *J. Anal. Math.*, 66:307–329, 1995.
- [Sem93] Stephen Semmes. Bi-Lipschitz mappings and strong A_∞ weights. *Ann. Acad. Sci. Fenn. Ser. A I Math.*, 18(2):211–248, 1993.
- [Sem96] Stephen Semmes. On the nonexistence of bi-Lipschitz parameterizations and geometric problems about A_∞ -weights. *Rev. Mat. Iberoamericana*, 12(2):337–410, 1996.
- [Tol03] Xavier Tolsa. Painlevé’s problem and the semiadditivity of analytic capacity. *Acta Math.*, 190(1):105–149, 2003.
- [Vod00] Serguei K. Vodop’yanov. P-differentiability on Carnot groups in different topologies and related topics. In *Proceedings on Analysis and Geometry (Russian) (Novosibirsk Akademgorodok, 1999)*, pages 603–670. Izdat. Ross. Akad. Nauk Sib. Otd. Inst. Mat., Novosibirsk, 2000.
- [Why64] Gordon Thomas Whyburn. *Topological analysis*. Second, revised edition. Princeton Mathematical Series, No. 23. Princeton University Press, Princeton, N.J., 1964.
- [Wil70] Stephen Willard. *General topology*. Addison-Wesley Publishing Co., Reading, Mass.-London-Don Mills, Ont., 1970.
- [Wil12] Marshall Williams. Geometric and analytic quasiconformality in metric measure spaces. *Proc. Amer. Math. Soc.*, 140(4):1251–1266, 2012.
- [You15] Malik Younsi. On removable sets for holomorphic functions. *EMS Surv. Math. Sci.*, 2(2):219–254, 2015.

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF JYVASKYLA, P.O. Box 35 (MAD), FI-40014, UNIVERSITY OF JYVASKYLA, FINLAND.

E-mail address: `toni.m.h.ikonen@jyu.fi`

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, UNIVERSITY OF COLOGNE, WEYERTAL 86-90, 50931 KÖLN, GERMANY.

E-mail address: `mromney@math.uni-koeln.de`