

**This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.**

**Author(s):** Eck, Jenalle L.; Kytöviita, Minna-Maarit; Laine, Anna-Liisa

**Title:** Arbuscular mycorrhizal fungi influence host infection during epidemics in a wild plant pathosystem

**Year:** 2022

**Version:** Accepted version (Final draft)

**Copyright:** © 2022 The Authors New Phytologist © 2022 New Phytologist Foundation

**Rights:** CC BY-NC-ND 4.0

**Rights url:** <https://creativecommons.org/licenses/by-nc-nd/4.0/>

**Please cite the original version:**

Eck, J. L., Kytöviita, M., & Laine, A. (2022). Arbuscular mycorrhizal fungi influence host infection during epidemics in a wild plant pathosystem. *New Phytologist*, 236(5), 1922-1935.

<https://doi.org/10.1111/nph.18481>

# Arbuscular mycorrhizal fungi influence host infection during epidemics in a wild plant pathosystem

Jenalle L. Eck<sup>1</sup> , Minna-Maarit Kytöviita<sup>2</sup>  and Anna-Liisa Laine<sup>1,3</sup> 

<sup>1</sup>Department of Evolutionary Biology and Environmental Studies, University of Zurich, 8057 Zurich, Switzerland; <sup>2</sup>Department of Biological and Environmental Science, University of Jyväskylä, 40014 Jyväskylä, Finland; <sup>3</sup>Organismal and Evolutionary Biology Research Program, Faculty of Biological and Environmental Sciences, University of Helsinki, 00790 Helsinki, Finland

Author for correspondence:

Jenalle L. Eck

Email: [jenalle.eck@ieu.uzh.ch](mailto:jenalle.eck@ieu.uzh.ch)

Received: 15 February 2022

Accepted: 15 August 2022

New Phytologist (2022)

doi: 10.1111/nph.18481

**Key words:** mutualism, mycorrhizal fungi, plant disease, plant pathogen, *Plantago lanceolata*, *Podosphaera plantaginis*, powdery mildew, protective symbiont.

## Summary

- While pathogenic and mutualistic microbes are ubiquitous across ecosystems and often occur within hosts, how they interact to determine patterns of disease in genetically diverse wild populations is unknown.
- To test whether microbial mutualists provide protection against pathogens, and whether this varies among host genotypes, we conducted a field experiment in three naturally occurring epidemics of a fungal pathogen, *Podosphaera plantaginis*, infecting a host plant, *Plantago lanceolata*, in the Åland Islands, Finland. In each population, we collected epidemiological data on experimental plants from six allopatric populations that had been inoculated with a mixture of mutualistic arbuscular mycorrhizal fungi or a nonmycorrhizal control.
- Inoculation with arbuscular mycorrhizal fungi increased growth in plants from every population, but also increased host infection rate. Mycorrhizal effects on disease severity varied among host genotypes and strengthened over time during the epidemic. Host genotypes that were more susceptible to the pathogen received stronger protective effects from inoculation.
- Our results show that arbuscular mycorrhizal fungi introduce both benefits and risks to host plants, and shift patterns of infection in host populations under pathogen attack. Understanding how mutualists alter host susceptibility to disease will be important for predicting infection outcomes in ecological communities and in agriculture.

## Introduction

Protective symbionts – species that provide defensive benefits to their hosts – help to determine the outcome of species interactions and, thus, shape ecological and evolutionary dynamics between hosts and parasites (Brownlie & Johnson, 2009; May & Nelson, 2014; King *et al.*, 2016; Sochard *et al.*, 2020). Despite their importance, ecological studies examining the role of protective symbionts in influencing host–parasite interactions in natural populations and communities are rare (Oliver *et al.*, 2014; Hafer-Hahmann & Vorbürger, 2021). Protection against infectious disease by mutualistic microbes, such as mycorrhizal fungi, has been demonstrated under controlled laboratory conditions in several economically important agricultural plant species (Norman *et al.*, 1996; Pozo *et al.*, 2002; Hao *et al.*, 2005; Li *et al.*, 2010; Song *et al.*, 2015; Berdeni *et al.*, 2018). Although mutualists may also affect disease under field conditions (Newsham *et al.*, 1995), it has not been verified whether protective symbionts mediate infection under natural epidemics, which are characterized by repeated pathogen encounters, as well as by environmental and genotypic diversity. Mycorrhizal associations are widespread among terrestrial plants (Öpik *et al.*, 2006; van der Heijden

*et al.*, 2008) and have important impacts on plant fitness and population dynamics (Barea *et al.*, 2002; Koide & Dickie, 2002), community composition (Hartnett & Wilson, 2002) and ecosystem functioning (Rillig, 2004). Understanding how mycorrhizal fungi and other mutualists influence patterns of plant disease is essential given that disease is a major factor shaping the abundance, diversity and distribution of species in plant communities (Bever *et al.*, 2015) and affecting food production (Johansson *et al.*, 2004; Gosling *et al.*, 2006; Pretty *et al.*, 2011; Hohmann & Messmer, 2017). Although both plant-associated pathogenic and mutualistic microbes are ubiquitous across ecosystems, how they interact to determine disease risk in natural, genetically diverse populations is not known.

Mycorrhizal fungi produce a suite of growth, nutritional and/or defensive effects that may help protect plants from co-occurring antagonists, such as pathogenic microbes and herbivores (Delavaux *et al.*, 2017). Association with mycorrhizal fungi often improves plant nutrient and water uptake (Smith & Read, 2008), although there is both intra- and interspecific variation among plants in their ability to form and benefit from mycorrhizal associations (Thrall *et al.*, 2011; Rasmussen *et al.*, 2019). Increases in host size and nutritional status as a result of

mycorrhizal association can improve host tolerance to parasites and abiotic stress (Azcón-Aguilar & Barea, 1996). Arbuscular mycorrhizal fungi can also influence host defenses directly, by upregulating defense gene expression in their host plants (Azcón-Aguilar & Barea, 1996; Pozo & Azcón-Aguilar, 2007; Jung *et al.*, 2012; Goddard *et al.*, 2021). This form of protection, known as defense priming, allows a more efficient activation of defense mechanisms in response to attack by potential enemies and has been shown to reduce the negative effects of interactions with a wide range of antagonist species (Jung *et al.*, 2012; Delavaux *et al.*, 2017). Although mycorrhizal associations occur belowground (at the root–soil interface), the induced resistance response in the host is systemic (Cameron *et al.*, 2013; Goddard *et al.*, 2021), meaning that even strictly aboveground parasites may be affected (Koricheva *et al.*, 2009). It is unclear how often mycorrhizal growth and defensive benefits are conferred together in hosts and how they operate simultaneously to determine the incidence and outcome of interactions between hosts and parasites.

Empirical studies in controlled environments have shown that host association with mutualists may also present risks that can influence infection dynamics (Polin *et al.*, 2014). For example, ecological costs may occur when combinations of host and mutualist species or genotypes are mismatched (Klironomos, 2003; Hoeksema *et al.*, 2010), resulting in inefficient mutualisms that fail to convert host resources into growth or defensive benefits (Johnson *et al.*, 1997; Jones & Smith, 2004; Grman, 2012). Furthermore, unfavorable abiotic conditions may reduce or negate potential mutualist benefits (Hoeksema *et al.*, 2010; Qu *et al.*, 2021). In addition, defensive benefits from mutualists may not be effective against all parasite species (e.g. depending on their life history) (Pozo & Azcón-Aguilar, 2007) or durable to changes in parasite traits and/or composition in the environment. Finally, mutualists could also affect patterns of host infection indirectly, for example, via changes in host size that influence parasite contact rates. Hence, the potential ecological risks and/or benefits of mutualist association in the presence of parasites may depend on host genotype and vary among or within host populations and environments; however, this has remained largely unexplored in natural populations.

In addition, it is unclear how mutualism-derived protection acts alongside innate host resistance to determine the outcome of host–pathogen interactions. Host genetic resistance can vary widely among and within natural populations (Salvaudon *et al.*, 2008; Laine *et al.*, 2011) – potentially as a result of costs associated with its maintenance (Brown, 2003; Susi & Laine, 2015) – and may be under a different set of selection pressures than mutualism (Thompson, 1994). Whether symbiosis presents benefits or costs to host individuals and populations could depend on their degree of resistance. In resistant hosts, resources provided to mutualists in return for defensive benefits could represent an unnecessary metabolic cost. However, in susceptible hosts, mutualist-derived protection could compensate for lack of genetic resistance, presenting a viable alternate strategy for coping with pathogens. Mutualism-derived protection could be especially beneficial when genetic resistance is costly to

maintain or when disease is ephemeral, as it often is in natural populations (Burdon & Thrall, 2014). How much of host resistance is derived from innate genetic defenses vs protective symbionts (e.g. defense priming or improved pathogen tolerance), and whether these types of defenses are linked in different species combinations and environmental contexts remain to be seen.

Despite general recognition for the impact of mycorrhizal fungi on plant fitness, how mycorrhizal association may impact host infection – and to what extent this varies among plant genotypes and populations – is poorly understood in natural populations. To examine this, we conducted a field experiment to test whether inoculation with arbuscular mycorrhizal fungi affects infection dynamics by a fungal pathogen in a shared host under natural epidemic conditions. Specifically, we ask the following questions: do growth effects resulting from inoculation with arbuscular mycorrhizal fungi vary among host populations and maternal genotypes; does inoculation with arbuscular mycorrhizal fungi influence host infection rate; upon infection, does prior inoculation with arbuscular mycorrhizal fungi affect disease severity; are growth and defensive effects from inoculation with arbuscular mycorrhizal fungi linked in host genotypes; and are disease susceptibility and defensive effects as a result of inoculation with arbuscular mycorrhizal fungi linked in host genotypes? To answer these questions within an ecologically relevant context, we placed mycorrhizal-inoculated and nonmycorrhizal-inoculated plants in wild host populations during a natural pathogen epidemic. The experiment was conducted in the long-term Åland Islands study site (Finland), where infection by a fungal pathogen (powdery mildew) has been surveyed on a large host population network of *Plantago lanceolata* L. (Plantaginaceae) since 2001 (Jousimo *et al.*, 2014). From prior studies in this pathosystem, we know that several factors, such as spatial context (Laine, 2006; Soubeyrand *et al.*, 2009; Jousimo *et al.*, 2014), pathogen genetic diversity (Eck *et al.*, 2022), local adaptation of pathogen strains to sympatric host populations (Laine, 2005, 2007a) and abiotic conditions (Laine, 2007b, 2008; Penczykowski *et al.*, 2015), are all critical in determining infection, but the impact of mutualistic interactions on infection dynamics has not been determined. To our knowledge, this is the first study of the impact of mycorrhizal fungi on infection by a plant pathogen under natural epidemics and across different host populations and genotypes.

## Materials and Methods

### Study system

Our study is focused on a fungal pathogen, *Podosphaera plantaginis* (Castagne) U. Braun & S. Takam., infecting *Plantago lanceolata* L. (Plantaginaceae) in the Åland Islands (60°08'53"N, 19°47'18"E), Finland. We carried out an experiment in three natural *P. lanceolata* populations that are part of a network of > 4000 mapped populations (Hanski, 1999). These populations have been surveyed for infection by *P. plantaginis* since 2001 (Laine & Hanski, 2006). *Plantago lanceolata* (ribwort plantain) is native to Åland and much of Eurasia; it occurs mainly in small

meadows and disturbed areas in Åland. It is monoecious, self-incompatible and reproduces either sexually (via wind-dispersed pollen and seeds; Bos, 1992) or asexually (via clonally produced side-rosettes) (Sagar & Harper, 1964). *Plantago lanceolata* associates commonly with mycorrhizal fungi and has been found in association with a wide variety of mycorrhizal species in Åland (Rasmussen *et al.*, 2018; J. L. Eck *et al.*, unpublished).

*Podosphaera plantaginis*, a powdery mildew fungus (order Erysiphales), is an obligate biotroph of foliar tissue and is host-specific in Åland, infecting only *P. lanceolata* (Laine, 2004). Fungal hyphae grow on the surface of *P. lanceolata* leaves, producing localized infections that mitigate host growth and reproduction (Bushnell, 2002), and may lead to mortality in the presence of abiotic stress, such as drought (Laine, 2004; Susi & Laine, 2015). Infections are transmitted via asexually produced, wind-dispersed spores (conidia) that are produced cyclically (*c.* every 2 wk) throughout an epidemic season (approximately June to September in Åland) (Ovaskainen & Laine, 2006). Resting structures (chasmothecia), produced via haploid selfing or outcrossing between strains (Tollenaere & Laine, 2013), allow the pathogen to overwinter (Tack & Laine, 2014). In Åland, *P. plantaginis* persists as a metapopulation, with frequent colonization and extinction events (Jousimo *et al.*, 2014). Resistance in *P. lanceolata* against *P. plantaginis* is strain-specific (Laine, 2004, 2007a). Previous studies have demonstrated high variation in resistance within and among host populations (Laine, 2004, 2007a; Jousimo *et al.*, 2014).

### Mycorrhizal inoculation of experimental plants


To measure the effects of association with arbuscular mycorrhizal fungi in genetically diverse host plants, seeds were collected from six geographically variable populations of *P. lanceolata* in the Åland Islands (Fig. 1a, right panel) in August 2007. Seeds were collected from five haphazardly chosen maternal plants in each population and were stored separately in paper envelopes (seeds from one maternal plant are half or full siblings). In April 2008, seeds from each of the 30 maternal plants were planted in separate 6 × 6 × 7 cm pots in a glasshouse at the University of Oulu (Oulu, Finland) in sterilized sand. Ten healthy 2-wk-old seedlings from each maternal genotype were transferred to individual pots (one seedling per 6 × 6 × 7 cm pot) filled with experimental substrate (a 5 : 4 : 1 mixture of heat-sterilized sand : heat-sterilized garden soil : perlite, combined with 1 g of bone meal and 3 g dolomite l<sup>-1</sup> substrate). At the time of transfer, five of the seedlings from each maternal genotype were inoculated with mycorrhizal fungi, while the other five received a nonmycorrhizal control treatment. The mycorrhizal inoculum consisted of a mixture of spores of three arbuscular mycorrhizal fungal species native to Finland (*Glomus hoi*, *Claroideoglomus clareoideum* and *Glomus mosseae* (BEG 29)), as *P. lanceolata* is colonized naturally by several mycorrhizal symbionts (Johnson *et al.*, 2003). Allopatric isolates of each species (originating from central Finland (61°10'N, 24°40'E) rather than Åland) were used so as not to confound the experiment with potential local adaptation between mycorrhizal fungi and their host populations (Hoeksema

*et al.*, 2010). Spore inocula were produced by growing the mycorrhizal species with nonexperimental *P. lanceolata* in a soil substrate identical to the experimental one. Spores were washed out of the nonexperimental substrate with water, and 15 spores of each species (45 spores in total) were pipetted on to the roots of each seedling in the mycorrhizal treatment in 2 ml of water (6 ml in total); seedlings in the nonmycorrhizal control treatment received 2 ml of filtered spore washing water from each fungal species (6 ml in total). Hereafter, plants in the mycorrhizal-inoculated treatment are abbreviated as 'AMF' (i.e. arbuscular mycorrhizal fungi), and plants in the nonmycorrhizal control treatment as 'NM' (i.e. nonmycorrhizal). All experimental seedlings were also inoculated with bacteria at this time: bacteria were filtered from a soil mix collected from the six Åland seed source populations to restore the native, nonmycorrhizal soil microbial community. Seedlings were fertilized weekly with a dilute nitrogen-based solution. Natural light in the glasshouse was supplemented with Osram HQI lamps to provide a photoperiod of 18 h : 6 h, light : dark. At 6 wk of age (4 wk post-inoculation), the plants were moved to an outdoor area at the University of Oulu and grown on tables under a transparent plastic roof for an additional 6 wk, to acclimatize to field conditions (as *P. lanceolata* and *P. plantaginis* do not inhabit this region, infection at this stage was highly unlikely).

### Natural epidemic field experiment

At 10 wk post-inoculation (mid-July 2008), 288 healthy 12-wk-old experimental plants were placed in to three naturally occurring populations of *P. lanceolata*, infected by populations of *P. plantaginis*, in Åland to gain infections naturally from the surrounding epidemic. The three field epidemic populations were allopatric to the six seed origin populations (Fig. 1a, right panel). Thus, they represent common-garden field sites, in which the host genotypes are not locally adapted to the local environmental conditions or to the local pathogen population. Presence of *P. plantaginis* at the sites was confirmed by surveying the populations for visible signs of infection before placement of the experimental plants. Each field site received 96 experimental plants: 16 plants from each of the six seed origin populations (eight AMF plants and eight NM plants). Within these subgroups, the five maternal genotypes within each seed origin population were allocated as evenly as possible to the three field sites; the individuals placed in each site were selected at random (full, pairwise replication of every maternal genotype × mycorrhizal treatment combination within every field site was not possible owing to lack of plants). At each field site, experimental plants were randomly fitted in to one of 96 plastic containers (14 × 10.5 × 4.5 cm) that were placed on the ground in a random array in the proximity of naturally infected, wild *P. lanceolata* individuals inhabiting the field population. The containers prevented contact between the bottom of the pots and the soil, reducing the likelihood that the experimental plants would acquire soil microbes from the environment.

At the time of placement in the field epidemic populations, we measured the initial size of each experimental plant: the total



**Fig. 1** Inoculation with arbuscular mycorrhizal fungi produced variable growth benefits in experimental plants from different genetic origins. Before exposure to *Podospheera plantaginis* in field conditions, negative binomial generalized linear models showed that the magnitude of the growth benefits in experimental *Plantago lanceolata* plants (following inoculation with a mixture of three arbuscular mycorrhizal fungal species) varied among 30 host maternal genotypes (a, left; Table S1; mycorrhizal inoculation (MYC)  $\times$  maternal genotype (GEN),  $P < 0.001$ ,  $n = 287$  plants). In the right panel of (a), colored dots represent seed origin populations and black dots represent field epidemic populations. After pathogen exposure in the field epidemic experiment, host growth continued to be linked to mycorrhizal inoculation (b, c; Tables S2, S3; MYC,  $P < 0.001$ ), seed origin population (b; Table S2; seed origin population,  $P < 0.001$ ), and maternal genotype (c; Table S3; GEN,  $P < 0.001$ ), but growth benefits no longer varied among maternal genotypes. In (a), error bars represent a 95% confidence interval. In (b, c), black dots represent outlier individuals, box notches represent a 95% confidence interval for comparing medians, box hinges correspond to the 1<sup>st</sup> and 3<sup>rd</sup> quartiles, and box whiskers extend to the largest and smallest values no further than 1.5 $\times$  the interquartile range from the hinges. Myc., mycorrhizal; Non-Myc., nonmycorrhizal.

number of leaves, as well as the length and width (in cm) of the longest leaf were recorded. Plants were considered infected if any leaf showed powdery, white spot(s) (i.e. characteristic signs of infection with *P. plantaginis* in this region; Fig. S1), forming a blotch or lesion of any size upon visual inspection (all leaves were inspected for signs of infection). All experimental plants were uninfected (i.e. had zero leaves infected by *P. plantaginis*) at the beginning of the experiment. Infection surveys were then conducted every 3 d at each field site, on a rotating survey schedule (1<sup>st</sup> day field population, 9051; 2<sup>nd</sup> day field population, 3484; and 3<sup>rd</sup> day field population, 9066). During each infection survey, the number of total leaves and leaves infected by *P. plantaginis* on each plant were counted (leaves that withered during the experiment were not counted). The plants were also re-randomized into a new position in the experimental array (this was done to minimize the effect of spatial positioning, with respect to distance to infected individuals and prevailing wind direction, on infection rate and severity), and were watered if necessary. Seven infection surveys were conducted in each field population (Datasets S1, S2). At the end of the experiment (in late August 2008) a haphazardly selected subset of 21 AMF plants and 20 NM plants were harvested, and their oven-dried above-ground biomass was measured. We created a metric approximating the leaf area of each plant at the beginning of the experiment by first applying the length and width of the plant's longest leaf to the equation yielding the area of an oval, then multiplying the area of the longest leaf by the total number of leaves on the plant. Heavy rains during the fourth and fifth surveys resulted in missing infection data for some field populations (identifying symptoms caused by *P. plantaginis* on wet leaves is challenging) and may have influenced infection in the sixth and seventh surveys in all populations (as heavy rain washes spores away from infected leaf tissue and damages spore viability; Sivapalan, 1993). Because of this, we focus here on infection data from the third survey (coinciding with peak infection rates and the completion of one 14 d initial pathogen infection and reproduction cycle), as well as the final, seventh survey at the end of the experiment (coinciding with reinfections because of pathogen reproduction and incorporating variable abiotic conditions). Using data from these two surveys (peak epidemic and end-of-experiment), we quantified the infection status (0/1, infected or uninfected) and calculated the infection severity (i.e. the proportion of infected leaves) of each experimental plant. Plants were considered infected if one or more leaves showed signs of infection. One plant that died during the experiment was excluded from all analyses.

## Statistical methods

**Do growth effects resulting from inoculation with arbuscular mycorrhizal fungi vary among host populations and maternal genotypes?** To test whether host growth was explained by inoculation with arbuscular mycorrhizal fungi, host genetic origin or an interaction between these factors, we built a series of generalized linear models (GLMs). All statistical tests were conducted in the R statistical environment (R Core Team, 2021). Negative binomial GLMs were constructed with the MASS package to

counter overdispersion (Venables & Ripley, 2002). To explore the effect of genetic origin on host growth at two levels of biological organization, seed origin population (POP) and maternal genotype (GEN) were included as explanatory factors in separate models. Host growth was modeled before epidemic exposure (at the time of placement in the field experiment) and post-epidemic exposure (during the last survey of the experiment) using the number of leaves on each host. When modeling initial host size, the explanatory factors were mycorrhizal treatment (MYC), POP/GEN and their interaction. When modeling final host size, field epidemic population (SITE), initial host size (SIZE) and a four-way interaction term between all factors were also included as explanatory factors whenever possible (without compromising model fit). We also tested whether a subset of AMF and NM plants varied in aboveground dry biomass at the end of the experiment using a Mann–Whitney *U*-test to counter nonnormality in the data. Throughout our study, nonsignificant interaction terms were removed from all final models. To address nonindependence of the mycorrhizal treatment and host size variables during the field experiment, all models including size were also tested with this variable omitted, and the results were compared.

**Does inoculation with arbuscular mycorrhizal fungi influence host infection rate?** To test whether host infection rate upon exposure to field epidemics is explained by inoculation with arbuscular mycorrhizal fungi, host genetic origin, field epidemic population or an interaction between these factors, we built a series of generalized logistic regression models. Host infection status was used as the binomial response variable in each model (with family set to quasibinomial to counter overdispersion). Infection data were analyzed at the peak of the epidemic and at the end of the experiment. In each model, MYC, POP/GEN and SITE (and SIZE at the corresponding time point, in applicable models) were included as explanatory factors, as well as all interaction terms.

**Upon infection, does previous inoculation with arbuscular mycorrhizal fungi affect disease severity?** To test whether disease severity in infected hosts is influenced by previous inoculation with arbuscular mycorrhizal fungi, host genetic origin, field epidemic population or an interaction between these factors, we built a series of generalized logistic regression models. These models explored two measures of disease severity (i.e. the proportion of leaves infected and the number of infected leaves in infected individuals) at two time points (i.e. at the peak of the epidemic and at the end of the experiment). In each model, weights were set to the total number of leaves on the individual; family was set to binomial in models of the proportion of leaves infected, and to quasipoisson in models of the number of leaves infected (to counter overdispersion). In each model, MYC, POP/GEN and SITE (and SIZE at the corresponding time point, in applicable models) were included as explanatory factors, as well as all interaction terms.

**Are growth and defensive effects from mycorrhizal inoculation linked in host genotypes?** To test for a relationship between

growth and defensive effects following inoculation with arbuscular mycorrhizal fungi in the maternal genotypes, we first used the negative binomial GLM examining differences in host size (i.e. the model built in the first section of the statistical methods) to calculate the estimated effect of mycorrhizal inoculation on host growth in each maternal genotype. Host growth effects were estimated at the beginning of the experiment to avoid the effects of pathogen exposure and variable field conditions. Effect sizes were obtained using the LSMEANS package (Lenth, 2016) and were averaged over the levels of field epidemic population. Second, we used the generalized logistic regression model examining differences in the number of infected leaves in infected hosts (i.e. the model built in the third section of the statistical methods) to calculate the estimated effect of mycorrhizal inoculation on host defense in each maternal genotype. Defensive effects were quantified at the epidemic peak to capture the highest infection rates. We then built a linear regression model linking defense effects (as the response variable) and growth effects in each genotype.

**Are disease susceptibility and defensive effects from mycorrhizal inoculation linked in host genotypes?** We tested for a relationship between host disease susceptibility and defensive effects following inoculation with arbuscular mycorrhizal fungi in the host genotypes. First, we quantified disease susceptibility in each genotype in the absence of the mutualist (i.e. in NM plants only) using the estimated coefficients from the model examining the number of infected leaves at the epidemic peak (i.e. the same model as used in the section earlier). Genotype-level coefficients were obtained using the LSMEANS package (Lenth, 2016) and were averaged over the levels of field epidemic population. Second, defensive effects resulting from inoculation with mycorrhizal fungi were quantified for each maternal genotype as the effect of mycorrhizal inoculation from these same models (i.e. as the change in the coefficient when the genotype was inoculated). Finally, we built a linear regression model linking defense effect sizes (as the response variable) and disease susceptibility in each genotype.

## Results

### Do growth effects resulting from mycorrhizal inoculation vary among host populations and maternal genotypes?

Inoculation with arbuscular mycorrhizal fungi produced growth benefits in experimental plants from nearly every host genetic origin. Before pathogen exposure in field conditions, host growth was determined by mycorrhizal inoculation and seed origin population (Fig. S2; Table S1; MYC,  $P < 0.001$ ; POP,  $P < 0.001$ ,  $n = 287$  plants), as well as by an interaction between mycorrhizal inoculation and host maternal genotype (Fig. 1a, left panel; Table S1; MYC  $\times$  GEN,  $P = 0.002$ ). At the end of field epidemic experiment, mycorrhizal inoculation (Fig. 1b,c; Tables S2, S3; MYC,  $P < 0.001$ ), seed origin population (Fig. 1b; Table S2; POP,  $P < 0.001$ ) and maternal genotype (Fig. 1c; Table S3; GEN,  $P < 0.001$ ) continued to be tightly linked to host growth, but the interaction between mycorrhizal inoculation and


maternal genotype had disappeared. Host initial size also predicted host final size (Table S2; SIZE,  $P < 0.001$ ). Mycorrhizal inoculation also increased the aboveground dry biomass of plants at the end of the experiment (Fig. S3; Table S4; MYC,  $P < 0.001$ ;  $W = 11$ ,  $n = 41$  plants).

### Does inoculation with mycorrhizal fungi influence host infection rate?

Upon pathogen exposure in the field epidemics, host infection rate was influenced by several factors whose importance shifted over time. Host infection rates reached 76% at the peak of the epidemic but fell to 34% by the end of the experiment (Fig. 2a). Host infection rate also varied among the three field epidemic populations, with the effect of field site strengthening over time (Fig. 2b–d; Tables S5–S8; SITE (peak of epidemic),  $P = 0.03$  (POP),  $P = 0.01$  (GEN); end of experiment,  $P < 0.001$  (POP/GEN)). Although inoculation with arbuscular mycorrhizal fungi marginally influenced host infection rate during the peak of the epidemic (Fig. 2c; Tables S5, S6; MYC,  $P = 0.09$  (POP),  $P = 0.11$  (GEN),  $n = 287$  plants), at the end of experiment AMF plants were slightly more likely to be infected than NM plants (Fig. 2d; Tables S7, S8; MYC,  $P = 0.04$  (POP),  $P = 0.05$  (GEN),  $n = 286$  plants). The effect of mycorrhizal inoculation on host infection rates was weaker than the effect on host growth. Host size also influenced host infection at the end of the experiment: larger plants were more likely to be infected (Tables S7, S8; SIZE,  $P = 0.06$  (POP),  $P = 0.04$  (GEN)).

### Upon infection, does previous inoculation with mycorrhizal fungi affect disease severity?

Among infected individuals, disease severity was influenced by several factors, including mycorrhizal inoculation and host genetic origin. AMF plants had lower proportions of their leaves infected relative to NM plants throughout the experiment (Fig. 3a,b; Tables S9–S12; MYC,  $P < 0.001$ ,  $n = 210$  plants (peak of epidemic) and  $P = 0.003$ ,  $n = 98$  plants (end of experiment)). The proportion of leaves infected also varied among seed origin populations (Fig. 3a; Table S9; POP,  $P = 0.004$ ) and maternal genotypes (Table S10; GEN,  $P = 0.02$ ) during the peak of the epidemic but was similar among host genetic origins by the end of the experiment (Tables S11, S12). At the epidemic peak, mycorrhizal inoculation and maternal genotype interacted to determine the number of infected leaves on infected hosts (Fig. 3c; Table S13; MYC  $\times$  GEN,  $P = 0.05$ ), with both positive and negative effects occurring; maternal genotype also interacted with field epidemic population (Table S13; MYC  $\times$  SITE,  $P = 0.004$ ). By contrast, seed origin population did not interact with other variables and had a stronger influence during the peak of the epidemic (Tables S14, S15; POP,  $P = 0.005$  (peak of epidemic) and  $P = 0.08$  (end of experiment)). By the end of the experiment, AMF plants had higher numbers of infected leaves than NM plants (Fig. 3d; Table S16; MYC,  $P = 0.04$ ). Like the effect on host infection rates, the effect of mycorrhizal inoculation on disease severity was weaker than the effect on host growth. Field



**Fig. 2** Host infection rate was increased in mycorrhizal-inoculated plants and varied among field epidemic populations. Infection rates by *Podosphaera plantaginis* in experimental *Plantago lanceolata* plants increased over time in each mycorrhizal treatment (a) and field epidemic population (b) until the peak of the epidemic, then fell near the end of the experiment. Generalized logistic regression models showed that though host infection rate was marginally increased in arbuscular mycorrhizal-inoculated plants (AMF) relative to nonmycorrhizal plants (NM) at the peak of the epidemic (c; Tables S5, S6; mycorrhizal inoculation (MYC),  $P = 0.09$ ,  $n = 287$  plants); at the end of the experiment host infection rates were increased in AMF relative to NM plants in all three field epidemic populations (Tables S7, S8; MYC,  $P = 0.04$ ,  $n = 286$  plants). Host infection rates also varied among the three field epidemic populations throughout the experiment (c, d; Tables S5–S8; field epidemic population,  $P = 0.01$  (peak) and  $P < 0.001$  (end),  $n = 286$  plants). In (b), missing data as a result of heavy rains on days 15–20 are indicated by light gray solid lines. In (c, d), predicted values resulting from generalized logistic models are plotted; black dots represent outlier individuals, box notches represent a 95% confidence interval for comparing group medians, box hinges correspond to the 1<sup>st</sup> and 3<sup>rd</sup> quartiles, and box whiskers extend to the largest and smallest value no further than  $1.5 \times$  the interquartile range from the hinges. Myc., mycorrhizal; Non-Myc., nonmycorrhizal.

epidemic population also influenced the number of infected leaves in hosts at the end of the experiment (Fig. 3d; Table S15; SITE,  $P = 0.02$ ). Host size predicted the proportion of infected leaves in infected individuals at the peak of the epidemic (Tables S9, S10; SIZE,  $P < 0.001$ ) and the number of infected leaves in infected individuals throughout the experiment (Tables S13–S16; SIZE,  $P \leq 0.01$ ).

### Are growth and defensive effects from mycorrhizal inoculation linked in host genotypes?


There was a marginally significant relationship between the growth and defensive effects conferred by inoculation with arbuscular mycorrhizal fungi across the host maternal genotypes at the end of the experiment (Fig. 4a; Table S17;  $P = 0.11$ ,  $n = 30$

maternal genotypes). Host genotypes that experienced more positive growth effects as a result of inoculation with mycorrhizal fungi (i.e. higher increases in leaf number in AMF relative to NM plants) experienced slightly more negative defensive outcomes (i.e. higher increases in the number of infected leaves in AMF relative to NM plants).


### Are disease susceptibility and defensive effects from mycorrhizal inoculation linked in host genotypes?

We found a strong relationship between host disease susceptibility and the magnitude of the defensive effect received from mycorrhizal inoculation in the maternal genotypes (Fig. 4b; Table S18;  $P < 0.001$ ,  $n = 30$  maternal genotypes). At the peak of the epidemic, host genotypes that suffered more severe infections





**Fig. 3** Among infected plants, disease severity varied among mycorrhizal treatments, seed origin populations, and field epidemic sites. Generalized logistic regression models show that previous inoculation with a three-species mixture of arbuscular mycorrhizal fungi was linked to reductions in the proportion of leaves infected by *Podosphaera plantaginis* in infected individuals of *Plantago lanceolata*, both at the peak of the epidemic (a; Tables S9, S10; mycorrhizal inoculation (MYC),  $P < 0.001$ ,  $n = 210$  plants) and at the end of the experiment (b; Tables S11, S12; MYC,  $P = 0.003$ ,  $n = 98$  plants). Mycorrhizal inoculation and maternal genotype also interacted to determine the number of infected leaves on infected plants at the peak of the epidemic, with both negative and positive effects of mycorrhizas on disease (c; Table S13; MYC  $\times$  maternal genotype,  $P = 0.05$ ). At the end of the experiment, infected mycorrhizal-inoculated plants had more infected leaves than did infected nonmycorrhizal plants (d; Table S16; MYC,  $P = 0.04$ ). Seed origin population (a; Table S9; seed origin population,  $P = 0.004$ ) and field epidemic population (d; Tables S13, S15, S16; field epidemic population,  $P < 0.02$ ) also influenced host disease severity. In (a, b, d), predicted values resulting from generalized logistic models are plotted; black dots represent outlier individuals, box hinges correspond to the 1<sup>st</sup> and 3<sup>rd</sup> quartiles, and box whiskers extend to the largest and smallest values no further than  $1.5 \times$  the interquartile range from the hinges. In (c), error bars represent 95% confidence interval. Myc., mycorrhizal; Non-Myc., nonmycorrhizal.



**Fig. 4** Growth and disease susceptibility are linked to defensive effects from arbuscular mycorrhizal fungi (AMF) in the host genotypes. (a) In a field epidemic experiment with *Plantago lanceolata* individuals from 30 maternal genotypes, linear regression models show that host genotypes that grew larger when inoculated with a three-species mixture of AMF experienced marginally more negative infection outcomes from exposure to *Podosphaera plantaginis* (Table S17;  $P = 0.11$ ,  $n = 30$  genotypes). (b) In the same experiment, host genotypes that were more susceptible to infection (when not inoculated with mycorrhizas) received stronger disease protection effects from inoculation with mycorrhizal fungi (Table S18;  $P < 0.001$ ,  $n = 30$  genotypes). In (a, b), each point represents one maternal genotype, originating from one of six seed origin populations. The y-axis represents defensive effects as a result of mycorrhizas, that is, the estimated change in the mean number of infected leaves in each genotype as a result of mycorrhizal inoculation. The shaded area represents a 95% confidence interval. In (a), the x-axis represents growth effects as a result of mycorrhizas, that is, the estimated change in mean host size (leaf number) in each genotype as a result of mycorrhizal inoculation. Changes in host size were estimated before pathogen exposure in variable field conditions. In (b), the x-axis represents host disease susceptibility, that is, the estimated mean number of infected leaves in infected nonmycorrhizal (NM) plants in each genotype at the peak of the epidemic: more positive x-axis values indicate more severe infections in the absence of the mutualist.

(i.e. had more infected leaves) when not inoculated with mycorrhizal fungi experienced greater reductions in disease severity (i.e. in the number of infected leaves) when inoculated with mycorrhizal fungi.

## Discussion

While both plant-associated pathogenic and mutualistic microbes are ubiquitous across ecosystems, how they interact to determine patterns of infection in genetically diverse host populations is not known. In a field experiment placing wild hosts of 30 maternal genotypes in naturally occurring pathogen epidemics, we found that inoculation with a three-species mixture of arbuscular mycorrhizal fungi produced benefits and risks that influenced above-ground patterns of host infection. Mycorrhizal inoculation increased growth in hosts of nearly every genotype, but also increased infection rates from a foliar fungal pathogen. The effects of mycorrhizal inoculation on disease severity varied over the course of the epidemic, with both protective and negative effects occurring among the host genotypes. Moreover, disease susceptibility and mycorrhiza-derived defense effects appeared to be linked in the host genotypes: more susceptible host genotypes (in the absence of the mutualists) received stronger protection against disease when inoculated with mycorrhizal fungi. Arbuscular mycorrhizal fungi have been shown to protect agricultural

plants against belowground (Azcón-Aguilar & Barea, 1996; Hao *et al.*, 2005) and foliar (Fiorilli *et al.*, 2018; Pozo de la Hoz *et al.*, 2021) pathogens in controlled laboratory conditions, but our results provide the first evidence of how microbial mutualists can shift patterns of host infection in genetically diverse wild populations under pathogen attack. Mycorrhizal fungi also appeared to be linked to a growth–defense trade-off in the hosts: mycorrhizal-inoculated plants grew larger and became more infected by the pathogen, with the host genotypes that obtained the greatest growth benefit from mycorrhizal fungi also suffering the largest increases in disease. Together, our results underscore that under natural ecological and epidemic conditions mycorrhizal fungi produce a complex and temporally variable array of positive and negative effects on host growth and infection.

Inoculation with mycorrhizal fungi provided growth benefits to host plants from every population and maternal genotype. The magnitude of these effects varied among hosts from different genetic origins before exposure to the field epidemics, but homogenized over time in field conditions. Owing to the positive relationship between plant size and survival, growth is often intrinsically linked to host fitness (Harper & White, 1974). Our results are consistent with studies reporting evidence for the importance of mycorrhizal fungi in determining host growth and/or fitness and showing variability in such effects among host genotypes (Rasmussen *et al.*, 2017; Qin *et al.*, 2021). In addition,

susceptibility to the pathogen epidemics in the field varied among hosts of different genetic origins, consistent with other studies in this pathosystem (Laine, 2005, 2007a; Tack *et al.*, 2014; Susi & Laine, 2015) and the broader wild plant disease literature (Carlsson-Granér, 1997; Price *et al.*, 2004). Together, these results provide evidence for the importance of genotype in mediating host–microbe interactions (Eck *et al.*, 2019; Sallinen *et al.*, 2020), although more generalist interactions can also certainly occur (Gilbert & Webb, 2007; Halbritter *et al.*, 2012; Hersh *et al.*, 2012).

Building upon such studies, we show evidence of changes in host–parasite interactions as a result of prior inoculation with microbial mutualists. Inoculation experiments with agricultural plant species and their pathogens have shown that arbuscular mycorrhizal fungi can reduce disease incidence or severity (Norman *et al.*, 1996; Pozo *et al.*, 2002; Hao *et al.*, 2005; Li *et al.*, 2010; Song *et al.*, 2015; Berdeni *et al.*, 2018). However, symbiotic relationships in cultivars may differ from those of wild plants (Xing *et al.*, 2012); thus, it is not straightforward to predict responses in wild populations from controlled agricultural trials. In this study, mycorrhizal inoculation increased infection rates in hosts of a wild plant species and produced variable effects on disease severity in hosts of different genotypes and over the course of the epidemic. By the end of the experiment, mycorrhizal inoculation was weakly linked to higher numbers of infected leaves in diseased hosts (though variation among genotypes remained). Our results suggest that environmental, temporal and genetic contexts may alter the potential defensive effects related to mycorrhizal association and are consistent with other experimental studies showing that mycorrhizal effects on host infection may vary among host genotypes (Mark & Cassells, 1996; Steinkellner *et al.*, 2012). However, in our experiment, the effects of mycorrhizal inoculation on infection were weaker than the effects on host growth. Additional studies that explicitly quantify mycorrhizal colonization are needed to confirm the contribution of the symbiont to host growth, infection and fitness. The potential risks of mycorrhizal association in wild plants are relevant for theoretical studies which speculate as to how mycorrhizal fungi might influence host fitness in the presence of pathogens and affect plant population and community dynamics (Bachelot *et al.*, 2015).

In many free-living organisms, an evolutionary trade-off exists between growth and defense (Herms & Mattson, 1992). In our experiment, additional insights can be gained by examining whether changes in host growth as a result of mutualism are related to changes in host defense across genotypes. In our experiment, mycorrhizal-inoculated hosts consistently grew larger and were more likely to become infected by the pathogen. This could occur if increases in host size increase pathogen encounter rates, as might be expected for pathogen species with wind- or passively dispersed spores (such as *P. plantaginis*). In addition, host genotypes that experienced larger growth benefits from mycorrhizal inoculation suffered marginally larger increases in disease severity (although there was considerable variation in this effect). That some host genotypes grew larger and experienced reductions in disease severity following mycorrhizal inoculation suggests that defense priming could occur occasionally. Infected AMF plants

also had lower proportions of infected leaves than did NM plants (relative to their size), although it is unclear whether this may offset the costs of having higher numbers of infected leaves in this pathosystem. Changes in host tolerance to pathogens following mycorrhizal inoculation could also explain increases in infected leaf numbers, although mycorrhizal fungi are generally expected to improve host nutritional status or increase leaf toughness (Meier & Hunter, 2018), making foliar pathogen spread more difficult. Thus, it is likely that differences in host infection between AMF and NM plants in our experiment were mediated by increases in host size (as host size also predicted some aspects of infection) or defense priming in some genotypes.

In addition, our findings indicate that arbuscular mycorrhizal fungi could help susceptible host genotypes to compensate for lack of innate resistance while placing costs on well-defended genotypes. We found that the magnitude of defense effects following mycorrhizal inoculation were linked to pathogen susceptibility in the host genotypes: host genotypes that were susceptible to more severe infections received stronger disease protection effects when inoculated with the mutualist. By contrast, more resistant host genotypes were likely to experience slight increases in disease load when inoculated. Thus, inoculation with mycorrhizal fungi tended to equalize disease severity between more resistant and more susceptible host genotypes, potentially reducing the relative importance of host genetic resistance in determining pathogen effects. However, the linkages between disease susceptibility and mycorrhizal defense effects in our study (as well as between host growth and these effects) were revealed *post hoc* and should be confirmed with experiments designed to test these hypotheses. If confirmed, it could indicate that host genotypes may experience trade-offs in investment in genetic resistance vs mycorrhizal-mediated resistance. It could also suggest that mycorrhizal association may have been selected for and maintained in host populations partially because it increases the fitness of susceptible host genotypes. In this way, mycorrhizal protective effects could contribute to the maintenance of diversity in host genetic resistance within and among populations (Laine, 2004, 2007a; Jousimo *et al.*, 2014).

Variation among host populations and genotypes in mycorrhizal benefits and risks could be a result of several factors. These factors include intraspecific differences in hosts' ability to form associations with and derive function from different mycorrhizal species, differences in mycorrhizal colonization rates or community composition, environmental variation or differences in host–pathogen population dynamics over time. Previous studies demonstrated variation in arbuscular mycorrhizal colonization rates among individuals within species – variation that is thought to have a partially genetic component (Plouznikoff *et al.*, 2019; Pawlowski *et al.*, 2020). Although we observed clear differences in host growth and infection as a result of the mycorrhizal inoculation treatment, data on mycorrhizal colonization are needed to confirm mycorrhizas as the mechanism underlying the observed effects. Environmental conditions may also impact plant–microbe interactions (Santoyo *et al.*, 2017). Consistent with other studies in this pathosystem, infection rates varied among field populations and changed over time (Eck *et al.*, 2022). By contrast, the effects of mycorrhizal inoculation on hosts were similar

among field populations, although changes over time also occurred. Growth benefits following inoculation with mycorrhizal fungi varied among host populations and genotypes before pathogen exposure but became homogenous over time in the field conditions. Changes in host infection rate and disease severity as a result of mycorrhizal inoculation were also similar among field populations but increased over time. Together, these results suggest that mycorrhizal effects on hosts are more temporally than environmentally sensitive. There is also some chance that environmental mycorrhizal spores could have come into contact with our experimental soils, such that true amounts of mycorrhizal colonization in the nonmycorrhizal treatment could be low (rather than none), and the mycorrhizal communities in the experimental pots could contain species that were not inoculated. The limited duration of the field experiment reduces the likelihood that this could cause strong effects (Sanders & Sheikh, 1983); however, if it occurred, it should have occurred evenly among treatments and seed sources. Future studies in which mycorrhizal composition, colonization rates and function are quantified in genetically diverse hosts and in variable environmental conditions over time are necessary to corroborate this work and disentangle the effect of mycorrhizal growth and defensive effects from intraspecific variation in host growth and resistance.

Together, our results suggest that symbiosis with arbuscular mycorrhizal fungi produces benefits and alters infection risks from a pathogen during natural epidemics in genetically variable host populations. Altered patterns of host growth and infection as a result of mutualist species may cascade to affect patterns of abundance, diversity and distribution of the associated organisms, as well as ecosystem processes (Brown *et al.*, 2001). Moreover, we are beginning to acknowledge the importance of direct and indirect microbial interactions within hosts in determining host fitness and parasite population dynamics (Kemen, 2014; Kemen *et al.*, 2015; Kroll *et al.*, 2017). Belowground soil and rhizosphere processes may also affect aboveground interactions with pathogens and herbivores, and vice versa (van der Putten *et al.*, 2001; Wardle *et al.*, 2004; Frew, 2021), and mutualists and parasites may act simultaneously to determine host fitness (Bezemer & van Dam, 2005). Our results also highlight the importance of characterizing host–microbe interactions under natural conditions and temporal interaction sequences, which will allow more precise modeling of epidemiological and ecological community dynamics. In addition, if growth and defensive effects as a result of mutualism are common, they should ultimately affect the coevolutionary trajectories of the associated organisms (van Dam & Heil, 2010). Understanding how mutualism alters host susceptibility and parasite interactions will be important for understanding, predicting and managing disease in ecological communities and in agriculture.

## Acknowledgements

We would like to thank the University of Oulu Botanical Gardens for providing glasshouse facilities, Riitta Ovaska for

assistance during the field experiment, Dr Mauritz Vestberg for the fungal isolates, and the anonymous reviewers for their constructive feedback. This work was funded by grants from the European Research Council (ERC-2016-COG RESISTANCE, LS8) and the Academy of Finland (MULTA #327222 and #296686) to A-LL, and the University of Zurich (Forschungskredit FK-20-108 and FK-21-118) to JLE. Open access funding provided by Universitat Zurich.

## Author contributions

A-LL and M-MK designed and conducted the experiment. JLE and A-LL analyzed the data. JLE wrote the first draft of the manuscript. All authors contributed to and approved the final version.

## ORCID

Jenalle L. Eck  <https://orcid.org/0000-0001-5664-7450>  
Minna-Maarit Kytöviita  <https://orcid.org/0000-0002-8928-6951>  
Anna-Liisa Laine  <https://orcid.org/0000-0002-0703-5850>

## Data availability

The data that support the findings of this study are available in the supplementary material of this article.

## References

- Azcón-Aguilar C, Barea JM. 1996. Arbuscular mycorrhizas and biological control of soil-borne plant pathogens – an overview of the mechanisms involved. *Mycorrhiza* 6: 457–464.
- Bachelot B, Uriarte M, McGuire K. 2015. Interactions among mutualism, competition, and predation foster species coexistence in diverse communities. *Theoretical Ecology* 8: 297–312.
- Barea JM, Azcón R, Azcón-Aguilar C. 2002. Mycorrhizosphere interactions to improve plant fitness and soil quality. *Antonie Van Leeuwenhoek* 81: 343–351.
- Berdeni D, Cotton TEA, Daniell TJ, Bidartondo MI, Cameron DD, Evans KL. 2018. The effects of arbuscular mycorrhizal fungal colonization on nutrient status, growth, productivity, and canker resistance of apple (*Malus pumila*). *Frontiers in Microbiology* 9: 1461.
- Bever JD, Mangan SA, Alexander HM. 2015. Maintenance of plant species diversity by pathogens. *Annual Review of Ecology, Evolution, and Systematics* 46: 305–325.
- Bezemer TM, van Dam NM. 2005. Linking aboveground and belowground interactions via induced plant defenses. *Trends in Ecology and Evolution* 20: 617–624.
- Bos M. 1992. Gene flow characters and population structure in *Plantago lanceolata*. In: Kuiper PJC, Bos M, eds. *PLANTAGO: a multidisciplinary study*. Berlin, Germany: Springer-Verlag, 222–231.
- Brown JH, Whitham TG, Ernest SKM, Gehring CA. 2001. Complex species interactions and the dynamics of ecological systems: long-term experiments. *Science* 293: 643–650.
- Brown JKM. 2003. A cost of disease resistance: paradigm or peculiarity? *Trends in Genetics* 19: 667–671.
- Brownlie JC, Johnson KN. 2009. Symbiont-mediated protection in insect hosts. *Trends in Microbiology* 17: 348–354.
- Burdon JJ, Thrall PH. 2014. What have we learned from studies of wild plant–pathogen associations? – the dynamic interplay of time, space and life-history. *European Journal of Plant Pathology* 138: 417–429.

- Bushnell WR. 2002. The role of powdery mildew research in understanding host–pathogen interaction: past, present, and future. In: Bélanger RR, Bushnell WR, Dik AJ, Carver LW, eds. *The powdery mildews – a comprehensive treatise*. St Paul, MN, USA: The American Phytopathology Society, 1–12.
- Cameron DD, Neal AL, van Wees SCM, Ton J. 2013. Mycorrhiza-induced resistance: more than the sum of its parts? *Trends in Plant Science* 18: 539–545.
- Carlsson-Granér U. 1997. Anther-smut disease in *Silene dioica*: variation in susceptibility among genotypes and populations, and patterns of disease within populations. *Evolution* 51: 1416–1426.
- van Dam NM, Heil M. 2010. Multitrophic interactions below and above ground: *en route* to the next level. *Journal of Ecology* 99: 77–88.
- Delavaux CS, Smith-Ramesh LM, Kuebbing SE. 2017. Beyond nutrients: a meta-analysis of the diverse effects of arbuscular mycorrhizal fungi on plants and soils. *Ecology* 98: 2111–2119.
- Eck JL, Barrès B, Soubeyrand S, Sirén J, Numminen E, Laine A-L. 2022. Strain diversity and spatial distribution are linked to epidemic dynamics in host populations. *The American Naturalist* 199: 59–74.
- Eck JL, Stump SM, Delavaux CS, Mangan SA, Comita LS. 2019. Evidence of within-species specialization by soil microbes and the implications for plant community diversity. *Proceedings of the National Academy of Sciences, USA* 116: 7371–7376.
- Fiorilli V, Vannini C, Ortolani F, Garcia-Seco D, Chiapello M, Novero M, Domingo G, Terzi V, Morcia C, Bagnaresi P *et al.* 2018. Omics approaches revealed how arbuscular mycorrhizal symbiosis enhances yield and resistance to leaf pathogen in wheat. *Scientific Reports* 8: 9625.
- Frew A. 2021. Aboveground herbivory suppresses the arbuscular mycorrhizal symbiosis, reducing phosphorus uptake. *Applied Soil Ecology* 168: 104133.
- Gilbert GS, Webb CO. 2007. Phylogenetic signal in plant pathogen–host range. *Proceedings of the National Academy of Sciences, USA* 104: 4979–4983.
- Goddard M-L, Belval L, Martin I, Roth L, Laloue H, Deglene-Benbrahim L, Valat L, Bertsch C, Chong J. 2021. Arbuscular mycorrhizal symbiosis triggers major changes in primary metabolism together with modification of defense responses and signaling in both roots and leaves of *Vitis vinifera*. *Frontiers in Plant Science* 12: 721614.
- Gosling P, Hodge A, Goodlass G, Bending GD. 2006. Arbuscular mycorrhizal fungi and organic farming. *Agriculture, Ecosystems, and Environment* 113: 17–35.
- Grman E. 2012. Plant species differ in their ability to reduce allocation to non-beneficial arbuscular mycorrhizal fungi. *Ecology* 93: 711–718.
- Hafer-Hahmann NC, Vorburger C. 2021. Positive association between the diversity of symbionts and parasitoids of aphids in field populations. *Ecosphere* 12: e3355.
- Halbritter AH, Carroll GC, Gusewell S, Roy BA. 2012. Testing assumptions of the enemy release hypothesis: generalist versus specialist enemies of the grass *Brachypodium sylvaticum*. *Mycologia* 104: 34–44.
- Hanski I. 1999. *Metapopulation ecology*. Oxford, UK: Oxford University Press.
- Hao Z, Christie P, Qin L, Wang C, Li X. 2005. Control of fusarium wilt of cucumber seedlings by inoculation with an arbuscular mycorrhizal fungus. *Journal of Plant Nutrition* 28: 1961–1974.
- Harper JL, White J. 1974. The demography of plants. *Annual Review of Ecology and Systematics* 5: 419–463.
- Hartnett DC, Wilson GWT. 2002. The role of mycorrhizas in plant community structure and dynamics: lessons from grasslands. In: Smith SE, Smith FA, eds. *Diversity and integration in mycorrhizas. Developments in plant and soil sciences, vol. 94*. Dordrecht, the Netherlands: Springer, 319–331.
- van der Heijden MGA, Bardgett RD, van Straalen NM. 2008. The unseen majority: soil microbes as drivers of plant diversity and productivity in terrestrial ecosystems. *Ecology Letters* 11: 296–310.
- Hermes DA, Mattson WJ. 1992. The dilemma of plants: to grow or defend. *The Quarterly Review of Biology* 67: 283–335.
- Hersh MH, Vilgalys R, Clark JS. 2012. Evaluating the impacts of multiple generalist pathogens on temperate tree seedling survival. *Ecology* 93: 511–520.
- Hoeksema JD, Chaudhary VB, Gehring CA, Johnson NC, Karst J, Koide RT, Pringle A, Zabinski C, Bever JD, Moore JC *et al.* 2010. A meta-analysis of context-dependency in plant response to inoculation with mycorrhizal fungi. *Ecology Letters* 13: 394–407.
- Hohmann P, Messmer MM. 2017. Breeding for mycorrhizal symbiosis: focus on disease resistance. *Euphytica* 213: 1–11.
- Johansson JF, Paul LR, Finlay RD. 2004. Microbial interactions in the mycorrhizosphere and their significance for sustainable agriculture. *FEMS Microbiology Ecology* 48: 1–13.
- Johnson D, Vandenkoornhuyse PJ, Leake JR, Gilbert L, Booth RE, Grime JP, Young JPW, Read DJ. 2003. Plant communities affect arbuscular mycorrhizal fungal diversity and community composition in grassland microcosms. *New Phytologist* 161: 503–515.
- Johnson NC, Graham JH, Smith FA. 1997. Functioning of mycorrhizal associations along the mutualism–parasitism continuum. *New Phytologist* 135: 575–585.
- Jones MD, Smith SE. 2004. Exploring functional definitions of mycorrhizas: are mycorrhizas always mutualisms? *Canadian Journal of Botany* 82: 1089–1109.
- Jousimo J, Tack AJM, Ouskainen O, Mononen T, Susi H, Tollenaere C, Laine A-L. 2014. Ecological and evolutionary effects of fragmentation on infectious disease dynamics. *Science* 344: 1289–1293.
- Jung SC, Martinez-Medina A, Lopez-Raez JA, Pozo MJ. 2012. Mycorrhiza-induced resistance and priming of plant defenses. *Journal of Chemical Ecology* 38: 651–664.
- Kemen AC, Agler MT, Kemen E. 2015. Host–microbe and microbe–microbe interactions in the evolution of obligate plant parasitism. *New Phytologist* 206: 1207–1228.
- Kemen E. 2014. Microbe–microbe interactions determine oomycete and fungal host colonization. *Current Opinion in Plant Biology* 20: 75–81.
- King KC, Brockhurst MA, Vasieva O, Paterson S, Betts A, Ford SA, Frost CL, Horsburgh MJ, Haldenby S, Hurst GDD. 2016. Rapid evolution of microbe-mediated protection against pathogens in a worm host. *The ISME Journal* 10: 1915–1924.
- Klironomos J. 2003. Variation in plant response to native and exotic arbuscular mycorrhizal fungi. *Ecology* 84: 2292–2301.
- Koide RT, Dickie IA. 2002. Effects of mycorrhizal fungi on plant populations. *Plant and Soil* 244: 307–317.
- Koricheva J, Gange AC, Jones T. 2009. Effects of mycorrhizal fungi on insect herbivores: a meta-analysis. *Ecology* 90: 2088–2097.
- Kroll S, Agler MT, Kemen E. 2017. Genomic dissection of host–microbe and microbe–microbe interactions for advanced plant breeding. *Current Opinion in Plant Biology* 36: 71–78.
- Laine A-L. 2004. Resistance variation within and among host populations in a plant pathogen metapopulation: implications for regional pathogen dynamics. *Journal of Ecology* 92: 990–1000.
- Laine A-L. 2005. Spatial scale of local adaptation in a plant–pathogen metapopulation. *Journal of Evolutionary Biology* 18: 930–938.
- Laine A-L. 2006. Evolution of host resistance: looking for coevolutionary hotspots at small spatial scales. *Proceedings: Biological Sciences* 273: 267–273.
- Laine A-L. 2007a. Detecting local adaptation in a natural plant–pathogen metapopulation: a laboratory vs. field transplant approach. *Journal of Evolutionary Biology* 20: 1665–1673.
- Laine A-L. 2007b. Pathogen fitness components and genotypes differ in their sensitivity to nutrient and temperature variation in a wild plant–pathogen association. *Journal of Evolutionary Biology* 20: 2371–2378.
- Laine A-L. 2008. Temperature-mediated patterns of local adaptation in a natural plant–pathogen metapopulation. *Ecology Letters* 11: 327–337.
- Laine A-L, Burdon JJ, Dodds PN, Thrall PH. 2011. Spatial variation in disease resistance: from molecules to metapopulations. *Journal of Ecology* 99: 96–112.
- Laine A-L, Hanski I. 2006. Large-scale dynamics of a specialist plant pathogen in a fragmented landscape. *Journal of Ecology* 94: 217–226.
- Lenth RV. 2016. Least-squares means: the R package LSMEANS. *Journal of Statistical Software* 69: 1–33.
- Li Y, Yanagi A, Miyawaki Y, Okada T, Matsubara Y. 2010. Disease tolerance and changes in antioxidative abilities in mycorrhizal strawberry plants. *Journal of the Japanese Society for Horticultural Science* 79: 174–178.
- Mark GL, Cassells AC. 1996. Genotype-dependence in the interaction between *Glomus fistulosum*, *Phytophthora fragariae* and the wild strawberry (*Fragaria vesca*). *Plant & Soil* 185: 233–239.
- May G, Nelson P. 2014. Defensive mutualisms: do microbial interactions within hosts drive the evolution of defensive traits? *Functional Ecology* 28: 356–363.

- Meier AR, Hunter MD. 2018. Arbuscular mycorrhizal fungi mediate herbivore-induction of plant defenses differently above and belowground. *Oikos* 127: 1759–1775.
- Newsham KK, Fitter AH, Watkinson AR. 1995. Arbuscular mycorrhiza protect an annual grass from root pathogenic fungi in the field. *Journal of Ecology* 83: 991–1000.
- Norman JR, Atkinson D, Hooker JE. 1996. Arbuscular mycorrhizal fungal-induced alteration to root architecture in strawberry and induced resistance to the root pathogen *Phytophthora fragariae*. *Plant and Soil* 185: 191–198.
- Oliver KM, Smith AH, Russell JA. 2014. Defensive symbiosis in the real world – advancing ecological studies of heritable, protective bacteria in aphids and beyond. *Functional Ecology* 28: 341–355.
- Öpik M, Moora M, Liira J, Zobel M. 2006. Composition of root-colonizing arbuscular mycorrhizal fungal communities in different ecosystems around the globe. *Journal of Ecology* 94: 778–790.
- Ovaskainen O, Laine A-L. 2006. Inferring evolutionary signals from ecological data in a plant–pathogen metapopulation. *Ecology* 87: 880–891.
- Pawlowski ML, Vuong TD, Valliyodan B, Nguyen HT, Hartman GL. 2020. Whole genome resequencing identifies quantitative trait loci associated with mycorrhizal colonization of soybean. *Theoretical and Applied Genetics* 133: 409–417.
- Penczykowski R, Walker E, Soubeyrand S, Laine A-L. 2015. Linking winter conditions to large-scale disease dynamics in a wild plant–pathogen metapopulation. *New Phytologist* 205: 1142–1152.
- Plouznikoff K, Asins MJ, Dupré de Boulois H, Carbonell EA, Declerck S. 2019. Genetic analysis of tomato root colonization by arbuscular mycorrhizal fungi. *Annals of Botany* 124: 933–946.
- Polin S, Simon J-C, Outreman Y. 2014. An ecological cost associated with protective symbionts of aphids. *Ecology and Evolution* 4: 836–840.
- Pozo de la Hoz J, Rivero J, Azcón-Aguilar C, Urrestarazu M, Pozo MJ. 2021. Mycorrhiza-induced resistance against foliar pathogens is uncoupled of nutritional effects under different light intensities. *Journal of Fungi* 7: 402.
- Pozo MJ, Azcón-Aguilar C. 2007. Unraveling mycorrhiza-induced resistance. *Current Opinion in Plant Biology* 10: 393–398.
- Pozo MJ, Cordier C, Dumas-Gaudot E, Gianinazzi S, Barea JM, Azcón-Aguilar C. 2002. Localized versus systemic effect of arbuscular mycorrhizal fungi on defence responses to *Phytophthora* infection in tomato plants. *Journal of Experimental Botany* 53: 525–534.
- Pretty J, Sutherland WJ, Ashby J, Auburn J, Baulcombe D, Bell M, Bentley J, Bickersteth S, Brown K, Burke J *et al.* 2011. The top 100 questions of importance to the future of global agriculture. *International Journal of Agricultural Sustainability* 8: 219–236.
- Price JS, Bever JD, Clay K. 2004. Genotype, environment, and genotype by environment interactions determine quantitative resistance to leaf rust (*Coleosporium asterum*) in *Euthamia graminifolia* (Asteraceae). *New Phytologist* 162: 729–743.
- van der Putten WH, Vet LEM, Harvey JA, Wäckers FL. 2001. Linking above- and belowground multitrophic interactions of plants, herbivores, pathogens, and their antagonists. *Trends in Ecology and Evolution* 16: 547–554.
- Qin J, Geng Y, Li X, Zhang C, Zhao X, von Gadow K. 2021. Mycorrhizal type and soil pathogenic fungi mediate tree survival and density dependence in a temperate forest. *Forest Ecology and Management* 496: 119459.
- Qu L, Wang M, Biere A. 2021. Interactive effects of mycorrhizae, soil phosphorus, and light on growth and induction and priming of defense in *Plantago lanceolata*. *Frontiers in Plant Science* 12: 647372.
- R Core Team. 2021. *R: a language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing. [WWW document] URL <https://www.R-project.org> [accessed 3 March 2021].
- Rasmussen PU, Amin T, Bennett AE, Karlssongreen K, Timonen S, van Nouhuys S, Tack AJM. 2017. Plant and insect genetic variation mediate the impact of arbuscular mycorrhizal fungi on a natural plant–herbivore interaction. *Ecological Entomology* 42: 793–802.
- Rasmussen PU, Chareesri A, Neilson R, Bennett AE, Tack AJM. 2019. The impact of dispersal, plant genotype and nematodes on arbuscular mycorrhizal fungal colonization. *Soil Biology and Biochemistry* 132: 28–35.
- Rasmussen PU, Hugerth LW, Blanchet FG, Andersson AF, Lindahl BD, Tack AJM. 2018. Multiscale patterns and drivers of arbuscular mycorrhizal fungal communities in the roots and root-associated soil of a wild perennial herb. *New Phytologist* 220: 1248–1261.
- Rillig MC. 2004. Arbuscular mycorrhizae and terrestrial ecosystem processes. *Ecology Letters* 7: 740–754.
- Sagar GR, Harper JL. 1964. *Plantago major* L., *P. media* L. and *P. lanceolata* L. *Journal of Ecology* 52: 189–221.
- Sallinen S, Norberg A, Susi H, Laine A-L. 2020. Intraspecific host variation plays a key role in virus community assembly. *Nature Communications* 11: 5610.
- Salvaudon L, Giraud T, Shykoff JA. 2008. Genetic diversity in natural populations: a fundamental component of plant–microbe interactions. *Current Opinion in Plant Biology* 11: 135–143.
- Sanders FE, Sheikh NA. 1983. The development of vesicular–arbuscular mycorrhizal infection in plant root systems. *Plant and Soil* 71: 223–246.
- Santoyo G, Hernández-Pacheco C, Hernández-Salmerón J, Hernández-León R. 2017. The role of abiotic factors modulating the plant–microbe–soil interactions: towards sustainable agriculture – a review. *Spanish Journal of Agricultural Research* 15: e03R01.
- Sivapalan A. 1993. Effects of water on germination of powdery mildew conidia. *Mycological Research* 97: 71–76.
- Smith SE, Read DJ. 2008. *Mycorrhizal symbiosis*, 3<sup>rd</sup> edn. Cambridge, UK: Academic Press.
- Sochard C, Bellec L, Simon J-C, Outreman Y. 2020. Influence of “protective” symbionts throughout the different steps of an aphid–parasitoid interaction. *Current Zoology* 67: 441–453.
- Song Y, Chen D, Lu K, Sun Z, Zeng R. 2015. Enhanced tomato disease resistance primed by arbuscular mycorrhizal fungus. *Frontiers in Plant Science* 6: 1–13.
- Soubeyrand S, Laine A-L, Hanski I, Penttinen A. 2009. Spatiotemporal structure of host–pathogen interactions in a metapopulation. *The American Naturalist* 174: 308–320.
- Steinkellner S, Hage-Ahmed K, García-Garrido JM, Illana A, Ocampo JA, Vierheilig H. 2012. A comparison of wild-type, old and modern tomato cultivars in the interaction with the arbuscular mycorrhizal fungus *Glomus mosseae* and the tomato pathogen *Fusarium oxysporum* f. sp. *lycopersici*. *Mycorrhiza* 22: 189–194.
- Susi H, Laine A-L. 2015. The effectiveness and costs of pathogen resistance strategies in a perennial plant. *Journal of Ecology* 103: 303–315.
- Tack AJM, Hakala J, Petäjä T, Kulmala M, Laine A-L. 2014. Genotype and spatial structure shape pathogen dispersal and disease dynamics at small spatial scales. *Ecology* 95: 703–714.
- Tack AJM, Laine A-L. 2014. Ecological and evolutionary implications of spatial heterogeneity during the off-season for a wild plant pathogen. *New Phytologist* 202: 297–308.
- Thompson JN. 1994. *The Coevolutionary Process*. Chicago, IL, USA: The University of Chicago Press.
- Thrall PH, Laine A-L, Broadhurst LM, Bagnall DJ, Brockwell J. 2011. Symbiotic effectiveness of rhizobial mutualists varies in interactions with native Australian legume genera. *PLoS ONE* 6: ge23545.
- Tollenaere C, Laine A-L. 2013. Investigating the production of sexual resting structures in a plant pathogen reveals unexpected self-fertility and genotype-by-environment effects. *Journal of Evolutionary Biology* 26: 1716–1726.
- Venables WN, Ripley BD. 2002. *Modern applied statistics with S*, 4<sup>th</sup> edn. New York, NY, USA: Springer.
- Wardle DA, Bardgett RD, Klironomos JN, Setälä H, van der Putten WH, Wall DH. 2004. Ecological linkages between aboveground and belowground biota. *Science* 304: 1629–1633.
- Xing X, Koch AM, Jones AMP, Ragone D, Murch S, Hart MM. 2012. Mutualism breakdown in breadfruit domestication. *Proceedings of the Royal Society B: Biological Sciences* 279: 1122–1130.

## Supporting Information

Additional Supporting Information may be found online in the Supporting Information section at the end of the article.

**Dataset S1** Experimental data.

**Dataset S2** Descriptions of experimental data variables.

**Fig. S1** Photograph of *Plantago lanceolata* showing signs of infection with *Podosphaera plantaginis*.

**Fig. S2** Growth of mycorrhizal-inoculated plants relative to non-mycorrhizal plants in each seed origin population at the beginning of the field epidemic experiment.

**Fig. S3** Biomass was increased in mycorrhizal-inoculated plants relative to nonmycorrhizal plants at the end of the field epidemic experiment.

**Table S1** Effects of mycorrhizal inoculation and host genetic origin on the growth of healthy individuals at the start of the field epidemic experiment.

**Table S2** Factors influencing the growth of six seed origin populations at the end of the field epidemic experiment.

**Table S3** Factors influencing the growth of 30 maternal genotypes at the end of the field epidemic experiment.

**Table S4** Effect of mycorrhizal association on final harvest biomass in a subset of the experimental individuals.

**Table S5** Factors influencing host infection rate in six seed origin populations during the peak of the field epidemic experiment.

**Table S6** Factors influencing host infection rate in 30 maternal genotypes during the peak of the field epidemic experiment.

**Table S7** Factors influencing host infection rate in six seed origin populations at the end of the field epidemic experiment.

**Table S8** Factors influencing host infection rate in 30 maternal genotypes at the end of the field epidemic experiment.

**Table S9** Factors influencing the proportion of infected leaves in hosts in six seed origin populations during the epidemic peak.

**Table S10** Factors influencing the proportion of infected leaves in hosts in 30 maternal genotypes at the epidemic peak.

**Table S11** Factors influencing the proportion of infected leaves in hosts in six seed origin populations at the end of the field epidemic experiment.

**Table S12** Factors influencing the proportion of infected leaves in hosts in 30 maternal genotypes at the end of the field epidemic experiment.

**Table S13** Factors influencing the number of infected leaves in hosts in 30 maternal genotypes at the peak of the field epidemic experiment.

**Table S14** Factors influencing the number of infected leaves in hosts in six seed origin populations during the epidemic peak.

**Table S15** Factors influencing the number of infected leaves in hosts in six seed origin populations at the end of the field epidemic experiment.

**Table S16** Factors influencing the number of infected leaves in hosts in 30 maternal genotypes at the end of the field epidemic experiment.

**Table S17** Relationship between host growth and defensive effects as a result of mycorrhizal inoculation in the maternal genotypes.

**Table S18** Relationship between disease susceptibility and defensive effects as a result of mycorrhizal inoculation in the maternal genotypes.

Please note: Wiley Blackwell are not responsible for the content or functionality of any Supporting Information supplied by the authors. Any queries (other than missing material) should be directed to the *New Phytologist* Central Office.