

JYX



JYVÄSKYLÄN YLIOPISTO
UNIVERSITY OF JYVÄSKYLÄ

This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Krupchyk, Katya; Salo, Mikko; Uhlmann, Gunther; Wang, Jenn-Nan

Title: On the scientific work of Victor Isakov

Year: 2022

Version: Accepted version (Final draft)

Copyright: © 2022 American Institute of Mathematical Sciences (AIMS)

Rights: In Copyright

Rights url: <http://rightsstatements.org/page/InC/1.0/?language=en>

Please cite the original version:

Krupchyk, K., Salo, M., Uhlmann, G., & Wang, J.-N. (2022). On the scientific work of Victor Isakov. *Inverse Problems and Imaging*, 16(6), 1429-1445. <https://doi.org/10.3934/ipi.2022058>



ON THE SCIENTIFIC WORK OF VICTOR ISAKOV

KATYA KRUPCHYK^{✉1}, MIKKO SALO^{✉2}, GUNTHER UHLMANN^{✉3,4}, AND JENN-NAN WANG^{✉5}

¹Department of Mathematics, University of California, Irvine, USA

²Department of Mathematics and Statistics, University of Jyväskylä, Finland

³Department of Mathematics, University of Washington, USA

⁴Institute for Advanced Study, Hong Kong University of Science and Technology, China

⁵Institute of Applied Mathematical Sciences, NCTS, National Taiwan University, Taiwan

1. Introduction. This special issue of Inverse Problems and Imaging with title “Analytical aspects of inverse problems in PDEs” is dedicated to the memory of Victor Isakov, a world leading expert in the field of inverse problems for partial differential equations (PDE). Over the course of his distinguished career, Isakov made numerous fundamental contributions to inverse scattering, the Calderón problem, inverse problems for elliptic, parabolic, and hyperbolic PDE, inverse problems of potential theory, as well as uniqueness in the Cauchy problem. He wrote two very influential research monographs [27] and [35].



Victor Isakov was born on November 4, 1947, in Stalinsk (later renamed Novokuznetsk), Russia. He received his PhD in 1973 from the (now Sobolev) Institute of Mathematics, Novosibirsk, Russia, where he was also appointed as a researcher from 1971 to 1983. He also served as a part time associate professor at the Novosibirsk State University from 1974 to 1982. In 1987 Victor Isakov moved to the US, after staying for two months as a visiting professor at the University of Florence,

2020 *Mathematics Subject Classification.* Primary: 35R30.

Key words and phrases. Inverse problems, singular solutions, nonlinear PDE, complex geometrical optics, Calderón problem, partial data, increasing stability.

Italy. He held visiting positions at the Courant Institute of New York University, Cornell University, and University of Minnesota before being appointed a Professor at Wichita State University in 1988. In 2006 he was designated a Distinguished Professor by the Kansas Board of Regents, a position he held until his untimely death in 2021.

Apart from being a leading mathematician, Victor Isakov had wide cultural interests and was an outstanding pianist.

Let us now proceed to discuss some of the significant scientific results established by Victor Isakov.

2. On the mathematics of Victor Isakov.

2.1. Applications of singular solutions to inverse problems. In the study of inverse problems, besides complex geometric optics solutions [4], [69], singular solutions are also useful in many cases. The use of singular solutions in solving inverse problems was first introduced by Isakov [26]. In [26], Isakov considered the problem of determining a discontinuous conductivity coefficient from the knowledge of the Dirichlet-to-Neumann map. Let Ω be a C^2 bounded domain in \mathbb{R}^n with $n \geq 2$. Consider the conductivity equation

$$\operatorname{div}(a_j \nabla u_j) = 0 \quad \text{in } \Omega, \quad (1)$$

where $a_j(x) = a_0(x) + \chi(\Omega_j) b_j(x)$ with $a_0 \in C^2(\bar{\Omega})$, $b_j \in C^2(\bar{\Omega}_j)$ satisfying $a_0 + b_j > 0$ on $\bar{\Omega}_j$ and $b_j \neq 0$ on $\partial\Omega_j$, $j = 1, 2$. Suppose that $\Omega_j \Subset \Omega$ is a Lipschitz domain and $\Omega \setminus \bar{\Omega}_j$ is connected. Let Γ_0 be a non-empty open subset of $\partial\Omega$. Associated to (1) and Γ_0 is the local Dirichlet-to-Neumann map $\Lambda_j^{\Gamma_0}$ given by

$$\Lambda_j^{\Gamma_0}(\varphi) = a_0 \frac{\partial u_j}{\partial \nu} \Big|_{\Gamma_0},$$

where u_j is the solution to (1) with Dirichlet data φ and $\operatorname{supp}(\varphi) \subset \Gamma_0$, and ν is the unit outer normal of $\partial\Omega$. Assume that a_0 is known.

Isakov in [26] showed that if $\Lambda_1^{\Gamma_0} = \Lambda_2^{\Gamma_0}$ then $\Omega_1 = \Omega_2$ and $b_1 = b_2$ on $\partial\Omega_1$. The result is proved by a contradiction argument. Assume that $\Omega_1 \neq \Omega_2$. Let $x_0 \in \partial\Omega_1 \setminus \bar{\Omega}_2$ and B be a ball centered at x_0 with $B \cap \bar{\Omega}_2 = \emptyset$. Using that $\Lambda_1^{\Gamma_0} = \Lambda_2^{\Gamma_0}$ and applying a Runge type approximation property, one can derive the following orthogonality relation,

$$\int_{\Omega_1} b_1 \nabla u_3 \cdot \nabla u_2 = \int_{\Omega_2} b_2 \nabla u_3 \cdot \nabla u_2, \quad (2)$$

where u_2 is any solution to (1) with $j = 2$ near $\bar{\Omega}_1 \cup \bar{\Omega}_2$ and u_3 solves $\operatorname{div}(a_3 \nabla u_3) = 0$ near $\bar{\Omega}_1 \cup \bar{\Omega}_2$, where a_3 is a C^1 extension of $a_0 + b_1$ in $B \cap \bar{\Omega}_1$. The key idea used by Isakov was to substitute $u_3(x) = K_3(y, x)$ and $u_2 = K_2(y, x)$ into (2), where K_3 and K_2 are Green's functions for the corresponding equations with singularity at a point y near x_0 . Next it follows from (2) that

$$\begin{aligned} & \int_{B_0 \cap \Omega_1} b_1 \nabla K_3(y, \cdot) \cdot \nabla K_2(y, \cdot) \\ &= - \int_{B_0^c \cap \Omega_1} b_1 \nabla K_3(y, \cdot) \cdot \nabla K_2(y, \cdot) + \int_{\Omega_2} b_2 \nabla K_3(y, \cdot) \cdot \nabla K_2(y, \cdot), \end{aligned} \quad (3)$$

where $B_0 \subset B$. As $y \rightarrow x_0$, the left-hand side of (3) blows up, while the right-hand side of (3) remains bounded. This contradiction implies that $\Omega_1 = \Omega_2$. The proof of the fact that $b_1 = b_2$ on $\partial\Omega_1$ follows by a similar argument.

Isakov later applied the method to an inverse transmission scattering problem [28]. By simplifying some arguments in [28], Kirsch and Kress [41] extended Schiffer's result [57] to the case of sound-hard obstacles in which a sound-hard obstacle is uniquely determined by the far-field pattern at one fixed wave number. In [33] Isakov used the method of singular and complex geometric optics solutions to recover a transparent obstacle, transmission coefficients, as well as an electric potential in the Schrödinger equation inside the obstacle from partial boundary measurements as well as from scattering data at a fixed frequency, see also [47] for an extension of this result to the case of magnetic Schrödinger equations. We also mention that Alessandrini [2] used suitable singular solutions in the study of Calderón's problem. By modifying Isakov's method, Ikehata [20] proposed a reconstruction method of determining the discontinuity of the medium by the boundary measurements. Ikehata named the reconstruction method the probe method. Isakov's idea also played a key role in the development of a reconstruction method by Potthast [67]. Furthermore, the linear sampling method introduced by Colton and Kirsch [10] has some resemblance to Isakov's method.

2.2. Complex geometric optics solutions and completeness of products of solutions. In [29] Isakov developed an original, powerful, and flexible approach to the construction of complex geometric optics solutions for multiplicative perturbations of a large class of constant coefficient PDE. Specifically, let $P = \sum_{|\alpha| \leq m} a_\alpha D^\alpha$ be a partial differential operator on \mathbb{R}^n of order m with constant coefficients, $a_\alpha \in \mathbb{C}$, and let $P(\xi) = \sum_{|\alpha| \leq m} a_\alpha \xi^\alpha$, $\xi \in \mathbb{R}^n$, be its full symbol. We also set $\tilde{P}(\xi) = (\sum_{|\beta| \geq 0} |P^{(\beta)}(\xi)|^2)^{1/2}$, where $P^{(\beta)}(\xi) = \partial_\xi^\beta P(\xi)$. Assuming that

$$\sup_{\xi \in \mathbb{R}^n} \frac{1}{\tilde{P}(\xi + \zeta)} \rightarrow 0, \text{ as } |\zeta| \rightarrow \infty, \zeta \in P^{-1}(0) \subset \mathbb{C}^n, \quad (4)$$

in [29] Isakov constructed solutions to the equation

$$(P + q)u = 0 \quad \text{in } \Omega, \quad (5)$$

that are of the form

$$u(x; \zeta) = e^{i\zeta \cdot x} (1 + r(x; \zeta)) \in L^2(\Omega). \quad (6)$$

Here $\Omega \subset \mathbb{R}^n$ is an open bounded set, $q \in L^\infty(\Omega)$, $\zeta \in \mathbb{C}^n$, $\zeta \cdot \zeta = 0$, and $\|r\|_{L^2(\Omega)} \rightarrow 0$, as $|\zeta| \rightarrow \infty$. The proof proceeds as follows. In order for u given by (6) to satisfy (5), the remainder term r should solve

$$P(D + \zeta)r = -q(1 + r). \quad (7)$$

To solve (7), a brilliant idea of Isakov is to employ a regular fundamental solution E_ζ of $P(D + \zeta)$, constructed by Hörmander [18], which gives a right inverse of $P(D + \zeta)$ on $L^2(\Omega)$ with the bound

$$\|E_\zeta\|_{\mathcal{L}(L^2(\Omega), L^2(\Omega))} \leq C \sup_{\xi \in \mathbb{R}^n} \frac{1}{\tilde{P}(\xi + \zeta)}.$$

Thanks to the assumption (4), we have $\|E_\zeta\|_{\mathcal{L}(L^2(\Omega), L^2(\Omega))} \rightarrow 0$, as $|\zeta| \rightarrow \infty$, $\zeta \in P^{-1}(0) \subset \mathbb{C}^n$. The solution $r \in L^2(\Omega)$ of (7) is then obtained as the unique fixed point of the contraction map $r \mapsto E_\zeta(-q(1 + r))$ on $L^2(\Omega)$ for $|\zeta|$ large.

Another significant result of the work [29] is a proof of the completeness of the products of solutions to (5) with two different potentials q_1 and q_2 , under an additional assumption on the operator P , which can be stated as follows: there

exists an open non-empty set $U \subset \mathbb{R}^n$ such that for every $\xi \in U$ and every positive $R > 0$, there are $\zeta_1, \zeta_2 \in P^{-1}(0) \subset \mathbb{C}^n$ such that $\zeta_1 + \zeta_2 = \xi$, $|\zeta_j| > R$. The proof of this result is based on the construction of complex geometric optics solutions of the form (6) to the equations (5) with two different potentials q_1 and q_2 , and two complex frequencies ζ_1 and ζ_2 as in the additional assumption above. As also shown in [29], the two conditions above are satisfied, in particular, when P is the biharmonic operator, the heat operator, as well as the ultrahyperbolic operator.

An account of the powerful method of [29] of constructing complex geometric optics solutions is given in the book [7]. This method turned out to be of crucial importance when establishing the Borg–Levinson theorem for elliptic operators of higher order with constant coefficients in [45]. It was also used, in particular, when solving an inverse problem of recovering a non-compactly supported potential in the Schrödinger equation in an unbounded waveguide from boundary measurements in [40] as well as when studying the inverse problem of determining a time dependent potential in the wave equation from partial boundary measurements in [44].

2.3. Inverse problems for nonlinear equations. Isakov began to study inverse boundary value problems for nonlinear equations in the early 90’s. In [30], Isakov considered the problem of determining the semilinear term $a(x, u)$ in the parabolic equation $\partial_t u - \Delta u + a(x, u) = 0$ given all possible lateral boundary measurements (the lateral Dirichlet-to-Neumann map). A key observation in Isakov’s approach was that the equality of the Dirichlet-to-Neumann maps corresponding to the semilinear parabolic equations implies the equality of the Dirichlet-to-Neumann maps for the linear parabolic equations obtained by the linearization technique. Precisely, let Λ_{a_j} be the lateral Dirichlet-to-Neumann map associated with the equation $\partial_t u_j - \Delta u_j + a_j(x, u_j) = 0$, $j = 1, 2$, and let $\Lambda_{a_j}^*$ be the lateral Dirichlet-to-Neumann map associated to the linearized equation

$$\partial_t v_j - \Delta v_j + a_j^* v_j = 0$$

with

$$a_j^*(x, t; g) = \frac{\partial a_j}{\partial u}(x, t; u_j(x, t; g)),$$

where $u_j(x, t; g)$ is the unique solution of the semilinear parabolic equation with the lateral Dirichlet condition g . Isakov then showed that if $\Lambda_{a_1} = \Lambda_{a_2}$ then $\Lambda_{a_1}^* = \Lambda_{a_2}^*$. Having established the equality of the Dirichlet-to-Neumann maps for the linear parabolic equation, the unique determination of $a(x, u)$ follows from the corresponding result for the linear equation. The idea of using the linearization technique in treating inverse problems for nonlinear equations is highly original.

The idea of linearization in [30] was later applied to inverse boundary value problems for semilinear elliptic equations [25], [23] and for quasilinear equations [70], [31]. The inverse boundary value problem for the Navier-Stokes equation studied in [61] was also based on the linearization technique, see also [59] and [38]. Subsequently, following [30], a second order linearization of the nonlinear Dirichlet-to-Neumann map has also been successfully exploited, see [3], [71], [70]. The work [43] developed the linearization idea further, by introducing higher order linearizations in the case of inverse problems for hyperbolic PDE. This allowed one to solve such inverse problems for nonlinear equations in situations where the corresponding inverse problems in the linear setting are open, see also [11], [60] and reference there for the use of higher order linearization in the hyperbolic setting. We refer to the works [16], [54] for the introduction of higher order linearization

technique in the case of inverse problems for elliptic PDE and to the works [55], [52], [51], [6], [56] and references there for some further developments in the elliptic case.

2.4. The Calderón problem with partial data. Isakov made a fundamental contribution to the celebrated Calderón problem with partial data in his work [32]. When stating his result, let $\Omega \subset \mathbb{R}^n$, $n \geq 3$, be a bounded domain with smooth boundary, and let $\Gamma \subset \partial\Omega$ be open nonempty with $\partial\Omega \setminus \Gamma$ being either part of a hyperplane or part of a sphere. Associated to Γ , we introduce the partial Cauchy data set for the Schrödinger equation,

$$C_q^\Gamma = \{(u|_\Gamma, \partial_\nu u|_\Gamma) : u \in H^1(\Omega) \text{ satisfying } (-\Delta + q)u = 0 \text{ in } \Omega, \text{ supp}(u|_{\partial\Omega}) \subset \Gamma\}.$$

Here $q \in L^\infty(\Omega)$ and $H^1(\Omega)$ is the standard Sobolev space on Ω . In [32] Isakov proved that the equality $C_{q_1}^\Gamma = C_{q_2}^\Gamma$ implies that $q_1 = q_2$ in Ω .

Let us proceed to explain the main ideas in the proof of this result in the case when $\Omega \subset \{x \in \mathbb{R}^n : x_n > 0\}$, $\partial\Omega \cap \{x_n = 0\} \neq \emptyset$, and $\Gamma = \partial\Omega \setminus (\partial\Omega \cap \{x_n = 0\})$. First, using that $C_{q_1}^\Gamma = C_{q_2}^\Gamma$, one can derive the following integral identity,

$$\int_{\Omega} (q_1 - q_2)u_1 u_2 \, dx = 0, \quad (8)$$

for all $u_j \in H^1(\Omega)$ satisfying $(-\Delta + q_j)u_j = 0$ in Ω and $\text{supp}(u_j|_{\partial\Omega}) \subset \Gamma$, $j = 1, 2$. To construct solutions u_j which vanish on the hyperplane $x_n = 0$, the beautiful idea of Isakov is to use complex geometric optics solutions in combination with a reflection along this hyperplane. Specifically, let $\Omega^* = \{x^* = (x', -x_n) : x = (x', x_n) \in \Omega\}$ be the reflection of Ω with respect to the hyperplane $x_n = 0$, and let $\tilde{q}_j \in L^\infty(\Omega \cup \Omega^*)$ be the even extension of the potential q_j , $j = 1, 2$. Let

$$\tilde{u}_j = e^{\zeta_j \cdot x} (1 + \tilde{r}_j) \in H^1(\Omega \cup \Omega^*)$$

be a complex geometric optics solution to the Schrödinger equation $(-\Delta + \tilde{q}_j)\tilde{u}_j = 0$ in $\Omega \cup \Omega^*$, $j = 1, 2$. Here $\zeta_j \in \mathbb{C}^n$ is such that $\zeta_j \cdot \zeta_j = 0$, $|\zeta_j| \rightarrow \infty$, and the remainder term r_j satisfies $\|r_j\|_{L^2(\Omega \cup \Omega^*)} \leq C/|\zeta_j|$. The main point of [32] is to set

$$u_j(x) = \tilde{u}_j(x) - \tilde{u}_j(x^*) \in H^1(\Omega),$$

and observe that u_j satisfies $(-\Delta + q_j)u_j = 0$ in Ω and $u_j|_{x_n=0} = 0$. Substituting the solutions u_j into the integral identity (8), we get

$$\begin{aligned} & \int_{\Omega} (q_1 - q_2) \left(e^{(\zeta_1 + \zeta_2) \cdot x} (1 + \tilde{r}_1(x))(1 + \tilde{r}_2(x)) + e^{(\zeta_1^* + \zeta_2^*) \cdot x} (1 + \tilde{r}_1(x^*))(1 + \tilde{r}_2(x^*)) \right) dx \\ &= \int_{\Omega} (q_1 - q_2) \left(e^{(\zeta_1 + \zeta_2^*) \cdot x} (1 + \tilde{r}_1(x))(1 + \tilde{r}_2(x^*)) + e^{(\zeta_1^* + \zeta_2) \cdot x} (1 + \tilde{r}_1(x^*))(1 + \tilde{r}_2(x)) \right) dx. \end{aligned} \quad (9)$$

Next, the idea is that for $\xi \in \mathbb{R}^n$ belonging to a suitable non-empty open set U , one can choose ζ_j so that $\zeta_j \cdot \zeta_j = 0$, $\zeta_1 + \zeta_2 = i\xi$, $\text{Re}(\zeta_1 + \zeta_2) = \text{Re}(\zeta_1^* + \zeta_2) = 0$, and $|\text{Im}(\zeta_1 + \zeta_2^*)|, |\text{Im}(\zeta_1^* + \zeta_2)| \rightarrow \infty$ as $|\zeta_j| \rightarrow \infty$. Letting $|\zeta_j| \rightarrow \infty$ and applying the Riemann–Lebesgue lemma in the right hand side of (9), we get

$$\int_{\Omega \cup \Omega^*} (\tilde{q}_1 - \tilde{q}_2) e^{ix \cdot \xi} \, dx = 0,$$

for $\xi \in U$. Therefore, $q_1 = q_2$ in Ω , showing the result.

In [19] logarithmic type stability estimates were obtained for Isakov's result [32], see also [53] and [64] for the increasing stability phenomenon. Isakov's reflection approach has been further employed for partial data inverse problems for other important PDE, in particular for Maxwell's systems in [13], [5], and magnetic Schrödinger equations in [42], [65]. Partial data results for the slab and half spaces geometries based on Isakov's reflection approach were obtained in [58], [42], [62], [12], and [66]. The work [48] unified the approaches [50] and [32] and extended both of them, see also [37]. We refer to [49] for a survey on partial data inverse problems.

2.5. Increasing stability in inverse problems. In his later career, Isakov was interested in the question of increasing stability in inverse problems. Most inverse problems are known to be ill-posed. The stability estimate is often an optimal log-type estimate, see for example [1], [63], [46]. However, numerical evidence [8] indicates that the stability increases with the wave number. Also, motivated by the result of John [39] where in general one can only expect a weak logarithmic stability in the Cauchy problem for the Helmholtz equation in the general geometry, Isakov [17] tried to provide a rigorous justification of the increasing stability phenomenon by considering the Cauchy problem for the Helmholtz equation under a convexity condition. The result of [17] shows that the stability of the continuation from the Cauchy data on the sub-domain of the boundary consists of two parts, a Lipschitz part and a Hölder part. The Hölder part decreases as the wave number increases. The argument used in [17] is elementary involving a suitable Carleman estimate. It is fair to say that Isakov initiated the investigation of increasing stability for inverse problems by carefully analyzing the effect of the wave number or the frequency on the stability estimate.

The result of [17] was later extended to the increasing stability for the Cauchy problem for the Helmholtz equation in the whole domain [68]. In this case, the stability estimate consists of two terms, a Lipschitz term and a logarithmic term. The interesting property is that the logarithmic part decreases as the wave number grows. This turns out to be a typical situation for some inverse boundary value problems, see [34], [24], [22], and for some inverse source problems, see [9], [21], [14], [15], [36]. The method used in the inverse source problems mentioned above is based on the observability bounds for the wave-type equations and an explicit bound in the analytic continuation.

2.6. Other contributions. It is impossible to cover all of Isakov's vast contributions to the field of inverse problems in this short editorial. For instance, the short review above does not address the fundamental work done by Isakov in the fields of the inverse problem of gravimetry, the inverse conductivity problem with a single measurement, Carleman estimates and applications to the unique continuation property and inverse problems, inverse parabolic problems, the inverse problem of option pricing, etc. For interested readers, we refer to Isakov's books [27], [35] and references therein for further details.

3. List of Victor Isakov's publications.

3.1. Research monographs.

1. Isakov, V., *Inverse source problems*, Mathematical Surveys and Monographs, 34. American Mathematical Society, Providence, RI, 1990.

2. Isakov, V., *Inverse problems for partial differential equations*, Applied Mathematical Sciences, 127. Springer-Verlag, New York, 1998. (Second edition 2006, third edition 2017).

3.2. Research papers.

1. Isakov, V., *The uniqueness of the solution of the exterior contact inverse problem of the theory of the logarithmic potential*, (Russian) *Differentsial'nye Uravneniya* **7** (1971), 34–39.
2. Isakov, V., *The uniqueness of the solution of the contact inverse problem of potential theory*, (Russian) *Differentsial'nye Uravneniya* **8** (1972), 30–40.
3. Isakov, V., *An inverse problem of potential theory for the string equation*, (Russian) *Differentsial'nye Uravneniya* **9** (1973), 74–83, 194.
4. Isakov, V., *The solvability of the inverse problem of potential theory*, (Russian) *Dokl. Akad. Nauk SSSR* **213** (1973), 286–288.
5. Isakov, V., *The solvability of the inverse problem of potential theory*, (Russian) *Sibirsk. Mat. Z.* **14** (1973), 987–1005, 1157.
6. Isakov, V., *The uniqueness of the reconstruction of a domain and the density from the exterior potential*, (Russian) *Sibirsk. Mat. Z.* **15** (1974), 683–686, 704.
7. Isakov, V., *The uniqueness of the solution of certain inverse hyperbolic problems*, (Russian) *Differentsial'nye Uravneniya* **10** (1974), 165–167, 183.
8. Isakov, V., *Inverse theorems on the smoothness of potentials*, (Russian) *Differentsial'nye Uravneniya* **11** (1975), 66–74, 202.
9. Isakov, V., *The reconstruction of a domain from scattering data*, (Russian) *Dokl. Akad. Nauk SSSR* **230** (1976), no. 3, 520–522.
10. Isakov, V., *Uniqueness theorems for inverse problems of thermal potentials*, (Russian) *Sibirsk. Mat. Z.* **17** (1976), no. 2, 259–272, 477.
11. Isakov, V., *Analyticity of the solutions of nonlinear transmission problems*, (Russian) *Differentsial'nye Uravneniya* **12** (1976), no. 1, 59–68, 187.
12. Isakov, V., *On the inverse problem of potential for hyperbolic equations*, (Russian) *Dokl. Akad. Nauk SSSR* **237** (1977), no. 2, 257–259.
13. Isakov, V., *The inverse problem of potential theory for hyperbolic equations*, (Russian) *Differentsial'nye Uravneniya* **13** (1977), no. 2, 305–313, 381–382.
14. Isakov, V., *General diffraction problems for hyperbolic equations*, (Russian) *Partial differential equations*, 32–56, 180, *Proc. Sobolev Sem.*, 77–2, *Akad. Nauk SSSR Sibirsk. Otdel., Inst. Mat., Novosibirsk*, 1977.
15. Isakov, V., *Strictly hyperbolic diffraction problems*, (Russian) *Differentsial'nye Uravneniya* **14** (1978), no. 1, 30–38, 188.
16. Isakov, V., *Uniqueness theorems for the inverse problem of potential theory*, (Russian) *Dokl. Akad. Nauk SSSR* **240** (1978), no. 3, 526–529.
17. Isakov, V., *The existence of solutions of inverse problems of potential theory*, (Russian) *Dokl. Akad. Nauk SSSR* **240** (1978), no. 4, 771–773.
18. Isakov, V., *Reconstruction of a domain from scattering data*, (Russian) *Direct and inverse problems for partial differential equations and their applications* (Russian), pp. 148–159, 187, "Fan", Tashkent, 1978.
19. Isakov, V., *On uniqueness of solutions for inverse problems of potential theory*, *Inverse and improperly posed problems in differential equations* (*Proc. Conf., Math. Numer. Methods, Halle, 1979*), pp. 135–139, *Math. Res.*, 1, *Akademie-Verlag, Berlin*, 1979.

20. Isakov, V., *Existence of solutions of an inverse problem of potential theory*, (Russian) *Differentsial'nye Uravneniya* **16** (1980), no. 6, 1097–1107, 1149.
21. Isakov, V., *On the uniqueness of the solution of the Cauchy problem*, (Russian) *Dokl. Akad. Nauk SSSR* **255** (1980), no. 1, 18–21.
22. Isakov, V., *A class of inverse problems for parabolic equations*, (Russian) *Dokl. Akad. Nauk SSSR* **263** (1982), no. 6, 1296–1299.
23. Isakov, V., *Uniqueness of the extension of solutions of hyperbolic equations*, (Russian) *Mat. Zametki* **32** (1982), no. 1, 75–82, 126.
24. Isakov, V., *A nonhyperbolic Cauchy problem for $\square_b \square_c$ and its applications to elasticity theory*, *Comm. Pure Appl. Math.* **39** (1986), no. 6, 747–767.
25. Isakov, V., *On uniqueness of recovery of a discontinuous conductivity coefficient*, *Comm. Pure Appl. Math.* **41** (1988), no. 7, 865–877.
26. Isakov, V., *Uniqueness of the continuation across a time-like hyperplane and related inverse problems for hyperbolic equations*, *Comm. Partial Differential Equations* **14** (1989), no. 4, 465–478.
27. Isakov, V., *Uniqueness for inverse parabolic problems with a lateral overdetermination*, *Comm. Partial Differential Equations* **14** (1989), no. 6, 681–689.
28. Friedman, A., Isakov, V., *On the uniqueness in the inverse conductivity problem with one measurement*, *Indiana Univ. Math. J.* **38** (1989), no. 3, 563–579.
29. Isakov, V., Powell, J., *On the inverse conductivity problem with one measurement*, *Inverse Problems* **6** (1990), no. 2, 311–318.
30. Isakov, V., *Some inverse problems for elliptic and parabolic equations*, *Inverse problems in partial differential equations* (Arcata, CA, 1989), 203–214, SIAM, Philadelphia, PA, 1990.
31. Isakov, V., *On uniqueness in the inverse transmission scattering problem*, *Comm. Partial Differential Equations* **15** (1990), no. 11, 1565–1587.
32. Isakov, V., *Inverse parabolic problems with the final overdetermination*, *Comm. Pure Appl. Math.* **44** (1991), no. 2, 185–209.
33. Ames, K. A., Isakov, V., *An explicit stability estimate for an ill-posed Cauchy problem for the wave equation*, *J. Math. Anal. Appl.* **156** (1991), no. 2, 597–610.
34. Isakov, V., *An inverse hyperbolic problem with many boundary measurements*, *Comm. Partial Differential Equations* **16** (1991), no. 6–7, 1183–1195.
35. Isakov, V., *Completeness of products of solutions and some inverse problems for PDE*, *J. Differential Equations* **92** (1991), no. 2, 305–316.
36. Bellout, H., Friedman, A., Isakov, V., *Stability for an inverse problem in potential theory*, *Trans. Amer. Math. Soc.* **332** (1992), no. 1, 271–296.
37. Isakov, V., Sun, Z. Q., *Stability estimates for hyperbolic inverse problems with local boundary data*, *Inverse Problems* **8** (1992), no. 2, 193–206.
38. Isakov, V., *On uniqueness of the discontinuous conductivity coefficient*, *Wave propagation and inversion*, 46–55, SIAM, Philadelphia, PA, 1992.
39. Isakov, V., *Stability estimates for obstacles in inverse scattering*, *J. Comput. Appl. Math.* **42** (1992), no. 1, 79–88.
40. Engl, H. W., Isakov, V., *On the identifiability of steel reinforcement bars in concrete from magnetostatic measurements*, *European J. Appl. Math.* **3** (1992), no. 3, 255–262.
41. Isakov, V., *On uniqueness in inverse problems for semilinear parabolic equations*, *Arch. Rational Mech. Anal.* **124** (1993), no. 1, 1–12.

42. Isakov, V., *Carleman type estimates in an anisotropic case and applications*, J. Differential Equations **105** (1993), no. 2, 217–238.
43. Isakov, V., *Prospecting discontinuities by boundary measurements*, Inverse problems: principles and applications in geophysics, technology, and medicine (Potsdam, 1993), 215–223, Math. Res., 74, Akademie-Verlag, Berlin, 1993.
44. Isakov, V., *New stability results for soft obstacles in inverse scattering*, Inverse Problems **9** (1993), no. 5, 535–543.
45. Isakov, V., *Uniqueness and stability in multi-dimensional inverse problems*, Inverse Problems **9** (1993), no. 6, 579–621.
46. Isakov, V., *Uniqueness for inverse problems in quasilinear differential equations*, Inverse problems in mathematical physics (Saariselkä, 1992), 71–86, Lecture Notes in Phys., 422, Springer, Berlin, 1993.
47. Isakov, V., Sylvester, J., *Global uniqueness for a semilinear elliptic inverse problem*, Comm. Pure Appl. Math. **47** (1994), no. 10, 1403–1410.
48. Isakov, V., *On uniqueness in the inverse conductivity problem with one boundary measurement*, Tomography, impedance imaging, and integral geometry (South Hadley, MA, 1993), 105–114, Lectures in Appl. Math., 30, Amer. Math. Soc., Providence, RI, 1994.
49. Alessandrini, G., Isakov, V., Powell, J., *Local uniqueness in the inverse conductivity problem with one measurement*, Trans. Amer. Math. Soc. **347** (1995), no. 8, 3031–3041.
50. Isakov, V., Nachman, A., *Global uniqueness for a two-dimensional semilinear elliptic inverse problem*, Trans. Amer. Math. Soc. **347** (1995), no. 9, 3375–3390.
51. Elcrat, A., Isakov, V., Neculoiu, O., *On finding a surface crack from boundary measurements*, Inverse Problems **11** (1995), no. 2, 343–351.
52. Isakov, V., *Uniqueness and stability in inverse parabolic problems*, Inverse problems in diffusion processes (Lake St. Wolfgang, 1994), 21–41, SIAM, Philadelphia, PA, 1995.
53. Isakov, V., Sun, Z., *The inverse scattering at fixed energies in two dimensions*, Indiana Univ. Math. J. **44** (1995), no. 3, 883–896.
54. Isakov, V., *On uniqueness for a discontinuity surface of the speed of propagation*, J. Inverse Ill-Posed Probl. **4** (1996), no. 1, 33–38.
55. Elayyan, A., Isakov, V., *On identification of a diffusion coefficient*, Distributed parameter systems: modelling and control (Warsaw, 1995). Control Cybernet. **25** (1996), no. 3, 441–450.
56. Isakov, V., Sever, A., *Numerical implementation of an integral equation method for the inverse conductivity problem*, Inverse Problems **12** (1996), no. 6, 939–951.
57. Alessandrini, G., Isakov, V., *Analyticity and uniqueness for the inverse conductivity problem*, Rend. Istit. Mat. Univ. Trieste **28** (1996), no. 1–2, 351–369 (1997).
58. Elayyan, A., Isakov, V., *On uniqueness of recovery of the discontinuous conductivity coefficient of a parabolic equation*, SIAM J. Math. Anal. **28** (1997), no. 1, 49–59.
59. Isakov, V., *On uniqueness in a lateral Cauchy problem with multiple characteristics*, J. Differential Equations **134** (1997), no. 1, 134–147.

60. Isakov, V., *On uniqueness and stability in the Cauchy problem*, Optimization methods in partial differential equations (South Hadley, MA, 1996), 131–146, Contemp. Math., 209, Amer. Math. Soc., Providence, RI, 1997.
61. Bouchouev, I., Isakov, V., *The inverse problem of option pricing*, Inverse Problems **13** (1997), no. 5, L11–L17.
62. Bukhgeim, A., Dyatlov, G., Isakov, V., *Stability of memory reconstruction from the Dirichlet-Neumann operator*, (Russian) Sibirsk. Mat. Zh. **38** (1997), no. 4, 738–749, i; translation in Siberian Math. J. **38** (1997), no. 4, 636–646.
63. Elayyan, A., Isakov, V., *On an inverse diffusion problem*, SIAM J. Appl. Math. **57** (1997), no. 6, 1737–1748.
64. Isakov, V., *On reconstruction of the diffusion and of the principal coefficient of a hyperbolic equation*, Inverse problems in wave propagation (Minneapolis, MN, 1995), 259–275, IMA Vol. Math. Appl., 90, Springer, New York, 1997.
65. Isakov, V., *Some inverse problems for the diffusion equation*, Conference on Inverse Problems, Control and Shape Optimization (Carthage, 1998). Inverse Problems **15** (1999), no. 1, 3–10.
66. Bouchouev, I., Isakov, V., *Uniqueness, stability and numerical methods for the inverse problem that arises in financial markets*, Inverse Problems **15** (1999), no. 3, R95–R116.
67. Isakov, V., Kindermann, S., *Identification of the diffusion coefficient in a one-dimensional parabolic equation*, Inverse Problems **16** (2000), no. 3, 665–680.
68. Belishev, M., Isakov, V., Pestov, L., Sharafutdinov, V., *On the reconstruction of a metric from external electromagnetic measurements*, (Russian) Dokl. Akad. Nauk **372** (2000), no. 3, 298–300.
69. Eller, M., Isakov, V., *Carleman estimates with two large parameters and applications*, Differential geometric methods in the control of partial differential equations (Boulder, CO, 1999), 117–136, Contemp. Math., 268, Amer. Math. Soc., Providence, RI, 2000.
70. Isakov, V., Yamamoto, M., *Carleman estimate with the Neumann boundary condition and its applications to the observability inequality and inverse hyperbolic problems*, Differential geometric methods in the control of partial differential equations (Boulder, CO, 1999), 191–225, Contemp. Math., 268, Amer. Math. Soc., Providence, RI, 2000.
71. Bukhgeim, A., Cheng, J., Isakov, V., Yamamoto, M., *Uniqueness in determining damping coefficients in hyperbolic equations*, Analytic extension formulas and their applications (Fukuoka, 1999/Kyoto, 2000), 27–46, Int. Soc. Anal. Appl. Comput., 9, Kluwer Acad. Publ., Dordrecht, 2001.
72. Delillo, T., Isakov, V., Valdivia, N., Wang, L., *The detection of the source of acoustical noise in two dimensions*, SIAM J. Appl. Math. **61** (2001), no. 6, 2104–2121.
73. Isakov, V., *Uniqueness of recovery of some systems of semilinear partial differential equations*, Special issue to celebrate Pierre Sabatier’s 65th birthday (Montpellier, 2000). Inverse Problems **17** (2001), no. 4, 607–618.
74. Isakov, Victor *On the uniqueness of the continuation for a thermoelasticity system*. SIAM J. Math. Anal. **33** (2001), no. 3, 509–522.
75. Isakov, V., *Uniqueness of recovery of some quasilinear partial differential equations*, Comm. Partial Differential Equations **26** (2001), no. 11–12, 1947–1973.
76. Bouchouev, I., Isakov, V., Valdivia, N., *Recovery of volatility coefficient by linearization*, Quant. Finance **2** (2002), no. 4, 257–263.

77. Isakov, V., Wu, S., *On theory and application of the Helmholtz equation least squares method in inverse acoustics*, Inverse Problems **18** (2002), no. 4, 1147–1159.
78. Eller, M., Isakov, V., Nakamura, G., Tataru, D., *Uniqueness and stability in the Cauchy problem for Maxwell and elasticity systems*, Nonlinear partial differential equations and their applications. Collège de France Seminar, Vol. XIV (Paris, 1997/1998), 329–349, Stud. Math. Appl., 31, North-Holland, Amsterdam, 2002.
79. Isakov, V., *Uniqueness of recovery of some systems of quasilinear elliptic and parabolic partial differential equations*, Nonlinear problems in mathematical physics and related topics, II, 201–212, Int. Math. Ser. (N. Y.), 2, Kluwer/Plenum, New York, 2002.
80. Isakov, V., *On detecting the source of acoustical noise*, Theoretical and computational acoustics, 2001 (Beijing), 615–628, World Sci. Publ., River Edge, NJ, 2002.
81. Imanuvilov, O., Isakov, V., Yamamoto, M., *An inverse problem for the dynamical Lamé system with two sets of boundary data*, Comm. Pure Appl. Math. **56** (2003), no. 9, 1366–1382.
82. DeLillo, T., Isakov, V., Valdivia, N., Wang, L., *The detection of surface vibrations from interior acoustical pressure*, Inverse Problems **19** (2003), no. 3, 507–524.
83. Isakov, V., Yamamoto, M., *Stability in a wave source problem by Dirichlet data on subboundary*, J. Inverse Ill-Posed Probl. **11** (2003), no. 4, 399–409.
84. Cheng, J., Isakov, V., Yamamoto, M., Zhou, Q., *Lipschitz stability in the lateral Cauchy problem for elasticity system*, J. Math. Kyoto Univ. **43** (2003), no. 3, 475–501.
85. Isakov, V., Nakamura, G., Wang, J.-N., *Uniqueness and stability in the Cauchy problem for the elasticity system with residual stress*, Inverse problems: theory and applications (Cortona/Pisa, 2002), 99–113, Contemp. Math., 333, Amer. Math. Soc., Providence, RI, 2003.
86. Isakov, V., *The inverse problem of option pricing*, Recent development in theories & numerics, 47–55, World Sci. Publ., River Edge, NJ, 2003.
87. Belishev, M., Isakov, V., *On the uniqueness of the reconstruction of the parameters of the Maxwell system from dynamic boundary data*, (Russian) Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI) 285 (2002), Mat. Vopr. Teor. Rasprostr. Voln. 31, 15–32, 224; translation in J. Math. Sci. (N.Y.) **122** (2004), no. 5, 3459–3469
88. Isakov, V., *Carleman type estimates and their applications*, New analytic and geometric methods in inverse problems, 93–125, Springer, Berlin, 2004.
89. Hrycak, T., Isakov, V., *Increased stability in the continuation of solutions to the Helmholtz equation*, Inverse Problems **20** (2004), no. 3, 697–712.
90. Isakov, V., *Carleman estimates and applications to inverse problems*, Milan J. Math. **72** (2004), 249–271.
91. Delillo, T., Hrycak, T., Isakov, V., *Theory and boundary element methods for near-field acoustic holography*, J. Comput. Acoust. **13** (2005), no. 1, 163–185.
92. Isakov, V., *An inverse problem for the dynamical Lamé system with two sets of local boundary data*, Control theory of partial differential equations, 101–109, Lect. Notes Pure Appl. Math., 242, Chapman & Hall/CRC, Boca Raton, FL, 2005.

93. Isakov, V., *On uniqueness in the inverse conductivity problem with local data*, Inverse Probl. Imaging **1** (2007), no. 1, 95–105.
94. Isakov, V., *Increased stability in the continuation for the Helmholtz equation with variable coefficient*, Control methods in PDE-dynamical systems, 255–267, Contemp. Math., 426, Amer. Math. Soc., Providence, RI, 2007.
95. Isakov, V., Wang, J.-N., Yamamoto, M., *Uniqueness and stability of determining the residual stress by one measurement*, Comm. Partial Differential Equations **32** (2007), no. 4-6, 833–848.
96. Subbarayappa, D., Isakov, V., *On increased stability in the continuation of the Helmholtz equation*, Inverse Problems **23** (2007), no. 4, 1689–1697.
97. Isakov, V., Wang, J.-N., Yamamoto, M., *An inverse problem for a dynamical Lamé system with residual stress*, SIAM J. Math. Anal. **39** (2007/08), no. 4, 1328–1343.
98. Isakov, V., *On uniqueness of obstacles and boundary conditions from restricted dynamical and scattering data*, Inverse Probl. Imaging **2** (2008), no. 1, 151–165.
99. Isakov, V., *On uniqueness in the general inverse transmission problem*, Comm. Math. Phys. **280** (2008), no. 3, 843–858.
100. Isakov, V., *On inverse problems in secondary oil recovery*, European J. Appl. Math. **19** (2008), no. 4, 459–478.
101. Isakov, V., Kim, N., *Carleman estimates with two large parameters for second order operators and applications to elasticity with residual stress*, Appl. Math. (Warsaw) **35** (2008), no. 4, 447–465.
102. Isakov, V., *Increased stability in the Cauchy problem for some elliptic equations*, Instability in models connected with fluid flows. I, 339–362, Int. Math. Ser. (N. Y.), 6, Springer, New York, 2008.
103. Isakov, V., Kim, N., *Global uniqueness and Lipschitz stability of residual stress from one boundary measurement*, Mathematical methods for imaging and inverse problems, 45–54, ESAIM Proc., 26, EDP Sci., Les Ulis, 2009.
104. Isakov, V., Kim, N., *Carleman estimates with second large parameter for second order operators*, Sobolev spaces in mathematics. III, 135–159, Int. Math. Ser. (N. Y.), 10, Springer, New York, 2009.
105. Imanuvilov, O., Isakov, V., Yamamoto, M., *New realization of the pseudoconvexity and its application to an inverse problem*, Appl. Anal. **88** (2009), no. 5, 637–652.
106. Isakov, V., *Inverse obstacle problems*, Inverse Problems **25** (2009), no. 12, 123002, 18 pp.
107. Isakov, V., *On identification of doping profile in semiconductors*, Imaging microstructures, 123–136, Contemp. Math., 494, Amer. Math. Soc., Providence, RI, 2009.
108. Isakov, V., Kim, N., *Weak Carleman estimates with two large parameters for second order operators and applications to elasticity with residual stress*, Discrete Contin. Dyn. Syst. **27** (2010), no. 2, 799–825.
109. Aralummalle Subbarayappa, D., Isakov, V., *Increasing stability of the continuation for the Maxwell system*, Inverse Problems **26** (2010), no. 7, 074005, 14 pp.
110. Isakov, V., Kim, K., Nakamura, G., *Reconstruction of an unknown inclusion by thermography*, Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) **9** (2010), no. 4, 725–758.

111. Isakov, V., *Increasing stability for the Schrödinger potential from the Dirichlet-to-Neumann map*, Discrete Contin. Dyn. Syst. Ser. S **4** (2011), no. 3, 631–640.
112. Isakov, V., Kindermann, S., *Subspaces of stability in the Cauchy problem for the Helmholtz equation*, Methods Appl. Anal. **18** (2011), no. 1, 1–29.
113. Isakov, V., Leung, S., Qian, J., *A fast local level set method for inverse gravimetry*, Commun. Comput. Phys. **10** (2011), no. 4, 1044–1070.
114. Elcrat, A., Isakov, V., Kropf, E., Stewart, D., *A stability analysis of the harmonic continuation*, Inverse Problems **28** (2012), no. 7, 075016, 13 pp.
115. Isakov, V., Myers, J., *On the inverse doping profile problem*, Inverse Probl. Imaging **6** (2012), no. 3, 465–486.
116. Isakov, V., *Carleman estimates for some anisotropic elasticity systems and applications*, Evol. Equ. Control Theory **1** (2012), no. 1, 141–154.
117. Isakov, V., Leung, S., Qian, J., *A three-dimensional inverse gravimetry problem for ice with snow caps*, Inverse Probl. Imaging **7** (2013), no. 2, 523–544.
118. Isakov, V., *Recovery of time dependent volatility coefficient by linearization*, Evol. Equ. Control Theory **3** (2014), no. 1, 119–134.
119. Isakov, V., Nagayasu, S., Uhlmann, G., Wang, J.-N., *Increasing stability of the inverse boundary value problem for the Schrödinger equation*, Inverse problems and applications, 131–141, Contemp. Math., 615, Amer. Math. Soc., Providence, RI, 2014.
120. Isakov, V., Wang, J.-N., *Increasing stability for determining the potential in the Schrödinger equation with attenuation from the Dirichlet-to-Neumann map*, Inverse Probl. Imaging **8** (2014), no. 4, 1139–1150.
121. Isakov, V., *Increasing stability of the continuation for general elliptic equations of second order*, New prospects in direct, inverse and control problems for evolution equations, 203–218, Springer INdAM Ser., 10, Springer, Cham, 2014.
122. Isakov, V., *Increasing stability for near field from the scattering amplitude*, Spectral theory and partial differential equations, 59–70, Contemp. Math., 640, Amer. Math. Soc., Providence, RI, 2015.
123. Cheng, J., Isakov, V., Lu, S., *Increasing stability in the inverse source problem with many frequencies*, J. Differential Equations **260** (2016), no. 5, 4786–4804.
124. Isakov, V., Lai, R.-Y., Wang, J.-N., *Increasing stability for the conductivity and attenuation coefficients*, SIAM J. Math. Anal. **48** (2016), no. 1, 569–594.
125. Isakov, V., *Carleman estimates with two large parameters for an anisotropic system of elasticity*, Rend. Istit. Mat. Univ. Trieste **48** (2016), 261–283.
126. Deng, Z.-C., Hon, Y. C., Isakov, V., *Recovery of time-dependent volatility in option pricing model*, Inverse Problems **32** (2016), no. 11, 115010, 30 pp.
127. Isakov, V., Lu, S., *Increasing stability in the inverse source problem with attenuation and many frequencies*, SIAM J. Appl. Math. **78** (2018), no. 1, 1–18.
128. Entekhabi, M. N., Isakov, V., *On increasing stability in the two dimensional inverse source scattering problem with many frequencies*, Inverse Problems **34** (2018), no. 5, 055005, 14 pp.
129. Isakov, V., Lu, S., *Inverse source problems without (pseudo) convexity assumptions*, Inverse Probl. Imaging **12** (2018), no. 4, 955–970.

130. Isakov, V., Kabanikhin, S., Shananin, A., Shishlenin, M., Zhang, S., *Algorithm for determining the volatility function in the Black-Scholes model*, Comput. Math. Math. Phys. **59** (2019), no. 10, 1753–1758.
131. Isakov, V., *On increasing stability of the continuation for elliptic equations of second order without (pseudo)convexity assumptions*, Inverse Probl. Imaging **13** (2019), no. 5, 983–1006.
132. Isakov, V., Lu, S., Xu, B., *Linearized inverse Schrödinger potential problem at a large wavenumber*, SIAM J. Appl. Math. **80** (2020), no. 1, 338–358.
133. Isakov, V., Lu, S., *On the inverse source problem with boundary data at many wave numbers*, Inverse problems and related topics, 59–80, Springer Proc. Math. Stat., 310, Springer, Singapore, 2020.
134. Entekhabi, M., Isakov, V., *Increasing stability in acoustic and elastic inverse source problems*, SIAM J. Math. Anal. **52** (2020), no. 5, 5232–5256.
135. Isakov, V., Wang, J.-N., *Uniqueness and increasing stability in electromagnetic inverse source problems*, J. Differential Equations **283** (2021), 110–135.
136. Isakov, V., Titi, A., *Stability and the inverse gravimetry problem with minimal data*, J. Inverse Ill-Posed Probl. **30** (2022), no. 1, 147–162.
137. Isakov, V., Lu, S., Xu, B., *A linearised inverse conductivity problem for the Maxwell system at a high frequency*, J. Comput. Phys. **455** (2022), Paper No. 111003, 18 pp.
138. Isakov, V., *On increasing stability in an inverse source problem with local boundary data at many wave numbers*, Appl. Anal. **101** (2022), no. 10, 3550–3562.

4. List of Victor Isakov’s students.

- Jeff Powell (Ph.D. 1991, Wichita State University)
- Ali Sever (Ph.D. 1995, Wichita State University)
- Alaeddin Elayyan (Ph.D. 1996, Wichita State University)
- Ilya Bouchouev (Ph.D. 1997, Wichita State University)
- Matthias Eller (Ph.D. 1997, Wichita State University)
- Nicolas Valdivia (Ph.D. 2002, Wichita State University)
- Joseph Myers (Ph.D. 2009, Wichita State University)
- Nanhee Kim (Ph.D. 2010, Wichita State University)
- Deepak Aralumallige Subbarayappa (Ph.D. 2010, Wichita State University)
- Li Liang (Ph.D. 2015, Wichita State University)
- Mozghan Nora Entekhabi (Ph.D. 2018, Wichita State University)
- Aseel Titi (Ph.D. 2021, Wichita State University)

REFERENCES

- [1] G. Alessandrini. Stable determination of conductivity by boundary measurements. *Appl. Anal.*, 27(1-3):153–172, 1988.
- [2] G. Alessandrini. Singular solutions of elliptic equations and the determination of conductivity by boundary measurements. *J. Differential Equations*, 84(2):252–272, 1990.
- [3] Y. Assylbekov and T. Zhou. Direct and inverse problems for the nonlinear time-harmonic Maxwell equations in Kerr-type media. *J. Spectr. Theory*, 11, 2021.
- [4] A. Calderón. Seminar on numerical analysis and its applications to continuum physics. *Soc. Brasileira de Matematica*, pages 65–73, 1980.
- [5] P. Caro. On an inverse problem in electromagnetism with local data: stability and uniqueness. *Inverse Probl. Imaging*, 5(2):297–322, 2011.

- [6] C. Cârstea, A. Feizmohammadi, Y. Kian, K. Krupchyk, and G. Uhlmann. The Calderón inverse problem for isotropic quasilinear conductivities. *Adv. Math.*, 391:Paper No. 107956, 31, 2021.
- [7] M. Choulli. *Une introduction aux problèmes inverses elliptiques et paraboliques*, volume 65 of *Mathématiques & Applications (Berlin) [Mathematics & Applications]*. Springer-Verlag, Berlin, 2009.
- [8] D. Colton, H. Haddar, and M. Piana. The linear sampling method in inverse electromagnetic scattering theory. *Inverse problems*, 19(6):S105–S137, 2003.
- [9] J. Cheng, V. Isakov, and S. Lu. Increasing stability in the inverse source problem with many frequencies. *J. Differential Equations*, 260(5):4786–4804, 2016.
- [10] D. Colton and A. Kirsch. A simple method for solving inverse scattering problems in the resonance region. *Inverse problems*, 12(4):383–393, 1996.
- [11] X. Chen, M. Lassas, L. Oksanen, and G. Paternain. Detection of Hermitian connections in wave equations with cubic non-linearity. *J. Eur. Math. Soc. (JEMS)*, 24(7):2191–2232, 2022.
- [12] P. Caro and K. Marinov. Stability of inverse problems in an infinite slab with partial data. *Comm. Partial Differential Equations*, 41(4):683–704, 2016.
- [13] P. Caro, P. Ola, and M. Salo. Inverse boundary value problem for Maxwell equations with local data. *Comm. Partial Differential Equations*, 34(10-12):1425–1464, 2009.
- [14] M. Entekhabi and V. Isakov. On increasing stability in the two dimensional inverse source scattering problem with many frequencies. *Inverse Problems*, 34(5):055005, 2018.
- [15] M. Entekhabi and V. Isakov. Increasing stability in acoustic and elastic inverse source problems. *SIAM J. Math. Anal.*, 52(5):5232–5256, 2020.
- [16] A. Feizmohammadi and L. Oksanen. An inverse problem for a semi-linear elliptic equation in Riemannian geometries. *J. Differential Equations*, 269(6):4683–4719, 2020.
- [17] T. Hrycak and V. Isakov. Increased stability in the continuation of solutions to the helmholtz equation. *Inverse problems*, 20(3):697–712, 2004.
- [18] L. Hörmander. *The analysis of linear partial differential operators. II. Classics in Mathematics*. Springer-Verlag, Berlin, 2005. Differential operators with constant coefficients, Reprint of the 1983 original.
- [19] H. Heck and J.-N. Wang. Optimal stability estimate of the inverse boundary value problem by partial measurements. *Rend. Istit. Mat. Univ. Trieste*, 48:369–383, 2016.
- [20] M. Ikehata. Reconstruction of the shape of the inclusion by boundary measurements. *Comm. Partial Differential Equations*, 23(7-8):1459–1474, 1998.
- [21] V. Isakov and S. Lu. Increasing stability in the inverse source problem with attenuation and many frequencies. *SIAM J. Appl. Math.*, 78(1):1–18, 2018.
- [22] V. Isakov, R.-Y. Lai, and J.-N. Wang. Increasing stability for the conductivity and attenuation coefficients. *SIAM J. Math. Anal.*, 48(1):569–594, 2016.
- [23] V. Isakov and A. Nachman. Global uniqueness for a two-dimensional semilinear elliptic inverse problem. *Trans. Amer. Math. Soc.*, 347(9):3375–3390, 1995.
- [24] V. Isakov, S. Nagayasu, G. Uhlmann, and J.-N. Wang. Increasing stability of the inverse boundary value problem for the schrödinger equation. *Contemp. Math*, 615:131–141, 2014.
- [25] V. Isakov and J. Sylvester. Global uniqueness for a semilinear elliptic inverse problem. *Comm. Pure Appl. Math.*, 47(10):1403–1410, 1994.
- [26] V. Isakov. On uniqueness of recovery of a discontinuous conductivity coefficient. *Comm. Pure Appl. Math.*, 41(7):865–877, 1988.
- [27] V. Isakov. *Inverse source problems*, volume 34 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 1990.
- [28] V. Isakov. On uniqueness in the inverse transmission scattering problem. *Comm. Partial Differential Equations*, 15(11):1565–1586, 1990.
- [29] V. Isakov. Completeness of products of solutions and some inverse problems for PDE. *J. Differential Equations*, 92(2):305–316, 1991.
- [30] V. Isakov. On uniqueness in inverse problems for semilinear parabolic equations. *Arch. Ration. Mech. Anal.*, 124(1):1–12, 1993.
- [31] V. Isakov. Uniqueness of recovery of some quasilinear partial differential equations. *Comm. Partial Differential Equations*, 26(11-12):1947–1973, 2001.

- [32] V. Isakov. On uniqueness in the inverse conductivity problem with local data. *Inverse Probl. Imaging*, 1(1):95–105, 2007.
- [33] V. Isakov. On uniqueness in the general inverse transmission problem. *Comm. Math. Phys.*, 280(3):843–858, 2008.
- [34] V. Isakov. Increasing stability for the schrödinger potential from the dirichlet-to neumann map. *Discrete Contin. Dyn. Syst. Ser. S*, 4(3):631–640, 2011.
- [35] V. Isakov. *Inverse problems for partial differential equations*, volume 127 of *Applied Mathematical Sciences*. Springer, Cham, third edition, 2017.
- [36] V. Isakov and J.-N. Wang. Uniqueness and increasing stability in electromagnetic inverse source problems. *J. Differential Equations*, 283:110–135, 2021.
- [37] O. Imanuvilov and M. Yamamoto. Inverse boundary value problem for the Schrödinger equation in a cylindrical domain by partial boundary data. *Inverse Problems*, 29(4):045002, 2013.
- [38] O. Imanuvilov and M. Yamamoto. Global uniqueness in inverse boundary value problems for the Navier–Stokes equations and Lamé system in two dimensions. *Inverse Problems*, 31(3):035004, 2015.
- [39] F. John. Continuous dependence on data for solutions of partial differential equations with a prescribed bound. *Comm. Pure Appl. Math.*, 13(4):551–585, 1960.
- [40] Y. Kian. Recovery of non-compactly supported coefficients of elliptic equations on an infinite waveguide. *J. Inst. Math. Jussieu*, 19(5):1573–1600, 2020.
- [41] A. Kirsch and R. Kress. Uniqueness in inverse obstacle scattering. *Inverse problems*, 9(2):285–299, 1993.
- [42] K. Krupchyk, M. Lassas, and G. Uhlmann. Inverse problems with partial data for a magnetic Schrödinger operator in an infinite slab and on a bounded domain. *Comm. Math. Phys.*, 312(1):87–126, 2012.
- [43] Y. Kurylev, M. Lassas, and G. Uhlmann. Inverse problems for Lorentzian manifolds and non-linear hyperbolic equations. *Invent. Math.*, 212(3):781–857, 2018.
- [44] Y. Kian and L. Oksanen. Recovery of time-dependent coefficient on Riemannian manifold for hyperbolic equations. *Int. Math. Res. Not. IMRN*, (16):5087–5126, 2019.
- [45] K. Krupchyk and L. Päivärinta. A Borg-Levinson theorem for higher order elliptic operators. *Int. Math. Res. Not. IMRN*, (6):1321–1351, 2012.
- [46] H. Koch, A. Rüländ, and M. Salo. On instability mechanisms for inverse problems. *Ars Inveniendi Analytica*, (7):93 pp., 2021.
- [47] K. Krupchyk. Inverse transmission problems for magnetic Schrödinger operators. *Int. Math. Res. Not. IMRN*, (1):65–164, 2014.
- [48] C. Kenig and M. Salo. The Calderón problem with partial data on manifolds and applications. *Anal. PDE*, 6(8):2003–2048, 2013.
- [49] C. Kenig and M. Salo. Recent progress in the Calderón problem with partial data. In *Inverse problems and applications*, volume 615 of *Contemp. Math.*, pages 193–222. Amer. Math. Soc., Providence, RI, 2014.
- [50] C. Kenig, J. Sjöstrand, and G. Uhlmann. The Calderón problem with partial data. *Ann. of Math. (2)*, 165(2):567–591, 2007.
- [51] K. Krupchyk and G. Uhlmann. Partial data inverse problems for semilinear elliptic equations with gradient nonlinearities. *Math. Res. Lett.*, 27(6):1801–1824, 2020.
- [52] K. Krupchyk and G. Uhlmann. A remark on partial data inverse problems for semilinear elliptic equations. *Proc. Amer. Math. Soc.*, 148(2):681–685, 2020.
- [53] L. Liang. Increasing stability for the inverse problem of the Schrödinger equation with the partial Cauchy data. *Inverse Probl. Imaging*, 9(2):469–478, 2015.
- [54] M. Lassas, T. Liimatainen, Y.-H. Lin, and M. Salo. Inverse problems for elliptic equations with power type nonlinearities. *J. Math. Pures Appl. (9)*, 145:44–82, 2021.
- [55] M. Lassas, T. Liimatainen, Y.-H. Lin, and M. Salo. Partial data inverse problems and simultaneous recovery of boundary and coefficients for semilinear elliptic equations. *Rev. Mat. Iberoam.*, 37(4):1553–1580, 2021.
- [56] T. Liimatainen, Y.-H. Lin, M. Salo, and T. Tyni. Inverse problems for elliptic equations with fractional power type nonlinearities. *J. Differential Equations*, 306:189–219, 2022.
- [57] P. Lax and R. Phillips. *Scattering theory*. Academic press, 1990.
- [58] X. Li and G. Uhlmann. Inverse problems with partial data in a slab. *Inverse Probl. Imaging*, 4(3):449–462, 2010.

- [59] R.-Y. Lai, G. Uhlmann, and J.-N. Wang. Inverse boundary value problem for the Stokes and the Navier–Stokes equations in the plane. *Arch. Ration. Mech. Anal.*, 215(3):811–829, 2015.
- [60] M. Lassas, G. Uhlmann, and Y. Wang. Inverse problems for semilinear wave equations on Lorentzian manifolds. *Comm. Math. Phys.*, 360(2):555–609, 2018.
- [61] X. Li and J.-N. Wang. Determination of viscosity in the stationary Navier–Stokes equations. *J. Differential Equations*, 242(1):24–39, 2007.
- [62] S. Liu and Y. Yang. Determine a magnetic Schrödinger operator with a bounded magnetic potential from partial data in a slab. *Appl. Math. Optim.*, 83(1):277–296, 2021.
- [63] N. Mandache. Exponential instability in an inverse problem for the Schrödinger equation. *Inverse Problems*, 17(5):1435–1444, 2001.
- [64] A. Pal Choudhury and H. Heck. Increasing stability for the inverse problem for the Schrödinger equation. *Math. Methods Appl. Sci.*, 41(2):606–614, 2018.
- [65] L. Potenciano-Machado. Optimal stability estimates for a magnetic Schrödinger operator with local data. *Inverse Problems*, 33(9):095001, 2017.
- [66] V. Pohjola. An inverse problem for the magnetic Schrödinger operator on a half space with partial data. *Inverse Probl. Imaging*, 8(4):1169–1189, 2014.
- [67] R. Potthast. *Point sources and multipoles in inverse scattering theory*. Chapman and Hall/CRC, 2001.
- [68] D. Subbarayappa and V. Isakov. On increased stability in the continuation of the Helmholtz equation. *Inverse Problems*, 23(4):1689–1697, 2007.
- [69] J. Sylvester and G. Uhlmann. A global uniqueness theorem for an inverse boundary value problem. *Ann. of Math.*, pages 153–169, 1987.
- [70] Z. Sun and G. Uhlmann. Inverse problems in quasilinear anisotropic media. *Amer. J. Math.*, 119(4):771–797, 1997.
- [71] Z. Sun. On a quasilinear inverse boundary value problem. *Math. Z.*, 221(2):293–305, 1996.