

JYX



This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Engelstein, Max; Kauranen, Aapo; Prats, Martí; Sakellaris, Georgios; Sire, Yannick

Title: Minimizers for the Thin One-Phase Free Boundary Problem

Year: 2021

Version: Accepted version (Final draft)

Copyright: © 2021 Wiley Periodicals LLC.

Rights: In Copyright

Rights url: <http://rightsstatements.org/page/InC/1.0/?language=en>

Please cite the original version:

Engelstein, M., Kauranen, A., Prats, M., Sakellaris, G., & Sire, Y. (2021). Minimizers for the Thin One-Phase Free Boundary Problem. *Communications on Pure and Applied Mathematics*, 74(9), 1971-2022. <https://doi.org/10.1002/cpa.22011>

Minimizers for the thin one-phase free boundary problem

Max Engelstein ^{*}, Aapo Kauranen [†], Martí Prats [‡], Georgios Sakellaris [§], Yannick Sire [¶]

February 1, 2023

Abstract

We consider the “thin one-phase” free boundary problem, associated to minimizing a weighted Dirichlet energy of the function in \mathbb{R}_+^{n+1} plus the area of the positivity set of that function in \mathbb{R}^n . We establish full regularity of the free boundary for dimensions $n \leq 2$, prove almost everywhere regularity of the free boundary in arbitrary dimension and provide content and structure estimates on the singular set of the free boundary when it exists. All of these results hold for the full range of the relevant weight.

While our results are typical for the calculus of variations, our approach does not follow the standard one first introduced in [AC81]. Instead, the nonlocal nature of the distributional measure associated to a minimizer necessitates arguments which are less reliant on the underlying PDE.

Contents

1	Introduction	2
2	Preliminaries	7
3	Compactness of minimizers	9
3.1	Caccioppoli Inequality	9
3.2	Compactness	11
4	Monotonicity formula and some immediate consequences	13
4.1	Dimension reduction	13
4.2	Upper semicontinuity	15
5	Measure-theoretic properties	16
5.1	Finite perimeter	16
5.2	Energy gap	19
6	Full regularity in \mathbb{R}^{2+1}	21

^{*}ME (University of Minnesota Twin Cities, Minnesota): maxe@mit.edu

[†]AK (Universitat Autònoma de Barcelona-BGSMath, Catalonia): aapo.p.kauranen@jyu.fi

[‡]MP (Universitat Autònoma de Barcelona, Catalonia): mprats@mat.uab.cat

[§]GS (Universitat Autònoma de Barcelona, Catalonia): gsakellaris@mat.uab.cat

[¶]YS (Johns Hopkins University, Maryland): sire@math.jhu.edu

7	Uniform bounds around the free boundary	22
7.1	Uniform non-degeneracy	22
7.2	Behavior of the distributional fractional Laplacian	24
7.3	Uniform Hölder character	28
7.4	Lower estimates for the distributional fractional Laplacian	30
8	Rectifiability of the singular set	31
8.1	Quantitative stratification for minimizers to \mathcal{J}	32
8.2	The Refined Covering Theorem	34
8.3	Tools for bad balls: key dichotomy	36
8.4	Tools for good balls: packing estimates and GMT	37
A	Relation with the nonlocal Bernoulli problem	39
B	Uniqueness of extensions	40

1 Introduction

This article is devoted to the study of the regularity properties of a weighted version of the thin one-phase problem. More precisely we investigate even, nonnegative minimizers of the following functionals: denote $x \in \mathbb{R}^{n+1}$ by $x = (x', y) \in \mathbb{R}^n \times \mathbb{R}$, and for $\beta \in (-1, 1)$ we define

$$\mathcal{J}(v, \Omega) := \int_{\Omega} |y|^{\beta} |\nabla v|^2 dx + m(\{v > 0\} \cap \mathbb{R}^n \cap \Omega), \quad (1.1)$$

where m stands for the n -dimensional Lebesgue measure. Here, and throughout the paper, the integration is done with respect to the $(n+1)$ -dimensional Lebesgue measure unless stated otherwise. This functional is finite for open sets, Ω , and functions in the weighted Hilbert space,

$$H^1(\beta, \Omega) := \{v \in L^2(\Omega; |y|^{\beta}) : \nabla v \in L^2(\Omega; |y|^{\beta})\},$$

equipped with the usual weighted norm.

Our main concern is to investigate fine regularity properties of the free boundary of minimizers v of (1.1), that is the set,

$$F(v) := \partial_{\mathbb{R}^n} \{v(x, 0) > 0\} \cap \Omega.$$

Since the free boundary lies on a codimension 1 subspace of the ambient space \mathbb{R}^{n+1} , such a problem is called a *thin one-phase free boundary problem*. This type of free boundary problem has been investigated for the first time by Caffarelli, Roquejoffre and the last author in [CRS10b] in relation with the theory of semi-permeable membranes (see, e.g., [DL76]). As we will describe later this is an analogue of the classical one-phase problem (also called the Bernoulli problem) but for the fractional Laplacian.

The Bernoulli problem was first treated in a rigorous mathematical way by Alt and Caffarelli in the seminal paper [AC81]: in the Bernoulli problem we consider minimizers of (1.1) where $\beta = 0$ and the second term is replaced by $\mathcal{L}^{n+1}(\{v > 0\} \cap \Omega)$ (where \mathcal{L}^{n+1} stands for the Lebesgue measure in \mathbb{R}^{n+1}). In particular, for the Bernoulli problem, the free boundary fully sits in the ambient space, \mathbb{R}^{n+1} . In [AC81], the authors provided a general strategy to attack this type of problem. Out of necessity we needed to modify this blueprint in several substantial ways (see below for a more detailed comparison). For more information on the one-phase problem (and some of its variants) we refer to the book of Caffarelli and Salsa (and references therein) [CS05], and to the more recent survey of De Silva, Ferrari and Salsa [DSFS19].

As noticed in [CRS10b], problem (1.1) is related in a tight way to the *standard* one-phase free boundary problem but with the Dirichlet energy replaced by the Gagliardo semi-norm $[u]_{\dot{H}^\alpha}$, for $\alpha = \frac{1-\beta}{2} \in (0, 1)$. This connection suggests that the thin one-phase problem is actually intrinsically a *nonlocal* problem, though the energy in (1.1) is clearly local.

Connection with the fractional one-phase problem

As previously mentioned, the functional \mathcal{J} introduced by Caffarelli, Roquejoffre and the last author in [CRS10b] is a local version of the following nonlocal free boundary problem: given a function $f \in L^1_{\text{loc}}(\mathbb{R}^n)$ with suitable decay at infinity, we can define its fractional Laplacian at $x \in \mathbb{R}^n$ by

$$(-\Delta)^\alpha f(x) = c_{n,\alpha} p.v. \int_{\mathbb{R}^n} \frac{f(x) - f(\xi)}{|x - \xi|^{n+2\alpha}} d\xi.$$

At the formal level, we are interested in solutions of the free boundary problem

$$\begin{cases} (-\Delta)^\alpha f = 0 & \text{in } \Omega \cap \{f > 0\}, \\ \partial_\nu^\alpha f = A & \text{on } \Omega \cap F(f), \end{cases} \quad (1.2)$$

where $\partial_\nu^\alpha f(x) := \lim_{\Omega \cap \{f > 0\} \ni \xi \rightarrow x} \frac{f(\xi) - f(x)}{((\xi - x) \cdot \nu(x))^\alpha}$ and where f satisfies a given ‘‘Dirichlet boundary condition’’ on the complement of Ω .

As in the case of the classical Laplacian (see [AC81]), we are interested in obtaining equation (1.2) as the Euler-Lagrange equation of a certain functional. Given a locally integrable function f , consider its fractional Sobolev energy

$$[f]_{\dot{H}^\alpha(\mathbb{R}^n)} := \iint_{\mathbb{R}^{2n}} \frac{|f(x) - f(\xi)|^2}{|x - \xi|^{n+2\alpha}} d\xi dx.$$

Since we want to study competitors which vary only in a certain domain Ω , it is natural to consider only the integration region which may suffer variations when changing candidates. Thus, we define the energy

$$J(f, \Omega) := c_{n,\alpha} \iint_{\mathbb{R}^{2n} \setminus (\Omega^c)^2} \frac{|f(x) - f(\xi)|^2}{|x - \xi|^{n+2\alpha}} d\xi dx + m(\{f > 0\} \cap \Omega). \quad (1.3)$$

We say that $f \in L^1_{\text{loc}}$ is a minimizer of J in Ω if $J(f, \Omega)$ is finite and $J(f, \Omega) \leq J(g, \Omega)$ for every g satisfying that $f - g \in \dot{H}^\alpha(\mathbb{R}^n)$ and such that $f(x) = g(x)$ for almost every $x \in \Omega^c$. We say that f is a global minimizer if it is a minimizer for every open set $\Omega \subset \mathbb{R}^n$. Note that both terms in (1.3) are in competition, since a minimizer of the fractional Sobolev energy in Ω is α -harmonic and, thus, if it is non-negative outside of Ω it is strictly positive inside of Ω , maximizing the second term.

Consider now the Poisson kernel for fixed $n \in \mathbb{N}$ and $0 < \alpha < 1$

$$P_y(\xi) := P_{n,\alpha}(\xi, y) = c_{n,\alpha} \frac{|y|^{2\alpha}}{|(\xi, y)|^{n+2\alpha}} \quad \text{for every } (\xi, y) \in \mathbb{R}^n \times \mathbb{R}. \quad (1.4)$$

The Poisson extension of $f \in L^1_{\text{loc}}(\mathbb{R}^n)$ is given by

$$u(x', y) := f * P_y(x') = \int_{\mathbb{R}^n} P_{n,\alpha}(\xi, y) f(x' - \xi) d\xi \quad \text{for every } (x', y) \in \mathbb{R}^n \times \mathbb{R}. \quad (1.5)$$

By [CS07], with a convenient choice of the constant one gets

$$\lim_{y \searrow 0} y^{1-2\alpha} u_y(x', y) = -(-\Delta)^\alpha f(x')$$

in every point where f is regular enough. Moreover, the extension satisfies the localized equation $\nabla \cdot (|y|^\beta \nabla u) = 0$ weakly, away from $\mathbb{R}^n \times \{0\}$. The whole point is that local minimizers of (1.3) can be extended via the previous Poisson kernel P_y to (even) minimizers of (1.1) (see the Appendix for a precise statement). Therefore, the thin one-phase problem appears as a “localization” of the one-phase problem for the fractional Laplacian. Notice that, and this is of major importance for us, this localization technique does not carry over to other types of nonlocal operators besides pure powers of second-order elliptic operators. This is a major drawback of the theory, in the sense that, at the moment, it seems to be impossible to tackle one-phase problems involving more general operators than the fractional Laplacian. The main point is we do not know how to prove any kind of monotonicity for general integral operators.

This connection between the nonlocal analogue of the Bernoulli problem and our thin one-phase problem allows us to simplify several arguments by working in the purely nonlocal setting. However, this underlying nonlocality is also the reason why several results, which came more easily in the setting of [AC81], are non-trivial or substantially harder for us. For example, perturbations of solutions need to take into account long range effects which makes classical, local, perturbation arguments much more difficult.

In the paper [CRS10b], the authors proved basic properties of the minimizers for the functional \mathcal{J} such as optimal regularity, non-degeneracy near the free boundary, and positive densities of phases. Also they provided an argument for $n = 2$ showing that Lipschitz free boundaries are C^1 . A feature of the functional \mathcal{J} is that the weight $|y|^\beta$ is either degenerate or singular at $\{y = 0\}$ (except in the case $\beta = 0$). Such weights belong to the Muckenhoupt class A_2 and the seminal paper of Fabes, Kenig and Serapioni [FKS82] investigated regularity issues for elliptic PDEs involving such weights (among other things). After that, [DSS14] proved an ε -regularity result and [All12] showed the existence of a monotonicity formula for this setting.

In the case $\beta = 0$, the problem is still degenerate in the sense that derivatives near the free boundary blow up. The case $\beta = 0$ has been thoroughly investigated in the series of papers by De Silva, Savin and Roquejoffre [DR12, DS12, DS15].

The main goal of our paper is to provide a full picture of the regularity of the free boundary for any power $\beta \in (-1, 1)$, both in terms of measure-theoretic statements and partial (or full) regularity results. From this point of view our contribution is a complement of the paper by De Silva and Savin [DS15] for $\beta = 0$. It has to be noticed that the standard approach to regularity of Lipschitz free boundaries as developed by Caffarelli (see the monograph [CS05]) does not seem to work in our setting.

Our approach to regularity

In [AC81] (and many subsequent works), the minimizing property of the solution is used to prove that the distributional Laplacian of that solution is an Ahlfors-regular measure supported on the free boundary. This implies (amongst other things) that the free boundary is a set of (locally) finite perimeter, and thus almost every point on the free boundary has a measure theoretic tangent. One can then work purely with the weak formula (i.e. the analogue of (1.2)) to prove a “flat implies smooth” result which, together with the existence almost everywhere of a measure theoretic tangent, has as a consequence that the free boundary is almost everywhere a smooth graph and the free boundary condition in (1.2) holds in a classical sense at the smooth points.

A similar “flat implies smooth” result exists in our context (this is essentially due to De Silva, Savin and the last author, [DSS14], see Theorem 2.4 below). However, showing that the free boundary is the boundary of a set of finite perimeter proves to be much more difficult. Due to the nonlocal nature of the problem, $-\operatorname{div}(|y|^\beta \nabla u)$ (considered as a distribution) is not supported on the free boundary. Furthermore, the scaling of this measure does not allow us to conclude that the free boundary has the correct dimension (much less that it is Ahlfors regular).

To prove finite perimeter, we take the following approach inspired by the work of de Silva and Savin: after establishing some preliminaries we prove crucial compactness results. This, along with a monotonicity formula originally due to Allen [All12] allows us to run a dimension reduction argument in the vein of Federer or (in the context of free boundary problems) Weiss [Wei99]. With this tool in hand, we show that the set of points at which no blow-up is flat is a set of lower dimension. Locally finite perimeter and regularity for the reduced boundary then follow from a covering argument and some standard techniques.

Here and throughout the paper, we will denote the ball of radius r in \mathbb{R}^{n+1} centered at the origin by B_r , and $B'_r := B_r \cap \mathbb{R}^n \times \{0\}$. Moreover, for the definition of \mathbf{H}^β , see Section 2. We may then summarize our regularity results in the following theorem.

Theorem 1.1. *[Main Regularity Theorem] Let $u \in \mathbf{H}^\beta(B_1)$ be a (non-negative, even) local minimizer of \mathcal{J} in $B_1 \subset \mathbb{R}^{n+1}$. Let $B'_{1,+}(u) := \{x = (x', 0) \in B_1 : u(x) > 0\}$, let $F(u)$ be the boundary of $B'_{1,+}(u)$ inside of B'_1 and assume that $0 \in F(u)$. Then,*

1. $B'_{1,+}(u)$ (as a subset of $\mathbb{R}^n \times \{0\}$) is a set of locally finite perimeter in B'_1 .
2. We can write the free boundary as a disjoint union $F(u) = \mathcal{R}(u) \cup \Sigma(u)$, where $\mathcal{R}(u)$ is open inside $F(u)$, and for $x \in \mathcal{R}(u)$ there exists an $r_x > 0$ such that $B(x, r_x) \cap F(u)$ can be written as the graph of a $C^{1,s}$ -continuous function.
3. Furthermore, the set $\Sigma(u)$ is of Hausdorff dimension $\leq n-3$ (and, therefore, of \mathcal{H}^{n-1} -measure zero). In particular, for $n \leq 2$, $\Sigma(u)$ is empty, and moreover, if $n = 3$ then $\Sigma(u)$ is discrete.

The constants (implicit in the set of finite perimeter, and the Hölder continuity of the functions whose graph gives the free boundary) depend on n and β but not on $\|u\|_{\mathbf{H}^\beta(B_1)}$.

As usual $\Sigma(u) \subset F(u)$ is called the *singular set* of the free boundary: the set of points around which $F(u)$ cannot be parameterized as a smooth graph and all the blow-ups will be non-trivial minimal cones, see Theorem 2.4.

Our second contribution concerns the structure and size of the singular set. It builds on recent major works on quantitative stratification [NV17], extended to free boundary problems (in particular the one-phase problem) by Edelen and the first author [EE19].

Theorem 1.2. *Let $u \in \mathbf{H}^\beta(B_1)$ be a (non-negative, even) local minimizer of \mathcal{J} in B_1 and $0 \in F(u)$. Let $B'_{1,+}(u) := \{x = (x', 0) \in B_1 : u(x) > 0\}$ and $F(u)$ be the boundary of $B'_{1,+}(u)$ inside B'_1 . Then, there exists a $k_\alpha^* \geq 3$ such that $\Sigma(u)$ is $(n - k_\alpha^*)$ -rectifiable and*

$$\mathcal{H}^{n-k_\alpha^*}(\Sigma(u) \cap D) \leq C_{n,\alpha,\text{dist}(D,\partial B_1)} \quad \text{for every } D \subset\subset B_1.$$

In [DJ09], De Silva and Jerison constructed a singular minimizer for the Alt-Caffarelli one-phase problem in dimension 7, giving the dimension bound $k^* \leq 8$ in the previous theorem in this case (see [EE19]). This result is not known for the thin one-phase problem. The reason is that the one-phase problem, seen from the nonlocal point of view involving the fractional Laplacian, is related to the so-called nonlocal minimal surfaces introduced by Caffarelli, Roquejoffre and Savin [CRS10a]. Indeed, in [SV12], the authors proved that a fractional version of Allen-Cahn equation converges variationally to the standard perimeter functional for $\alpha \geq 1/2$ and to the so-called nonlocal minimal surfaces for $\alpha < 1/2$. We can then conjecture the bound $k_\alpha^* \leq 8$ for $\alpha \geq 1/2$ by analogy with the result for the standard one-phase problem but the bound for $\alpha < 1/2$ is not clear at all. However, one knows that there is no singular cone in dimension 2 for nonlocal minimal surfaces [SV13] and that the Bernstein problem is known for those in dimensions 2 and 3 [FV17].

We would like also to make a last remark about a result which is of purely nonlocal nature. In the case of the one-phase problem, one can show that the distributional Laplacian is a Radon

measure along the free boundary. In the case of the thin one-phase free boundary problem, due to the nonlocality of the problem, such a behavior does not happen in the sense that we will show that the fractional Laplacian is an absolutely continuous measure with respect to n -dimensional Lebesgue measure with a precise behavior. This phenomenon is of purely nonlocal nature and similar to the fact that the fractional harmonic measure is of trivial nature. More precisely, every minimizer u satisfies $\nabla \cdot (|y|^\beta \nabla u) = 0$ weakly, away from $\mathbb{R}^n \cap \{u \leq 0\}$. Thus, equation (1.2) above can be understood as an Euler-Lagrange equation for the functional \mathcal{J} in the sense that the restriction to \mathbb{R}^n of a given minimizer u in $\Omega \subset \mathbb{R}^{n+1}$, harmonic away from $\mathbb{R}^n \times \{0\}$ and with asymptotic behavior $u(x, y) = \mathcal{O}(|(x, y)|^\alpha)$ is always a solution to (1.2) for $A = A(\alpha)$ at “nice” points of the free boundary.

A brief summary of this paper follows. In Sections 3 and 4 we discuss compactness of minimizers and we recall Allen’s monotonicity formula to derive some immediate consequences. In Section 5 we show that the positive phase is a set of locally finite perimeter, establishing the first part of Theorem 1.1 (modulo energy bounds), and we show that the singular set can be identified using the Allen-Weiss density. Section 6 is devoted to deducing full regularity of minimizers in \mathbb{R}^{2+1} concluding the proof of Theorem 1.1.

Once we have established the finite perimeter, in Section 7 we remove the dependence of the estimates on the energy of the minimizer in the previous theorems, using a rather subtle argument which combines results from all the previous sections. A crucial step is to analyze some basic properties of the distributional fractional Laplacian of our minimizer. As stated above this analysis will not be enough to establish that the positivity set of the minimizer is a set of locally finite perimeter. We believe that many of these results may be of independent interest. For example, corresponding results for the classical Bernoulli problem have been used to understand free boundary problems for harmonic measure (see [KT03]).

Finally, Section 8 is devoted to the proof of Theorem 1.2.

Notation

We denote the constants that depend on the dimension n , α and perhaps some other fixed parameters which are clear from the context by C . Their value may change from an occurrence to another. On the other hand, constants with subscripts as C_0 retain their values along the text. For $a, b \geq 0$, we write $a \lesssim b$ if there is $C > 0$ such that $a \leq Cb$. We write $a \approx b$ to mean $a \lesssim b \lesssim a$.

Let u be a continuous function in \mathbb{R}^{n+1} . Then we write $\Omega_+(u) := \Omega \cap \{u > 0\}$, and we denote the zero phase, the positive phase and the free boundary by

$$\begin{aligned}\Omega_0(u) &:= \{x \in \mathbb{R}^n \times \{0\} : u(x) = 0\}^\circ, \\ \Omega'_+(u) &:= \Omega_+ \cap (\mathbb{R}^n \times \{0\}) = \{x \in \mathbb{R}^n \times \{0\} : u(x) > 0\}, \text{ and} \\ F(u) &:= F_\Omega(u) = \partial(\Omega_+(u) \cap \mathbb{R}^n \times \{0\}) \cap \Omega,\end{aligned}$$

respectively. Here both the boundary and the interior are taken with respect to the standard topology in \mathbb{R}^n . Note that $\mathbb{R}^n \times \{0\}$ is the disjoint union of $\Omega_0(u)$, $\Omega'_+(u)$ and $F(u)$ whenever u is non-negative. We also call $F_{\text{red}}(u) = F_{\text{red}, \Omega}(u)$ the points of $F_\Omega(u)$ where the free boundary is expressed locally as a C^1 surface. Finally, let $\Sigma(u) = \Sigma_\Omega(u) = F_\Omega(u) \setminus F_{\text{red}, \Omega}(u)$. In general we will write $\Omega' := \Omega \cap (\mathbb{R}^n \times \{0\})$.

Throughout the paper we will often fix $\beta \in (-1, 1)$ but then refer to $\alpha \in (0, 1)$ or vice versa. These two numbers are always connected by the relationship $\alpha = \frac{1-\beta}{2}$.

2 Preliminaries

In this section, we provide the known results concerning the problem under consideration. We say that a function u is *even* if it is symmetric with respect to the hyperplane $\mathbb{R}^n \times \{0\}$, that is, $u(x', y) = u(x', -y)$. The function spaces that we will consider are the following

$$\mathbf{H}^\beta(\Omega) := \{u \in H^1(\beta, \Omega) : u \text{ is even and non-negative}\}$$

and

$$\mathbf{H}_{\text{loc}}^\beta(\Omega) := \{u \in L_{\text{loc}}^2(\Omega) : u \in \mathbf{H}^\beta(B) \text{ for every ball } B \subset\subset \Omega\}.$$

We will omit Ω in the notation when it is clear from the context.

Definition 2.1. *We say that a function $u \in \mathbf{H}_{\text{loc}}^\beta(\Omega)$ is a (local) minimizer of \mathcal{J} in a domain Ω if for every ball $B \subset\subset \Omega$ and for every function $v \in \mathbf{H}^\beta(B)$ such that the traces $v|_{\partial B} \equiv u|_{\partial B}$, the inequality*

$$\mathcal{J}(u, B) \leq \mathcal{J}(v, B)$$

holds.

As usual for several free boundary problems, it is a natural question to exhibit a particular (global) solution so that one gets an idea of the qualitative properties of general solutions. Let us consider the following function: for every $x \in \mathbb{R}^n$ let

$$f_{n,\alpha}(x) := c_{n,\alpha}(x_n)_+^\alpha,$$

where $a_+ = \max\{0, a\}$. If $n = 1$, $f_{1,\alpha}$ is a solution to (1.2) for a convenient choice of $c_{1,\alpha}$ (see [BV16, Theorem 3.1.4]). In fact one can see that the same is true for $n \geq 1$ using Fubini's Theorem conveniently, with

$$-(-\Delta)^\alpha f_{n,\alpha}(x) = c_{n,\alpha}(x_n)_-^{-\alpha}, \quad (2.1)$$

where $a_- = \max\{0, -a\}$.

As a toy question we wonder whether the trivial solutions are minimizers. Indeed, this is the case, as we will see later in Section 4.1.

Proposition 2.2. *Let $n \in \mathbb{N}$ and $0 < \alpha < 1$. Then the trivial solution $u_{n,\alpha} := f_{n,\alpha} * P_y$ is a minimizer of \mathcal{J} in every ball $B \subset \mathbb{R}^{n+1}$.*

Next we collect the main properties of minimizers in the unit ball proven in [CRS10b, Theorems 1.1-1.4, Proposition 3.3 and Corollary 3.4].

Theorem 2.3. *If $u \in \mathbf{H}^\beta(B_1)$ is a minimizer of \mathcal{J} in $\Omega = B_1$ with $\|u\|_{\mathbf{H}^\beta(B_1)} := \|\nabla u\|_{L^2(B_1, |y|^\beta)} \leq E_0$ and $x_0 \in F(u) \cap B_{\frac{1}{2}}$, then it satisfies*

P1: Optimal regularity (see [CRS10b, Theorem 1.1]): $\|u\|_{\dot{C}^\alpha(B_{1/2})} \leq C(1 + E_0)$.

P2: Nondegeneracy (see [CRS10b, Theorem 1.2]): $u(x) \geq C \text{dist}(x, F(u))^\alpha$ for $x \in B'_{\frac{1}{2},+}$.

P3: Interior corkscrew condition (see [CRS10b, Proposition 3.3]): there exists $x_+ \in B'_r(x_0)$ so that $B'(x_+, C_0 r) \subset \Omega'_+(u)$.

P4: Positive density (see [CRS10b, Theorem 1.3]): $|\Omega_0 \cap B'_r(x_0)| \gtrsim r^n$.

P5: Blow-ups are minimizers (see [CRS10b, Corollary 3.4]): The limit of a blow-up sequence $u_k(x) := \frac{u(x_0 + \rho_k x)}{\rho_k^\alpha}$ converging weakly in $H^1(\beta, B_1)$ and uniformly is a global minimizer.

P6: Normal behavior at the free boundary (see [CRS10b, Theorem 1.4]): the boundary condition in (1.2) is satisfied at every point on the free boundary with a measure theoretic normal (see [EG15]) for a prescribed value of A .

All the constants depend on n and α ; and also on E_0 except for the ones in P1 and P2.

A major tool in the present paper is an ϵ -regularity result, i.e. in the language of free boundaries a statement of the type “flatness implies smoothness”. In [DSS14], the authors proved such an ϵ -regularity result for viscosity solutions to the overdetermined system associated to minimizers of \mathcal{J} . Here we establish that all local minimizers are in fact viscosity solutions. While this verification may be standard for experts in the field, we include it here for the sake of completeness.

Theorem 2.4 (ϵ -regularity). *There exists $\epsilon > 0$ depending only on n , α and E_0 such that for every non-negative, even minimizer u of the energy (1.1) on a ball $B \subset \mathbb{R}^{n+1}$ with $\|u\|_{\mathbf{H}^\beta(B)} \leq E_0 r(B)^{\frac{n}{2}}$ and*

$$\{(x, 0) \in B : x_n \leq -\epsilon\} \subset B_0(u) \subset \{(x, 0) \in B : x_n \leq \epsilon\}, \quad (2.2)$$

we have that $F(u) \in C_{\text{loc}}^{1,\gamma}(\frac{1}{2}B)$, with $0 < \gamma < 1$.

Note that the dependence on E_0 will be removed in Section 7.

Proof. We say that u is a *viscosity solution* to

$$\begin{cases} \nabla \cdot (|y|^\beta \nabla u) = 0 & \text{in } B_1^+(u), \\ \lim_{t \rightarrow 0^+} \frac{u(x_0 + t\nu(x_0), 0)}{t^\alpha} = 1, & \text{for } (x_0, 0) \in F(u), \end{cases} \quad (2.3)$$

if

- i) $u \in C(B_1)$, $u \geq 0$,
- ii) $u \in C_{\text{loc}}^{1,1}(B_{1,+}(u))$, u is even and it solves $\nabla \cdot (|y|^\beta \nabla u) = 0$ in the viscosity sense, and
- iii) any strict *comparison subsolution* (resp. *supersolution*) cannot touch from below (resp. from above) at a point $(x_0, 0) \in F(u)$.

We claim that

$$\text{every non-negative even minimizer is a viscosity solution.} \quad (2.4)$$

Conditions (i) and (ii) have been verified in [DSS14, Vit18]. To verify our claim it suffices to prove condition (iii) above: that any strict comparison subsolution cannot touch u from below at a point $(x_0, 0) \in F(u)$. The analogous claim for strict comparison supersolutions will follow in the same way.

Let us recall (see, e.g. Definition 2.2 in [DSS14]), that $w \in C(B_1)$ is a strict comparison subsolution (resp. supersolution) to (2.3) if

- a) $w \geq 0$,
- b) w is even with respect to $\{y = 0\}$,
- c) $w \in C^2(\{w > 0\})$,
- d) $\text{div}(|z|^\beta \nabla w) \geq 0$ in $B_1 \setminus \{y = 0\}$,

e) $F(w)$ is locally given by the graph of a C^2 function and for any $x_0 \in F(w)$ we may write

$$w(x, y) = aU((x - x_0) \cdot \nu(x_0), y) + o(\|(x - x_0, y)\|^\alpha), \quad (x, y) \rightarrow (x_0, 0). \quad (2.5)$$

Here U is the extension of the trivial solution (see [DSS14]), $\nu(x_0)$ is the unit normal to $F(w)$ considered as a subset of \mathbb{R}^n pointing into $\{w > 0\}$ and $a \geq 1$.

f) Furthermore, either the inequality is strict in d), or $a > 1$ in e).

So assume that $w \geq u$ where w is a strict comparison subsolution and u is some minimizer and that $w = u$ at $(x_0, 0) \in F(u)$. Since $u(x_0, 0) = 0$ it follows that $(x_0, 0) \in F(w)$ and with a harmless rotation we can guarantee that $\nu((x_0, 0)) = e_n$. We want to show that e_n is also the measure theoretic unit normal to $F(u)$. Indeed, since $F(w)$ is C^2 there must exist a ball $B \subset \{w > 0\}$ which is tangent to $F(w)$ at $(x_0, 0)$. It must then be that case that $B \subset \{u > 0\}$ as well. Thus $(x_0, 0) \in F(u)$ has a tangent ball from the inside which, by [CRS10b] Proposition 4.5 implies that u has the asymptotic expansion

$$u(x, y) = U((x - x_0) \cdot \nu(x_0), y) + o(\|(x - x_0, y)\|^\alpha), \quad (x, y) \rightarrow (x_0, 0).$$

If $u \geq w$ this implies that w must satisfy the expansion in (2.5) with $a = 1$ at the point x_0 . This, in turn, implies that $\operatorname{div}(|z|^\beta \nabla w) > 0$ in $B_1 \setminus \{y = 0\}$ (by the definition of a strict subsolution). Furthermore, since $w \in C^2$ where $\{w > 0\}$ we can guarantee that $\operatorname{div}(|z|^\beta \nabla w) \geq 0$ in all of $B_1 \cap \{w > 0\}$.

Let us return to the ball B which is a subset of $\{u > 0\}$ and $\{w > 0\}$ and for which $(x_0, 0) \in \bar{B}$. We know that $w - u \neq 0$ in $B \setminus \{y = 0\}$ (this is because w strictly satisfies the differential inequality in B away from $\{y = 0\}$) and we know that $w - u$ is a subsolution in B . Furthermore $(x_0, 0) \in B$ is a strict maximum, so by the Hopf lemma in [CS14, Proposition 4.11] it must be that

$$\lim_{t \downarrow 0^+} \frac{(w - u)(x_0 + t\nu(x_0), 0)}{t^\alpha} > 0.$$

This contradicts the fact u and w both satisfy (2.5) at $(x_0, 0)$ with $a = 1$. Therefore, $(x_0, 0)$ must not have been a touching point and u is indeed a viscosity solution.

Since, u is a viscosity solution, [DSS14, Theorem 1.1] applies and we get the desired ε -regularity. \square

3 Compactness of minimizers

In this section we prove important results on the compactness of minimizers. As we mentioned above, our contribution is that convergent sequences of minimizers also converge in the relevant weighted Sobolev spaces strongly rather than just weakly. This will prove essential to the compactness arguments used in the later sections of this paper.

3.1 Caccioppoli Inequality

First we want to show that the distribution $\lambda := \nabla \cdot (|y|^\beta \nabla u)$ is in fact a Radon measure with support in the complement of the positive phase as long as u is a minimizer. In Section 7 we will come back to this measure to understand its behavior around the free boundary.

Lemma 3.1. *Let $\Omega \subset \mathbb{R}^{n+1}$ be an open set, and let $u \in W_{\text{loc}}^{1,2}(\Omega, |y|^\beta)$ be such that $\nabla \cdot (|y|^\beta \nabla u) = 0$ weakly in $\Omega_+(u)$, i.e., for every $\eta \in C_c^\infty(\Omega_+(u))$,*

$$\langle \nabla \cdot (|y|^\beta \nabla u), \eta \rangle := - \int (|y|^\beta \nabla u) \nabla \eta = 0. \quad (3.1)$$

Then $\lambda := \nabla \cdot (|y|^\beta \nabla u)$ is a positive Radon measure supported on $\{u = 0\}$ and for every $v \in W^{1,2}(\Omega, |y|^\beta) \cap C_c(\Omega)$

$$\int v d\lambda = - \int |y|^\beta \nabla u \cdot \nabla v. \quad (3.2)$$

Proof. Indeed, by (3.1) the quantity

$$- \int |y|^\beta \nabla u \cdot \nabla \zeta = - \int |y|^\beta \nabla u \cdot \nabla \left(\zeta \max \left\{ \min \left\{ 2 - \frac{u}{\varepsilon}, 1 \right\}, 0 \right\} \right) \geq - \int_{\Omega \cap \{0 < u < 2\varepsilon\}} |y|^\beta |\nabla u| |\nabla \zeta| \xrightarrow{\varepsilon \rightarrow 0} 0$$

defines a positive functional on positive $\zeta \in C_c^{0,1}(\Omega)$. Moreover, for compact $K \subset \Omega$, consider a Lipschitz function f_K such that $\chi_K \leq f_K \leq \chi_\Omega$. If $\zeta \in C_c^{0,1}(K)$, by the positivity shown above we obtain

$$- \int |y|^\beta \nabla u \cdot \nabla \zeta \leq - \|\zeta\|_{L^\infty} \int |y|^\beta \nabla u \cdot \nabla f_K \leq C_{K,u} \|\zeta\|_{L^\infty}$$

and, by Hahn-Banach's theorem, we can extend the functional to a positive functional in $C_c(\Omega)$, that is given by integration against a positive Radon measure by the Riesz representation theorem.

The fact that (3.2) holds for all functions in $W^{1,2}(\Omega, |y|^\beta) \cap C_c(\Omega)$ follows by a standard density argument. \square

The Caccioppoli inequality is the first step to proving convergence in a Sobolev sense. It will also be useful when we remove the *a priori* dependence of our results on the Sobolev norm of the minimizer.

Lemma 3.2 (Caccioppoli Inequality). *Let $B \subset \mathbb{R}^{n+1}$ be a ball of radius r centered on $\mathbb{R}^n \times \{0\}$, and let $u \in W^{1,2}(B, |y|^\beta)$ be such that $\nabla \cdot (|y|^\beta \nabla u) = 0$ weakly in $B \cap \{u > 0\}$. Then*

$$\int_{\frac{1}{2}B} |y|^\beta |\nabla u|^2 \leq \frac{4}{r^2} \int_{B \setminus \frac{1}{2}B} |y|^\beta u^2.$$

Proof. Let η be a Lipschitz function such that $\chi_{\frac{1}{2}B} \leq \eta \leq \chi_B$ and with $|\nabla \eta| \leq \frac{1}{r}$. By Lemma 3.1

$$0 = \int_B u \eta^2 d\lambda = \int_B |y|^\beta \nabla u \cdot \nabla (u \eta^2).$$

By the Leibniz rule

$$\int_B |y|^\beta \eta^2 |\nabla u|^2 = - \int_B |y|^\beta 2u \eta \nabla u \cdot \nabla \eta,$$

and using Hölder's inequality we get

$$\int_{\frac{1}{2}B} |y|^\beta |\nabla u|^2 \leq \int_B |y|^\beta \eta^2 |\nabla u|^2 \leq \int_B |y|^\beta 4u^2 |\nabla \eta|^2 \leq \frac{4}{r^2} \int_{B \setminus \frac{1}{2}B} |y|^\beta u^2.$$

\square

Lemma 3.3. *Let $u \in \mathbf{H}^\beta(B_r)$ be a minimizer of \mathcal{J} in B_{2r} and $0 \in F(u)$. Then*

$$r^{-n/2} \|\nabla u\|_{L^2(\frac{1}{2}B_r; |y|^\beta)} \leq r^{-\alpha} \|u\|_{L^\infty(B_r)} \leq \|u\|_{\dot{C}^\alpha(B_r)} \leq C \left(1 + r^{-n/2} \|\nabla u\|_{L^2(B_{2r}; |y|^\beta)} \right).$$

Proof. The first inequality is an immediate consequence of Caccioppoli, the middle estimate is trivial and the last follows from P1 in Theorem 2.3. \square

3.2 Compactness

In the following lemma we prove the compactness of minimizers in the relevant Sobolev spaces. For convenience, we also detail several compactness results which were either already proven in [CRS10b] or are standard consequences of the non-degeneracy estimates in Theorem 2.3. Nevertheless, we include full proofs here for the sake of completeness. We note here (as we did above and will do again below) that while we currently need to assume the uniform bound on the Hölder norm of the functions u_k we can get rid of this assumption in the light of the results of Section 7.

Lemma 3.4 (Compactness results). *Let $\{u_k\}_{k=1}^\infty \subset \mathbf{H}_{\text{loc}}^\beta(\Omega)$ be a sequence of minimizers in a domain $\Omega \subset \mathbb{R}^{n+1}$ with $\|u_k\|_{\dot{C}^\alpha(\Omega)} \leq E_0$ with non-empty free boundary. Then there exists a subsequence converging to some $u_0 \in \mathbf{H}_{\text{loc}}^\beta(\Omega)$ such that for every bounded open set $G \subset\subset \Omega$ we have*

1. $u_k \rightarrow u_0$ in $C^\beta(G)$ for every $\beta < \alpha$,
2. $u_k \rightarrow u_0$ in $L^p(G)$ for every $p \leq \infty$,
3. $\partial\{u_k > 0\} \cap \bar{G} \rightarrow \partial\{u_0 > 0\} \cap \bar{G}$ in the Hausdorff distance,
4. $\chi_{\{u_k > 0\}} \rightarrow \chi_{\{u_0 > 0\}}$ in $L^1(G')$, and
5. $\nabla u_k \rightarrow \nabla u_0$ in $L^p(G; |y|^\beta)$ for every $p \leq 2$.

Proof. The first claim follows from uniform Hölder continuity and compact embeddings of Hölder spaces. The claim (2) follows from (1) easily.

We now prove the third claim. Let $\epsilon > 0$. We will first show that for $x \in \mathbb{R}^n \times \{0\}$ we have

$$d(x, F(u_0)) > \epsilon \Rightarrow d(x, F(u_k)) > \frac{\epsilon}{2} \quad (3.3)$$

for large k . This implies that $F(u_k) \subset \{x: d(F(u_0), x) < 2\epsilon\}$ for k large enough.

Let $B(x, \epsilon) \subset F(u_0)^c$. If u_0 is positive in $B(x, \epsilon)$ then it is bounded from below by a positive number in $B(x, \epsilon/2)$. In this case u_k are also positive in $B(x, \epsilon/2)$ for large k due to uniform convergence in G . Thus $B(x, \epsilon/2) \subset F(u_k)^c$ for large k . If $u \equiv 0$ in $B'(x, \epsilon)$ then due to the uniform convergence we know that for k large enough $u_k < C\epsilon^\alpha$ in $B'(x, \epsilon)$, where C is a constant given by P2 in Theorem 2.3 so that u_k has no free boundary points in $B(x, \epsilon/2)$ for all large k . This proves (3.3).

Next we will show that for all large k

$$F(u_0) \subset \{x: d(F(u_k), x) < \epsilon\}. \quad (3.4)$$

If this was not true we could find a point $x \in F(u_0)$ and a subsequence of u_k such that $B'(x, \epsilon) \subset F(u_k)^c$ for every k in the subsequence. If the subsequence contains infinitely many u_k such that $u_k \equiv 0$ in $B(x, \epsilon)$ then also $u_0 \equiv 0$ due to uniform convergence. Otherwise, the sequence contains infinitely many u_k for which $B(x, \epsilon)$ is contained in the positive phase. In this case the non-degeneracy implies that in $B(x, \epsilon/2)$ we have $u_k > C\epsilon^\alpha$, with C independent of k . Again uniform convergence implies the same lower bound for u_0 , which contradicts our choice $x \in F(u_0)$.

To show the fourth claim we notice that $F(u_0)$ has zero n -dimensional Lebesgue measure by the Lebesgue differentiation Theorem and the positive density of the zero phase. Take an open set $V \supset F(u_0)$ with $m(V \cap G') < \epsilon$. For large k we have $F(u_k) \cup F(u_0) \subset V \cap G'$, so $\|\chi_{\{u_k > 0\}} - \chi_{\{u_0 > 0\}}\|_{L^1(G')} < \epsilon$.

Also the sequence is uniformly bounded in $H^{1,p}(G; |y|^\beta)$ by the Caccioppoli inequality. This implies by compactness [HKM06, 1.31 Theorem] the weak convergence of ∇u_k in $L^p(G; |y|^\beta)$. To obtain strong convergence, use Lemma 3.5 below. □

It remains to show that weak convergence implies strong convergence.

Lemma 3.5. *Any sequence of minimizers $\{u_k\}_{k=0}^\infty$ in $\Omega \subset \mathbb{R}^{n+1}$ with $u_k \rightarrow u_0$ uniformly and $\nabla u_k \rightarrow \nabla u_0$ weakly in $L^2_{\text{loc}}(\Omega, |y|^\beta)$ satisfies that $\nabla u_k \rightarrow \nabla u_0$ in $L^2_{\text{loc}}(\Omega, |y|^\beta)$.*

Proof. Let $\eta \in C_c^{0,1}(\Omega)$ be a non-negative function. We claim that for every $\varepsilon > 0$ there exists j_0 so that

$$\int |y|^\beta \eta |\nabla u - \nabla u_j|^2 \leq \varepsilon$$

for $j \geq j_0$.

First we isolate the main difficulty

$$\int |y|^\beta \eta |\nabla u_0 - \nabla u_j|^2 = \int |y|^\beta \eta (\nabla u_0 - \nabla u_j) \cdot \nabla u_0 - \int |y|^\beta \eta (\nabla u_0 - \nabla u_j) \cdot \nabla u_j.$$

By weak convergence,

$$\left| \int |y|^\beta \eta (\nabla u_0 - \nabla u_j) \cdot \nabla u_0 \right| \leq \varepsilon/4$$

for j big enough. Note that this is true even if the u_j are not minimizers. The bound on the second term, however, needs the minimization property.

We observe that

$$\int |y|^\beta \eta (\nabla u_0 - \nabla u_j) \cdot \nabla u_j = \underbrace{\int |y|^\beta (\nabla u_0 - \nabla u_j) \cdot \nabla (\eta u_j)}_{=:I} - \underbrace{\int |y|^\beta u_j (\nabla u_0 - \nabla u_j) \cdot \nabla \eta}_{=:II}. \quad (3.5)$$

To estimate I in (3.5), let λ_j be the measures corresponding to u_j from Lemma 3.1. By (3.2) we get that

$$\int |y|^\beta (\nabla u_0 - \nabla u_j) \cdot \nabla (\eta u_j) = \int \eta u_j d\lambda_0 - \int \eta u_j d\lambda_j.$$

Since λ_j is supported on $\{u_j = 0\}$ we have that

$$\int \eta u_j d\lambda_j = 0$$

for every j (including $j = 0$ as u_0 is also a minimizer to \mathcal{J} , see Corollary 3.4 in [CRS10b]).

To finish the estimate on I in (3.5) we observe that

$$\int \eta u_j d\lambda_0 = \int \eta (u_j - u_0) d\lambda_0 \leq \sup_{\text{supp } \eta} |u_j - u_0| \int \eta d\lambda_0.$$

By uniform convergence on compact subsets, for j big enough, $\sup_{\text{supp } \eta} |u_j - u_0| \leq \frac{\varepsilon}{4\|\eta\|_{L^1(\lambda_0)}}$.

We turn towards estimating II in (3.5):

$$\begin{aligned} |II| &= \left| \int |y|^\beta u_j (\nabla u_0 - \nabla u_j) \cdot \nabla \eta \right| \leq \left| \int |y|^\beta (\nabla u_0 - \nabla u_j) \cdot (u_0 \nabla \eta) \right| \\ &\quad + \sup_{\text{supp } \eta} |u_j - u_0| \|\nabla u_0 - \nabla u_j\|_{L^2(\Omega, |y|^\beta)} \|\nabla \eta\|_{L^2(\Omega, |y|^\beta)}. \end{aligned} \quad (3.6)$$

The first term goes to zero by weak convergence of ∇u_j to ∇u_0 . The second term satisfies

$$\sup_{\text{supp } \eta} |u_j - u_0| \|\nabla u_0 - \nabla u_j\|_{L^2(\text{supp } \eta, |y|^\beta)} \|\nabla \eta\|_{L^2(\Omega, |y|^\beta)} \leq \varepsilon/4$$

for j big enough, by uniform convergence and the uniform bound of $\|\nabla u_j\|_{L^2(\text{supp } \eta, |y|^\beta)}$ derived from the Caccioppoli inequality in Lemma 3.2 together with uniform convergence. \square

Lemma 3.4 implies that minimizers converge to minimizers (which was observed in Corollary 3.4 in [CRS10b]), but also implies the stronger fact that the energy is continuous under this convergence:

Corollary 3.6. *Let u_k be a sequence of minimizers in $\Omega \subset \mathbb{R}^{n+1}$ with $u_k \rightarrow u_0$ locally uniformly and $\sup_k \|u_k\|_{\mathbf{H}^\beta} < \infty$. Then u_0 is also a minimizer to \mathcal{J} in Ω and for any $B \subset\subset \Omega$ we have $\mathcal{J}(u_k, B) \rightarrow \mathcal{J}(u_0, B)$ after passing to a subsequence.*

4 Monotonicity formula and some immediate consequences

From [All12] we have the following monotonicity formula:

Theorem 4.1 (Monotonicity formula, see [All12, Theorem 4.3]). *Let $u \in \mathbf{H}^\beta(B_\delta(x_0))$ be a minimizer in $B_\delta(x_0)$ for the functional \mathcal{J} with $x_0 \in F(u)$. Then the function*

$$r \mapsto \Psi_r^u(x) := \Psi(r) = \frac{\mathcal{J}(u, B_r(x_0))}{r^n} - \frac{\alpha}{r^{n+1}} \int_{\partial B_r(x_0)} |y|^\beta u^2 d\mathcal{H}^n$$

is defined and nondecreasing in $(0, \delta)$, and for $0 < \rho < \sigma < \delta$, it satisfies

$$\Psi(\sigma) - \Psi(\rho) = \int_{B_\sigma(x_0) \setminus B_\rho(x_0)} |y|^\beta \frac{2|\alpha u(x) - (x - x_0) \cdot \nabla u(x)|^2}{|x_0 - x|^{n+2}} dx \geq 0.$$

As a consequence, the blow-up limits are cones, in the sense of the following corollary.

Corollary 4.2. *Let $u \in \mathbf{H}^\beta(B_\delta(x_0))$ be a minimizer in $B_\delta(x_0)$ with $x_0 = (x'_0, 0)$. Consider a decreasing sequence $0 < \rho_k \xrightarrow{k \rightarrow \infty} 0$ and the associated rescalings $u_k(x) := \frac{u(x_0 + \rho_k x)}{\rho_k^\alpha}$. Then the Allen-Weiss density*

$$\Psi_0^u(x_0) := \lim_{r \searrow 0} \Psi_r^u(x_0)$$

is well defined. Furthermore, for every bounded open set $D \subset \mathbb{R}^{n+1}$ and $k \geq k(D)$ this subsequence u_k is bounded in $H^{1,2}(D; |y|^\beta)$ and, passing to a subsequence u_{k_j} , converges (in the sense of Lemma 3.4) to u_0 which is a globally defined minimizer of \mathcal{J} that is homogeneous of degree α .

The proof is the same as in [Wei99, Theorem 2.8]

Remark 4.3 (Non-uniqueness of blow-ups). *We call the function u_0 appearing in Corollary 4.2 a blow-up of u at x_0 . A priori, the function u_0 may depend on the subsequence u_{k_j} . However, a simple scaling argument shows that for all radii $r \geq 0$ and all blow-ups u_0 to u at x_0 we have*

$$\Psi_r^{u_0}(0) \equiv \Psi_0^u(x_0).$$

4.1 Dimension reduction

We use the homogeneity of the blow-ups to obtain dimension estimates on the points in the free boundary for which there exists a non-flat blow-up. This process is known as “dimension reduction” and has been applied to a variety of situations (see [Wei99] for its application to the Bernoulli problem).

The first lemma shows that blow-up limits of blow-up limits have additional symmetry:

Lemma 4.4. *Let $u \in \mathbf{H}_{\text{loc}}^\beta(\mathbb{R}^{n+1})$ be an α -homogeneous minimizer of \mathcal{J} and let $x_0 \in F(u) \setminus \{0\}$. Then any blow-up limit u_0 at x_0 is invariant in the direction of x_0 , i.e., for every $x \in \mathbb{R}^{n+1}$ and every $\lambda \in \mathbb{R}$,*

$$u_0(x + \lambda x_0) = u_0(x).$$

Proof. Let $x \in \mathbb{R}^{n+1}$, and consider its decomposition $x = \tilde{x} + \lambda x_0$ with $\tilde{x} \in \langle x_0 \rangle^\perp$. We only need to check that

$$u_0(x) = u_0(\tilde{x}). \quad (4.1)$$




Figure 4.1: The distance $\text{dist}(P_1, P_3) = \mathcal{O}(\rho_k^2)$.

Consider a ball $B = B(0, r) \subset \mathbb{R}^{n+1}$ so that $\tilde{x}, x \in B$. Let $\{\rho_k\}$ be a sequence of radii converging to zero and such that $u_k(x) := \frac{u(x_0 + \rho_k x)}{\rho_k^\alpha}$ converges to u_0 uniformly on B_r . For k big enough, $\|u_k - u_0\|_{L^\infty(B_r)} < \varepsilon$. Then,

$$|u_0(x) - u_0(\tilde{x})| \leq 2\varepsilon + |u_k(x) - u_k(\tilde{x})|. \quad (4.2)$$

To control the last term above, we use the homogeneity of u . Writing $P_1 := x_0 + \rho_k \tilde{x}$ and $P_2 := x_0 + \rho_k x$ we have $\rho_k^\alpha u_k(\tilde{x}) = u(P_1)$ and $\rho_k^\alpha u_k(x) = u(P_2)$. Let P_3 be the intersection between the line through P_1 and x_0 and the line through the origin and P_2 (see Figure 4.1). By homogeneity of u

$$u(P_2) = u(P_3) \left(\frac{|P_2|}{|P_3|} \right)^\alpha = u(P_3) \left(1 \pm \frac{|P_2 - P_3|}{|P_3|} \right)^\alpha.$$

Thus,

$$\rho_k^\alpha |u_k(x) - u_k(\tilde{x})| \leq |u(P_1) - u(P_3)| (1 + \mathcal{O}(\rho_k))^\alpha \leq |u(P_1) - u(P_3)| + |u(P_3)| \mathcal{O}(\rho_k).$$

By Thales' Theorem, $|P_1 - P_3| = \frac{|P_1 - P_2||P_3 - x_0|}{|x_0|} = \mathcal{O}(\rho_k^2)$ and using the \dot{C}^α character of u and the fact that $u(x_0) = 0$, we get

$$\rho_k^\alpha |u_k(x) - u_k(\tilde{x})| \leq \|u\|_{\dot{C}^\alpha} (|P_1 - P_3|^\alpha + |P_3|^\alpha \mathcal{O}(\rho_k)) = \mathcal{O}(\rho_k^{2\alpha}) + \mathcal{O}(\rho_k),$$

and (4.1) follows by (4.2) since $\rho_k \rightarrow 0$. \square

We then recall that a minimizer with a translational symmetry is actually a minimizer without that symmetry in one dimension less. This is known as ‘‘cone splitting’’:

Lemma 4.5. *Let $u \in \mathbf{H}_{\text{loc}}^\beta(\mathbb{R}^{n+1})$ be an α -homogeneous minimizer of \mathcal{J} in \mathbb{R}^{n+1} which is invariant in the direction e_n . Then $\tilde{u}(x', y) := u(x', 0, y)$ is a minimizer of \mathcal{J} in one dimension less.*

Proof. The proof is a slight variation of [Wei99, Proof of Lemma 3.2]. \square

Next we provide a non-standard proof of Proposition 2.2, that is, to show that the trivial solution is a minimizer. We use *P5* in a sequence of conveniently chosen blow-ups and a dimension reduction argument, based on the following lemma. Note that the proposition could also be proven via a classical dimension reduction argument.

Proof of Proposition 2.2. Consider a non-zero minimizer u with non-empty free boundary (see [CRS10b, Proposition 3.2] for its existence), choose a free boundary point $x_0 \in F(u)$ and consider u_0 to be a blow-up weak limit at this point, which exists and is α -homogeneous by Lemma 4.2. Then u_0 is also a global minimizer by P5 and not null by the nondegeneracy condition.

Next we argue by induction: given $0 \leq j \leq n-2$ let u_j be an α -homogeneous global minimizer different from 0 such that it is invariant in a j -dimensional linear subspace $H_j \subset \mathbb{R}^n$, i.e., for every $v \in H_j$ and every $x' \in \mathbb{R}^n$,

$$u_j(x', y) = u(x' + v, y).$$

Consider a point $x_j \in F(u_j) \setminus (H_j \times \{0\})$ which exists as long as $j < n-1$ by the interior corkscrew condition and positive density, and let u_{j+1} be a blow-up limit at this point, which is again an α -homogeneous global minimizer. We claim that u_{j+1} is invariant in fact in the $(j+1)$ -dimensional subspace $H_j + \langle x'_j \rangle$.

Indeed u_{j+1} is invariant in $\langle x'_j \rangle$ by Lemma 4.4. On the other hand, since u_j is invariant in H_j , so are the functions in the blow-up sequence and, thus, u_{j+1} is invariant in H_j . Thus, for $v \in H_j$, $v_0 \in \langle x'_j \rangle$ and $x \in \mathbb{R}^{n+1}$ we get

$$u(x + v + x'_j) = u(x + v) = u(x),$$

and the claim follows.

Thus, after $n-1$ steps, we obtain u_{n-1} which is an α -homogeneous global minimizer invariant in an $(n-1)$ -dimensional space H_{n-1} , with non-empty free boundary. Thus,

$$u_{n-1}(x', 0) = C_{n,\alpha}(x'_n)_+^\alpha,$$

where the constant is given by P6. The proposition follows by Proposition A.1. \square

4.2 Upper semicontinuity

Next we show that Allen-Weiss' energy at a fixed radius is continuous both with respect to the minimizer and with respect to the point:

Lemma 4.6. *Let $u_j \in \mathbf{H}_{\text{loc}}^\beta(\Omega)$ be minimizers of \mathcal{J} in Ω and $u_j \rightarrow u_0$ in the sense of Lemma 3.4. Then, for $x_j \rightarrow x_0$ and $r < \text{dist}(x_0, \partial\Omega)$,*

$$\Psi_r^{u_j}(x_j) \xrightarrow{j \rightarrow \infty} \Psi_r^{u_0}(x_0).$$

Proof. Let $\varepsilon > 0$. We want to check that for j big enough,

$$|\Psi_r^{u_j}(x_j) - \Psi_r^{u_0}(x_0)| \leq \varepsilon.$$

We will consider the three terms of the energy separately. For the first term,

$$\int_{B_r(x_j)} |y|^\beta |\nabla u_j|^2 - \int_{B_r(x_0)} |y|^\beta |\nabla u_0|^2 \leq r^n \varepsilon / 3$$

follows from the L^2 convergence of the gradients. Indeed, if $\delta_j := |x_j - x_0| \leq \delta$ for j big enough and $B_{r+\delta} \subset \Omega$, then

$$\begin{aligned} \int_{B_r(x_j)} |y|^\beta |\nabla u_j|^2 - \int_{B_r(x_0)} |y|^\beta |\nabla u_0|^2 &\leq \int_{B_r(x_j)} |y|^\beta (|\nabla u_j|^2 - |\nabla u_0|^2) + \int_{B_r(x_j) \Delta B_r(x_0)} |y|^\beta |\nabla u_0|^2 \\ &\leq \int_{B_{r+\delta}(x_j)} |y|^\beta (|\nabla u_j|^2 - |\nabla u_0|^2) + \int_{(B_{r+\delta_j} \setminus B_{r-\delta_j})(x_0)} |y|^\beta |\nabla u_0|^2 \\ &\leq r^n \varepsilon / 3. \end{aligned}$$

For the measure, we estimate

$$\left| \int_{B_r(x_j)'} \chi_{\Omega_+(u_j)} dm - \int_{B_r(x_0)'} \chi_{\Omega_+(u_0)} dm \right| \leq r^n \varepsilon / 3$$

for j big enough as a consequence of $\chi_{\Omega_+(u_j)} \rightarrow \chi_{\Omega_+(u_0)}$ in L^1_{loc} as before. The fact that

$$\alpha \left| \int_{\partial B_r(x_j)} u_j^2 - \int_{\partial B_r(x_0)} u_0^2 \right| \leq r^{n+1} \varepsilon / 3$$

for j big enough is a straight consequence of the uniform convergence and the continuity of u_0 . \square

It is well known that the limit of a decreasing sequence of continuous functions is upper semi-continuous (see [Dal12, Theorem 1.8]). The monotonicity formula also implies the following result.

Lemma 4.7. *Let $u_j \in \mathbf{H}_{\text{loc}}^\beta(\Omega)$ be minimizers of \mathcal{J} in Ω and $u_j \xrightarrow{j \rightarrow \infty} u_0$ in the sense of Lemma 3.4, with $x_j \in F(u_j)$ for $j \in \mathbb{N}$. Then, if $x_j \rightarrow x_0$ and $r_j \rightarrow 0$,*

$$\limsup_j \Psi_0^{u_j}(x_j) \leq \limsup_j \Psi_{r_j}^{u_j}(x_j) \leq \Psi_0^{u_0}(x_0).$$

Proof. The first inequality comes from monotonicity.

To see that

$$\limsup_j \Psi_{r_j}^{u_j}(x_j) \leq \Psi_0^{u_0}(x_0),$$

it is enough to check that for every $r > 0$

$$\limsup_j \Psi_r^{u_j}(x_j) \leq \Psi_r^{u_0}(x_0),$$

or using monotonicity it suffices to show that for every $\varepsilon > 0$ and j big enough,

$$\Psi_r^{u_j}(x_j) - \Psi_r^{u_0}(x_0) \leq \varepsilon.$$

But this is true for j big enough because the left-hand side converges to 0 by the continuity of the energy from Lemma 4.6. \square

5 Measure-theoretic properties

5.1 Finite perimeter

We will show that $\Omega'_+(u)$ is a set of locally finite perimeter. Then $F_{\text{red}}(u)$ will coincide with the measure-theoretic reduced boundary by the ϵ -regularity theorem, see [AC81, Sections 4.6 and 4.7].

Definition 5.1. *For every $0 < \alpha < 1$ we can define*

$$k_\alpha^* := \inf \left\{ k \in \mathbb{N} : \exists \text{ an } \alpha\text{-homogeneous minimizer } u \in \mathbf{H}_{\text{loc}}^\beta(\mathbb{R}^{k+1}) \text{ such that } \Sigma(u) = \{0\} \right\}.$$

Note that, to the best of our knowledge, there is no result showing that k_α^* needs to be finite.

Lemma 5.2. *Let u be an α -homogeneous minimizer of \mathcal{J} in \mathbb{R}^{n+1} with $n < k_\alpha^*$. Then u is a rotation of the trivial solution.*

See [Wei99, Section 3] for the proof.

From the positive density properties, we know that $k_\alpha^* \geq 2$. From the homogeneity of the blow-ups we find out that the free boundary in \mathbb{R}^{1+1} is in fact a collection of isolated points. Later in Theorem 6.1 we will show that in fact $k_\alpha^* \geq 3$.

Lemma 5.3 (Isolated singularities). *Let $u \in \mathbf{H}_{\text{loc}}^\beta(\Omega)$ for $\Omega \subset \mathbb{R}^{1+1}$ be a minimizer of \mathcal{J} in Ω . Then $F(u)$ has no accumulation points in Ω .*

Proof. Arguing by contradiction, we assume that $F(u)$ has an interior accumulation point which, without loss of generality, we assume to be the origin.

Let $(x_k, 0)$ be a sequence of singular points converging to 0 with $x_k > 0$. Consider the blow-up rescaling $u_k(x) := \frac{u(x_k x)}{x_k^\alpha}$. Note that $u_k(0, 0) = u_k(1, 0) = 0$. Moreover, by the interior corkscrew condition, there exist $z_k \in (1/2, 3/2)$ such that $u_k|_{B_c(z_k, 0)} > 0$, so $u(z_k, 0) \gtrsim C$ by the non-degeneracy condition.

Choosing a subsequence, we may assume that $z_k \rightarrow z_0 \geq 1/2$, and $u_k \rightarrow u_0$ in the sense of Lemma 3.4. In particular u_0 is homogeneous by Corollary 4.2, reaching a contradiction with the fact that $u_0(1, 0) = 0$ and $u_0(z_0, 0) \gtrsim C$. \square

We will prove the local finiteness of the perimeter of the free boundary adapting a proof of De Silva and Savin in [DS15]. Our proof is essentially the same, but we repeat it for the sake of completeness.

As in [DS15] we say that a set $A \subset \mathbb{R}^n$ satisfies the property (P_t) if the following holds: for every $x \in A$ there exists an $r_x > 0$ such that for every $0 < r < r_x$, every subset S of $B(x, r) \cap A$ can be covered with a finite number of balls $B(x_i, r_i)$ with $x_i \in S$ such that

$$\sum_i r_i \leq r^t/2. \quad (5.1)$$

Lemma 5.4. *If $\mathcal{H}^t(\Sigma(U)) = 0$ for every minimal cone U in \mathbb{R}^{n+1} then $\mathcal{H}^t(\Sigma(u)) = 0$ for every minimizer u of \mathcal{J} defined on $\Omega \subset \mathbb{R}^{n+1}$*

Proof. We first show that $\Sigma(u)$ satisfies the property (P_t) . If (P_t) does not hold we find a point $y \in \Sigma(u)$ for (P_t) is violated for a sequence $r_k \rightarrow 0$. We consider the blow-up sequence

$$u_{r_k}(x) = r_k^{-\alpha} u(y + r_k x). \quad (5.2)$$

By Corollary 3.6 we may assume, by taking a subsequence, that u_{r_k} converges to a minimal cone U . By our assumptions we may cover $\Sigma(U) \cap B(0, 1)$ with a finite collection of balls $\{B(x_i, \frac{\rho_i}{10})\}_{i=1}^k$ with

$$\sum_i \rho_i^t \leq \frac{1}{2}.$$

By Lemma 3.4 we know that free boundaries converge in Hausdorff sense and thus the set $F(u_{r_k}) \cap B(0, 1) \setminus \cup_i B(x_i, \rho_i/5)$ is flat for all large k . From Theorem 2.4 we infer that all singularities must be covered by the same balls, that is, for all $k \geq k_0$

$$\Sigma(u_{r_k}) \cap B(0, 1) \subset \bigcup_i B(x_i, \rho_i/5). \quad (5.3)$$

After rescaling we see that u satisfies the condition for property (P_t) in the ball $B(y, r_k)$, which is a contradiction. Therefore the property (P_t) holds as claimed.

Consider the set $D_k := \{y \in \Sigma(u) : r_y \geq 1/k\}$. Fix a point $y_0 \in D_k$. By property (P_t) applied to $r_0 = 1/k$ we find a finite cover of $D_k \cap B(y_0, r_0)$ with balls $B(y_i, r_i)$, $y_i \in D_k$, satisfying

$$\sum_i r_i^t \leq r_0^t/2.$$

Similarly, for each ball $B(y_i, r_i)$ in the cover we use the property (P_t) to find a finite number of balls $B(y_{ij}, r_{ij})$, $y_{ij} \in D_k$, which cover $D_k \cap B(y_i, r_i)$ and satisfy

$$\sum_j r_{ij}^t \leq r_i^t/2,$$

and thus $\sum_{i,j} r_{ij} \leq r_0^t/4$. By repeating the argument N times we obtain a cover of $D_k \cap B(y_0, r_0)$ by balls $B(z_l, r_l)$ which satisfies

$$\sum_l r_l^t \leq 2^{-N} r_0^t.$$

This implies that $\mathcal{H}^t(B(y_0, r_0) \cap D_k) = 0$ and thus $\mathcal{H}^t(D_k) = 0$. By countable additivity we obtain the claim. \square

Lemma 5.5. *If $\mathcal{H}^t(\Sigma(U)) = 0$ for some $t > 0$ and for every minimal cone in \mathbb{R}^{n+1} , we then have that $\mathcal{H}^{t+1}(\Sigma(V)) = 0$ for every minimal cone V in $\mathbb{R}^{(n+1)+1}$.*

Proof. Without loss of generality we may assume $\Sigma(V) \neq \{0\}$. Let $x \in \Sigma(V) \setminus \{0\}$. By Corollary 3.6 the blow-ups at any point of $\Sigma(V) \cap \partial B$ converge to a minimal cone in dimension $(n+1)+1$ up to a subsequence. Let V_x be a blow-up at x . By Lemma 4.5 V_x is a minimal cone which is invariant in at least one direction. By Lemma 4.5, using our assumption this implies that $\mathcal{H}^{t+1}(\Sigma(V_x)) = 0$, and thus the singular set of every possible blow-up cone of any minimizer V has zero \mathcal{H}^{t+1} -measure.

Arguing as in Lemma 5.4 we obtain $\mathcal{H}^{t+1}(\Sigma(V)) = 0$. \square

Combining Lemmas 5.3, 5.4 and 5.5 we obtain the following corollary. Notice that we will be able to replace $n-1$ by $n-2$ by Theorem 6.1.

Corollary 5.6. *Every minimizer satisfies*

$$\mathcal{H}^{n-1}(\Sigma(u)) = 0.$$

Lemma 5.7. *Let $u \in \mathbf{H}^\beta(2B)$ be a minimizer of \mathcal{J} in $2B$ with $\|u\|_{\dot{C}^\alpha(2B)} < E_0$. Then there exists a constant C depending on n, α and E_0 and a finite collection of balls $\{B(X_i, r_i)\}$ s.t.*

$$\mathcal{H}^{n-1} \left((F(u) \cap B) \setminus \bigcap_{i=1}^m B(X_i, r_i) \right) \leq C \quad (5.4)$$

and

$$\sum_{i=1}^m r_i^{n-1} \leq \frac{1}{2}. \quad (5.5)$$

Proof. Proof is by contradiction. For $k \in \mathbb{N}$ assume we have $\|u_k\|_{\dot{C}^\alpha(2B)} < E_0$ and the left-hand side of (5.4) is bounded below by $k > 0$ for every collection of balls satisfying (5.5). By Lemma 3.2 we know the sequence u_k is bounded in $\mathbf{H}^\beta(B)$. Taking a subsequence we may assume that u_k converges locally uniformly to a minimizer u (see Corollary 3.6).

By Corollary 5.6 the set of singularities $\Sigma(u)$ has \mathcal{H}^{n-1} -measure zero and thus they can be covered with finitely many balls B_i satisfying (5.5).

Since $F(u) \setminus \Sigma(u)$ is a $C^{1,\gamma}$ -surface by Theorem 2.4, using the Hausdorff convergence of the free boundaries we apply again Theorem 2.4 to see that $F(u_k) \cap B_1 \setminus \bigcup_{i=1}^M B_i$ are also $C^{1,\gamma}$ -surfaces converging to $F(u) \cap B_1 \setminus \bigcup_{i=1}^M B_i$ uniformly in the C^1 -norm. This is a contradiction with the assumption that the Hausdorff measure blows up as k goes to ∞ . \square

The fact that the free boundary has finite perimeter follows now from the same iteration argument as [DS15, Lemma 5.10].

Lemma 5.8. *Let u be as in Lemma 5.7. Then for some constant C depending only on E_0 ,*

$$\mathcal{H}^{n-1}(F(u) \cap B) \leq C. \quad (5.6)$$

Proof. By Lemma 5.7 we find a finite collection of balls B_{r_i} such that

$$F(u) \cap B \subset \Gamma \cup \bigcup B_{r_i}, \quad (5.7)$$

with $\mathcal{H}^{n-1}(\Gamma) \leq C$ and $\sum r_i^{n-1} \leq \frac{1}{2}$.

Applying Lemma 5.7 again for each ball B_{r_i} we have

$$F(u) \cap B_{r_i} \subset \Gamma_i \cup \bigcup B_{r_{ij}}, \quad (5.8)$$

with $\mathcal{H}^{n-1}(\Gamma_i) \leq C r_i^{n-1}$ and $\sum r_{ij}^{n-1} \leq \frac{1}{2} r_i^{n-1}$. Moreover, we have

$$\mathcal{H}^{n-1}\left((F(u) \cap B_1) \cap \bigcup_{i,j} B_{r_{ij}}\right) \leq \mathcal{H}^{n-1}(\Gamma) + \sum \mathcal{H}^{n-1}(\Gamma_i) \leq C \left(1 + \sum_{i,j} r_{ij}^{n-1}\right) \leq C \left(1 + \frac{1}{2}\right).$$

Continuing inductively, after k steps we have that

$$F(u) \cap B_1 \subset \Gamma' \cup \bigcup_{q=1}^N B_{r_q}, \quad (5.9)$$

with

$$\mathcal{H}^{n-1}(\Gamma') \leq C \left(\sum_{i=0}^k 2^{-i}\right) \leq 2C,$$

and $\sum r_q^{n-1} \leq 2^{-k}$. This gives the claim. \square

Finally the fact that $\{u > 0\} \cup \Omega$ has locally finite perimeter in Ω follows from the previous lemma and well-known results of Federer, see for example [AFP00, Prop. 3.62] or [Fed69, 4.5.11].

5.2 Energy gap

Next we will check that the Allen-Weiss density can also be used to identify singular points. First let us state a useful identity for minimizers (which is also valid in the context of variational solutions in the sense of [Wei98]).

Lemma 5.9 (See [All12, Proposition 3.4]). *Let $u \in \mathbf{H}_{\text{loc}}^\beta(\Omega)$ be a minimizer to (1.1) in Ω . For every $B \subset\subset \Omega$ we have*

$$\int_B |y|^\beta |\nabla u|^2 = \int_{\partial B} |y|^\beta u \nabla u \cdot \nu d\mathcal{H}^n. \quad (5.10)$$

Let u be a minimizer and $x_0 \in F(u)$. If we consider a blow-up u_0 at x_0 , then

$$\Psi_0^u(x_0) = \Psi_1^{u_0}(0) = \int_{B_1} |y|^\beta |\nabla u_0|^2 + m(\{u_0 > 0\} \cap \mathbb{R}^n \cap B_1) - \alpha \int_{\partial B_1} |y|^\beta u_0^2 d\mathcal{H}^n.$$

By Lemma 5.9 we get

$$\Psi_1^{u_0}(0) = \int_{\partial B_1(x_0)} |y|^\beta u_0 \nabla u_0 \cdot \nu d\mathcal{H}^n + m(\{u_0 > 0\} \cap \mathbb{R}^n \cap B_1) - \alpha \int_{\partial B_1} |y|^\beta u_0^2 d\mathcal{H}^n.$$

Since $\nabla u_0(x) \cdot \nu(x) = \frac{\alpha}{|x|} u_0(x)$ almost everywhere on the sphere, the first and the third terms cancel out and we obtain

$$\Psi_1^{u_0}(0) = m(\{u_0 > 0\} \cap B'_1).$$

Thus, the density Ψ_0^u at a free boundary point is given by the area of the positive phase of any blow-up at the same point.

We write $\omega_n := m(B'_1)$ for the volume of the n -dimensional ball.

Proposition 5.10. *Every homogeneous minimizer $u \in \mathbf{H}_{\text{loc}}^\beta(\mathbb{R}^{n+1})$ has density*

$$\Psi_1^u(0) = m(\{u > 0\} \cap B'_1) \geq \frac{\omega_n}{2},$$

and equality is only attained when u is the trivial minimizer.

Proof. Let u be a minimizer such that $\Psi_1^u(0) \leq \frac{\omega_n}{2}$.

Let $x_1 \in F_{\text{red}}(u)$. Being a regular point, $\Psi_0^u(x_1) = \frac{\omega_n}{2}$. On the other hand, by the homogeneity and the continuity in Lemma 4.6,

$$\lim_{r \rightarrow \infty} \Psi_r^u(x_1) = \lim_{r \rightarrow \infty} \Psi_1^u(x_1/r) = \Psi_1^u(0) \leq \frac{\omega_n}{2}.$$

Combining both assertions with the monotonicity of Ψ we get that $\Psi_r^u(x_1) \equiv \frac{\omega_n}{2}$. But using the second formula in Theorem 4.1, one can see that this is true only whenever Ψ is α -homogeneous with respect to x_1 . Thus, u is 1-symmetric and invariant in the direction of $\langle x_1 \rangle$.

By Corollary 5.6 $F_{\text{red}}(u)$ has full \mathcal{H}^{n-1} measure on $F(u)$. Thus, we can find $x_1, \dots, x_{n-1} \in F_{\text{red}}(u)$ linearly independent. By the previous discussion u is invariant on an $(n-1)$ -dimensional affine manifold and, thus, it is the trivial solution. \square

Corollary 5.11 (Energy Gap). *There exists $\bar{\epsilon} > 0$ depending only on n and α such that every minimizer $u \in \mathbf{H}_{\text{loc}}^\beta(\Omega)$ and every singular point $x_0 \in \Sigma(u)$ satisfy*

$$\Psi_1^u(x_0) - \frac{\omega_n}{2} \geq \bar{\epsilon}.$$

Proof. Assume the conclusion to be false. Then there exist u_j minimizers in B_1 with

$$\Psi_1^{u_j}(0) \leq \frac{\omega_n}{2} + 1/j.$$

Passing to a subsequence, $u_j \rightarrow u_0$ as in Lemma 3.4. Using Lemma 4.7 we get that

$$\Psi_1^{u_0}(0) = \lim_j \Psi_1^{u_j}(0) \leq \frac{\omega_n}{2}.$$

But then u_0 is the trivial cone by Proposition 5.10. Since $F(u_j) \rightarrow F(u)$ in the Hausdorff distance, using ϵ -regularity (see Theorem 2.4) we get that u_j is the trivial cone for j big enough. \square

The value $\bar{\epsilon}$ above depends on the constants and on $\|u\|_{\dot{C}^\alpha}$ in a neighborhood of x_0 . In Section 7 we will show that $\bar{\epsilon}$ does not depend on u at all.

6 Full regularity in \mathbb{R}^{2+1}

In the case of $n = 2$, we prove full regularity of the free boundary for minimizers of our functional. Note that this result does not depend on the previous sections except that we use dimension reduction and blow-ups to deduce regularity of the free boundary.

Theorem 6.1. *Let $n = 2$. Then there is no singular minimal cone. In particular, the free boundary $F(u)$ of every minimizer u is $C^{1,\alpha}$ everywhere.*

Proof. We follow closely the arguments in [DS15, Theorem 5.5], building on [SV13]. The case $\beta = 0$ has been considered in [DS15]. The idea is to construct a competitor by a perturbation argument. We note at this point that the argument is two dimensional in nature and does not generalize to higher dimensions. Recall the functional under consideration:

$$\mathcal{J}(u, \Omega) = \int_{\Omega} |y|^{\beta} |\nabla u|^2 + m(\{u > 0\} \cap \mathbb{R}^n \cap \Omega).$$

Let V be a non trivial minimal cone. Define, as in [DS15], the Lipschitz continuous function

$$\psi_R(t) = \begin{cases} 1, & 0 \leq t \leq R, \\ 2 - \frac{\ln(t)}{\ln(R)}, & R \leq t \leq R^2, \\ 0, & t \geq R^2 \end{cases} \quad (6.1)$$

Define now the bi-Lipschitz change of coordinates

$$Z(x', y) = (x', y) + \psi_R(|(x', y)|)e_1$$

and set $V_R^+(Z) = V(x', y)$. Clearly, one has

$$D_{(x', y)} Z = \text{Id} + A$$

where $\|A\| \leq |\psi'_R(|(x', y)|)| \ll 1$. Defining now V_R^- exactly as V_R^+ changing ψ_R into $-\psi_R$, the very same computation as in [DS15] gives

$$\mathcal{J}(V_R^+, B_{R^2}) + \mathcal{J}(V_R^-, B_{R^2}) \leq 2\mathcal{J}(V, B_{R^2}) + \int_{B_{R^2}} |y|^{\beta} |\nabla V|^2 \|A\|^2.$$

Now, we have

$$\int_{B_{R^2}} |y|^{\beta} |\nabla V|^2 \|A\|^2 = \int_R^{R^2} \int_{\partial B_r} |y|^{\beta} |\nabla V|^2 \|A\|^2 d\mathcal{H}^n dr.$$

Now since V is homogeneous of degree α by assumption, the function $g(x, y) = |y|^{\beta} |\nabla V|^2$ is homogeneous of degree $\beta + 2\alpha - 2 = -1$. Therefore by a trivial change of variables on the sphere of radius r and using the fact that $n = 2$, we get the very same estimate

$$\int_{B_{R^2}} |y|^{\beta} |\nabla V|^2 \|A\|^2 \leq \frac{C}{\ln(R)} \xrightarrow{R \rightarrow \infty} 0.$$

The rest of the proof follows verbatim [DS15], page 1318 since this is only based on energy considerations and we refer the reader to it. \square

7 Uniform bounds around the free boundary

The optimal regularity bound and the non-degeneracy described in Theorem 2.3 were obtained in [CRS10b] with bounds that depend on the seminorm $\|u\|_{\dot{H}^\beta(B_1)}$. As a consequence, this dependence propagates to many of our estimates above. In this chapter we use the semi-norm dependent estimates (e.g. Lemma 5.8) to prove semi-norm *independent* non-degeneracy estimates. Re-running the arguments above yields the semi-norm independent results presented in our main Theorem 1.1.

The question of semi-norm independence may seem purely technical; however, independence allows the compactness arguments of the next section to work without additional assumptions on the minimizers involved.

7.1 Uniform non-degeneracy

We will begin by showing uniform non-degeneracy from scratch to deduce uniform Hölder character from this fact, reversing the usual arguments in the literature.

The following lemma was shown in [All12, Corollary 4.2] in a more general setting. Here we give a more basic approach based on [AC81, Lemma 3.4]. The main difference is that where Alt and Caffarelli could use the energy to directly control the H^1 norm of the minimizer, in our case we need to find an alternative because the measure term of the functional is computed on the thin phase (as opposed to the H^1 norm which is computed on the whole space). To bypass this difficulty we will use Allen's monotonicity formula.

The drawback of our approach is that we need the ball to be centered on the free boundary, while in the original lemma, Alt and Caffarelli could center the ball in the zero phase, allowing for a slightly better result.

Lemma 7.1. *Let u be a minimizer in B_r with $0 \in F(u)$. Then $\sup_{\partial B_r} u \geq Cr^\alpha$ with C depending only on n and α .*

Proof. By rescaling we can assume that $r = 1$.

Let $\mathcal{L}u := -\nabla \cdot (|y|^\beta \nabla u)$, consider $\Gamma(x) = \frac{1}{|x|^{n-2\alpha}}$ which is a solution of $\mathcal{L}\Gamma = 0$ away from the origin (or $\Gamma(x) = \log|x|$ if $n = 1$ and $\alpha = 1/2$), and let

$$\tilde{v}(x) := \ell \frac{\max\{1 - \Gamma(2x), 0\}}{1 - \Gamma(2)},$$

where

$$\ell := \sup_{\partial B_1} u.$$

It follows that $u \leq \tilde{v}$ on ∂B_1 and thus

$$\mathcal{J}(u, B_1) \leq \mathcal{J}(\min\{u, \tilde{v}\}, B_1),$$

and observing that $\tilde{v} = 0$ on $B_{1/2}$ and $\tilde{v} > 0$ on the annulus $A := B_1 \setminus B_{1/2}$, we get

$$\begin{aligned} \int_{B_{\frac{1}{2}}} |y|^\beta |\nabla u|^2 + m(B'_{\frac{1}{2},+}(u)) &\leq \int_A |y|^\beta (|\nabla(\min\{u, \tilde{v}\})|^2 - |\nabla u|^2) + m(A'_+(\min\{u, \tilde{v}\})) - m(A'_+(u)) \\ &\leq -2 \int_A |y|^\beta \nabla \max\{u - \tilde{v}, 0\} \cdot \nabla \tilde{v}. \end{aligned}$$

By Green's theorem, writing $d\sigma = |y|^\beta d\mathcal{H}^n$ we get

$$\int_{B_{\frac{1}{2}}} |y|^\beta |\nabla u|^2 + m(B'_{\frac{1}{2},+}(u)) \leq -2 \int_{\partial B_{\frac{1}{2}}} u \partial_\nu \tilde{v} d\sigma = C_{n,\alpha} \ell \int_{\partial B_{\frac{1}{2}}} u d\sigma, \quad (7.1)$$

with $C_{n,\alpha} > 0$.

Using the monotonicity formula and Proposition 5.10, we get that $\psi^u(r) \geq \psi^u(0) \geq \frac{\omega(B_1)}{2}$ and, therefore

$$\frac{\alpha}{r} \int_{\partial B_r} u^2 d\sigma + \frac{\omega(B_1)r^d}{2} \leq \mathcal{J}_r(u), \quad (7.2)$$

so using Hölder's inequality and the AM-GM inequality we obtain

$$\int_{\partial B_{\frac{1}{2}}} u d\sigma \leq \left(\int_{\partial B_{\frac{1}{2}}} u^2 d\sigma \right)^{\frac{1}{2}} C_{n,\alpha}^{\frac{1}{2}} \leq \frac{1}{2} \int_{\partial B_{\frac{1}{2}}} u^2 d\sigma + \frac{1}{2} C_{n,\alpha} \leq C_{n,\alpha} \mathcal{J}_{\frac{1}{2}}(u). \quad (7.3)$$

Combining (7.1), (7.2) and (7.3) we obtain

$$0 < \mathcal{J}_{\frac{1}{2}}(u) \leq C_{n,\alpha} \ell \mathcal{J}_{\frac{1}{2}}(u),$$

and therefore $\ell \geq C_{n,\alpha}^{-1}$. □

To show averaged non-degeneracy we need a mean value principle which is well-known, but we include its proof for the sake of completeness.

Lemma 7.2 (Mean value principle). *Let $u \in H^1(\beta, \Omega)$ be a weak solution to $\mathcal{L}u := \nabla \cdot (|y|^\beta \nabla u) = 0$ in Ω , and let $x_0 \in \mathbb{R}^n \times \{0\}$ with $B_r(x_0) \subset \Omega$. Then*

$$u(x_0) = \int_{B_r} u d\omega$$

where the mean is taken with respect to the measure $d\omega := |y|^\beta dx$.

Proof. Changing variables, we have that

$$A(\rho) := \frac{1}{\rho^{\beta+n+1}} \int_{B_\rho(x_0)} |y|^\beta u(x) dx = \int_{B_1} |y|^\beta u(\rho x + x_0) dx.$$

On the other hand, set

$$\begin{aligned} \tilde{A}(\rho) &:= \int_{B_1} |y|^\beta \nabla u(\rho x + x_0) \cdot x dx \\ &= \int_{B_\rho(x_0)} \left(\frac{|y|}{\rho} \right)^\beta \frac{\nabla u(x) \cdot (x - x_0)}{\rho} \frac{dx}{\rho^{n+1}} = \frac{1}{2\rho^{\beta+n+2}} \int_{B_\rho(x_0)} |y|^\beta \nabla u(x) \cdot \nabla |x - x_0|^2 dx. \end{aligned}$$

Since u is a weak solution to $\nabla \cdot (|y|^\beta \nabla u) = 0$ in Ω , we can apply Green's formula twice to obtain

$$\tilde{A}(\rho) = \frac{1}{2\rho^{\beta+n+2}} \int_{\partial B_\rho(x_0)} |x - x_0|^2 |y|^\beta \nabla u(x) \cdot \nu dx = \frac{1}{2\rho^{\beta+n}} \int_{\partial B_\rho(x_0)} |y|^\beta \nabla u(x) \cdot \nu dx = 0.$$

Since u is absolutely continuous on lines (see [EG15, Theorem 4.21]), for almost every x we have $\int_\rho^r \nabla u(tx + x_0) \cdot x dt = u(rx + x_0) - u(\rho x + x_0)$. Applying Fubini's Theorem we get

$$\int_\rho^r \tilde{A}(t) dt = \int_{B_1} |y|^\beta \int_\rho^r \nabla u(tx + x_0) \cdot x dt dx = \int_{B_1} |y|^\beta (u(rx + x_0) - u(\rho x + x_0)) dx = A(r) - A(\rho).$$

So $A(r) - A(\rho) = 0$ for all $\rho < r$.

On the other hand, taking the mean with respect to the measure $d\omega := |y|^\beta dx$ and using the continuity of u (see [FKS82, Theorem 2.3.12]) we obtain

$$\left| u(x_0) - \frac{1}{\omega(B_1)} \lim_{\rho \rightarrow 0} A(\rho) \right| = \lim_{\rho \rightarrow 0} \frac{1}{\omega(B_\rho(x_0))} \left| \int_{B_\rho(x_0)} (u(x_0) - u(x)) d\omega(x) \right| \leq \lim_{\rho \rightarrow 0} o_{\rho \rightarrow 0}(1) = 0.$$

□

Corollary 7.3. *Let u be a minimizer in B_r with $0 \in F(u)$ and let $d\sigma = |y|^\beta d\mathcal{H}^n$. Then $\int_{\partial B_r} u d\sigma \geq Cr^\alpha$ with C depending only on n and α .*

Proof. Let v be the \mathcal{L} -harmonic replacement of u in B_r , that is, the solution to

$$\begin{cases} \mathcal{L}v = 0 & \text{in } B_r, \\ v \equiv u & \text{on } \partial B_r, \end{cases} \quad (7.4)$$

see [HKM06, Theorem 3.17]. After differentiating with respect to the radius, by the mean value principle we get that $v(0) = \int_{\partial B_r} u d\sigma$. By the comparison principle and the Harnack inequality we get that

$$Cr^\alpha \leq \sup_{B_{r/2}} u \leq \sup_{B_{r/2}} v \leq C \int_{\partial B_r} u d\sigma. \quad (7.5)$$

□

7.2 Behavior of the distributional fractional Laplacian

Next we use an idea of [AC81] and investigate the behavior of the distributional α -Laplacian of the minimizer introduced in Section 3. As mentioned in the introduction, in [AC81] this investigation immediately yields that the positivity set is a set of locally finite perimeter, and more precisely, that it is Ahlfors regular of the correct dimension. However, the nonlocal nature of this problem indicates that the distributional fractional Laplacian may not be supported on the free boundary and thus we cannot expect to immediately gain such strong geometric information.

First we can bound the growth of the fractional Laplacian measure around a free boundary point. Note that this growth is the natural counterpart to the upper Ahlfors regularity in the case of Alt-Caffarelli minimizers.

Theorem 7.4. *Let $u \in \mathbf{H}^\beta(B_{2r}(x_0))$ be a minimizer of \mathcal{J} in $B_{2r}(x_0)$, and let $x_0 \in F(u)$. Then, we have*

$$\lambda(B_r(x_0)) \leq Cr^{n-\alpha}.$$

In particular $\lambda(F(u)) = 0$.

A glance at (2.1) will convince the reader that these estimates are sharp, for they cannot be improved even in the case of the trivial solution.

Proof. Without loss of generality we may assume that $x_0 = 0$. Let $\mathcal{L}u := -\nabla \cdot (|y|^\beta \nabla u)$ and let v be the \mathcal{L} -harmonic replacement of u in B_{2r} , see (7.4). Write $d\sigma = |y|^\beta d\mathcal{H}^n$ and $M := \int_{\partial B_{2r}} u d\sigma$. By Harnack's inequality (see [CRS10b], for instance) and the mean value principle in Lemma 7.2,

$$\inf_{B_r} v \geq Cv(0) = CM.$$

We have that

$$\lambda(B_r) = \int_{B_r} d\lambda \leq \frac{1}{CM} \int_{B_r} v d\lambda.$$

Since $u \equiv 0$ in the support of λ and u is \mathcal{L} -subharmonic (see [AC81, Lemma 2.2]) we get

$$\int_{B_r} v d\lambda = \int_{B_r} (v - u) d\lambda \leq \int_{B_{2r}} (v - u) d\lambda.$$

By the properties of the measure λ , we obtain

$$\int_{B_{2r}} (v - u) d\lambda = - \int_{B_{2r}} |y|^\beta \nabla(v - u) \cdot \nabla u = \int_{B_{2r}} |y|^\beta (|\nabla u|^2 - |\nabla v|^2),$$

and using the definition of the functional and the fact that u is a minimizer, we get

$$\int_{B_{2r}} |y|^\beta (|\nabla u|^2 - |\nabla v|^2) = \mathcal{J}(u, B_{2r}) - m(B_{2r}^+(u)) - \mathcal{J}(v, B_{2r}) + m(B_{2r}') \leq Cr^n.$$

All together, we have that

$$\lambda(B_r) \leq \frac{1}{CM} Cr^n,$$

and, since uniform non-degeneracy (see Corollary 7.3) implies that $M \geq Cr^\alpha$ we can conclude the proof of the first statement.

To show the second one, note that since the free boundary has locally finite $(n-1)$ -dimensional Hausdorff measure, given a set $E \subset F(u)$ and $k \in \mathbb{N}$ we can find a collection of balls $I_k = \{B_i^k\}_i$ such that

$$E \subset \bigcup_{B \in I_k} B, \quad \sup_{B \in I_k} r(B) \leq 1/k \quad \text{and} \quad \sum_{B \in I_k} r(B)^{n-1} \leq 2\mathcal{H}^{n-1}(E).$$

Thus,

$$\lambda(E) \leq \sum_{B \in I_k} \lambda(B) \lesssim \sum_{B \in I_k} r(B)^{n-\alpha} \leq \sup_{B \in I_k} r(B)^{1-\alpha} \sum_{B \in I_k} r(B)^{n-1} \xrightarrow{k \rightarrow \infty} 0.$$

□

Next we study the measure away from the free boundary. We should emphasize here that even though the estimates in Lemma 7.5 and Theorem 7.6 depend on E_0 , they will be used to remove the dependence of our other estimates on E_0 . More precisely, Theorem 7.6 will play a role in establishing the continuity of the Green function in Lemma 7.9. This qualitative fact is used to prove the quantitative uniform Hölder character in Theorem 7.8.

After proving Theorem 7.8, we may drop the hypothesis $\|u\|_{\mathbf{H}^\beta(B_2)} \leq E_0$ from both Lemma 7.5 and Theorem 7.6.

Lemma 7.5. *If $u \in \mathbf{H}_{\text{loc}}^\beta(B_2)$ is a minimizer of \mathcal{J} in the ball B_2 with $\|u\|_{\mathbf{H}^\beta(B_2)} \leq E_0$ and $0 \in F(u)$, then for every $x_0 = (x', 0) \in B_{1,0}(u)$ we get*

$$\lim_{y \rightarrow 0} |y|^\beta |u_y(x', y)| \approx C \text{dist}(x_0, F(u))^{-\alpha}.$$

Moreover, for every ball B centered at $\mathbb{R}^n \times \{0\}$ with $B' \subset\subset B_{1,0}(u)$, we have that

$$|y|^\beta |u_y(x', y)| \leq C \text{dist}(x_0, F(u))^{-\alpha},$$

for $|y| < C_B \text{dist}(x, F(u))$, where the constant C_B may depend on B .

Proof. Let u be a minimizer, and let $B := B_r(x_0)$ with $B' \subset B_{1,0}(u)$. By [Sil12, Lemma 2.2], we can write $u(x', y) = |y|^{1-\beta}g(x') + \mathcal{O}(y^2)$, where g is a $C^{1+\beta}(\frac{1}{2}B')$ function, with a uniform control on the error term in terms of $\|u\|_{L^2(B, |y|^\beta)}$. In particular, $\lim_{y \rightarrow 0} |y|^{\beta-1}u(x', y) = g(x')$.

Let us define

$$\tilde{u}(x', y) := \begin{cases} u(x', y) & \text{if } y \geq 0 \\ -u(x', -y) & \text{if } y < 0. \end{cases} \quad (7.6)$$

It is clear that $\mathcal{L}\tilde{u} \equiv 0$ in B . According to [Vit18, Lemma 3.26, Corollary 3.29] $v(x', y) = |y|^\beta y^{-1} \tilde{u}(x', y)$ is an even $C^\infty(\frac{1}{2}B)$ function in $\mathbf{H}^{2-\beta}(B)$ (note that $1 < 2 - \beta < 3$ is out of the usual range of β) and satisfying $\nabla \cdot (|y|^{2-\beta} \nabla v) = 0$. The mean value principle (see Lemma 7.2) applies also to this case, so

$$g(x'_0) = v(x_0) = \frac{1}{\int_{\frac{1}{2}B} |y|^{2-\beta}} \int_{\frac{1}{2}B} |y|^{2-\beta} v(x) = C \frac{1}{r^{2-\beta+n+1}} \int_{\frac{1}{2}B} |y| u(x),$$

and using *P1-P3*, if $r = \text{dist}(x_0, F(u))$ we get

$$g(x'_0) = v(x_0) \approx Cr^{\beta-2+1+\alpha} = Cr^{-\alpha}.$$

On the other hand, on the upper half plane we have $u_y = (y^{1-\beta}v)_y = (1-\beta)y^{-\beta}v + y^{1-\beta}v_y$, so

$$y^\beta u_y(x', y) = (1-\beta)v(x', y) + yv_y(x', y),$$

and

$$\lim_{y \rightarrow 0^+} y^\beta u_y(x', y) = (1-\beta)g(x') \approx r^{-\alpha},$$

the limit being uniform on compact subsets of B . □

Theorem 7.6. *If $u \in \mathbf{H}_{\text{loc}}^\beta(B_2)$ is a minimizer of \mathcal{J} in the ball B_2 with $\|u\|_{\mathbf{H}^\beta(B_2)} \leq E_0$, then the measure λ is absolutely continuous with respect to the Lebesgue measure, and for m -almost every $x \in B'_1(u)$ we have that*

$$\frac{d\lambda}{dm}(x) = 2 \lim_{y \rightarrow 0} |y|^\beta u_y(x', y) \approx \chi_{B_{1,0}(u)}(x) \text{dist}(x, F(u))^{-\alpha},$$

with constants depending on n, α and E_0 .

Proof. By Theorem 7.4 we only need to show absolute continuity in $B_{1,0}(u) \cup B'_{1,+}(u)$. For $x = (x', 0) \in B'_{1,+}(u)$ by [CS07, Lemma 4.2] we have that

$$\lim_{y \rightarrow 0} |y|^\beta u_y(x', y) = 0,$$

and, for $x \in B_{1,0}(u)$ we have seen in Lemma 7.5 that

$$\lim_{y \rightarrow 0} |y|^\beta u_y(x', y) \approx \text{dist}(x, F(u))^{-\alpha},$$

showing the second part of the statement.

Consider a ball $B_r(x_0)$ with $x_0 \in \mathbb{R}^n \times \{0\}$ and a collection of even smooth functions $\chi_{B_r} \leq \psi_k \leq \chi_{B_{r+\frac{1}{k}}}$. Then

$$\lambda(B_r) \leq - \int |y|^\beta \nabla u \cdot \nabla \psi_k \leq \lambda(B_{r+\frac{1}{k}}), \quad (7.7)$$

and for every $\varepsilon > 0$ we use the Green's theorem to get

$$-\int |y|^\beta \nabla u \cdot \nabla \psi_k = -\int_{|y| \leq \varepsilon} |y|^\beta \nabla u \cdot \nabla \psi_k - \int_{|y| = \varepsilon} |y|^\beta \psi_k \nabla u \cdot \nu \, dm.$$

Using the symmetry properties and taking limits,

$$-\int |y|^\beta \nabla u \cdot \nabla \psi_k = 2 \lim_{\varepsilon \rightarrow 0} \int \varepsilon^\beta \psi_k(x', \varepsilon) u_y(x', \varepsilon) \, dm(x'). \quad (7.8)$$

Next we want to apply the dominated convergence theorem. Let us begin by considering a ball $B_r(x_0) \subset B_1$ centered in the zero phase, with $\text{dist}(B'_r(x_0), F(u)) \geq 2r$. In this case, by Lemma 7.5 we have

$$\varepsilon^\beta u_y(x', \varepsilon) \lesssim r^{-\alpha}, \quad (7.9)$$

with constants depending perhaps on u and B_r as well.

If instead $B'_r(x_0) \subset B'_{1,+}(u)$, by [Vit18, Theorem 3.28] u is an even C^∞ function on $B'_r(x_0)$, so $|y|^\beta u_y = \mathcal{O}(|y|^{1+\beta})$. Thus

$$\varepsilon^\beta u_y(x', \varepsilon) \lesssim r^{2-2\alpha}. \quad (7.10)$$

In both cases, the dominated convergence theorem applies and

$$\lim_{\varepsilon \rightarrow 0} \int_{B_{r+\frac{1}{k}} \cap \{y=\varepsilon\}} \varepsilon^\beta \psi_k u_y \, dm = \int_{B'_{r+\frac{1}{k}}} \psi_k \lim_{\varepsilon \rightarrow 0} (\varepsilon^\beta u_y(\cdot, \varepsilon)) \, dm,$$

and by (7.7) and (7.8) we obtain

$$\lambda(B_r) \leq 2 \int_{B'_{r+\frac{1}{k}}} \psi_k \lim_{\varepsilon \rightarrow 0} (\varepsilon^\beta u_y(\cdot, \varepsilon)) \, dm \leq \lambda(B_{r+\frac{1}{k}}).$$

In particular $\lim_{\varepsilon \rightarrow 0} (\varepsilon^\beta u_y(\cdot, \varepsilon)) \in L^1_{\text{loc}}(B_{1,0}(u) \cup B'_{1,+}(u))$ and taking limits in k we get

$$\lambda(B_r) = 2 \int_{B'_r} \lim_{\varepsilon \rightarrow 0} (\varepsilon^\beta u_y(\cdot, \varepsilon)) \, dm.$$

□

A consequence of our control of the behavior of λ is that we can establish the existence of exterior corkscrews. We should note that exterior corkscrews can be also obtained by a purely geometric argument given the non-degeneracy and positive density of Theorem 2.3 (see, e.g. the proof of Proposition 10.3 in [DT15]).

Corollary 7.7. *If $u \in \mathcal{I}$ is a minimizer in B_2 with $\|u\|_{\mathbf{H}^\beta(B_2)} \leq E_0$, then $B'_{1,+}(u)$ satisfies the exterior corkscrew condition, i.e. there exists a constant C_1 such that for every $x \in F(u)$ and every $0 < r < \text{dist}(x, \partial B_1)$ one can find $x_0 \in B_r(x)$ so that*

$$B(x_0, C_1 r) \cap B'_{1,+}(u) = \emptyset.$$

Proof. This is a consequence of Theorems 7.4 and 7.6, and the positive density condition for the zero phase. Indeed, given a ball $B_r \subset \mathbb{R}^{n+1}$, combining both theorems we get

$$\begin{aligned} r^{n-\alpha} &\gtrsim \lambda(B_{1,0}(u) \cap B_r) \geq C_{E_0} \int_{B_{1,0}(u) \cap B_r} \text{dist}(x, \partial B_1)^{-\alpha} \\ &\geq C \left(\sup_{B_{1,0}(u) \cap B_r} \text{dist}(x, \partial B_1) \right)^{-\alpha} |B_{1,0}(u) \cap B_r|, \end{aligned}$$

and the positive density condition implies that

$$|B_{1,0}(u) \cap B_r| \geq C_{E_0} r^n.$$

Thus,

$$\sup_{B_{1,0}(u) \cap B_r} \text{dist}(x, \partial B_1) \geq C_{E_0} r,$$

which is equivalent to the exterior corkscrew condition. \square

7.3 Uniform Hölder character

The uniform non-degeneracy of Section 7.1 lets us conclude uniform control on the Hölder norm of u .

Theorem 7.8. *Let u be a minimizer of \mathcal{J} in B_r with $0 \in F(u)$. Then $|u(x)| \leq C|x|^\alpha$ for every $x \in \partial B_{r/2}$ with C depending only on n and α .*

Proof. Again we set v to be the \mathcal{L} -harmonic replacement of u inside of B_r as in (7.4). Let $\tilde{u} := v - u$, so that

$$\mathcal{L}\tilde{u} = \mathcal{L}v - \mathcal{L}u = -\lambda = -\nabla \cdot (|y|^\beta \nabla u),$$

and $\tilde{u} \in H_0^{1,2}(B_r; |y|^\beta)$.

Consider the Green function $G : B_r \times B_r \rightarrow \mathbb{R}$ such that $\mathcal{L}G(\cdot, z) = \delta_z$, and $G(\cdot, z) \in H_{\text{loc}}^{1,2}(\overline{B_r} \setminus \{z\})$ with null trace on ∂B_r (see [FJK82, Proposition 2.4]). By [FJK82, Proposition 2.1, Lemma 2.7] there exists $p_0 > 1$ so that \tilde{u} is the unique function in $H_0^{1,p_0}(B_r; |y|^\beta)$ such that $\mathcal{L}\tilde{u} = \lambda$, and moreover

$$\tilde{u}(z) = \int_{B_r} G(z, x) d\lambda(x), \quad (7.11)$$

for almost every $z \in B_r$.

Below, in Lemma 7.9, we will see that the equality (7.11) is in fact valid for every $z \in B_{r/4}$, that is, $\tilde{u} = \int_{B_r} G(\cdot, x) d\lambda(x)$. In particular

$$v(0) = \tilde{u}(0) = \int_{B_r} G(0, x) d\lambda(x).$$

Next we use the following estimate (see [FJK82, Theorem 3.3]): let $z, x \in B_{r/4}$. Then

$$G(z, x) \approx \int_{|x-z|}^r \frac{s ds}{w(B(x, s))},$$

where w is the A_2 weight $w(x) = |y|^\beta$. Computing, for $x = (x', y)$ we obtain

$$w(B(x, s)) \approx s^n \int_{y-s}^{y+s} |t|^\beta dt \approx s^n \max\{|y|, s\}^{\beta+1}.$$

First we assume that $n - 2\alpha > 0$. Thus, if $x \in B'_{r/4}$ then

$$G(z, x) \approx \int_{|x-z|}^r s^{-n-\beta} ds \approx |x-z|^{-n-\beta+1} = |x-z|^{2\alpha-n}. \quad (7.12)$$

Note that $\lambda(B_r) \leq Cr^{n-\alpha}$ by Theorem 7.4. Thus, writing $A_{t,s} := B_s \setminus B_t$, we have that

$$v(0) = \int_{B_r} G(0, x) d\lambda(x) \leq \int_{Cr^{2\alpha-n}}^\infty \lambda(\{x \in B_{r/4} : G(0, x) > t\}) dt + \int_{A_{r/4,r}} G(0, x) d\lambda(x).$$

By the strong maximum principle, the Green function in the annulus is bounded by $Cr^{n-2\alpha}$. This fact, together with Theorem 7.4, implies that

$$v(0) \leq \int_{Cr^{2\alpha-n}}^{\infty} \lambda \left(B_{Ct^{\frac{-1}{(n-2\alpha)}}} \right) dt + Cr^\alpha \leq C \int_{Cr^{2\alpha-n}}^{\infty} t^{-\frac{n-\alpha}{n-2\alpha}} dt + Cr^\alpha = Cr^\alpha.$$

By the mean value theorem we conclude that

$$\int_{\partial B_r} v d\sigma \leq Cr^\alpha,$$

where $d\sigma = |y|^\beta d\mathcal{H}^n$. The theorem follows by observing that, as in (7.5), the mean of v dominates u by $\sup_{\partial B_{r/2}} u \leq \sup_{\partial B_{r/2}} v \leq C \int_{\partial B_r} v d\sigma$.

In case $n - 2\alpha = 0$, which could only happen for $n = 1$ and $\alpha = 1/2$, estimate (7.12) reads as

$$G(z, x) \approx \log \left(\frac{r}{|x - z|} \right),$$

and the proof follows the same steps.

In case $n - 2\alpha < 0$, then estimate (7.12) reads as

$$G(z, x) \approx r^{n-2\alpha},$$

and the estimate is even better compared to the above. □

Lemma 7.9. $\int_{B_r} G(z, x) d\lambda(x)$ is continuous in $z \in B_{r/4}$.

Proof. Let $\varepsilon < r/2$ and let $z_1, z_2 \in B_{r/4}$, with $|z_1 - z_2| \leq \varepsilon/2$. Then

$$\begin{aligned} \int_{B_r} |G(z_1, x) - G(z_2, x)| d\lambda(x) &\leq \int_{B_r \setminus B_\varepsilon(z_1)} |G(z_1, x) - G(z_2, x)| d\lambda(x) \\ &\quad + \int_{B_\varepsilon(z_1)} G(z_1, x) d\lambda(x) + \int_{B_\varepsilon(z_1)} G(z_2, x) d\lambda(x). \end{aligned} \quad (7.13)$$

Next we use (7.12) and Theorems 7.4 and 7.6. By decomposing the domain on dyadic annuli, in case $n - 2\alpha > 0$ we get

$$\int_{B_\varepsilon(z_1)} G(z_1, x) d\lambda(x) \leq \sum_{j \leq 0} \int_{A_{2^{j-1}\varepsilon, 2^j\varepsilon}(z_1)} G(z_1, x) d\lambda(x) \lesssim \sum_{j \leq 0} \lambda(B_{2^j\varepsilon}(z_1)) (2^{j-1}\varepsilon)^{2\alpha-n} \lesssim \varepsilon^\alpha \sum_{j \leq 0} 2^{j\alpha}.$$

In case $n - 2\alpha = 0$ we obtain $\varepsilon^\alpha \sum_{j \leq 0} 2^{j\alpha} \log \left(\frac{r}{2^j\varepsilon} \right)$ on the right-hand side instead, and in case $n - 2\alpha < 0$ we obtain $\varepsilon^{n-\alpha} r^{2\alpha-n} \sum_{j \leq 0} 2^{j(n-\alpha)}$. In every case, fixing ε small enough this term can be as small as wanted. The same will happen with the last term on the right-hand side of (7.13).

On the other hand, by [FKS82, Theorem 2.3.12] Green's function is uniformly continuous on the set $\{(z, x) \in B_r \times B_r : |z - x| > \varepsilon\}$ so $|G(z_1, x_1) - G(z_2, x_2)| \leq \delta_\varepsilon(|z_1 - z_2| + |x_1 - x_2|)$ with $\delta_\varepsilon(t) \xrightarrow{t \rightarrow 0} 0$. Thus,

$$\int_{B_r \setminus B_\varepsilon(z_1)} |G(z_1, x) - G(z_2, x)| d\lambda(x) \leq \delta_\varepsilon(|z_1 - z_2|) \lambda(B_r) \rightarrow 0.$$

Assuming that $|z_1 - z_2|$ is small enough, we obtain that $\int_{B_r} |G(z_1, x) - G(z_2, x)| d\lambda(x)$ is as small as wanted and the claim follows. □

Remark 7.10. *In light of Theorem 7.8 and Caccioppoli inequality (see Section 3.1), arguing as in [CRS10b, Theorem 1.1] we obtain that every minimizer u in a ball B_r with $0 \in F(u)$ has uniform C^α character in $B_{r/2}$ and the same for the \mathbf{H}^β norm. Moreover, using [CRS10b, Theorem 1.2] we can find interior corkscrew points with constants not depending on these norms. This allows us to remove the a priori dependence on $\|u\|_{\mathbf{H}^\beta}$ from all of our results above.*

7.4 Lower estimates for the distributional fractional Laplacian

Next we bound the growth of the measure around a free boundary point from below. None of these results will be used in the present paper, but we include them to give a complete picture of the tools under consideration.

Theorem 7.11. *Let $u \in \mathbf{H}^\beta(B_{2r})$ be a minimizer of \mathcal{J} in B_{2r} such that $0 \in F(u)$. Then we have*

$$\lambda(B_r) \geq Cr^{n-\alpha}.$$

Proof. Let $\mathcal{L}u := -\nabla \cdot (|y|^\beta \nabla u)$ and let v be the \mathcal{L} -harmonic replacement of u in B_r (see (7.4)). Let $\tilde{u} := v - u$ and consider the Green function $G : B_r \times B_r \rightarrow \mathbb{R}$ as in the proof of Theorem 7.8.

Let $0 < \kappa < 1$ to be fixed later. By P1-P3 in Theorem 2.3 there exists a point $z_0 \in B_{\kappa r}$ with

$$u(z_0) \approx (\kappa r)^\alpha, \tag{7.14}$$

with constants depending only on n and α by Remark 7.10. By P1 there is a constant c such that for every $z \in B(z_0, c\kappa r)$ we have that $u(z) \approx (\kappa r)^\alpha$. Since λ is supported on the zero phase of u , the ball $B(z_0, c\kappa r)$ is away from its support, and

$$\tilde{u}(z) = \int_{B_r \setminus B(z_0, c\kappa r)} G(z, x) d\lambda(x).$$

Using the strong maximum principle (see [HKM06, Theorem 6.5]) and (7.12), for almost every $z \in B(z_0, c\kappa r/2)$ we get

$$\begin{aligned} \tilde{u}(z) &\leq \lambda(B_r) \sup_{x \in B_r \setminus B(z_0, c\kappa r)} G(z, x) = \lambda(B_r) \sup_{x \in B_{r/4} \setminus B(z_0, c\kappa r)} G(z, x) \\ &\approx \lambda(B_r) \sup_{x \notin B(z_0, c\kappa r)} |x - z|^{2\alpha-n} = \lambda(B_r)(c\kappa r)^{2\alpha-n}. \end{aligned}$$

That is,

$$\tilde{u}(z) \lesssim \lambda(B_r)(c\kappa r)^{2\alpha-n}. \tag{7.15}$$

On the other hand, note that u is continuous. By the Riesz representation theorem, there exists a probability measure $\omega_{\mathcal{L}}^z$ such that

$$v(z) = \int_{\partial B_r} u(x) d\omega_{\mathcal{L}}^z(x).$$

We can choose r so that ∂B_r intersects a big part of a corkscrew ball, i.e., assume that there exists a point $\xi_0 \in \partial B_r'$ which is the center of a ball $B'(\xi_0, cr)$ where u has positive values. This can be done by the interior corkscrew condition, with all the constants involved depending only on n and α . Then, changing the constant if necessary, all points $\xi \in B(\xi_0, cr)$ satisfy that $u(\xi) \geq Cr^\alpha$ by the non-degeneracy condition and the optimal regularity. Call $U := \partial B_r \cap B(\xi_0, cr)$. Then

$$v(z) \gtrsim r^\alpha \omega_{\mathcal{L}}^z(U).$$

But $\omega_{\mathcal{L}}^z(U)$ is bounded below by a constant by [HKM06, Lemma 11.21] and the Harnack inequality (use a convenient Harnack chain). All in all, we have that

$$v(z) \gtrsim r^\alpha. \quad (7.16)$$

Combining (7.15), (7.14) and (7.16) and choosing κ small enough, depending in n and α , we get

$$\lambda(B_r) \gtrsim \frac{\tilde{u}(z_0)}{(c\kappa r)^{2\alpha-n}} \geq \frac{Cr^\alpha - C'(\kappa r)^\alpha}{(c\kappa r)^{2\alpha-n}} \geq C_{n,\alpha} r^{n-\alpha},$$

for κ small enough.

In case $n-2\alpha = 0$, that is for $n = 1$ and $\alpha = 1/2$, using similar changes as in the proof of Theorem 7.8 we get $\tilde{u}(z) \lesssim \lambda(B_r) \sup_{x \in B(z_0, c\kappa r)} \log\left(\frac{r}{|x-z|}\right) \approx \lambda(B_r) |\log \kappa|$ instead of (7.15). In case $n-2\alpha < 0$, the proof is even easier than before. \square

Remark 7.12. *Theorem 7.11 implies that the $(n-\alpha)$ -Hausdorff measure of the free boundary is locally finite. This does not suffice to show finite perimeter of the positive phase and, therefore, we had to use the approach in Section 5.*

The following theorem summarizes the information that we have gathered so far about the measure λ .

Theorem 7.13. *If $u \in \mathbf{H}_{\text{loc}}^\beta(\Omega)$ is a minimizer of \mathcal{J} in Ω , then the measure λ is absolutely continuous with respect to the Lebesgue measure in $\Omega'(u)$. Moreover, given $x_0 \in F(u)$ and $r > 0$ such that $B_{2r}(x_0) \subset \Omega$, then*

$$\lambda(B_r(x_0)) \approx r^{n-\alpha}, \quad (7.17)$$

and for almost every $x \in B_r^i(x_0)$ we have that

$$\frac{d\lambda}{dm}(x) = 2 \lim_{y \rightarrow 0} |y|^\beta u_y(x', y) \approx \chi_{\Omega_0(u)}(x) \text{dist}(x, F(u))^{-\alpha},$$

with constants depending only on n and α .

8 Rectifiability of the singular set

In this section we use the Rectifiable-Reifenberg and quantitative stratification framework of Naber-Valtorta [NV17] to prove Hausdorff measure and structure results for the singular set. Recall that k_α^* is the first dimension in which there exists non-trivial α -homogeneous global minimizers to (1.1) defined in Section 5.

Theorem 8.1. *Let $u \in \mathbf{H}_{\text{loc}}^\beta(\Omega)$ be a minimizer of (1.1) in a domain Ω . Then $\Sigma(u)$ is $(n - k_\alpha^*)$ -rectifiable and for every $D \subset \subset \Omega$, we have*

$$\mathcal{H}^{n-k_\alpha^*}(\Sigma(u) \cap D) \leq C_{n,\alpha, \text{dist}(D, \partial\Omega)}.$$

Part of the power of this framework is that it is very general. One needs certain compactness properties on the minimizers and a connection between the drop in the monotonicity formula and the local flatness of the singular set (see Theorem 8.14 below). To avoid redundancy and highlight the original contributions of this article, we omit many details here and try to focus on the estimates needed to apply this framework to minimizers of (1.1). Whenever we omit details we will refer the interested reader to the relevant parts of [EE19].

The key first step is to introduce the appropriate formulation of quantitative stratification. First introduced by Cheeger and Naber [CN13] in the context of manifolds with Ricci curvature bounded from below, this is a way to quantify the intuitive fact that $F(u)$ should “look” $(n - k_\alpha^*)$ -dimensional near a point $x_0 \in F(u)$ at which the blow-ups have $(n - k_\alpha^*)$ -linearly independent translational symmetries.

8.1 Quantitative stratification for minimizers to \mathcal{J}

We have seen in Section 4.1 that homogeneous functions have linear spaces of translational symmetry. Here we want to quantify (both in terms of size and stability) how far a function is from having no more than k directions of translational symmetry.

Definition 8.2. *We write V^k for the collection of linear k -dimensional subspaces of \mathbb{R}^n . A function u is said to be k -symmetric if it is α -homogeneous with respect to some point, and there exists a $L \in V^k$ so that*

$$u(x+v) = u(x), \quad \text{for every } v \in L.$$

A function u is said to be (k, ϵ) -symmetric in a ball B if for some k -symmetric \tilde{u} we have

$$r^{-2-n} \int_B |y|^\beta |u - \tilde{u}|^2 dy < \epsilon.$$

Next we define the k -stratum $S^k(u)$, the (k, ϵ) -stratum $S_\epsilon^k(u)$ and the (k, ϵ, r) -stratum $S_{\epsilon, r}^k(u)$. A key insight here is to define these strata by the blow-ups having k or fewer symmetries as opposed to exactly k symmetries.

Definition 8.3. *Let $0 \leq k \leq n$, $0 < \epsilon < \infty$ and $0 < r < \text{dist}(x, \partial\Omega)$, let u be a continuous function in Ω and let $x \in F(u)$. We say that:*

- $x \in S^k(u)$ if u has no $(k+1)$ -symmetric blow-ups at x .
- $x \in S_\epsilon^k(u)$ if u is not $(k+1, \epsilon)$ -symmetric in $B_s(x)$ for every $0 < s \leq \min\{1, \text{dist}(x, \partial\Omega)\}$.
- $x \in S_{\epsilon, r}^k(u)$ if u is not $(k+1, \epsilon)$ -symmetric in $B_s(x)$ for every $r \leq s \leq \min\{1, \text{dist}(x, \partial\Omega)\}$.

If it is clear from the context we will omit u from the notation.

We now detail some standard properties of the strata defined above and how they interact with the free boundary $F(u)$. While the proofs are mostly standard, we give the details as the scaling associated to the problem (1.1) adds some technical difficulties. This proof also provides a blueprint for fleshing out the details in Sections 8.3 and 8.4.

Lemma 8.4. *Let $0 \leq j \leq k \leq n$, $0 < \epsilon \leq \tau < \infty$, $0 < r \leq s < \text{dist}(x, \partial\Omega)$, and let $u \in \mathbf{H}_{\text{loc}}^\beta(\Omega)$ be a minimizer in Ω . Then:*

1. $S^0 \subset S^1 \subset \dots \subset S^{n-1} = S^n = F(u)$. Moreover, for the reduced boundary, we have that $F_{\text{red}}(u) \subset S^{n-1} \setminus S^{n-2}$ and $\Sigma(u) \subset S^{n-k_\alpha^*}$.
2. We have $S_\tau^j \subset S_\epsilon^k \subset S^k$ and, moreover, $S^k = \bigcup_{\epsilon > 0} S_\epsilon^k$.
3. Also $S_\tau^j \subset S_{\tau, r}^j \subset S_{\epsilon, s}^k$ and, moreover, $S_\epsilon^k = \bigcap_{r > 0} S_{\epsilon, r}^k$.
4. The sets S_ϵ^k are closed, in both x and u : if $u_i \xrightarrow{L_{\text{loc}}^2(\Omega; |y|^\beta)} u$ and $x_i \rightarrow x$ with $x_i \in S_\epsilon^k(u_i)$, then $x \in S_\epsilon^k(u)$.

5. If $u_i \xrightarrow{L^2_{\text{loc}}(\Omega; |y|^\beta)} u$, $\epsilon_i \rightarrow 0$, and u_i are (k, ϵ_i) -symmetric in B_1 , then u is k -symmetric in B_1 .

Proof. 1. The inclusions $S^k \subset S^{k+1}$ of the first property are trivial. The last equalities are consequences of the non-degeneracy. The fact that $F_{\text{red}}(u) \cap S^{n-2} = \emptyset$ can be deduced from the Hausdorff convergence of the free boundaries described in Lemma 3.4 and Theorem 2.4. Finally, $\Sigma(u) \subset S^{n-k^*}$ is a consequence of Lemmas 4.5 and 5.2.

2. The inclusions $S^j_\tau \subset S^k_\epsilon$ of the second property come from the definitions: if $x \notin S^k_\epsilon$ then there exist a ball $B \subset \Omega$ centered at x and a $(k+1)$ -symmetric \tilde{u} so that $r(B)^{-2-n} \int_B |y|^\beta |u - \tilde{u}|^2 dy < \epsilon \leq \tau$. But \tilde{u} is also $(j+1)$ -symmetric. Thus, $x \notin S^j_\tau$.

The fact that $S^k_\epsilon \subset S^k$ is a consequence of the uniform convergence on Lemma 3.4: if $x \notin S^k$, then u has a $(k+1)$ -symmetric blow-up sequence $u_i \rightarrow u_0$ at x converging uniformly. Thus,

$$\begin{aligned} \int_{B_{\rho_i}(x)} |y|^\beta \left| u(x) - \rho_i^\alpha u_0 \left(\frac{x-x_0}{\rho_i} \right) \right|^2 dx &= \rho_i^{\beta+2\alpha+n+1} \int_{B_1(x)} |y|^\beta \left| \frac{u(x_0 + \rho_i x)}{\rho_i^\alpha} - u_0(x) \right|^2 dx \\ &\leq \rho_i^{n+2} \omega(B_1) \|u_i - u_0\|_{L^\infty}. \end{aligned}$$

That is,

$$\rho_i^{-n-2} \int_{B_{\rho_i}(x)} |y|^\beta \left| u(x) - \rho_i^\alpha u_0 \left(\frac{x-x_0}{\rho_i} \right) \right|^2 dx \xrightarrow{i \rightarrow \infty} 0,$$

and therefore, for every ϵ there exists a ball small enough so that u is $(k+1, \epsilon)$ -symmetric in it. In particular $S^k \supset \bigcup_{\epsilon > 0} S^k_\epsilon$.

To see the converse, assume that $x \notin \bigcup_\epsilon S^k_\epsilon$. Then for every $i \in \mathbb{N}$ there exist a $(k+1)$ -symmetric function \tilde{u}_i , invariant with respect to $L_i \in V^{k+1}$ and $r_i < \min\{1, \text{dist}(x, \partial\Omega)\}$ such that

$$\frac{1}{r_i^{n+2}} \int_{B_{r_i}} |y|^\beta |u(x) - \tilde{u}_i(x)|^2 dx < \frac{1}{i}.$$

In the case when r_i stays away from zero, since $r_i < 1$, we can take a subsequence converging to $r_0 \in (0, 1)$, and one can see that u is $(k+1)$ -symmetric in the ball $B_{r_0}(x_0)$. Otherwise, consider $u_i := \frac{u(x_0 + r_i x)}{r_i^\alpha}$ and $\tilde{u}_{i,i} = \frac{\tilde{u}_i(x_0 + r_i x)}{r_i^\alpha}$. Taking subsequences, we can assume that $L_i \rightarrow L_0$ locally in the Hausdorff distance, and that $u_i \rightarrow u_0$ locally uniformly. One can check also using the Hölder character of u that $\{\tilde{u}_{i,i}\}$ is uniformly bounded in $L^2(B; |y|^\beta)$, so taking subsequences again, we can assume the existence of \tilde{u}_0 so that $\tilde{u}_{i,i} \rightarrow \tilde{u}_0$ in $L^2(B; |y|^\beta)$. This function will be $(k+1)$ -symmetric, being invariant in the directions of L_0 . By the triangle inequality we get

$$\int_{B_1} |y|^\beta |u_0 - \tilde{u}_0|^2 dx \lesssim \int_{B_1} |y|^\beta |u_0 - u_i|^2 dx + \int_{B_1} |y|^\beta |u_i - \tilde{u}_{i,i}|^2 dx + \int_{B_1} |y|^\beta |\tilde{u}_{i,i} - \tilde{u}_0|^2 dx.$$

The first and the last integrals converge to zero by our choice of the subsequence. For the middle term just change variables as before:

$$\int_{B_1} |y|^\beta |u_i - \tilde{u}_{i,i}|^2 dx = \frac{1}{r_i^{n+2}} \int_{B_{r_i}} |y|^\beta |u(x) - \tilde{u}_i(x)|^2 dx \rightarrow 0.$$

Thus we have that $u_0 = \tilde{u}_0$ and, therefore, $x \notin S^k$.

3. The inclusions $S^j_\tau \subset S^j_{r,r} \subset S^k_{\epsilon,s}$ of the third property come from the definitions and thus, $S^k_\epsilon \subset \bigcap_{r>0} S^k_{\epsilon,r}$. The converse implication is also trivial.

4. The closedness is obtained by a contradiction argument again. It is straightforward but we write it here for the sake of completeness.

Assume by contradiction that $x \notin S_\epsilon^k(u)$. Then there exist a $(k+1)$ -symmetric function \tilde{u} and a radius r such that

$$\epsilon_0 := \frac{1}{r^{n+2}} \int_{B_r(x)} |y|^\beta |u(x) - \tilde{u}(x)|^2 dx < \epsilon.$$

Let $\tau < 1$ to be fixed and consider $i_0 \in \mathbb{N}$ so that $B_{\tau r}(x_i) \subset B_r(x)$ for every $i \geq i_0$. By the triangle inequality

$$\frac{1}{(\tau r)^{n+2}} \int_{B_{\tau r}(x_i)} |y|^\beta |u_i(x) - \tilde{u}(x)|^2 dx \leq \frac{1}{(\tau r)^{n+2}} \|u_i - u\|_{L^2(B_{\tau r}(x_i); |y|^\beta)}^2 + \frac{\epsilon_0}{\tau^{n+2}}.$$

We define τ so that $\frac{\epsilon_0}{\tau^{n+2}} = \frac{\epsilon + \epsilon_0}{2}$. Choose i_0 big enough so that every $i \geq i_0$ satisfies that $\|u_i - u\|_{L^2(B_{\tau r}(x_i); |y|^\beta)}^2 < (\tau r)^{n+2} \frac{\epsilon - \epsilon_0}{2}$. Then $x_i \notin S_\epsilon^k(u_i)$, contradicting the hypothesis.

5. Assume that \tilde{u}_i is invariant with respect to $L_i \in V^{k+1}$ and

$$\int |y|^\beta |u_i - \tilde{u}_i|^2 \leq \epsilon_i.$$

Consider a subsequence $\{u_i\}$ so that the varieties $L_i \rightarrow L$ locally in the Hausdorff distance. Using the triangle inequality as in 4 it follows that u is (k, δ_i) -symmetric with $\delta_i \rightarrow 0$. \square

Proposition 8.5. *There exists $\epsilon(n, \alpha) > 0$ such that if $u \in \mathbf{H}_{\text{loc}}^\beta(\Omega)$ is a minimizer of \mathcal{J} in a domain $\Omega \subset \mathbb{R}^{n+1}$, then $\Sigma(u) \subset S_\epsilon^{n-k_\alpha^*}(u)$.*

Proof. It is enough to show that if u is a minimizer of \mathcal{J} in $B_2(0)$, then $\Sigma(u) \cap B_1(0) \subset S_\epsilon^{n-k_\alpha^*}(u)$.

By contradiction, let us assume that there is a sequence of positive numbers $\epsilon_i \xrightarrow{i \rightarrow \infty} 0$, functions u_i minimizing \mathcal{J} in $B_2(0)$ and $x_i \in \Sigma(u_i) \cap B_1(0)$, $r_i \in (0, 1]$, with u_i being $(n - k_\alpha^* + 1, \epsilon_i)$ -symmetric in $B_{r_i}(x_i)$, and let L_i be an $(n - k_\alpha^* + 1)$ -dimensional subspace that leaves invariant one of the admissible $(n - k_\alpha^* + 1)$ -symmetric approximants. By rescaling we can assume that $r_i = 1$.

Passing to a subsequence we can assume that $L_i \rightarrow L_0 \in V^{n-k_\alpha^*+1}$ locally in the Hausdorff distance and $x_i \rightarrow x_0$. By the compactness results in Lemma 3.4 we have a uniform limit u_0 which is a minimizer as well, and it is $(n - k_\alpha^* + 1)$ -symmetric with invariant manifold L_0 . By Lemma 4.4 any blow-up $u_{0,0}$ at x_0 will be $(n - k_\alpha^* + 1)$ -symmetric as well. Applying Lemma 4.5 $(n - k_\alpha^* + 1)$ times we find that the restriction of $u_{0,0}$ to the orthogonal manifold L_0^\perp is a $(k_\alpha^* - 1)$ -dimensional minimal cone which, by Lemma 5.2 is the trivial solution, and so is $u_{0,0}$. Thus, x_0 is a regular point for u_0 .

On the other hand, the Hausdorff convergence of Lemma 3.4 together with the improvement of flatness of Theorem 2.4 imply that for i big enough $x_i \in F_{\text{red}}(u_i)$, reaching a contradiction. \square

8.2 The Refined Covering Theorem

Our estimates on the size and structure of the singular set $\Sigma(u)$ come from similar results concerning the $S_\epsilon^k(u)$. In particular, we prove the following covering result:

Theorem 8.6. *Let $u \in \mathbf{H}^\beta(B_5)$ be a minimizer to (1.1) in B_5 with $0 \in F(u)$. For given real numbers $\epsilon > 0$, $0 < r \leq 1$ and every natural number $1 \leq k \leq n - 1$, we can find a collection of balls $\{B_r(x_i)\}_{i=1}^N$ with $N \leq C_{n,\alpha,\epsilon} r^{-k}$ such that*

$$S_{\epsilon,r}^k(u) \cap B_1 \subset \bigcup_i B_r(x_i).$$

In particular, $|B'_r(S_{\epsilon,r}^k \cap B_1)| \leq C_{n,\alpha,\epsilon} r^{n-k}$ for every $0 < r \leq 1$ and

$$\mathcal{H}^k(S_\epsilon^k(u) \cap B_1) \leq C_{n,\alpha,\epsilon}.$$

From Proposition 8.5 and Theorem 8.6 we can conclude the following corollary which comprises the second part of Theorem 8.1 above.

Corollary 8.7. *If $u \in \mathbf{H}^\beta(B_5)$ is a minimizer to (1.1) in B_5 with $0 \in F(u)$, then $\Sigma(u)$ is $(n - k_\alpha^*)$ -rectifiable and for every $0 < r \leq 1$ we have*

$$|B_r(\Sigma(u) \cap B_1)| \leq C_{n,\alpha} r^{k_\alpha^*}.$$

In particular,

$$\mathcal{H}^{n-k_\alpha^*}(\Sigma(u) \cap B_1) \leq C_{n,\alpha}.$$

Rectifiability is encoded in the following result. We omit the details of proof here but it is a consequence of the packing result above, the Rectifiable-Reifenberg theorem of [NV17] and Theorem 8.14 below. For more details see Sections 2 and 8 of [EE19] (particularly Theorem 2.2 in the former and the proof of Theorem 1.12 in the latter).

Theorem 8.8. *Let u be a non-negative, even minimizer to (1.1) in a domain Ω . Then $S_\epsilon^k(u)$ is k -rectifiable for every ϵ and, hence, each stratum $S^k(u)$ is k -rectifiable as well.*

The proof of Theorem 8.6 follows from inductively applying the following, slightly more technical, packing result (for details see Section 4 of [EE19]).

Theorem 8.9. *Let $\epsilon > 0$. There exists $\eta(n, \alpha, \epsilon)$ such that, for every minimizer $u \in \mathbf{H}^\beta(B_5)$ of \mathcal{J} in B_5 with $0 \in F(u)$ and $0 < R < 1/10$, there is a finite collection \mathcal{U} of balls B with center $x_B \in S_{\epsilon, \eta R}^k$ and radius $R \leq r_B \leq 1/10$ which satisfy the following properties:*

A) *Covering control:*

$$S_{\epsilon, \eta R}^k \cap B_1 \subset \bigcup_{B \in \mathcal{U}} B.$$

B) *Energy drop: For every $B \in \mathcal{U}$,*

$$\text{either } r_B = R, \quad \text{or } \sup_{2B} \Psi_{2r_B}^u \leq \sup_{B_2} \Psi_2^u - \eta.$$

C) *Packing:*

$$\sum_{B \in \mathcal{U}} r_B^k \leq c(n, \alpha, \epsilon).$$

We construct the balls of Theorem 8.9 using a “stopping time” or “good ball/bad ball” argument. Much of this argument uses harmonic analysis and geometric measure theory and is completely independent of the original problem (1.1). However, there are a few places in which we need to connect the behavior of minimizers to the geometric structure of the singular set. Here we will sketch the “good ball/bad ball” argument, taking for granted the estimates needed to apply this argument to our functional. In the next few subsection we will provide these estimates. For more details on the construction itself we refer the reader to Section 7 in [EE19].

Outline of the Construction in Theorem 8.9 To find this covering we define good and bad balls as follows: imagine our ball, B , has radius 1. We say that B is a *good ball*, if at every point in $x \in S_\epsilon^k(u) \cap B$ the monotone quantity centered at that point at some small scale, ρ , is not much smaller than the monotone quantity on ball B (we say these points have “small density drop”). A ball B is a *bad ball* if all the points in $S_\epsilon^k(u) \cap B$ with small density drop are contained in a small neighborhood of a $(k - 1)$ -plane. This dichotomy follows from Theorem 8.10 in Section 8.3.

In a good ball of radius r we cover $S_\epsilon^k(u)$ with balls of radius ρr iterating the construction until we find a bad ball or until the radius of the ball becomes very small. In a bad ball, we cover $S_\epsilon^k(u)$

away from the $(k-1)$ -plane without much care. Close to the $(k-1)$ -plane we cover $S_\varepsilon^k(u)$ with balls of radius ρr iterating the construction until we reach a good ball or until the radius of the ball becomes very small.

Inside long strings of good balls, the packing estimates follow from powerful tools in geometric measure theory (see Theorem 8.13 below) and the connection between the drop in monotonicity and the local flatness of the singular strata (see Theorem 8.14 below). We give more details in Section 8.4.

Inside long strings of bad balls each of which is near the $(k-1)$ -plane of the previous bad ball, we have even better packing estimates than expected (as we are effectively well approximated by planes which are lower dimensional). This leaves only points which are in many bad balls and in most of those balls they are far away from the $(k-1)$ -plane. However, at these points the monotone quantity drops a definite amount many times, which contradicts either finiteness or monotonicity. This implies that the points and scales inside the bad balls which are not close to the $(k-1)$ -plane form a negligible set (the technical term is a Carleson set). We give more information about the bad balls in Section 8.3.

8.3 Tools for bad balls: key dichotomy

Theorem 8.10 (Key dichotomy). *Let $\varepsilon, \rho, \gamma, \eta' > 0$ be fixed numbers with $\rho\gamma < 2$. There exists an $\eta_0(n, \alpha, \varepsilon, \rho, \gamma, \eta') < \rho/100$ such that for every $\eta \leq \eta_0$, every $r > 0$, every $E > 0$ and every minimizer $u \in \mathbf{H}^\beta(B_{4r})$ of \mathcal{J} in B_{4r} with $0 \in F(u)$ and $\sup_{B_r} \Psi_{2r}^u \leq E$, then either*

- $\Psi_{\gamma\rho r}^u \geq E - \eta'$ on $S_{\varepsilon, \eta r}^k \cap B_r$, or
- there exists $\ell \in L^{k-1}$ so that $\{x \in B_r : \Psi_{2\eta r}^u(x) \geq E - \eta\} \subset B_{\rho r}(\ell)$.

The key dichotomy is a direct consequence of the Lemma 8.11 below. The core idea is to make effective the following assertion: if u is k -symmetric, then along the invariant manifold the Allen-Weiss density is constant, and every point away from the manifold will have $(k+1)$ -symmetric blow-ups by Lemma 4.4.

Lemma 8.11. *Let $\varepsilon, \rho, \gamma, \eta' > 0$ be fixed numbers with $\gamma\rho < 2$. There exist $\eta_0, \theta > 0$ such that for every $\eta < \eta_0$, every $E > 0$ and every minimizer u of \mathcal{J} in B_4 with $0 \in F(u)$ and $\sup_{B_1} \Psi_2^u \leq E$, if there exist $w_0, \dots, w_k \in B_1$ and affine manifolds $L^i := \langle w_0, \dots, w_i \rangle \in V^i$ with*

$$w_i \notin B_\rho(L^{i-1}), \quad \text{and} \quad \Psi_{2\eta}^u(w_i) \geq E - \eta \quad \text{for every } i \in \{0, \dots, k\},$$

then,

$$\Psi_{\gamma\rho}^u(x) \geq E - \eta' \quad \text{on } B_\theta(L^k) \cap B_1 \tag{8.1}$$

and

$$S_{\varepsilon, \eta}^k \cap B_1 \subset B_\theta(L^k) \tag{8.2}$$

The proof follows (with only minor modifications) the proof in [EE19, Lemma 3.3]. We end this subsection by formally defining the good/bad balls alluded to above:

Definition 8.12. *Let $x \in B_2$, $0 < R < r < 2$ and u be a minimizer to \mathcal{J} in B_5 . We say that the ball $B_r(x)$ is good if*

$$\Psi_{\gamma\rho r}^u \geq E - \eta' \quad \text{on } S_{\varepsilon, \eta R}^k \cap B_r(x),$$

and otherwise we say that $B_r(x)$ is bad.

By Theorem 8.10 in any bad ball B there exists an affine $(k-1)$ -manifold ℓ_B with

$$\{w \in B : \Psi_{2\eta r}^u(w) \geq E - \eta\} \subset B_{\rho r}(\ell_B^{k-1}). \tag{8.3}$$

8.4 Tools for good balls: packing estimates and GMT

In this section we control the local flatness of the singular strata by the drop in monotonicity. To do this we introduce a key tool from geometric measure theory which estimates the flatness of a set. Given a Borel measure μ , a point x and a radius r , the beta coefficient is defined as follows:

$$\beta_{\mu,2}^k(B_r(x))^2 := \beta_{\mu,2}^k(x,r)^2 = \inf_{L \in V_a^k} \frac{1}{r^k} \int_{B_r(x)} \frac{\text{dist}(z,L)^2}{r^2} d\mu(z) \quad (8.4)$$

where V_a^k stands for the collection of k -dimensional affine sets of \mathbb{R}^n . The beta coefficients are meant to measure in a scale invariant way how far is a measure from being flat, in this case in the L^2 distance, although other L^p versions have been used in the literature for $1 \leq p \leq \infty$ quite often, dating back to [Jon90] (for the L^∞ version) and David-Semmes [DS93] (for the L^p version).

If we control the size of the β^k 's we can conclude size and structure estimates on the measure μ . The following theorem says exactly this and represents a major technical achievement. It differs (importantly) from prior work in this area by the lack of *a priori* assumptions on the upper or lower densities of the measure involved.

Theorem 8.13 (Discrete-Reifenberg Theorem, see [NV17, Theorem 3.4]). *Let $\{B_{r_q}(q)\}_q$ be a collection of disjoint balls, with $q \in B_1(0)$ and $0 < r_q \leq 1$, and let μ be the packing measure $\mu := \sum_q r_q^k \delta_q$, where δ_q stands for the Dirac delta at q . There exist constants $\tau_{DR}, C_{DR} > 0$ depending only on the dimension such that if*

$$\int_0^{2r} \int_{B_r(x)} \beta_{\mu,2}^k(z,s)^2 d\mu(z) \frac{ds}{s} \leq \tau_{DR} r^k \quad \text{for every } x \in B_1(0), 0 < r \leq 1,$$

then

$$\mu(B_1(0)) = \sum_q r_q^k \leq C_{DR}.$$

To obtain the packing estimates required for the Discrete-Reifenberg Theorem, we need to control the beta coefficients. The key estimate of this entire framework lies in the following theorem, which shows the drop in monotonicity at a given point and a given scale controls the beta coefficient at a comparable scale.

Theorem 8.14. *Let $\epsilon > 0$ be given. There exist $\delta(n, \alpha, \epsilon)$ and $c(n, \alpha, \epsilon)$ such that for every $u \in \mathbf{H}^\beta(B_{5r})$ minimizing \mathcal{J} in $B_{5r}(x)$ with $x \in F(u)$ and*

$$\begin{cases} u \text{ is } (0, \delta)\text{-symmetric in } B_{4r}(x) \\ u \text{ is not } (k+1, \epsilon)\text{-symmetric in } B_{4r}(x), \end{cases} \quad (8.5)$$

and every Borel measure μ , we have that

$$\beta_{\mu,2}^k(B_r(x))^2 \leq \frac{c(n, \alpha, \epsilon)}{r^k} \int_{B_r(x)} (\Psi_{4r}^u(w) - \Psi_r^u(w)) d\mu(w). \quad (8.6)$$

We follow the proof of [EE19, Theorem 5.1] closely. First the authors give an explicit formula for the beta coefficients.

Lemma 8.15. *Let X be the center of mass of a Borel measure μ on $B = B_r(x)$. Let $\{\lambda_i\}_{i=1}^n$ be the decreasing sequence of eigenvalues of the non-negative bilinear form*

$$Q(v, w) := \int_B (v \cdot (z - X))(w \cdot (z - X)) d\mu(z),$$

and let $\{v_i\}_{i=1}^n$ be a corresponding orthonormal sequence of eigenvectors, that is $v_i \cdot v_j = \delta^{ij}$ and $Q(v_i, v) = \lambda_i v_i \cdot v$. Then

$$\beta_{\mu,2}^k(B)^2 = \frac{1}{r^k} \int_B \frac{\text{dist}(z, L^k)^2}{r^2} d\mu(z) = \frac{\mu(B)}{r^k} \frac{(\lambda_{k+1} + \dots + \lambda_n)}{r^2},$$

where $L^k := X + \text{span}\{v_1, \dots, v_k\}$.

Next we find a relation between the eigenvalues of Q and Allen-Weiss' energy.

Lemma 8.16. *Under the hypothesis of Lemma 8.15, for every $u \in \mathbf{H}^\beta(B_{5r})$ minimizing \mathcal{J} in $B_{5r}(x)$ and every $i \leq n$, we have that*

$$\lambda_i \frac{2}{r^{n+2}} \int_{A_{2r,3r}(x)} |y|^\beta (v_i \cdot Du(z))^2 dz \leq C \int_{B_r(x)} (\Psi_{4r}^u(w) - \Psi_r^u(w)) d\mu(w). \quad (8.7)$$

Proof. The argument follows as in [EE19, (18) and below]. In formula (18) one needs to change $u(z)$ by $\alpha u(z)$, which can be done with exactly the same argument. \square

Finally, using compactness, we bound the left-hand side of (8.16) from below.

Lemma 8.17. *Let $\epsilon > 0$ be given. There exists a $\delta(n, \alpha, \epsilon)$ and $c(n, \alpha, \epsilon)$ such that, for every orthonormal basis $\{v_i\}_{i=1}^n$ and every $u \in \mathbf{H}^\beta(B_{5r})$ minimizing \mathcal{J} in $B_{5r}(x)$ with $x \in F(u)$ and satisfying (8.5), we have that*

$$\frac{1}{c(n, \alpha, \epsilon)} \leq r^{-n} \int_{A_{2r,3r}(x)} |y|^\beta \sum_{i=1}^{k+1} (v_i \cdot Du(z))^2 dz. \quad (8.8)$$

Proof. The proof follows that of [EE19, (19)] and we omit it. \square

Proof of Theorem 8.14. By Lemmas 8.15, 8.17 and 8.16 we get that

$$\begin{aligned} \beta_{\mu,2}^k(B)^2 &\leq \frac{\mu(B)}{r^{k+2}} (n-k) \lambda_{k+1} \\ &\leq \frac{\mu(B)}{r^k} (n-k) c(n, \alpha, \epsilon) \sum_{i=1}^{k+1} \frac{\lambda_i}{r^{n+2}} \int_{A_{2r,3r}(x)} |y|^\beta (v_i \cdot Du(z))^2 dz \\ &\leq \frac{c(n, \alpha, \epsilon)}{r^k} \int_{B_r(x)} (\Psi_{4r}^u(w) - \Psi_r^u(w)) d\mu(w). \end{aligned}$$

\square

Acknowledgements

M.E. was partially supported by an NSF postdoctoral fellowship, NSF DMS 1703306 and by David Jerison's grant DMS 1500771. A.K. acknowledges Financial support from the Spanish Ministry of Economy and Competitiveness, through the María de Maeztu Programme for Units of Excellence in R&D (MDM-2014-0445). M.P. was funded by the European Research Council under the grant agreements 320501-ANGEOM and 834728-QUAMAP. G.S. has received funding from the European Union's Horizon 2020 research and innovation programme under Marie Skłodowska-Curie grant agreement No 665919. A.K., M.P. and G.S. were also partially funded by 2017-SGR-0395 (Catalonia) and MTM-2016-77635-P (MINECO, Spain). Y.S. is partially supported by the Simons foundation.

M.E. would also like to thank Nick Edelen for many fruitful conversations regarding the quantitative stratification and rectifiable Reifenberg framework.

A.K., M.P., G.S. would like to thank Xavier Cabré, Tomás Sanz, Matteo Cozzi, Albert Mas, Maria del Mar González, Luis Silvestre and Stefano Vita for some conversations around [Vit18]. They would also like to thank Mihalis Mouroglou for some conversations regarding the degenerate elliptic measure.

Appendix

A Relation with the nonlocal Bernoulli problem

As in [DV17, Lemma 2.1], we see that the study of minimizers of \mathcal{J} includes the study of minimizers of J .

Proposition A.1. *If f is a minimizer of J in the unit ball of \mathbb{R}^n then $f * P_y$ is a minimizer of \mathcal{J} in every ball B such that $B' \subset B_1$.*

*If $u = f * P_y$ is a minimizer of \mathcal{J} , then f is a minimizer for J . In particular, if u is a minimizer of \mathcal{J} in every ball, positive outside the hyperplane $\{y = 0\}$, and $u(x, y) = \mathcal{O}(|(x, y)|^\alpha)$, then $u|_{\mathbb{R}^n \times \{0\}}$ is a minimizer for J in every ball.*

We follow [DV17, Lemma 2.1], that is, we use the following result from [CRS10a, Section 7].

Lemma A.2 (see [CRS10a, Section 7]). *Let f, g satisfy that $J_0(f, B_1), J_0(g, B_1) < \infty$, and suppose that $f - g$ is compactly supported in $B_1 \subset \mathbb{R}^n$. Then we have that*

$$J_0(g, B_1) - J_0(f, B_1) = c_{n,\alpha} \inf \int_{\Omega} |y|^\beta (|\nabla v(x, u)|^2 - |\nabla(f * P_y)(x)|^2),$$

where the infimum is taken among all the symmetric bounded Lipschitz domains Ω with the property that $\Omega \cap (\mathbb{R}^n \times \{0\}) \subset B_1$ and among all symmetric functions v with trace g satisfying that $v - f * P_y$ is compactly supported on Ω .

Proof of Proposition A.1. Let f be a minimizer of J in the unit ball of \mathbb{R}^n and let B_r be a ball such that $B'_r \subset B_1$. We want to show that $u := f * P_y$ is a minimizer of \mathcal{J} in B_r .

Let $v : \mathbb{R}^{n+1} \rightarrow \mathbb{R}$ so that $v \equiv u$ in $\mathbb{R}^{n+1} \setminus B_r$ and $v \in H^1(\beta, B_r)$. Let g be the trace of v in $\mathbb{R}^n \times \{0\}$. By Lemma A.2 we have that

$$J_0(g, B_1) - J_0(f, B_1) \leq c_{n,\alpha} \int_{B_{r+\varepsilon}} |y|^\beta (|\nabla v|^2 - |\nabla u|^2) \quad (\text{A.1})$$

for every $\varepsilon > 0$.

In particular, since $g|_{(B_r)^\varepsilon} \equiv 0$, g is an admissible competitor for f and $J(f, B_1) \leq J(g, B_1)$, i.e.,

$$\begin{aligned} J_0(g, B_1) - J_0(f, B_1) &\geq -m(\{g > 0\} \cap B_1) + m(\{f > 0\} \cap B_1) \\ &= m(\{u > 0\} \cap B'_r) - m(\{v > 0\} \cap B'_r). \end{aligned} \quad (\text{A.2})$$

The proposition follows combining (A.1) and (A.2) and letting $\varepsilon \rightarrow 0$.

The converse follows the same sketch: every global minimizer can be expressed as the Poisson extension of its restriction to the hyperplane by Proposition B.1 and it is left to the reader. \square

As a consequence of the previous proposition, all the results that we have proven for minimizers of \mathcal{J} also apply to minimizers of J :

Corollary A.3. *If $u : \mathbb{R}^n \rightarrow \mathbb{R}$ is a minimizer to J in $B_2 \subset \mathbb{R}^n$ and $0 \in F(u)$, then $\|u\|_{C^\alpha(B_1)} \leq C$, it satisfies the nondegeneracy condition $u(x) \geq C \text{dist}(x, F(u))^\alpha$ for $x \in B_1$, the positive phase satisfies the corkscrew condition, every blow-up limit is α -homogeneous, and the boundary condition in (1.2) is satisfied at $F_{\text{red}}(u)$.*

Moreover, the positive phase $\{u > 0\} \cap B_1$ is a set of finite perimeter, the singular set is an $(n-3)$ -rectifiable set, it is discrete whenever $n = 3$ and it is empty if $n \leq 2$.

All the constants depend only on n and α .

B Uniqueness of extensions

In Proposition A.1 we have used the following result, included in [CRS10b, Proposition 3.1]. Here we provide a proof which is different than the one appearing in [CRS10b].

Proposition B.1. *Let $\alpha \in (0, 1)$, $\beta = 1 - 2\alpha$, and set $\mathcal{L}u = -\operatorname{div}(|y|^\beta \nabla u)$ in \mathbb{R}^{n+1} . Suppose that $v : \overline{\mathbb{R}_+^{n+1}} \rightarrow \mathbb{R}$ is nonnegative outside \mathbb{R}^n , it is a solution to $\mathcal{L}v = 0$ in \mathbb{R}_+^{n+1} with $v(x', 0) = 0$ for all $x' \in \mathbb{R}^n$ and $|v(x)| \leq C|x|^\alpha$. Then $v \equiv 0$.*

Proof. First, since $|y|^\beta$ is C^∞ away from the hyperplane \mathbb{R}^n , $v \in C_{\text{loc}}^\infty(\mathbb{R}_+^{n+1})$. Let now $i \in \{1, \dots, n\}$, and set

$$f_m(x) = \frac{v\left(x + \frac{1}{m}e_i\right) - v(x)}{1/m}.$$

Let $B_r = B_r(x', 0)$ be a ball centered at $(x', 0) \in \mathbb{R}^n \times \{0\}$ with radius r , and let B_{2r} be its double ball. Set also $w(x) = w(x', y) = y^\beta$ for $y > 0$. Since f_m is a solution of $\mathcal{L}f_m = 0$ in $B_r^+ = B \cap \mathbb{R}_+^{n+1}$, [FKS82, Theorem 2.4.3] shows that

$$\max_{B_r^+} |f_m(x)| \leq C \left(\frac{1}{w(B_{2r}^+)} \int_{B_{2r}^+} |f_m|^2 w \right)^{1/2}.$$

From convergence of difference quotients (similarly to [Eva98, Theorem 3, page 277]), if $v \in H^1(\beta, B_{2r}^+)$, the last estimate will imply that f_m is uniformly bounded in B_r^+ by a constant C_r . Therefore, from the boundary Caccioppoli estimate ([FKS82, (2.4.2)]) we have that

$$\int_{B_{r/2}^+} |\nabla f_m|^2 w \leq \frac{C}{r^2} \int_{B_r^+} |f_m|^2 w \leq \frac{C}{r^2} \int_{B_r^+} C_r^2 w \leq C_{n,r,w} < \infty,$$

hence $\{f_m\}$ is bounded in $H^1(\beta, B_{r/2}^+)$. From weak compactness, a subsequence of $\{f_m\}$ converges to a solution of $\mathcal{L}u = 0$ in $B_{r/2}^+$, and since $f_m \rightarrow \partial_i v$ pointwise, we obtain that $\partial_i v$ is an $H^1(\beta, B_{r/2}^+)$ solution in $B_{r/2}^+$. Hence $\partial_i v$ is a solution to $\mathcal{L}u = 0$ in \mathbb{R}_+^{n+1} .

Now, for $x = (x', y) \in \mathbb{R}_+^{n+1}$, let $R = |x|$. We distinguish between two cases: $y > R/16$, and $y < R/16$.

In the first case, set B_R to be the ball of radius R , centered at x . Note then that $B_{R/16} \subseteq \mathbb{R}_+^{n+1}$. Then, from [FKS82, Theorem 2.3.1], Caccioppoli's estimate and the assumption $|v(x)| \leq C|x|^\alpha$,

$$|\partial_i v(x)|^2 \leq \frac{C}{w(B_{R/32})} \int_{B_{R/32}} |\partial_i v|^2 w \leq \frac{C}{w(B_{R/32})} \frac{C}{R^2} \int_{B_{R/16}} |v|^2 w \leq \frac{C}{R^2} \frac{w(B_{R/16})}{w(B_{R/32})} \sup_{B_{R/16}} |v| \leq CR^{2\alpha-2}.$$

In the second case, let B_R be the ball centered at $(x', 0)$ with radius R , and denote $B_R^+ = B_R \cap \mathbb{R}_+^{n+1}$. Then $x \in B_{R/8}^+$, therefore from [FKS82, Theorem 2.4.3] and the boundary Caccioppoli estimate,

$$|\partial_i v(x)|^2 \leq \frac{C}{w(B_{R/8}^+)} \int_{B_{R/8}^+} |\partial_i v|^2 w \leq \frac{C}{w(B_{R/8}^+)} \frac{C}{R^2} \int_{B_{R/4}^+} |v|^2 w \leq \frac{C}{R^2} \frac{w(B_{R/4}^+)}{w(B_{R/8}^+)} \sup_{B_{R/4}^+} |v| \leq CR^{2\alpha-2}.$$

So, in all cases, $|\partial_i v(x)| \leq C|x|^{\alpha-1}$. Letting $R \rightarrow \infty$ and using the maximum principle, we find that $\partial_i v = 0$ for any $i = 1, \dots, n$. Therefore v does not depend on the first n variables, so $v(x', y) = v(y)$. Hence, in \mathbb{R}_+^{n+1} ,

$$0 = -\operatorname{div}(y^\beta \nabla v(y)) = -\partial_y(y^\beta v'(y)) \Rightarrow y^\beta v'(y) = \tilde{c},$$

for some constant \tilde{c} . From [FKS82, Theorem 2.4.6], v is Hölder continuous up to the boundary, therefore for any $y > 0$,

$$v(y) = v(y) - v(0) = \int_0^y v' = \int_0^y \tilde{c}s^{-\beta} ds = \frac{\tilde{c}}{1-\beta}y^{1-\beta},$$

which implies that

$$|\tilde{c}| = (1-\beta)y^{\beta-1}|v(y)| = (1-\beta)y^{\beta-1}|v(0,y)| \leq (1-\beta)y^{\beta-1}y^\alpha = (1-\beta)y^{-\alpha},$$

for any $y > 0$. Letting $y \rightarrow \infty$ we obtain that $\tilde{c} = 0$, hence $v'(y) = 0$ as well, which implies that v is a constant. Since v vanishes on \mathbb{R}^n , this implies that $v \equiv 0$. \square

References

- [AC81] Hans Wilhem Alt and Luis A. Caffarelli. Existence and regularity for a minimum problem with free boundary. *Journal für die reine und angewandte Mathematik*, 325:105–144, 1981.
- [AFP00] Luigi Ambrosio, Nicola Fusco, and Diego Pallara. *Functions of bounded variation and free discontinuity problems*. Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, New York, 2000.
- [All12] Mark Allen. Separation of a lower dimensional free boundary in a two-phase problem. *Mathematical research letters*, 19(5):1055–1074, 2012.
- [BV16] Claudia Bucur and Enrico Valdinoci. *Nonlocal diffusion and applications*, volume 20. Springer, 2016.
- [CN13] Jeff Cheeger and Aaron Naber. Lower bounds on ricci curvature and quantitative behavior of singular sets. *Invent. Math.*, 191(2):321–339, 2013.
- [CRS10a] Luis A. Caffarelli, Jean-Michel Roquejoffre, and Ovidiu Savin. Nonlocal minimal surfaces. *Communications on Pure and Applied Mathematics*, 63(9):1111–1144, 2010.
- [CRS10b] Luis A. Caffarelli, Jean-Michel Roquejoffre, and Yannick Sire. Variational problems with free boundaries for the fractional laplacian. *Journal of the European Mathematical Society*, 12(5):1151–1179, 2010.
- [CS05] Luis A. Caffarelli and Sandro Salsa. *A geometric approach to free boundary problems*, volume 68. American Mathematical Society Providence, RI, 2005.
- [CS07] Luis A. Caffarelli and Luis Silvestre. An extension problem related to the fractional laplacian. *Communications in partial differential equations*, 32(8):1245–1260, 2007.
- [CS14] Xavier Cabré and Yannick Sire. Nonlinear equations for fractional Laplacians, I: Regularity, maximum principles, and Hamiltonian estimates. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 31(1):23–53, 2014.
- [Dal12] Gianni Dal Maso. *An introduction to Γ -convergence*, volume 8. Springer Science & Business Media, 2012.
- [DJ09] Daniela De Silva and David Jerison. A singular energy minimizing free boundary. *J. Reine Angew. Math.*, 635:1–21, 2009.

- [DL76] Georges Duvaut and Jacques-Louis Lions. *Inequalities in mechanics and physics*. Springer-Verlag, Berlin-New York, 1976. Translated from the French by C. W. John, Grundlehren der Mathematischen Wissenschaften, 219.
- [DR12] Daniela De Silva and Jean-Michel Roquejoffre. Regularity in a one-phase free boundary problem for the fractional Laplacian. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 29(3):335–367, 2012.
- [DS93] Guy David and Stephen Semmes. *Analysis of and on uniformly rectifiable sets*, volume 38. American Mathematical Soc., 1993.
- [DS12] Daniela De Silva and Ovidiu Savin. $C^{2,\alpha}$ regularity of flat free boundaries for the thin one-phase problem. *J. Differential Equations*, 253(8):2420–2459, 2012.
- [DS15] Daniela De Silva and Ovidiu Savin. Regularity of lipschitz free boundaries for the thin one-phase problem. *Journal of the European Mathematical Society*, 17(6):1293–1326, 2015.
- [DSFS19] Daniela De Silva, Fausto Ferrari, and Sandro Salsa. Recent progresses on elliptic two-phase free boundary problems. to appear in *Disc. Cont. Dyn. Systems*, 2019.
- [DSS14] Daniela De Silva, Ovidiu Savin, and Yannick Sire. A one-phase problem for the fractional laplacian: regularity of flat free boundaries. *arXiv preprint arXiv:1401.6443*, 2014.
- [DT15] Guy David and Tatiana Toro. Regularity of almost minimizers with free boundary. *Calc. Var. Partial Differential Equations*, 54(1):455–524, 2015.
- [DV17] Serena Dipierro and Enrico Valdinoci. Continuity and density results for a one-phase nonlocal free boundary problem. In *Annales de l’Institut Henri Poincaré (C) Non Linear Analysis*, volume 34, pages 1387–1428. Elsevier, 2017.
- [EE19] Nick Edelen and Max Engelstein. Quantitative stratification for some free-boundary problems. *Trans. Amer. Math. Soc.*, 371(3):2043–2072, 2019.
- [EG15] Lawrence Craig Evans and Ronald F. Gariepy. *Measure theory and fine properties of functions*. CRC press, 2015.
- [Eva98] Lawrence Craig Evans. *Partial differential equations*, volume 19 of *Graduate Studies in Mathematics*. Oxford University Press, 1998.
- [Fed69] Herbert Federer. *Geometric measure theory*. Springer, 1969.
- [FJK82] Eugene Barry Fabes, David Jerison, and Carlos E. Kenig. The wiener test for degenerate elliptic equations. *Ann. Inst. Fourier (Grenoble)*, 32(3):151–182, 1982.
- [FKS82] Eugene Barry Fabes, Carlos E. Kenig, and Raul Paolo Serapioni. The local regularity of solutions of degenerate elliptic equations. *Comm. Partial Differential Equations*, 7(1):77–116, 1982.
- [FV17] Alessio Figalli and Enrico Valdinoci. Regularity and Bernstein-type results for nonlocal minimal surfaces. *J. Reine Angew. Math.*, 729:263–273, 2017.
- [HKM06] Juha Heinonen, Tero Kilpeläinen, and Olli Martio. *Nonlinear potential theory of degenerate elliptic equations*. Courier Corporation, 2006.

- [Jon90] Peter W. Jones. Rectifiable sets and the traveling salesman problem. *Invent. Math.*, 102(1):1–15, 1990.
- [KT03] Carlos E. Kenig and Tatiana Toro. On the free boundary regularity theorem of alt and caffarelli. *Discrete and Continuous Dynamical Systems*, 10:397–422, 2003.
- [NV17] Aaron Naber and Daniele Valtorta. Rectifiable-reifenberg and the regularity of stationary and minimizing harmonic maps. *Annals of Mathematics*, pages 131–227, 2017.
- [Sil12] Luis Silvestre. On the differentiability of the solution to an equation with drift and fractional diffusion. *Indiana University Mathematics Journal*, pages 557–584, 2012.
- [SV12] Ovidiu Savin and Enrico Valdinoci. Γ -convergence for nonlocal phase transitions. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 29(4):479–500, 2012.
- [SV13] Ovidiu Savin and Enrico Valdinoci. Regularity of nonlocal minimal cones in dimension 2. *Calc. Var. Partial Differential Equations*, 48(1-2):33–39, 2013.
- [Vit18] Stefano Vita. *Strong competition systems ruled by anomalous diffusions*. PhD thesis, Università degli Studi di Torino, 2018.
- [Wei98] Georg Sebastian Weiss. Partial regularity for weak solutions of an elliptic free boundary problem. *Communications in partial differential equations*, 23(3-4):439–455, 1998.
- [Wei99] Georg Sebastian Weiss. Partial regularity for a minimum problem with free boundary. *Journal of Geometric Analysis*, 9(2):317–326, 1999.