

This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Alanko, Lauri; Laukkanen, Jari A.; Rottensteiner, Mirva; Rasmus, Salla; Kuha, Tero; Valtonen, Maarit; Kujala, Urho M.

Title: Sports and exercise medicine clinic in public hospital settings : a real-life concept and experiences of the treatment of the first 1151 patients

Year: 2023

Version: Accepted version (Final draft)

Copyright: © Taylor & Francis 2022

Rights: CC BY-NC-ND 4.0

Rights url: <https://creativecommons.org/licenses/by-nc-nd/4.0/>

Please cite the original version:

Alanko, L., Laukkanen, J. A., Rottensteiner, M., Rasmus, S., Kuha, T., Valtonen, M., & Kujala, U. M. (2023). Sports and exercise medicine clinic in public hospital settings : a real-life concept and experiences of the treatment of the first 1151 patients. *Postgraduate Medicine*, 135(3), 283-289. <https://doi.org/10.1080/00325481.2022.2135894>

Sports and exercise medicine clinic in public hospital settings: A real-life concept and experiences of the treatment of the first 1151 patients

Lauri Alanko^{1,2,3}, Jari A. Laukkanen (ORCID ID 0000-0002-3738-1586)^{4,5}, Mirva Rottensteiner (ORCID ID 0000-0003-1988-0952)^{2,3}, Salla Rasmus^{1,3}, Tero Kuha¹, Maarit Valtonen³, Urho M. Kujala (ORCID ID 000-0002-9262-1992)²

¹Central Finland Health Care District, Sports and Exercise Medicine Clinic, Jyväskylä, Finland.

²Faculty of Sport and Health Sciences, University of Jyväskylä, Jyväskylä, Finland

³Finnish Institute of High Performance Sport KIHU, Jyväskylä, Finland

⁴Institute of Clinical Medicine, Department of Medicine, University of Eastern Finland, Kuopio, Finland

⁵Central Finland Health Care District, Department of Medicine, Jyväskylä, Finland

Correspondence to:

Jari A. Laukkanen, Faculty of Sport and Health Sciences, University of Jyväskylä, P.O. Box 35, FI-40014 Jyväskylä, Finland e-mail: jariantero.laukkanen@uef.fi

Words (text only): 3871

Abstract

Objective: Physical exercise has been shown to have a variety of health-promoting effects, including improvements in cardiorespiratory and muscular fitness, symptoms and risk factors such as LDL and HDL cholesterol, triglyceride and glycosylated haemoglobin concentrations. Regular physical exercise may slow down or even reverse the progression of various non-communicable diseases (NCDs). Despite the overwhelming evidence, physical exercise is not comprehensively used as a treatment component either in primary care or in hospital settings.

Methods: The outpatient Sports and Exercise Medicine Clinic (SEMC) is the first specialized clinic in Finland to use physical exercise as a part of the public health care system. Patients needing specialist attention due to NCDs, usually combined with sedentary lifestyles, are referred to the clinic.

Results: The prerequisites for patient referral are the known efficacy of physical exercise intervention in the treatment of disease and the need for sports and exercise medicine expertise. The focus of the clinic is to implement physical activity into daily life with other health-promoting habits such as diet, rest and the reduction of substance use. In addition, SEMC promotes the inclusion of physical exercise in several local treatment guidelines in the hospital district. The advisory treatment protocol of SEMC consists of a baseline evaluation, face-to-face visits with a physician and/or physiotherapist at 3, 6 and 9–12 months, and contacts via phone between hospital visits. Laboratory tests, body composition, walking tests, and measurements of muscle strength and balance are performed at baseline, and body composition and physical tests are repeated after 6 and 9–12 months.

Conclusions: At the core of the treatment is individualization, using motivational interviewing, considering the patients' personal interests and resources, and encouraging the patient to be an active member of our multi-professional team. We reported the first results in the SEMC with future development plans for the clinic.

Keywords: body composition cardiorespiratory fitness, exercise, physical activity, prevention

1. Introduction

A sedentary lifestyle has been shown to increase the prevalence of several chronic diseases and risk factor levels, such as type 2 diabetes, elevated blood pressure, obesity and sleep apnoea (<https://www.acsm.org/education-resources/pronouncements-scientific-communications/official-statements>). Yet, there is strong evidence from randomized controlled trials (RCTs) that physical exercise therapy leads to improved physical fitness and function in patients with chronic diseases (1) and has other health benefits as well (2, 3). Physical exercise training leads to improvements in body composition, including body weight control and the reduction of visceral fat (4,5). There are meta-analyses of RCTs in patients with NCDs summarizing the evidence of other multi-dimensional benefits of physical exercise therapy in the secondary prevention and treatment of chronic diseases (6,7,8). These meta-analyses show that various cardiometabolic risk factor levels are improved in most of the common cardiometabolic diseases, pain is reduced in musculoskeletal diseases, mood (depression and anxiety) and health-related quality of life are improved in various disease categories, and disease-specific 'proxy' indicators of disease progression are improved in diseases such as type 2 diabetes, hypertension, coronary heart disease, heart failure, claudication, chronic obstructive pulmonary disease, rheumatoid arthritis, fibromyalgia, depression, anxiety and schizophrenia (6,7,8).

Despite the overwhelming evidence, the systematic implementation of physical exercise therapy in medical care system has been limited. Healthcare professionals may have been reluctant to recognize physical exercise as a part of medical treatment. Systematic use of physical exercise training has been thought to belong more to the world of sports, and its wider implementation from younger to adult populations is thought to be the mission/duty of organizations other than the healthcare system. There are different types of sports and exercise medicine clinics, but usually they are not linked to public health care, which would give all patients with NCDs the opportunity to access physical exercise therapy, and clinics for specific diseases have only randomly included physical exercise in their treatment protocols. This paper describes the protocol and preliminary results of the recently developed Sports and Exercise Medicine Clinic (SEMC) in the Central Finland Central Hospital (CFCH), discusses experiences and offers suggestions for further development.

2. Methodological development and concept of the sports and exercise medicine clinic

2.1. Patient scope and referral

The SEMC is open to all patients living in the Central Finland Health Care District (total population: 252,676), and they are referred to SEMC by both primary health care and CFCH physicians and also from private health care services. The referral criteria are the evidence-based effect of physical exercise in the treatment of patients' disease, the need for specialist-led physical exercise interventions for patients with

physical exercise disabling NCDs, and patients' own even mild to moderate motivation to implement physical exercise into their everyday lives. A typical patient has several NCDs and a sedentary lifestyle.

2.2. Steps in the clinic development

From the beginning, SEMC has been collaborating actively with the Faculty of Sports and Health Sciences of the University of Jyväskylä. The first step in the development of the SEMC in the CFCH was a questionnaire study offered to all specialist physicians at the hospital. All respondents supported the development of the clinic and indicated that physical inactivity with obesity was the most common challenge needing attention (9).

After piloting the project, the clinic was started in 2016 with a part-time specialist physician in Sports and Exercise Medicine (SEM) and a part-time physician specializing in SEM; later, a part-time physiotherapist was added. Throughout the years, the protocol has been tested and modified in order to improve it. Currently, the personnel consist of a part-time chief physician (specialist in SEM), a full-time physician in training and a full-time physiotherapist.

The ethics committee of the Central Finland Health Care District, Jyväskylä, Finland, has evaluated the reporting of the treatment results of the SEMC patients and gave permission to the study through commenting that there is no need for a specific ethics committee statement as the study reports results of the hospital's regular treatment praxis and only the physician has access to patient data using personal identification codes. The Central Finland Health Care District, Jyväskylä, Finland gave permission to the SEMC study.

2.3 Current concept of the SEMC

The physical exercise intervention protocol usually lasts 9–12 months and consists of a primary visit and follow-ups at 3, 6 and 9–12 (the last one is decided individually) months after the baseline visit. In between visits to the SEMC, patients are contacted by phone.

The baseline visit lasts 60 minutes, during which the patient's overall situation is evaluated. In addition to possible routine disease-specific medical examinations, prior to the first and 6-month visits to the clinic, patients complete both the lifestyle questionnaire (including questions on physical exercise, sitting time, dietary habits, sleep, alcohol consumption, smoking) and the quality-of-life questionnaire RAND-36 (10). The medical conditions are evaluated, as well as the possibilities for increasing physical exercise and developing a healthier lifestyle. Laboratory tests (including an ECG) are collected prior to the first visit to assist in the medical evaluation of the patient's risk factors. By the end of the visit, goals are set and recorded. The goals, which are tailored to each patient individually, include the amount of increase in physical activity/exercise, possible changes in diet and decrease in substance use and screen time. The forms of activity are decided based on the interests of the patient, considering possible medical/economic/social hindrances. The patient helps set these goals and is thus included in the treatment decision-making. The principles of motivational interviewing are applied throughout the process.

The American College of Sport Medicine's (ACSM) recommendation for physical exercise is 2.5 hours of aerobic training and two strength training sessions a week ([Physical Activity Guidelines Resources \(acsm.org\)](#)). At the start of the physical exercise intervention this recommendation is often unrealistic for SEMC patients, as most of them have previously engaged in little or no physical activity. In general, the main goal at the beginning of the intervention is getting any physical exercise at all to become a regular part of a patient's everyday life with later progression of the physical exercise amount. The key in physical exercise prescription was taking into account the health status, patient needs, initial physical fitness level, practical possibilities and motivational preferences of the patient to tailor a progressive physical exercise therapy program with individually adapted physical exercise intensities, frequencies and modes with checking the need of supervision and speed of progression at follow-up visits (see later).

The physician's appointment is followed by two hours of detailed assessment and the evaluation of functional capacity and body composition by an experienced physiotherapist. The tests include a 6-minute walking test (occasionally, a stress-ECG test is done prior to a walking test for patient safety), body composition analysis (bioimpedance, Biospace InBody 770), sit-ups from a chair (with patients over 55 years), squats (with patients 35–55 years), hand-grip strength and one-leg stance. In the case of a specific need, either a medical or psychological or another appointment with the physiotherapist is scheduled. These visits are usually either to treat a musculoskeletal disease (i.e. osteoarthritis, chronic lower back pain, etc.) or to develop a personal exercise and/or strength programme, especially in cases when attending a gym is not possible (due to distance, financial or social issues, etc.).

To further help the patient in the beginning of treatment, SEMC has its own strength-training group. This group is a very low-level entry for those who have no previous experience in weight training and who usually find it very hard to go to an open gym. There are six patients in this group at a time, enabling personal supervision by our physiotherapist. The participants of this strength-training group were provided individualized advice on the training loads and other aspects of their physical exercise therapy from the experienced physiotherapist. Patients can take part in this group six times, after which they are guided to communal and/or private gyms that suit the patient in question. In addition, SEMC collaborates with municipal recreation services, taking into account the restrictions in physical exercising. In the communal services, patients receive personal assistance when starting to exercise, information about public physical exercise facilities/opportunities and even follow-up visits.

During control visits, patients' progress is evaluated. Positive feedback is given even for patients with minor advances and possible problems are assessed. The goals are re-evaluated and, if necessary, re-adjusted accordingly. At 6 months, the effects of intervention are evaluated, and if it is very obvious that no progress is forthcoming or that the goals of the intervention have been achieved, the intervention is terminated. It is not common that after 6 months, the patient has implemented so much exercise into his/her everyday life that the ACSM guidelines have been fulfilled and is mentally ready to continue the physically active lifestyle on her/his own. However, usually, the follow-up ends after 9–12 months, and even if the positive changes are excellent after 6 months, the patient still needs psychological support. Sometimes the measurable progress

begins to happen later in the intervention, and in these cases, the intervention can be prolonged. The decision to end the contacts at SEMC is made together with the patient, not solely by the physician.

Additionally, there are available nutritional therapists both in the hospital and in the primary health care, and patients with a BMI over 35 can be directed for consultation. The overweight and obese patients were given general nutritional recommendations and for selected patients, such as those who needed bariatric surgery, were forwarded to nutritionist for individualized diet prescription. The current SEMC protocol included a joint project with LIKES Research Centre on Physical Activity and Health, in which patients with the lowest scores in the RAND36 survey have the opportunity to meet with a certified psychologist.

3. Experiences and comments

Since the start of autumn 2016 to end-December 2021, the SEMC has treated 1151 patients (**Figure 1**). The patient flow rose in the first years, reaching the patient visit numbers. **Table 1** lists the 20 most common diagnoses of our 2016–2020 patients. Overweight is present in approximately two-thirds of all patients, followed by (co-existing) sleep apnoea, hypertension, non-insulin dependent diabetes, and asthma. The first musculoskeletal disorder in which symptoms are worsened by physical inactivity and attenuated with exercise – osteoarthritis – is ranked sixth. Cardiovascular disease-related diagnoses are less common than expected, as a separate physiotherapy-led rehabilitation protocol for cardiac patients already exists in the hospital district.

In its first five years (from 2016 to 2020) SEMC treated 874 patients, with 457 different diagnoses adding to the total number of 4,158 diagnoses, corresponding to 4.8 diagnoses per patient. Thus, the SEMC patient sample deviates from typical patients in exercise therapy RCTs, who commonly have only one disease. The limitations of the RCTs include that the patients are selected groups according to their disease status, motivation to participate in the trial, non-individualized training programs as well as coverage of the diverse patient population.

Personalized physical exercise therapy has been the core protocol, adjusted to patients' needs, abilities and opportunities. Patients have given positive feedback, and many of them have succeeded in changing their sedentary lifestyle into a physically active one. A rough estimate based on three master's theses produced by SEMC patients (11,12,13) and the personal views of the personnel working in the SEMC is that slightly more than half of the patients succeeded in increasing physical exercise clinically significantly. Among the 106 first patients who often were physically inactive and overweight or obese at baseline were in particular able to increase strength training frequency during the first six months of the intervention (13). Thus, while our real-life experience indicates that this is somewhat lower than what has been reported in RCTs, the patients at SEMC have not been excluded on virtually any criteria such as low motivation, and the volume, intensity and type of exercise has been tailored to severely incapacitated patients as well.

Individuals have different interests regarding types of physical exercises, and our experience is that it is critical to discuss the likes/dislikes of the patient, ask for preferences and be supportive in the choice of physical activities. The most common physical activity is walking/Nordic walking, followed by swimming/water running, gym training, (electric-)cycling and cross-country skiing. But there are patients who participate in dancing and even virtual reality gaming. In short, if a patient likes an activity that is not a health risk, he/she should be encouraged to continue. The starting activity levels vary from none to several hours of weekly physical exercise to several hours but are usually below the national guidelines. Another typical feature in SEMC patients is that there can be some aerobic exercise or strength training at the beginning of the intervention, but very rarely both. The level of effort in aerobic training is low, that is, slow walking with no/little effort. One of the goals during the intervention is to guide the patient into the existing exercise facilities, non-supervised and supervised, instead of organizing separate training programmes. In this way, we try to reach a high cost-benefit ratio and promote long-term lifestyle change.

Furthermore, according to our experiences at SEMC, there are patients who start increasing physical activity during their waiting time for the clinic or are very responsive to the physician's simple advice. These patients have often been physically active during childhood/adolescence but changed later to sedentary lifestyles. These patients often start sports type activities which they have participated in at younger age. On the other end, there are patients who have never been physically active in their entire lives and need more support both psychologically and in starting physical exercise. At the core of personalized physical exercise therapy is that the provision of therapy can and should be intensified or changed during follow-up appointments if the patient has not been able to implement physical exercise into his/her everyday life. In our experience, this kind of treatment protocol may lead to better overall long-term success than that reported in RCTs with strictly standardized programmes, especially with patients with several NCDs and disabilities.

SEMC relies on the physical activity level reported by the patient (through interviews and questionnaires). Most SEMC patients do not have sport watches or activity trackers. Some patients find them a positive challenge, while others consider them more of a threat, causing psychological pressure. On the other hand, an old-fashioned physical exercise diary has proven to be an easy and cost-effective way of monitoring and encouraging daily activities. Different types of technology are needed to meet individual preferences.

RCTs are common method of evaluating the effects of treatment, but they have some serious downsides from the real-life perspective. First, participants for RCTs are usually selected and highly motivated for exercise training, whereas those with the gravest need for increased physical activity are probably the least likely to participate RCTs such as some of the SEMC patients with increased age and multiple NCDs. Reasons for being physically inactive are multitude including such as having several NCDs, social/financial difficulties. Second, in reported RCTs, all intervention group participants have usually followed the same pre-determined intervention, thus neglecting the personal interests and what may or may not be possible for the patient – given the circumstances/health/age and so on. In our experience, the physician/physiotherapist must first agree with the patient on personalized ways of increasing physical activity to motivate the individual. Identifying the type of intervention according to personal motives, hopes, needs and possibilities for

increasing physical activity may lead to a better success rate, cost effectiveness and long-term results in real life compared to a strict one-programme-fits-all principle.

More critical analyses of the effectiveness of SEMC treatment protocol are to come, but we have monitored the effectiveness of the treatment to get feedback for further development. Among other things, patients have been cured of sleep apnoea, have had their type 2 diabetes put into remission, have had their anxiety disorder/depression attenuated, and have increased their aerobic capacity, balance and strength, with or without weight loss. Among the sleep apnoea patients all symptoms were reduced ($p < 0.001$) after six months intervention (11). Among the first 106 patients who had not undergone obesity surgery body composition improved for weight, BMI, and muscle mass (13). Among those who increased physical activity, body composition also improved statistically significantly for visceral fat area ($p = 0.004$) (13). At the beginning of the SEMC development, there was close collaboration with the clinic for pulmonary diseases, and follow-ups show that sleep apnoea patients benefit from physical exercise therapy, also independent of weight reduction (11). As expected, reduction of body weight and fat percentage are associated with the increases in physical activity levels of the patients (12), but it should be noted that the health benefits of regular physical exercising also occur independent of weight reduction in most patient groups.

4. Integration of physical exercise into usual care treatment paths

Due to the wide range of effects of physical exercise on several non-communicable diseases, SEMC has been actively involved in the development of several local treatment guidelines for CFCH. In the treatment path for sleep apnoea patients with several NCDs, for whom the CPAP treatment is not successful/patient does not want CPAP treatment or when the overall situation requires specialist-led intervention, they are offered the opportunity to receive an SEMC physical exercise intervention protocol. In the type 2 diabetes treatment path, the SEMC physical exercise intervention is offered to patients with comorbid disease that is hindering physical exercise. Patients with coronary heart disease, atrial fibrillation and heart failure which are the most common cardiac patient groups have benefitted considerably from individualised exercise training. SEMC has taken care of those cardiac patients who need special exercise training and more support with coexisting disease conditions. Cardiac patients are advised to follow home-based exercise training, taking into account individualized training possibilities at home and surrounding facilities. SEMC patients will be given more detailed and individualized counselling for physical exercise than in usual cardiac rehabilitation. All patients with morbid obesity indicating bariatric surgery are directed to a slightly modified physical exercise intervention. In addition, all mammary cancer patients with post-operative hormonal treatments are offered the opportunity of an SEMC consultation, a single visit, a full protocol, or a personally adjusted one. SEMC is collaborating with primary health care to develop treatment guidelines and protocols to include physical exercise as part of the treatment of all patients with insufficient physical exercise levels.

5. Future considerations

The SEMC protocol has been developed and modified throughout its existence, with the ultimate goal of becoming effective, cost-effective and monitorable and able to be implemented in other health care districts in Finland (and even worldwide). A large meta-analysis suggest that guideline bodies should consider the inclusion of cardiorespiratory fitness in standard risk assessment methods (14). Cardiorespiratory and muscle fitness tests and body composition analyses are validated methods, easy to perform and repeatable. In future it is interesting to analyze how improvements in fitness tests are associated with improvements in health benefits as high fitness levels have been shown to predict good prognosis in observational studies (14,15,16). The addition of questionnaires to the protocol made it easier to follow changes in physical exercise and activity, diet and subjective quality of life. Currently, we are working to standardize the way patient reports are written. We are planning to collect automated feedback from the patients to gain more insight into which parts of the protocol the patients consider most valuable and what changes they would prefer.

During the COVID pandemic some of the baseline physician contacts were organized as teleconsultations with good success as the patients had previously participated in clinical examinations and some laboratory screenings at other clinics. Also, during the pandemic higher number of follow-up contacts were organized as teleconsultations. This knowledge supports the useability of digital approaches.

SEMC is collaborating with the University of Jyväskylä in doing research on the effectiveness of the SEMC protocol. As mentioned above, so far, three master's theses have been written by patients of SEMC. To create a sound platform for future research and a tool to develop the protocol, a detailed real-life database that includes all our patients is under development. This database will be used both to assess the efficacy of the treatment and to evaluate the protocol, thus leading to development in local clinical practices.

In the long term, we would like to increase the SEMC multi-professional team to include new specialists, above all, a full-time psychologist. The responses from the project with the local Research Centre on Physical Activity and Health have been positive, and there would be a great demand for psychological support. In addition to patient work, the psychologist would be a much-needed support for the other staff members to increase their skills in supporting the patient. Another much-needed specialist is a nutritional therapist; the need at the hospital exceeds capacity at the moment. Here, too, the therapist could be consulted by the rest of the team more easily than is now possible. A physiologist would be needed to take at least part of the measurements, to conduct testing with the patients and to develop the methods. This would free up time for the physiotherapist. Something that is very much needed and hoped for by the patients is more strength training groups, which would require more physiologist/physiotherapy capacity. An easy transition from the SEMC training group to public gyms and exercise facilities still needs an improved protocol, a matter to be worked on with the communal exercise professionals. Additionally, it is vital that continuous monitoring of the process and results, combined with a constant drive to evolve SEMC, is further developed.

This will be done using upcoming patient register studies in collaboration with the Faculty of Sports and Health Sciences, University of Jyväskylä.

We will also continue to collaborate with primary health care personnel in Central Finland to actively screen for patient's physical activity levels and take action by delivering personalized physical exercise therapy if the individual's physical activity does not meet the national recommendations. Currently, primary care physicians have had their training periods in SEMC. In some primary health care centres, there are physical exercise instructors working within the health care system. An important up-coming step in the development of collaboration has been the addition of a physical exercise report to the electronic patient record in which both health care and exercise professionals can document and follow the progress of the physical exercise therapy.

6. Conclusions

Personalized physical exercise therapy is not being used yet as a medical treatment to its full potential. Developing and implementing personalized exercise therapy as part of the healthcare system is of major interest, and it may have a worldwide impact, be cost-efficient for the community and effectively improve an individual's health and quality of life.

The SEMC protocol is based on individual physical exercise implementation, developed by both the patient and a multi-professional team. In the SEMC experience, it seems that it is possible to increase a patient's physical activity level, subjective wellbeing and quality of life, even if the patient has no history of physical activity and has a multitude of disabling NCDs. We have gained encouraging experience as we seek to realize the goal of integrating and linking personalized physical exercise closely to patient treatment in both hospital and primary health care settings.

Declaration of funding:

None

Disclosure of financial/other conflicts of interest:

None

7. References

1. Pasanen T, Tolvanen S, Heinonen A, Kujala UM. Exercise therapy for physical function in chronic diseases: a systematic review of meta-analyses of randomized controlled trials. *Br J Sports Med* 2017;51:1459-65.
2. Kujala UM. Evidence of the effects of exercise therapy in the treatment of chronic disease. *Br J Sports Med* 2009;43:550-55.
3. Pedersen BK, Saltin B. Exercise as medicine – evidence for prescribing exercise as therapy in 26 different chronic diseases. *Scand J Med Sci Sports* 2015;25 (Suppl. 3):1-72.
4. Ohkawara K, Tanaka S, Miyachi M, et al. A dose-response relation between aerobic exercise and visceral fat reduction: systematic review of clinical trials. *Int J Obes (Lond)* 2007;31:1786–97.
5. Kim KB, Kim K, Kim C, et al. Effects of exercise on the body composition and lipid profile of individuals with obesity: A systematic review and meta-analysis. *J Obes Metab Syndr* 2019;28:278–94.
6. Kujala UM. Summary of the effects of exercise therapy in non-communicable diseases: Clinically relevant evidence from meta-analyses of randomized controlled trials. *MedRxiv* 2021; doi: <https://doi.org/10.1101/2021.02.11.21251608>.
7. Anderson E, Durstine JL. Physical activity, exercise, and chronic diseases: A brief review. *Sports Med Health Sci* 2019;1:3-10.
8. Winnige P, Vysoky R, Dosbaba F, et al. Cardiac rehabilitation and its essential role in the secondary prevention of cardiovascular diseases. *World J Clin Cases* 2021;9:1761-84.
9. Backman K. Physicians' views on the need of an exercise clinic based in the Central Finland Central Hospital (In Finnish with an abstract in English). Faculty of Sport and Health Sciences, University of Jyväskylä, Master's thesis, 2005. <http://urn.fi/URN:NBN:fi:jyu-200540>
10. Hayes RD, Sherbourne CD, Mazel RM. The RAND 36-item Health Survey 1.0. *Health Economics* 1993;2:217–27.
11. Renkola H-K. The impact of individualized exercise program on overweight sleep apnea patient's symptoms, anthropometry and functional capacity: Results of the sports and exercise medicine outpatient clinic (In Finnish with an abstract in English). Faculty of Sport and Health Sciences, University of Jyväskylä, Master's thesis, 2020. <http://urn.fi/URN:NBN:fi:jyu-202004272902>.
12. Laaksonen M. Effects of exercise treatment on anthropometry, physical functioning and perceived health among obese patients in the Sport and Exercise Medicine outpatient clinic (In Finnish with an abstract in English). Faculty of Sport and Health Sciences, University of Jyväskylä, Master's thesis, 2020. <http://urn.fi/URN:NBN:fi:jyu-202007165342>.

13. Ben-Khalifa J. Effect of exercise therapy on physical function, body composition and increased physical activity directed at the Sports and Exercise Medicine Clinic (In Finnish with an abstract in English). Faculty of Sport and Health Sciences, University of Jyväskylä, Master's thesis, 2021.

<https://jyx.jyu.fi/handle/123456789/76354>.

14. Laukkanen JA, Isiozor NM, Kunutsor SK. Objectively assessed cardiorespiratory fitness and all-cause mortality risk: An updated meta-analysis of 37 cohort studies involving 2,258,029 participants. *Mayo Clin Proc.* 2022;97(6):1054-1073.

15. Laukkanen JA, Kujala U. Low cardiorespiratory fitness is a risk factor for death: Exercise intervention may lower mortality? *J Am Coll Cardiol.* 2018;72(19):2293-2296.

16. Laukkanen JA, Kunutsor SK, Yates T, et al. Prognostic relevance of cardiorespiratory fitness as assessed by submaximal exercise testing for all-cause mortality: A UK Biobank prospective study. *Mayo Clin Proc.* 2020;95(5):867-878.

Table 1. Typical patient groups which have been forwarded to the Sports and Exercise Medicine clinic, with the main goals of treatment and simple follow-up parameters in use or recommended.

Disease	Main goals of the exercise therapy	Easy-to-use follow-up measures*	N†
Obesity, including overweight in children	Reduction of/controlling body weight, prevention of metabolic disturbances, maintaining muscle mass during weight reduction	WT, WM, body composition, muscular strength tests, glucose and lipid risk factors	1011
Sleep apnoea	Reduction of daytime tiredness, improving sleep quality, reduction of weight, improving fitness	WT, WM	637
Hypertension	Reduction of blood pressure levels and weight, treatment of metabolic disturbances	Blood pressure, body composition	384
Type 2 diabetes	Improving glucose metabolism, preventing complications	WM, glucose and lipid levels	228
Asthma	Improvement in exercise capacity and quality of life, reduction of dyspnoea symptoms	WT, clinical exercise test, symptom questionnaire, PEF follow-up, flow-volume spirometry	186
Arthrosis and other degenerative joint diseases	Pain reduction and function improvement	Pain-VAS, WT, muscular strength tests	127
Metabolic syndrome	Improvements in metabolism, reduction of waist circumference, improvement in cardio-metabolic risk factors and physical fitness	WT, WM, blood glucose and lipids, blood pressure	120
Chronic pain in any part of the spine, with or without neural compression, spondylolisthesis, stenosis etc	Pain reduction, function improvement	Pain-VAS, muscular strength tests	103
Hypercholesterolemia	Improvement in lipid profile	blood lipids	94
Neurologic diseases (MS, Parkinson's disease, etc.)	Maintaining or improving mobility, motor control and balance	WT, balance tests	84
Coronary heart disease	Prevention of disease progression, reduction of symptoms	WT, WM, glucose and lipid risk factors	81
Hypothyroidism	Reduction of symptoms	WT, blood test (TSH, T4v)	80
Atrial fibrillation	Control of symptoms, maintaining exercise- and functional capacity	WT, heart rate response during exercise test	76
Depression	Reduction of depression symptoms	Mood questionnaire	48
Tendinopathies, bursitis, tendon overuse injuries	Pain reduction and function improvement	Pain-VAS, WT, muscular strength tests	38
Fibromyalgia	Maintaining physical fitness, mood, and function	WT, symptom questionnaire	32
Acute myocardial infarction, rehabilitation	Improving physical fitness and functional capacity Reducing symptoms/symptom control and possible fears of physical activity	WT, WM, heart rate responses during exercise and assessment of CVD risk factors	31
Cancer	Maintaining or improving physical and mental function, muscle mass and quality of life	WT, muscular strength tests	26
Chronic heart failure	Symptom control, improvement or maintenance of function, prevention of disease progression	WT, heart rate response during exercise, blood tests (pro-BNP)	24
Bi-polar disorder	Reduction of symptoms	Mood questionnaire	21

*Disease-specific quality of life questionnaires are important in the evaluation of treatment results.

†Diagnosis number among 1,535 first patients visiting the clinic (one patient can have several diagnoses)

WT=walking test, WM=measurement of waist circumference, VAS=visual analogue scale.

Figure 1. The number of yearly new patient visits (blue), all yearly patient visits (maroon), yearly phone calls (green), all yearly patient contacts (violet), and cumulative number of treated patients (light blue line) from 2016 through 2021.

