

This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Liu, Jiayin; Zhou, Yuan

Title: A Rademacher type theorem for Hamiltonians $H(x, p)$ and an application to absolute minimizers

Year: 2023

Version: Published version

Copyright: © 2023 the Authors

Rights: CC BY 4.0

Rights url: <https://creativecommons.org/licenses/by/4.0/>

Please cite the original version:

Liu, J., & Zhou, Y. (2023). A Rademacher type theorem for Hamiltonians $H(x, p)$ and an application to absolute minimizers. *Calculus of Variations and Partial Differential Equations*, 62(5), Article 144. <https://doi.org/10.1007/s00526-023-02484-9>



A Rademacher type theorem for Hamiltonians $H(x, p)$ and an application to absolute minimizers

Jiayin Liu^{1,2} · Yuan Zhou³

Received: 21 June 2022 / Accepted: 31 March 2023
© The Author(s) 2023

Abstract

We establish a Rademacher type theorem involving Hamiltonians $H(x, p)$ under very weak conditions in both of Euclidean and Carnot-Carathéodory spaces. In particular, $H(x, p)$ is assumed to be only measurable in the variable x , and to be quasiconvex and lower-semicontinuous in the variable p . Without the lower-semicontinuity in the variable p , we provide a counter example showing the failure of such a Rademacher type theorem. Moreover, by applying such a Rademacher type theorem we build up an existence result of absolute minimizers for the corresponding L^∞ -functional. These improve or extend several known results in the literature.

Mathematics Subject Classification Primary 49J10 · 49J52; Secondary 35F21 · 51K05

Contents

1	Introduction
2	Preliminaries

Communicated by Andrea Mondino.

The first author is supported by the Academy of Finland via the projects: Quantitative rectifiability in Euclidean and non-Euclidean spaces, Grant No. 314172, and Singular integrals, harmonic functions, and boundary regularity in Heisenberg groups, Grant No. 328846. The second author is supported by the National Natural Science Foundation of China (No. 12025102 & No. 11871088) and by the Fundamental Research Funds for the Central Universities. Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

✉ Jiayin Liu
jiayin.mat.liu@jyu.fi

Yuan Zhou
yuan.zhou@bnu.edu.cn

¹ School of Mathematical Science, Beihang University, Changping District Shahe Higher Education Park South Third Street No. 9, Beijing 102206, People's Republic of China

² Department of Mathematics and Statistics, University of Jyväskylä, P.O. Box 35 (MaD), FI-40014 Jyväskylä, Finland

³ School of Mathematical Sciences, Beijing Normal University, Haidian District Xijiekou Waidajie No.19, Beijing 100875, People's Republic of China

3	Proof of Theorem 1.3
4	Proof of Theorem 1.4
5	McShane extensions and minimizers
6	Proof of Theorem 1.6
7	Further discussion
	Appendix A: Rademacher's theorem in Euclidean domains—revisit
	References

1 Introduction

Let $\Omega \subset \mathbb{R}^n$ be a domain (that is, an open and connected subset) of \mathbb{R}^n with $n \geq 2$. We first recall Rademacher's theorem in Euclidean spaces. See Appendix for some of its consequence related to Sobolev and Lipschitz spaces.

Theorem 1.1 *If $u : \Omega \rightarrow \mathbb{R}$ is a Lipschitz function, that is,*

$$|u(x) - u(y)| \leq \lambda |x - y| \quad \forall x, y \in \Omega \text{ for some } 0 \leq \lambda < \infty, \quad (1.1)$$

then, at almost all $x \in \Omega$, u is differentiable and $|\nabla u(x)| = \text{Lipu}(x)$. Here $|\nabla u(x)|$ is the Euclidean length of the derivative $\nabla u(x)$ at x , and $\text{Lipu}(x)$ is the pointwise Lipschitz constant at x defined by

$$\text{Lipu}(x) := \limsup_{y \rightarrow x} \frac{|u(y) - u(x)|}{|y - x|}. \quad (1.2)$$

The above Rademacher's theorem was extended to Carnot-Carathéodory spaces (Ω, X) , where X is a family of smooth vector fields in Ω satisfying the Hörmander condition (See Sect. 2). Denote by Xu the distributional horizontal derivative of $u \in L^1_{\text{loc}}(\Omega)$. Write d_{CC} as the Carnot-Carathéodory distance with respect to X . One then has the following; see [19, 21, 24, 36] and, for the better result in Carnot group and Carnot type vector field, see [40, 42].

Theorem 1.2 *If $u : \Omega \rightarrow \mathbb{R}$ is a Lipschitz function with respect to d_{CC} , that is,*

$$|u(x) - u(y)| \leq \lambda d_{CC}(x, y) \quad \forall x, y \in \Omega \text{ for some } 0 \leq \lambda < \infty, \quad (1.3)$$

then, $Xu \in L^\infty(\Omega, \mathbb{R}^m)$ and, for almost all $x \in \Omega$, the length $|Xu(x)| \leq \lambda$.

Under the additional assumption that X is a Carnot type vector field in Ω , or in particular, (Ω, X) is a domain in some Carnot group, one further has $|Xu(x)| = \text{Lip}_{d_{CC}} u(x)$ for almost all $x \in \Omega$, where $\text{Lip}_{d_{CC}} u(x)$ is defined by (1.2) with $|y - x|$ replaced by $d_{CC}(y, x)$.

This paper aims to build up some Rademacher type theorem involving Hamiltonians $H(x, p)$ in both of Euclidean and Carnot-Carathéodory spaces. Throughout this paper, the following assumptions are always held for $H(x, p)$.

Assumption 1 Suppose that $H : \Omega \times \mathbb{R}^m \rightarrow [0, +\infty)$ is measurable and satisfies

(H1) For each $x \in \Omega$, $H(x, \cdot)$ is quasi-convex, that is,

$$H(x, tp + (1-t)q) \leq \max\{H(x, p), H(x, q)\}, \quad \forall p, q \in \mathbb{R}^m, \forall t \in [0, 1] \text{ and } \forall x \in \Omega.$$

(H2) For each $x \in \Omega$, $H(x, 0) = \min_{p \in \mathbb{R}^m} H(x, p) = 0$.

(H3) It holds that $R_\lambda < \infty$ for all $\lambda \geq 0$, and $\lim_{\lambda \rightarrow \infty} R'_\lambda = \infty$, where and in below,

$$R_\lambda := \sup\{|p| \mid (x, p) \in \Omega \times \mathbb{R}^m, H(x, p) \leq \lambda\}$$

and

$$R'_\lambda := \inf\{|p| \mid (x, p) \in \Omega \times \mathbb{R}^m, H(x, p) \geq \lambda\}.$$

For any $\lambda \geq 0$, we define

$$d_\lambda(x, y) := \sup\{u(y) - u(x) \mid u \in \dot{W}_X^{1,\infty}(\Omega) \text{ with } \|H(\cdot, Xu)\|_{L^\infty(\Omega)} \leq \lambda\} \quad \forall x, y \in \Omega. \quad (1.4)$$

Recall that $\dot{W}_X^{1,\infty}(\Omega)$ denotes the set of all functions $u \in L^\infty(\Omega)$ whose distributional horizontal derivatives $Xu \in L^\infty(\Omega; \mathbb{R}^m)$. It was known that any function $u \in \dot{W}_X^{1,\infty}(\Omega)$ admits a continuous representative \tilde{u} ; see [21, Theorem 1.4] and also Theorem 2.3 and Remark 2.4 below. In this paper, in particular, in (1.4) above, we always identify functions in $\dot{W}_X^{1,\infty}(\Omega)$ as their continuous representatives. We remark that d_λ is not a distance necessarily, but Lemma 2.9 says that d_λ is always a pseudo-distance as defined in Definition 2.8 below.

Given any $\lambda \geq 0$, by the definition (1.4), if $u \in \dot{W}_X^{1,\infty}(U)$ and $\|H(\cdot, Xu)\|_{L^\infty(U)} \leq \lambda$, then $u(y) - u(x) \leq d_\lambda(x, y) \quad \forall x, y \in \Omega$. It is natural to ask whether the converse is true or not. However, such converse is not necessarily true as witnessed by the Hamiltonian

$$\lfloor |p| \rfloor = \max\{t \in \mathbb{N} \mid |p| - t \geq 0\} \quad \forall x \in \Omega, p \in \mathbb{R}^m; \quad (1.5)$$

for details see Remark 1.9 below. The point is that $\lfloor |p| \rfloor$ is not lower-semicontinuous. Below, the converse is shown to be true if $H(x, p)$ is assumed additionally to be lower-semicontinuous in the variable p , that is,

(H0) For almost all $x \in \Omega$, $H(x, p) \leq \liminf_{q \rightarrow p} H(x, q) \quad \forall p \in \mathbb{R}^m$.

Theorem 1.3 *Suppose that H satisfies (H0)-(H3). Given any $\lambda \geq 0$ and any function $u : \Omega \rightarrow \mathbb{R}$, the following are equivalent:*

- (i) $u \in \dot{W}_X^{1,\infty}(\Omega)$ and $\|H(\cdot, Xu)\|_{L^\infty(\Omega)} \leq \lambda$;
- (ii) $u(y) - u(x) \leq d_\lambda(x, y) \quad \forall x, y \in \Omega$;
- (iii) For any $x \in \Omega$, there exists a neighborhood $N(x) \subset \Omega$ such that

$$u(y) - u(z) \leq d_\lambda(z, y) \quad \forall y, z \in N(x).$$

In particular, if $u : \Omega \rightarrow \mathbb{R}$ satisfies any one of (i)-(iii), then

$$\|H(\cdot, Xu)\|_{L^\infty(\Omega)} = \inf\{\lambda \geq 0 \mid \lambda \text{ satisfies (ii)}\} = \inf\{\lambda \geq 0 \mid \lambda \text{ satisfies (iii)}\}. \quad (1.6)$$

Using Theorem 1.3, when $\lambda \geq \lambda_H$ we prove that d_λ has a pseudo-length property, which allows us to get the following. Here and below, define

$$\lambda_H := \inf\{\lambda \geq 0 \mid R'_\lambda > 0\}. \quad (1.7)$$

Since R'_λ defined in (H3) of Assumption 1 is always nonnegative and increasing in $\lambda \geq 0$ and tends to ∞ as $\lambda \rightarrow \infty$, we know that $0 \leq \lambda_H < \infty$, and moreover, $\lambda > \lambda_H$ if and only if $R'_\lambda > 0$.

Theorem 1.4 *Suppose that H satisfies (H0)-(H3). Given any $\lambda \geq \lambda_H$ and any function $u : \Omega \rightarrow \mathbb{R}$, the statement (i) in Theorem 1.3 is equivalent to the following*

- (iv) For any $x \in \Omega$, there exists a neighborhood $N(x) \subset \Omega$ such that

$$u(y) - u(x) \leq d_\lambda(x, y) \quad \forall y \in N(x).$$

In particular, if $u : \Omega \rightarrow \mathbb{R}$ satisfies (iv), then

$$\max\{\lambda_H, \|H(\cdot, Xu)\|_{L^\infty(\Omega)}\} = \min\{\lambda \geq \lambda_H \mid \lambda \text{ satisfies (iv)}\}. \quad (1.8)$$

As a consequence of Theorem 1.3 and Theorem 1.4, we have the following Corollary 1.5. Associated to Hamiltonian $H(x, p)$, we introduce some notion and notations. Denote by $\dot{W}_H^{1,\infty}(\Omega)$ the collection of all $u \in \dot{W}_X^{1,\infty}(\Omega)$ with $\|H(\cdot, Xu)\|_{L^\infty(\Omega)} < \infty$. Denote by $\text{Lip}_H(\Omega, X)$ the class of functions $u : \Omega \rightarrow \mathbb{R}$ satisfying (ii) for some $\lambda > 0$ equipped with (semi-)norms

$$\text{Lip}_H(u, \Omega) = \inf\{\lambda \geq \lambda_H \mid \lambda \text{ satisfies (ii)}\}. \quad (1.9)$$

Denote by $\text{Lip}_H^*(\Omega)$ the collection of all functions u with

$$\text{Lip}_H^*(u, \Omega) = \sup_{x \in \Omega} \text{Lip}_H u(x) < \infty,$$

where we write the pointwise “Lipschitz” constant

$$\text{Lip}_H u(x) = \inf\{\lambda \geq \lambda_H \mid \lambda \text{ satisfies (iv)}\}. \quad (1.10)$$

Thanks to the right continuity of the map $\lambda \in [\lambda_H, \infty) \mapsto d_\lambda(x, y)$ as given in Lemma 4.3, the infima in (1.9) and (1.10) are actually minima.

Corollary 1.5 *Suppose that H satisfies (H0)–(H3) with $\lambda_H = 0$. Then $\dot{W}_H^{1,\infty}(\Omega) = \text{Lip}_H(\Omega) = \text{Lip}_H^*(\Omega)$ and*

$$\|H(\cdot, Xu)\|_{L^\infty(\Omega)} = \text{Lip}_H(u, \Omega) = \text{Lip}_H^*(u, \Omega).$$

Next, we apply the above Rademacher type property to study a minimization problem for L^∞ -functionals corresponding to the above Hamiltonian $H(x, p)$ in both Euclidean and Carathéodory spaces:

$$\mathcal{F}(u, U) := \|H(\cdot, Xu)\|_{L^\infty(U)} \text{ for any } u \in W_{X,\text{loc}}^{1,\infty}(U) \text{ and domain } U \subset \Omega.$$

Aronsson [1–3, 5] in 1960’s initiated the study in this direction via introducing absolute minimizers. A function $u \in W_{X,\text{loc}}^{1,\infty}(U)$ is called an absolute minimizer in U for H and X (write $u \in AM(U; H, X)$ for short) if for any domain $V \Subset U$, it holds that

$$\mathcal{F}(u, V) \leq \mathcal{F}(v, V) \text{ whenever } v \in \dot{W}_X^{1,\infty}(V) \cap C(\bar{V}) \text{ and } u|_{\partial V} = v|_{\partial V}.$$

Here and throughout this paper, for domains A and B , the notation $A \Subset B$ stands for that A is a bounded subdomain of B and its closure $\bar{A} \subset B$.

The existence of absolute minimizers with a given boundary value has been extensively studied. Apart from the pioneering work by Aronsson mentioned above, we refer the readers to [7, 9, 12, 14, 15, 26, 33] and the references therein in the Euclidean setting. For the existence results in Heisenberg groups, Carrot–Carathéodory spaces and general metric spaces with special type of Hamiltonians, we refer the readers to [10, 34, 36–38, 47]. Usually, there are two major approaches to obtain the existence of absolute minimizers. When dealing with C^2 Hamiltonians, one usually transfers the study of absolute minimizers into the study of viscosity solutions of the Aronsson equation (the Euler–Lagrange equation of the L^∞ -functional \mathcal{F}). Thus, to get the existence of absolute minimizers, it suffices to show the existence of the corresponding viscosity solutions. This approach was employed, for instance, in [3, 4, 9, 15, 25, 33, 47]. To study the existence of absolute minimizers for Hamiltonians $H(x, p)$ with less regularity, one efficient way is to use Perron’s method to first get the existence of absolute minimizing Lipschitz extensions (ALME), and then show the equivalence between

ALMEs and absolute minimizers. This idea was adopted in [1, 2, 7, 26, 34, 37, 38]. To see the close connection between ALMEs and absolute minimizers, we refer the readers to [17, 35, 36] and references therein.

Theorem 1.3, Theorem 1.4 and Corollary 1.5 allow us to apply Perron's method directly and then to establish the following existence result of absolute minimizers. This is partially motivated by [12]. However, since we are faced with measurable Hamiltonians, there are several new barriers to be overcome as illustrated at the end of this section.

Theorem 1.6 *Suppose that H satisfies (H0)-(H3) with $\lambda_H = 0$. Given any domain $U \Subset \Omega$ and $g \in \text{Lip}_{d_{CC}}(\partial U)$, there must be a function $u \in AM(U; H, X) \cap \text{Lip}_{d_{CC}}(\bar{U})$ so that $u|_{\partial U} = g$.*

Theorem 1.3, Theorem 1.4, Corollary 1.5 and Theorem 1.6 improve or extend several previous studies in the literature including Theorem 1.1 and Theorem 1.2 above; see Remark 1.7 and Remark 1.8 below.

- Remark 1.7** (i) In Euclidean spaces, that is, $X = \{\frac{\partial}{\partial x_i}\}_{1 \leq i \leq n}$, if $H(x, p) = |p|$, then Corollary 1.5 coincides with Lemma 6, which is a consequence of Theorem 1.1. In Carnot-Carathéodory spaces (Ω, X) , if $H(x, p) = |p|$, then Corollary 1.5 coincides with Lemma 2.7, which is a consequence of Theorem 1.2.
- (ii) In Euclidean spaces, if $H(x, p)$ is lower semi-continuous in $U \times \mathbb{R}^n$, $H(x, \cdot)$ is quasi-convex for each $x \in U$, and satisfies (H2) and (H3), (ii) \Leftrightarrow (i) in Theorem 1.3 was proved in Champion-De Pascale [14] (see also [6, 15] for convex $H(p)$ in Euclidean spaces). The proof in [14] relies on the lower semi-continuity in both of x and p heavily, which allows for approximation $H(x, p)$ via a continuous Hamiltonian in x and p . But such an approach fails under the weaker assumptions (H0)&(H1) here. We refer to Sect. 7 for more details and related further discussions.
- (iii) In both Euclidean and Carrot-Carathéodory spaces, if $H(x, p) = \sqrt{\langle A(x)p, p \rangle}$, where $A(x)$ is a measurable symmetric matrix-valued function satisfying uniform ellipticity, then Theorem 1.3 was established in [36, 38]. The proofs therein rely on the inner product structure, and also do not work here. For measure spaces endowed with strongly regular nonlocal Dirichlet energy forms, where the Hamiltonian is given by the square root of Dirichlet form, we refer to [22, 37, 38, 44] for some corresponding Rademacher type property.
- (iv) Under merely (H0)-(H3), one can not expect that $\text{Lip}_H(x) = H(x, Xu(x))$ almost everywhere. Recall that in Euclidean spaces, there does exist $A(x)$, which satisfying Remark 1.7(iii) above, so that such pointwise property fails for the Hamiltonian $\sqrt{\langle A(x)p, p \rangle}$. For more details see [36, 46].

- Remark 1.8** (i) In Euclidean spaces, that is, $X = \{\frac{\partial}{\partial x_i}\}_{1 \leq i \leq n}$, if $H(x, p)$ is given by Euclidean norm and also any Banach norm, the existence of absolute minimizers was established in Aronsson [1, 2] and Aronsson et al [7]. If $H(x, p)$ is given by $\sqrt{\langle A(x)p, p \rangle}$ with A being as in Remark 1.7 (iii) above, existence of absolute minimizers is given by [36] with the aid of [35]. In a similar way, with the aid of [35], Guo et al [26] also obtained the existence of absolute minimizers if $H(x, p)$ is a measurable function in $\Omega \times \mathbb{R}^n$, and satisfies that $\frac{1}{C} < H(x, p) < C$ for all $x \in \Omega$ and $p \in S^{n-1}$, where $C \geq 1$ is a constant, and that $H(x, \eta p) = |\eta|H(x, p)$ for all $x \in \Omega$, $p \in \mathbb{R}^n$ and $\eta \in \mathbb{R}$.
- (ii) In Euclidean spaces, if $H(x, p)$ is continuous in both variables x, p and quasi-convex in the variable p , with an additional growth assumption in p , Barron et al [9] built up

the existence of absolute minimizers. If $H(x, p)$ is lower semi-continuous in x, p and quasi-convex in p and satisfies (H1)-(H3), Champion-De Pascale [14] established the existence of absolute minimizers with the help of their Rademacher type theorem in Remark 1.7(ii). Recall that the lower semi-continuity of H plays a key role in [14] to obtain the pseudo-length property for d_λ .

- (iii) In Heisenberg groups, if $H(x, p) = \frac{1}{2}|p|^2$ we refer to [10] for the existence of absolute minimizers. In any Carnot group, if $H \in C^2(\Omega \times \mathbb{R}^m)$, $D_{pp}^2 H(x, \cdot)$ is positive definite, and there exists $\alpha \geq 1$ such that

$$H(x, \eta p) = \eta^\alpha H(x, p) \quad \forall x \in \Omega, \eta > 0, p \in \mathbb{R}^n, \quad (1.11)$$

then the existence of absolute minimizers was obtained by Wang [47] via considering viscosity solutions to the corresponding Aronsson equations.

The following remark explains that, without the assumption (H0), Theorem 1.3 does not necessarily hold.

Remark 1.9 The Hamiltonian $H(x, p) = [|p|]$ given in (1.5) satisfies (H1)-(H3) but does not satisfy (H0). Given any $\lambda \in (0, 1)$, we have

$$\begin{aligned} d_\lambda(x, y) &= \sup\{u(y) - u(x) \mid u \in \dot{W}_X^{1,\infty}(\Omega) \text{ with } \|H(\cdot, Xu)\|_{L^\infty(\Omega)} = \| [|Xu|] \|_{L^\infty(\Omega)} \leq \lambda\} \\ &= \sup\{u(y) - u(x) \mid u \in \dot{W}_X^{1,\infty}(\Omega) \text{ with } \| |Xu| \|_{L^\infty(\Omega)} \leq 1\} \\ &= d_{CC}(x, y) \quad \forall x, y \in \Omega. \end{aligned}$$

Fix any $z \in \Omega$ and write $u(x) = d_\lambda(z, x) \quad \forall x \in \Omega$. By the triangle inequality we have

$$u(y) - u(x) = d_\lambda(z, y) - d_\lambda(z, x) \leq d_\lambda(x, y)$$

Recall that, when $X = \{\frac{\partial}{\partial x_j}\}_{1 \leq j \leq n}$ or when X is given by Carnot type Hörmander vector fields, one always has $|Xd_{CC}(z, \cdot)| = 1$ almost everywhere; see [40]. For such X , we conclude that

$$\|H(\cdot, Xu)\|_{L^\infty(\Omega)} = 1 > \lambda.$$

Thus Theorem 1.3 fails.

The following remark explains the reasons why we need $\lambda \geq \lambda_H$ in Theorem 1.4, and why we assume $\lambda_H = 0$ in Theorem 1.6. Note that, in Theorem 1.3 where λ_H maybe not 0, we do get the equivalence among (i), (ii) and (iii) for any $\lambda \geq 0$.

Remark 1.10 (i) To prove (iv) in Theorem 1.4 \Rightarrow (i) in Theorem 1.3, we need a pseudo-length property for d_λ as in Proposition 4.1. When $\lambda > \lambda_H$ (equivalently, $R'_\lambda > 0$), to get such pseudo-length property for d_λ , our proof does use $R'_\lambda > 0$ so to guarantee that the topology induced by $\{d_\lambda(x, \cdot)\}_{x \in \Omega}$ (See Definition 2.8) is the same as the Euclidean topology; see Remark 4.4. When $\lambda = \lambda_H$, we get such pseudo-length property for d_{λ_H} via approximating by $d_{\lambda_H + \epsilon}$ with sufficiently small $\epsilon > 0$.

- (ii) When $\lambda_H > 0$ and $0 \leq \lambda < \lambda_H$, we do not know whether d_λ enjoys such pseudo-length property. We remark that there does exist some Hamiltonian $H(x, p)$ which satisfies the assumptions (H0)-(H3) with $\lambda_H > 0$ (that is, for some $\lambda > 0$, $R'_\lambda = 0$); but for $0 < \lambda < \lambda_H$, the topology induced by $\{d_\lambda(x, \cdot)\}_{x \in \Omega}$ does not coincide with the Euclidean topology; see Remark 2.11 (ii).

- (iii) To get the existence of absolute minimizers, our approach does need Theorem 1.4 and also several properties of d_λ^U , whose proof relies heavily on the pseudo-length property for d_λ and $R'_\lambda > 0$. In Theorem 1.6, we assume $\lambda_H = 0$ so that we can work with all Lipschitz boundary g so to get existence of absolute minimizer.

In the case $\lambda_H > 0$, our approach will give the existence of absolute minimizer when the boundary $g : \partial U \rightarrow \mathbb{R}$ satisfies $\mu(g, \partial U) > \lambda_H$, but does not work when $\mu(g, \partial U) \leq \lambda_H$. Here $\mu(g, \partial U)$ is the infimum of λ so that $g(y) - d(x) \leq d_\lambda^U(x, y)$ for all $x, y \in \partial U$.

The paper is organized as below, where we also clarify the ideas and main novelties to prove Theorem 1.3, Theorem 1.4 and also Theorem 1.6. We emphasize that in our results from Sects 2, 3, 4, 5 and 6, X is a fixed smooth vector field in a domain Ω and satisfies the Hörmander condition; and that the Hamiltonian $H(x, p)$ always enjoys (H0)-(H3). In all results from Sects. 5 and 6, we further assume $\lambda_H = 0$.

In Sect. refsp1, we state several facts about the analysis and geometry in Carnot-Carathéodory spaces employed in the proof.

In Sect. 3, we prove (i) \Leftrightarrow (ii) \Leftrightarrow (iii) in Theorem 1.3. Since (i) \Rightarrow (ii) follows from the definition and that (ii) \Rightarrow (iii) is obvious, it suffices to prove (iii) \Rightarrow (i). To this end, we borrow some ideas from [22, 36, 44, 45], which were designed for nonlocal Dirichlet energy forms originally.

The key is that, by employing assumptions (H0), (H1) and Mazur's theorem, we are able to prove that if $v_j \in \dot{W}_X^{1,\infty}(\Omega)$ with $\|H(\cdot, Xv_j)\|_{L^\infty(\Omega)} \leq \lambda$ and $v_j \rightarrow v$ in $C^0(\Omega)$ as $j \rightarrow \infty$, then $v \in \dot{W}_X^{1,\infty}(\Omega)$ with $\|H(\cdot, Xv)\|_{L^\infty(\Omega)} \leq \lambda$; see Lemma 3.1 for details. Thanks to this, choosing a suitable sequence of approximation functions via the definition of d_λ , we then show that

$$d_\lambda(x, \cdot) \in \dot{W}_X^{1,\infty}(\Omega) \text{ and } \|H(\cdot, Xd_\lambda(x, \cdot))\|_{L^\infty(\Omega)} \leq \lambda \text{ for all } \lambda > 0 \text{ and } x \in \Omega.$$

See Lemma 3.3. Given any u satisfying (iii), we construct approximation functions u_j from d_λ and use Lemma 3.4 to show $H(x, Du_j) \leq \lambda$. That is (i) holds.

In Sect. 4, we prove (iii) in Theorem 1.3 \Leftrightarrow (iv) in Theorem 1.4. Since (iii) \Rightarrow (iv) is obvious, it suffices to show (iv) \Rightarrow (iii).

This follows from the pseudo-length property of pseudo metric d_λ in Proposition 4.1. To get such a length property we find some special functions which fulfill the assumption Theorem 1.3(iii), and hence we show that the pseudo metric d_λ has a pseudo-length property.

In Sect. 5, we introduce McShane extensions and minimizers, and then gather several properties of them and pseudo-distance in Lemma 5.2 to Lemma 5.9, which are required in Sect. 6. These properties also have their own interests.

Given any domain $U \Subset \Omega$, via the intrinsic distance d_λ^U induced from U , we introduce McShane extensions $\mathcal{S}_{g;V}^\pm$ of any $g \in \text{Lip}_{d_{CC}}(\partial U)$ in U . There are several reasons to use d_λ^U other than d_λ , for example, d_λ^U has the pseudo-length property in U but the restriction of d_λ may not have; moreover, Theorem 1.3, and Theorem 1.4 holds if (Ω, d_λ) therein is replaced by (U, d_λ^U) , but not necessarily hold if (Ω, d_λ) therein is replaced by (U, d_λ) . However, the use of d_λ^U causes several difficulties. For example, d_λ^U may be infinity when extended to \bar{U} . This makes it quite implicit to see the continuity of McShane extensions around ∂U from the definition. In Lemma 5.6, we get such continuity by analyzing the behaviour of d_λ^U near ∂U . Moreover, as required in Sect. 6, we have to study the relations between d_λ^U and d_λ^V for subdomains V of U in Lemma 5.3 and Lemma 5.4.

In Sect. 6, we prove Theorem 1.6 in a constructive way by using above Rademacher type property and Perron's approach, where we borrow some ideas from [9, 12, 14].

The proof consists of crucial Lemma 6.2, Lemma 6.4 and Proposition 6.5. Lemma 6.2 says that McShane extensions $\mathcal{S}_{g;U}^\pm$ in U of function g in ∂U are local super/sub minimizers in U . Since $\mathcal{S}_{g;U}^\pm$ are the maximum/minimum minimizers, the proof of Lemma 6.2 is reduced to showing that for any subdomain $V \subset U$, the McShane extensions $\mathcal{S}_{h^\pm,V}^\pm$ in V with boundary $h^\pm = \mathcal{S}_{g;U}^\pm|_{\partial V}$ satisfy

$$\mathcal{S}_{h^\pm,V}^\pm(y) - \mathcal{S}_{h^\pm,V}^\pm(x) \leq d_\lambda^U(x, y) \quad \text{for all } x, y \in \overline{V};$$

see Lemma 5.8 and Lemma 5.9 and the proof of Lemma 6.2. However, since Lemma 5.6 only gives

$$\mathcal{S}_{h^\pm,V}^\pm(y) - \mathcal{S}_{h^\pm,V}^\pm(x) \leq d_\lambda^V(x, y) \quad \text{for all } x, y \in \overline{V},$$

we must improve $d_\lambda^V(x, y)$ here to the smaller one $d_\lambda^U(x, y)$. To this end, we show $d_\lambda^V = d_\lambda^U$ locally in Lemma 5.3, and also use the pseudo-length property of d_λ^U heavily. Lemma 6.4 says that a function which is both of local superminimizers and subminimizer must be an absolute minimizer. To get the required local minimizing property, we use McShane extensions to construct approximation functions and also need the fact $d_\lambda^V = d_\lambda^{V \setminus \{x_i\}_{1 \leq i \leq m}}$ in $\overline{V} \times \overline{V}$ as in Lemma 5.4. Proposition 6.5 says that the supremum of local subminimizers are absolute minimizers. Due to Lemma 6.4, it suffices to prove the local super/sub minimizing property of such a supremum. We do prove this via using Lemma 6.2 and Lemma 5.9 repeatedly and a contradiction argument.

In Sect. 7, we aim at explaining some obstacles in using previous approach to establish the Rademacher type theorem and the existence of absolute minimizers. Indeed, in the literature to study Hamiltonians with better regularity or homogeneity, for instance, [14, 26], another intrinsic distance \bar{d}_λ is more common used in those proof which is hard to fit our setting.

In Appendix, we revisit the Rademacher's theorem in the Euclidean space to show that Theorem 1.3 and Corollary 1.5 are indeed an extension of the Rademacher's theorem.

2 Preliminaries

In this section, we introduce the background and some known results related to Carnot-Carathéodory spaces employed in the proof.

Let $X := \{X_1, \dots, X_m\}$ for some $m \leq n$ be a family of smooth vector fields in Ω which satisfies the Hörmander condition, that is, there is a step $k \geq 1$ such that, at each point, $\{X_i\}_{i=1}^m$ and all their commutators up to at most order k generate the whole \mathbb{R}^n . Then for each $i = 1, \dots, m$, X_i can be written as

$$X_i = \sum_{l=1}^n b_{il} \frac{\partial}{\partial x_l} \quad \text{in } \Omega$$

with $b_{il} \in C^\infty(\Omega)$ for all $i = 1, \dots, m$ and $l = 1, \dots, n$.

Define the Carnot-Carathéodory distance corresponding to X by

$$d_{CC}(x, y) := \inf\{\ell_{d_{CC}}(\gamma) \mid \gamma \in \mathcal{ACH}(0, 1; x, y; \Omega)\} \quad (2.1)$$

Here and below, we write $\gamma \in \mathcal{ACH}(0, 1; x, y; \Omega)$ if $\gamma: [0, 1] \rightarrow \Omega$ is absolutely continuous, $\gamma(0) = x$, $\gamma(1) = y$, and there exists measurable functions $c_i: [0, 1] \rightarrow \mathbb{R}$ with $1 \leq i \leq m$

such that $\dot{\gamma}(t) = \sum_{i=1}^m c_i(t) X_i(\gamma(t))$ whenever $\dot{\gamma}(t)$ exists. The length of γ is

$$\ell_{d_{CC}}(\gamma) := \int_0^1 |\dot{\gamma}(t)| dt = \int_0^1 \sqrt{\sum_{i=1}^m c_i^2(t)} dt.$$

In the Euclidean case, we have the following remark.

Remark 2.1 In Euclidean case, that is, $X = \{\frac{\partial}{\partial x_i}\}_{1 \leq i \leq n}$, one has d_{CC} coincides with the intrinsic distance d_E^Ω as given in (A.2). In particular, $d_{CC}(x, y) = d_E^\Omega(x, y)$ for all $x, y \in \Omega$ with $|x - y| < \text{dist}(x, \partial\Omega)$. When Ω is convex, one further have and $d_{CC}(x, y) = |x - y|$ for all $x, y \in \Omega$; however, when Ω is not convex, this is not necessarily true. See Lemma A.4 in Appendix for more details.

Since X is a Hörmander vector field in Ω , for any compact set $K \subset \Omega$, there exists a constant $C(K) \geq 1$ such that

$$C(K)^{-1} |x - y| \leq d_{CC}(x, y) \leq C(K) |x - y|^{\frac{1}{k}} \quad \forall x, y \in K,$$

see for example [41] and [27, Chapter 11]. This shows that the topology induces by (Ω, d_{CC}) is exactly the Euclidean topology.

Given a function $u \in L_{\text{loc}}^1(\Omega)$, its distributional derivative along X_i is defined by the identity

$$\langle X_i u, \phi \rangle = \int_{\Omega} u X_i^* \phi dx \quad \text{for all } \phi \in C_0^\infty(\Omega),$$

where $X_i^* = -\sum_{l=1}^n \frac{\partial}{\partial x_l} (b_{il} \cdot)$ denotes the formal adjoint of X_i . Write $X^* = (X_1^*, \dots, X_m^*)$. We call $Xu := (X_1 u, \dots, X_m u)$ the horizontal distributional derivative for $u \in L_{\text{loc}}^1(\Omega)$ and the norm $|Xu|$ is defined by

$$|Xu| = \sqrt{\sum_{i=1}^m |X_i u|^2}.$$

For $1 \leq p \leq \infty$, denote by $\dot{W}_X^{1,p}(\Omega)$ the p -th integrable horizontal Sobolev space, that is, the collection of all functions $u \in L_{\text{loc}}^1(\Omega)$ with its distributional derivative $Xu \in L^p(\Omega)$.

Equip $\dot{W}_X^{1,p}(\Omega)$ with the semi-norm $\|u\|_{\dot{W}_X^{1,p}(\Omega)} = \|Xu\|_{L^p(\Omega)}$.

The following was proved in [23, Lemma 3.5 (II)].

Lemma 2.2 If $u \in \dot{W}_X^{1,p}(U)$ with $1 \leq p < \infty$ and $U \Subset \Omega$, then $u^+ = \max\{u, 0\} \in \dot{W}_X^{1,p}(U)$ with $Xu^+ = (Xu)\chi_{\{x \in U | u > 0\}}$ almost everywhere.

We recall the following imbedding of horizontal Sobolev spaces from [24, Theorem 1.4]. For any set $U \subset \Omega$, the Lipschitz class $\text{Lip}_{d_{CC}}(U)$ is the collection of all functions $u : U \rightarrow \mathbb{R}$ with its seminorm

$$\text{Lip}_{d_{CC}}(u, U) := \sup_{x \neq y, x, y \in U} \frac{|u(x) - u(y)|}{d_{CC}(x, y)} < \infty.$$

Theorem 2.3 For any subdomain $U \Subset \Omega$, if $u \in \dot{W}_X^{1,\infty}(U)$, then there is a continuous function $\tilde{u} \in \text{Lip}_{d_{CC}}(U)$ with $\tilde{u} = u$ almost everywhere and

$$\text{Lip}_{d_{CC}}(\tilde{u}, U) \leq C(U, \Omega) [\|u\|_{L^\infty(U)} + \|u\|_{\dot{W}_X^{1,\infty}(U)}].$$

Remark 2.4 For any $u \in \dot{W}_X^{1,\infty}(U)$, we call above \tilde{u} given in Theorem 2.3 as the continuous representative of u . Up to considering \tilde{u} , in this paper we always assume that u itself is continuous.

We have the following dual formula of d_{CC} .

Lemma 2.5 For any $x, y \in \Omega$, we have

$$d_{CC}(x, y) = \sup\{u(y) - u(x) \mid u \in \dot{W}_X^{1,\infty}(\Omega) \text{ with } \|Xu\|_{L^\infty(\Omega)} \leq 1\}. \quad (2.2)$$

To prove this we need the following bound for the norm of horizontal derivative of smooth approximation of functions in $\dot{W}_X^{1,\infty}(\Omega)$, see for example [27, Proposition 11.10]. Denote by $\{\eta_\epsilon\}_{\epsilon \in (0,1)}$ the standard smooth mollifier, that is, $\eta_\epsilon(x) = \epsilon^{-n} \eta(\frac{x}{\epsilon}) \quad \forall x \in \mathbb{R}^n$, where $\eta \in C^\infty(\mathbb{R}^n)$ is supported in unit ball of \mathbb{R}^n (with Euclidean distance), $\eta \geq 0$ and $\int_{\mathbb{R}^n} \eta dx = 1$.

Proposition 2.6 Given any compact set $K \subset \Omega$, there is $\epsilon_K \in (0, 1)$ such that for any $\epsilon < \epsilon_K$ and $u \in \dot{W}_X^{1,\infty}(\Omega)$ one has

$$|X(u * \eta_\epsilon)(x)| \leq \|Xu\|_{L^\infty(\Omega)} + A_\epsilon(u) \quad \forall x \in K, \quad (2.3)$$

where $A_\epsilon(u) \geq 0$ and $\lim_{\epsilon \rightarrow 0} A_\epsilon(u) \rightarrow 0$ in K .

Proof of Lemma 2.5 Recall that it was shown by [32, Proposition 3.1] that

$$d_{CC}(x, y) = \sup\{u(y) - u(x) \mid u \in C^\infty(\Omega) \text{ with } \|Xu\|_{L^\infty(\Omega)} \leq 1\} \quad \forall x, y \in \Omega. \quad (2.4)$$

It then suffices to show that for any $u \in \dot{W}_X^{1,\infty}(\Omega)$ with $\|Xu\|_{L^\infty(\Omega)} \leq 1$, we have

$$u(y) - u(x) \leq d_{CC}(x, y) \quad \forall x, y \in \Omega.$$

Note that u is assumed to be continuous as in Remark 2.4.

To this end, given any $x, y \in \Omega$, for any $\epsilon > 0$ there exists a curve $\gamma_\epsilon \subset \mathcal{ACH}(0, 1; x, y; \Omega)$ such that $\ell_{d_{CC}}(\gamma_\epsilon) \leq (1 + \epsilon)d_{CC}(x, y)$. We can find a domain $U \Subset \Omega$ such that $\gamma_\epsilon \subset U$. It is standard that $u * \eta_t \rightarrow u$ uniformly in \overline{U} and hence

$$u(y) - u(x) = \lim_{t \rightarrow 0} [u * \eta_t(y) - u * \eta_t(x)].$$

Next, by Proposition 2.6, for $0 < t < t_{\overline{U}}$ one has

$$|X(u * \eta_t)(z)| \leq \|Xu\|_{L^\infty(\Omega)} + A_t u(z) \quad \forall z \in \overline{U},$$

and moreover, $A_t u(z) \rightarrow 0$ uniformly in \overline{U} as $t \rightarrow 0$. Obviously, we can find $t_{\epsilon, \overline{U}} < t_{\overline{U}}$ such that for any $0 < t < t_{\epsilon, \overline{U}}$, we have $A_t u(x) \leq \epsilon$ and hence, by $\|Xu\|_{L^\infty(\Omega)} \leq 1$, $|X(u * \eta_t)(z)| \leq 1 + \epsilon$, for all $z \in \overline{U}$. Therefore

$$\begin{aligned} u * \eta_t(y) - u * \eta_t(x) &= \int_0^1 [(u * \eta_t) \circ \gamma_t]'(s) ds \\ &= \int_0^1 X(u * \eta_t)(\gamma_t(s)) \cdot \dot{\gamma}_t(s) ds \\ &\leq (1 + \epsilon) \ell_{d_{CC}}(\gamma_\epsilon) \\ &\leq (1 + \epsilon)(1 + \epsilon) d_{CC}(x, y). \end{aligned}$$

Sending $t \rightarrow 0$ and $\epsilon \rightarrow 0$, one concludes $u(y) - u(x) \leq d_{CC}(x, y)$ as desired. \square

As a consequence of Rademacher type theorem (that is, Theorem 1.2), we have the following, which is an analogue of Lemma 6. Denote by $\text{Lip}_{d_{CC}}^*(\Omega)$ the collection of all functions u in Ω

with

$$\text{Lip}_{d_{CC}}^*(u, \Omega) := \sup_{x \in \Omega} \text{Lip}_{d_{CC}} u(x) < \infty. \quad (2.5)$$

Lemma 2.7 We have $\dot{W}_X^{1,\infty}(\Omega) = \text{Lip}_{d_{CC}}(\Omega) = \text{Lip}_{d_{CC}}^*(u, \Omega)$ with

$$\|Xu\|_{L^\infty(\Omega)} = \text{Lip}_{d_{CC}}(u, \Omega) = \text{Lip}_{d_{CC}}^*(u, \Omega) \quad (2.6)$$

Proof First, we show $\text{Lip}_{d_{CC}}(\Omega) = \text{Lip}_{d_{CC}}^*(\Omega)$ and $\text{Lip}_{d_{CC}}(u, \Omega) = \text{Lip}_{d_{CC}}^*(u, \Omega)$. Notice that $\text{Lip}_{d_{CC}}(u, \Omega) \subset \text{Lip}_{d_{CC}}^*(u, \Omega)$ and $\text{Lip}_{d_{CC}}^*(u, \Omega) \leq \text{Lip}_{d_{CC}}(u, \Omega)$ are obvious. We prove

$$\text{Lip}_{d_{CC}}^*(u, \Omega) \subset \text{Lip}_{d_{CC}}(u, \Omega) \text{ and } \text{Lip}_{d_{CC}}(u, \Omega) \leq \text{Lip}_{d_{CC}}^*(u, \Omega).$$

Let $u \in \text{Lip}_{d_{CC}}^*(u, \Omega)$. Given any $x, y \in \Omega$, and $\gamma \in \mathcal{ACH}(0, 1; x, y; \Omega)$, parameterise γ such that $|\dot{\gamma}(t)| = \ell_{d_{CC}}(\gamma)$ for almost every $t \in [0, 1]$. Since

$$A_{x,y} := \sup_{t \in [0,1]} \text{Lip} u(\gamma(t)) < \infty,$$

for each $t \in [0, 1]$ we can find $r_t > 0$ such that

$$|u(\gamma(s)) - u(\gamma(t))| \leq A_{x,y} |\gamma(s) - \gamma(t)| = A_{x,y} \ell_{d_{CC}}(\gamma) |s - t|$$

whenever $|s - t| \leq r_t$ and $s \in [0, 1]$.

Since $[0, 1] \subset \cup_{t \in [0,1]} (t - r_t, t + r_t)$, we can find an increasing sequence $t_i \in [0, 1]$ with $t_0 = 0$ and $t_N = 1$ such that

$$[0, 1] \subset \cup_{i=1}^N (t_i - \frac{1}{2}r_{t_i}, t_i + \frac{1}{2}r_{t_i}).$$

Write $x_i = \gamma(t_i)$ for $i = 0, \dots, N$. We have

$$\begin{aligned} |u(x) - u(y)| &= \left| \sum_{i=0}^{N-1} [u(x_i) - u(x_{i+1})] \right| \\ &\leq \sum_{i=0}^{N-1} |u(x_i) - u(x_{i+1})| \\ &\leq A_{x,y} \ell_{d_{CC}}(\gamma) \sum_{i=0}^{N-1} |t_i - t_{i+1}| \\ &= A_{x,y} \ell_{d_{CC}}(\gamma). \end{aligned}$$

Noticing that $A_{x,y} \leq \text{Lip}^*(u, \Omega) < \infty$ for all $x, y \in \Omega$, we deduce that

$$|u(y) - u(x)| \leq \text{Lip}_{d_{CC}}^*(u, \Omega) \ell_{d_{CC}}(\gamma) \quad \forall x, y \in \Omega. \quad (2.7)$$

For any $\epsilon > 0$, recalling the definition of d_{CC} in (2.1), there exists $\{\gamma_\epsilon\}_{\epsilon>0} \subset \mathcal{ACH}(0, 1; x, y; \Omega)$ such that

$$\ell_{d_{CC}}(\gamma_\epsilon) \leq (1 + \epsilon) d_{CC}(x, y) \quad \forall x, y \in \Omega. \quad (2.8)$$

Combining (2.7) and (2.8), we have

$$\frac{|u(y) - u(x)|}{d_{CC}(x, y)} \leq \lim_{\epsilon \rightarrow 0} \frac{|u(y) - u(x)|}{(1 + \epsilon)d_{CC}(x, y)} \leq \text{Lip}_{d_{CC}}^*(u, \Omega) \quad \forall x, y \in \Omega.$$

Taking supremum among all $x, y \in \Omega$ in the above inequality, we deduce that $u \in \text{Lip}_{d_{CC}}(u, \Omega)$ and $\text{Lip}_{d_{CC}}(u, \Omega) \leq \text{Lip}_{d_{CC}}^*(u, \Omega)$. Hence the second equality in (2.6) holds.

Next, we show $\dot{W}_X^{1,\infty}(\Omega) = \text{Lip}_{d_{CC}}(\Omega)$ and $\text{Lip}_{d_{CC}}(u, \Omega) = \|Xu\|_{L^\infty(\Omega)}$. By Theorem 1.2, we know $\text{Lip}_{d_{CC}}(\Omega) \subset \dot{W}_X^{1,\infty}(\Omega)$ and $\|Xu\|_{L^\infty(\Omega)} \leq \text{Lip}_{d_{CC}}(u, \Omega)$.

To see $\dot{W}_X^{1,\infty}(\Omega) \subset \text{Lip}_{d_{CC}}(\Omega)$ and $\text{Lip}_{d_{CC}}(u, \Omega) \leq \|Xu\|_{L^\infty(\Omega)}$, let $u \in \dot{W}_X^{1,\infty}(\Omega)$. Then $\|Xu\|_{L^\infty(\Omega)} =: \lambda < \infty$. If $\lambda > 0$, then $\lambda^{-1}u \in \dot{W}_X^{1,\infty}(\Omega)$ and $\|X(\lambda^{-1}u)\|_{L^\infty(\Omega)} = 1$. Hence $\lambda^{-1}u$ could be the test function in (2.2), which implies

$$\lambda^{-1}u(y) - \lambda^{-1}u(x) \leq d_{CC}(x, y) \quad \forall x, y \in \Omega,$$

or equivalently,

$$\frac{|u(y) - u(x)|}{\|Xu\|_{L^\infty(\Omega)}} \leq d_{CC}(x, y) \quad \forall x, y \in \Omega.$$

Therefore, $u \in \text{Lip}_{d_{CC}}(\Omega)$ and $\text{Lip}_{d_{CC}}(u, \Omega) \leq \|Xu\|_{L^\infty(\Omega)}$. If $\lambda = 0$, then similar as the above discussion, we have for any $\lambda' > 0$

$$\frac{|u(y) - u(x)|}{\lambda'} \leq d_{CC}(x, y) \quad \forall x, y \in \Omega.$$

Therefore, $u \in \text{Lip}_{d_{CC}}(\Omega)$ and $\text{Lip}_{d_{CC}}(u, \Omega) \leq \lambda'$ for any $\lambda' > 0$. Hence $\text{Lip}_{d_{CC}}(u, \Omega) = 0 = \|Xu\|_{L^\infty(\Omega)}$ and we complete the proof. \square

Next, we recall some concepts from metric geometry. First we recall the notion of pseudo-distance.

Definition 2.8 We say that ρ is a pseudo-distance in a set $\Omega \subset \mathbb{R}^n$ if ρ is a function in $\Omega \times \Omega$ such that

- (i) $\rho(x, x) = 0$ for all $x \in \Omega$ and $\rho(x, y) \geq 0$ for all $x, y \in \Omega$;
- (ii) $\rho(x, z) \leq \rho(x, y) + \rho(y, z)$ for all $x, y, z \in \Omega$.

We call (Ω, ρ) as a pseudo-metric space. The topology induced by $\{\rho(x, \cdot)\}_{x \in \Omega}$ (resp. $\{\rho(\cdot, x)\}_{x \in \Omega}$) is the weakest topology on Ω such that $\rho(x, \cdot)$ (resp. $\rho(\cdot, x)$) is continuous for all $x \in \Omega$.

We remark that since the above pseudo-distance ρ may not have symmetry, the topology induced by $\{\rho(x, \cdot)\}_{x \in \Omega}$ in Ω may be different from that induced by $\{\rho(\cdot, x)\}_{x \in \Omega}$.

Suppose that $H(x, p)$ is an Hamiltonian in Ω satisfying assumptions (H0)-(H3). Let $\{d_\lambda\}_{\lambda \geq 0}$ be defined as in (1.4). Thanks to the convention in Remark 2.4, one has

$$d_\lambda(x, y) := \sup\{u(y) - u(x) \mid u \in \dot{W}_X^{1,\infty}(\Omega) \text{ with } \|H(\cdot, Xu)\|_{L^\infty(\Omega)} \leq \lambda\} \quad \forall x, y \in \Omega. \quad (2.9)$$

The following properties holds for d_λ .

Lemma 2.9 *The following holds.*

- (i) For any $\lambda \geq 0$, d_λ is a pseudo distance on Ω .

(ii) For any $\lambda \geq 0$,

$$R'_\lambda d_{CC}(x, y) \leq d_\lambda(x, y) \leq R_\lambda d_{CC}(x, y) < \infty \quad \forall x, y \in \Omega. \quad (2.10)$$

(iii) If $H(x, p) = H(x, -p)$ for all $p \in \mathbb{R}^m$ and almost all $x \in \Omega$, then $d_\lambda(x, y) = d_\lambda(y, x)$ for all $x, y \in \Omega$.

Proof To see Lemma 2.9 (i), by choosing constant functions as test functions in (2.9), one has $\rho(x, y) \geq 0$ for all $x, y \in \Omega$. Obviously, one has $d_\lambda(x, x) = 0$ for all $x \in \Omega$. Besides,

$$\begin{aligned} d_\lambda(x, z) &= \sup\{u(z) - u(x) : u \in \dot{W}_X^{1,\infty}(\Omega), \|H(\cdot, Xu)\|_{L^\infty(\Omega)} \leq \lambda\} \\ &\leq \sup\{u(y) - u(x) : u \in \dot{W}_X^{1,\infty}(\Omega), \|H(\cdot, Xu)\|_{L^\infty(\Omega)} \leq \lambda\} \\ &\quad + \sup\{u(z) - u(y) : u \in \dot{W}_X^{1,\infty}(\Omega), \|H(\cdot, Xu)\|_{L^\infty(\Omega)} \leq \lambda\} \\ &= d_\lambda(x, y) + d_\lambda(y, z). \end{aligned}$$

By Definition 2.8, d_λ is a pseudo distance.

To see Lemma 2.9 (ii), by (H3), we have

$$\begin{aligned} \{u \in \dot{W}_X^{1,\infty}(\Omega) \mid \|Xu\|_{L^\infty(\Omega)} \leq R'_\lambda\} &\subset \{u \in \dot{W}_X^{1,\infty}(\Omega) \mid \|H(x, Xu)\|_{L^\infty(\Omega)} \leq \lambda\} \\ &\subset \{u \in \dot{W}_X^{1,\infty}(\Omega) \mid \|Xu\|_{L^\infty(\Omega)} \leq R_\lambda\}. \end{aligned}$$

From this and the definitions of d_{CC} in (2.2) and d_λ in (2.9), we deduce (2.10) as desired.

Finally we show Lemma 2.9 (iii), since $H(x, p) = H(x, -p)$ for all $p \in \mathbb{R}^m$ and almost all $x \in \Omega$, then

$$\|H(x, Xu)\|_{L^\infty(\Omega)} = \|H(x, X(-u))\|_{L^\infty(\Omega)} \quad \text{for all } u \in \dot{W}_X^{1,\infty}(\Omega).$$

Hence for any $x, y \in \Omega$, u can be a test function for $d_\lambda(x, y)$ in the right hand side of (2.9) if and only if $-u$ can be a test function for $d_\lambda(y, x)$ in the right hand side of (2.9). As a result, $d_\lambda(x, y) = d_\lambda(y, x)$, which completes the proof. \square

As a consequence of Lemma 2.9, we obtain the following.

Corollary 2.10 For any $\lambda > \lambda_H$, d_λ is comparable with d_{CC} , that is ,

$$0 < R'_\lambda \leq \frac{d_\lambda(x, y)}{d_{CC}(x, y)} \leq R_\lambda < \infty \quad \forall x, y \in \Omega. \quad (2.11)$$

Consequently, the topology induced by $\{d_\lambda(x, \cdot)\}_{x \in \Omega}$ and $\{d_\lambda(\cdot, x)\}_{x \in \Omega}$ coincides with the one induced by d_{CC} in Ω , and hence, is the Euclidean topology.

Remark 2.11 (i) We remark that in Lemma 2.9 (iii), without the assumption $H(x, p) = H(x, -p)$ for all $p \in \mathbb{R}^m$ and almost all $x \in \Omega$, d_λ may not be symmetric, that is, $d_\lambda(x, y) = d_\lambda(y, x)$ may not hold for all $x, y \in \Omega$.

(ii) If $R'_\lambda = 0$ for some $\lambda > 0$, then the topology induced by $\{d_\lambda(x, \cdot)\}_{x \in \Omega}$ may be different from the Euclidean topology. To wit this, we construct an Hamiltonian $H(p)$ in Euclidean disk $\Omega = \{x \in \mathbb{R}^2 \mid |x| < 1\}$ with $X = \{\frac{\partial}{\partial x_1}, \frac{\partial}{\partial x_2}\}$, which satisfies (H0)-(H3) with $\lambda_H > 0$. Define $H : \Omega \times \mathbb{R}^2 \rightarrow [0, \infty)$ by

$$H(p) = H(p_1, p_2) = \max\{|p|, 2\} \chi_{\{p \in \mathbb{R}^2 \mid p_1 < 0\}} + |p| \chi_{\{p \in \mathbb{R}^2 \mid p_1 \geq 0\}}$$

where χ_E is the characteristic function of the set E . One can check that $H(p)$ satisfies (H0)-(H3). We omit the details.

Now we show

$$R'_1 = \inf\{|p| \mid H(p) \geq 1\} = 0,$$

and thus $\lambda_H \geq 1 > 0$. Indeed, for any $p = (p_1, p_2)$ and $p_1 \geq 0$, one has $H(p) = |p|$ and hence $H(p) \geq 1$ implies $|p| \geq 1$. On the other hand, for any $p = (p_1, p_2)$ and $p_1 < 0$, we always have $H(p) = \max\{|p|, 2\} \geq 2$, and hence

$$R'_1 = \inf\{|p| \mid p = (p_1, p_2) \in \mathbb{R}^2 \text{ with } |p| < 1 \text{ and } p_1 < 0\} = 0.$$

Writing $e_1 = (1, 0)$, we claim that

$$d_1(x, x + ae_1) = 0 \quad \forall x, x + ae_1 \in \Omega \text{ with } a \in (-1, 0]. \quad (2.12)$$

This claim implies that the topology induced by $\{d_1(x, \cdot)\}_{x \in \Omega}$ is different with the Euclidean topology.

To see the claim (2.12), writing $o = (0, 0)$, we only need to show that

$$d_1(o, ae_1) = 0 \quad \forall a \in (-1, 0). \quad (2.13)$$

It then suffices to show that $u(ae_1) - u(0) \leq 0$ for all $u \in W^{1,\infty}(\Omega)$ with $\|H(\nabla u)\|_{L^\infty(\Omega)} \leq 1$. Given such a function u , observe that $\|H(\nabla u)\|_{L^\infty(\Omega)} \leq 1$ implies $\frac{\partial u}{\partial x_1}(x) \geq 0$ and $|\nabla u(x)| \leq 1$ for almost all $x \in \mathbb{R}^2$. Let $\{\gamma_\delta\}_{0 \leq \delta \leq 1}$ be the line segment joining δe_2 and $ae_1 + \delta e_2$ with $e_2 = (0, 1)$, that is,

$$\gamma_\delta(t) := t(a, \delta) + (1-t)(0, \delta) \quad \forall t \in [0, 1].$$

Since $u \in W^{1,\infty}(\Omega)$ implies that u is ACL (see [28, Section 6.1]), there exists $\{\delta_n\}_{n \in \mathbb{N}}$ depending on u such that u is differentiable almost everywhere on γ_{δ_n} . Noting that $\dot{\gamma}_{\delta_n}(t) = -e_1$ and by $\frac{\partial u}{\partial x_1}(x) \geq 0$, one has

$$\nabla u(\gamma_{\delta_n}(t)) \cdot \dot{\gamma}_{\delta_n}(t) = -\frac{\partial u}{\partial x_1}(\gamma_{\delta_n}(t)) \leq 0 \quad \forall t \in [0, 1],$$

and hence

$$u((a, \delta_n)) - u((0, \delta_n)) = \int_0^1 \nabla u(\gamma_{\delta_n}(t)) \cdot \dot{\gamma}_{\delta_n}(t) dt = \int_0^1 -\partial_1 u(\gamma_{\delta_n}(t)) dt \leq 0.$$

Thus

$$u(ae_1) - u(0) = \lim_{n \rightarrow \infty} [u((a, \delta_n)) - u((0, \delta_n))] \leq 0$$

as desired.

Finally, we introduce the pseudo-length property.

Definition 2.12 We say a pseudo-metric space (Ω, ρ) is a pseudo-length space if for all $x, y \in \Omega$,

$$\rho(x, y) := \inf\{\ell_\rho(\gamma) \mid \gamma \in \mathcal{C}(a, b; x, y; \Omega)\}$$

where $\mathcal{C}(a, b; x, y; \Omega)$ denotes the class of all continuous curves $\gamma : [a, b] \rightarrow \Omega$ with $\gamma(a) = x$ and $\gamma(b) = y$, and

$$\ell_\rho(\gamma) := \sup \left\{ \sum_{i=0}^{N-1} \rho(\gamma(t_i), \gamma(t_{i+1})) \mid a = t_0 < t_1 < \cdots < t_N = b \right\}.$$

3 Proof of Theorem 1.3

In this section, we always suppose that the Hamiltonian $H(x, p)$ enjoys assumptions (H0)-(H3). To prove Theorem 1.3, we first need several auxiliary lemmas.

Lemma 3.1 *Suppose that $\{u_j\}_{j \in \mathbb{N}} \subset \dot{W}_X^{1,\infty}(\Omega)$, and there exists $\lambda \geq 0$ such that*

$$\|H(x, Xu_j)\|_{L^\infty(\Omega)} \leq \lambda \text{ for all } j \in \mathbb{N}.$$

If $u_j \rightarrow u$ in $C^0(\Omega)$, then $u \in \dot{W}_X^{1,\infty}(\Omega)$ and $\|H(x, Xu)\|_{L^\infty(\Omega)} \leq \lambda$. Here and in below, for any open set $V \subset \Omega$, $u_j \rightarrow u$ in $C^0(V)$ refers to for any $K \Subset V$, $u_j \rightarrow u$ in $C^0(\overline{K})$.

Proof By Lemma 2.7, one has

$$|u(x) - u(y)| = \lim_{j \rightarrow \infty} |u_j(x) - u_j(y)| \leq \limsup_{j \rightarrow \infty} \|Xu_j\|_{L^\infty(\Omega)} d_{CC}(x, y).$$

By (H3) and $\|H(\cdot, Xu_j)\|_{L^\infty(\Omega)} \leq \lambda$, we have $\|Xu_j\|_{L^\infty(\Omega)} \leq R_\lambda$ for all j , and hence

$$|u(x) - u(y)| \leq R_\lambda d_{CC}(x, y),$$

that is, $u \in \text{Lip}_{d_{CC}}(\Omega)$. By Lemma 2.7 again, we have $u \in \dot{W}_X^{1,\infty}(\Omega)$.

Next we show that $\|H(x, Xu)\|_{L^\infty(\Omega)} \leq \lambda$. It suffices to show that $\|H(x, X(u|_U))\|_{L^\infty(U)} \leq \lambda$ for any $U \Subset \Omega$. Given any $U \Subset \Omega$, we claim that $X(u_j|_U)$ converges to $X(u|_U)$ weakly in $L^2(U, \mathbb{R}^m)$, that is, for all $1 \leq i \leq m$, one has

$$\lim_{j \rightarrow \infty} \int_U u_j X_i^* \phi \, dx = \int_U \phi Xu \, dx \quad \forall \phi \in L^2(U).$$

To see this claim, note that $\|Xu_j\|_{L^\infty(\Omega)} \leq R_\lambda$ for all $j \in \mathbb{N}$, and hence $\|Xu_j\|_{L^2(U)} \leq R_\lambda |U|^{1/2}$ for all $j \in \mathbb{N}$. In other words, for each $k \in \mathbb{N}$, the set $\{X(u_j|_U)\}_{j \in \mathbb{N}}$ is bounded in $L^2(U, \mathbb{R}^m)$. By the weak compactness of $L^2(U, \mathbb{R}^m)$, any subsequence of $\{Xu_j\}_{j \in \mathbb{N}}$ admits a subsubsequence which converges weakly in $L^2(U, \mathbb{R}^m)$. Therefore, to get the above claim, by a contradiction argument we only need to show that for any subsequence $\{Xu_{j_s}\}_{s \in \mathbb{N}}$ of $\{Xu_j\}_{j \in \mathbb{N}}$, if $Xu_{j_s} \rightarrow q_k$ weakly in $L^2(U, \mathbb{R}^m)$ as $s \rightarrow \infty$, then $X(u|_U) = q_k$. For such $\{Xu_{j_s}\}_{s \in \mathbb{N}}$, recalling that $u_j \rightarrow u$ in $C^0(\Omega)$ as $j \rightarrow \infty$, for all $1 \leq i \leq m$ one has

$$\int_U u X_i^* \phi \, dx = \lim_{j \rightarrow \infty} \int_U u_j X_i^* \phi \, dx = \lim_{s \rightarrow \infty} \int_U u_{j_s} X_i^* \phi \, dx = \lim_{s \rightarrow \infty} \int_U (X_i u_{j_s}) \phi \, dx = \int_U q_k \phi \, dx$$

for any $\phi \in C_c^\infty(U)$. This implies that $Xu|_U = q_k$ as desired.

By Mazur's Theorem, for any $l > 0$, we can find a finite convex combination w_l of $\{X(u_j|_U)\}_{j=1}^\infty$ so that $\|w_l - X(u|_U)\|_{L^2(U)} \rightarrow 0$ as $l \rightarrow \infty$. Here w_l is a finite convex combination of $\{X(u_j|_U)\}_{j=1}^\infty$ if there exist $\{\eta_j\}_{j=1}^{k_l}$ for some k_l such that

$$\sum_{i=1}^{k_l} \eta_i = 1 \quad \text{and} \quad w_l = \sum_{j=1}^{k_l} X(u_j|_U)$$

By the quasi-convexity of $H(x, \cdot)$ as in (H1), we have

$$H(x, w_l) = H(x, \sum_{j=1}^{k_l} \eta_j X(u_j|_U)) \leq \sup_{1 \leq j \leq k_l} H(x, X(u_j|_U)) \leq \lambda \quad \text{for almost all } x \in U.$$

Up to considering subsequence we may assume that $w_l \rightarrow X(u|_U)$ almost everywhere in U . By the lower-semicontinuity of $H(x, \cdot)$ as in (H0), we conclude that

$$H(x, X(u|_U)) \leq \liminf_{l \rightarrow \infty} H(x, w_l) \leq \lambda \quad \text{for almost all } x \in U.$$

The proof is complete. \square

Lemma 3.2 *If $v \in \dot{W}_X^{1,\infty}(\Omega)$, then*

$$v^+ = \max\{v, 0\} \in \dot{W}_X^{1,\infty}(\Omega) \quad \text{and} \quad Xv^+ = (Xv)\chi_{\{x \in \Omega, v > 0\}} \quad \text{almost everywhere.} \quad (3.1)$$

Consequently, let $\{v_i\}_{1 \leq i \leq j} \subset \dot{W}_X^{1,\infty}(\Omega)$ for some $j \in \mathbb{N}$. If $u = \max_{1 \leq i \leq j} \{v_i\}$ or $u = \min_{1 \leq i \leq j} \{v_i\}$, then

$$u \in \dot{W}_X^{1,\infty}(\Omega) \quad \text{and} \quad \|H(x, Xu)\|_{L^\infty(\Omega)} \leq \max_{1 \leq i \leq j} \{\|H(x, Xv_i)\|_{L^\infty(\Omega)}\}. \quad (3.2)$$

Proof First we prove (3.1). Let $v \in \dot{W}_X^{1,\infty}(\Omega)$. By Lemma 2.7, $v \in \text{Lip}_{d_{CC}}(\Omega)$. Observe that

$$|v^+(x) - v^+(y)| \leq |v(x) - v(y)| \leq \text{Lip}_{d_{CC}}(v, \Omega) d_{CC}(x, y) \quad \forall x, y \in \Omega,$$

that is, $v^+ \in \text{Lip}_{d_{CC}}(\Omega)$. By Lemma 2.7 again, $v^+ \in \dot{W}_X^{1,\infty}(\Omega)$. To get $Xv^+ = (Xv)\chi_{\{x \in \Omega, v > 0\}}$ almost everywhere, it suffices to consider the restriction $v|_U$ of v in any bounded domain $U \Subset \Omega$, that is, to prove $X(v|_U)^+ = (Xv|_U)\chi_{\{x \in U, v > 0\}}$ almost everywhere. But this always holds thanks to Lemma 2.2 and the fact $v|_U \in \dot{W}_X^{1,p}(U)$ for any $1 \leq p < \infty$.

Next we prove (3.2). If $u = \max\{v_1, v_2\}$, where $v_i \in \dot{W}_X^{1,\infty}(\Omega)$ for $i = 1, 2$, then $u = v_2 + (v_1 - v_2)^+$. By (3.1), $u \in \dot{W}_X^{1,\infty}(\Omega)$ and

$$\begin{aligned} Xu &= Xv_2 + X(v_1 - v_2)^+ \\ &= Xv_2 + [(X(v_1 - v_2))\chi_{\{x \in \Omega, v_1 > v_2\}}] \\ &= (Xv_2)\chi_{\{x \in \Omega, v_1 \leq v_2\}} + (Xv_1)\chi_{\{x \in \Omega, v_1 > v_2\}}. \end{aligned}$$

Thus

$$H(x, Xu(x)) = H(x, Xv_2)\chi_{\{x \in \Omega, v_1 \leq v_2\}} + H(x, Xv_1)\chi_{\{x \in \Omega, v_1 > v_2\}} \quad \text{for almost all } x \in \Omega.$$

A similar argument holds for $u = \min\{v_1, v_2\}$. This gives (3.2) when $j = 2$. By an induction argument, we get (3.2) for all $j \geq 2$. \square

Lemma 3.3 *For any $\lambda \geq 0$ and $x \in \Omega$, we have $d_\lambda(x, \cdot), d_\lambda(\cdot, x) \in \dot{W}_X^{1,\infty}(\Omega)$ and*

$$\|H(\cdot, Xd_\lambda(x, \cdot))\|_{L^\infty(\Omega)} \leq \lambda \quad \text{and} \quad \|H(\cdot, -Xd_\lambda(\cdot, x))\|_{L^\infty(\Omega)} \leq \lambda.$$

Proof Given any $x \in \Omega$ and $\lambda \geq 0$, write $v(z) = d_\lambda(x, z)$ for all $z \in \Omega$. To see $H(\cdot, Xv) \leq \lambda$ almost everywhere, by Lemma 3.1, it suffices to find a sequence of function $u_j \in \dot{W}_X^{1,\infty}(\Omega)$ so that $H(\cdot, Xu_j) \leq \lambda$ almost everywhere and $u_j \rightarrow v$ in $C^0(\Omega)$ as $j \rightarrow \infty$.

To this end, let $\{K_j\}_{j \in \mathbb{N}}$ be a sequence of compact subsets in Ω with

$$\Omega = \bigcup_{j \in \mathbb{N}} K_j \quad \text{and} \quad K_j \subset K_{j+1}^\circ.$$

For $j \in \mathbb{N}$ and $y \in K_j$, by definition of d_λ we can find a function $v_{y,j} \in \dot{W}_X^{1,\infty}(\Omega)$ such that $H(\cdot, Xv_{y,j}) \leq \lambda$ almost everywhere,

$$d_\lambda(x, y) - \frac{1}{2j} \leq v_{y,j}(y) - v_{y,j}(x).$$

Since Lemma 2.9 implies that $d_\lambda(x, \cdot)$ is continuous, there exists an open neighbourhood $N_{y,j}$ of y with

$$d_\lambda(x, z) - \frac{1}{j} \leq v_{y,j}(z) - v_{y,j}(x), \quad \forall z \in N_{y,j}.$$

Thanks to the compactness of K_j , there exist $y_1, \dots, y_l \in K_j$ such that $K_j \subset \bigcup_{i=1}^l N_{y_i,j}$. Write

$$u_j(z) := \max\{v_{y_i,j}(z) - v_{y_i,j}(x) : i = 1, \dots, l\} \quad \forall z \in K_j.$$

Then $u_j(x) = 0$, and

$$d_\lambda(x, z) \leq u_j(z) + \frac{1}{j} \text{ for all } z \in K_j.$$

Moreover by Lemma 3.2 we have

$$H(\cdot, Xu_j) \leq \lambda \quad \text{in } \Omega.$$

Since

$$d_\lambda(x, z) \geq u_j(z) \text{ for all } z \in K_j$$

is clear, we conclude that $u_j \rightarrow v$ in $C^0(K_i)$ as $j \rightarrow \infty$ for all i , and hence $u_j \rightarrow v$ uniformly in any compact subset of Ω as $j \rightarrow \infty$.

Similarly, we can show $\|H(x, -Xd_\lambda(\cdot, x))\|_{L^\infty(\Omega)} \leq \lambda$ which finishes the proof. \square

In general, for any $E \subset \Omega$, we define

$$d_{\lambda,E}(z) := \inf_{x \in E} \{d_\lambda(x, z)\}.$$

Lemma 3.4 For any set $E \subset \Omega$, we have $d_{\lambda,E} \in \dot{W}_X^{1,\infty}(\Omega)$ and $\|H(x, Xd_{\lambda,E})\|_{L^\infty(\Omega)} \leq \lambda$.

Proof Let $\{K_j\}_{j \in \mathbb{N}}$ be a sequence of compact subsets in Ω with $\Omega = \bigcup_{j \in \mathbb{N}} K_j$ and $K_j \subset K_{j+1}^\circ$. For each j and $y \in K_j$, we can find $z_{y,j} \in E$ such that

$$d_{\lambda,E}(y) \leq d_{\lambda,z_{y,j}}(y) \leq d_{\lambda,E}(y) + 1/2j.$$

Thus there exists a neighborhood $N(y)$ of y such that

$$d_{\lambda,E}(y) \leq d_{\lambda,z_{y,j}}(y) \leq d_{\lambda,E}(z) + 1/j \quad \forall z \in N(y).$$

So we can find $\{y_1, \dots, y_{l_j}\}$ such that $K_j = \bigcup_{i=1}^{l_j} N(y_i)$ for all $i = 1, \dots, l_j$. Write

$$u_j(z) = \min_{i=1, \dots, l_j} \{d_{\lambda,z_{y_i,j}}(z)\} \quad \forall z \in K_j.$$

$$d_{\lambda,E} \leq u_j(z) \leq d_{\lambda,E} + 1/j \quad \text{in } K_j.$$

This means that $u_j \rightarrow d_{\lambda,E}$ in $C^0(\Omega)$ as $j \rightarrow \infty$. Note that

$$\|H(\cdot, Xu_j)\|_{L^\infty(\Omega)} \leq \lambda.$$

By Lemma 3.1 we have $d_{\lambda,E} \in \dot{W}_X^{1,\infty}(\Omega)$ and $\|H(\cdot, Xd_{\lambda,E})\|_{L^\infty(\Omega)} \leq \lambda$ as desired. \square

We are able to prove (i) \Rightarrow (ii) \Rightarrow (iii) \Rightarrow (i) in Theorem 1.3 as below.

Proof (Proofs of (i) \Rightarrow (ii) \Rightarrow (iii) \Rightarrow (i) in Theorem 1.3) The definition of d_λ directly gives (i) \Rightarrow (ii). Obviously, (ii) \Rightarrow (iii). Below we prove (iii) \Rightarrow (i). Recall that (iii) says that $u(y) - u(z) \leq d_\lambda(z, y)$ for all $x \in \Omega$ and $y, z \in N(x)$, where $N(x) \subset \Omega$ is a neighbourhood of x . To get (i), since $\Omega = \bigcup_{x \in \Omega} N(x)$, it suffices to show that for all $x \in \Omega$, one has $u \in W_X^{1,\infty}(N(x))$ and $\|H(\cdot, Xu)\|_{L^\infty(N(x))} \leq \lambda$.

Fix any x , and write $U = N(x)$. Without loss of generality we assume that U is bounded.

Notice that $u \in L^\infty(U)$. Let $M \in \mathbb{N}$ so that $M \geq \sup_U |u|$. For each $k \in \mathbb{N}$ and $l \in \{-Mk, \dots, Mk\}$, set

$$u_{k,l}(z) := l/k + d_{\lambda, F_{k,l}}(z) \quad \forall z \in U$$

where

$$F_{k,l} := \{y \in U \mid u(y) \leq \frac{l}{k}\}.$$

By Lemma 3.4, one has

$$\|H(\cdot, Xu_{k,l})\|_{L^\infty(U)} \leq \lambda.$$

Set

$$u_k(z) := \min_{l \in \{-Mk, \dots, Mk\}} u_{k,l}(z) \quad \forall z \in U.$$

To get $u \in \dot{W}_X^{1,\infty}(U)$ and $\|H(\cdot, Xu)\|_{L^\infty(U)} \leq \lambda$, thanks to Lemma 3.1 with $\Omega = U$, we only need to show $u_k \rightarrow u$ in $C^0(U)$ as $k \rightarrow \infty$.

To see $u_k \rightarrow u$ in $C^0(U)$ as $k \rightarrow \infty$, note that, for any $k \in \mathbb{N}$, $-M \leq u \leq M$ in U implies $-Mk \leq ku \leq Mk$ in U . Thus, at any $z \in U$, we can find $j \in \mathbb{N}$ with $-k \leq j \leq k$, which depends on z , such that $Mj \leq ku(z) \leq M(j+1)$. Letting $l = Mj$, we have $\frac{l}{k} \leq u(z) \leq \frac{l+M}{k}$. We claim that $u_k(z) \in [u(z), \frac{l+M}{k}]$. Obviously, this claim gives

$$|u(z) - u_k(z)| \leq \frac{M}{k} \quad \forall z \in U,$$

and hence, the desired convergence $u_k \rightarrow u$ in $C^0(U)$ as $k \rightarrow \infty$.

Below we prove the above claim $u_k(z) \in [u(z), \frac{l+M}{k}]$. Recall that $u(z) \in [\frac{l}{k}, \frac{l+M}{k}]$ for some $l = Mj$ with $-k \leq j \leq k$.

First, we prove $u_k(z) \leq \frac{l+M}{k}$. If $l + M > Mk$, then $M < (l + M)/k$. Since

$$F_{k,Mk} = \{y \in U \mid u(y) \leq M\} = U,$$

we have $d_{\lambda, F_{k,Mk}}(z) = 0$ and hence,

$$u_{k,Mk}(z) = M + d_{\lambda, F_{k,Mk}}(z) = M < \frac{l + M}{k}.$$

Therefore,

$$u_k(z) \leq u_{k,Mk}(z) < \frac{l + M}{k}. \quad (3.3)$$

If $l + M \leq Mk$, then $u(z) \in [\frac{l}{k}, \frac{l+M}{k}]$ implies $z \in F_{k,l+M}$ and hence $d_\lambda(z, F_{k,l+M}) = 0$. Thus

$$u_k(z) \leq u_{k,l+M}(z) = \frac{l + M}{k} + d_\lambda(z, F_{k,l+M}) = \frac{l + M}{k}. \quad (3.4)$$

Combining (3.4) and (3.3), we have $u_k(z) \leq \frac{l+M}{k}$ as desired.

Next, we prove $u(z) \leq u_k(z)$. For any $-k \leq j \leq k$ with $Mj \leq l$, since $u(z) \geq \frac{l}{k} \geq \frac{Mj}{k}$, we can find $w \in \partial F_{k,Mj}$ such that $d_{\lambda, F_{k,Mj}}(z) = d_{\lambda}(w, z)$. Since $w \in \partial F_{k,Mj}$, we deduce that $u(w) = \frac{Mj}{k}$ and

$$u_{k,Mj}(z) = \frac{Mj}{k} + d_{\lambda, F_{k,Mj}}(z) = u(w) + d_{\lambda}(w, z). \quad (3.5)$$

Note that $w \in \overline{U}$, hence there exists a sequence $\{w_s\}_{s \in \mathbb{N}} \subset U$ such that $w_s \rightarrow w$ as $s \rightarrow \infty$.

Thus by the assumption (iii),

$$u(z) - u(w) = \lim_{s \rightarrow \infty} [u(z) - u(w_s)] \leq \lim_{s \rightarrow \infty} d_{\lambda}(w_s, z)$$

By the triangle inequality and $d_{\lambda}(w_s, w) \leq R_{\lambda} d_{CC}(w_s, w)$ given in Lemma 2.9, we then obtain

$$u(z) - u(w) \leq \lim_{s \rightarrow \infty} [d_{\lambda}(w_s, w) + d_{\lambda}(w, z)] = d_{\lambda}(w, z). \quad (3.6)$$

Combining (3.5) and (3.6), we have

$$u(z) \leq u(w) + d_{\lambda}(w, z) = u_{k,Mj}(z). \quad (3.7)$$

On the other hand, for any $-k \leq j \leq k$ with $Mj > l$, we have

$$u_{k,Mj}(z) \geq \frac{Mj}{k} \geq \frac{M(l+1)}{k} > u(z).$$

From this and (3.7), it follows that

$$u_k(z) = \min_{j \in \{-k, \dots, l/M\}} u_{k,Mj}(z) \geq u(z)$$

as desired. \square

4 Proof of Theorem 1.4

In this section, we always suppose that the Hamiltonian $H(x, p)$ enjoys assumptions (H0)-(H3).

To prove Theorem 1.4, we need to show that (Ω, d_{λ}) is a pseudo-length space for all $\lambda \geq \lambda_H$ in the sense of Definition 2.12. In other words, define

$$\rho_{\lambda}(x, y) := \inf\{\ell_{d_{\lambda}}(\gamma) \mid \gamma \in \mathcal{C}(a, b; x, y; \Omega)\},$$

where we recall the pseudo-length $\ell_{d_{\lambda}}(\gamma)$ induced by d_{λ} defined in Definition 2.12 and λ_H in (1.7). We have the following.

Proposition 4.1 *For any $\lambda \geq \lambda_H$, we have $d_{\lambda} = \rho_{\lambda}$.*

To prove Proposition 4.1, we need the following approximation midpoint property of d_{λ} .

Proposition 4.2 *For any $\lambda \geq 0$, we have*

$$\inf_{z \in \Omega} \max\{d_{\lambda}(x, z), d_{\lambda}(z, y)\} \leq \frac{1}{2} d_{\lambda}(x, y) \quad \text{for all } x, y \in \Omega. \quad (4.1)$$

Proof We prove by contradiction. Suppose that (4.1) were not true. There exists $x_0, y_0 \in \Omega$ such that

$$\inf_{z \in \Omega} \max\{d_\lambda(x_0, z), d_\lambda(z, y_0)\} \geq \frac{1}{2}d_\lambda(x_0, y_0) + \epsilon_0 := r_0 \quad (4.2)$$

for some $\epsilon_0 > 0$.

Given any $\delta \in (0, \epsilon_0)$, define $f(z) := f_1(z) + f_2(z)$ with

$$f_1(z) := \min\{d_\lambda(x_0, z) - (r_0 - \delta), 0\}, \quad f_2(z) := \max\{(r_0 - \delta) - d_\lambda(z, y_0), 0\} \quad \forall z \in \Omega.$$

We claim that f satisfies Theorem 1.3(iii), that is, for any $z \in \Omega$, there is an open neighborhood $N(z)$ such that

$$f(y) - f(w) \leq d_\lambda(w, y) \quad \forall w, y \in N(z). \quad (4.3)$$

Assume the claim (4.3) holds for the moment. Since we have already shown the equivalence between (ii) and (iii) in Theorem 1.3, we know that f satisfies Theorem 1.3(ii), that is,

$$f(y) - f(w) \leq d_\lambda(w, y) \quad \forall w, y \in \Omega.$$

In particular,

$$f(y_0) - f(x_0) \leq d_\lambda(x_0, y_0). \quad (4.4)$$

On the other hand, we have $f_1(x_0) = -(r_0 - \delta)$ and $f_2(y_0) = r_0 - \delta$. Since (4.2) implies

$$d_\lambda(x_0, y_0) = \max\{d_\lambda(x_0, x_0), d_\lambda(x_0, y_0)\} \geq \frac{1}{2}d_\lambda(x_0, y_0) + \epsilon_0 = r_0$$

and $f_2(x_0) = 0$ and $f_1(y_0) = 0$. Therefore,

$$f(y_0) - f(x_0) = f_2(y_0) - f_1(x_0) = 2r_0 - 2\delta = d_\lambda(x_0, y_0) + 2\epsilon_0 - 2\delta,$$

By $\delta < \epsilon_0$, one has

$$f(y_0) - f(x_0) > d_\lambda(x_0, y_0),$$

which contradicts to (4.4).

Finally we prove the above claim (4.3). Firstly, thanks to Lemma 3.2 and 3.3, $H(x, Xf_1) \leq \lambda$ and $H(x, Xf_2) \leq \lambda$ almost everywhere in Ω , and hence, by the definition of d_λ ,

$$f_1(y) - f_1(w) \leq d_\lambda(w, y) \text{ and } f_2(y) - f_2(w) \leq d_\lambda(w, y) \quad \forall w, y \in \Omega.$$

Next, set

$$\Lambda_1 := \{z \in \Omega \mid d_\lambda(x_0, z) < r_0\}, \quad \Lambda_2 := \{z \in \Omega \mid d_\lambda(z, y_0) < r_0\}.$$

and

$$\Lambda_3 := \{z \in \Omega \mid d_\lambda(x_0, z) > r_0 - \delta \text{ and } d_\lambda(z, y_0) > r_0 - \delta\}$$

For any $z \in \Lambda_1$, that is, $d_\lambda(x_0, z) < r_0$, (4.2) implies $d_\lambda(z, y_0) \geq r_0$. Consequently,

$$f_2(z) = \max\{(r_0 - \delta) - d_\lambda(z, y_0), 0\} = 0$$

and hence, $f(z) = f_1(z)$. Consequently,

$$f(w) - f(y) = f_1(w) - f_1(y) \leq d_\lambda(y, w) \quad \forall w, y \in \Lambda_1. \quad (4.5)$$

Similarly, for any $z \in \Lambda_2$, that is, $d_\lambda(z, y_0) < r_0$, (4.2) implies $d_\lambda(x_0, z) \geq r_0$. Consequently,

$$f_1(z) = \min\{d_\lambda(x_0, z) - (r_0 - \delta), 0\} = 0$$

and hence, $f(z) = f_2(z)$. Consequently,

$$f(w) - f(y) = f_2(w) - f_2(y) \leq d_\lambda(y, w) \quad \forall w, y \in \Lambda_2. \quad (4.6)$$

For any $z \in \Lambda_3$, that is, $d_\lambda(x_0, z) > r_0 - \delta$ and $d_\lambda(z, y_0) > r_0 - \delta$, we have $f_1(z) = 0 = f_2(z)$ and hence $f(z) = 0$. Consequently

$$f(w) - f(y) = 0 \leq d_\lambda(y, w) \quad \forall w, y \in \Lambda_3. \quad (4.7)$$

Noticing that $\{\Lambda_i\}_{i=1,2,3}$ forms an open cover of Ω , for any $z \in \Omega$, we choose

$$N(z) = \begin{cases} \Lambda_1 & \text{if } z \in \Lambda_1, \\ \Lambda_2 & \text{if } z \in \Lambda_2 \setminus \Lambda_1, \\ \Lambda_3 & \text{if } z \in \Omega \setminus (\Lambda_1 \cup \Lambda_2). \end{cases} \quad (4.8)$$

From (4.5), (4.6) and (4.7), we obtain (4.3) as desired with the choice of $N(z)$ in (4.8). The proof is complete. \square

Lemma 4.3 *Given any $x, y \in \Omega$, the map $\lambda \in [\lambda_H, \infty) \mapsto d_\lambda(x, y) \in [0, \infty)$ is nondecreasing and right continuous.*

Proof The fact that $d_\lambda(x, y)$ is non-decreasing with respect to λ is obvious for any $x, y \in \Omega$ from the definition of d_λ . Given any $x, y \in \Omega$, we show the right-continuity the map $\lambda \in [\lambda_H, \infty) \mapsto d_\lambda(x, y) \in [0, \infty)$. We argue by contradiction. Assume there exists $\lambda_0 \geq \lambda_H$ and $x, y \in \Omega$ such that

$$\liminf_{\lambda \rightarrow \lambda_0+} d_\lambda(x, y) = \lim_{\lambda \rightarrow \lambda_0+} d_\lambda(x, y) = c > d_{\lambda_0}(x, y). \quad (4.9)$$

Let $w_\lambda(\cdot) := d_\lambda(x, \cdot)$. By Lemma 3.3, we know $\|H(\cdot, Xw_\lambda)\|_{L^\infty(\Omega)} \leq \lambda$. Since $\{w_\lambda\}_{\lambda > \lambda_0}$ is a non-decreasing sequence with respect to λ , $\{w_\lambda\}$ converges pointwise to a function w as $\lambda \rightarrow \lambda_0+$ and for any set $V \subseteq \Omega$ and $x, y \in \bar{V}$, we have $w_\lambda \rightarrow w$ in $C^0(\bar{V})$. Then applying Lemma 3.1, we have

$$\|H(\cdot, Xw)\|_{L^\infty(\bar{V})} \leq \lambda$$

for any $\lambda > \lambda_0$, which implies

$$\|H(\cdot, Xw)\|_{L^\infty(\bar{V})} \leq \lambda_0. \quad (4.10)$$

By the definition of w , we have

$$w(x) = \lim_{\lambda \rightarrow \lambda_0+} d_\lambda(x, x) = 0, \text{ and } w(y) = \lim_{\lambda \rightarrow \lambda_0+} d_\lambda(x, y) = c \quad (4.11)$$

Combining (4.10) and (4.11) and applying Theorem 1.3, we have

$$c - 0 = w(y) - w(x) \leq d_{\lambda_0}(x, y),$$

which contradicts to (4.9). The proof is complete. \square

We are in the position to show

Proof of Proposition 4.1 We consider the cases $\lambda > \lambda_H$ and $\lambda = \lambda_H$ separately.

Case 1. $\lambda > \lambda_H$. First, $d_\lambda \leq \rho_\lambda$ follows from the triangle inequality for d_λ .

To see $\rho_\lambda \leq d_\lambda$, it suffices to prove that for any $z \in \Omega$, the function $\rho_\lambda(z, \cdot) : \Omega \rightarrow \mathbb{R}$ satisfies Theorem 1.3(iii), that is, for any $x \in \Omega$ we can find a neighborhood $N(x)$ of x such that

$$\rho_\lambda(z, y) - \rho_\lambda(z, w) \leq d_\lambda(w, y) \quad \forall w, y \in N(x). \quad (4.12)$$

Indeed, since we have already shown the equivalence of (i) and (iii) in Theorem 1.3, (4.12) implies that $\rho_\lambda(z, \cdot)$ satisfies Theorem 1.3(i), that is, $\rho_\lambda(z, \cdot) \in \dot{W}_X^{1,\infty}(\Omega)$ and $\|H(\cdot, X\rho_\lambda(z, \cdot))\|_{L^\infty(\Omega)} \leq \lambda$. Taking $\rho_\lambda(z, \cdot)$ as the test function in the definition of $d_\lambda(z, x)$, one has

$$\rho_\lambda(z, x) \leq d_\lambda(z, x) \quad \forall x \in \Omega$$

as desired.

To prove (4.12), let $z \in \Omega$ be fixed. For any $x \in \Omega$ and any $t > 0$, write

$$B_{d_\lambda}^+(x, t) := \{y \in \Omega \mid d_\lambda(x, y) < t \text{ or } d_\lambda(y, x) < t\}$$

and

$$B_{d_\lambda}^-(x, t) := \{y \in \Omega \mid d_\lambda(x, y) < t \text{ and } d_\lambda(y, x) < t\}.$$

For any $x \in \Omega$, letting $r_x = \min\{\frac{R'_\lambda}{10}d_{CC}(x, \partial\Omega), 1\}$, by Corollary 2.10, we have

$$B_{d_\lambda}^+(x, 6r_x) \subset B_{d_{CC}}(x, \frac{6r_x}{R'_\lambda}) \Subset \Omega, \quad (4.13)$$

where $R'_\lambda > 0$ thanks to (H3). Write $N(x) = B_{d_\lambda}^-(x, r_x)$. Given any $w, y \in N(x)$, it then suffices to prove $\rho_\lambda(z, y) - \rho_\lambda(z, w) \leq d_\lambda(w, y)$. To this end, for any $0 < \epsilon < \frac{1}{2}d_\lambda(w, y)$, we will construct a curve

$$\gamma_\epsilon : [0, 1] \rightarrow B_{d_\lambda}^+(x, 6r_x) \text{ with } \gamma_\epsilon(0) = w, \gamma_\epsilon(1) = y \text{ and } \ell_{d_\lambda}(\gamma_\epsilon) \leq d_\lambda(w, y) + \epsilon. \quad (4.14)$$

Assume the existence of γ_ϵ for the moment. By the triangle inequality for ρ_λ , we have

$$\rho_\lambda(z, y) - \rho_\lambda(z, w) \leq \ell_{d_\lambda}(\gamma_\epsilon) \leq d_\lambda(w, y) + \epsilon$$

By sending $\epsilon \rightarrow 0$, this yields $\rho_\lambda(z, y) - \rho_\lambda(z, w) \leq d_\lambda(w, y)$ as desired.

Construction of a curve γ_ϵ satisfying (4.14). For each $t \in \mathbb{N}$, set

$$D_t := \{k2^{-t} \mid k \in \mathbb{N}, 0 \leq k \leq 2^t\}.$$

We will use induction and Proposition 4.2 to construct a set

$$Y_t = \{y_s\}_{s \in D_t} \subset B_{d_\lambda}^+(x, 5r_x) \quad (4.15)$$

with $y_0 = w$ and $y_1 = y$ so that $Y_t \subset Y_{t+1}$, and that

$$d_\lambda(y_{j2^{-t}}, y_{(j+1)2^{-t}}) \leq 2^{-t}(d_\lambda(w, y) + \epsilon) \text{ for any } 0 \leq j \leq 2^t - 1. \quad (4.16)$$

The construction of $\{Y_t\}_{t \in \mathbb{N}}$ is postponed to the end of this proof. Assuming that $\{Y_t\}_{t \in \mathbb{N}}$ are constructed, we are able to construct γ_ϵ as below.

Firstly, set $D := \cup_{t \in \mathbb{N}} D_t$ and $Y := \cup_{t \in \mathbb{N}} Y_t$. Given any $s_1, s_2 \in D$ with $s_1 < s_2$, there exists $t \in \mathbb{N}$ such that $s_1 = l2^{-t} \in D_t$, $s_2 = k2^{-t} \in D_t$ for some $l < k$ and hence $y_{s_1}, y_{s_2} \in Y_t$. Using (4.16) and the triangle inequality for d_λ , we have

$$d_\lambda(y_{s_1}, y_{s_2}) \leq \sum_{j=l}^{k-1} d_\lambda(y_{j2^{-t}}, y_{(j+1)2^{-t}}) \leq |k-l|2^{-t}(d_\lambda(w, y) + \epsilon) = |s_1 - s_2|(d_\lambda(w, y) + \epsilon). \quad (4.17)$$

Next, define a map $\gamma_\epsilon^0 : D \rightarrow Y$ by $\gamma_\epsilon^0(s) = y_s$ for all $s \in D$. The above inequality (4.17) implies that

$$\lim_{D \ni s' \rightarrow s} \gamma_\epsilon^0(s') = \gamma_\epsilon^0(s) \text{ for all } s \in D.$$

Since D is dense in $[0, 1]$ and $\overline{B_{d_\lambda}^+(y_0, 5r_x)}$ is complete, it is standard to extend γ_ϵ^0 uniquely to a continuous map $\gamma_\epsilon : [0, 1] \rightarrow B_{d_\lambda}^+(y_0, 6r_x)$, that is, $\gamma_\epsilon(s) = \gamma_\epsilon^0(s)$ for any $s \in D$ and

$$\gamma_\epsilon(s) = \lim_{D \ni s' \rightarrow s} \gamma_\epsilon(s') = \lim_{D \ni s' \rightarrow s} y_{s'} \text{ for any } s \in [0, 1] \setminus D.$$

Recalling (4.17), one therefore has

$$d_\lambda(\gamma_\epsilon(s_1), \gamma_\epsilon(s_2)) \leq |s_1 - s_2|(d_\lambda(x, y) + \epsilon) \quad \forall s_1, s_2 \in [0, 1], s_1 \leq s_2,$$

which gives $\ell_{d_\lambda}(\gamma_\epsilon) \leq \delta + \epsilon$. Thus the curve γ_ϵ satisfies (4.14) as desired.

Construction of $\{Y_t\}_{t \in \mathbb{N}}$ via induction and Proposition 4.2. Since $y, w \in B_{d_\lambda}^-(x, r_x)$, we have $d_\lambda(w, x) < r_x$ and $d_\lambda(x, y) < r_x$, which implies

$$\delta := d_\lambda(w, y) \leq d_\lambda(w, x) + d_\lambda(x, y) < 2r_x.$$

We construct $Y_1 = \{y_0, y_{1/2}, y_1\}$ which satisfies (4.16) with $t = 1$ and

$$Y_1 \subset B_{d_\lambda}^+(x, 3r_x) \subset B_{d_\lambda}^+(x, 5r_x). \quad (4.18)$$

We set $y_0 = w$ and $y_1 = y$. Noting that Proposition 4.2 gives

$$\inf_{z \in \Omega} \max\{d_\lambda(y_0, z), d_\lambda(z, y_1)\} \leq \frac{1}{2}d_\lambda(y_0, y_1)$$

we choose $y_{1/2} \in \Omega$ so that

$$\max\{d_\lambda(y_0, y_{1/2}), d_\lambda(y_{1/2}, y_1)\} \leq \frac{1}{2}\delta + \frac{1}{4}\epsilon. \quad (4.19)$$

Obviously, (4.19) gives (4.16). To see (4.18), obviously,

$$y_0, y_1 \in B_{d_\lambda}^-(x, r_x) \subset B_{d_\lambda}^+(x, 3r_x).$$

Moreover, noting that $0 < \epsilon < \frac{1}{2}\delta < r_x$ implies

$$\frac{1}{2}\delta + \frac{1}{4}\epsilon < \delta < 2r_x$$

and that $y \in B_{d_\lambda}^-(x, r_x)$ implies $d_\lambda(y, x) \leq r_x$, we have

$$d_\lambda(y_{1/2}, x) \leq d_\lambda(y_{1/2}, y_1) + d_\lambda(y_1, x) \leq \frac{1}{2}\delta + \frac{1}{4}\epsilon + d_\lambda(y, x) \leq 3r_x,$$

which gives $y_{1/2} \in B_{d_\lambda}^+(x, 3r_x)$.

In general, by induction given any $t \geq 2$, assume that $Y_{t-1} = \{y_s\}_{s \in D_{t-1}}$ is constructed so that

$$Y_{t-1} \subset B_{d_\lambda}^+(x, (3 + \sum_{l=1}^{t-2} 2^{-l})r_x + \epsilon \sum_{l=1}^{t-1} 2^{-l}) \quad (4.20)$$

and that

$$d_\lambda(y_{j2^{-(t-1)}}, y_{(j+1)2^{-(t-1)}}) \leq 2^{-(t-1)} \left(\delta + \epsilon \sum_{l=1}^{t-1} 2^{-l} \right) \text{ for any } 0 \leq j \leq 2^{t-1} - 1. \quad (4.21)$$

Here and in what follows, we make the convention that $\sum_{l=1}^{t-2} 2^{-l} = 0$ if $t = 2$. Since

$$(3 + \sum_{l=1}^{\infty} 2^{-l})r_x + \epsilon \sum_{l=1}^{\infty} 2^{-l} \leq 4r_x + \epsilon < 4r_x + \delta \leq 5r_x, \quad (4.22)$$

the inclusion (4.20) implies $Y_{t-1} \subset B_{d_\lambda}^+(x, 5r_x)$ and hence (4.15).

Below, we construct $Y_t = \{y_s\}_{s \in D_t}$ satisfying (4.16) and

$$Y_t \subset B_{d_\lambda}^+(x, (3 + \sum_{l=1}^{t-1} 2^{-l})r_x + \epsilon \sum_{l=1}^t 2^{-l}), \quad (4.23)$$

Note that (4.22) and (4.23) imply $Y_t \subset B_{d_\lambda}^+(x, 5r_x)$ and hence (4.15).

We define $Y_t = \{y_s\}_{s \in D_t}$ first. Since $D_{t-1} \subsetneq D_t$, for $s \in D_{t-1}$, $y_s \in Y_{t-1}$ is defined. It is left to define y_s for $s \in D_t \setminus D_{t-1}$. Given any $s \in D_t \setminus D_{t-1}$, we know that $s = j2^{-t} \in D_t \setminus D_{t-1}$ for some odd j with $1 \leq j \leq 2^t - 1$. Write $s' = (j-1)2^{-t}$ and $s'' = (j+1)2^{-t}$. Then $s', s'' \in D_{t-1}$ and hence $y_{s'} \in Y_{t-1}$ and $y_{s''} \in Y_{t-1}$ are defined. Since Proposition 4.2 gives

$$\inf_{z \in \Omega} \max\{d_\lambda(y_{s'}, z), d_\lambda(z, y_{s''})\} \leq \frac{1}{2}d_\lambda(y_{s'}, y_{s''}) \quad (4.24)$$

we choose $y_s \in \Omega$ such that

$$\max\{d_\lambda(y_{s'}, y_s), d_\lambda(y_s, y_{s''})\} \leq \frac{1}{2}d_\lambda(y_{s'}, y_{s''}) + 2^{-2t}\epsilon. \quad (4.25)$$

Note that (4.25) and (4.21) gives (4.16) directly. Indeed, for any $0 \leq j \leq 2^t - 1$, if j is odd, applying (4.25) with $s = j2^{-t}$, $s' = (j-1)2^{-t}$ and $s'' = (j+1)2^{-t}$, we deduce that

$$d_\lambda(y_{j2^{-t}}, y_{(j+1)2^{-t}}) = d_\lambda(y_s, y_{s''}) \leq \frac{1}{2}d_\lambda(y_{s'}, y_{s''}) + 2^{-2t}\epsilon.$$

Since $s', s'' \in D_{t-1}$ and $s'' = s' + 2^{-(t-1)}$, applying (4.21) to $y_{s'}, y_{s''}$ we have

$$d_\lambda(y_{j2^{-t}}, y_{(j+1)2^{-t}}) \leq \frac{1}{2}2^{-(t-1)}\left(\delta + \epsilon \sum_{l=1}^{t-1} 2^{-l}\right) + 2^{-t}2^{-t}\epsilon = 2^{-t}\left(\delta + \epsilon \sum_{l=1}^t 2^{-l}\right). \quad (4.26)$$

If j is even, then $j \leq 2^t - 2$. Applying (4.25) with $s = (j+1)2^{-t}$, $s' = j2^{-t}$ and $s'' = (j+2)2^{-t}$, we deduce that

$$d_\lambda(y_{j2^{-t}}, y_{(j+1)2^{-t}}) = d_\lambda(y_{s'}, y_s) \leq \frac{1}{2}d_\lambda(y_{s'}, y_{s''}) + 2^{-2t}\epsilon.$$

Similarly, we also have (4.26).

To see (4.23), since (4.20) gives

$$Y_{t-1} \subset B_{d_\lambda}^+(x, (3 + \sum_{l=1}^{t-2} 2^{-l})r_x + \epsilon \sum_{l=1}^{t-1} 2^{-l}) \subset B_{d_\lambda}^+(x, (3 + \sum_{l=1}^{t-1} 2^{-l})r_x + \epsilon \sum_{l=1}^t 2^{-l}), \quad (4.27)$$

it suffices to check

$$\begin{aligned} Y_t \setminus Y_{t-1} &= \{y_{j2^{-t}} \mid 1 \leq j \leq 2^t - 1, \\ j \text{ is odd}\} &\subset B_{d_\lambda}^+(x, (3 + \sum_{l=1}^{t-1} 2^{-l})r_x + \epsilon \sum_{l=1}^t 2^{-l}). \end{aligned}$$

For any odd number j with $1 \leq j \leq 2^t - 1$, since $y_{(j-1)2^{-t}} \in Y_{t-1}$, combining (4.26) and (4.27) and noting $\epsilon < \delta < 2r_x$, we obtain

$$d_\lambda(y_{j2^{-t}}, x) \leq d_\lambda(y_{j2^{-t}}, y_{(j-1)2^{-t}}) + d_\lambda(y_{(j-1)2^{-t}}, x)$$

$$\begin{aligned}
&\leq 2^{-t} \left(2r_x + \epsilon \sum_{l=1}^t 2^{-l} \right) + \left(3 + \sum_{l=1}^{t-2} 2^{-l} \right) r_x + \epsilon \sum_{l=1}^{t-1} 2^{-l} \\
&\leq \left(3 + \sum_{l=1}^{t-1} 2^{-l} \right) r_x + \epsilon \sum_{l=1}^t 2^{-l}
\end{aligned}$$

which implies (4.23). We finish the proof of Case 1.

Case 2. $\lambda = \lambda_H$. Fix $x, y \in \Omega$. For any $\epsilon > 0$ sufficiently small, by the right continuity of the map $\lambda \mapsto d_\lambda(x, y)$ at $\lambda = \lambda_H$ from Lemma 4.3, there exists $\mu > \lambda_H$ such that

$$d_\mu(x, y) < d_{\lambda_H}(x, y) + \frac{\epsilon}{2}. \quad (4.28)$$

By Case (i), there exists $\gamma : [0, 1] \rightarrow \Omega$ joining x and y such that

$$\ell_{d_\mu}(\gamma) < d_\mu(x, y) + \frac{\epsilon}{2}. \quad (4.29)$$

By the definition of the pseudo-length and recalling from Lemma 4.3 that the map $\lambda \mapsto d_\lambda(z, w)$ is non-decreasing for all $z, w \in \Omega$, we have

$$\ell_{d_{\lambda_H}}(\gamma) \leq \ell_{d_\mu}(\gamma). \quad (4.30)$$

Combining (4.28), (4.29) and (4.30), we conclude

$$\ell_{d_{\lambda_H}}(\gamma) < d_{\lambda_H}(x, y) + \epsilon.$$

The proof is complete. \square

We are ready to prove Theorem 1.4.

Proof of Theorem 1.4 Obviously, (iii) in Theorem 1.3 \Rightarrow (iv) in Theorem 1.4. To see the converse, let $\lambda \geq 0$ be as in (iv). Given any x and $y, z \in N(x)$, where $N(x)$ is given in (iv) we need to show

$$u(y) - u(z) \leq d_\lambda(z, y).$$

By Proposition 4.1, we know (Ω, d_λ) is a pseudo-length space. Hence for any $\epsilon > 0$, there exists a curve $\gamma_\epsilon : [0, 1] \rightarrow \Omega$ joining z and y such that

$$\ell_{d_\lambda}(\gamma_\epsilon) \leq d_\lambda(z, y) + \epsilon. \quad (4.31)$$

Since $\gamma_\epsilon \subset \Omega$ is compact, we can find a finite set $\{t_i\}_{i=0}^n \subset [0, 1]$ satisfying

$$t_0 = 0, \quad t_n = 1, \quad \text{and } \gamma_\epsilon(t_{i+1}) \in N(\gamma_\epsilon(t_i)), \quad i = 0, \dots, n-1$$

where $N(\gamma_\epsilon(t_i))$ is the neighbourhood of $\gamma_\epsilon(t_i)$ in (iv). Hence by (iv) we have

$$u(\gamma_\epsilon(w_{i+1})) - u(\gamma_\epsilon(w_i)) \leq d_\lambda(\gamma_\epsilon(w_i), \gamma_\epsilon(w_{i+1})), \quad i = 0, \dots, i-1.$$

Summing the above inequalities from 0 to $n-1$, we have

$$\begin{aligned}
u(y) - u(z) &= u(\gamma_\epsilon(w_n)) - u(\gamma_\epsilon(w_0)) \leq \sum_{i=0}^{n-1} d_\lambda(\gamma_\epsilon(w_i), \gamma_\epsilon(w_{i+1})) \leq \sum_{i=0}^{n-1} \ell_{d_\lambda}(\gamma_\epsilon|_{[w_i, w_{i+1}]}) \\
&= \ell_{d_\lambda}(\gamma_\epsilon) \leq d_\lambda(y, z) + \epsilon
\end{aligned}$$

where in the last inequality we applied (4.31). Letting $\epsilon \rightarrow 0$ in the above inequality, we obtain (iii) in Theorem 1.3.

Finally, (1.8) is a direct consequence of (iv) \Leftrightarrow (i) and thanks to Lemma 4.3, the minimum in (1.8) is achieved. \square

Remark 4.4 The assumption $R'_\lambda > 0$ is needed in the proof of Proposition 4.1. Indeed, recall the construction of γ_ϵ in the proof of Proposition 4.1 below (4.17). To guarantee γ_ϵ is a continuous map, especially $\gamma_\epsilon([0, 1])$ is compact under the topology induced by d_{CC} , we need $\{d_\lambda(x, \cdot)\}_{x \in \Omega}$ induces the same topology as the one by d_{CC} . By Corollary 2.10, $R'_\lambda > 0$ can guarantee this.

Moreover, to show $\gamma_\epsilon \in \Omega$ in (4.13) in the proof of Proposition 4.1, for each $x \in \Omega$, we need the existence of $r_x > 0$, such that

$$B_{d_\lambda}^+(x, r_x) \subseteq \Omega. \quad (4.32)$$

Again, by Corollary 2.10, $R'_\lambda > 0$ can guarantee this. If $R'_\lambda > 0$ does not hold for some $\lambda > 0$, in Remark 2.11 (ii), the example shows that (4.32) may fail for some $x \in \Omega$.

5 McShane extensions and minimizers

In this section, we always suppose that the Hamiltonian $H(x, p)$ enjoys assumptions (H0)-(H3) and further that $\lambda_H = 0$.

Let $U \subseteq \Omega$ be any domain. Note that the restriction of d_λ in U may not have the pseudo-length property in U , and moreover, Theorem 1.3 with Ω replaced by U may not hold for the restriction of d_λ in U . Thus instead of d_λ , below we use intrinsic pseudo metrics $\{d_\lambda^U\}_{\lambda > 0}$ in U , which are defined via (1.4) with Ω replaced by U , that is,

$$d_\lambda^U(x, y) := \sup\{u(y) - u(x) : u \in \dot{W}_X^{1,\infty}(U), \|H(\cdot, Xu)\|_{L^\infty(U)} \leq \lambda\} \quad \forall x, y \in U \text{ and } \lambda \geq 0. \quad (5.1)$$

Obviously we have proved the following.

Corollary 5.1 *Theorem 1.3, Theorem 1.4 and Proposition 4.1 hold with Ω replaced by U and d_λ replaced by d_λ^U . In particular, d_λ^U has the pseudo-length property in U for all $\lambda \geq 0$.*

Observe that, apriori, d_λ^U is only defined in U but not in \bar{U} . Naturally, we extend $d_\lambda^U : U \times U \rightarrow [0, \infty)$ as a function $\tilde{d}_\lambda^U : \bar{U} \times \bar{U} \rightarrow [0, \infty]$ by

$$\tilde{d}_\lambda^U(x, y) = \liminf_{r \rightarrow 0} \{d_\lambda^U(z, w) \mid (z, w) \in U \times U, |(z, w) - (x, y)| \leq r\}.$$

Obviously, $\tilde{d}_\lambda^U(x, y) = d_\lambda^U(x, y)$ for all $(x, y) \in U \times U$, and \tilde{d}_λ^U is lower semicontinuous in $\bar{U} \times \bar{U}$, that is, for any $a \in \mathbb{R}$, the set

$$\{(x, y) \in \bar{U} \times \bar{U} \mid \tilde{d}_\lambda^U(x, y) > a\}$$

is open in \bar{U} . One may also note that it may happen that $d_\lambda^U(x, y) = +\infty$ for some $(x, y) \notin U \times U$. Below, for the sake of simplicity, we write \tilde{d}_λ^U as d_λ^U . We define d_{CC}^U by letting $H(x, p) = |p|$ in U and $\lambda = 1$ in (5.1).

The following property will be used later.

Lemma 5.2 *Let $U \subseteq \Omega$ be a subdomain and $\lambda \geq 0$.*

- (i) *For any $x, y \in \bar{U}$, we have $d_\lambda^U(x, y) \geq d_\lambda(x, y) \geq R'_\lambda d_{CC}(x, y)$.*
- (ii) *For any $x \in U$ and $y \in U$ with $d_{CC}(x, y) < d_{CC}(x, \partial U)$, we have $d_\lambda^U(x, y) \leq R_\lambda d_{CC}(x, y)$.*

(iii) For any $x \in U$, let $x^* \in \partial U$ be the point such that $d_{CC}(x, x^*) = d_{CC}(x, \partial U)$. Then

$$d_\lambda^U(x, x^*) \leq R_\lambda d_{CC}(x, x^*) < \infty.$$

(iv) For any $x \in \overline{U}$ and $y \in U$ we have

$$d_\lambda^U(x, z) \leq d_\lambda^U(x, y) + d_\lambda^U(y, z) \text{ and } d_\lambda^U(z, x) \leq d_\lambda^U(z, y) + d_\lambda^U(y, x) \quad \forall z \in \partial U.$$

(v) Given any $z \in \partial U$, if $d_\lambda^U(x, z) = \infty$ for some $x \in \overline{U}$, then $d_\lambda^U(y, z) = +\infty$ for any $y \in \overline{U}$.

(vi) Given any $x, y \in \overline{U} \times \overline{U}$ the map $\lambda \in [0, \infty) \mapsto d_\lambda^U(x, y) \in [0, \infty]$ is nondecreasing and for $0 < \lambda < \mu < \infty$,

$$d_\lambda^U(x, y) < \infty \text{ if and only if } d_\mu^U(x, y) < \infty.$$

As a consequence,

$$\overline{U}^* := \{y \in \overline{U} \mid d_\lambda^U(x, y) < \infty \text{ for some } x \in U \text{ and } \lambda > 0\}$$

is well-defined independent of the choice of $\lambda > 0$.

(vii) Given any $x, y \in \overline{U}^* \times \overline{U}^*$, the map $\lambda \in [0, \infty) \mapsto d_\lambda^U(x, y) \in [0, \infty]$ is right continuous.

Proof To see (i), for any $x, y \in U$, since the restriction $u|_U$ is a test function in the definition of $d_\lambda^U(x, y)$ whenever u is a test function in the definition of $d_\lambda(x, y)$, we have $d_\lambda^U(x, y) \geq d_\lambda(x, y)$. In general, given any $(x, y) \in (\overline{U} \times \overline{U}) \setminus (U \times U)$, for any $r > 0$ sufficiently small, we have $d_\lambda^U(z, w) \geq d_\lambda(z, w)$ whenever $z, w \in U$ and $|(z, w) - (x, y)| \leq r$. By the continuity of d_λ in $\Omega \times \Omega$, we have

$$\liminf_{r \rightarrow 0} \{d_\lambda^U(z, w) \mid z, w \in U \text{ and } |(z, w) - (x, y)| \leq r\} \geq d_\lambda(x, y),$$

that is, $d_\lambda^U(x, y) \geq d_\lambda(x, y)$. Recall that $d_\lambda(x, y) \geq R' d_{CC}(x, y)$ comes from Lemma 2.1.

To see (ii), given any $y \in U$ with $d_{CC}(x, y) < d_{CC}(x, \partial U)$, there is a geodesic γ with respect to d_{CC} connecting x and y so that $\gamma \subset B_{d_{CC}}(x, d_{CC}(x, \partial U))$. For any function $u \in \dot{W}_X^{1, \infty}(U)$ with $\|H(\cdot, Xu)\|_{L^\infty(U)} \leq \lambda$, we know that $\|Xu\|_{L^\infty(U)} \leq R_\lambda$. Let $U' \Subset U$ and $x, y \in U'$. Thanks to Proposition 2.6, we can find a sequence $\{u_k\}_{k \in \mathbb{N}} \subset C^\infty(U')$ such that $u_k \rightarrow u$ everywhere as $k \rightarrow \infty$ and $\|Xu_k\|_{L^\infty(U')} \leq R_\lambda + A_k(u)$ with $\lim_{k \rightarrow \infty} A_k(u) \rightarrow 0$. Since

$$\begin{aligned} u_k(y) - u_k(x) &= \int_\gamma Xu_k \cdot \dot{\gamma} \leq \int_\gamma |Xu_k| |\dot{\gamma}| \leq \int_\gamma (R_\lambda + A_k(u)) |\dot{\gamma}| \\ &= (R_\lambda + A_k(u)) d_{CC}(x, y) \text{ for all } k \in \mathbb{N}, \end{aligned}$$

we have

$$u(y) - u(x) = \lim_{k \rightarrow \infty} \{u_k(y) - u_k(x)\} \leq \lim_{k \rightarrow \infty} [R_\lambda + A_k(u)] d_{CC}(x, y) = R_\lambda d_{CC}(x, y).$$

Taking supremum in the above inequality over all such u , we have $d_\lambda^U(x, y) \leq R_\lambda d_{CC}(x, y)$.

To see (iii), given any $x \in U$, there exists $x^* \in \partial U$ such that $d_{CC}(x, x^*) = d_{CC}(x, \partial U)$. By (ii) and the definition of $d_\lambda^U(x, x^*)$, we know that $d_\lambda^U(x, x^*) \leq R_\lambda d_{CC}(x, x^*) < \infty$.

To get (iv), for any $x \in \overline{U}$, $y \in U$ and $z \in \partial U$, choose $x_k, z_k \in U$ such that $d_\lambda^U(x_k, y) \rightarrow d_\lambda^U(x, y)$ and $d_\lambda^U(y, z_k) \rightarrow d_\lambda^U(y, z)$ as $k \rightarrow \infty$. Since

$$d_\lambda^U(x_k, z_k) \leq d_\lambda^U(x_k, y) + d_\lambda^U(y, z_k),$$

letting $k \rightarrow \infty$ and by the lower-semicontinuous of d_λ^U we get

$$d_\lambda^U(x, z) \leq d_\lambda^U(x, y) + d_\lambda^U(y, z)$$

In a similar way, we also have

$$d_\lambda^U(z, x) \leq d_\lambda^U(z, y) + d_\lambda^U(y, x)$$

Note that (v) is a direct consequence of (iv).

We show (vi). The fact that $d_\lambda^U(x, y)$ is non-decreasing with respect to λ is obvious for any $x, y \in \bar{U}$. Assume $0 < \lambda < \mu < \infty$. Then $d_\mu^U(x, y) < +\infty$ implies $d_\lambda^U(x, y) < +\infty$. Conversely, if $d_\mu^U(x, y) = +\infty$, we show $d_\lambda^U(x, y) = +\infty$. This may happen if at least one of x and y lie in ∂U . Then for any $\{x_k\}_{k \in \mathbb{N}} \subset U$ and $\{y_k\}_{k \in \mathbb{N}} \subset U$ converging to x and y , it holds

$$\liminf_{k \rightarrow \infty} d_\mu^U(x_k, y_k) = +\infty$$

where we let $x_k \equiv x$ (resp. $y_k \equiv y$) if $x \in U$ (resp. $y \in U$). By (i) and (ii), we have for any $\lambda \leq \mu$

$$\liminf_{k \rightarrow \infty} d_\lambda^U(x_k, y_k) \geq R'_\lambda \liminf_{k \rightarrow \infty} d_{CC}(x_k, y_k) \geq \frac{R'_\lambda}{R_\mu} \liminf_{k \rightarrow \infty} d_\mu^U(x_k, y_k) = +\infty$$

where we recall that $R'_\lambda > 0$ for all $\lambda > 0$.

Finally, we show (vii). Since we only consider $x, y \in \bar{U}^*$, by an approximation argument, it is enough to show the right-continuity for $x, y \in U$. The proof is similar to the one of Lemma 4.3. We omit the details. The proof of Lemma 5.2 is complete. \square

Lemma 5.3 Suppose that $U \Subset \Omega$ and that V is a subdomain of U . For any $\lambda \geq 0$, one has

$$d_\lambda^U(x, y) \leq d_\lambda^V(x, y) \quad \forall x, y \in \bar{V}.$$

Conversely, given any $\lambda > 0$ and $x \in V$, there exists a neighborhood $N_\lambda(x) \Subset V$ of x such that

$$d_\lambda^U(x, y) = d_\lambda^V(x, y) \text{ and } d_\lambda^U(y, x) = d_\lambda^V(y, x) \text{ for any } y \in N_\lambda(x).$$

Proof For any $u \in \dot{W}_X^{1,\infty}(U)$ with $\|H(\cdot, Xu)\|_{L^\infty(U)} \leq \lambda$, we know that the restriction $u|_V \in \dot{W}_X^{1,\infty}(V)$ with $\|H(\cdot, Xu|_V)\|_{L^\infty(V)} \leq \lambda$. Hence by the definition of d_λ^U and d_λ^V , $d_\lambda^U(x, y) \leq d_\lambda^V(x, y)$ for all $x, y \in V$ and then for all $x, y \in \bar{V}$.

Conversely, we just show $d_\lambda^U(x, \cdot) = d_\lambda^V(x, \cdot)$ in some neighborhood $N(x)$. In a similar way, we can prove $d_\lambda^U(\cdot, x) = d_\lambda^V(\cdot, x)$ in some neighborhood $N(x)$.

By Lemma 5.2 (i), one has $d_\lambda^V(x, y) \geq R'_\lambda d_{CC}(x, y)$ for any $x, y \in V$. Thus for any $r > 0$, $d_\lambda^V(x, y) \leq r$ implies $d_{CC}(x, y) \leq r/R'_\lambda$. Given any $x \in V$ and $0 < r < d_{CC}(x, \partial V)/R'_\lambda$, we therefore have

$$N_r(x) := \{y \in V \mid d_\lambda^V(x, y) \leq r\} \subset B_{d_{CC}}(x, r/R'_\lambda) \Subset V.$$

Define $u_{x,r} : \Omega \rightarrow \mathbb{R}$ by

$$u_{x,r}(z) := \min\{d_\lambda^V(x, z), r\} \quad \forall z \in \Omega.$$

If $r < d_{CC}(x, \partial V)R'_\lambda/4$, we claim that

$$u_{x,r} \in \text{Lip}_{d_{CC}}(\Omega) \quad \text{with} \quad H(z, Xu_{x,r}(z)) \leq \lambda \quad \text{for almost all } z \in \Omega. \quad (5.2)$$

Assume claim (5.2) holds for the moment. By (5.2), we are able to take $u_{x,r}|_U$ as a test function in d_λ^U so that

$$u_{x,r}(z) - u_{x,r}(w) \leq d_\lambda^U(w, z) \quad \forall (w, z) \in U \times U.$$

On the other hand, for any $y \in N_r(x)$, since $d_\lambda^V(x, y) = u_{x,r}(y) - u_{x,r}(x)$, we get $d_\lambda^V(x, y) \leq d_\lambda^U(x, y)$ as desired.

Finally, we prove the claim (5.2).

Proof of the claim (5.2). First, by Lemma 3.3 and Lemma 3.4, the restriction $u_{x,r}|_V$ of $u_{x,r}$ in V belongs to $\dot{W}_X^{1,\infty}(V)$ with $H(z, Xu_{x,r}|_V(z)) \leq \lambda$ almost everywhere in V , and hence

$$u_{x,r}|_V(z) - u_{x,r}|_V(w) \leq d_\lambda^V(w, z) \quad \forall (w, z) \in V \times V. \quad (5.3)$$

Next, we show $u \in \text{Lip}_{d_{CC}}(\Omega)$. Given any $w, z \in \Omega$, we consider 3 cases separately.

Case 1. $w \in \overline{B_{d_{CC}}(x, r/R'_\lambda)}$ and $z \in \overline{B_{d_{CC}}(x, r/R'_\lambda)}$. We have $z \in \overline{B_{d_{CC}}(w, 2r/R'_\lambda)}$. Since

$$d_{CC}(w, \partial V) \geq d_{CC}(x, \partial V) - d_{CC}(x, w) \geq \frac{4r}{R'_\lambda} - \frac{r}{R'_\lambda} = \frac{3r}{R'_\lambda} > \frac{2r}{R'_\lambda} \geq d_{CC}(w, z),$$

by Lemma 5.2 (ii), $d_\lambda^V(w, z) \leq R_\lambda d_{CC}(w, z)$, which combined with (5.3), gives

$$u_{x,r}(z) - u_{x,r}(w) \leq R_\lambda d_{CC}(w, z).$$

Case 2. $w, z \notin B_{d_{CC}}(x, r/R'_\lambda)$. Then $w, z \in \Omega \setminus B_{d_{CC}}(x, r/R'_\lambda)$, since $u_{x,r}$ is constant r in $\Omega \setminus B_{d_{CC}}(x, r/R'_\lambda)$, we know that

$$u_{x,r}(z) - u_{x,r}(w) = r - r = 0 \leq R_\lambda d_{CC}(w, z).$$

Case 3. $w \in \overline{B_{d_{CC}}(x, r/R'_\lambda)}$ and $z \notin \overline{B_{d_{CC}}(x, r/R'_\lambda)}$. Then for any $\epsilon > 0$, there exists a curve $\gamma_\epsilon : [0, 1] \rightarrow \Omega$ joining z and w such that

$$\ell_{d_{CC}}(\gamma_\epsilon) \leq d_{CC}(w, z) + \epsilon$$

and there exists $t \in [0, 1]$ such that $\gamma_\epsilon(t) \in \partial B_{d_{CC}}(x, r/R'_\lambda)$. Thus using Case 1 and Case 2, we deduce

$$\begin{aligned} u_{x,r}(z) - u_{x,r}(w) &= u_{x,r}(z) - u_{x,r}(\gamma_\epsilon(t)) + u_{x,r}(\gamma_\epsilon(t)) - u_{x,r}(w) \\ &\leq R_\lambda d_{CC}(z, \gamma_\epsilon(t)) + R_\lambda d_{CC}(\gamma_\epsilon(t), w) \\ &\leq R_\lambda \ell_{d_{CC}}(\gamma_\epsilon|_{[0,t]}) + R_\lambda \ell_{d_{CC}}(\gamma_\epsilon|_{[t,1]}) \\ &\leq R_\lambda [d_{CC}(w, z) + \epsilon]. \end{aligned}$$

Letting $\epsilon \rightarrow 0$ in the above inequality, we conclude

$$u_{x,r}(z) - u_{x,r}(w) \leq R_\lambda d_{CC}(w, z).$$

Finally, by Lemma 2.7, $u_{x,r} \in \dot{W}_X^{1,\infty}(\Omega)$. Note that $Xu_{x,r} = 0$ in $\Omega \setminus \overline{B_{d_{CC}}(x, r/R'_\lambda)}$ implies $H(z, Xu_{x,r}(z)) = 0$ almost everywhere. Therefore recalling $H(z, Xu_{x,r}(z)) \leq \lambda$ almost everywhere in V and $\Omega = V \cup (\Omega \setminus \overline{B_{d_{CC}}(x, r/R'_\lambda)})$, we conclude $H(z, Xu_{x,r}(z)) \leq \lambda$ almost everywhere in Ω . \square

Lemma 5.4 Suppose that $U \Subset \Omega$ and that $V = U \setminus \{x_i\}_{1 \leq i \leq m}$ for some $m \in \mathbb{N}$ and $\{x_i\}_{1 \leq i \leq m} \subset U$. Then for any $\lambda \geq 0$, one has

$$d_\lambda^V(x, y) = d_\lambda^U(x, y) \text{ for all } x, y \in \overline{U}.$$

Proof Obviously $d_\lambda^U \leq d_\lambda^V$ in \overline{U} . Conversely, we show $d_\lambda^V \leq d_\lambda^U$ in \overline{U} . First, by an approximation argument, it suffices to consider $x, y \in U$. By the right continuity of $\lambda \in [0, \infty) \rightarrow d_\lambda^U(x, y)$ for any $x, y \in \overline{U}^*$, up to considering $d_{\mu+\epsilon}^U$ for sufficiently small $\epsilon > 0$, we may assume that $\mu > 0$. For any $\lambda > 0$, by the pseudo-length property of d_λ^U as in Proposition 4.1, it suffices to prove that for any curve $\gamma : [a, b] \rightarrow U$, one has

$$d_\lambda^V(\gamma(a), \gamma(b)) \leq \ell_{d_\lambda^U}(\gamma). \quad (5.4)$$

We consider the following 3 cases.

Case 1. $\gamma((a, b)) \subset V$ and $\gamma(\{a, b\}) \subset V$, that is, $\gamma([a, b]) \subset V$. Recall from Lemma 5.3 that, for each $x \in V$, there exists a neighborhood $N(x)$ such that

$$d_\lambda^V(x, y) = d_\lambda^U(x, y) \quad \forall y \in N(x).$$

Since $\gamma \subset \cup_{t \in [a, b]} N(\gamma(t))$, we can find $a = t_0 = 0 < t_1 < \dots < t_m = b$ such that $\gamma \subset \cup_{i=0}^m N(\gamma(t_i))$ and $\gamma([t_i, t_{i+1}]) \subset N(\gamma(t_i))$. By the triangle inequality, one has

$$d_\lambda^V(\gamma(a), \gamma(b)) \leq \sum_{i=0}^{m-1} d_\lambda^U(\gamma(t_i), \gamma(t_{i+1})) \leq \ell_{d_\lambda^U}(\gamma).$$

Case 2. $\gamma((a, b)) \subset V$ and $\gamma(\{a, b\}) \not\subset V$. Applying Case 1 to $\gamma|_{[a+\epsilon, b-\epsilon]}$ for sufficiently small $\epsilon > 0$, we get

$$d_\lambda^V(\gamma(a), \gamma(b)) \leq \lim_{\epsilon \rightarrow 0} d_\lambda^V(\gamma(a+\epsilon), \gamma(b-\epsilon)) \leq \liminf_{\epsilon \rightarrow 0} \ell_{d_\lambda^U}(\gamma|_{[a+\epsilon, b-\epsilon]}) \leq \ell_{d_\lambda^U}(\gamma).$$

Case 3. $\gamma((a, b)) \not\subset V$. Without loss of generality, for each x_i , there is at most one $t \in (a, b)$ such that $\gamma(t) = x_i$. Indeed, let t^\pm as the maximum/minimum $s \in (a, b)$ such that $\gamma(s) = x_i$. Then $a \leq t^- \leq t^+ \leq b$. If $t^- < t^+$, we consider $\gamma_1 : [a, b - (t^+ - t^-)] \rightarrow U$ with $\gamma_1(t) = \gamma(t)$ for $t \in [a, t^-]$, and $\gamma_1(t) = \gamma(t - (t^+ - t^-))$ for $t \in [t^+, b]$. Then $\ell_{d_\lambda^U}(\gamma_1) \leq \ell_{d_\lambda^U}(\gamma)$. Repeating this procedure for x_2, \dots, x_m in order, we may get a new curve η such that for each x_i , there is at most one $t \in (a, b)$ such that $\gamma(t) = x_i$ and $\ell_{d_\lambda^U}(\eta) \leq \ell_{d_\lambda^U}(\gamma)$.

Denote by $\{a_j\}_{j=0}^s$ with $a = a_0 < a_1 < \dots < a_s = b$ such that $\gamma(\{a_1, \dots, a_{s-1}\}) \subset U \setminus V = \{x_i\}_{1 \leq i \leq m}$ and $\gamma([a, b] \setminus \{a_1, \dots, a_{s-1}\}) \subset V$. Applying Case 2 to $\gamma|_{[a_j, a_{j+1}]}$ for all $0 \leq j \leq s-1$, we obtain

$$d_\lambda^V(\gamma(a), \gamma(b)) \leq \sum_{j=0}^{s-1} d_\lambda^V(\gamma(a_j), \gamma(a_{j+1})) \leq \sum_{j=0}^{s-1} \ell_{d_\lambda^U}(\gamma|_{[a_j, a_{j+1}]}) \leq \ell_{d_\lambda^U}(\gamma)$$

as desired. The proof is complete. \square

For any $g \in C^0(\partial U)$, write

$$\mu(g, \partial U) := \inf \{ \lambda \geq 0 \mid g(y) - g(x) \leq d_\lambda^U(x, y) \quad \forall x, y \in \partial U \}.$$

The following lemma says the infimum can be reached.

Lemma 5.5 *We have*

$$\mu(g, \partial U) = \min \{ \lambda \geq 0 \mid g(y) - g(x) \leq d_\lambda^U(x, y) \quad \forall x, y \in \partial U \cap \overline{U}^* \}.$$

Proof First, if $x \in \partial U \setminus \overline{U}^*$ or $y \in \partial U \setminus \overline{U}^*$, we have $d_\lambda^U(x, y) = \infty$. Hence $g(y) - g(x) \leq d_\lambda^U(x, y)$ holds trivially, which implies that

$$\mu(g, \partial U) = \inf \{ \lambda \geq 0 \mid g(y) - g(x) \leq d_\lambda^U(x, y) \quad \forall x, y \in \partial U \cap \overline{U}^* \}.$$

Thanks to Lemma 5.2(vii), we finish the proof. \square

If $\mu = \mu(g, \partial U) < \infty$, we define

$$\mathcal{S}_{g;U}^+(x) := \inf_{y \in \partial U} \{g(y) + d_\mu^U(y, x)\} \quad \text{and} \quad \mathcal{S}_{g;U}^-(x) := \sup_{y \in \partial U} \{g(y) - d_\mu^U(x, y)\} \quad \forall x \in \overline{U}.$$

Note that $\mathcal{S}_{g;U}^\pm$ serve as “McShane” extensions of g in U .

Lemma 5.6 *If $\mu = \mu(g, \partial U) < \infty$, then we have*

- (i) $\mathcal{S}_{g;U}^\pm \in \dot{W}_X^{1,\infty}(U) \cap C^0(\overline{U})$ with $\mathcal{S}_{g;U}^\pm = g$ on ∂U ;
- (ii) for any $x, y \in \overline{U}$,
$$\mathcal{S}_{g;U}^\pm(y) - \mathcal{S}_{g;U}^\pm(x) \leq d_\mu^U(x, y); \tag{5.5}$$
- (iii) $\|H(\cdot, X\mathcal{S}_{g;U}^\pm)\|_{L^\infty(U)} \leq \mu$.

Proof By Corollary 5.1, (ii) implies (iii) and $\mathcal{S}_{g;U}^\pm \in \dot{W}_X^{1,\infty}(U)$. Below we show $\mathcal{S}_{g;U}^\pm = g$ on ∂U , (ii) and $\mathcal{S}_{g;U}^\pm \in C^0(\overline{U})$ in order.

Proof for $\mathcal{S}_{g;U}^\pm = g$ on ∂U . For any $x \in \partial U$, by definition we have

$$\mathcal{S}_{g;U}^-(x) \geq g(x) \geq \mathcal{S}_{g;U}^+(x).$$

Conversely, for $y \in \partial U$, one has

$$g(y) - d_\mu^U(x, y) \leq g(x) \leq g(y) + d_\mu^U(y, x)$$

and hence $\mathcal{S}_{g;U}^-(x) \leq g(x) \leq \mathcal{S}_{g;U}^+(x)$ as desired.

Proof of (ii). We only prove (ii) for $\mathcal{S}_{g;U}^-$; the proof for $\mathcal{S}_{g;U}^+$ is similar. For $x, y \in \partial U$, by the definition of μ one has

$$\mathcal{S}_{g;U}^-(y) - \mathcal{S}_{g;U}^-(x) = g(y) - g(x) \leq d_\mu^U(x, y).$$

For $x \in U$ and $y \in \partial U$, by definition

$$\mathcal{S}_{g;U}^-(x) \geq g(y) - d_\mu^U(x, y) = \mathcal{S}_{g;U}^-(y) - d_\mu^U(x, y)$$

and hence

$$\mathcal{S}_{g;U}^-(y) - \mathcal{S}_{g;U}^-(x) \leq d_\mu^U(x, y).$$

For $x \in \overline{U}$ and $y \in U$, by Lemma 5.2(iv), we have

$$d_\mu^U(x, z) \leq d_\mu^U(x, y) + d_\mu^U(y, z) \quad \forall z \in \partial U.$$

One then has

$$\begin{aligned} \mathcal{S}_{g;U}^-(y) &= \sup_{z \in \partial U} \{g(z) - d_\mu^U(y, z)\} \\ &\leq \sup_{z \in \partial U} \{g(z) - d_\mu^U(x, z) + d_\mu^U(x, y)\} \end{aligned}$$

$$\leq \mathcal{S}_{g;U}^-(x) + d_\mu^U(x, y),$$

and hence

$$\mathcal{S}_{g;U}^-(y) - \mathcal{S}_{g;U}^-(x) \leq d_\mu^U(x, y).$$

Proof of $\mathcal{S}_{g;U}^\pm \in C^0(\overline{U})$. We only consider $\mathcal{S}_{g;U}^- \in C^0(\overline{U})$; the proof for $\mathcal{S}_{g;U}^+ \in C^0(\overline{U})$ is similar. It suffices to show that for any $x \in \partial U$ and a sequence $\{x_j\} \subset U$ converging to x ,

$$\lim_{j \rightarrow \infty} \mathcal{S}_{g;U}^-(x_j) = \mathcal{S}_{g;U}^-(x) = g(x).$$

Choosing $x_j^* \in \partial U$ such that $d_{CC}(x_j, x_j^*) = \text{dist}_{d_{CC}}(x_j, \partial U)$, one has

$$d_{CC}(x_j, x_j^*) \leq d_{CC}(x_j, x) \rightarrow 0, \quad \text{and hence } d_{CC}(x_j^*, x) \rightarrow 0.$$

Thanks to Lemma 5.2(iii) with $x = x_j$ and $x^* = x_j^*$ therein, we deduce

$$d_\mu^U(x_j, x_j^*) \leq R_\mu d_{CC}(x_j, x_j^*) \rightarrow 0.$$

Since

$$\mathcal{S}_{g;U}^-(x_j) \geq g(x_j^*) - d_\mu^U(x_j, x_j^*),$$

by the continuity of g , we have

$$\liminf_{j \rightarrow \infty} \mathcal{S}_{g;U}^-(x_j) \geq g(x).$$

Assume that

$$\liminf_{j \rightarrow \infty} \mathcal{S}_{g;U}^-(w_j) \geq g(x) + 2\epsilon \quad \text{for some } \{w_j\}_{j \in \mathbb{N}} \subset U \text{ with } w_j \rightarrow x \text{ and some } \epsilon > 0.$$

By the definition we can find $z_j \in \partial U$ such that

$$\mathcal{S}_{g;U}^-(w_j) - \epsilon \leq g(z_j) - d_\mu^U(w_j, z_j).$$

Thus for $j \in \mathbb{N}$ sufficiently large, we have

$$g(z_j) - d_\mu^U(w_j, z_j) \geq g(x) + \epsilon.$$

Up to some subsequence, we assume that $z_j \rightarrow z \in \partial U$. Note that

$$d_\mu^U(x, z) \leq \liminf_{j \rightarrow \infty} d_\mu^U(w_j, z_j)$$

By the continuity of g , we conclude

$$g(z) - d_\mu^U(x, z) \geq g(x) + \epsilon,$$

which is a contradiction with the definition of μ and $\mu < \infty$. \square

Write

$$\mathbf{I}(g, U) = \inf \{ \|H(\cdot, Xu)\|_{L^\infty(U)} \mid u \in \dot{W}_X^{1,\infty}(U) \cap C^0(\overline{U}), u|_{\partial U} = g \}. \quad (5.6)$$

A function $u : \overline{U} \rightarrow \mathbb{R}$ is called as a minimizer for $\mathbf{I}(g, U)$ if

$$u \in \dot{W}_X^{1,\infty}(U) \cap C^0(\overline{U}), u|_{\partial U} = g \quad \text{and} \quad \|H(\cdot, Xu)\|_{L^\infty(U)} = \mathbf{I}(g, U).$$

We have the following existence and properties for minimizers.

Lemma 5.7 For any $g \in C(\partial U)$ with $\mu(g, \partial U) < \infty$, we have the following:

- (i) We have $\mu(g, \partial U) = \mathbf{I}(g, U)$. Both of $\mathcal{S}_{g;U}^{\pm}$ are minimizers for $\mathbf{I}(g, U)$.
- (ii) If u is a minimizer for $\mathbf{I}(g, U)$ then

$$\mathcal{S}_{g;U}^{-} \leq u \leq \mathcal{S}_{g;U}^{+} \text{ in } \overline{U} \text{ and } \|H(x, Xu)\|_{L^{\infty}(U)} = \mathbf{I}(g, U) = \mu(g, \partial U).$$

- (iii) If u, v are minimizer for $\mathbf{I}(g, U)$, then $tu + (1 - t)v$ with $t \in (0, 1)$, $\max\{u, v\}$ and $\min\{u, v\}$ are minimizers for $\mathbf{I}(g, U)$.

Proof (i) Since $\mu(g, \partial U) < \infty$, then by Lemma 5.6, we know that $\mathcal{S}_{g;U}^{\pm}$ satisfies the condition required in (5.6) and hence

$$\mathbf{I}(g, U) \leq \|H(\cdot, X\mathcal{S}_{g;U}^{\pm})\|_{L^{\infty}(U)} \leq \mu. \quad (5.7)$$

Below we show that $\mu(g, \partial U) \leq \mathbf{I}(g, U)$. Note that combining this and (5.7) we know that

$$\mathbf{I}(g, U) = \|H(\cdot, X\mathcal{S}_{g;U}^{\pm})\|_{L^{\infty}(U)} = \mu$$

and moreover, $\mathcal{S}_{g;U}^{\pm}$ are minimizers for $\mathbf{I}(g, U)$.

For any $\lambda > \mathbf{I}(g, U)$, there is a function $u \in \dot{W}_X^{1,\infty}(U) \cap C^0(\overline{U})$ with $u = g$ on ∂U such that $\|H(x, Xu)\|_{L^{\infty}(U)} \leq \lambda$. By Corollary 5.1,

$$u(y) - u(x) \leq d_{\lambda}^U(x, y), \quad \forall x, y \in U.$$

By the continuity of u in \overline{U} and the definition of d_{λ}^U in $\overline{U} \times \overline{U}$ we have

$$g(x) - g(y) \leq d_{\lambda}^U(x, y) \text{ for all } x, y \in \partial U.$$

Thus $\mu(g, \partial U) \leq \mathbf{I}(g, U)$.

- (ii) If u is a minimizer for $\mathbf{I}(g, U)$, one has

$$\|H(\cdot, Xu)\|_{L^{\infty}(U)} = \mathbf{I}(g, U) = \mu.$$

By Corollary 5.1, $u(y) - u(x) \leq d_{\mu}^U(x, y)$ for any $x, y \in U$ and hence, by the continuity of u and the definition of d_{μ}^U , for all $x, y \in \overline{U}$. Since $u = g$ on ∂U , for any $x \in U$, one has $g(y) - d_{\mu}^U(x, y) \leq u(x)$ for any $y \in \partial U$, which yields $\mathcal{S}_{g;U}^{-}(x) \leq u(x)$. By a similar argument, one also has $u \leq \mathcal{S}_{g;U}^{+}$ in U as desired.

- (iii) Suppose that u_1, u_2 are minimizers for $\mathbf{I}(g, U)$. Set

$$u_{\eta} := (1 - \eta)u_1 + \eta u_2 \text{ for any } \eta \in [0, 1].$$

Then $u_{\eta} \in \dot{W}_X^{1,\infty}(U) \cap C^0(\overline{U})$, and $u_{\eta} = g$ on ∂U , and by (H1),

$$H(\cdot, Xu_{\eta}(\cdot)) \leq \max_{i=1,2} \{H(\cdot, Xu_i(\cdot))\} \leq \mu \text{ a.e. on } U.$$

This, combined with the definition of $\mathbf{I}(g, U)$, implies that u_{η} is also a minimizer for $\mathbf{I}(g, U)$.

Finally, note that

$$\max\{u_1, u_2\} \in \dot{W}_X^{1,\infty}(U) \cap C^0(\overline{U}), \max\{u_1, u_2\} = g \text{ on } \partial U.$$

By Lemma 3.2, one has

$$H(\cdot, X \max\{u_1, u_2\}(\cdot)) \leq \max_{i=1,2} \{H(\cdot, Xu_i(\cdot))\} \leq \mu \text{ a.e. on } U.$$

We know that

$$\mathbf{I}(g, U) \leq \mathcal{F}(\max\{u_1, u_2\}, U) \leq \max\{\mathcal{F}(u, U), \mathcal{F}(v, U)\} = \mathbf{I}(g, U),$$

that is, $\max\{u_1, u_2\}$ is a minimizer for $\mathbf{I}(g, U)$. Similarly, $\min\{u_1, u_2\}$ is a minimizer for $\mathbf{I}(g, U)$. \square

We have the following improved regularity for Mschane extension via d_λ .

Lemma 5.8 Suppose that $U \Subset \Omega$ and that $V \subset U$ is a subdomain. If $g : \partial V \rightarrow \mathbb{R}$ satisfies

$$g(y) - g(x) \leq d_\lambda^U(x, y) \quad \forall x, y \in \partial V \quad (5.8)$$

for some $\lambda \geq 0$, then $\mu(g, \partial U) \leq \lambda$ and

$$S_{g,V}^\pm(y) - S_{g,V}^\pm(x) \leq d_\lambda^U(x, y) \quad \forall x, y \in \overline{V}. \quad (5.9)$$

Proof Since $d_\lambda^U \leq d_\lambda^V$ in $\overline{V} \times \overline{V}$, we know that

$$g(y) - g(x) \leq d_\lambda^V(x, y) \quad \forall x, y \in \partial V$$

and hence

$$\mu(g, \partial V) = \min\{\eta \geq 0 \mid g(y) - g(x) \leq d_\eta^V(x, y) \quad \forall x, y \in \partial V\} \leq \lambda.$$

To prove (5.9), by the pseudo-length property of d_λ^U as in Corollary 5.1, it suffices to prove that for any curve $\gamma : [a, b] \rightarrow \Omega$ with $\gamma(a), \gamma(b) \in \overline{V}$, one has

$$S_{g,V}^\pm(\gamma(b)) - S_{g,V}^\pm(\gamma(a)) \leq \ell_{d_\lambda^U}(\gamma). \quad (5.10)$$

We consider the following 4 cases.

Case 1. $\gamma((a, b)) \subset V$ and $\gamma(\{a, b\}) \subset V$, that is, $\gamma([a, b]) \subset V$. Noting $\mu = \mu(g, \partial V) \leq \lambda$, one then has $d_\mu^U \leq d_\lambda^V$ in $V \times V$. From the definition of $S_{g,V}^\pm$, it follows

$$S_{g,V}^\pm(y) - S_{g,V}^\pm(x) \leq d_\mu^V(x, y) \quad \forall x, y \in V.$$

Recall from Lemma 5.3 that, for each $x \in V$, there exists a neighborhood $N(x) \Subset V$ such that

$$d_\lambda^V(x, y) = d_\lambda^U(x, y) \quad \forall y \in N(x).$$

We therefore have

$$S_{g,V}^\pm(y) - S_{g,V}^\pm(x) \leq d_\lambda^U(x, y) \quad \forall x \in U, y \in N(x). \quad (5.11)$$

Since $\gamma \subset \cup_{t \in [a, b]} N(\gamma(t))$, we can find $a = t_0 = 0 < t_1 < \dots < t_m = b$ such that $\gamma \subset \cup_{i=0}^m N(\gamma(t_i))$ and $\gamma([t_i, t_{i+1}]) \subset N(\gamma(t_i))$. Applying (5.11) to $\gamma(t_i)$ and $\gamma(t_{i+1})$, we have

$$S_{g,V}^\pm(\gamma(t_{i+1})) - S_{g,V}^\pm(\gamma(t_i)) \leq d_\lambda^U(\gamma(t_i), \gamma(t_{i+1})).$$

Thus

$$\begin{aligned} S_{g,V}^\pm(\gamma(b)) - S_{g,V}^\pm(\gamma(a)) &= \sum_{i=0}^{m-1} [S_{g,V}^\pm(\gamma(t_{i+1})) - S_{g,V}^\pm(\gamma(t_i))] \\ &\leq \sum_{i=0}^{m-1} d_\lambda^U(\gamma(t_i), \gamma(t_{i+1})) \leq \ell_{d_\lambda^U}(\gamma). \end{aligned}$$

Case 2. $\gamma((a, b)) \subset V$ and $\gamma(\{a, b\}) \not\subset V$. Applying Case 1 to $\gamma|_{[a+\epsilon, b-\epsilon]}$ for sufficiently small $\epsilon > 0$, we get

$$\begin{aligned} S_{g,V}^{\pm}(\gamma(b)) - S_{g,V}^{\pm}(\gamma(a)) &= \lim_{\epsilon \rightarrow 0} [S_{g,V}^{\pm}(\gamma(b-\epsilon)) - S_{g,V}^{\pm}(\gamma(a+\epsilon))] \\ &\leq \liminf_{\epsilon \rightarrow 0} \ell_{d_{\lambda}^U}(\gamma|_{[a+\epsilon, b-\epsilon]}) \leq \ell_{d_{\lambda}^U}(\gamma). \end{aligned}$$

Case 3. $\gamma((a, b)) \not\subset V$ and $\gamma(\{a, b\}) \subset V$. Set

$$t_* = \min\{t \in [a, b], \gamma(t) \notin V\} \text{ and } t^* = \max\{t \in [0, 1], \gamma(t) \notin V\}.$$

Then $\gamma(t_*) \in \partial V$ and $\gamma(t^*) \in \partial V$. Write

$$\begin{aligned} S_{g,V}^{\pm}(\gamma(b)) - S_{g,V}^{\pm}(\gamma(a)) &= S_{g,V}^{\pm}(\gamma(b)) - S_{g,V}^{\pm}(\gamma(t^*)) + S_{g,V}^{\pm}(\gamma(t^*)) \\ &\quad - S_{g,V}^{\pm}(\gamma(t_*)) + S_{g,V}^{\pm}(\gamma(t_*)) - S_{g,V}^{\pm}(\gamma(a)). \end{aligned}$$

Note that

$$S_{g,V}^{\pm}(\gamma(t^*)) - S_{g,V}^{\pm}(\gamma(t_*)) = g(\gamma(t^*)) - g(\gamma(t_*)) \leq d_{\lambda}^U(\gamma(t_*), \gamma(t^*)) \leq \ell_{d_{\lambda}^U}(\gamma|_{[a, t_*]}).$$

By this, and applying Case 2 to $\gamma|_{[a, t_*]}$ and $\gamma|_{[t^*, b]}$, we obtain

$$S_{g,V}^{\pm}(\gamma(b)) - S_{g,V}^{\pm}(\gamma(a)) \leq \ell_{d_{\lambda}^U}(\gamma|_{[t^*, b]}) + \ell_{d_{\lambda}^U}(\gamma|_{[t_*, t^*]}) + \ell_{d_{\lambda}^U}(\gamma|_{[a, t_*]}) = \ell_{d_{\lambda}^U}(\gamma).$$

Case 4. $\gamma((a, b)) \not\subset V$ and $\gamma(\{a, b\}) \not\subset V$. If $\gamma(\{a, b\}) \subset \partial V$, then

$$S_{g,V}^{\pm}(\gamma(b)) - S_{g,V}^{\pm}(\gamma(a)) = g(\gamma(b)) - g(\gamma(a)) \leq d_{\lambda}^U(\gamma(b), \gamma(a)) \leq \ell_{d_{\lambda}^U}(\gamma).$$

If $\gamma(a) \in V$ and $\gamma(b) \in \partial V$, set $s^* = \min\{s \in [a, b] \mid \gamma(s) \in \partial V\}$. Obviously $a < s^* \leq b$, and we can find a sequence of $\epsilon_i > 0$ so that $\epsilon_i \rightarrow 0$ as $i \rightarrow \infty$ and $\gamma(s^* - \epsilon_i) \in V$. Write

$$S_{g,V}^{\pm}(\gamma(b)) - S_{g,V}^{\pm}(\gamma(a)) = S_{g,V}^{\pm}(\gamma(b)) - S_{g,V}^{\pm}(\gamma(s^*)) + S_{g,V}^{\pm}(\gamma(s^*)) - S_{g,V}^{\pm}(\gamma(a)).$$

Since $\gamma(s^*), \gamma(b) \in \partial V$, we have

$$S_{g,V}^{\pm}(\gamma(b)) - S_{g,V}^{\pm}(\gamma(s^*)) = g(\gamma(b)) - g(\gamma(s^*)) \leq d_{\lambda}^U(\gamma(b), \gamma(s^*)) \leq \ell_{d_{\lambda}^U}(\gamma|_{[s^*, b]}).$$

Applying Case 3 to $\gamma|_{[a, s^* - \epsilon_i]}$, one has

$$\begin{aligned} S_{g,V}^{\pm}(\gamma(s^*)) - S_{g,V}^{\pm}(\gamma(a)) &= \lim_{i \rightarrow \infty} [S_{g,V}^{\pm}(\gamma(s^* - \epsilon_i)) - S_{g,V}^{\pm}(\gamma(a))] \\ &\leq \lim_{i \rightarrow \infty} \ell_{d_{\lambda}^U}(\gamma|_{[a, s^* - \epsilon_i]}) \leq \ell_{d_{\lambda}^U}(\gamma|_{[a, s^*]}). \end{aligned}$$

We therefore have

$$S_{g,V}^{\pm}(\gamma(b)) - S_{g,V}^{\pm}(\gamma(a)) \leq \ell_{d_{\lambda}^U}(\gamma|_{[s^*, b]}) + \ell_{d_{\lambda}^U}(\gamma|_{[a, s^*]}) = \ell_{d_{\lambda}^U}(\gamma).$$

If $\gamma(a) \in \partial V$ and $\gamma(b) \in V$, we could prove in a similar way. Thus (5.10) holds and the proof is complete. \square

The following will be used in Sect. 6. Let $U \Subset \Omega$ and $u \in \dot{W}_X^{1,\infty}(U) \cap C^0(\overline{U})$ satisfying

$$u(y) - u(x) \leq d_{\mu}^U(x, y) \quad \forall x, y \in \overline{U} \quad (5.12)$$

for some $0 \leq \mu < \infty$. Given any subdomain $V \subset U$, write $h = u|_{\partial V}$ as the restriction of u in \overline{V} . Since $d_{\mu}^U \leq d_{\lambda}^V$ in $\overline{V} \times \overline{V}$ as given in Lemma 5.3, one has

$$u(y) - u(x) \leq d_{\mu}^V(x, y) \quad \forall x, y \in \partial V$$

and hence $\mu(u|_{\partial V}, \partial V) \leq \mu$. Denote by $\mathcal{S}_{u|_{\partial V}, V}^{\mp}$ the McShane extension of $u|_{\partial V}$ in V . Define

$$u^{\pm} := \begin{cases} \mathcal{S}_{u|_{\partial V}, V}^{\pm} & \text{in } V \\ u & \text{in } \overline{U} \setminus V. \end{cases} \quad (5.13)$$

Lemma 5.9 *Under the assumption (5.12) for some $0 \leq \mu < \infty$, the functions u^{\pm} defined in (5.13) are continuous in \overline{U} and satisfy*

$$u^{\pm}(y) - u^{\pm}(x) \leq d_{\mu}^U(x, y) \quad \forall x, y \in \overline{U}. \quad (5.14)$$

In particular, $u^{\pm} \in \dot{W}_X^{1, \infty}(U) \cap C^0(\overline{U})$ and $\|H(\cdot, Xu^{\pm})\|_{L^{\infty}(U)} \leq \mu$.

Proof We only prove Lemma 5.9 for u^+ ; the proof of Lemma 5.9 for u^- is similar. By Lemma 5.6, $\mathcal{S}_{h, V}^+ \in C^0(\overline{V})$ and $\mathcal{S}_{h, V}^+ = u$ on ∂V . These, together with $u \in C^0(\overline{U})$ imply that $u^+ \in C^0(\overline{U})$. Moreover, by Corollary 5.1, if u^+ satisfies (5.14), then $u^+ \in \dot{W}_X^{1, \infty}(U)$ and $\|H(\cdot, Xu^{\pm})\|_{L^{\infty}(U)} \leq \mu$. Below we prove (5.14) for u^+ via the following 3 cases. By the right continuity of $\lambda \in [0, \infty) \rightarrow d_{\lambda}^U(x, y)$ for any $x, y \in \overline{U}$, up to considering $d_{\mu+\epsilon}^U$ for sufficiently small $\epsilon > 0$, we may assume that $\mu > 0$.

Case 1. $x, y \in \overline{U} \setminus V$. By (5.12) we have

$$u^+(y) - u^+(x) = u(y) - u(x) \leq d_{\mu}^U(x, y).$$

Case 2. $x \in \overline{V}$ and $y \in V$ or $x \in V$ and $y \in \overline{V}$. Applying Lemma 5.8 with $(U, d_{\lambda}^U, V, d_{\lambda}^V, g)$ replaced by $(U, d_{\mu}^U, V, d_{\mu}^V, u|_{\partial V})$, one has

$$u^+(y) - u^+(x) = \mathcal{S}_{h, V}^+(y) - \mathcal{S}_{h, V}^+(x) \leq d_{\mu}^U(x, y) \quad \forall x, y \in \overline{V}.$$

Case 3. $x \in V$ and $y \in \overline{U} \setminus V$ or $x \in \overline{U} \setminus V$ and $y \in V$. For any $\epsilon > 0$, by Corollary 5.1, there exists a curve γ joining x, y such that

$$\ell_{d_{\mu}^U}(\gamma) \leq d_{\mu}^U(x, y) + \epsilon.$$

Let $z \in \gamma \cap \partial V$. Applying Case 2 and Case 1, we have

$$\begin{aligned} u^+(y) - u^+(x) &= u^+(y) - u^+(z) + u^+(z) - u^+(x) \leq d_{\mu}^U(z, y) + d_{\mu}^U(x, z) \\ &\leq \ell_{d_{\mu}^U}(\gamma) \leq d_{\mu}^U(x, y) + \epsilon. \end{aligned}$$

By the arbitrariness of $\epsilon > 0$, we have $u^+(y) - u^+(x) \leq d_{\mu}^U(x, y)$ as desired. \square

6 Proof of Theorem 1.6

In this section, we always suppose that the Hamiltonian $H(x, p)$ enjoys assumptions (H0)-(H3) and further that $\lambda_H = 0$.

Definition 6.1 Let $U \Subset \Omega$ be a domain and $g \in C^0(\partial U)$ with $\mu(g, \partial U) < \infty$.

- (i) A minimizer u for $\mathbf{I}(g, U)$ is called a local superminimizer for $\mathbf{I}(g, U)$ if $u \geq \mathcal{S}_{u|_{\partial V}, V}^-$ in V for any subdomain $V \subset U$.
- (ii) A minimizer u for $\mathbf{I}(g, U)$ is called a local subminimizer for $\mathbf{I}(g, U)$ if $u \leq \mathcal{S}_{u|_{\partial V}, V}^+$ in V for any subdomain $V \subset U$.

The next lemma shows McShane extensions are local super/sub minimizers.

Lemma 6.2 Let $U \subseteq \Omega$ and $g \in C^0(\partial U)$ with $\mu(g, \partial U) < \infty$.

(i) For any subdomain $V \subset U$, we have

$$\mathcal{S}_{h^+,V}^- \leq \mathcal{S}_{h^+,V}^+ \leq \mathcal{S}_{g;U}^+ \text{ in } V, \quad \text{where } h^+ = \mathcal{S}_{g;U}^+|_{\partial V}. \quad (6.1)$$

In particular, $\mathcal{S}_{g;U}^+$ is a local superminimizer for $\mathbf{I}(g, U)$.

(ii) For any subdomain $V \subset U$, we have

$$\mathcal{S}_{g;U}^- \leq \mathcal{S}_{h^-,V}^- \leq \mathcal{S}_{h^-,V}^+ \text{ in } V, \quad \text{where } h^- = \mathcal{S}_{g;U}^-|_{\partial V}. \quad (6.2)$$

In particular, $\mathcal{S}_{g;U}^-$ is a local subminimizer for $\mathbf{I}(g, U)$.

Proof We only prove (i); the proof for (ii) is similar. Write $\mu = \mu(g, \partial U)$. By Lemma 5.7, $\mathcal{S}_{g;U}^+$ is a minimizer for $\mathbf{I}(g, U)$, $\mu = \mathbf{I}(g, U) = \|H(\cdot, X\mathcal{S}_{g;U}^+)\|_{L^\infty(U)}$, and

$$\mathcal{S}_{g;U}^+(y) - \mathcal{S}_{g;U}^+(x) \leq d_\mu^U(x, y) \quad \forall x, y \in \overline{U} \quad (6.3)$$

Fix any subdomain $V \subset U$. Denote by $\mathcal{S}_{h^+,V}^\pm$ the McShane extension of $h^+ = \mathcal{S}_{g;U}^+|_{\partial V}$ in V . Note that $\mathcal{S}_{h^+,V}^- \leq \mathcal{S}_{h^+,V}^+$ in \overline{V} , that is, the first inequality in (6.1) holds.

Below we show $\mathcal{S}_{h^+,V}^+ \leq \mathcal{S}_{g;U}^+$ in V , that is, the second inequality in (6.1). Let u^+ be as in the (5.13) with $u = \mathcal{S}_{g;U}^+$, that is,

$$u^+ := \begin{cases} \mathcal{S}_{h^+,V}^+ & \text{in } V \\ \mathcal{S}_{g;U}^+ & \text{in } \overline{U} \setminus V. \end{cases}$$

By (6.3), we apply Lemma 5.9 to conclude that $u^+ \in \dot{W}_X^{1,\infty}(U) \cap C^0(\overline{U})$ and $\|H(\cdot, Xu^+)\|_{L^\infty(U)} \leq \mu$. Note that $u^+ = \mathcal{S}_{g;U}^+$ on ∂U and hence, by definition of $\mathbf{I}(g, U)$, one has $\mathbf{I}(g, U) \leq \|H(\cdot, Xu^+)\|_{L^\infty(U)}$. Recalling $\mu = \mathbf{I}(g, U)$, one obtains that $\mathbf{I}(g, U) = \|H(\cdot, Xu^+)\|_{L^\infty(U)}$, that is, u^+ is a minimizer for $\mathbf{I}(g, U)$. By Lemma 5.7(i) again, $u^+ \leq \mathcal{S}_{g;U}^+$ in U . Since $\mathcal{S}_{h^+,V}^+ = u^+$ in V , we conclude that $\mathcal{S}_{h^+,V}^+ \leq \mathcal{S}_{g;U}^+$ in V as desired. The proof is complete. \square

Lemma 6.3 Let $V \subseteq \Omega$ be a domain and $P = \{x_j\}_{j \in \mathbb{N}} \subset V$ be a dense subset of V . Assume $u \in C^0(\overline{V})$ and $\{u_j\}_{j \in \mathbb{N}} \subset \dot{W}_X^{1,\infty}(V) \cap C^0(\overline{V})$ such that for any $j \in \mathbb{N}$,

$$|u_j(x_i) - u(x_i)| \leq \frac{1}{j} \text{ for any } i = 1, \dots, j, \quad (6.4)$$

and

$$\|H(\cdot, Xu_j(\cdot))\|_{L^\infty(V)} \leq \lambda < \infty. \quad (6.5)$$

Then $u_j \rightarrow u$ in $C^0(V)$, and moreover,

$$u \in \dot{W}_X^{1,\infty}(V) \text{ and } \|H(x, Xu)\|_{L^\infty(V)} \leq \lambda. \quad (6.6)$$

Proof We only need to prove $u_j \rightarrow u$ in $C^0(V)$. Note that (6.6) follows from this and Lemma 3.1.

Since $u \in C^0(\overline{V})$ and \overline{V} is compact, u is uniform continuous in \overline{V} , that is, for any $\epsilon > 0$, there exists $h_\epsilon \in (0, \epsilon)$ such that for all

$$|u(x) - u(y)| \leq \epsilon \quad \text{whenever } x, y \in \overline{V} \quad \text{with } |x - y| < h_\epsilon. \quad (6.7)$$

Recalling the assumption (H3), by (6.5) one has

$$\|Xu_j(\cdot)\|_{L^\infty(V)} \leq R_\lambda \quad \forall j \in \mathbb{N}. \quad (6.8)$$

By Lemma 2.7,

$$u_j(y) - u_j(x) \leq R_\lambda d_{CC}^V(x, y) \quad \forall x, y \in V.$$

Given any $K \Subset V$, recall from [41] that

$$d_{CC}^V(x, y) \leq C(K, V)|x - y|^{1/k} \quad \forall x, y \in \overline{K}.$$

It then follows

$$|u_j(y) - u_j(x)| \leq R_\lambda C(K, V)|x - y|^{1/k} \quad \forall x, y \in \overline{K}.$$

Given any $\epsilon > 0$, thanks to the density of $\{x_i\}_{i \in \mathbb{N}}$ in V , one has $\overline{K} \subset \cup_{x_i \in \overline{K}} B(x_i, h_\epsilon)$. By the compactness of \overline{K} , we have

$$\overline{K} \subset \cup\{B(x_i, h_\epsilon) \mid 1 \leq i \leq i_K \text{ and } x_i \in \overline{K}\}$$

for some $i_K \in \mathbb{N}$. For any $j \geq \max\{i_K, 1/\epsilon\}$ and for any $x \in \overline{K}$, choose $1 \leq i \leq i_K$ such that $x_i \in \overline{K}$ and $|x - x_i| \leq h_\epsilon < \epsilon$. Thus $|u(x_i) - u(x)| \leq \epsilon$. By (6.4) we have $|u_j(x_i) - u(x_i)| \leq \frac{1}{j} \leq \epsilon$. Thus

$$\begin{aligned} |u_j(x) - u(x)| &\leq |u_j(x) - u_j(x_i)| + |u_j(x_i) - u(x_i)| + |u(x_i) - u(x)| \\ &\leq R_\lambda C(K, V)\epsilon^{1/k} + 2\epsilon. \end{aligned}$$

This implies that $u_j \rightarrow u$ in $C^0(\overline{K})$ as $j \rightarrow \infty$. The proof is complete. \square

The following clarifies the relations between absolute minimizers and local super/subminimizers.

Lemma 6.4 *Let $U \Subset \Omega$ and $g \in C^0(\partial U)$ with $\mu(g, \partial U) < \infty$. Then a function $u : \overline{U} \rightarrow \mathbb{R}$ is an absolute minimizer for $\mathbf{I}(g, U)$ if and only if it is both a local superminimizer and a local subminimizer for $\mathbf{I}(g, U)$.*

Proof If u is an absolute minimizer for $\mathbf{I}(g, U)$, then for every subdomain $V \subset U$, u is a minimizer for $\mathbf{I}(u|_{\partial V}, V)$. By Lemma 5.6, $\mathcal{S}_{u|_{\partial V}, V}^- \leq u \leq \mathcal{S}_{u|_{\partial V}, V}^+$, that is, u is both a local superminimizer and a local subminimizer for $\mathbf{I}(g, U)$.

Conversely, suppose that u is both a local superminimizer and a local subminimizer for $\mathbf{I}(g, U)$. We need to show that u is absolute minimizer for $\mathbf{I}(g, U)$. It suffices to prove that for any domain $V \Subset U$, u is a minimizer for $\mathbf{I}(u, V)$, in particular, $\|H(\cdot, Xu(\cdot))\|_{L^\infty(V)} \leq \mathbf{I}(u, V)$.

The proof consists of 3 steps.

Step 1. Given any subdomain $V \subset U$, choose a dense subset $\{x_j\}_{j \in \mathbb{N}}$ of V . Set $V_j = V \setminus \{x_i\}_{1 \leq i \leq j}$ and $V_0 = V$. Note that

$$\partial V_j = \partial V_{j-1} \cup \{x_j\} = \partial V_0 \cup \{x_i\}_{1 \leq i \leq j} \quad \forall j \in \mathbb{N}.$$

For each $j \geq 0$, set

$$\mu_j = \mu(u|_{\partial V_j}, \partial V_j) = \inf\{\lambda \geq 0 \mid u(y) - u(x) \leq d_\lambda^{V_j}(x, y) \quad \forall x, y \in \partial V_j\}.$$

Since $\overline{V} \subset \overline{U}$, we have $d_\mu^U(x, y) \leq d_\mu^V(x, y)$ for all $x, y \in \overline{V}^*$, we have

$$u(y) - u(x) \leq d_\mu^V(x, y) \quad \forall x, y \in \overline{V}.$$

and hence $\mu_0 \leq \mu$. By Lemma 5.7 (i), $\mathbf{I}(u, V_0) = \mu_0$. In a similar way and by induction, for all $j \geq 0$, since $V_{j+1} \subset V_j$, we have

$$\mathbf{I}(u, V_{j+1}) = \mu_{j+1} \leq \mu_j = \mathbf{I}(u, V_j) \leq \mu_0 = \mathbf{I}(u, V_0) \leq \mu = \mathbf{I}(u, U). \quad (6.9)$$

Step 2. We construct a sequence $\{u_j\}_{j \in \mathbb{N}}$ of functions so that, for each $j \in \mathbb{N}$,

$$u_j \in \dot{W}_X^{1,\infty}(V_j) \cap C^0(\bar{V}) \text{ and } u_j(x) = u(x) \text{ for any } x \in \partial V_j = \partial V \cup \{x_i\}_{1 \leq i \leq j} \quad (6.10)$$

and

$$\|H(\cdot, Xu_j(\cdot))\|_{L^\infty(V_{j-1})} = \mu_{j-1} \text{ for any } j \in \mathbb{N}. \quad (6.11)$$

For any $j \geq 1$, since u is both a local superminimizer and a local subminimizer for $\mathbf{I}(g, U)$, by Definition 6.1,

$$\mathcal{S}_{u|_{\partial V_{j-1}}, V_{j-1}}^- \leq u \leq \mathcal{S}_{u|_{\partial V_{j-1}}, V_{j-1}}^+ \text{ in } V_{j-1}.$$

At x_j , we have

$$a_j := \mathcal{S}_{u|_{\partial V_{j-1}}, V_{j-1}}^-(x_j) \leq u(x_j) \leq b_j := \mathcal{S}_{u|_{\partial V_{j-1}}, V_{j-1}}^+(x_j) \quad (6.12)$$

Define $u_j : \bar{V}_{j-1} = \bar{V} \rightarrow \mathbb{R}$ by

$$u_j := \begin{cases} \mathcal{S}_{u|_{\partial V_{j-1}}, V_{j-1}}^+ & \text{if } a_j = b_j, \\ \frac{u(x_j) - a_j}{b_j - a_j} \mathcal{S}_{u|_{\partial V_{j-1}}, V_{j-1}}^+ + (1 - \frac{u(x_j) - a_j}{b_j - a_j}) \mathcal{S}_{u|_{\partial V_{j-1}}, V_{j-1}}^- & \text{if } a_j < b_j. \end{cases}$$

To see (6.10), observe that Lemma 5.7 gives $u_j \in \dot{W}_X^{1,\infty}(V_j) \cap C^0(\bar{V})$. Moreover, for any $x \in \partial V_j$, one has either $x \in \partial V_{j-1}$ or $x = x_j$. In the case $x \in \partial V_{j-1}$, by Lemma 5.7 one has

$$u_j(x) = \mathcal{S}_{u|_{\partial V_{j-1}}, V_{j-1}}^+(x) = \mathcal{S}_{u|_{\partial V_{j-1}}, V_{j-1}}^-(x) = u(x).$$

In the case $x = x_j$, if $a_j = b_j$, then (6.12) implies

$$u(x_j) = \mathcal{S}_{u|_{\partial V_{j-1}}, V_{j-1}}^+(x_j) = b_j = u(x_j);$$

if $a_j < b_j$, then

$$\begin{aligned} u_j(x_j) &= \frac{u(x_j) - a_j}{b_j - a_j} \mathcal{S}_{u|_{\partial V_{j-1}}, V_{j-1}}^+(x_j) + (1 - \frac{u(x_j) - a_j}{b_j - a_j}) \mathcal{S}_{u|_{\partial V_{j-1}}, V_{j-1}}^-(x_j) \\ &= \frac{u(x_j) - a_j}{b_j - a_j} b_j + (1 - \frac{u(x_j) - a_j}{b_j - a_j}) a_j \\ &= u(x_j). \end{aligned}$$

To see (6.11), applying Lemma 5.7(iii) with $t = \frac{u(x_j) - a_j}{b_j - a_j}$, we deduce that u_j is a minimizer for $\mathbf{I}(u|_{\partial V_{j-1}}, V_{j-1})$, that is,

$$\|H(\cdot, Xu_j(\cdot))\|_{L^\infty(V_{j-1})} = \mathbf{I}(u|_{\partial V_{j-1}}, V_{j-1}) = \mu_{j-1}. \quad (6.13)$$

Step 3. We show that, for all $j \in \mathbb{N}$,

$$u_j(z) - u_j(y) \leq d_\mu^V(y, z) \quad \forall y, z \in V. \quad (6.14)$$

Note that, by Corollary 5.1 in V , (6.14) yields that $u_j \in \dot{W}_X^{1,\infty}(V)$ and $\|H(x, Xu_j)\|_{L^\infty(V)} \leq \mu$. Applying Lemma 6.3 to $\{u_j\}_{j \in \mathbb{N}}$ and u , we conclude that $u \in \dot{W}_X^{1,\infty}(V)$ and $\|H(x, Xu)\|_{L^\infty(V)} \leq \mu$ as desired.

To see (6.14), using (6.11) and Corollary 5.1 in V_{j-1} , we have

$$u_j(z) - u_j(y) \leq d_{\mu_{j-1}}^{V_{j-1}}(y, z) \quad \forall y, z \in V_{j-1}.$$

Thus (6.9) implies

$$u_j(z) - u_j(y) \leq d_\mu^{V_{j-1}}(y, z) \quad \forall y, z \in V_{j-1}.$$

Thanks to Lemma 5.4 we have $d_\mu^{V_{j-1}} = d_\mu^V$ in $V \times V$ and hence

$$u_j(z) - u_j(y) \leq d_\mu^V(y, z) \quad \forall y, z \in V_{j-1}.$$

By the continuity of u_j in \bar{V} we have (6.14) and hence finish the proof. \square

Finally, using Perron's approach, we obtain the existence of absolute minimizers.

Proposition 6.5 *Let $U \Subset \Omega$ and $g \in C^0(\partial U)$ with $\mu(g, \partial U) < \infty$. Define*

$$U_g^+(x) := \sup\{u(x) \mid u : \bar{U} \rightarrow \mathbb{R} \text{ is a local subminimizer for } \mathbf{I}(g, U)\} \quad \forall x \in \bar{U} \quad (6.15)$$

and

$$U_g^-(x) := \inf\{u(x) \mid u : \bar{U} \rightarrow \mathbb{R} \text{ is a local superminimizer for } \mathbf{I}(g, U)\} \quad \forall x \in \bar{U}.$$

Then U_g^\pm are absolute minimizers for $\mathbf{I}(g, U)$.

Proof We only show that U_g^+ is an absolute minimizer for $\mathbf{I}(g, U)$; similarly one can prove that U_g^- is also an absolute minimizer for $\mathbf{I}(g, U)$. Thanks to Lemma 6.4, it suffices to show that U_g^+ is a minimizer for $\mathbf{I}(g, U)$, a local subminimizer for $\mathbf{I}(g, U)$ and a local superminimizer for $\mathbf{I}(g, U)$. Note that $\mathbf{I}(g, U) = \mu(g, \partial U) < \infty$.

Prove that U_g^+ is a minimizer for $\mathbf{I}(g, U)$. Firstly, since any local subminimizer w for $\mathbf{I}(g, U)$ is a minimizer for $\mathbf{I}(g, U)$. We know

$$w \in \dot{W}_X^{1,\infty}(U) \cap C^0(\bar{U}), \quad w|_{\partial U} = g, \quad \text{and} \quad \|H(\cdot, Xw(\cdot))\|_{L^\infty(U)} = \mathbf{I}(g, U) < \infty.$$

Recalling the assumption (H3), we have $\|Xw\|_{L^\infty(U)} \leq R_{\mathbf{I}(g, U)}$ and hence $w \in \text{Lip}_{d_{CC}}(\bar{U})$ with $\text{Lip}_{d_{CC}}(w, \bar{U}) \leq R_{\mathbf{I}(g, U)}$. By a direct calculation, one has

$$U_g^+ \in \text{Lip}_{d_{CC}}(\bar{U}) \quad \text{with} \quad \text{Lip}_{d_{CC}}(U_g^+, \bar{U}) \leq R_{\mathbf{I}(g, U)} \quad \text{and} \quad U_g^+|_{\partial U} = g. \quad (6.16)$$

Next, let $\{x_i\}_{i \in \mathbb{N}}$ be a dense set of U . For any $i, j \in \mathbb{N}$, by the definition of U_g^+ , there exists a local subminimizer u_{ij} for $\mathbf{I}(g, U)$ such that

$$u_{ij}(x_i) \geq U_g^+(x_i) - \frac{1}{j}.$$

Note that, by Definition 6.1, u_{ij} is also a minimizer for $\mathbf{I}(g, U)$.

Moreover, for each $j \in \mathbb{N}$, write

$$u_j := \max\{u_{ij}\}_{1 \leq i \leq j}.$$

Lemma 5.7(iii) implies that u_j is a minimizer for $\mathbf{I}(g, U)$ and hence

$$u_j \in \dot{W}_X^{1,\infty}(U) \quad \text{and} \quad \|H(\cdot, Xu_j(\cdot))\|_{L^\infty(U)} \leq \mathbf{I}(g, U) \quad \forall j \in \mathbb{N}. \quad (6.17)$$

For $1 \leq i \leq j$, from the definition of U_g^+ , it follows that

$$U_g^+(x_i) \geq u_j(x_i) \geq U_g^+(x_i) - \frac{1}{j}. \quad (6.18)$$

Finally, combining (6.16), (6.17) and (6.18), we are able to apply Lemma 6.3 to U_g^+ and $\{u_j\}_{j \in \mathbb{N}}$ so to get

$$u_j \rightarrow U_g^+ \text{ in } C^0(U), \quad U_g^+ \in \dot{W}_X^{1,\infty}(U) \text{ and } \|H(\cdot, XU_g^+)\|_{L^\infty(U)} \leq \mathbf{I}(g, U). \quad (6.19)$$

Hence

$$\|H(\cdot, XU_g^+)\|_{L^\infty(U)} = \mathbf{I}(g, U).$$

Together with (6.16) yields that U_g^+ is a minimizer for $\mathbf{I}(g, U)$.

Prove that U_g^+ is a local subminimizer for $\mathbf{I}(g, U)$.

We argue by contradiction. Assume on the contrary that U_g^+ is not a local subminimizer for $\mathbf{I}(g, U)$. Then, by definition, there exists a subdomain $V \subset U$, and some x_0 in V such that

$$U_g^+(x_0) > \mathcal{S}_{h^+,V}^+(x_0),$$

where $\mathcal{S}_{h^+,V}$ is the McShane extension in V of $h^+ = U_g^+|_{\partial V}$. By the definition of U_g^+ , there exists a local subminimizer u for $\mathbf{I}(g, U)$ such that

$$U_g^+(x_0) \geq u(x_0) > \mathcal{S}_{h^+,V}^+(x_0). \quad (6.20)$$

The definition of U_g^+ also gives

$$u \leq U_g^+ \text{ in } U. \quad (6.21)$$

Define

$$E := \{x \in \overline{V} \mid u(x) > \mathcal{S}_{h^+,V}^+(x)\}.$$

Since both u and $\mathcal{S}_{h^+,V}^+$ are continuous, E is an open subset of \overline{V} . Since

$$U_g^+ = \mathcal{S}_{h^+,V}^+ \text{ on } \partial V,$$

by (6.21), we infer that $u \leq \mathcal{S}_{h^+,V}^+$ on ∂V and hence $E \subset V$. Obviously, $x_0 \in E$. Denote by E_0 the component of E containing x_0 .

Recalling that u is a local subminimizer for $\mathbf{I}(g, U)$, by Definition 6.1, we have $u \leq \mathcal{S}_{u|_{\partial E_0}, E_0}^+$ in E . Since $x_0 \in E_0$, we have $\mathcal{S}_{u|_{\partial E_0}, E_0}^+(x_0) \geq u(x_0)$, which, combined with (6.20), gives

$$\mathcal{S}_{u|_{\partial E_0}, E_0}^+(x_0) \geq u(x_0) > \mathcal{S}_{h^+,V}^+(x_0). \quad (6.22)$$

On the other hand, we are able to apply Lemma 6.2 with $(U, V, g, h = \mathcal{S}_{g;U|_{\partial V}}^+)$ therein replaced by $(V, E_0, h^+, h = \mathcal{S}_{h^+,V|_{E_0}}^+)$ here and then obtain

$$\mathcal{S}_{h^+,E_0}^+ \leq \mathcal{S}_{h^+,V}^+ \text{ in } E_0.$$

Since $u|_{E_0} = \mathcal{S}_{h^+,V|_{E_0}}^+ = h$, at $x_0 \in E$, we arrive at

$$\mathcal{S}_{u|_{\partial E_0}, E_0}^+(x_0) \leq \mathcal{S}_{h^+,V}^+(x_0). \quad (6.23)$$

Note that (6.22) contradicts with (6.23) as desired.

Prove that U_g^+ is a local superminimizer for $\mathbf{I}(g, U)$. By definition, it suffices to prove that, for any given subdomain $V \subset U$, we have $\mathcal{S}_{h^+, V}^- \leq U_g^+$ in V , where we write $h^+ = U_g^+|_{\partial V}$.

To this end, define u^+ as in (5.13) with u therein replaced by U_g^+ , that is,

$$u^+ := \begin{cases} \mathcal{S}_{h^+, V}^- & \text{in } V \\ U_g^+ & \text{in } \overline{U} \setminus V. \end{cases} \quad (6.24)$$

Then u^+ is a minimizer for $\mathbf{I}(g, U)$. Indeed, since U_g^+ is a minimizer for $\mathbf{I}(g, U)$, we know that U_g^+ satisfies (5.12) with $\mu = \mathbf{I}(g, U)$ therein. This allows us to apply Lemma 5.9 with $u = U_g^+$ therein and then conclude that $u^+ \in \dot{W}_X^{1, \infty}(U) \cap C^0(U)$, $u^+ = U_g^+ = g$ on ∂U , and $\|H(x, Xu^+)\| \leq \mathbf{I}(g, U)$. Therefore, by definition of $\mathbf{I}(g, U)$, $\|H(x, Xu^+)\| = \mathbf{I}(g, U)$, and hence u^+ is a minimizer for $\mathbf{I}(g, U)$.

We further claim that

$$u^+ \text{ is a local subminimizer for } \mathbf{I}(g, U). \quad (6.25)$$

Assume that this claim holds. Choosing u^+ as a test function in the definition of U_g^+ in (6.15), we know that

$$U_g^+ \geq u^+ \text{ in } U.$$

and in particular $U_g^+ \geq \mathcal{S}_{h^+, V}^-$ in V as desired.

Proof the claim (6.25).

To get (6.25), by Definition 6.1, we still need to show for any subdomain $B \subset U$,

$$u^+ \leq \mathcal{S}_{u^+|_{\partial B}, B}^+ \text{ in } B. \quad (6.26)$$

To prove (6.26), we argue by contradiction. Assume that (6.26) is not correct, that is,

$$W := \{x \in B \mid u^+(x) > \mathcal{S}_{u^+|_{\partial B}, B}^+(x)\} \neq \emptyset. \quad (6.27)$$

Up to considering some connected component of W , we may assume that W is connected. Note that

$$u^+ = \mathcal{S}_{u^+|_{\partial B}, B}^+ \text{ on } \partial W. \quad (6.28)$$

Consider the set

$$D := \{x \in B \mid U_g^+(x) > \mathcal{S}_{u^+|_{\partial B}, B}^+(x)\}. \quad (6.29)$$

By continuity, both of W and D are open.

Below, we consider two cases: $D = \emptyset$ and $D \neq \emptyset$.

Case $D = \emptyset$. If D is empty, then we always have $U_g^+ \leq \mathcal{S}_{u^+|_{\partial B}, B}^+$ in B . Thus

$$U_g^+ \leq \mathcal{S}_{u^+|_{\partial B}, B}^+ < u^+ \text{ in } W.$$

Since $U_g^+ = u^+ \in \overline{U} \setminus V$, this implies

$$W \subset V. \quad (6.30)$$

Since $u^+ = \mathcal{S}_{h^+, V}^-$ in \overline{V} and $W \subset V$ gives $\partial W \subset \overline{V}$, we have

$$u^+ = \mathcal{S}_{h^+, V}^- \text{ on } \partial W. \quad (6.31)$$

Moreover, by Lemma 6.2, we know that $\mathcal{S}_{h^+,V}^-$ with $h^+ = U_g^+|_{\partial V}$ is a local subminimizer for $\mathbf{I}(U_g^+, V)$. By the definition of local subminimizer, and by $W \subset V$, we have

$$\mathcal{S}_{h^+,V}^- \leq \mathcal{S}_{u^+|_{\partial W},W}^+ \quad \text{in } W, \quad (6.32)$$

where we recall $\mathcal{S}_{h^+,V}^-|_{\partial W} = u^+|_{\partial W}$ from (6.31).

Applying Lemma 6.2 with (U, V, g, h^+) therein replaced by $(B, W, u^+|_{\partial B}, \mathcal{S}_{u^+|_{\partial B},B}^+|_{\partial W})$ here, recalling $\mathcal{S}_{u^+|_{\partial B},B}^+|_{\partial W} = u^+|_{\partial W}$ from (6.28), we obtain

$$\mathcal{S}_{u^+|_{\partial W},W}^+ \leq \mathcal{S}_{u^+|_{\partial B},B}^+ \quad \text{in } W. \quad (6.33)$$

Combining (6.32) and (6.33), by $W \subset V$, one arrives at

$$u^+ = \mathcal{S}_{h^+,V}^- \leq \mathcal{S}_{u^+|_{\partial B},B}^+ \quad \text{in } W,$$

which contradicts with (6.27).

Case $D \neq \emptyset$. Up to considering some connected component of D , we may assume that D is connected. By the definition of D in (6.29), we infer that

$$U_g^+ = \mathcal{S}_{u^+|_{\partial B},B}^+ \quad \text{on } \partial D. \quad (6.34)$$

Since U_g^+ is a local subminimizer for $\mathbf{I}(g, U)$ as proved above, we know

$$U_g^+ \leq \mathcal{S}_{U_g^+|_{\partial D},D}^+ \quad \text{in } D. \quad (6.35)$$

Applying Lemma 6.2 with (U, V, g, h^+) therein replaced by $(B, D, u^+|_{\partial B}, \mathcal{S}_{u^+|_{\partial B},B}^+|_{\partial D})$, recalling $\mathcal{S}_{u^+|_{\partial B},B}^+|_{\partial D} = U_g^+|_{\partial D}$ from (6.34), we obtain

$$\mathcal{S}_{U_g^+|_{\partial D},D}^+ \leq \mathcal{S}_{u^+|_{\partial B},B}^+ \quad \text{in } D. \quad (6.36)$$

Combining (6.35) and (6.36), we deduce

$$U_g^+ \leq \mathcal{S}_{u^+|_{\partial B},B}^+ \quad \text{in } D.$$

Recalling (6.29), this contradicts with $D \neq \emptyset$. The proof is complete. \square

Theorem 1.6 is now a direct consequence of the above series of results.

Proof of Theorem 1.6 Let $g \in \text{Lip}_{d_{CC}}(\partial U)$. It suffices to show that $\mu(g, \partial U) < \infty$, which allows us to use Proposition 6.5 and then conclude the desired absolute minimizer U_g^+ therein.

Taking $0 < \lambda < \infty$ such that $R'_\lambda \geq \text{Lip}_{d_{CC}}(g, \partial U)$, we have

$$g(y) - g(x) \leq R'_\lambda d_{CC}(x, y) \quad \forall x, y \in \partial U.$$

From Lemma 2.9 (ii), that is, $R'_\lambda d_{CC} \leq d_\lambda^U$, it follows that

$$g(y) - g(x) \leq d_\lambda^U(x, y) \quad \forall x, y \in \partial U.$$

and hence that

$$\mu(g, \partial U) = \inf\{\mu \geq 0 \mid g(y) - g(x) \leq d_\mu(x, y)\} \leq \lambda < \infty.$$

The proof is complete. \square

7 Further discussion

Note that Rademacher type Theorem 1.3 is a cornerstone when showing the existence of absolute minimizers. Indeed, Champion and Pascale [14] and Guo-Xiang-Yang [26] established partial results similar to Theorem 1.3 for a special class of Hamiltonians considered in this paper to show the existence of absolute minimizers. However, their method seems to be invalid for more general Hamiltonians considered in this paper. We briefly explain the reason below.

Remark 7.1 Champion and Pascale [14] showed the McShane extension is a minimizer for H when H is lower semi-continuous on $U \times \mathbb{R}^n$. In fact, they defined another intrinsic distance induced by $H(x, p)$. For every $\lambda \geq 0$,

$$L_\lambda(x, q) := \sup_{\{p \in H_\lambda(x)\}} p \cdot q, \quad \forall x \in \overline{U} \text{ and } q \in \mathbb{R}^n.$$

where $H_\lambda(x)$ is the sub-level set at x , namely, $H_\lambda(x) = \{p \in \mathbb{R}^n \mid H(x, p) \leq \lambda\}$.

For $0 \leq a < b \leq +\infty$, let $\gamma : [a, b] \rightarrow \overline{U}$ be a Lipschitz curve, that is, there exists a constant $C > 0$ such that $|\gamma(s) - \gamma(t)| \leq C|s - t|$ whenever $s, t \in [a, b]$. The L_λ -length of γ is defined by

$$\ell_\lambda(\gamma) := \int_a^b L_\lambda(\gamma(\theta), \gamma'(\theta)) d\theta,$$

which is nonnegative, since $L_\lambda(x, q) \geq 0$ for any $x \in \overline{U}$ and $q \in \mathbb{R}^n$. For a pair of points $x, y \in \overline{U}$, the \bar{d}_λ -distance from x to y is defined by

$$\bar{d}_\lambda(x, y) := \inf \left\{ \ell_\lambda(\gamma) \mid \gamma \in \mathcal{C}(a, b, x, y, \overline{U}) \right\}.$$

Then, they prove two intrinsic pseudo-distance are equal, that is

$$d_\lambda(x, y) = \bar{d}_\lambda(x, y) \quad \text{for any } \lambda > 0 \text{ and for any } x, y \in \overline{U}. \quad (7.1)$$

Thanks to the definition of \bar{d}_λ , they can justify (i) \Leftrightarrow (ii) in Rademacher type Theorem 1.3.

However, when asserting (7.1), we will meet obstacles in generalizing [14] Proposition A.2 since we are faced with measurable H .

On the other hand, Guo-Xiang-Yang [26] provided another method to identify a weak version of (7.1) for measurable Finsler metrics H , that is

$$\lim_{y \rightarrow x} \frac{d_\lambda(x, y)}{\bar{d}_\lambda(x, y)} = 1 \quad \text{for any } \lambda > 0 \text{ and for any } x, y \in \overline{U}. \quad (7.2)$$

Here \tilde{d}_λ induced by measurable Finsler metrics H is defined in the following way.

$$\tilde{d}_\lambda(x, y) := \sup_N \inf \left\{ \ell_\lambda(\gamma) \mid \gamma \in \Gamma_N(a, b, x, y, \overline{U}) \right\} \quad (7.3)$$

where the supremum is taken over all subsets N of \overline{U} such that $|N| = 0$ and $\Gamma_N(a, b, x, y, U)$ denotes the set of all Lipschitz continuous curves γ in \overline{U} with end points x and y such that $\mathcal{H}^1(N \cap \gamma) = 0$ with \mathcal{H}^1 being the one dimensional Hausdorff measure.

In fact, (7.2), combined with the method in [14] will be sufficient for validating (i) \Leftrightarrow (ii) in Rademacher type Theorem 1.3. Unfortunately, since we are coping with Hörmander vector field, a barrier arises when modifying their proofs. Indeed, their method uses a result by [16] that every x -measurable Hamiltonian H can be approximated by a sequence of smooth Hamiltonians $\{H_n\}_n$ such that two intrinsic distances $\tilde{d}_\lambda^{H_n}$ and $d_\lambda^{H_n}$ induced by H_n by means

of (7.3) and (1.4) satisfy $\lim_{n \rightarrow \infty} d_\lambda^{H_n} = d_\lambda$ and $\limsup_{n \rightarrow \infty} \tilde{d}_\lambda^{H_n} \leq \tilde{d}_\lambda$ uniformly on $U \times U$ respectively. The process of the proof of [16] is based on a C^1 Lusin approximation property for curves. Namely, given a Lipschitz curve $\gamma : [0, 1] \rightarrow U$ joining x and y , for any $\epsilon > 0$, there exists a C^1 curve $\tilde{\gamma} : [0, 1] \rightarrow U$ with the same endpoints such that

$$\mathcal{L}^1(\{t \in [0, 1] \mid \tilde{\gamma}(t) \neq \gamma(t) \text{ or } \tilde{\gamma}'(t) \neq \gamma'(t)\}) < \epsilon$$

where \mathcal{L}^1 denotes the one dimensional Lebesgue measure. Besides,

$$\|\tilde{\gamma}\|_{L^\infty} \leq c \|\gamma\|_{L^\infty},$$

for some constant c depending only on n . Although this version of C^1 Lusin approximation property holds for horizontal curves in Heisenberg groups ([43]) and step 2 Carnot groups ([39]), it fails for some horizontal curve in Engel group ([43]).

In summary, it is difficult to generalize the properties of the pseudo metric \tilde{d}_λ not only from Euclidean space to the case of Hörmander vector fields but also from lower-semicontinuous $H(x, p)$ to measurable $H(x, p)$. Hence we would like to pose the following open problem.

Problem 7.2 Under the assumptions (H0)-(H3), does (7.2) holds?

Acknowledgements The authors would like to thank Katrin Fässler for the reference [39, 43] and her comment on this paper. The authors are also grateful to the anonymous referee for many valuable suggestions which make the paper more self-contained and improve the final presentation of the paper.

Funding Open Access funding provided by University of Jyväskylä (JYU).

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Appendix A: Rademacher's theorem in Euclidean domains—revisit

In this appendix, we state some consequence of Rademacher's theorem (Theorem 1.1) for Sobolev and Lipschitz classes, see Lemma 3 and Lemma 6 below. They were well-known in the literature and also partially motivated our Theorem 1.3 and Corollary 1.5. For reader's convenience, we give the details.

Recall that $\Omega \subset \mathbb{R}^n$ is always a domain. The homogeneous Sobolev space $\dot{W}^{1,\infty}(\Omega)$ is the collection of all functions $u \in L^1_{\text{loc}}(\Omega)$ with its distributional derivative $\nabla u = (\frac{\partial u}{\partial x_i})_{1 \leq i \leq n} \in L^\infty(\Omega)$. We equip $\dot{W}^{1,\infty}(\Omega)$ with the semi-norm

$$\|u\|_{\dot{W}^{1,\infty}(\Omega)} = \|\nabla u\|_{L^\infty(\Omega)}.$$

Write $\dot{W}^{1,\infty}_{\text{loc}}(\Omega)$ as the collection of all functions u in Ω so that $u \in \dot{W}^{1,\infty}(V)$ whenever $V \Subset \Omega$. Here and below, $V \Subset \Omega$ means that V is bounded domain with $\bar{V} \subset \Omega$. On the other hand, denote by $\text{Lip}(\Omega)$ the collection of all Lipschitz functions u in Ω , that is, all functions u satisfying (1.1). We equip $\text{Lip}(\Omega)$ with the semi-norm

$$\text{Lip}(u, \Omega) = \sup_{x, y \in \Omega, x \neq y} \frac{|u(y) - u(x)|}{|x - y|} = \inf\{\lambda \geq 0 \text{ satisfying (1.1)}\}.$$

Denote by $\text{Lip}_{\text{loc}}(\Omega)$ the collection of all functions u in Ω so that $u \in \text{Lip}(V)$ for any $V \Subset \Omega$. Moreover, denote by $\text{Lip}^*(\Omega)$ the collection of all functions u in Ω with

$$\text{Lip}^*(u, \Omega) := \sup_{x \in \Omega} \text{Lip}u(x) < \infty. \quad (1)$$

Obviously, $\text{Lip}(\Omega) \subset \text{Lip}^*(\Omega)$ with the seminorm bound $\text{Lip}^*(u, \Omega) \leq \text{Lip}(u, \Omega)$. Next, we have the following relation.

Lemma 1 *We have $\text{Lip}^*(\Omega) \subset \text{Lip}_{\text{loc}}(\Omega)$. For any convex subdomain $V \subset \Omega$ and $u \in \text{Lip}^*(\Omega)$, we have*

$$|u(x) - u(y)| \leq \|u\|_{\text{Lip}^*(V)} |x - y| \quad \forall x, y \in V. \quad (2)$$

Proof Let $u \in \text{Lip}^*(\Omega)$. To see $u \in \text{Lip}_{\text{loc}}(\Omega)$, it suffices to prove $u \in \text{Lip}(B)$ for any ball $B \Subset \Omega$. Given any $x, y \in B$, denote by $\gamma(t) = x + t(y - x)$. Since $A_{x,y} = \sup_{t \in [0,1]} \text{Lip}u(\gamma(t)) < \infty$, for each $t \in [0, 1]$ we can find $r_t > 0$ such that $|u(\gamma(s)) - u(\gamma(t))| \leq A_{x,y} |\gamma(s) - \gamma(t)| = A_{x,y} |s - t| |x - y|$ whenever $|s - t| \leq r_t$ and $s \in [0, 1]$. Since $[0, 1] \subset \bigcup_{t \in [0,1]} (t - r_t, t + r_t)$ we can find an increasing sequence $t_i \in [0, 1]$ with $t_0 = 0$ and $t_N = 1$ such that $[0, 1] \subset \bigcup_{i=1}^N (t_i - \frac{1}{2}r_{t_i}, t_i + \frac{1}{2}r_{t_i})$. Write $x_i = \gamma(t_i)$ for $i = 0, \dots, N$. We have

$$\begin{aligned} |u(x) - u(y)| &= \left| \sum_{i=0}^{N-1} [u(x_i) - u(x_{i+1})] \right| \leq \sum_{i=0}^{N-1} |u(x_i) - u(x_{i+1})| \\ &\leq A_{x,y} \sum_{i=0}^{N-1} |x_i - x_{i+1}| = A_{x,y} |x - y|. \end{aligned}$$

Noticing that $A_{x,y} \leq \text{Lip}^*(u, B) \leq \text{Lip}^*(u, \Omega) < \infty$ for all $x, y \in B$, we deduce that $u \in \text{Lip}(B)$ and hence $u \in \text{Lip}_{\text{loc}}(\Omega)$.

If $V \Subset \Omega$ is convex, then for any $x, y \in \Omega$, the line-segment joining x and y lies in V . Hence similar to the above discussion, we have

$$|u(x) - u(y)| \leq A_{x,y} |x - y|, \quad \forall x, y \in V$$

and $A_{x,y} \leq \|u\|_{\text{Lip}^*(V)} < \infty$ for all $x, y \in V$. Therefore, (2) holds and the proof is complete. \square

On the other hand, functions $\dot{W}_{\text{loc}}^{1,\infty}(\Omega)$ admit continuous representatives.

Lemma 2 (i) *Each $u \in \dot{W}_{\text{loc}}^{1,\infty}(\Omega)$ admits a unique continuous representative \tilde{u} , that is, $\tilde{u} \in \dot{W}_{\text{loc}}^{1,\infty}(\Omega)$ with $\tilde{u} = u$ almost everywhere in Ω . Moreover, $\tilde{u} \in \text{Lip}_{\text{loc}}(\Omega)$, and for any convex subdomain $V \subset \Omega$, we have*

$$|\tilde{u}(x) - \tilde{u}(y)| \leq \|u\|_{\dot{W}^{1,\infty}(\Omega)} |x - y| \quad \forall x, y \in V.$$

(ii) *Each $u \in \dot{W}^{1,\infty}(\Omega)$ admits a unique continuous representative \tilde{u} , that is, $\tilde{u} \in \dot{W}^{1,\infty}(\Omega)$ with $\tilde{u} = u$ almost everywhere in Ω . Moreover, $\tilde{u} \in \text{Lip}^*(\Omega)$ with $\text{Lip}^*(\tilde{u}, \Omega) \leq \|u\|_{\dot{W}^{1,\infty}(\Omega)}$.*

Proof Since (ii) can be shown in a similar way as (i), we only prove (i). Given any convex domain $V \Subset \Omega$, for any pair x, y of Lebesgue points of u , we have

$$u(y) - u(x) = \lim_{\delta \rightarrow 0} [u * \eta_\delta(y) - u * \eta_\delta(x)] = \lim_{\delta \rightarrow 0} \nabla(u * \eta_\delta)(x + t_\delta(y - x)) \cdot (y - x)$$

where η_δ is the standard mollifier in \mathbb{R}^n with its support $\text{spt}\eta_\delta \subset B(0, \delta)$ and $t_\delta \in [0, 1]$. Also, since for any $z \in V$,

$$|\nabla(u * \eta_\delta)(z)| = |(\nabla u) * \eta_\delta(z)| \leq \|\nabla u\|_{L^\infty(B(z, \delta))},$$

we deduce that for any pair x, y of Lebesgue points of u ,

$$|u(y) - u(x)| \leq \|\nabla u\|_{L^\infty(V)}|y - x|.$$

If $z \in V$ is not a Lebesgue point of u , let $\{z_i\}_{i \in \mathbb{N}} \subset V$ be a sequence of Lebesgue points of u converging to z . We have

$$\lim_{i \rightarrow \infty} |u(z_i) - u(z_{i+1})| \leq \lim_{i \rightarrow \infty} \|\nabla u\|_{L^\infty(V)}|z_i - z_{i+1}| = 0,$$

which implies $\{u(z_i)\}_{i \in \mathbb{N}} \subset \mathbb{R}$ is a Cauchy sequence. Since $\|u\|_{L^\infty(V)} < \infty$, we know $\{u(z_i)\}_{i \in \mathbb{N}}$ has a limit in \mathbb{R} independent of the choice of the sequence $\{u(z_i)\}_{i \in \mathbb{N}}$. Define $\tilde{u}(z) := u(z)$ if $z \in V$ is a Lebesgue point of u and $\tilde{u}(z) = \lim_{i \rightarrow \infty} u(z_i)$ if $z \in V$ is not a Lebesgue point of u where $\{z_i\}_{i \in \mathbb{N}} \subset V$ is a sequence of Lebesgue points of u converging to z . We know $\tilde{u} : V \rightarrow \mathbb{R}$ is well-defined and moreover,

$$|\tilde{u}(y) - \tilde{u}(x)| \leq \|\nabla u\|_{L^\infty(V)}|y - x| \quad \forall x, y \in V.$$

Thus $\tilde{u} \in \text{Lip}(V)$ with $\sup_{x \in V} \text{Lip}\tilde{u}(x) \leq \text{Lip}(\tilde{u}, V) \leq \|\nabla u\|_{L^\infty(V)}$. In particular, \tilde{u} is continuous, which shows (i). \square

Thanks to lemma 2, below for any function $u \in \dot{W}_{\text{loc}}^{1,\infty}(\Omega)$ or $u \in \dot{W}^{1,\infty}(\Omega)$, up to considering its continuous representative \tilde{u} , we may assume that u is continuous. Under this assumption, Lemma 2 further gives $\dot{W}_{\text{loc}}^{1,\infty}(\Omega) \subset \text{Lip}_{\text{loc}}(\Omega)$, and $\dot{W}^{1,\infty}(\Omega) \subset \text{Lip}^*(\Omega)$ with a norm bound $\text{Lip}^*(u, \Omega) \leq \|u\|_{\dot{W}^{1,\infty}(\Omega)}$. Rademacher's theorem (Theorem 1.1) tells that their converse are also true. Indeed, we have the following.

Lemma 3 (i) We have $\dot{W}_{\text{loc}}^{1,\infty}(\Omega) = \text{Lip}_{\text{loc}}(\Omega)$ and $\text{Lip}(\Omega) \subset \dot{W}^{1,\infty}(\Omega) = \text{Lip}^*(\Omega)$ with $\text{Lip}^*(u, \Omega) = \|u\|_{\dot{W}^{1,\infty}(\Omega)} \leq \text{Lip}(u, \Omega)$.

(ii) If Ω is convex, then $\text{Lip}(\Omega) = \dot{W}^{1,\infty}(\Omega) = \text{Lip}^*(\Omega)$ with $\text{Lip}^*(u, \Omega) = \|u\|_{\dot{W}^{1,\infty}(\Omega)} = \text{Lip}(u, \Omega)$.

Proof (i) If $u \in \text{Lip}_{\text{loc}}(\Omega)$, applying Rademacher's theorem (Theorem 1.1) to all subdomains $V \Subset \Omega$, one has $u \in \dot{W}_{\text{loc}}^{1,\infty}(\Omega)$ and $|\nabla u(x)| = \text{Lip}u(x)$ for almost all $x \in \Omega$ (whenever u is differentiable at x). Hence $\text{Lip}_{\text{loc}}(\Omega) \subset \dot{W}_{\text{loc}}^{1,\infty}(\Omega)$. Combining Lemma 2(i), we know $\text{Lip}_{\text{loc}}(\Omega) = \dot{W}_{\text{loc}}^{1,\infty}(\Omega)$.

If $u \in \text{Lip}^*(\Omega)$, that is, $\text{Lip}^*(u, \Omega) = \sup_{x \in \Omega} \text{Lip}u(x) < \infty$. We have $u \in \text{Lip}_{\text{loc}}(\Omega)$ and hence $u \in \dot{W}_{\text{loc}}^{1,\infty}(\Omega)$ and $|\nabla u(x)| = \text{Lip}u(x) \leq \text{Lip}^*(u, \Omega) < \infty$ for almost all $x \in \Omega$. Thus $u \in \dot{W}^{1,\infty}(\Omega)$.

By definition, it is obvious that $\text{Lip}(\Omega) \subset \text{Lip}^*(\Omega)$. Hence Lemma 3 (i) holds.

(ii) By Lemma 3 (i), we only need to show $\text{Lip}^*(\Omega) \subset \text{Lip}(\Omega)$. Applying Lemma 1 with $V = \Omega$ therein, (2) becomes

$$\frac{|u(x) - u(y)|}{|x - y|} \leq \|u\|_{\text{Lip}^*(\Omega)} \quad \forall x, y \in \Omega.$$

Taking supremum among all $x, y \in \Omega$ in the left hand side of the above inequality, we arrive at

$$\|u\|_{\text{Lip}(\Omega)} \leq \|u\|_{\text{Lip}^*(\Omega)},$$

which gives the desired result. \square

Remark 4 (i) Lemma 1 and Lemma 3 fail if we relax $\sup_{x \in \Omega} \text{Lip} u(x)$ in the definition (1) to be $\|\text{Lip} u\|_{L^\infty(\Omega)} = \text{esssup}_{x \in \Omega} \text{Lip} u(x)$. This is witnessed by the standard Cantor function w in $[0, 1]$. Denote by E the standard Cantor set. It is well-known that w is continuous but not absolutely continuous in $[0, 1]$. Since Lipschitz functions are always absolutely continuous, we know that w is neither Lipschitz nor locally Lipschitz in $\Omega = (0, 1)$, and hence $w \notin \text{Lip}_{\text{loc}}(\Omega)$. On the other hand, observe that $\Omega \setminus E$ consists of a sequence of open intervals which mutually disjoint, and w is a constant in each such intervals and hence $\text{Lip} w(x) = 0$ therein. So we know that $\text{Lip} w(x) = 0$ in $\Omega \setminus E$. Since $|E| = 0$, we have $\|\text{Lip} u\|_{L^\infty(\Omega)} = 0$.

(ii) In general, if Ω is not convex, one cannot expect that $\dot{W}^{1,\infty}(\Omega) \subset \text{Lip}(\Omega)$ with a norm bound. Indeed, consider the planar domain

$$U := \{x = (x_1, x_2) \in \mathbb{R}^2 \mid |x| < 1\} \setminus [0, 1) \times \{0\}. \quad (3)$$

Indeed, in the polar coordinate (r, θ) , let $w : U \rightarrow \mathbb{R}$ be

$$w(r, \theta) := r\theta \text{ for all } 0 < r < 1 \text{ and } 0 < \theta < 2\pi.$$

One can show that $w \in \dot{W}^{1,\infty}(U)$ so that $w(x_1, x_2) < \pi/3$ when $1/2 \leq x_1 < 1$ and $0 < x_2 < 1/10$, and $w(x_1, x_2) > 5\pi/6$ when $1/2 \leq x_1 < 1$ and $-1/10 < x_2 < 0$. One has $\text{Lip}(w, \Omega) = \infty$ and hence $w \notin \text{Lip}(\Omega)$.

The example in Remark 4 (ii) also indicates that the Euclidean distance does not match the geometry of domains and hence $\text{Lip}(\Omega)$ defined via Euclidean distance is not the perfect one to understand $\dot{W}^{1,\infty}(\Omega)$.

Instead of Euclidean distance, for any domain Ω , we consider the intrinsic distance

$$d_E^\Omega(x, y) = \inf\{\ell(\gamma) \mid \gamma : [0, 1] \rightarrow \Omega \text{ is absolute continuous curve joining } x, y\}, \quad (4)$$

where $\ell(\gamma) := \int_0^1 |\dot{\gamma}(t)| dt$ is the Euclidean length. We have the dual formula.

Lemma 5 (i) For any $x, y \in \Omega$,

$$d_E^\Omega(x, y) = \sup\{u(y) - u(x) \mid u \in \dot{W}^{1,\infty}(\Omega), \|\nabla u\|_{L^\infty(\Omega)} \leq 1\}. \quad (5)$$

(ii) If $x, y \in \Omega$ with $|x - y| \leq \text{dist}(x, \partial\Omega)$, then $d_E^\Omega(x, y) = |x - y|$.

(iii) If Ω is convex, then $d_E^\Omega(x, y) = |x - y|$ for all $x, y \in \Omega$.

Proof (i) Set

$$\tilde{d}_E^\Omega(x, y) = \sup\{u(y) - u(x) \mid u \in \dot{W}^{1,\infty}(\Omega), \|\nabla u\|_{L^\infty(\Omega)} \leq 1\}. \quad (6)$$

Notice that $d_E^\Omega(x, \cdot) \in \text{Lip}^*(\Omega) = \dot{W}^{1,\infty}(\Omega)$ (Lemma 3 (i)) and $\|\nabla d_E^\Omega(x, \cdot)\|_{L^\infty(\Omega)} \leq 1$ for all $x \in \Omega$. Hence letting $d_E^\Omega(x, \cdot)$ be the test function in (6), we see

$$d_E^\Omega(x, y) \leq \tilde{d}_E^\Omega(x, y) \quad \forall x, y \in \Omega.$$

To see the contrary, fix $x, y \in \Omega$. Let $\{u_i\}_{i \in \mathbb{N}}$ be a sequence of test functions in (6) such that

$$\tilde{d}_E^\Omega(x, y) = \lim_{i \rightarrow \infty} (u_i(y) - u_i(x)).$$

Let $\gamma : [0, 1] \rightarrow \Omega$ be an arbitrary absolute continuous curve joining x and y . Then there exists a domain $U \Subset \Omega$ with $\gamma \subset U$. Let $\{\eta_\delta\}_{\delta > 0}$ be the standard mollifiers in \mathbb{R}^n . For each

$i \in \mathbb{N}$, we know $u_i * \eta_\delta \in C^\infty(\Omega)$ and $\|\nabla(u_i * \eta_\delta)\|_{L^\infty(U)} \leq \|\nabla u_i\|_{L^\infty(U)} \leq 1$. Then we have

$$\begin{aligned}\tilde{d}_E^\Omega(x, y) &= \lim_{i \rightarrow \infty} [u_i(y) - u_i(x)] \\ &= \lim_{i \rightarrow \infty} \lim_{\delta \rightarrow 0} [u_i * \eta_\delta(y) - u_i * \eta_\delta(x)] \\ &= \lim_{i \rightarrow \infty} \lim_{\delta \rightarrow 0} \int_0^1 \nabla(u_i * \eta_\delta)(\gamma(t)) \cdot \dot{\gamma}(t) dt \\ &\leq \lim_{i \rightarrow \infty} \lim_{\delta \rightarrow 0} \int_0^1 |\nabla(u_i * \eta_\delta)(\gamma(t))| |\dot{\gamma}(t)| dt \\ &\leq \int_0^1 |\dot{\gamma}(t)| dt \\ &= \ell(\gamma).\end{aligned}$$

Finally, taking infimum among all absolute continuous curves joining x and y in the above inequality, we conclude

$$\tilde{d}_E^\Omega(x, y) \leq d_E^\Omega(x, y) \quad \forall x, y \in \Omega.$$

(ii) If $|x - y| \leq \text{dist}(x, \partial\Omega)$, then the line-segment γ joining x and y is contained in Ω . Letting γ be the absolute continuous curve in (4),

$$|x - y| \leq d_E^\Omega(x, y) \leq \ell(\gamma) = |x - y| \quad \forall x, y \in \Omega.$$

(iii) If Ω is convex, for all $x, y \in \Omega$, since the line-segment joining them is contained in Ω , similarly to (ii), we have $d_E^\Omega(x, y) = |x - y|$. The proof is complete. \square

Note that if Ω is not convex, one cannot expect $d_E^\Omega(x, y) = |x - y|$ for all $x, y \in \Omega$. Indeed, if Ω is given by the domain U as in (3), for points $(1/2, \epsilon)$ and $(1/2, -\epsilon)$ with $\epsilon \in (0, 1/10)$, the Euclidean distance between them is 2ϵ . However, note that any curve $\gamma : [0, 1] \rightarrow \Omega$ joining them must have intersection with $(-1, 0) \times \{0\}$, which is call z . One then deduce that

$$\ell(\gamma) \geq |(1/2, \epsilon) - z| + |(-1/2, \epsilon) - z| \geq 1/2 + 1/2 = 1 + 2\epsilon.$$

Thus the intrinsic distance between $(1/2, \epsilon)$ and $(1/2, -\epsilon)$ is always larger than or equals to $1 + 2\epsilon$.

With in Lemma 5, we show that the Lipschitz spaces defined via the intrinsic distance perfectly match with the Sobolev space $\dot{W}^{1,\infty}(\Omega)$, see Lemma 6 below. Denote by $\text{Lip}_{d_E^\Omega}(\Omega)$ the collection of all Lipschitz functions u in Ω with respect to d_E^Ω , that is,

$$\text{Lip}_{d_E^\Omega}(u, \Omega) := \sup \frac{|u(x) - u(y)|}{d_E^\Omega(x, y)} < \infty.$$

We also denote by $\text{Lip}_{d_E^\Omega}^*(\Omega)$ the collection of all functions u in Ω with

$$\text{Lip}_{d_E^\Omega}^*(u, \Omega) := \sup_{x \in \Omega} \text{Lip}_{d_E^\Omega} u(x) < \infty.$$

Lemma 6 We have $\text{Lip}_{d_E^\Omega}(\Omega) = \dot{W}^{1,\infty}(\Omega) = \text{Lip}^*(\Omega)$ and

$$\|\nabla u\|_{L^\infty(\Omega)} = \text{Lip}_{d_E^\Omega}(u, \Omega) = \text{Lip}^*(u, \Omega) = \sup_{x \in \Omega} \text{Lip}_{d_E^\Omega} u(x).$$

Proof Recall that Lemma 6 gives $d_E^\Omega(x, y) = |x - y|$ whenever $|x - y| \leq \text{dist}(x, \partial\Omega)$. One then has $\text{Lip}_{d_E^\Omega}(\Omega) \subset \text{Lip}_{\text{loc}}(\Omega)$, and moreover, $\text{Lip}u(x) = \text{Lip}_{d_E^\Omega}u(x)$ for all $x \in \Omega$, which gives $\text{Lip}^*_{d_E^\Omega}(\Omega) = \text{Lip}^*(\Omega)$.

Next, we show $\dot{W}^{1,\infty}(\Omega) \subset \text{Lip}_{d_E^\Omega}(\Omega)$ and $\text{Lip}_{d_E^\Omega}(u, \Omega) \leq \|\nabla u\|_{L^\infty(\Omega)}$. Let $u \in \dot{W}^{1,\infty}(\Omega)$. Then $\|\nabla u\|_{L^\infty(\Omega)} =: \lambda < \infty$. If $\lambda > 0$, then $\lambda^{-1}u \in \dot{W}^{1,\infty}(\Omega)$ and $\|\nabla(\lambda^{-1}u)\|_{L^\infty(\Omega)} = 1$. Hence $\lambda^{-1}u$ could be the test function in (5), which implies

$$\lambda^{-1}u(y) - \lambda^{-1}u(x) \leq d_E^\Omega(x, y) \quad \forall x, y \in \Omega,$$

or equivalently,

$$\frac{|u(y) - u(x)|}{\|\nabla u\|_{L^\infty(\Omega)}} \leq d_E^\Omega(x, y) \quad \forall x, y \in \Omega.$$

Therefore, $u \in \text{Lip}_{d_E^\Omega}(\Omega)$ and $\text{Lip}_{d_E^\Omega}(u, \Omega) \leq \|\nabla u\|_{L^\infty(\Omega)}$. If $\lambda = 0$, then similar as the above discussion, we have for any $\lambda' > 0$

$$\frac{|u(y) - u(x)|}{\lambda'} \leq d_E^\Omega(x, y) \quad \forall x, y \in \Omega.$$

Therefore, $u \in \text{Lip}_{d_E^\Omega}(\Omega)$ and $\text{Lip}_{d_E^\Omega}(u, \Omega) \leq \lambda'$ for any $\lambda' > 0$. Hence $\text{Lip}_{d_E^\Omega}(u, \Omega) = 0 = \|\nabla u\|_{L^\infty(\Omega)}$.

Moreover, we show $\text{Lip}_{d_E^\Omega}(\Omega) \subset \text{Lip}^*(\Omega)$ and $\text{Lip}^*(u, \Omega) \leq \text{Lip}_{d_E^\Omega}(u, \Omega)$. Let $u \in \text{Lip}_{d_E^\Omega}(\Omega)$. Then $\text{Lip}_{d_E^\Omega}(u, \Omega) < \infty$. Since $\text{Lip}^*_{d_E^\Omega}(u, \Omega) \leq \text{Lip}_{d_E^\Omega}(u, \Omega)$ and $\text{Lip}^*_{d_E^\Omega}(u, \Omega) = \text{Lip}(u, \Omega)$, we arrive at

$$\text{Lip}(u, \Omega) \leq \text{Lip}_{d_E^\Omega}(u, \Omega) < \infty.$$

Therefore, $u \in \text{Lip}^*(\Omega)$.

Finally, recalling that $\dot{W}^{1,\infty}(\Omega) = \text{Lip}^*(\Omega)$ and $\|\nabla u\|_{L^\infty(\Omega)} = \text{Lip}(u, \Omega)$ in Lemma 3, we finish the proof. \square

References

1. Aronsson, G.: Minimization problems for the functional $\sup_x F(x, f(x), f'(x))$. Ark. Mat. **6**, 33–53 (1965)
2. Aronsson, G.: Minimization problems for the functional $\sup_x F(x, f(x), f'(x))$. II. Ark. Mat. **6**, 409–431 (1966)
3. Aronsson, G.: Extension of functions satisfying Lipschitz conditions. Ark. Mat. **6**, 551–561 (1967)
4. G. Aronsson, On the partial differential equation $u_x^2 u_{xx} + 2u_x u_y u_{xy} + u_y^2 u_{yy} = 0$. Ark. Mat. **7**, (1968), 395–425
5. Aronsson, G.: Minimization problems for the functional $\sup_x F(x, f(x), f'(x))$. III. Ark. Mat. **7**, 509–512 (1969)
6. Armstrong, S.N., Crandall, M.G., Julin, V., Smart, C.K.: Convexity criteria and uniqueness of absolutely minimizing functions. Arch. Ration. Mech. Anal. **200**(2), 405–443 (2011)
7. Aronsson, G., Crandall, M.G., Juutinen, P.: A tour of the theory of absolutely minimizing functions. Bull. Am. Math. Soc. (N.S.) **41**, 439–505 (2004)
8. N. Barron, Viscosity solutions and analysis in L^∞ . In: Nonlinear Analysis, Differential Equations and Control (Montreal, QC, 1998). NATO Sci. Ser. C Math. Phys. Sci. 528. Dordrecht: Kluwer Acad. Publ., pp. 1–60. (1999)
9. Barron, E.N., Jensen, R.R., Wang, C.Y.: The Euler equation and absolute minimizers of L^∞ functionals. Arch. Ration. Mech. Anal. **157**, 255–283 (2001)
10. Bieske, T.: On ∞ -harmonic functions on the Heisenberg group. Comm. Partial Diff. Eq. **27**(3–4), 727–761 (2002)

11. Boutet de Monvel, A., Lenz, D., Stollmann, P.: Schnol's theorem for strongly local forms. *Israel J. Math.* **173**, 189–211 (2009)
12. Champion, T., De Pascale, L., Prinari, F.: Γ -Convergence and absolute minimizers for supremal functionals. *ESAIM Control Optim. Calc. Var.* **10**, 14–27 (2004)
13. V. M. Chernikov, S. K. Vodop'yanov, Sobolev Spaces and hypoelliptic equations I,II. *Siberian Advances in Mathematics*. 6 (1996) no. 3, 27–67, no. 4, 64–96. Translation from: *Trudy In-ta matematiki RAN. Sib. otd-nie*. 29 (1995), 7–62
14. Champion, T., De Pascale, L.: Principles of comparison with distance functions for absolute minimizers. *J. Convex Anal.* **14**, 515–541 (2007)
15. Crandall, M.: An efficient derivation of the Aronsson equation. *Arch. Ration. Mech. Anal.* **167**, 271–279 (2003)
16. Davini, A.: Smooth approximation of weak Finsler metrics. *Diff. Integr. Eq.* **18**(5), 509–530 (2005)
17. Dragoni, F., Manfredi, J.J., Vittone, D.: Weak Fubini property and infinity harmonic functions in Riemannian and sub-Riemannian manifolds. *Trans. Am. Math. Soc.* **365**(2), 837–859 (2013)
18. Friedrichs, K.O.: The identity of weak and strong extensions of differential operators. *Trans. Am. Math. Soc.* **55**, 132–151 (1944)
19. Franchi, B., Hajlasz, P., Koskela, P.: Definitions of Sobolev classes on metric spaces. *Annales de l'Institut Fourier* **49**(6), 1903–1924 (1999)
20. Franchi, B., Serapioni, R., Serra Cassano, F.: Meyers-Serrin type theorems and relaxation of variational integrals depending on vector fields. *Houston J. Math.* **22**, 859–890 (1996)
21. Franchi, B., Serapioni, R., Serra Cassano, F.: Approximation and imbedding theorems for weighted Sobolev spaces associated with Lipschitz continuous vector fields. *Boll. Un. Mat. Ital.* **7**, 83–117 (1997)
22. Frank, R.L., Lenz, D., Wingert, D.: Intrinsic metrics for non-local symmetric Dirichlet forms and applications to spectral theory. *J. Funct. Anal.* **266**(8), 4765–4808 (2014)
23. Garofalo, N., Nhieu, D.M.: Isoperimetric and Sobolev inequalities for Carnot-Carathéodory spaces and the existence of minimal surfaces. *Commun. Pure Appl. Math.* **49**, 1081–1144 (1996)
24. Garofalo, N., Nhieu, D.: Lipschitz continuity, global smooth approximations and extension theorems for Sobolev functions in Carnot-Carathéodory spaces. *J. Anal. Math.* **74**, 67–97 (1998)
25. Gariepy, R., Wang, C.Y., Yu, Y.: Generalized cone comparison principle for viscosity solutions of the Aronsson equation and absolute minimizers. *Commun. Partial Diff. Eq.* **31**, 1027–1046 (2006)
26. Guo, C.Y., Xiang, C., Yang, D.: L^∞ -variational problems associated to measurable Finsler structures. *Nonlinear Anal.* **132**, 126–140 (2015)
27. Hajlasz, P., Koskela, P.: Sobolev met Poincaré. *Mem. Am. Math. Soc.* **145**, 688 (2000)
28. J. Heinonen, P. Koskela, N. Shanmugalingam and J. T. Tyson. Sobolev spaces on metric measure spaces: an approach based on upper gradients, Cambridge Studies in Advanced Mathematics Series, Cambridge University Press, 2015
29. Jensen, R.: Uniqueness of Lipschitz extensions: minimizing the sup norm of the gradient. *Arch. Ration. Mech. Anal.* **123**, 51–74 (1993)
30. Jensen, R., Wang, C.Y., Yu, Y.: Uniqueness and nonuniqueness of viscosity solutions to Aronsson's equation. *Arch. Ration. Mech. Anal.* **190**(2), 347–370 (2008)
31. Jerison, D.: The Poincaré inequality for vector fields satisfying Hörmander's condition. *Duke Math. J.* **53**, 503–523 (1986)
32. D. Jerison, A. Sanchez-Calle, Subelliptic, second order differential operators. In: *Complex analysis, III* (College Park, Md., 1985–86). pp. 46–77, Lecture Notes in Math., 1277, Springer, 1987
33. Juutinen, P.: Minimization problems for Lipschitz functions via viscosity solutions. *Ann. Acad. Sci. Fenn. Math. Diss. No.* **115**, 53 (1998)
34. Juutinen, P.: Absolutely minimizing Lipschitz extensions on a metric space. *Ann. Acad. Sci. Fenn. Math.* **27**(1), 57–67 (2002)
35. Juutinen, P., Shanmugalingam, N.: Equivalence of AMLE, strong AMLE, and comparison with cones in metric measure space. *Math. Nachr.* **279**, 1083–1098 (2006)
36. Koskela, P., Shanmugalingam, N., Zhou, Y.: L^∞ -Variational problem associated to Dirichlet forms. *Math. Res. Lett.* **19**, 1263–1275 (2012)
37. Koskela, P., Zhou, Y.: Geometry and analysis of Dirichlet forms. *Adv. Math.* **231**, 2755–2801 (2012)
38. Koskela, P., Shanmugalingam, N., Zhou, Y.: Intrinsic geometry and analysis of diffusion process and L^∞ -variational problem. *Arch. Rational Mech. Anal.* **214**(1), 99–142 (2014)
39. Le Donne, E., Speight, G.: Lusin approximation for horizontal curves in step 2 carnot groups. *Calculus Var. Partial Diff. Eq.* **55**, 1–22 (2016)
40. Monti, R., Cassano, F.S.: Surface measures in Carnot- Caratheodory spaces. *Calc. Var. Partial Diff. Eq.* **13**, 339–376 (2001)

41. Nagel, A., Stein, E.M., S, Wainger,: Balls and metrics defined by vectorfields I. basic properties. *Acta Math.* **155**, 103–147 (1985)
42. Pansu, P.: Métriques de Carnot-Carathéodory et quasiisométries des espaces symétriques de rang un. *Annals Math.* **129**, 1–60 (1989)
43. Speight, G.: Approximation and Horizontal Curves in Carnot Groups. *Revista Matematica Iberoamericana* **32**, 1425–1446 (2016)
44. Stollmann, P.: A dual characterization of length spaces with application to Dirichlet metric spaces. *Stud. Math.* **198**, 221–233 (2010)
45. Sturm, K.T.: Analysis on local Dirichlet spaces. I. Recurrence, conservativeness and L^p -Liouville properties. *J. Reine Angew. Math.* **456**, 173–196 (1994)
46. Sturm, K.T.: Is a diffusion process determined by its intrinsic metric? *Chaos Solitons Fractals* **8**, 1855–1860 (1997)
47. Wang, C.Y.: The Aronsson equation for absolute minimizers of L^∞ functionals associated with vector fields satisfying Hörmander’s conditions. *Trans. AMS.* **359**, 91–113 (2007)

Publisher’s Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.