

This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Ikonen, Jenni N.; Törmäkangas, Timo; von Bonsdorff, Mikaela B.; Mikkola, Tuija M.; Eriksson, Johan G.; Haapanen, Markus J.

Title: Physical and mental functioning trajectory classes among older adults and their association with specialized healthcare use

Year: 2023

Version: Published version

Copyright: © The Author(s) 2023

Rights: CC BY 4.0

Rights url: <https://creativecommons.org/licenses/by/4.0/>

Please cite the original version:

Ikonen, J. N., Törmäkangas, T., von Bonsdorff, M. B., Mikkola, T. M., Eriksson, J. G., & Haapanen, M. J. (2023). Physical and mental functioning trajectory classes among older adults and their association with specialized healthcare use. *BMC Geriatrics*, 23, Article 448.

<https://doi.org/10.1186/s12877-023-04157-w>

RESEARCH

Open Access



Physical and mental functioning trajectory classes among older adults and their association with specialized healthcare use

Jenni N. Ikonen^{1,2,3}, Timo Törmäkangas⁴, Mikaela B. von Bonsdorff^{2,4}, Tuija M. Mikkola^{2,5}, Johan G. Eriksson^{1,2,6,7} and Markus J. Haapanen^{1,2,8*}

Abstract

Background Sex-specific physical and mental functioning trajectory classification could offer a way of understanding the differences in healthcare use at older age.

Methods Using latent growth mixture models, sex-specific physical and mental functioning trajectory classes were formed for 1991 participants (mean age 61.5 years) of the Helsinki Birth Cohort Study. Physical and mental functioning were evaluated with the SF-36 survey conducted in clinical examinations in 2001–2004, 2011–2013, and 2017–2018. First and follow-up outpatient visits, emergency visits, and hospital days were extracted from a national register between the first clinical examination and the year 2017. We used regression models to examine the associations between healthcare use and trajectory classes.

Results Two physical and mental functioning trajectory classes, *high* and *intermediate*, were observed for both sexes. The *intermediate* physical functioning trajectory class was associated with higher utilization rates of all examined specialized healthcare services (fully-adjusted IRRs varying 1.36–1.58; 95% CI = 1.03–1.79, 95% CI = 1.21–2.05) compared to the *high* trajectory class. Relative to the *high* trajectory class, the *intermediate* mental trajectory class was associated with the use of first outpatient visits (fully-adjusted IRRs 1.17, 95% CI = 1.03–1.33 for men, and 1.16, 95% CI = 1.04–1.30 for women). The findings were similar among both sexes.

Conclusions Compared to the *high* trajectory class, the *intermediate* physical functioning trajectory class was associated with greater specialized healthcare use and the *intermediate* mental trajectory class with first outpatient visits. Public health interventions should be considered to support functioning with aging.

Keywords Growth mixture model, Hospital service, Health service

*Correspondence:

Markus J. Haapanen
markus.haapanen@helsinki.fi

Full list of author information is available at the end of the article



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Introduction

Globally, healthcare systems are facing challenges due to the increasing demands of rapidly aging populations. As most healthcare expenses in high-income countries result from specialized healthcare services provided mainly at hospitals [1], there is a need to understand factors associated with healthcare service use among older adults.

Healthy aging requires physical functioning that enables wellbeing at older age [2]. Decline in physical functioning, however, is inevitable at older age. Yet the rate of decline varies depending on environmental, sociodemographic, and lifestyle factors [2]. Decline in physical functioning might also be temporal or permanent making the declining process dynamic [2–4]. Therefore, distinguishable long-term patterns, trajectories, can be recognised in a population [2–4] and might be more reliable to understand declining physical functioning at older age. Moreover, although functional impairments have been associated with adverse health outcomes like mortality [5], and acute and long-term care use [6, 7], little is known on the association between physical functioning trajectory classes and specialized healthcare use [8, 9]. Further, since older women are reported to have poorer physical functioning than older men [10, 11], sex-specific physical functioning trajectory classification might offer valuable information on the possible sex differences in healthcare use.

In addition to changes in physical functioning, possible changes in social circles and chronic diseases at an older age may increase the likelihood of deterioration of mental health [12]. However, although the literature shows a higher likelihood of worse mental health among women compared to men [13, 14], no studies have examined the association between sex-specific mental functioning trajectory classes and healthcare use.

The aim of the current study is to examine the association between physical and mental functioning trajectory classes and specialized healthcare use among older men and women. We hypothesized that worse physical and mental functioning classes would be associated with greater specialized healthcare use compared to those with more stable classes.

Methods

Study population

The participants of the present study are a subsample of the Helsinki Birth Cohort Study of 8760 individuals born in Helsinki University Hospital between 1934 and 1944 [15]. A random sample of 2003 individuals participated in clinical examinations between the years 2001 and 2004. Follow-up examinations were performed

between the years 2011 and 2013 and 2017 and 2018 as seen in the flowchart in Supplementary Fig. 1. Physical and mental functioning were assessed with the validated 36-Item Short Form Health Survey [16, 17] at all three study visits for those who participated (Supplementary Figure 1). Specialized healthcare use was obtained from a national specialized healthcare register starting from the first clinical examination date and ending at the end of the year 2017. Overall, 1999 participants had data on physical and mental functioning from at least one clinical visit to determine their physical and mental functioning. Eight individuals were excluded due to death in less than a year after the first clinical examination. The final study population consisted of 1991 individuals. All study participants gave a written informed consent before participating in any clinical procedures. The clinical study was performed in accordance with the Declaration of Helsinki and approved by the Coordinating Ethics Committee of the Hospital District of Helsinki and Uusimaa (THL/101/6.02.00/2019).

Physical and mental functioning

Physical and mental functioning were assessed with the Finnish validated version of the 36-Item Short Form Health Survey Version 1.0 (SF-36), which measures self-reported general health and wellbeing [17]. The SF-36 consists of eight sub-domains which each include 2–10 items. The questionnaire has shown to be valid and practical instrument for older community dwelling older adults [18], and unidimensionality and reliability of the sub-scales have been proved among older adults [19]. The sub-domains are fundamentally associated with healthy functioning and are named as physical functioning (10 items), role limitation due to physical problems (4 items), bodily pain (2 items), general health perception (5 items), vitality (4 items), social functioning (2 items), role limitation due to emotional problems (3 items), and mental health (5 items). Each item was coded to range from 0 to 100 with higher scores indicating better functioning or well-being. Sub-domain scores were formed by calculating means of the items, and coded missing if less than half of the items were available. The scores of the sub-domains were then standardized [20] using the means and standard deviations of a US reference population from 1990, and then weighted using factor score coefficients from the US population. Next, the sub-domain scores were aggregated to create physical and mental component scores. Finally, both component scores were standardized using mean of 50 and standard deviation of 10 [21]. The same research assistant nurses carried out the procedures at all three clinical examinations. The functioning scores were further divided by 10 for analysis in mixture models.

Healthcare utilization data

Healthcare utilization data were retrieved from the Care Register of Health Care, a national specialized healthcare register maintained by the Finnish Institute for Health and Welfare [22]. The register includes data on outpatient and inpatient healthcare visits from all hospitals in Finland. The data were retrieved starting from the first clinical examination date organized during 2001–2004 and continued until the end of the year 2017 or time of death where applicable. Emergency visits and first and follow-up outpatient visits were directly extracted from the register whereas hospital days were calculated by subtracting the admission date from the discharge date. If inpatient care exceeded 365 days, it was no longer treated as acute care and coded as 365 days in the dataset. Visits or hospital days were further summed for each study participant. First outpatient visits included all visits where a patient care relationship started with a new referral to a specific specialty while follow-up outpatient visits included all visits regarding the check-ups of the previously examined chronic diseases or injuries.

Covariates

Covariates were assessed in the first clinical examination. Self-reported physician diagnosed chronic diseases or symptoms included myocardial infarction, angina pectoris, congestive heart failure, claudication, osteoporosis, stroke, depression, asthma, and emphysema. Self-reported chronic diseases were divided into three categories including 'no diseases', 'one chronic disease', and 'two or more chronic diseases'. Frequency of smoking was assessed with a questionnaire and categorized into 'never', 'former', and 'current smoker'. Self-reported alcohol consumption was categorized into 'does not consume', 'two times a month at most', and 'three times a month or more'. Information on the participants' highest educational attainment was obtained from Statistics Finland [23] in the year 2000 and divided into four groups: 'basic or less or unknown', 'upper secondary', 'lower tertiary including polytechnic, vocational, and bachelor's degree', and 'upper tertiary including master's degree or higher'. Information on leisure time physical activity (LTPA) was assessed with the Kuopio Ischemic Heart Disease Risk Factor Study questionnaire [24]. Participants were asked about the duration, frequency, and intensity of LTPA during the past 12 months. Based on activity-specific metabolic equivalent (MET) values for each intensity grade, total leisure-time physical activity was calculated, and the results were expressed as METhours per week.

Statistical analyses

To form physical and mental functioning trajectory classes, latent growth mixture models (LGMM) were fitted to the SF-36 summary component scores available at the clinical examinations in 2001–2004, 2011–2013 and 2017–2018. The analyses were performed separately for women and men with Mplus version 7 (1998–2015). Assuming non-response was missing at random, we used LGMM with Full Information Maximum Likelihood [25] to capture unobserved latent classes using all available data with similar physical and mental functioning trajectories, but which were relatively distinct across the latent classes over follow-up time. Within the latent classes, the component scores were used as outcomes in a growth model with time indexed through individually varying observation times corresponding to the chronological age of participants. Each latent class had their own growth parameters, intercept (indicating participant's level of physical or mental functioning scores) and slope (indicating participant's rate of linear changes in physical or mental functioning scores over time). For the classes, the average trajectory was based on the means of the intercept and slope parameters calculated at age means at each time point. Final classification of the participants was assigned by the highest group membership posterior probability which was based on data from each participant's observed trajectory. The optimal number of latent classes was determined using the Bayesian Information Criterion (BIC), with lower values indicating a better fit of the model, and clarity of the participants' classification. Clarity of classification into trajectory classes was assessed with percentage of individuals falling into latent classes based on membership posterior probabilities (high probabilities indicates clear separation of trajectory classes) and summarized by model entropy (an aggregate function of posterior probabilities), which ranges between zero and one, with higher values one indicating better class separation. Average trajectory classes were named based on the starting level and the slope of the graphical average trajectory: trajectory classes with better functioning between the numbers were named 'high' and the lower average trajectory classes 'intermediate' due to graphical presentation approximately in the middle of the chart. Trajectory classes with no or little slope were named 'stable' since a slight decrease in functioning was considered to be related to normal aging. Trajectories with more declining slopes were named 'declining'.

Means and standard deviations for continuous variables and proportions for categorical variables were calculated to compare baseline characteristics. Statistical differences in baseline characteristics were examined

with the independent sample t-test, the Mann–Whitney U test, or the chi-square test. We used negative binomial regression models to examine the associations between physical and mental functioning trajectory classes and specialized healthcare use. Two models were created: Model 1 was adjusted for age, and Model 2 additionally for LTPA, education, number of chronic diseases, smoking, and alcohol consumption. The analyses were performed separately for men and women, and individual exposure times (offsets) were adjusted for the models. The results are shown as incidence rate ratios (IRRs). Significance level was set at 0.05, and p-values were not corrected for multiple testing. Further, negative binomial regression models were used to perform sensitivity analyses where we examined the associations between specialized healthcare use and physical and mental functioning trajectory classes with membership probability of 0.85 or higher. The analyses were performed in SPSS 26.0 for Windows (version 26.0, 1989–2020, SPSS Inc., Armonk, NY, USA).

Results

Physical and mental functioning trajectory classes

Figures 1 and 2 show physical and mental functioning trajectories in the two classes identified for men and women. The two physical functioning trajectory classes for an average participant among men and women

were named *high declining* (65.6% of men, 55.5% of women) and *intermediate declining* (34.4% of men, 44.5% of women). As presented in Figs. 1 and 2, two mental functioning average trajectory classes were identified for both men and women: *high stable* (77.5% of men and 65.9% of women) and *intermediate stable* (22.5% of men and 34.1% of women). We refer to these trajectory classes as *high* and *intermediate* hereon in this manuscript. Supplementary Figs. 2 and 3 in the Supplement show all observed individual physical and mental functioning trajectories by sex.

Characteristics of the study participants

Table 1 presents the characteristics of the study population. Men were more likely to have higher education, were more likely ex- or current smokers, were characterised by higher alcohol consumption, and had slightly higher annual utilization rates of emergency visits, first outpatient visits, and hospital days compared to women (Table 1). There were no differences between sexes in age, mean number of chronic diseases, and LTPA (Table 1). Supplementary Tables 1 and 2 in the Supplement show the characteristics of the study population according to the physical and mental functioning trajectory classes and sex.

MEN




Fig. 1 Model-expectation trajectories and 95% confidence intervals for physical (PCS) and mental (MCS) component score at means of individually varying age in three measurement time points from two groups (intermediate declining and high declining or intermediate stable or high stable) of growth mixture models among men. (Note: PCS and MCS were divided by 10 for the analyses resulting in the expected average trajectory scores seen in y-axis)

WOMEN




Fig. 2 Model-expectation trajectories and 95% confidence intervals for physical (PCS) and mental (MCS) component scores at means of individually varying age in three measurement time points from two groups (intermediate declining and high declining or intermediate stable or high stable) of growth mixture models among women. (Note: PCS and MCS were divided by 10 for the analyses resulting in the expected average trajectory scores seen in y-axis)

Physical and mental functioning trajectory classes and specialized healthcare use

Table 2 displays the results of regression analyses regarding physical functioning trajectory classes and specialized healthcare service use. Both men and women in the *intermediate* physical functioning trajectory class had higher utilization rates of any specialized healthcare service than those in the *high* trajectory class (Table 2). After all adjustments, the IRRs of both sexes were similar (Table 2).

Table 3 shows the IRRs for specialized healthcare service use by mental functioning trajectory classes. Men and women classified into the *intermediate* trajectory class utilized first outpatient visits at higher rates in comparison to the *high* trajectory class. The utilization rates of first outpatient visits in the *intermediate* trajectory class were similar among both sexes (Table 3). In fully-adjusted models, no other statistically significant associations were observed (Table 3). Overall, the IRRs for specialized healthcare service use by mental functioning trajectory classes were smaller compared to the IRRs of physical functioning trajectory classes. The results were further confirmed in sensitivity analyses as seen in Supplementary Tables 3 and 4 in the Supplement.

Discussion

In the present study, men and women in the *intermediate* physical functioning trajectory class had higher utilization rates of any specialized healthcare service compared with high functioning. Relative to the *high* mental trajectory class, the *intermediate* mental functioning trajectory class was associated with higher utilization rate of first outpatient visits. The findings were similar among both sexes.

We identified two physical and mental functioning trajectory classes for men and women. More than half of both men (65.5%) and women (55.5%) were included in the *high* physical functioning trajectory class suggesting relatively good physical functioning. In contrast, approximately one third of men (34.4%) and almost half of women (45.5%) belonged to the *intermediate* physical functioning trajectory class. Therefore, although decline over age was observed in both high and intermediate trajectory classes, a substantial proportion of the study population had an overall lower physical functioning level at older age. Among men and women, the observed individual trajectories in the *intermediate* class were less precisely captured by the trajectory model due to large variation in trajectory shapes suggesting more, possibly health-related variation in functioning patterns.

Table 1 Baseline characteristics of the participants at the clinical examinations between the years 2001 and 2004

	All		Men		Women		P-value
	n	Mean (SD)	n	Mean (SD)	n	Mean (SD)	
Age	1991	61.5 (2.9)	921	61.5 (2.8)	1070	61.5 (3.0)	0.96
Education							<0.001
Basic or less or unknown, %	709	35.6	273	29.6	436	40.7	
Upper secondary, %	511	25.7	225	24.4	286	26.7	
Lower level tertiary, %	528	87.8	272	29.5	256	23.9	
Upper level tertiary, %	243	12.2	151	16.4	92	8.6	
Chronic diseases							0.30
No diseases, %	1370	68.8	648	70.4	722	67.5	
1 chronic disease, %	430	21.6	185	20.1	245	22.9	
2 or more chronic diseases, %	191	9.6	88	9.6	103	9.6	
Smoking							<0.001
Never, %	834	41.9	245	26.8	589	55.6	
Quitted earlier, %	669	33.6	419	45.8	250	23.6	
Current smoker, %	472	23.7	251	27.4	221	20.8	
Alcohol use							<0.001
Does not use	146	7.3	69	7.5	77	7.2	
2 times/month at most	820	41.2	255	27.8	565	53.2	
3 times/month or more	1013	50.9	592	64.6	421	39.6	
Physical activity (METHours/week)	1955	46.0 (39.2)	900	45.2 (35.6)	1055	46.7 (41.1)	0.61
Specialized healthcare use during the follow-up							
Emergency visits/year	1991	0.29 (0.44)	921	0.35 (0.50)	1070	0.25 (0.39)	<0.001
First outpatient visits/year	1991	0.36 (0.33)	921	0.37 (0.33)	1070	0.36 (0.34)	0.37
Follow-up outpatient visits/year	1991	1.9 (4.2)	921	2.2 (5.6)	1070	1.62 (2.6)	0.016
Hospital days/year	1991	2.5 (7.3)	921	3.1 (8.1)	1070	1.9 (6.5)	<0.001

SD Standard deviation, MET Metabolic equivalent

Table 2 Incidence rate ratios (IRRs) for healthcare service use across physical functioning trajectory classes and sex

	Men		Women	
	Model 1 ^a	Model 2 ^b	Model 1 ^a	Model 2 ^b
	IRR (95% CI)	IRR (95% CI)	IRR (95% CI)	IRR (95% CI)
Emergency visit				
High declining	Ref	Ref	Ref	Ref
Intermediate declining	1.73 (1.47, 2.04) ^{***}	1.47 (1.25, 1.74) ^{***}	1.59 (1.35, 1.88) ^{***}	1.41 (1.18, 1.67) ^{***}
First outpatient visit				
High declining	Ref	Ref	Ref	Ref
Intermediate declining	1.52 (1.36, 1.70) ^{***}	1.43 (1.28, 1.60) ^{***}	1.70 (1.54, 1.88) ^{***}	1.56 (1.41, 1.73) ^{***}
Follow-up outpatient visit				
High declining	Ref	Ref	Ref	Ref
Intermediate declining	1.68 (1.21, 2.31) ^{**}	1.58 (1.21, 2.05) ^{***}	1.61 (1.36, 1.92) ^{***}	1.43 (1.20, 1.70) ^{***}
Hospital days				
High declining	Ref	Ref	Ref	Ref
Intermediate declining	1.76 (1.32, 2.35) ^{***}	1.38 (1.02, 1.87) [*]	1.70 (1.20, 2.40) ^{**}	1.36 (1.03, 1.79) [*]

CI Confidence interval

* = $p < 0.05$, ** = $p < 0.01$, *** = $p < 0.001$ ^a Adjusted for age, $n = 921$ men and $n = 1070$ women^b Adjusted for age, physical activity, education, number of chronic diseases, smoking, and alcohol consumption, $n = 894$ men and $n = 1043$ women

Table 3 Incidence rate ratios (IRRs) for healthcare service use across mental functioning trajectory classes and sex

	Men		Women	
	Model 1 ^a	Model 2 ^b	Model 1 ^a	Model 2 ^b
	IRR (95% CI)	IRR (95% CI)	IRR (95% CI)	IRR (95% CI)
Emergency visit				
High declining	Ref	Ref	Ref	Ref
Intermediate declining	1.25 (1.03, 1.53)*	1.07 (0.90, 1.27)	1.21 (1.02, 1.44)*	1.08 (0.92, 1.28)
First outpatient visit				
High declining	Ref	Ref	Ref	Ref
Intermediate declining	1.21 (1.07, 1.38)**	1.17 (1.03, 1.33)*	1.24 (1.11, 1.39)**	1.16 (1.04, 1.30)**
Follow-up outpatient visit				
High declining	Ref	Ref	Ref	Ref
Intermediate declining	1.08 (0.84, 1.39)	1.10 (0.88, 1.37)	1.22 (1.02, 1.48)*	1.08 (0.90, 1.31)
Hospital days				
High declining	Ref	Ref	Ref	Ref
Intermediate declining	1.58 (1.14, 2.18)**	1.32 (0.92, 1.91)	1.09 (0.79, 1.52)	0.96 (0.72, 1.28)

CI Confidence interval

* = $p < 0.05$, ** = $p < 0.01$, *** = $p < 0.001$

^a Adjusted for age, $n = 921$ men and $n = 1070$ women

^b Adjusted for age, physical activity, education, number of chronic diseases, smoking, and alcohol consumption, $n = 894$ men and $n = 1043$ women

Most men (77.5%) and most women (65.9%) belonged to the *high* mental trajectory classes. In contrast, the *intermediate* mental functioning trajectory class was presented by approximately one fifth of men (22.5%) and one third of women (34.1%). Although specific interpretations cannot be done based on the proportions, poorer physical performance [10, 11] and mental health [13, 14] have been reported among older women compared to men.

The *intermediate* physical functioning trajectory class was consistently associated with specialized healthcare use among both sexes relative to the *high* physical functioning trajectory class. As far as we are aware, no other studies have examined the association between specialized healthcare use and physical functioning trajectory classes assessed as a function of the participants' chronological age and separately for both sexes. However, other studies using latent growth mixture models have found associations between trajectory classes and disability [8, 9]. A Chinese study reported an association between late-onset disability trajectory classes and inpatient and outpatient care [8]. Further, late-onset and progressive disability trajectory classes were associated with expenses of inpatient and outpatient care [8]. Another study found progressive disability trajectory classification to have direct effects on healthcare service use including long-term care, inpatient care, and emergency services, and mediating effects from age and education to the use of same healthcare services among Taiwanese older adults [9]. Moreover, our results can be considered consistent

with several previous studies which have concluded disability [6, 26, 27] and poor physical functioning [28] to increase the utilization of healthcare services. Some evidence suggests for a two-way association between physical functioning and chronic diseases. Chronic diseases are a known risk factor for physical functioning [29]. Therefore, in addition to age, chronic diseases may have contributed to decline in physical functioning trajectory and healthcare use. On the other hand, declining physical functioning might predispose individuals for chronic diseases [30, 31], possibly requiring hospital care. Possibly supporting the association between chronic diseases and physical functioning, we found a higher prevalence of specific chronic diseases in the intermediate physical functioning class compared to the high trajectory class in Supplementary Table 1.

Further, we found quite uniformly elevated utilization rates of specialized healthcare services among participants in the intermediate physical functioning trajectory class. We examined four types of visits including emergency visits, first and follow-up outpatient visits, and hospital days, and found the estimated IRRs to vary between 1.36 and 1.58 among men and women in the intermediate physical functioning trajectory class. The lowest increases in utilization rates were found for hospital days. However, the differences in the rates across these four types of visits were negligible suggesting similar increases in specialized healthcare utilization in all four types of visits among those in intermediate physical functioning trajectory class. This aligns with our hypothesis

of physical functioning being an important factor to consider when studying healthcare service use among older adults.

We are not aware of studies to have examined the association between mental functioning trajectory classes and specialized healthcare use among older adults. In comparison to the *high* trajectory class, we found the *intermediate* mental functioning trajectory class to be associated with higher utilization rate of first outpatient visits among both men and women. The utilization rate was similar among both sexes. Although the literature regarding this topic is limited, it suggests that mental disorders, especially depression which is one of the most common mental health problems among older adults [32], are associated with outpatient [33–35] and inpatient care [34–36]. However, although both sexes in the *intermediate* mental functioning trajectory class had higher prevalence of depression and scored higher in Beck's Depression Inventory [37] compared to the *high* trajectory in Supplementary Table 2, we found no association between trajectory classification and inpatient care. It must be noted, however, that a lower level of mental functioning does not necessarily mean a mental disease. Thus, the results of previous studies regarding healthcare use and mental disorders may not be directly comparable to our study. Future studies are needed to examine the associations between mental health trajectory classification and specialized healthcare use.

Overall, our study highlights the importance of maintaining higher levels of functioning, which may also result in lower utilization rates of specialized healthcare services and, therefore, help maintaining the sustainability of specialized healthcare services while populations age. Although physiological processes in aging make decline in physical functioning inevitable, physical activity improves or slows the process of decline [38]. In general, physical activity is thought to change the trajectory of decline [38]. Therefore, exercise promotion programs to all older adults to support physical functioning may lead to cost-savings in healthcare. Additionally, lifestyle factors, like poor diet, obesity, and smoking, have been associated with poor physical functioning emphasizing the role of public health interventions in the promotion of healthy lifestyle [39, 40]. Since the differences between trajectory classes were observed since the beginning of the follow-up, the interventions should start early enough to prevent decline in physical functioning at older age. Further, good mental functioning in long-term might decrease the need for first outpatient visits while aging. Thus, supporting mental functioning at older age may be beneficial to both older adults and to the healthcare system. More research, however, is needed to understand

the associations between mental functioning trajectory classes and healthcare use before further conclusions for public health interventions can be reliably done.

The strengths of the current study are well collected data from a unique birth cohort and reliable register data. The study has, however, some limitations. First, since the clinical examinations were based on voluntariness, those with poor physical functioning may not have participated due to health reasons. This may have affected the formation of the trajectory classes. Second, the latent class mixture model is data-driven method, and we cannot address causality between physical functioning and increased specialized healthcare use, so it might be possible that greater healthcare use is linked to factors associated with poorer physical functioning. Third, the results may not be entirely generalizable outside Nordic countries since the healthcare systems of other countries differ. Finally, we did not have healthcare expenses available for our study.

In conclusion, poorer physical functioning trajectory classification among older adults was associated with greater specialized healthcare use. The *intermediate* mental health trajectory class, in turn, was only associated with the higher use of first outpatient visits. The associations found in the current study were similar among both sexes. Public health interventions should be considered to support physical functioning with aging. However, more research, especially regarding mental health trajectories and specialized healthcare use, is needed.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12877-023-04157-w>.

Additional file 1: Supplementary Table 1. Baseline characteristics by physical functioning trajectory classes among men and women. **Supplementary Table 2.** Baseline characteristics by mental functioning trajectory classes among men and women. **Supplementary Table 3.** Incidence rate ratios (IRRs) for healthcare service use across sex and physical functioning trajectory classes with membership probability of 0.85 or higher. **Supplementary Table 4.** Incidence rate ratios (IRRs) for healthcare service use across sex and mental functioning trajectory classes with membership probability of 0.85 or higher. **Supplementary Fig. 1.** Selection of the study participants. **Supplementary Fig. 2.** Observed (gray line), model-expectation (black line) and average (blue line) model-based trajectories for physical component score (PCS) in the intermediate and high declining classes among men and women. **Supplementary Fig. 3.** Observed (gray line), model-expectation (black line) and average (blue line) model-based trajectories for mental component score (MCS) in the intermediate and high stable classes among men and women.

Acknowledgements

Not applicable.

Authors' contributions

JNI designed the study, analyzed, and interpreted the data, and drafted the article. TT designed the study, analyzed, and interpreted the data, and revised the article critically for important intellectual content. MBVB, TAT, JGE, and MJH

designed the study, interpreted the data, and revised the article critically for important intellectual content. All authors approved the final version of the manuscript before its submission.

Funding

Open Access funding provided by University of Helsinki including Helsinki University Central Hospital. This study was supported by Avohoidon Tutkimussäätiö. The sponsor had no role in the design, methods, analysis, and preparation of the manuscript of the article.

Availability of data and materials

The datasets generated and/or analyzed during the current study are available from the corresponding author on reasonable request.

Declarations

Ethics approval and consent to participate

All study participants gave a written informed consent before participating in any clinical procedures. The procedures were approved by the Coordinating Ethics Committee of the Hospital District of Helsinki and Uusimaa.

Consent for publication

Not applicable.

Competing interests

The authors declare no competing interests.

Author details

¹Department of General Practice and Primary Health Care, University of Helsinki, Helsinki, Finland. ²Folkhälsan Research Center, Helsinki, Finland. ³Population Health Unit, Finnish Institute for Health and Welfare, Helsinki, Finland. ⁴Gerontology Research Center and Faculty of Sport and Health Sciences, University of Jyväskylä, Jyväskylä, Finland. ⁵Clinicum, Faculty of Medicine, University of Helsinki, Helsinki, Finland. ⁶Department of Obstetrics and Gynecology and Human Translational Research Programme, Yong Loo Lin School of Medicine, National University of Singapore, Singapore, Singapore. ⁷Agency for Science, Technology and Research (A*STAR), Singapore Institute for Clinical Sciences (SICS), Singapore, Singapore. ⁸Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden.

Received: 24 October 2022 Accepted: 6 July 2023

Published online: 21 July 2023

References

- OECD. Health at a Glance 2019: OECD Indicators. Paris: OECD Publishing; 2019. <https://doi.org/10.1787/4dd50c09-en>.
- World Health Organization. Decade of healthy ageing: baseline report. Geneva: World Health Organization; 2020. <https://www.who.int/publications/i/item/9789240017900>. Accessed 14 Jun 2023.
- Han L, Allore H, Murphy T, Gill T, Peduzzi P, Lin H. Dynamics of functional aging based on latent-class trajectories of activities of daily living. *Ann Epidemiol*. 2013;23:87. <https://doi.org/10.1016/J.ANNEPIDEM.2012.11.010>.
- Liang J, Xu X, Bennett JM, Ye W, Quiñones AR. Ethnicity and changing functional health in middle and late life: a person-centered approach. *J Gerontol Ser B Psychol Sci Soc Sci*. 2010;65B:470. <https://doi.org/10.1093/GERONB/GBP114>.
- Carey EC, Walter LC, Lindquist K, Covinsky KE. Development and validation of a functional morbidity index to predict mortality in community-dwelling elders. *J Gen Intern Med*. 2004;19:1027. <https://doi.org/10.1111/J.1525-1497.2004.40016.X>.
- Fried TR, Bradley EH, Williams CS, Tinetti ME. Functional disability and health care expenditures for older persons. *Arch Intern Med*. 2001;161:2602–7. <https://doi.org/10.1001/ARCHINTE.161.21.2602>.
- Gaugler JE, Duval S, Anderson KA, Kane RL. Predicting nursing home admission in the U. S: a meta-analysis. *BMC Geriatr*. 2007;7:1–14. <https://doi.org/10.1186/1471-2318-7-13/FIGURES/3>.
- Xiao J, Shi Z, Fang Y. Association between disability trajectory and health care service utilization among older adults in China. *J Am Med Dir Assoc*. 2021;22:2169–2176.e4. <https://doi.org/10.1016/J.JAMDA.2020.12.041>.
- Yu HW, Tu YK, Chen YM. Sociodemographic characteristics, disability trajectory, and health care and long-term care utilization among middle-old and older adults in Taiwan. *Arch Gerontol Geriatr*. 2019;82:161–6. <https://doi.org/10.1016/J.ARCHGER.2019.01.019>.
- Sialino LD, Schaap LA, Van Oostrom SH, Nooyens ACJ, Picavet HSJ, Twisk JWR, et al. Sex differences in physical performance by age, educational level, ethnic groups and birth cohort: the Longitudinal Aging Study Amsterdam. *PLoS ONE*. 14(12): e0226342. <https://doi.org/10.1371/journal.pone.0226342>.
- Sialino LD, Schaap LA, van Oostrom SH, Picavet HSJ, Twisk JWR, Verschuren WMM, et al. The sex difference in gait speed among older adults: how do sociodemographic, lifestyle, social and health determinants contribute? *BMC Geriatr*. 2021;21:1–11. <https://doi.org/10.1186/S12877-021-02279-7/TABLES/3>.
- World Health Organization. Mental health of older adults. 2023. <https://www.who.int/news-room/fact-sheets/detail/mental-health-of-older-adults>. Accessed 24 Aug 2021.
- Sialino LD, van Oostrom SH, Wijnhoven HAH, Picavet S, Verschuren WMM, Visser M, et al. Sex differences in mental health among older adults: investigating time trends and possible risk groups with regard to age, educational level and ethnicity. *Aging Ment Health*. 2020;25:2355–64. <https://doi.org/10.1080/1360786320201847248>.
- Luppa M, Sikorski C, Luck T, Ehreke L, Konnopka A, Wiese B, et al. Age- and gender-specific prevalence of depression in latest-life—systematic review and meta-analysis. *J Affect Disord*. 2012;136:212–21. <https://doi.org/10.1016/J.JAD.2010.11.033>.
- Barker DJP, Osmond C, Forsén TJ, Kajantie E, Eriksson JG. Trajectories of growth among children who have coronary events as adults. *N Engl J Med*. 2005;353:1802–9.
- Fayers PM, Machin D. Quality of Life: the assessment, analysis and interpretation of patient-reported outcomes: second edition. John Wiley and Sons. 2007. <https://doi.org/10.1002/9780470024522>.
- Ware John E JR, Sherbourne CD. The MOS 36-item short-form health survey (SF-36). I. Conceptual framework and item selection *PubMed*. *Med Care*. 1992;30:473–83.
- Walters SJ, Munro JF, Brazier JE. Using the SF-36 with older adults: a cross-sectional community-based survey. *Age Ageing*. 2001;30:337–43. <https://doi.org/10.1093/AGEING/30.4.337>.
- Mishra GD, Gale CR, Sayer AA, Cooper C, Dennison EM, Whalley LJ, et al. How useful are the SF-36 sub-scales in older people? Mokken scaling of data from the HALCYON programme. *Qual Life Res*. 2011;20:1005. <https://doi.org/10.1007/S11136-010-9838-7>.
- Ware JE, Kosinski M. SF-36 physical and mental health summary scales: A manual for users of version 1. 2nd ed. Lincoln, RI: Quality Metric; 2001.
- Taft C, Karlsson J, Sullivan M. Do SF-36 summary component scores accurately summarize subscale scores? *Qual Life Res*. 2001;10:395–404. <https://doi.org/10.1023/A:1012552211996>.
- National Institute for health and welfare. Care Register for Health Care. Register description. 2016. <https://www.thl.fi/en/web/thlfi-en/statistics/information-on-statistics/register-descriptions/care-register-for-health-care>. Accessed 1 Nov 2020.
- Statistics Finland. Education. 2021. https://www.stat.fi/til/kou_en.html. Accessed 18 May 2021.
- Lakka TA, Salonen JT. Intra-person variability of various physical activity assessments in the kuopio ischaemic heart disease risk factor study. *Int J Epidemiol*. 1992;21:467–72. <https://doi.org/10.1093/ije/21.3.467>.
- Wang M, Bodner TE. Growth Mixture Modeling: Identifying and Predicting Unobserved Subpopulations With Longitudinal Data. *Organ Res Methods*. 2007;10:635–56. <https://doi.org/10.1177/1094428106289397>.
- Laudisio A, Marzetti E, Franceschi F, Bernabei R, Zuccalà G. Disability is associated with emergency room visits in the elderly: a population-based study. *Aging Clin Exp Res*. 2015;27:663–71. <https://doi.org/10.1007/s40520-015-0324-5>.
- Wu CY, Hu HY, Li CP, Fang YT, Huang N, Chou YJ. The association between functional disability and acute care utilization among the elderly in Taiwan. *Arch Gerontol Geriatr*. 2013;57:177–83. <https://doi.org/10.1016/j.archger.2013.04.011>.

28. Cheng Y, Goodin AJ, Pahor M, Manini T, Brown JD. healthcare utilization and physical functioning in older adults in the United States. *J Am Geriatr Soc.* 2020;68:266–71. <https://doi.org/10.1111/jgs.16260>.
29. Manini T. Development of physical disability in older adults. *Curr Aging Sci.* 2011;4:184. <https://doi.org/10.2174/1874609811104030184>.
30. Ibarra AD, Johnson WH. Disability Status as an Antecedent to Chronic Conditions: National Health Interview Survey, 2006–2012. *Prev Chronic Dis.* 2014;11:130251. <https://doi.org/10.5888/PCD11.130251>.
31. Rasch EK, Magder L, Hochberg MC, Magaziner J, Altman BM. Health of Community-Dwelling Adults With Mobility Limitations in the United States: Incidence of Secondary Health Conditions. Part II *Arch Phys Med Rehabil.* 2008;89:219–30.
32. Zenebe Y, Akele B, Selassie MW/, Necho M. Prevalence and determinants of depression among old age: a systematic review and meta-analysis. *Ann Gen Psychiatry* 2021. 2021;20:1–19. <https://doi.org/10.1186/S12991-021-00375-X>.
33. Adepoju O, Lin SH, Mileski M, Kruse CS, Mask A. Mental health status and healthcare utilization among community dwelling older adults. *J Ment Health.* 2018;27:511–9. <https://doi.org/10.1080/09638237.2018.1466030>.
34. Huang H, Menezes PR, Da Silva SA, Tabb K, Barkil-Oteo A, Scazufca M. The association between depressive disorders and health care utilization: results from the São Paulo ageing and health study (SPAH). *Gen Hosp Psychiatry.* 2014;36:199–202. <https://doi.org/10.1016/J.GENHOSPPSYCH.2013.11.003>.
35. Guo J, Kong D, Fang L, Zhu Y, Zhang B. Depressive symptoms and health service utilization among Chinese middle-aged and older adults: a national population-based longitudinal survey. *Int J Ment Health Syst.* 2021;15:1–9. <https://doi.org/10.1186/S13033-020-00421-3/TABLES/4>.
36. Prina AM, Deeg D, Brayne C, Beekman A, Huisman M. The Association between Depressive Symptoms and Non-Psychiatric Hospitalisation in Older Adults. *PLoS One.* 7(4):e34821. <https://doi.org/10.1371/JOURNAL.PONE.0034821>.
37. Beck AT, Ward CH, Mendelson M, Mock J, Erbaugh J. An inventory for measuring depression. *Arch Gen Psychiatry.* 1961;4:561–71. <https://doi.org/10.1001/ARCHPSYC.1961.01710120031004>.
38. O'Neill D, Forman DE. The importance of physical function as a clinical outcome: assessment and enhancement. *Clin Cardiol.* 2020;43:108. <https://doi.org/10.1002/CLC.23311>.
39. Li D, Jia Y, Yu J, Liu Y, Li F, Liu Y, et al. Adherence to healthy lifestyle and the risk of function limitations in late life: the atherosclerosis risk in communities study. *Front Aging Neurosci.* 2021;13:446.
40. Robinson SM, Jameson KA, Syddall HE, Dennison EM, Cooper C, Aihie SA. Clustering of lifestyle risk factors and poor physical function in older adults: The Hertfordshire Cohort Study. *J Am Geriatr Soc.* 2013;61:1684. <https://doi.org/10.1111/JGS.12457>.

Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

