

JYX



This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): KDK Collaboration

Title: Evidence for ground-state electron capture of 40K

Year: 2023

Version: Published version

Copyright: © 2023 American Physical Society (APS)

Rights: In Copyright

Rights url: <http://rightsstatements.org/page/InC/1.0/?language=en>

Please cite the original version:

KDK Collaboration. (2023). Evidence for ground-state electron capture of 40K. Physical Review C, 108, Article 014327. <https://doi.org/10.1103/PhysRevC.108.014327>

Evidence for ground-state electron capture of ^{40}K

L. Hariasz¹, M. Stukel¹, P. C. F. Di Stefano^{1,*}, B. C. Rasco², K. P. Rykaczewski², N. T. Brewer^{2,3}, D. W. Stracener², Y. Liu², Z. Gai⁴, C. Rouleau⁴, J. Carter⁵, J. Kostensalo⁶, J. Suhonen⁷, H. Davis^{8,9}, E. D. Lukosi^{8,9}, K. C. Goetz¹⁰, R. K. Grzywacz^{2,3,11}, M. Mancuso¹², F. Petricca¹², A. Fijałkowska¹³, M. Wolińska-Cichočka^{2,3,14}, J. Ninković¹⁵, P. Lechner¹⁵, R. B. Ickert¹⁶, L. E. Morgan¹⁷, P. R. Renne^{5,18} and I. Yavin[†]

(KDK Collaboration)

¹Department of Physics, Engineering Physics & Astronomy, Queen's University, Kingston, Ontario K7L 3N6, Canada

²Physics Division, Oak Ridge National Laboratory, Oak Ridge, Tennessee 37831, USA

³Joint Institute for Nuclear Physics and Application, Oak Ridge National Laboratory, Oak Ridge, Tennessee 37831, USA

⁴Center for Nanophase Materials Sciences, Oak Ridge National Laboratory, Oak Ridge, Tennessee 37831, USA

⁵Berkeley Geochronology Center, Berkeley, California 94709, USA

⁶Natural Resources Institute Finland, Joensuu FI-80100, Finland

⁷Department of Physics, University of Jyväskylä, Jyväskylä FI-40014, Finland

⁸Department of Nuclear Engineering, University of Tennessee, Knoxville, Tennessee 37996, USA

⁹Joint Institute for Advanced Materials, University of Tennessee, Knoxville, Tennessee 37996, USA

¹⁰Nuclear and Extreme Environments Measurement Group, Oak Ridge National Laboratory, Oak Ridge, Tennessee 37831, USA

¹¹Department of Physics and Astronomy, University of Tennessee, Knoxville, Tennessee 37996, USA

¹²Max-Planck-Institut für Physik, Munich D-80805, Germany

¹³Faculty of Physics, University of Warsaw, Warsaw PL-02-093, Poland

¹⁴Heavy Ion Laboratory, University of Warsaw, Warsaw PL-02-093, Poland

¹⁵MPG Semiconductor Laboratory, Munich D-80805, Germany

¹⁶Department of Earth, Atmospheric, and Planetary Sciences, Purdue University, West Lafayette, Indiana 47907, USA

¹⁷U.S. Geological Survey, Geology, Geophysics, and Geochemistry Science Center, Denver, Colorado 80225, USA

¹⁸Department of Earth and Planetary Science, University of California, Berkeley 94720, USA



(Received 22 November 2022; accepted 23 May 2023; published 31 July 2023)

Potassium-40 is a widespread, naturally occurring isotope whose radioactivity impacts estimated geological ages spanning billions of years, nuclear structure theory, and subatomic rare-event searches—including those for dark matter and neutrinoless double-beta decay. The decays of this long-lived isotope must be precisely known for its use as a geochronometer, and to account for its presence in low-background experiments. There are several known decay modes for potassium-40, but a predicted electron-capture decay directly to the ground state of argon-40 has never been observed. The existence of this decay mode impacts several fields, while theoretical predictions span an order of magnitude. Here we report on the first, successful observation of this rare decay mode, obtained by the KDK (potassium decay) Collaboration using a novel combination of a low-threshold x-ray detector surrounded by a tonne-scale, high-efficiency γ -ray tagger at Oak Ridge National Laboratory. A blinded analysis reveals a distinctly nonzero ratio of intensities of ground-state electron-captures (I_{EC^0}) over excited-state ones (I_{EC^*}) of $I_{\text{EC}^0}/I_{\text{EC}^*} = 0.0095 \pm 0.0022 \pm 0.0010$ (68% CL), with the null hypothesis rejected at 4σ [Stukel *et al.*, *Phys. Rev. Lett.* **131**, 052503 (2023)]. In terms of branching ratio, this unambiguous signal yields $I_{\text{EC}^0} = 0.098\% \pm 0.023\% \pm 0.010\%$, roughly half of the commonly used prediction. This first observation of a third-forbidden unique electron capture improves our understanding of low-energy backgrounds in dark-matter searches and has implications for nuclear-structure calculations. For example, a shell-model based theoretical estimate for the neutrinoless double-beta decay half-life of calcium-48 is increased by a factor of 7_{-2}^{+3} . Our nonzero measurement shifts geochronological ages by up to a percent; implications are illustrated for Earth and solar system chronologies.

DOI: [10.1103/PhysRevC.108.014327](https://doi.org/10.1103/PhysRevC.108.014327)

I. INTRODUCTION

Potassium-40 is a common, natural isotope. It decays mainly by β^- decay, less frequently by electron capture to

an excited state of argon-40 (EC^*), and very rarely by β^+ decay (see Fig. 1). It is a frequent contaminant in various particle detectors, and a source of radioactive background in rare-event searches for dark matter [1–6] and neutrinoless double-beta decay [7]. Several experiments looking for dark matter are taking draconian steps to purify their NaI detectors of K, and/or deploy vetoes to tag the problematic low-energy

*distefan@queensu.ca

†yavin.itay@gmail.com




FIG. 1. ^{40}K decay scheme, with branching ratios and half-life calculated from our determination of $I_{\text{EC}^0}/I_{\text{EC}^*}$ (this work and [14]) and from literature values for T^- (partial half-life of the β^- decay) and T^* (partial half-life of EC^*) [15] (also shown: transition energy [16], Q_{EC^0} and Q^- [17]).

radiation from ^{40}K electron capture which falls in the expected dark matter signal region [3,4,6]. This veto method relies on identification of the high-energy γ ray from the de-excitation of ^{40}Ar following an EC^* decay. In addition, the long half-life (slightly over a billion years) of this isotope makes it a useful geochronometer via the K/Ar and $^{40}\text{Ar}/^{39}\text{Ar}$ dating techniques [8–11]. Finally, the presence of all three modes of β decay, some of which are extremely rare third-forbidden unique transitions, make this isotope of particular interest to nuclear structure theory [7,12,13].

Aside from the branches mentioned previously, an electron-capture decay directly to the ground state of ^{40}Ar (EC^0) has also been predicted by some [9,13,15,18], ignored or disputed by others [10,11], but never observed. This branch forms a particularly challenging background in rare-event searches as there is no high-energy γ ray from deexcitation that could be used to tag the low-energy radiation from the electron capture. Such a background would evade the veto technique mentioned previously. In addition, this added

background has been proposed [2] as a way to constrain the dark-matter interpretation of the longstanding, but controversial, DAMA/LIBRA Collaboration claim for discovery of dark matter [19]. From the standpoint of geochronology, the existence of this decay to ground state could mean that samples are up to tens of millions of years older than commonly believed [9]. An empirical frequency of this branch would also inform calculations in nuclear structure theory, including those for neutrinoless double-beta-decay half-lives. The KDK (potassium decay) Collaboration [20] has carried out the first measurement of EC^0 [14], using a novel detector configuration [21], as detailed in what follows.

II. DETECTOR AND ANALYSIS

The fully characterized KDK experimental setup [21] consists of a ^{40}K source, a sensitive x-ray silicon drift detector (SDD) and a near-100% efficient γ tagger (the Modular Total Absorption Spectrometer, MTAS [22]), as illustrated in Fig. 2.

The enriched ^{40}K source was made of KCl [16.1(6)% ^{40}K abundance in K] thermally deposited over 1 cm diameter onto a graphite substrate (all uncertainties in this paper correspond to a 68% confidence level). The $\approx 9 \times 10^{17}$ atoms of ^{40}K in the source have an activity of ≈ 16 Bq, equivalent to two bananas [23], and the source is 5.1(9) μm thin to allow the x rays to escape from it. The source rests directly in front of the SDD and is centered inside MTAS.

MTAS is a 1 tonne array of NaI scintillators with $\approx 98\%$ tagging efficiency for the 1.46 MeV γ rays of interest (Table 7 in [21]). Data with the ^{40}K source were acquired over 44 days. Following offline determination of coincidences between the SDD and MTAS for three nominal time windows of (1, 2, 4) μs , SDD pulses were fit and energy calibrations were performed. To avoid biases during the analysis, the anticoincident SDD spectrum was blinded from (0.88–1.4)




FIG. 2. Schematic displaying the cross-section of MTAS, along with the SDD. The SDD housing is centered in MTAS.




FIG. 3. ^{40}K coincidence spectrum between the SDD and MTAS. SDD and MTAS data obtained with the enriched ^{40}K source using a nominal $2\ \mu\text{s}$ coincident window is binned into a 250×250 grid over the displayed energy ranges.

keV (silicon escape peak region) and (2.0–3.8) keV (electron capture signal region) while cuts and analysis methods were established.

Over the course of the run, two types of SDD instabilities appeared: gain drops due to voltage supply failures that were reset by an operator, and noise bursts attributed to power fluctuations in the laboratory. Both can be identified in the SDD energy range below the blinded region; after this cut, 76% of the live time remains. Minor gain drifts in the SDD were corrected by tracking the coincidence x-ray lines.

Gain drifts of a few percent were also observed in MTAS. A Kolmogorov-Smirnov test comparing the arrival times of the open anticoincident and coincident SDD events in the 2–5 keV region returned a p value of 0.63. This implies both data sets are consistent with the same underlying time distribution, and is consistent with the tagging efficiency of MTAS not changing over the run. In addition, before opening the data set, five time regions with MTAS gain changes were identified. After opening the data, the full ρ analysis (detailed in later sections) was carried out on each of these regions, and the values were compared. A fit to a single common value of ρ yielded a χ^2 of 3.6 for four degrees of freedom, i.e., $p = 0.45$, implying the data are consistent with a constant tagging efficiency. Consistent results were observed over all three coincidence windows.

A. Physical phenomena visible in the SDD and MTAS spectra

A coincidence spectrum of the ^{40}K source with a $2\ \mu\text{s}$ nominal window can be seen in Fig. 3, which bins events by SDD and MTAS energies, resulting in various bands and peaks. The foci are near (3 keV, 1.46 MeV), corresponding to EC^* decays involving x-ray detection in the SDD and complete capture of the deexcitation γ in MTAS. Additional features in Fig. 3




FIG. 4. SDD coincidence and anticoincidence spectra. Results of simultaneous fit to coincident (top) and anticoincident (bottom) SDD spectra at a $2\ \mu\text{s}$ coincidence window. Signal counts are shown in green. Various fluorescence peaks and an exponential background model are included. The total minimization has an associated goodness of fit of $p = 0.4$.

involve partial energy depositions of EC^* decay products. The projection of this figure, with nonzero MTAS energy, onto the SDD energy, is the coincident SDD data spectrum in Fig. 4.

In addition to the Ar signal, characteristic x rays of Cl, K, and Ca are seen in the signal region (see Table I for the number of events, as well as Fig. 2 of [14]). K and Cl atoms are numerous enough in the source to be fluoresced by source β^- . Most β^- -fluoresced events are anticoincident, with a small coincident contribution primarily from the β^- which make it into MTAS. Relative coincident and anticoincident K counts are consistent with this description. Though Cl is also fluoresced by β^- , Ar x rays (generally from EC^*) are energetic enough to fluoresce Cl, contributing additional coincident events.

The Ar and Ca atoms in the source come from the slow decay of ^{40}K . There are therefore 9–10 orders of magnitude fewer Ar and Ca atoms than K or Cl ones in the source.

TABLE I. Fluorescence counts observed in SDD spectra. Fluorescence may occur via various sources, described in the text, which affect the coincident sorting of the resulting x-ray detection. The quoted values are obtained using a $2\ \mu\text{s}$ window between the SDD and MTAS.

Element	Coincident counts	Anticoincident counts
Cl	$3.48(2) \times 10^3$	$8.85(5) \times 10^3$
K	$2.9(3) \times 10^2$	$5.89(7) \times 10^3$
Ca	$1.0(5) \times 10$	$1.36(5) \times 10^3$

Based on the limited K fluorescence visible in KDK data, fluorescence of an Ar or Ca atom caused by something other than the atom itself is highly negligible.

Interestingly, there is a small contribution of characteristic Ca x rays visible in the anticoincident spectrum. Having ruled out external fluorescence, a product of the $^{40}\text{K} \rightarrow ^{40}\text{Ca}$ decay itself must produce these events; the decay to Ca must occasionally involve *self-fluorescence* of the daughter via the produced β^- . The exact low-order processes involved in such interactions are beyond the scope of this work, and we note that the Ca contribution has no effect on analysis of the Ar x rays.

Outside of the SDD signal region, an additional presentation of source EC* decays is contained at (1.2 keV, 1.46 MeV). This silicon escape peak is formed in the event that a source x-ray fluoresces a detector Si atom prior to detection. The remaining energy of the source x ray is equivalent to the difference between its initial energy and that of the fluoresced Si x ray (1.7 keV), resulting in a detectable 1.2 keV event.

Lastly, though it is possible for source γ rays to deposit energy in the SDD, this is modeled to occur in <0.5% of cases. This contribution is considered in the systematic analysis described further, though the effect is negligible relative to the statistical error of our measurement. β^+ from the source would eventually provide a small contribution to the continuous SDD background, though over the KDK runtime any such events are negligible due to the minute branching ratio of this mode.

B. Coincident and anticoincident events

In the KDK dataset, we expect a total of σ^* counts from EC* decays, and σ counts from EC⁰ decays present in our signal (Ar x-ray) peaks. These are expanded to

$$\begin{aligned}\sigma^* &= AT I_{\text{EC}^*} P_K^* \omega_K \eta (1 - \eta_\gamma), \\ \sigma &= AT I_{\text{EC}^0} P_K \omega_K \eta,\end{aligned}$$

where AT are total source decays over the run duration. K -shell capture probabilities $P_K^* = 0.7564(4)$, $P_K = 0.8908(7)$ using BETASHAPE code V.2.2 [24] differ for the two modes, though the fluorescence probability ω_K is the same. Both modes emit the same x rays, whose detection probability in the SDD is η . The γ -ray accompanying the EC* decay could deposit some energy in the SDD, with a small probability, thus shifting the event out of the signal region. From simulations, we estimate this probability to be $\eta_\gamma = 0.0048(48)$.

Various factors will sort σ and σ^* into coincident and anticoincident events; the main ones are

- (1) The efficiency with which the γ 's from EC* are tagged by MTAS is not perfect. We previously studied this parameter ϵ and determined it with high precision to be $\epsilon = 0.9792 \pm 0.006$ at a 2 μs coincidence window [21] using data and GEANT4 [25] simulations. This will reduce the expected number of EC* events that are properly tagged to $\sigma^* \epsilon$.
- (2) Conversely, the nonperfect tagging efficiency will lead to $\sigma^*(1 - \epsilon)$ EC*s being untagged (false positives).

- (3) In addition, some EC⁰ events may be in spurious coincidence with the MTAS background; the probability that this occurs in the $\mathcal{O}(\mu\text{s})$ coincidence window is $\beta_M \bar{t} \ll 1$ [21] (false negatives).
- (4) The remaining EC⁰ events are anticoincident.

These subsets of signal counts are summarized below:

$$\begin{aligned}\Sigma^* &= \overbrace{\sigma^* \epsilon}^{1. \text{ Observed EC}^*} + \overbrace{\sigma \beta_M \bar{t}}^{3. \text{ False negatives}} \\ \Sigma &= \overbrace{\sigma^*(1 - \epsilon)}^{2. \text{ False positives}} + \overbrace{\sigma}^{4. \text{ Observed EC}^0}\end{aligned}$$

where Σ^* are expected coincident counts, and Σ are expected anticoincident counts. We introduce a parameter for total signal counts, $\nu \equiv \sigma^* + \sigma$, and a useful term

$$\rho' \equiv \frac{\sigma}{\sigma^*} = \rho \frac{P_K}{P_K^*} \frac{1}{(1 - \eta_\gamma)}.$$

Using these two new expressions, we obtain the final likelihood terms relating ρ to expected coincident and anticoincident counts:

$$\begin{aligned}\Sigma^* &= \frac{\nu}{1 + \rho'} (\epsilon + \rho' \beta_M \bar{t}), \\ \Sigma &= \frac{\nu}{1 + \rho'} (1 - \epsilon + \rho').\end{aligned}\quad (1)$$

C. Likelihood method

SDD data are sorted by coincidence, and both subsets are fit simultaneously through minimization of the sum of the negative log-likelihoods:

$$-\ln \mathcal{L} = -(\ln \mathcal{L}_{\text{coin}} + \ln \mathcal{L}_{\text{anti}})$$

Each of the two terms is a binned Poisson likelihood ratio [26]:

$$-\ln \mathcal{L}_j = \sum_i \left[f(x_{i,j}; \boldsymbol{\theta}) - n_{i,j} + n_{i,j} \ln \left(\frac{n_{i,j}}{f(x_{i,j}; \boldsymbol{\theta})} \right) \right].$$

Above, index j represents either coincident or anticoincident data, $n_{i,j}$ are total events in bin i and $f(x_{i,j})$ are the model-predicted counts in that bin. In addition to providing estimators of the parameters and confidence intervals, this technique returns goodness of fit [26]. Some of the parameters in $\boldsymbol{\theta}$, such as our main one $\rho = I_{\text{EC}^0}/I_{\text{EC}^*}$ and those pertaining to the shapes of the lines, are shared between the coincident and anticoincident data, while others are not.

The spectra contain several fluorescent contributions (Cl, K, Ca). For each, we model the associated K_α and K_β x rays with Gaussian distributions, the means of which are fixed to known values [27]. For each such $K_\alpha + K_\beta$ pair, the free Gaussian width is shared across the two peaks, and is the same in both coincident and anticoincident spectra. A parameter for total $K_\alpha + K_\beta$ counts, not shared across the coincidence-sorted data, is left free. The ratio of intensities I_{K_α}/I_{K_β} is generally fixed to values in [28].

The continuous background model, consisting of decaying exponential and flat components, has all associated parameters left free. These parameters are not shared across the

TABLE II. Systematic, 68% C.L. errors on ρ . All sources of error, described in the text, are smaller than the statistical (0.0022).

Source	Systematic error
Fit range	9×10^{-4}
MTAS γ -ray-tagging efficiency	5×10^{-4}
Binning	1×10^{-4}
SDD γ -ray-tagging efficiency	4×10^{-5}
K -shell capture probabilities	8×10^{-6}
Expected MTAS background counts	3×10^{-6}

coincident and anticoincident data, as this background contribution has a different shape in each subset.

The Ar K_α and K_β x rays of interest are modeled in a similar manner as the fluorescence lines. In order for these components to directly inform ρ , we insert the expression for expected coincident and anticoincident Ar counts of Eq. (1) directly into the likelihood. This introduces free parameters ρ and total Ar counts, along with fixed terms including efficiencies, as described earlier in Sec. II B.

We note that the result ρ is stable to the choice of fixing or freeing the ratio of K_α to K_β intensities in the Ar and fluorescent components. Modeling of the latter generally has a negligible effect on the result, which is informed only by the Ar lines. An additional test modelled the signal x rays with a Voigt profile [29], which yields essentially identical results since it approaches the limiting Gaussian case.

Initial opening of the data led to a value for ρ of 0.008 ± 0.002 on the single chosen energy range before evaluation of systematic errors. A thorough analysis of systematics leads to our reported value of $\rho = I_{EC^0}/I_{EC^*} = 0.0095 \pm 0.0022 \pm 0.0010$. An example of a fit is shown in Fig. 4.

D. Systematics

The systematic errors pertinent to the experiment fall into two categories: fit characteristics and physical limitations. The choice of fit range (canonical fit range is 1.5–5.0 keV) pertaining to the mathematical background model is the dominant source of systematic error. The dominant physical error arises from the imperfect γ -ray tagging efficiency of MTAS. Table II contains a summary of each source of error and its effect.

To account for possible covariances between parameters which contribute to the overall systematic error, their effect is tested simultaneously by randomly drawing their value prior to a likelihood fit. Fit characteristics are drawn from a uniform distribution over the considered range. Physical parameters are drawn from a Gaussian whose width corresponds to the parameter's known uncertainty. This process is repeated 10,000 times to obtain a distribution of ρ , whose mean corresponds to our final measurement, and width is the systematic error, as shown in Fig. 5.

Though the systematic check described above uses a background model consisting of exponential and flat components, other models have been explored in depth. We have tested polynomials up to the third degree over various ranges, and




FIG. 5. Distributions of ρ and its statistical error from systematic checks. The value of ρ and its associated statistical error $\sigma_\rho^{\text{stat}}$ obtained from 10 000 systematic-varying fits are displayed. The width of the ρ distribution is equivalent to the systematic error. Contour lines of the fit function correspond to the color bar on the right.

find them to either be too simple or to find unphysical minima and maxima in the data. Overall, we find other models to either be too simple, find unphysical artefacts, or supply extraneous degrees of freedom to the canonical (exponential) case.

Lastly, our analysis was performed at 3 coincidence windows: (1, 2, and 4 μs , for which we find consistent results of $\rho = (0.0091, 0.0095, 0.0095)$ respectively, all with the same error $\pm 0.0022 \pm 0.0010$. These values are consistent with the value initially found upon opening the data and reported in the previous section. Ar counts obtained in the systematic check at 2 μs are summarized in Table III.

III. IMPLICATIONS

The following re-evaluation of the ^{40}K decay scheme using our measurement affects a variety of fields, including nuclear structure theory and geochronology. This is discussed below, illustrating impacts on neutrinoless double-beta decay of ^{48}Ca and the thermal history of the Acaapulco meteorite.

TABLE III. $^{40}\text{K} \rightarrow ^{40}\text{Ar}$ electron capture decays visible in 2 μs KDK data. The total ≈ 3 keV signal counts pertaining to this transition are sorted by coincidence, with ground-state decays contributing primarily to the anticoincident subset. Notations in parentheses refer to Sec. II B.

Component	Visible counts
Total Ar ($\Sigma^* + \Sigma$)	4.78×10^4
Coincident Ar (Σ^*)	4.63×10^4
Anticoincident Ar (Σ)	1.50×10^3
EC^0 decay (σ)	5.00×10^2

TABLE IV. Re-evaluation of the ^{40}K decay scheme. Branching ratios I and total half-life $T_{1/2}$ are calculated from our measurement of $\rho = I_{\text{EC}^0}/I_{\text{EC}^*}$, evaluation of measured partial EC* and β^- half-lives [15], and either the measured relative β^+/β^- feeding [18] or the predicted value of $I_{\text{EC}^0}/I_{\beta^+}$ [12]. The choice only affects the β^+ branching.

Quantity	Value	Uncertainty (68%C.L.)
I_{EC^0} (%)	0.098	$\pm 0.023(\text{stat}) \pm 0.010(\text{syst})$
I_{EC^*} (%)	10.31	± 0.04
I_{β^-} (%)	89.59	± 0.05
I_{β^+} (%) (expt)	0.00100	± 0.00013
I_{β^+} (%) (theory)	0.00045	± 0.00012
$T_{1/2}(\text{Ga})$	1.266	± 0.004

A. Construction of the decay scheme of ^{40}K

Constructing the full decay scheme for ^{40}K requires 4 parameters. Two of these are partial decay constants for the β^- and EC* branches, $\lambda^- = 0.4904 \pm 0.0019 \text{ Ga}^{-1}$ and $\lambda^* = 0.05646 \pm 0.00016 \text{ Ga}^{-1}$ (where Ga denotes 10^9 yr), taken from the most recent data evaluation (Sec. 5.2 of [15]). These absolute measurements are independent of EC⁰, though they depend on factors like the precise ^{40}K content of the source, and the efficiency of the detectors. Generally speaking, the other two parameters used are an experimental determination of I_{β^+}/I_{β^-} [18] and a theoretical value for $I_{\text{EC}^0}/I_{\beta^+}$ [12].

The former directly leads to $\lambda^+ = \frac{I_{\beta^+}}{I_{\beta^-}} \lambda^-$, while the latter then provides $\lambda^0 = \frac{I_{\text{EC}^0}}{I_{\beta^+}} \lambda^+$. From this complete set of partial decay constants, the total decay constant can be obtained: $\lambda = \sum_i \lambda_i$. Partial and total half-lives are then determined ($T_i = \frac{\ln 2}{\lambda_i}$). Lastly, branching ratios ($P_i = \frac{\lambda_i}{\sum_j \lambda_j}$) are obtained.

Our novel measurement provides an additional, experimental, value: $I_{\text{EC}^0}/I_{\text{EC}^*}$. In conjunction with λ^* , it leads directly to $\lambda^0 = \frac{I_{\text{EC}^0}}{I_{\text{EC}^*}} \lambda^*$. The value of λ^0 is the same, within uncertainties, for various commonly used sets of decay constants [10,15,27]. We keep λ^- , as before. To complete the set of four parameters with λ^+ , we need to choose between I_{β^+}/I_{β^-} and $I_{\text{EC}^0}/I_{\beta^+}$. This choice only affects the decay scheme at the level of the small β^+ branch, which varies by a factor of 2, as Table IV shows. This discrepancy advocates for further experimental and theoretical work on the β^+ branch.

B. Nuclear shell-model calculations

We obtain a theoretical estimate for the third-forbidden unique I_{EC^0} transition, that is complementary to the main experimental result, as described further. Moreover, the (in)frequency of this decay informs the extent to which such forbidden modes are suppressed, which is applicable to calculations of neutrinoless double-beta decay half-lives including that of ^{48}Ca . This commonly overlooked suppression, quantified as quenching of the weak axial-vector coupling, can significantly increase calculated $0\nu\beta\beta$ half-lives.

We calculate a value of $I_{\text{EC}^0} = 0.058 \pm 0.022$ using the Behrens-Bühring formalism ([30] for the full theory and




FIG. 6. Measured ground-state electron-capture branching ratio compared to predictions from Engelkemeir (1962) [18], log ft (1999) [34,35], Mougeot (2019) [12], Carter (2020) [9], Kossert (2022) [15], and this work (Sec. III B). All error bars correspond to a 68% C.L., with no uncertainty reported for Engelkemeir.

[31] for a streamlined version) with the nuclear matrix elements calculated in the shell-model framework using the code NUSHELLX@MSU [32] with the Hamiltonian $sdpfk$ [33]. It is consistent within uncertainties with a value obtained prior to the experiment. Our theoretical estimate is compared to other predictions along with the KDK measurement of this work in Fig. 6.

Using this Hamiltonian the half-lives of the three decay branches, β^- , EC*, and EC⁰, could be calculated. The corresponding decay amplitudes are proportional to the weak axial-vector coupling g_A , the value of which is known to be quenched for a wide range of nuclear masses [36]. Here the effective value of it can be determined by comparison of the computed and experimental half-lives, giving for the first-forbidden unique transition EC* the value $g_A^{\text{eff}} = 0.34$ and for the two third-forbidden unique transitions a value $g_A^{\text{eff}} = 0.43$ for β^- branch and $g_A^{\text{eff}} = 0.53$ for the EC⁰ branch. These values of g_A^{eff} are very well in line with the values $g_A^{\text{eff}} = 0.43\text{--}0.66$ obtained in the mass range $A = 74\text{--}126$ for the $2^- \leftrightarrow 0^+$ first-forbidden unique β and EC transitions in the framework of the proton-neutron quasiparticle random-phase approximation [37]. These results suggest that the forbidden contributions to the nuclear matrix elements of the $0\nu\beta\beta$ decay are very much suppressed and the resulting half-lives are much longer than expected based on the bare value $g_A^{\text{bare}} = 1.27$ of the axial-vector coupling. Using values in [38], we expect the quenching of this axial-vector coupling strength to increase the neutrinoless double-beta decay half-life of ^{48}Ca by a factor of 7_{-2}^{+3} .

C. Effect on geochronology

The branched decay of ^{40}K to ^{40}Ar is the basis of K/Ar dating and its variant the $^{40}\text{Ar}/^{39}\text{Ar}$ technique. The ubiquity of K and the $\approx 1.27 \text{ Ga}$ half-life of ^{40}K make these geochronometers some of the most useful and versatile isotopic methods available to date geological samples.

TABLE V. Effect of adding EC^0 to a commonly used set of decay constants [10] and the latest evaluation (Sec. 5.2 of [15]). λ_T is the total decay constant of ^{40}K , and λ_{Ar} is its partial decay constant to ^{40}Ar .

	$\lambda_T, 1\sigma$ (Ga^{-1})	$\lambda_{\text{Ar}}, 1\sigma$ (Ga^{-1})
Min <i>et al.</i> [10]	0.546 ± 0.005	0.0580 ± 0.0007
KDK with Min <i>et al.</i> [10]	0.547 ± 0.005	0.0585 ± 0.0007
Kossert <i>et al.</i> [15]	0.5468 ± 0.0019	0.0564 ± 0.00016
KDK with Kossert <i>et al.</i> [15]	0.5474 ± 0.0019	0.0570 ± 0.00021

The age equation for the $^{40}\text{K}/^{40}\text{Ar}$ isotope system is

$$t_{\text{K/Ar}} = \frac{1}{\lambda_T} \ln \left[\frac{N_{^{40}\text{Ar}} \lambda_T}{N_{^{40}\text{K}} \lambda_{\text{Ar}}} + 1 \right],$$

where $t_{\text{K/Ar}}$ is the age of the sample, N_x is the number of atoms of a given isotope in the sample, λ_T is the total decay constant of ^{40}K , and λ_{Ar} is the decay constant of ^{40}K to ^{40}Ar , including EC^* , EC^0 , and β^+ branches. In the K/Ar technique, the ratio of ^{40}K to ^{40}Ar isotopes is measured, leading to an estimation of the age of the sample. Precision of this technique currently reaches 0.5% [39].

In the $^{40}\text{Ar}/^{39}\text{Ar}$ technique, neutron activation is used to transmute ^{39}K to ^{39}Ar . This allows mass-spectrometric measurements of the parent proxy (^{39}Ar) and the daughter (^{40}Ar) on the same sample. The efficiency of the activation is hard to estimate, therefore the activation is generally carried out in parallel on a reference sample of known age, providing the age of the sample relative to the reference. The age of the reference must be established independently of the $^{40}\text{Ar}/^{39}\text{Ar}$ technique. Commonly used references include Fish Canyon ($t_m = 28.201 \pm 0.023$ Ma for a 68% CL) [40]. The age of the sample is then

$$t_{^{40}\text{Ar}/^{39}\text{Ar}} = \frac{1}{\lambda_T} \ln \left[\frac{N_{^{40}\text{Ar}}}{N_{^{39}\text{Ar}}} J + 1 \right],$$

where the irradiation parameter (J) is given by

$$J = \frac{e^{\lambda_T t_m} - 1}{M_{^{40}\text{Ar}}/M_{^{39}\text{Ar}}},$$

where $M_{^{40}\text{Ar}}/M_{^{39}\text{Ar}}$ is the measured isotopic ratio in the monitor. Unlike K/Ar, the $^{40}\text{Ar}/^{39}\text{Ar}$ technique no longer depends on the argon branching ratio. This technique currently reaches precisions of 0.1% [41].

Certain sets of decay constants widely used in the geological community question the existence of, or ignore, the EC^0 decay branch [10,11]. Using data in Table V, we illustrate how adding the EC^0 branch to various sets of decay constants [10,15] affects K/Ar dates throughout geologic time in Fig. 4(a) of [14]. In Fig. 4(b) of [14] we display the impact of this reevaluation on the commonly-used Fish Canyon sanidine standard, when the K/Ar age is recalculated from the $^{40}\text{K}/^{40}\text{Ar}$ ratio [42] and various decay constants [10,15,43]. Using these updated Fish Canyon ages with the same set of decay constants, we recalculate the $^{40}\text{Ar}/^{39}\text{Ar}$ ages of the Acapulco meteorite [44] in Fig. 4(c) of [14]. Also shown is the Pb/Pb age for phosphates from Acapulco [45], updated




FIG. 7. Effect of KDK on the thermal history of the Acapulco meteorite. $^{40}\text{Ar}/^{39}\text{Ar}$ ages of the Acapulco meteorite [44] calculated for various sets of decay constants [10,15,43] (Fig. 4(c) in [14]) are plotted along with the corresponding estimated closure temperature of 300 ± 25 °C (temperatures shifted for visibility). Age-temperature data obtained using Pb/Pb (updated from [45]) and Sm/Nd [48] also shown. For reference, the Pb/Pb age of the oldest dated solid in the solar system, a calcium-aluminum inclusion from the CV chondrite Efremovka, is 4567.35 ± 4.65 Ma [49]. All uncertainties correspond to a 68% confidence level, and include decay constant uncertainties.

to include uncertainties in the uranium isotopic composition [46], and the uranium decay constants [47] (4.555 ± 0.005 Ga).

Figure 7 shows the thermal history of the Acapulco meteorite using the various $^{40}\text{Ar}/^{39}\text{Ar}$ ages, along with those from Pb/Pb and Sm/Nd [48], with closure temperatures, and updated uncertainties for Sm/Nd, from [44]. Pb/Pb and Sm/Nd ages are statistically indistinguishable from $^{40}\text{Ar}/^{39}\text{Ar}$ ages calculated using recent decay constants [10,15] updated by our measurement of EC^0 (χ^2 fits to a common age respectively yield $p = 0.37$ for updated Kossert [15] and $p = 0.51$ for updated Min [10]), consistent with rapid cooling. The KDK measurement itself tends to decrease $^{40}\text{Ar}/^{39}\text{Ar}$ ages and therefore reduce apparent cooling rates. The systematic nature of this change may affect studies of past heat flow in Earth's crust.

IV. CONCLUSION

As detailed in this document and elsewhere [14], the KDK Collaboration has successfully measured the branching ratio of the elusive electron-capture decay of ^{40}K to the ground state of ^{40}Ar . The measured branching ratio is in agreement with the theoretical value calculated in this work. When compared to the traditionally used branching ratio, which is a factor of 2 larger, a factor of 5 increase in precision has been achieved. The improved precision will allow rare-event searches to better understand their low-energy backgrounds. Additionally, this measurement represents the first experimental verification of a third-forbidden unique transition

informing nuclear structure theory. Finally, our measurement affects K-decay based geochronological estimations by up to a percent.

ACKNOWLEDGMENTS

Xavier Mougeot of LNHB drew our attention to his latest evaluation of the decay scheme of ^{40}K . Engineering support has been contributed by Miles Constable and Fabrice Rétière of TRIUMF, as well as by Koby Dering through the NSERC/Queen's MRS. Funding in Canada was provided by NSERC through SAPIN and SAP RTI grants, as well as by the Faculty of Arts and Science of Queen's University, and by the McDonald Institute. Work was performed at Oak Ridge National Laboratory, managed by UT-Battelle, LLC, for the U.S. Department of Energy under Contract No. DE-AC05-00OR22725. Thermal deposition was conducted at the Center for Nanophase Materials Sciences, which is a DOE Office of Science User Facility. This manuscript has been authored by UT-Battelle, LLC under Contract No. DE-AC05-00OR22725

with the U.S. Department of Energy. U.S. support has also been supplied by the Joint Institute for Nuclear Physics and Applications, and by NSF Grant EAR-2102788. This material is based upon work supported by the U.S. Department of Homeland Security under Grant No. 2014-DN-077-ARI088-01. J.C., L.E.M., and P.R.R. acknowledge support from NSF Grant 2102788.

The U.S. Government retains and the publisher, by accepting the article for publication, acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this manuscript, or allow others to do so, for United States Government purposes. The Department of Energy will provide public access to these results of federally sponsored research in accordance with the DOE Public Access Plan [50]. The views and conclusions contained in this document are those of the authors and should not be interpreted as necessarily representing the official policies, either expressed or implied, of the U.S. Department of Homeland Security. Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

-
- [1] M. Schumann, Direct detection of WIMP dark matter: Concepts and status, *J. Phys. G: Nucl. Part. Phys.* **46**, 103003 (2019).
- [2] J. Pradler, B. Singh, and I. Yavin, On an unverified nuclear decay and its role in the DAMA experiment, *Phys. Lett. B* **720**, 399 (2013).
- [3] M. Antonello *et al.* (SABRE Collaboration), Monte Carlo simulation of the SABRE PoP background, *Astropart. Phys.* **106**, 1 (2019).
- [4] P. Adhikari *et al.* (COSINE-100 Collaboration), Background model for the NaI (TI) crystals in COSINE-100, *Eur. Phys. J. C* **78**, 490 (2018).
- [5] G. Angloher *et al.* (COSINUS Collaboration), Simulation-based design study for the passive shielding of the COSINUS dark matter experiment, *Eur. Phys. J. C* **82**, 248 (2022).
- [6] J. Amaré *et al.* (ANAIS-112 Collaboration), Annual modulation results from three-year exposure of ANAIS-112, *Phys. Rev. D* **103**, 102005 (2021).
- [7] H. Ejiri, J. Suhonen, and K. Zuber, Neutrino–nuclear responses for astro-neutrinos, single beta decays and double beta decays, *Phys. Rep.* **797**, 1 (2019).
- [8] F. Begemann, K. R. Ludwig, G. W. Lugmair, K. Min, L. E. Nyquist, P. J. Patchett, P. R. Renne, C.-Y. Shih, I. M. Villa, and R. J. Walker, Call for an improved set of decay constants for geochronological use, *Geochim. Cosmochim. Acta* **65**, 111 (2001).
- [9] J. Carter, R. B. Ickert, D. F. Mark, M. M. Tremblay, A. J. Cresswell, and D. C. W. Sanderson, Production of ^{40}Ar by an overlooked mode of ^{40}K decay with implications for K-Ar geochronology, *Geochronology* **2**, 355 (2020).
- [10] K. Min, R. Mundil, P. R. Renne, and K. R. Ludwig, A test for systematic errors in $^{40}\text{Ar}/^{39}\text{Ar}$ geochronology through comparison with U/Pb analysis of a 1.1-Ga rhyolite, *Geochim. Cosmochim. Acta* **64**, 73 (2000).
- [11] P. R. Renne, R. Mundil, G. Balco, K. Min, and K. R. Ludwig, Joint determination of ^{40}K decay constants and $^{40}\text{Ar}^*/^{40}\text{K}$ for the Fish Canyon sanidine standard, and improved accuracy for $^{40}\text{Ar}/^{39}\text{Ar}$ geochronology, *Geochim. Cosmochim. Acta* **74**, 5349 (2010).
- [12] X. Mougeot, Improved calculations of electron capture transitions for decay data and radionuclide metrology, *Appl. Radiat. Isot.* **134**, 225 (2018).
- [13] X. Mougeot, Towards high-precision calculation of electron capture decays, *Appl. Radiat. Isot.* **154**, 108884 (2019).
- [14] M. Stukel *et al.*, companion paper, Rare ^{40}K Decay with Implications for Fundamental Physics and Geochronology, *Phys. Rev. Lett.* **131**, 052503 (2023).
- [15] K. Kossert, Y. Amelin, D. Arnold, R. Merle, X. Mougeot, M. Schmiedel, and D. Zapata-García, Activity standardization of two enriched ^{40}K solutions for the determination of decay scheme parameters and the half-life, *Appl. Radiat. Isot.* **188**, 110362 (2022).
- [16] J. Chen, Nuclear data sheets for $A = 40$, *Nucl. Data Sheets* **140**, 1 (2017).
- [17] M. Wang, W. J. Huang, F. G. Kondev, G. Audi, and S. Naimi, The AME 2020 atomic mass evaluation (II). Tables, graphs and references, *Chin. Phys. C* **45**, 030003 (2021).
- [18] D. W. Engelkemeir, K. F. Flynn, and L. E. Glendenin, Positron emission in the decay of K^{40} , *Phys. Rev.* **126**, 1818 (1962).
- [19] R. Bernabei *et al.* (DAMA/LIBRA Collaboration), First model independent results from DAMA/LIBRA–Phase2, *Universe* **4**, 116 (2018).
- [20] P. C. F. Di Stefano (KDK Collaboration), The KDK (potassium decay) experiment, *J. Phys.: Conf. Ser.* **1342**, 012062 (2020).
- [21] M. Stukel *et al.* (KDK Collaboration), A novel experimental system for the KDK measurement of the ^{40}K decay scheme relevant for rare event searches, *Nucl. Instrum. Methods Phys. Res., Sect. A* **1012**, 165593 (2021).
- [22] M. Karny, K. P. Rykaczewski, A. Fijałkowska, B. C. Rasco, M. Wolińska-Cichocka, R. K. Grzywacz, K. C. Goetz, D. Miller, and E. F. Zganjar, Modular total absorption spectrometer, *Nucl. Instrum. Methods Phys. Res. A* **836**, 83 (2016).

- [23] B. Hoeling, D. Reed, and P. B. Siegel, Going bananas in the radiation laboratory, *Am. J. Phys.* **67**, 440 (1999).
- [24] X. Mougeot, BetaShape: A new code for improved analytical calculations of beta spectra, *EPJ Web Conf.* **146**, 12015 (2017).
- [25] S. Agostinelli *et al.* (GEANT4 Collaboration), GEANT4—a simulation toolkit, *Nucl. Instrum. Methods A* **506**, 250 (2003).
- [26] S. Baker and R. D. Cousins, Clarification of the use of chi-square and likelihood functions in fits to histograms, *Nucl. Instrum. Methods Phys. Res.* **221**, 437 (1984).
- [27] M. M. Bé *et al.*, *Table of Radionuclides* (BIPM, Sèvres, France, 2010), Vol. 5, A = 22–244, <https://hal-cea.archives-ouvertes.fr/cea-02476352v1>.
- [28] E. Schönfeld and H. Janßen, Evaluation of atomic shell data, *Nucl. Instrum. Methods Phys. Res., Sect. A* **369**, 527 (1996).
- [29] G. K. Wertheim, M. A. Butler, K. W. West, and D. N. E. Buchanan, Determination of the Gaussian and Lorentzian content of experimental line shapes, *Rev. Sci. Instrum.* **45**, 1369 (1974).
- [30] H. Behrens and W. Bühring, *Electron Radial Wave Functions and Nuclear Beta-Decay* (Clarendon, Oxford, 1982).
- [31] M. Haaranen, J. Kotila, and J. Suhonen, Spectrum-shape method and the next-to-leading-order terms of the β -decay shape factor, *Phys. Rev. C* **95**, 024327 (2017).
- [32] B. A. Brown and W. D. M. Rae, The shell-model code NuShellX@MSU, *Nucl. Data Sheets* **120**, 115 (2014).
- [33] K. Kaneko, Y. Sun, T. Mizusaki, and M. Hasegawa, Shell-model study for neutron-rich *sd*-shell nuclei, *Phys. Rev. C* **83**, 014320 (2011).
- [34] N. B. Gove and M. J. Martin, log-*f* tables for beta decay, *At. Data Nucl. Data Tables* **10**, 205 (1971).
- [35] LOGFT (accessed August 28th, 2020), with original code from Ref. [34]. Available at <https://www.nndc.bnl.gov/logft/>.
- [36] J. T. Suhonen, Value of the axial-vector coupling strength in β and $\beta\beta$ decays: A review, *Front. Physics* **5**, 55 (2017).
- [37] H. Ejiri, N. Soukouti, and J. Suhonen, Spin-dipole nuclear matrix elements for double beta decays and astro-neutrinos, *Phys. Lett. B* **729**, 27 (2014).
- [38] M. Kortelainen and J. Suhonen, Nuclear muon capture as a powerful probe of double-beta decays in light nuclei, *J. Phys. G Nucl. Phys.* **30**, 2003 (2004).
- [39] I. McDougall and P. Wellman, Calibration of GA1550 biotite standard for K/Ar and $^{40}\text{Ar}/^{39}\text{Ar}$ dating, *Chem. Geol.* **280**, 19 (2011).
- [40] K. F. Kuiper, A. Deino, F. J. Hilgen, W. Krijgsman, P. R. Renne, and J. R. Wijbrans, Synchronizing rock clocks of earth history, *Science* **320**, 500 (2008).
- [41] E. M. Niespolo, D. Rutte, A. L. Deino, and P. R. Renne, Intercalibration and age of the Alder Creek sanidine $^{40}\text{Ar}/^{39}\text{Ar}$ standard, *Quat. Geochronol.* **39**, 205 (2017).
- [42] F. Jourdan and P. R. Renne, Age calibration of the Fish Canyon sanidine $^{40}\text{Ar}/^{39}\text{Ar}$ dating standard using primary K–Ar standards, *Geochim. Cosmochim. Acta.* **71**, 387 (2007).
- [43] R. H. Steiger and E. Jäger, Subcommittee on geochronology: Convention on the use of decay constants in geo- and cosmochronology, *Earth Planet. Sci. Lett.* **36**, 359 (1977).
- [44] P. R. Renne, $^{40}\text{Ar}/^{39}\text{Ar}$ age of plagioclase from Acapulco meteorite and the problem of systematic errors in cosmochronology, *Earth Planet. Sci. Lett.* **175**, 13 (2000).
- [45] C. Göpel and G. Manhès, The thermal history of the Acapulco meteorite and its parent body deduced from U/Pb systematics in mineral separates and bulk rock fragments, *C. R. Geosci.* **342**, 53 (2010).
- [46] A. Goldmann, G. Brennecka, J. Noordmann, S. Weyer, and M. Wadhwa, The uranium isotopic composition of the Earth and the Solar System, *Geochim. Cosmochim. Acta* **148**, 145 (2015).
- [47] A. H. Jaffey, K. F. Flynn, L. E. Glendenin, W. C. t. Bentley, and A. M. Essling, Precision measurement of half-lives and specific activities of ^{235}U and ^{238}U , *Phys. Rev. C* **4**, 1889 (1971).
- [48] A. Prinzhofer, D. A. Papanastassiou, and G. J. Wasserburg, Samarium-neodymium evolution of meteorites, *Geochim. Cosmochim. Acta.* **56**, 797 (1992).
- [49] J. N. Connelly, M. Bizzarro, A. N. Krot, Å. Nordlund, D. Wielandt, and M. A. Ivanova, The absolute chronology and thermal processing of solids in the solar protoplanetary disk, *Science* **338**, 651 (2012).
- [50] <http://energy.gov/downloads/doe-public-access-plan>.