

JYX



This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): García-Bravo, Miguel; Rajala, Tapio; Takanen, Jyrki

Title: Two-Sided Boundary Points of Sobolev Extension Domains on Euclidean Spaces

Year: 2024

Version: Published version

Copyright: © The Author(s) 2023

Rights: CC BY 4.0

Rights url: <https://creativecommons.org/licenses/by/4.0/>

Please cite the original version:

García-Bravo, M., Rajala, T., & Takanen, J. (2024). Two-Sided Boundary Points of Sobolev Extension Domains on Euclidean Spaces. *Potential Analysis*, 60(3), 1249-1270.
<https://doi.org/10.1007/s11118-023-10086-5>



Two-Sided Boundary Points of Sobolev Extension Domains on Euclidean Spaces

Miguel García-Bravo¹ · Tapio Rajala¹ · Jyrki Takanen¹

Received: 3 September 2022 / Accepted: 11 June 2023
© The Author(s) 2023

Abstract

We prove an estimate on the Hausdorff dimension of the set of two-sided boundary points of general Sobolev extension domains on Euclidean spaces. We also present examples showing lower bounds on possible dimension estimates of this type.

Keywords Sobolev spaces · Extension domains · Two-sided points · Hausdorff dimension

Mathematics subject classification (2010) Primary 30L99. Secondary 46E35 · 26B30

1 Introduction

We continue the investigation of the geometric properties of Sobolev extension domains. In this paper, the space of Sobolev functions we use on a domain $\Omega \subset \mathbb{R}^n$ is the homogeneous Sobolev space $L^{1,p}(\Omega)$, which is the space of locally integrable functions whose weak derivatives belong to $L^p(\Omega)$. We endow this space with the homogeneous seminorm

$$\|f\|_{L^{1,p}(\Omega)} = \|\nabla f\|_{L^p(\Omega)} = \left(\int_{\Omega} |\nabla f(x)|^p dx \right)^{1/p}.$$

The reason for working with the homogeneous Sobolev space is simply to make our dimension estimates scaling invariant. We will comment on the non-homogeneous spaces after stating our main result.

We say that $E: L^{1,p}(\Omega) \rightarrow L^{1,p}(\mathbb{R}^n)$ is an extension operator if $Eu(x) = u(x)$ for all $u \in L^{1,p}(\Omega)$ and $x \in \Omega$, and if there exists a constant $C \geq 1$ so that for every $u \in L^{1,p}(\Omega)$ we have $\|Eu\|_{L^{1,p}(\mathbb{R}^n)} \leq C\|u\|_{L^{1,p}(\Omega)}$. We name the infimum over such possible constants C by $\|E\|$, and call it the norm of the extension operator. We say that a domain $\Omega \subset \mathbb{R}^n$

✉ Tapio Rajala
tapio.m.rajala@jyu.fi

Miguel García-Bravo
miguel05@ucm.es

Jyrki Takanen
jyrki.j.takanen@jyu.fi

¹ Department of Mathematics and Statistics, University of Jyväskylä, FI-40014 P.O. Box 35 (MaD), Jyväskylä, Finland

is an $L^{1,p}$ -extension domain if such an operator exists. The same definition applies for the non-homogeneous spaces $W^{1,p}(\Omega)$.

Throughout this manuscript each time we refer to a Sobolev extension domain we mean it with respect to the homogeneous norm, unless otherwise stated.

Already from the work of Calderón and Stein [16] we know that Lipschitz domains are $W^{1,p}$ -extension domains. However, much more complicated domains admit an extension operator. For instance, the Koch snowflake domains are extension domains and in some sense serve as sharp examples of extension domains in terms of the Hausdorff dimension of the boundary, see [13]. In [13], the question of the possible size of the boundary for simply connected planar Sobolev extension domains was studied. In particular, for these domains there is an upper bound on the Hausdorff dimension of the boundary in terms of the norm of the extension operator (although in [13] the bound was expressed in terms of a constant in a characterizing curve-condition property provided in [9]). Note that for the boundary of a general extension domain we cannot have a dimension estimate: take $\Omega = [0, 1]^n \setminus C^n$ with $C \subset \mathbb{R}$ a Cantor set of $\dim_{\mathcal{H}}(C) = 1$ but Lebesgue measure zero. Then Ω is a Sobolev $L^{1,p}$ -extension domain, but $\dim_{\mathcal{H}}(\partial\Omega) = n$.

With a bound on the dimension of the boundary, one might wonder what other geometric limitations does the existence of an extension operator imply. Let us approach this with a basic example of a domain that is not an $L^{1,p}$ -extension domain for any p , the slit disc: $\Omega := \mathbb{D} \setminus [0, 1] \times \{0\} \subset \mathbb{R}^2$. A continuous Sobolev function in $L^{1,p}(\Omega)$ which is one above $[1/2, 1] \times \{0\}$ and zero below it serves as an example of a function that cannot be extended to a global Sobolev function since no extension would be absolutely continuous on almost every vertical line segment. Notice however, that the slit disc is an example of a BV -extension domain because its complement is quasicontinuous (see [8]).

The slit disc example can be modified to a more delicate one by replacing the removed line segment $[0, 1] \times \{0\}$ by a larger set where the two-sided points are at a Cantor set on the previously removed line segment, see Figure 1 in Section 4. This will give a domain where the extendability of Sobolev functions depends on the exponent p . Such constructions will also play a role in this paper, see Section 4.2 (also for the precise definitions of these domains). By removing small neighbourhoods of the two-sided points from the domain, one can actually make the example into a Jordan domain and still retain the property of being a Sobolev extension domain for some p 's, but not for other, see [11] for a similar construction (and also the earlier works [14, 15]).

The slit disc and its variations have boundary points that can be approached from two different sides in the domain. In this paper we study the question of how large this set of two-sided points can be for a Sobolev extension domain. This question was already investigated in [17] by the third named author in the case of planar simply connected domains. Before continuing, let us give the definition of two-sidedness that we will use in this paper. In the case of simply connected planar domains, the definition can also be reformulated in various ways using conformal mappings, see [17].

Definition 1.1 (Two-sided points of the boundary of a domain) Let $\Omega \subset \mathbb{R}^n$ be a domain. A point $x \in \partial\Omega$ is called *two-sided*, if there exists $R > 0$ such that for all $r \in (0, R)$ there exist disjoint connected components Ω_r^1 and Ω_r^2 of $\Omega \cap B(x, r)$ that are nested: $\Omega_s^i \subset \Omega_r^i$ for $0 < s < r < R$ and $i \in \{1, 2\}$.

We denote the set of two-sided points of $\partial\Omega$ by \mathcal{T}_Ω , or simply by \mathcal{T} , if there is no possibility for confusion. Notice that the set \mathcal{T} need not be closed.

For $p \geq n$, we know that $L^{1,p}$ extension domains are quasiconvex (see [7, Theorem 3.1]). Therefore, for an $L^{1,p}$ -extension domain with $p \geq n$, we have $\mathcal{T} = \emptyset$. The interesting case is thus $1 \leq p < n$. For this range we prove the following estimate on the size of \mathcal{T} :

Theorem 1.2 *Let $n \geq 2$ and $p \in [1, n)$ and let $\Omega \subset \mathbb{R}^n$ be a Sobolev $L^{1,p}$ -extension domain. Then*

- (1) *If $p = 1$, then $\mathcal{H}^{n-1}(\mathcal{T}_\Omega) = 0$.*
- (2) *If $p > 1$ there exists a constant $C(n, p) > 0$ so that*

$$\dim_{\mathcal{H}}(\mathcal{T}_\Omega) \leq n - p - \frac{C(n, p)}{\|E\|^n \log(\|E\|)},$$

where $\|E\|$ is the operator norm of the homogeneous Sobolev extension operator.

Here we use the convention that $\mathcal{T}_\Omega = \emptyset$ whenever the bound on the right-hand side of the estimate is strictly less than 0.¹

Let us now comment on the non-homogeneous Sobolev spaces. For bounded domains Ω it is known that Sobolev $L^{1,p}$ -extension domains are the same as Sobolev $W^{1,p}$ -extension domains (see [7]), so even though our main result is stated for homogeneous Sobolev extension domains, it can be applied to $W^{1,p}$ -extension domains in the case that Ω is bounded. Let us note that there exist unbounded Sobolev $W^{1,p}$ -extension domains which are not $L^{1,p}$ -extension domains (see [7, Example 6.7]). However, one might expect that our result still applies for this unbounded case because having a dimension bound relies on local properties. Indeed, our method of proof will show that we can handle also with unbounded $W^{1,p}$ -extension domains because the measure density condition (see Proposition 2.2) is still true for every $r \in (0, 1)$ and the proof of Theorem 1.2 studies locally the set of two-sided points to estimate its dimension. We prefer to state our main theorem only for $L^{1,p}$ -extension domains because of their homogeneous norm. If we stated it for $W^{1,p}$ -extension domains, then a scaling of the domain Ω would perturb the norm of the operator E , and hence our estimate in the dimension of the two-sided points. Obviously, a scaling of a set will never change its dimension.

We will also give a size estimate on the two sided-points of BV -extension domains.

Theorem 1.3 *Let $n \geq 2$ and let $\Omega \subset \mathbb{R}^n$ be a BV -extension domain. Then \mathcal{T}_Ω has σ -finite $(n - 1)$ -dimensional Hausdorff measure.*

Observe that taking Ω to be a slit disc shows the sharpness of this result.

We will present the proof of Theorem 1.2 in Section 2 and the proof of Theorem 1.3 in Section 3. After that, in Section 4, we show that Theorem 1.2 (1) is sharp: there exist even a planar simply connected $L^{1,1}$ -extension domain with $\dim_{\mathcal{H}}(\mathcal{T}) = 1$. We also give a class of domains Ω_λ for each $n \geq 2$ with the sets of two-sided points $\mathcal{T}_\lambda = C_\lambda$ being Cantor sets, so that for every $p \in (1, n)$ there exists a constant $C(n, p)$ for which, with the explicit extension operators $E_\lambda : L^{1,p}(\Omega_\lambda) \rightarrow L^{1,p}(\mathbb{R}^n)$ we construct, we have $\|E_\lambda\| \rightarrow \infty$ as $\dim_{\mathcal{H}}(C_\lambda) \rightarrow n - p$ and the estimate

$$\dim_{\mathcal{H}}(C_\lambda) \geq n - p - \frac{C(n, p)}{\|E_\lambda\|} \tag{1.1}$$

is satisfied.

¹ To make this formally correct we adopt the notational convention $\dim_{\mathcal{H}}(\emptyset) = -\infty$.

This set of examples together with Theorem 1.2 shows that the possible optimal asymptotic behaviour for the dimension bound of the two-sided points in terms of the norm of the extension operator is between $n - p - C/\|E\|$ and $n - p - C/(\|E\|^n \log(\|E\|))$. We note that in [17] the exponents for the dimension bound and examples agreed, thus providing a possibly sharper estimate. However, as the study in [17] was done in terms of a constant in a characterizing curve condition, and since the dependence between this constant and the norm of the extension operator has not been clarified, the estimate in [17] does not yet translate to a sharp dimension estimate in terms of the norm of the extension operator in the planar simply connected case.

2 Dimension Estimate for the Set of Two-Sided Points

In this section we will prove Theorem 1.2. Before doing so, we go through some notation and lemmata.

We often denote by $C(\cdot)$ a computable constant depending only on the parameters listed in the parenthesis. The constant may differ between appearances, even within a chain of (in)equalities. By $a \lesssim b$ we mean that $a \leq Cb$ for some constant $C \geq 1$, that could depend on the dimension n . Similarly for $a \gtrsim b$. Then $a \sim b$ means that both $a \lesssim b$ and $a \gtrsim b$ hold. We denote by \mathbf{m}_n the n -dimensional Lebesgue measure on \mathbb{R}^n . We will also denote by $Q(x, s)$ the cube of center x and side length $s > 0$ and for a given cube $Q = Q(x, s)$ and some positive $K > 0$ we write $KQ = Q(x, Ks)$. We denote the side length of a cube by $\ell(Q)$.

We will use the following basic lemma, similar to [17, Lemma 3.2].

Lemma 2.1 *Let $F \subset \mathbb{R}^n$, $0 < \lambda < 1$, $s \geq 0$ and $i_0 \in \mathbb{N}$. For every $i \geq i_0$ let $\{x_k^i\}_{k \in I_i}$ be a maximal λ^i -separated net in F . Assume that for each $i \geq i_0$ and $k \in I_i$ there exists $j > i$ such that*

$$N_j < \lambda^{-(j-i)s},$$

where $N_j = \#\{l \in I_j : B(x_l^j, \lambda^j) \cap B(x_k^i, \lambda^i) \neq \emptyset\}$. Then $\dim_{\mathcal{H}}(F) \leq s$.

A measure density condition for Sobolev extension domains was proven in [4]. We will need to make the dependence of the parameters more explicit, so we modify slightly the proofs of [4, Lemma 11] and [4, Theorem 1] to obtain the following version of their measure density condition.

Proposition 2.2 *[Measure density condition] Let $1 \leq p < \infty$ and let $\Omega \subset \mathbb{R}^n$ be a Sobolev $L^{1,p}$ -extension domain with an extension operator E .*

(1) *If $1 \leq p < n$ then for all $x \in \overline{\Omega}$ and $r \in \left(0, \min \left\{1, \left(\frac{\mathbf{m}_n(\Omega)}{2\mathbf{m}_n(B(0,1))}\right)^{1/n}\right\}\right)$, denoting by Ω' a connected component of $\Omega \cap B(x, r)$ with $x \in \overline{\Omega'}$, we have*

$$\mathbf{m}_n(\Omega') \geq C(n, p)\|E\|^{-n}r^n.$$

(2) *If $p > n - 1$ then for all $x \in \overline{\Omega}$ and $r \in (0, \text{diam}(\Omega))$, denoting by Ω' a connected component of $\Omega \cap B(x, r)$ with $x \in \overline{\Omega'}$, we have*

$$\mathbf{m}_n(\Omega') \geq C(n, p)\|E\|^{-p}r^n.$$

Proof The case (2) follows by Theorem 2.2 and the proof of Theorem 4.1 from Koskela’s dissertation [7], where he uses the concept of variational p -capacity. Notice that in the proof of Theorem 4.1 the support of the test function u is contained in Ω' .

We look now at the case $1 \leq p < n$.

Let us denote $r_0 = r$. By induction, we define for every $i \in \mathbb{N}$ the radius $r_i \in (0, r_{i-1})$ by the equality

$$\mathbf{m}_n(\Omega' \cap B(x, r_i)) = \frac{1}{2} \mathbf{m}_n(\Omega' \cap B(x, r_{i-1})) = 2^{-i} \mathbf{m}_n(\Omega').$$

Since $x \in \overline{\Omega'}$, we have that $r_i \searrow 0$ as $i \rightarrow \infty$.

For each $i \in \mathbb{N}$, consider the function $f_i : \Omega \rightarrow \mathbb{R}$

$$f_i(y) = \begin{cases} 1, & \text{for } y \in B(x, r_i) \cap \Omega', \\ \frac{r_{i-1}-|x-y|}{r_{i-1}-r_i}, & \text{for } y \in (B(x, r_{i-1}) \setminus B(x, r_i)) \cap \Omega', \\ 0, & \text{otherwise.} \end{cases}$$

For the homogeneous Sobolev-norm of f_i we can estimate

$$\begin{aligned} \|f_i\|_{L^{1,p}(\Omega)}^p &= \int_{\Omega} |\nabla f_i|^p \leq |r_i - r_{i-1}|^{-p} \mathbf{m}_n((B(x, r_{i-1}) \setminus B(x, r_i)) \cap \Omega') \\ &= |r_i - r_{i-1}|^{-p} 2^{-i} \mathbf{m}_n(\Omega'). \end{aligned} \tag{2.1}$$

Call $p^* = \frac{np}{n-p}$. For any $Ef_i \in L^{1,p}(\mathbb{R}^n)$, by the Sobolev-Poincaré-inequality (we will also prove a variant of this later, see (2.3)), we know the existence of a constant $c_i \in \mathbb{R}$ (that can be assumed to be between 0 and 1) so that

$$\|Ef_i - c_i\|_{L^{p^*}(\mathbb{R}^n)} \leq C(n, p) \|Ef_i\|_{L^{1,p}(\mathbb{R}^n)}.$$

Hence we have the following chain of inequalities

$$\|f_i - c_i\|_{L^{p^*}(\Omega)} \leq \|Ef_i - c_i\|_{L^{p^*}(\mathbb{R}^n)} \leq C(n, p) \|Ef_i\|_{L^{1,p}(\mathbb{R}^n)} \leq C(n, p) \|E\| \|f_i\|_{L^{1,p}(\Omega)}.$$

Recall that by our choice of $r = r_0$ we always have

$$\mathbf{m}_n(\Omega \setminus B(x, r_{i-1})) \geq \mathbf{m}_n(\Omega \setminus B(x, r_0)) \geq \mathbf{m}_n(\Omega) - \mathbf{m}_n(B(x, r_0)) \geq \frac{\mathbf{m}_n(\Omega)}{2}$$

and

$$\mathbf{m}_n(B(x, r_i) \cap \Omega') \leq \frac{\mathbf{m}_n(\Omega)}{2}$$

for every $i \geq 1$. Then

$$\begin{aligned} \int_{\Omega} |f_i(y) - c_i|^{p^*} dy &\geq \max \left\{ \int_{\{y: f_i(y)=0\}} |c_i|^{p^*} dy, \int_{\{y: f_i(y)=1\}} |1 - c_i|^{p^*} dy \right\} \\ &\geq \max \left\{ |c_i|^{p^*} \mathbf{m}_n(\Omega \setminus B(x, r_{i-1})), |1 - c_i|^{p^*} \mathbf{m}_n(B(x, r_i) \cap \Omega') \right\} \\ &\geq \mathbf{m}_n(B(x, r_i) \cap \Omega') \cdot \max \left\{ |c_i|^{p^*}, |1 - c_i|^{p^*} \right\} \\ &\geq \mathbf{m}_n(B(x, r_i) \cap \Omega') \cdot 2^{-p^*}, \end{aligned}$$

so we write, using (2.1),

$$\begin{aligned} 2^{-p^*-i} \mathbf{m}_n(\Omega') &= 2^{-p^*} \mathbf{m}_n(B(x, r_i) \cap \Omega') \leq \|f_i - c_i\|_{L^{p^*}(\Omega)}^{p^*} \leq C(n, p) \|E\|^{p^*} \|f_i\|_{L^{1,p}(\Omega)}^{p^*} \\ &= C(n, p) \|E\|^{p^*} \left(\int_{\Omega} |\nabla f_i(y)|^p dy \right)^{p^*/p} \\ &\leq C(n, p) \|E\|^{p^*} \left(|r_i - r_{i-1}|^{-p} 2^{-i} \mathbf{m}_n(\Omega') \right)^{p^*/p} \\ &\leq C(n, p) \|E\|^{p^*} 2^{-ip^*/p} \mathbf{m}_n(\Omega')^{p^*/p} |r_{i-1} - r_i|^{-p^*}. \end{aligned}$$

Consequently,

$$\begin{aligned} r_{i-1} - r_i &\leq C(n, p) \|E\| 2^{i(1/p^*-1/p)} \mathbf{m}_n(\Omega')^{1/p-1/p^*} \\ &= C(n, p) \|E\| 2^{-i/n} \mathbf{m}_n(\Omega')^{1/n}. \end{aligned}$$

By summing up all these quantities we conclude that

$$r = r_0 = \sum_{i=1}^{\infty} (r_{i-1} - r_i) \leq C(n, p) \|E\| \sum_{i=1}^{\infty} 2^{-i/n} \mathbf{m}_n(\Omega')^{1/n} = \frac{C(n, p) \|E\|}{2^{1/n} - 1} \mathbf{m}_n(\Omega')^{1/n}.$$

This gives the claimed inequality.

Observe that the measure density condition only holds for $1 \leq p < \infty$. For $W^{1,\infty}$ -extension domains this is not true. Take for instance $C \subset [0, 1]$ a fat Cantor set with $\mathbf{m}_1(C) > 0$. Then almost every point of C is of density 1 on C , so $[0, 1] \setminus C$, whose closure is the whole interval $[0, 1]$, cannot satisfy any measure density condition. Then take $\Omega = \mathbb{R}^n \setminus C^n$ which will be quasiconvex by [5, Theorem A], and consequently a $W^{1,\infty}$ -extension domain by [4, Theorem 7], but does not satisfy any measure density condition either.

In the proof of Theorem 1.2 we will use the following consequence of a Sobolev-Poincaré type inequality (2.3). The proof of the lemma follows the proof for the classical Sobolev-Poincaré inequality that can be found in many text books. However, for our application of the lemma we need to include a set F that is removed when integrating the gradient of the Sobolev function. This fact forces us to be more cautious. For the convenience of the reader, we provide here the proof with the needed modifications.

Lemma 2.3 *Let $1 \leq p < n$, $Q \subset \mathbb{R}^n$ be a cube, $\delta \in (0, 1)$ and $F \subset Q$ such that for any $i \in \{1, \dots, n\}$ we have*

$$\mathbf{m}_{n-1}(P_i(F)) \leq \frac{\delta}{2n \cdot 2^n} \mathbf{m}_{n-1}(P_i(Q))$$

with P_i the projection $P_i: (x_1, \dots, x_n) \mapsto (x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n)$. Then for any $f \in W^{1,p}(Q)$ so that $0 \leq f \leq 1$ and

$$\min \left(\mathbf{m}_n(\{y \in \frac{1}{2}Q : f(y) = 0\}), \mathbf{m}_n(\{y \in \frac{1}{2}Q : f(y) = 1\}) \right) > \delta \frac{\ell(Q)^n}{2^n},$$

we have

$$\int_{Q \setminus F} |\nabla f(y)|^p dy \geq C(n, p) \delta^{\frac{n-p}{n}} \ell(Q)^{n-p}. \tag{2.2}$$

Remark. Observe that for the conclusion of Lemma 2.3 it is not enough to only require $\mathbf{m}_n(F)$ to be small. Consider for instance the cube minus a very thin central band which separates the cube in two connected components.

Proof We will show that the following version of Sobolev-Poincaré inequality holds for our function f :

$$\left(\int_A |f(y) - f_A|^{\frac{pn}{n-p}} dy \right)^{\frac{n-p}{pn}} \leq C(n, p) \left(\int_{Q \setminus F} |\nabla f(y)|^p dy \right)^{1/p}, \tag{2.3}$$

where $A = \{x \in \frac{1}{2}Q : P_i(x) \notin P_i(F) \text{ for every } i\}$ and

$$f_A = \frac{1}{\mathbf{m}_n(A)} \int_A f(y) dy.$$

Let us first observe that this implies

$$\begin{aligned} \int_{Q \setminus F} |\nabla f(y)|^p dy &\gtrsim \left(\int_A |f(y) - f_A|^{\frac{pn}{n-p}} dy \right)^{\frac{n-p}{n}} \\ &\gtrsim \max \left(\mathbf{m}_n(\{y \in A : f(y)=1\})^{\frac{n-p}{n}} |1 - f_A|^p, \mathbf{m}_n(\{y \in A : f(y)=0\})^{\frac{n-p}{n}} |f_A|^p \right) \\ &\gtrsim \delta^{\frac{n-p}{n}} \ell(Q)^{n-p} \max(|1 - f_A|^p, |f_A|^p) \\ &\gtrsim \delta^{\frac{n-p}{n}} \ell(Q)^{n-p}, \end{aligned}$$

thus giving the inequality (2.2). Above we used the simple observation that

$$\mathbf{m}_n\left(\frac{1}{2}Q \setminus A\right) \leq \frac{\delta}{4 \cdot 2^n} \mathbf{m}_n(Q). \tag{2.4}$$

To prove (2.3) we start by presenting the Sobolev embedding in the form

$$\left(\int_{A'} |g(y)|^{\frac{pn}{n-p}} dy \right)^{\frac{n-p}{pn}} \leq C(n, p, K) \left(\int_{Q \setminus F} |\nabla g(y)|^p dy \right)^{1/p}, \tag{2.5}$$

for all $g \in W_0^{1,p}(Q)$ with $|g| \leq 1$ and $\mathbf{m}_n(\{x \in A' : |g(x)| \geq 1/2\}) \geq K \delta \ell(Q)^n$ for some positive constant $K > 0$, and where $A' = \{x \in Q : P_i(x) \notin P_i(F) \text{ for every } i\}$. Following the proof of [2, Theorem 4.8] what we first get is

$$\left(\int_{A'} |g(y)|^{\frac{pn}{n-p}} dy \right)^{\frac{n-1}{n}} \leq C(n, p) \left(\int_{Q \setminus F} |g(y)|^{\frac{pn}{n-p}} dy \right)^{\frac{p-1}{p}} \left(\int_{Q \setminus F} |\nabla g(y)|^p dy \right)^{1/p}.$$

Note that by the properties of g and by definition of A'

$$\int_{Q \setminus A'} |g(y)|^{\frac{pn}{n-p}} dy \leq \mathbf{m}_n(Q \setminus A') < \frac{n\delta}{2n \cdot 2^n} \ell(Q)^n$$

and

$$\int_{A'} |g(y)|^{\frac{pn}{n-p}} dy \geq \left(\frac{1}{2}\right)^{\frac{pn}{n-p}} K \delta \ell(Q)^n.$$

Therefore,

$$\begin{aligned} \int_{Q \setminus F} |g(y)|^{\frac{pn}{n-p}} dy &\leq \int_Q |g(y)|^{\frac{pn}{n-p}} dy = \int_{A'} |g(y)|^{\frac{pn}{n-p}} dy + \int_{Q \setminus A'} |g(y)|^{\frac{pn}{n-p}} dy \\ &\leq (1 + C(n, p, K)) \int_{A'} |g(y)|^{\frac{pn}{n-p}} dy, \end{aligned} \tag{2.6}$$

and finally we can get (2.5).

Secondly, we apply the inequality (2.5) to the function $g(y) = (f(y) - f_{A'})\phi(y)$, where $\phi \in C_0^\infty(\mathbb{R}^n)$ is supported in Q , is equal to 1 on $\frac{1}{2}Q$ and $|\nabla\phi| \lesssim \frac{1}{\ell(Q)}$. We get

$$\begin{aligned} \left(\int_A |f(y) - f_A|^{\frac{pn}{n-p}} dy\right)^{\frac{n-p}{pn}} &\leq \left(\int_{A'} |(f(y) - f_A)\phi(y)|^{\frac{pn}{n-p}} dy\right)^{\frac{n-p}{pn}} \\ &\leq C(n, p) \left(\int_{Q \setminus F} |\nabla f(y)|^p dy\right)^{1/p} \\ &\quad + \frac{C(n, p)}{\ell(Q)} \left(\int_{Q \setminus F} |f(y) - f_A|^p dy\right)^{1/p}. \end{aligned} \tag{2.7}$$

To handle the last term above, we first prove that

$$\begin{aligned} \left(\int_{Q \setminus F} |f(y) - f_A|^p dy\right)^{1/p} &\leq C(n, p) \left(\int_{A'} |f(y) - f_A|^p dy\right)^{1/p} \\ &\leq C(n, p) \left(\left(\int_{A'} |f(y) - f_{A'}|^p dy\right)^{1/p} \right. \\ &\quad \left. + \left(\int_{A'} |f_{A'} - f_A|^p dy\right)^{1/p}\right) \\ &\leq C(n, p) \left(\int_{A'} |f(y) - f_{A'}|^p dy\right)^{1/p}. \end{aligned}$$

In the first inequality we are using a similar trick like in (2.6) (that $0 \leq f \leq 1$ and that $f = 1$ and $f = 0$ in large enough sets). In the last inequality we use Hölder inequality and the fact that by (2.4) we have

$$\mathbf{m}_n(A) \geq \mathbf{m}_n\left(\frac{1}{2}Q\right) - \mathbf{m}_n\left(\frac{1}{2}Q \setminus A\right) \geq \frac{\mathbf{m}_n(Q)}{2^n} - \frac{\delta \mathbf{m}_n(Q)}{4 \cdot 2^n} \geq 2^{-n-1} \mathbf{m}_n(Q) \geq 2^{-n-1} \mathbf{m}_n(A').$$

Finally, by modifying the standard proof for the Poincaré inequality (see [2, Section 4.5.2]) by first writing

$$|f(y) - f(x)| \leq \sum_{i=1}^n |f(z_i) - f(z_{i-1})|,$$

with $z_i = (y_1, \dots, y_i, x_{i+1}, \dots, x_n)$ so that z_i and z_{i-1} differ only in one coordinate, we are able to consider absolute continuity only along lines going in the coordinate directions. Thus, we obtain

$$\int_{A'} |f(y) - f_{A'}|^p dy \leq C(n, p) \ell(Q)^p \int_{Q \setminus F} |\nabla f(y)|^p dy.$$

Combining the above with (2.7) concludes the proof.

We are now ready to prove Theorem 1.2.

Proof (Proof of Theorem 1.2) Let us first make some initial reductions. By the definition of two-sided points we can write $\mathcal{T}_\Omega = \bigcup_{i \in \mathbb{N}} \mathcal{T}_i$, where

$$\begin{aligned} \mathcal{T}_i &= \{x \in \partial\Omega : \text{for every } r < 2^{-i}, \text{ there exist two different connected components } \Omega_r^1, \Omega_r^2 \\ &\quad \text{of } \Omega \cap B(x, r) \text{ that are nested, that is } \Omega_s^j \subset \Omega_r^j \text{ for } 0 < s < r, j = 1, 2\}. \end{aligned}$$

Observe that if $x \in \mathcal{T}_i$ and Ω_r^1, Ω_r^2 are the associated nested connected components of $\Omega \cap B(x, r)$ for each $r \in (0, 2^{-i})$, then $x \in \overline{\Omega_r^1} \cap \overline{\Omega_r^2}$ for all $r \in (0, 2^{-i})$.

It is clear that it is enough to estimate $\dim_{\mathcal{H}}(\mathcal{T}_i)$ for a fixed $i \in \mathbb{N}$. We now cover \mathcal{T}_i by countably many balls $B(z_k, 2^{-i}/6)$, where $z_k \in \mathcal{T}_i$. Then, for every $k \in \mathbb{N}$ we introduce the family of pairwise disjoint connected components of $B(z_k, 2^{-i}/2) \cap \Omega$, which we denote by $\{O_l^k\}_{l \in I}$. Let us check now that

$$\mathcal{T}_i \cap B(z_k, 2^{-i}/6) \subseteq \bigcup_{l \neq \tilde{l}} \partial O_l^k \cap \partial O_{\tilde{l}}^k. \tag{2.8}$$

Take $x \in \mathcal{T}_i \cap B(z_k, 2^{-i}/6)$. Since $x \in \mathcal{T}_i$ there exist two different connected components of $\Omega \cap B(x, 2^{-i})$, which we call U_1, U_2 , so that $x \in \partial U_1 \cap \partial U_2$. Therefore, using that

$$B(z_k, 2^{-i}/2) \cap \Omega \subset B(x, 2^{-i}) \cap \Omega,$$

the sets $U_1 \cap B(z_k, 2^{-i}/2)$ and $U_2 \cap B(z_k, 2^{-i}/2)$ will have connected components, which we call $O_l^k, O_{\tilde{l}}^k$, so that $x \in \partial O_l^k \cap \partial O_{\tilde{l}}^k$. We have then proved (2.8). Observe that we can write

$$\mathcal{T}_\Omega = \bigcup_{i,k} \bigcup_{l \neq \tilde{l}} \mathcal{T}_i \cap B(z_k, 2^{-i}/6) \cap (\partial O_l^k \cap \partial O_{\tilde{l}}^k).$$

Therefore, it is enough to just estimate the Hausdorff dimension of $\mathcal{T}_i \cap B(z_k, 2^{-i}/6) \cap (\partial O_l^k \cap \partial O_{\tilde{l}}^k)$ for fixed i, k, l, \tilde{l} with $l \neq \tilde{l}$. Each set of this type, that we call from now on G , has the following properties: there is some $x_0 \in \partial\Omega$ and some radius $r \in (0, 1)$ so that

$$G \subset \partial\Omega \cap B(x_0, r),$$

and there exist connected components $\Omega_1, \Omega_2 \subset \Omega \cap B(x_0, 3r)$ for which

$$G \subset \partial\Omega_1 \cap \partial\Omega_2.$$

We will now proceed to estimate the Hausdorff dimension of such a set G .

- (1) Let us first prove that $\mathcal{H}^{n-p}(G) = 0$ for all $1 \leq p < n$. In particular, this will handle the case $p = 1$ in the claim (1) of the theorem. We will use the well-known fact that for any given $h \in L^1_{\text{loc}}(\mathbb{R}^n)$ and $0 \leq s < n$ we have

$$\mathcal{H}^s \left(\left\{ x \in \mathbb{R}^n : \limsup_{\varepsilon \searrow 0} \frac{1}{\varepsilon^s} \int_{B(x,\varepsilon)} |h(y)| dy > 0 \right\} \right) = 0. \tag{2.9}$$

See for instance [2, Theorem 2.10] for a proof of this assertion.

We start by defining a function $u \in L^{1,p}(\Omega)$,

$$u(x) = \max \left(0, \min \left(1, 3 - r^{-1} \text{dist}(x, x_0) \right) \chi_{\Omega_1}(x) \right),$$

where χ_{Ω_1} denotes the indicator function of the set Ω_1 . Notice that u is locally Lipschitz in Ω by the fact that Ω_1 is a connected component of $\Omega \cap B(x_0, 3r)$. By Proposition 2.2, for every $x \in G$ and every $0 < \varepsilon \leq r$ sufficiently small

$$\min \left(\mathbf{m}_n(\Omega_1 \cap B(x, \varepsilon/2\sqrt{n})), \mathbf{m}_n(\Omega_2 \cap B(x, \varepsilon/2\sqrt{n})) \right) \geq C(n, p) \|E\|^{-n} \varepsilon^n.$$

Now, we apply Lemma 2.3 with the removed set $F = \emptyset$ and the cube $Q(x, 2\varepsilon/\sqrt{n})$ centered at x and with side length $2\varepsilon/\sqrt{n}$. Notice that $Q(x, 2\varepsilon/\sqrt{n})$ contains the ball

$B(x, \varepsilon/\sqrt{n})$ and is contained in the ball $B(x, \varepsilon)$, and so the Lemma 2.3 gives

$$\int_{B(x,\varepsilon)} |\nabla Eu(y)|^p dy \geq \int_{Q(x,2\varepsilon/\sqrt{n})} |\nabla Eu(y)|^p dy \geq C(n, p)\|E\|^{p-n}\varepsilon^{n-p}.$$

Therefore,

$$\limsup_{\varepsilon \searrow 0} \frac{1}{\varepsilon^{n-p}} \int_{B(x,\varepsilon)} |\nabla Eu(y)|^p dy \geq C(n, p)\|E\|^{p-n} > 0$$

for every $x \in G$, and using (2.9) we conclude $\mathcal{H}^{n-p}(G) = 0$.

We are done with the case $p = 1$ of Theorem 1.2. For the case $p > 1$ we will be able to be more precise in the estimation of the Hausdorff dimension in terms of the norm of the extension operator $E: L^{1,p}(\Omega) \rightarrow L^{1,p}(\mathbb{R}^n)$. For this we will follow a different approach.

- (2) Let us now focus on the case $p > 1$. First of all, call $C_1(n, p)$ and $C_2(n, p)$ the constants given by Proposition 2.2 and Lemma 2.3 respectively. Now we choose $0 < \lambda < 1$ so that,

$$\frac{\lambda^{p-1}}{1 - \lambda^{p-1}} = \frac{C_1(n, p)^{1+\frac{n-p}{n}} C_2(n, p)}{2^{2n+1} 3^n n^{(2n+1-p)/2} \mathbf{m}_n(B(0, 1))} \|E\|^{-2n}.$$

Then we choose $i_0 \in \mathbb{N}$ so that $0 < \lambda^{i_0} < r$ is small enough and such that Proposition 2.2 is satisfied.

We can do this because the term on the left hand side tends to zero as $\lambda \rightarrow 0$.

For every $i \in \mathbb{N}$, let $\{x_k^i\}_{k \in I_i}$ be a maximal λ^i -separated net of points in G . For every $i \in \mathbb{N}$ and $k \in I_i$ define

$$\mathcal{B}_j^{i,k} = \{B(x_l^{i+j}, \lambda^{i+j}) : B(x_l^{i+j}, \lambda^{i+j}) \cap B(x_k^i, \lambda^i) \neq \emptyset\},$$

$N_j^{i,k} = \#\mathcal{B}_j^{i,k}$ for $j \geq 0$, and

$$A_k^i = B(x_k^i, \lambda^i) \setminus \left(\bigcup_{j=1}^{\infty} \bigcup_{l \in I_{i+j}} B(x_l^{i+j}, \lambda^{i+j}) \right).$$

Now define $u_{i,k} = u \in L^{1,p}(\Omega)$ by

$$u(x) = \max \left(0, \min \left(1, 3 - \lambda^{-i} \text{dist} \left(x, x_k^i \right) \right) \chi_{\Omega_1}(x) \right).$$

Without loss of generality we can assume that the extension operator applied to any function $0 \leq u \leq 1$ also satisfies $0 \leq Eu \leq 1$. We then have

$$\|u\|_{L^{1,p}(\Omega)}^p \leq \int_{B(x_k^i, 3\lambda^i)} |\nabla u(x)|^p dx \leq (3^n \mathbf{m}_n(B(0, 1))) \lambda^{i(n-p)}. \tag{2.10}$$

By Proposition 2.2 and because $\lambda^{i_0} < r$, we have for $i \geq i_0$

$$\begin{aligned} & \min \left(\mathbf{m}_n(\Omega_1 \cap B(x_l^{i+j}, \lambda^{i+j}/2\sqrt{n})), \mathbf{m}_n(\Omega_2 \cap B(x_l^{i+j}, \lambda^{i+j}/2\sqrt{n})) \right) \\ & \geq C_1(n, p)\|E\|^{-n} \frac{\lambda^{n(i+j)}}{2^n n^{n/2}} \end{aligned}$$

for every $B(x_l^{i+j}, \lambda^{i+j}) \in \mathcal{B}_j^{i,k}$. (In the case $n - 1 < p < n$ Proposition 2.2 will give a better estimate with $\|E\|^{-p}$ in the above estimate. We shall comment about this case in a remark

at the end of the proof.) Applying Lemma 2.3 where again the removed set $F = \emptyset$, for the corresponding cube $Q(x_l^{i+j}, 2\lambda^{i+j}/\sqrt{n})$ centered at x_l^{i+j} and side length $2\lambda^{i+j}/\sqrt{n}$ (thus containing the ball $B(x_l^{i+j}, \lambda^{i+j}/\sqrt{n})$ and contained in the ball $B(x_l^{i+j}, \lambda^{i+j})$), we have

$$\begin{aligned} \int_{B(x_l^{i+j}, \lambda^{i+j})} |\nabla Eu(y)|^p dy &\geq \int_{Q(x_l^{i+j}, 2\lambda^{i+j}/\sqrt{n})} |\nabla Eu(y)|^p dy \\ &\geq \frac{C_2(n, p)C_1(n, p)^{(n-p)/n}}{n^{(n-p)/2}} \|E\|^{p-n} \lambda^{(n-p)(i+j)}. \end{aligned} \tag{2.11}$$

Thus, since $\sum_{B \in \mathcal{B}_j^{i,k}} \chi_B(x) \leq 3^n$ for all $x \in B(x_k^{i+j}, \lambda^{i+j})$, and by using (2.11) and (2.10), we get the estimate

$$\begin{aligned} \frac{C_2(n, p)C_1(n, p)^{(n-p)/n}}{n^{(n-p)/2}} N_j^{i,k} \|E\|^{p-n} \lambda^{(n-p)(i+j)} &\leq \sum_{B \in \mathcal{B}_j^{i,k}} \int_B |\nabla Eu(y)|^p dy \\ &\leq 3^n \int_{\mathbb{R}^n} |\nabla Eu(y)|^p dy \\ &\leq 3^n \|E\|^p \|u\|_{L^{1,p}(\Omega)}^p \\ &\leq (\mathbf{m}_n(B(0, 1))3^{2n}) \|E\|^p \lambda^{i(n-p)}. \end{aligned}$$

This implies the bound

$$N_j^{i,k} \leq \frac{\mathbf{m}_n(B(0, 1))3^{2n}n^{(n-p)/2}}{C_2(n, p)C_1(n, p)^{(n-p)/n}} \|E\|^n \lambda^{-j(n-p)} \tag{2.12}$$

for every $i, j \in \mathbb{N}$ with $i \geq i_0$ and $k \in I_i$.

Let us next estimate the \mathcal{H}^{n-1} -measure of the $(n - 1)$ -projections of the sets

$$F_{i,k} = \bigcup_{j=1}^{\infty} \bigcup_{B \in \mathcal{B}_j^{i,k}} B$$

for all $i \geq i_0$ and $k \in I_i$. By applying the estimate (2.12) and by the choice of λ , for every $i \geq i_0$ and $m = 1, \dots, n$,

$$\begin{aligned} \mathcal{H}^{n-1}(P_m(F_{i,k})) &\leq \sum_{j=1}^{\infty} N_j^{i,k} (2\lambda^{i+j})^{n-1} \\ &\leq 2^{n-1} \left(\frac{\mathbf{m}_n(B(0, 1))3^{2n}n^{(n-p)/2}}{C_2(n, p)C_1(n, p)^{(n-p)/n}} \right) \|E\|^n \lambda^{i(n-1)} \sum_{j=1}^{\infty} \lambda^{j(p-1)} \\ &= 2^{n-1} \left(\frac{\mathbf{m}_n(B(0, 1))3^{2n}n^{(n-p)/2}}{C_2(n, p)C_1(n, p)^{(n-p)/n}} \right) \|E\|^n \frac{\lambda^{p-1}}{1 - \lambda^{p-1}} \lambda^{i(n-1)} \\ &\leq \frac{C_1(n, p)\|E\|^{-n}}{2n \cdot 4^n} \left(\frac{2\lambda^i}{\sqrt{n}} \right)^{n-1}. \end{aligned}$$

Note that in Proposition 2.2 one can always assume $C_1(n, p)\|E\|^{-n} < 1$.

Suppose now that $s < \dim_{\mathcal{H}}(G)$. By Lemma 2.1 there exist $m_0 \geq i_0$ and $k_0 \in I_{m_0}$ such that $N_j^{m_0, k_0} \geq \lambda^{-js}$ for all $j \geq 0$.

For those fixed values m_0, k_0 and using the above estimate on the \mathcal{H}^{n-1} -measure of $P_m(F_{i,\ell})$, for the case $i = m_0 + j, \ell \in I_i, j \geq 0$, we can apply Lemma 2.3 to the function $u_{m_0, k_0} = u$, that was defined before. That is,

$$\begin{aligned} \int_{A_l^{m_0+j}} |\nabla Eu(y)|^p dy &\geq \int_{A_l^{m_0+j} \cap Q(x_l^{m_0+j}, 2\lambda^{m_0+j}/\sqrt{n})} |\nabla Eu(y)|^p dy \\ &= \int_{Q(x_l^{m_0+j}, 2\lambda^{m_0+j}/\sqrt{n}) \setminus F_{m_0+j,\ell}} |\nabla Eu(y)|^p dy \\ &\geq C(n, p) \|E\|^{p-n} \lambda^{(n-p)(m_0+j)}, \end{aligned}$$

where $A_l^{m_0+j} \subset B(x_l^{m_0+j}, \lambda^{m_0+j}) \in \mathcal{B}_j^{m_0, k_0}$. Now, by (2.10), and by summing over all the scales $j \geq 0$, we get

$$\begin{aligned} C(n) \|E\|^p \lambda^{m_0(n-p)} &\geq C(n) \|E\|^p \|u\|_{L^{1,p}(\Omega)}^p \geq C(n) \int_{\mathbb{R}^n} |\nabla Eu(y)|^p dy \\ &\geq \sum_{j=0}^{\infty} \sum_{\{l \in I_{m_0+j} : B(x_l^{m_0+j}, \lambda^{m_0+j}) \in \mathcal{B}_j^{m_0, k_0}\}} \int_{A_l^{m_0+j}} |\nabla Eu(y)|^p dy \\ &\geq \sum_{j=0}^{\infty} N_j^{m_0, k_0} C(n, p) \|E\|^{p-n} \lambda^{(n-p)(m_0+j)} \\ &\geq \sum_{j=0}^{\infty} \lambda^{-js} C(n, p) \|E\|^{p-n} \lambda^{(n-p)(m_0+j)} \\ &= C(n, p) \|E\|^{p-n} \frac{\lambda^{m_0(n-p)}}{1 - \lambda^{n-p-s}}. \end{aligned}$$

This implies (observe that by the choice of λ we have $\lambda \geq C(n, p) \|E\|^{\frac{-2n}{p-1}}$)

$$s \leq n - p - \frac{\log(1 - C(n, p) \|E\|^{-n})}{\log(\lambda)} \leq n - p - \frac{C(n, p)}{\|E\|^n \log(\|E\|)}.$$

Since $s < \dim_{\mathcal{H}}(G)$ was chosen arbitrarily, this concludes the proof of (2).

Remark 2.4 Let us make a remark on the case $n - 1 < p < n$. In this case, by applying Proposition 2.2 (2) we could slightly improve the estimates in the previous theorem. We would have that for the function u defined above,

$$\int_{A_l^{i+j}} |\nabla Eu(y)|^p dy \geq C(n, p) \|E\|^{-p\left(\frac{n-p}{n}\right)} \lambda^{(n-p)(i+j)},$$

and therefore

$$s \leq n - p - \frac{\log\left(1 - C(n, p) \|E\|^{-p\left(\frac{n-p}{n}\right)-p}\right)}{\log(\lambda)} \leq n - p - \frac{C(n, p)}{\|E\|^{2p-\frac{p^2}{n}} \log(\|E\|)}.$$

3 Two-Sided Points of BV-Extension Domains

For a given domain $\Omega \subset \mathbb{R}^n$ the space of functions of bounded variation in Ω is

$$BV(\Omega) = \{u \in L^1(\Omega) : \|Du\|(\Omega) < \infty\},$$

where

$$\|Du\|(\Omega) = \sup \left\{ \int_{\Omega} u \operatorname{div}(v) \, dx : v \in C_0^\infty(\Omega; \mathbb{R}^n), |v| \leq 1 \right\}$$

denotes the total variation of u on Ω . We endow this space with the norm $\|u\|_{BV(\Omega)} = \|u\|_{L^1(\Omega)} + \|Du\|(\Omega)$. We say that Ω is a BV-extension domain if there exists a constant $C > 0$ and a (not necessarily linear) extension operator $T: BV(\Omega) \rightarrow BV(\mathbb{R}^n)$ so that $Tu|_{\Omega} = u$ and

$$\|Tu\|_{BV(\mathbb{R}^n)} \leq C\|u\|_{BV(\Omega)}$$

for all $u \in BV(\Omega)$ and where $C > 0$ is an absolute constant, independent of u .

Let us point out that Ω being a $W^{1,1}$ -extension domain always implies that it is also a BV-extension domain (see [8, Lemma 2.4]).

A Lebesgue measurable subset $E \subset \mathbb{R}^n$ has finite perimeter in Ω if $\chi_E \in BV(\Omega)$, where χ_E denotes the indicator function of the set E . We set $P(E, \Omega) = \|D\chi_E\|(\Omega)$ and call it the perimeter of E in Ω . Moreover, the measure theoretic boundary of a set $E \subset \mathbb{R}^n$ is defined as

$$\partial^M E = \left\{ x \in \mathbb{R}^n : \limsup_{r \searrow 0} \frac{|E \cap B(x, r)|}{|B(x, r)|} > 0 \text{ and } \limsup_{r \searrow 0} \frac{|(\mathbb{R}^n \setminus E) \cap B(x, r)|}{|B(x, r)|} > 0 \right\},$$

and for a set of finite perimeter in Ω one always has $P(E, \Omega) = \mathcal{H}^{n-1}(\partial^M E \cap \Omega)$. Finally, let us recall the useful coarea formula for BV functions. Namely, for a given a function $u \in BV(\Omega)$, the superlevel sets $u_t = \{x \in \Omega : u(x) \geq t\}$ have finite perimeter in Ω for almost every $t \in \mathbb{R}$ and

$$\|Du\|(\Omega) = \int_{-\infty}^{\infty} P(u_t, \Omega) \, dt. \tag{3.1}$$

Proof (*Proof of Theorem 1.3*) We want to prove that \mathcal{T}_{Ω} has σ -finite $(n - 1)$ -dimensional Hausdorff measure. Similarly to the beginning part of the proof of Theorem 1.2 and reasoning by contradiction assume that there exists a set $G \subset \partial\Omega \cap B(x_0, r_0)$, with $r_0 \in (0, 1)$, $x_0 \in G$, and two connected components $\Omega_1, \Omega_2 \subset B(x_0, 3r_0) \cap \Omega$ for which $G \subset \partial\Omega_1 \cap \partial\Omega_2$ such that $\mathcal{H}^{n-1}(G) = \infty$.

Consider the set $E = B(x_0, r_0) \cap \Omega_1$ for which we have $\chi_E \in BV(\Omega)$. Take any measurable function v in \mathbb{R}^n so that $v|_{\Omega} = \chi_E$. Note that $\tilde{E}_t \cap \Omega = E$ for every $t \in (0, 1)$ for the superlevel sets $\tilde{E}_t = \{x \in \mathbb{R}^n : v(x) \geq t\}$. By using the measure density condition proved in [3, Proposition 2.3] applied to both connected components Ω_1 and Ω_2 , we get that there exists $c > 0$ so that

$$\mathbf{m}_n(\Omega_i \cap B(x, r)) \geq cr^n$$

for $i = 1, 2$ and all $x \in G, r \in (0, r_0)$. In particular, for every $x \in G$ we have

$$\limsup_{r \searrow 0} \frac{\mathbf{m}_n(B(x, r) \cap \tilde{E}_t)}{\mathbf{m}_n(B(x, r))} \geq \limsup_{r \searrow 0} \frac{\mathbf{m}_n(B(x, r) \cap \Omega_1)}{\mathbf{m}_n(B(x, r))} > 0$$

and

$$\limsup_{r \searrow 0} \frac{\mathbf{m}_n(B(x, r) \cap (\mathbb{R}^n \setminus \tilde{E}_t))}{\mathbf{m}_n(B(x, r))} \geq \limsup_{r \searrow 0} \frac{\mathbf{m}_n(B(x, r) \cap \Omega_2)}{\mathbf{m}_n(B(x, r))} > 0.$$

This means that $G \subset \partial^M \tilde{E}_t$. Hence, $\mathcal{H}^{n-1}(\partial^M \tilde{E}_t) \geq \mathcal{H}^{n-1}(G) = \infty$, so \tilde{E}_t does not have finite perimeter in \mathbb{R}^n for any $t \in (0, 1)$. Hence, by the coarea formula (3.1), $v \notin BV(\mathbb{R}^n)$.

4 Examples

4.1 Sharpness of the Estimate for $p = 1$

The following example shows the sharpness of Theorem 1.2 (1). In this case, where $p = 1$, we do not need to care about the norm of the extension operator and consequently, we can rely on previous non-quantitative characterizations of $W^{1,1}$ -extension domains.

Example 4.1 Let us define

$$\Omega_2 = (-1, 1)^2 \setminus \{(x, y) : |y| \leq \text{dist}(x, C), 0 \leq x \leq 1\}$$

with $C \subset [0, 1]$ a Cantor set with $\dim_{\mathcal{H}}(C) = 1$ and $\mathcal{H}^1(C) = 0$. See Fig. 1 for an illustration of the domain Ω_2 .

We claim that Ω_2 is a $W^{1,1}$ -extension domain and that

$$\dim_{\mathcal{H}}(\mathcal{T}) = 1.$$




Fig. 1 The domain showing the sharpness of Theorem 1.2 (1). The set \mathcal{T} here is the fat Cantor set without its left-most point

It is easy to see that $\dim_{\mathcal{H}}(\mathcal{T}) = 1$, since $\mathcal{T} = (C \times \{0\}) \setminus \{(0, 0)\}$. In order to see that Ω_2 is a $W^{1,1}$ -extension domain, one can use the following characterization from [10] for bounded planar simply connected domains: Ω is a $W^{1,1}$ -extension domain if and only if

$$\begin{aligned} &\text{there exists a constant } K \text{ so that for every } x, y \in \Omega^c \text{ there exists a curve} \\ &\gamma \subset \Omega^c \text{ with } x, y \in \gamma, \ell(\gamma) \leq K|x - y|, \text{ and } \mathcal{H}^1(\gamma \cap \partial\Omega) = 0. \end{aligned} \tag{4.1}$$

Now, the domain Ω_2 clearly satisfies (4.1) and is thus a $W^{1,1}$ -extension domain.

Let us remark that Example 4.1 can also be generalized to higher dimensions $n > 2$ by defining $\Omega \subset \mathbb{R}^n$ as a product $\Omega_2 \times (-1, 1)^{n-2}$. It is then clear that

$$\dim_{\mathcal{H}}(\mathcal{T}) = n - 1.$$

The fact that Ω is a $W^{1,1}$ -extension domain does not seem to immediately follow from known explicit results. One way to see that Ω is a $W^{1,1}$ -extension domain is the following. Observe that the proof in [12] of the fact that a product of $W^{1,p}$ -extension domains, with $p > 1$, is again a $W^{1,p}$ -extension domain relies on the explicit form of the extension operators (which in that case can always be assumed to be a Whitney extension operators). In the case $p = 1$ it is unknown if the extension can always be done with a Whitney-type extension. However, the extension operator constructed in [10] for simply connected planar domains, and in particular for Ω_2 is of Whitney-type. Thus, the argument in [12] goes through for our product domain Ω .

4.2 A Bound for the Estimates for $1 < p < n$

The case $1 < p < n$ requires more work than the case $p = 1$, since the estimate in Theorem 1.2 depends on the norm of the extension operator. The assignment of reflected cubes in the construction of the extension operator, and the estimate of the norm of the extension operator follow roughly the proof of the sufficiency of the characterizing curve condition of planar simply connected $W^{1,p}$ -extension domains [9].

Let us describe the family of domains Ω_λ we consider, where $\lambda \in (0, 1/2)$ refers to the contraction ratio of the Cantor set $C_\lambda \subset \mathbb{R}^{n-1}$. The Cantor sets C_λ we use are the standard ones obtained as $C_\lambda = \prod_{i=1}^{n-1} K_\lambda$ with K_λ being the Cantor set on the unit interval given as the attractor of the iterated function system $\{f_1(x) = \lambda x, f_2(x) = \lambda x + 1 - \lambda\}$.

We define first a set

$$D = (0, 1)^{n-2} \times ((-2, 1) \times (-3/2, 3/2) \setminus [-1, 0] \times [-1, 1])$$

and then the actual domain by carving out part of D :

$$\Omega_\lambda = D \setminus N_\lambda,$$

where

$$N_\lambda = \{(x_1, \dots, x_n) \in [0, 1]^n : |x_n| \leq \text{dist}((x_1, \dots, x_{n-1}), C_\lambda)\}.$$

Then, the set of two-sided points for Ω_λ is

$$\mathcal{T}_{\Omega_\lambda} = C_\lambda \times \{0\}$$

and so it has dimension

$$\dim_{\mathcal{H}}(\mathcal{T}_{\Omega_\lambda}) = \dim_{\mathcal{H}}(C_\lambda) = -\frac{(n - 1) \log 2}{\log \lambda}. \tag{4.2}$$

Our aim is to build an extension operator E_λ from $L^{1,p}(\Omega_\lambda)$ to $L^{1,p}(\mathbb{R}^n)$ for which we have

$$\dim_{\mathcal{H}}(C_\lambda) \geq n - p - \frac{C(n, p)}{\|E_\lambda\|}.$$

As will be explicit in (4.5), the constructed operator E_λ will be bounded only for p in the range

$$1 < p < n - \dim_{\mathcal{H}}(C_\lambda).$$

It is enough to construct an extension operator $E_\lambda: L^{1,p}(\Omega_\lambda) \rightarrow L^{1,p}(D)$, since the extension from $L^{1,p}(D)$ to $L^{1,p}(\mathbb{R}^n)$ is independent of λ , and exists since D is a Lipschitz domain. Moreover, our definition of E_λ will be independent of p and will give a bounded operator between the Sobolev spaces $W^{1,p}(\Omega_\lambda)$ and $W^{1,p}(D)$. From now on we consider $\lambda \in (0, 1/2)$ fixed and we denote the extension operator by E instead of E_λ to simplify the notation.

Below by a dyadic cube we mean a set of the form $Q = [0, 2^{-k}]^n + j \subset \mathbb{R}^n$ for some $k \in \mathbb{Z}$ and $j \in 2^{-k}\mathbb{Z}^n$. Let $\mathcal{W} = \{Q_i\}_{i \in \mathbb{N}}$ be a Whitney decomposition of the interior of N_λ and $\tilde{\mathcal{W}} = \{\tilde{Q}_i\}_{i \in \mathbb{N}}$ a Whitney decomposition of $\mathbb{R}^n \setminus N_\lambda$. This is

- (W1) Each Q_i is a closed dyadic cube inside N_λ .
- (W2) $N_\lambda = \bigcup_i Q_i$ and for every $i \neq j$ we have $\text{int}(Q_i) \cap \text{int}(Q_j) = \emptyset$.
- (W3) For every i we have $\sqrt{n}\ell(Q_i) \leq \text{dist}(Q_i, \partial N_\lambda) \leq 4\sqrt{n}\ell(Q_i)$.
- (W4) If $Q_i \cap Q_j \neq \emptyset$, we have $\frac{1}{4}\ell(Q_i) \leq \ell(Q_j) \leq 4\ell(Q_i)$.

The definition of $\tilde{\mathcal{W}}$ goes parallel. See [16, Chapter VI] for the existence of such Whitney decompositions. Consider also the subfamily of Whitney cubes

$$\mathcal{V} = \{Q \in \mathcal{W} : Q \cap ([0, 1]^{n-1} \times \{0\}) \neq \emptyset\}.$$

Let us also distinguish an important subset of Ω_λ , that we call

$$\tilde{Q}_0 = (0, 1)^{n-2} \times ((-2, 1) \times (-3/2, 3/2) \setminus [-1, 1] \times [-1, 1]).$$

Note that for every $\tilde{Q} \in \tilde{\mathcal{W}}$ we have $\partial((0, 1)^{n-1} \times (-1, 1)) \cap \text{int}(\tilde{Q}) = \emptyset$. Next we choose $\{\psi_i\}_{i \in \mathbb{N}}$ a partition of unity subordinate to the open cover $\{(9/8)\text{int}(Q_i)\}_{i \in \mathbb{N}}$ and so that $|\nabla \psi_i(x)| \lesssim \ell(Q_i)^{-1}$.

Now, given $u \in L^{1,p}(\Omega_\lambda)$ we assign a value

$$a_i = \frac{1}{\mathbf{m}_n(\tilde{Q}_{R(i)})} \int_{\tilde{Q}_{R(i)}} u(x) dx$$

for every $i \in \mathbb{N}$, where the function $R: \mathbb{N} \rightarrow \mathbb{N}$ is defined as follows. If $Q_i \in \mathcal{V}$, then $R(i) = 0$. If $Q_i \notin \mathcal{V}$ we assign $R(i)$ to be the unique index so that Q_i and $\tilde{Q}_{R(i)}$ belong to the same half-space $\{x_n < 0\}$ or $\{x_n > 0\}$, $P_n(Q_i) \subset P_n(\tilde{Q}_{R(i)})$, $\ell(\tilde{Q}_{R(i)}) \leq 2\ell(Q_i)$ where $P_n: \mathbb{R}^n \rightarrow \mathbb{R}^{n-1}: (x_1, \dots, x_n) \mapsto (x_1, \dots, x_{n-1})$, and $\tilde{Q}_{R(i)}$ is the closest cube to Q_i with the first three properties.

Now, we define the extension of the function u by

$$Eu(x) = \begin{cases} u(x), & \text{if } x \in \Omega_\lambda \\ \sum_{i=1}^\infty a_i \psi_i(x), & \text{if } x \in \text{int}(N_\lambda), \\ 0, & \text{if } x \in \partial N_\lambda \cap D, \end{cases} \tag{4.3}$$

Let us explain first why $Eu \in L^{1,p}(D)$. On the one hand, $Eu \in L^{1,p}(\Omega_\lambda)$ and, on the other hand, we will see later that $Eu \in L^{1,p}(\text{int}(N_\lambda))$. Since C_λ has Hausdorff-dimension

strictly less than $n - 1$, we have that almost every line parallel to the coordinate axis does not meet the set C_λ . Consequently, any function in $L^{1,p}(D \setminus C_\lambda)$ is also ACL on D , implying that $L^{1,p}(D) = L^{1,p}(D \setminus C_\lambda)$ as sets. So, in order to then have $Eu \in L^{1,p}(D)$, it suffices to show that $Eu \in L^{1,p}(D \setminus C_\lambda)$. This follows by noticing that by the definition of the operator E , the trace of u

$$Tu(x) = \lim_{r \searrow 0} \frac{1}{\mathbf{m}_n(B(x, r) \cap \Omega_\lambda)} \int_{B(x,r) \cap \Omega_\lambda} u(y) dy$$

on $\partial N_\lambda \setminus C_\lambda$ coincides with that of $Eu|_{\text{int}(N_\lambda)}$

$$TEu(x) = \lim_{r \searrow 0} \frac{1}{\mathbf{m}_n(B(x, r) \cap N_\lambda)} \int_{B(x,r) \cap \text{int}(N_\lambda)} Eu(y) dy.$$

To conclude that E is an extension operator it remains to control the L^p -norm of the gradient of the extension on $\text{int}(N_\lambda)$ by the L^p -norm of the gradient of the initial function.

We know that $\text{supp}(\psi_i) \subseteq \frac{9}{8}Q_i$ and that $|\nabla\psi_i(x)| \lesssim \ell(Q_i)^{-1}$ for every x and $i \in \mathbb{N}$, so it is clear that for every $x \in Q_i$,

$$|\nabla Eu(x)| \leq \left| \sum_{Q_j \cap Q_i \neq \emptyset} \nabla\psi_j(x)(a_j - a_i) \right| \lesssim \sum_{Q_j \cap Q_i \neq \emptyset} \ell(Q_j)^{-1}|a_j - a_i|.$$

Now, if we take a cube $Q_i \in \mathcal{W}$, using that at most $C(n)$ other cubes of the Whitney decomposition are intersecting it and that $\ell(Q_i) \sim \ell(Q_j)$ if $Q_i \cap Q_j \neq \emptyset$ we write

$$\begin{aligned} \|\nabla Eu\|_{L^p(Q_i)}^p &= \int_{Q_i} |\nabla Eu(x)|^p dx \lesssim \int_{Q_i} \sum_{Q_j \cap Q_i \neq \emptyset} \ell(Q_j)^{-p}|a_j - a_i|^p dx \\ &\lesssim \ell(Q_i)^{n-p} \sum_{Q_j \cap Q_i \neq \emptyset} |a_i - a_j|^p. \end{aligned} \tag{4.4}$$

It will be useful to work with chains of Whitney cubes that we next define. Given i, j so that $Q_i \cap Q_j \neq \emptyset$ and $Q_i, Q_j \notin \mathcal{V}$ we define the chain of cubes joining $\tilde{Q}_{R(i)}$ with $\tilde{Q}_{R(j)}$, and denote it by $C(\tilde{Q}_{R(i)}, \tilde{Q}_{R(j)})$, to be a minimal family of Whitney cubes whose union's interior is a connected set containing both the interiors of $\tilde{Q}_{R(i)}$ and $\tilde{Q}_{R(j)}$. Note that we always have $\#C(\tilde{Q}_{R(i)}, \tilde{Q}_{R(j)}) \leq C_0(n)$. Suppose $Q_i \notin \mathcal{V}$ is a cube so that there exists $Q_j \in \mathcal{V}$ with $Q_i \cap Q_j \neq \emptyset$. For the associated cube $\tilde{Q}_{R(i)}$ we define $C(\tilde{Q}_{R(i)}, \tilde{Q}_0)$ as a minimal family of sets in $\tilde{W} \cup \{\tilde{Q}_0\}$ whose union's interior is a connected set containing both the interiors of $\tilde{Q}_{R(i)}$ and \tilde{Q}_0 and so that every $\tilde{Q} \in C(\tilde{Q}_{R(i)}, \tilde{Q}_0)$ satisfies $P_n(\tilde{Q}_{R(i)}) \subset P_n(\tilde{Q})$.

We can assume there is an order in the chain when moving from $\tilde{Q}_{R(i)}$ to $\tilde{Q}_{R(j)}$ and call \tilde{Q}_{next} the next cube in the chain after \tilde{Q} . We write

$$C(\tilde{Q}_{R(i)}, \tilde{Q}_{R(j)}) = \{\tilde{Q}_{R(i)}, (\tilde{Q}_{R(i)})_{\text{next}}, \dots, \tilde{Q}_{R(j)}\}.$$

To ease the notation in the following sums from now on we write

$$C_{i,j} = C(\tilde{Q}_{R(i)}, \tilde{Q}_{R(j)}) \setminus \{\tilde{Q}_{R(j)}\} \text{ and } C_{i,0} = C(\tilde{Q}_{R(i)}, \tilde{Q}_0) \setminus \{\tilde{Q}_0\}.$$

Note that if $Q_i \notin \mathcal{V}$ and there does not exist Q_j such that $Q_j \in \mathcal{V}$ and $Q_i \cap Q_j \neq \emptyset$ we define $C_{i,0} = \emptyset$.

Let also write

$$\mathcal{I} = \{\tilde{Q} \in \tilde{W} : \tilde{Q} = \tilde{Q}_{R(i)} \text{ for some } i \geq 1\}.$$

We assert that the following claim holds.

Claim 4.2 With the above notation and for every $r > 0$ we have the following.

(i) For every $Q_i \notin \mathcal{V}$

$$\begin{aligned} \|\nabla Eu\|_{L^p(Q_i)}^p &\lesssim \sum_{\{\tilde{Q} \in \mathcal{I} : \#C(\tilde{Q}_{R(i)}, \tilde{Q}) \leq C_0(n)\}} \int_{\tilde{Q}} |\nabla u(x)|^p dx \\ &\quad + \ell(Q_i)^{n-p-rp} D(r, p) \sum_{\tilde{Q} \in C_{i,0}} \ell(\tilde{Q})^{p-n+rp} \int_{\tilde{Q} \cup \tilde{Q}_{\text{next}}} |\nabla u(x)|^p dx, \end{aligned}$$

where $D(r, p) = (1 - 2^{-\frac{rp}{p-1}})^{1-p}$, and for every $Q_i \in \mathcal{V}$, we have

$$\|\nabla Eu\|_{L^p(Q_i)}^p \lesssim \ell(Q_i)^{n-p-rp} \sum_{\substack{Q_j \cap Q_i \neq \emptyset \\ Q_j \notin \mathcal{V}}} D(r, p) \sum_{\tilde{Q} \in C_{j,0}} \ell(\tilde{Q})^{p-n+rp} \int_{\tilde{Q} \cup \tilde{Q}_{\text{next}}} |\nabla u(x)|^p dx.$$

(ii) For a given $\tilde{Q} \in \tilde{\mathcal{W}}$ and $k \in \mathbb{Z}$ we have

$$\begin{aligned} &\#\{Q_i \in \mathcal{W} \setminus \mathcal{V} : Q_i \text{ has a neighbouring cube in } \mathcal{V}, \ell(\tilde{Q}) = 2^k \ell(Q_i), \tilde{Q} \in C_{i,0}\} \\ &\lesssim 2^{-(n-1) \frac{k \log 2}{\log \lambda}}. \end{aligned}$$

Assuming for a moment that the claim is true let us show how one can estimate the full norm $\|\nabla Eu\|_{L^p(N_\lambda)}^p$. We first use Claim 4.2 (i) and change the order of summation to get

$$\begin{aligned} \|\nabla Eu\|_{L^p(N_\lambda)}^p &= \sum_{Q_i \in \mathcal{W}} \|\nabla Eu\|_{L^p(Q_i)}^p = \sum_{Q_i \notin \mathcal{V}} \|\nabla Eu\|_{L^p(Q_i)}^p + \sum_{Q_i \in \mathcal{V}} \|\nabla Eu\|_{L^p(Q_i)}^p \\ &\lesssim \sum_{\tilde{Q} \in \mathcal{I}} \sum_{\{i : \#C(\tilde{Q}_{R(i)}, \tilde{Q}) \leq C_0(n)\}} \int_{\tilde{Q}} |\nabla u(x)|^p dx \\ &\quad + 2 \sum_{\tilde{Q} \in \tilde{\mathcal{W}}} \sum_{\tilde{Q} \in C_{i,0}} \ell(Q_i)^{n-p-rp} D(r, p) \ell(\tilde{Q})^{p-n+rp} \int_{\tilde{Q} \cup \tilde{Q}_{\text{next}}} |\nabla u(x)|^p dx \\ &\lesssim \|\nabla u\|_{L^p(\Omega_\lambda)}^p + \sum_{\tilde{Q} \in \tilde{\mathcal{W}}} \sum_{\tilde{Q} \in C_{i,0}} \ell(Q_i)^{n-p-rp} D(r, p) \ell(\tilde{Q})^{p-n+rp} \\ &\quad \times \int_{\tilde{Q} \cup \tilde{Q}_{\text{next}}} |\nabla u(x)|^p dx. \end{aligned}$$

Moreover, by Claim 4.2 (ii) it follows that

$$\sum_{\{i : \tilde{Q} \in C_{i,0}\}} \ell(Q_i)^{n-p-rp} \lesssim \sum_{k=0}^{\infty} 2^{-(n-1) \frac{k \log 2}{\log \lambda}} (2^{-k} \ell(\tilde{Q}))^{n-p-rp} = \frac{\ell(\tilde{Q})^{n-p-rp}}{1 - 2^{-n+p-(n-1) \frac{\log 2}{\log \lambda} + rp}},$$

under the assumption

$$-n + p - (n - 1) \frac{\log 2}{\log \lambda} + rp < 0. \tag{4.5}$$

So, joining these facts together we get

$$\begin{aligned} \|\nabla Eu\|_{L^p(N_\lambda)}^p &\lesssim \|\nabla u\|_{L^p(\Omega_\lambda)}^p \\ &+ \sum_{\tilde{Q} \in \tilde{\mathcal{W}}_1} D(r, p) \left(\frac{1}{1 - 2^{-n+p-(n-1)\frac{\log 2}{\log \lambda} + rp}} \right) \int_{\tilde{Q} \cup \tilde{Q}_{next}} |\nabla u(x)|^p dx \\ &\lesssim \left(\frac{1}{1 - 2^{-\frac{rp}{p-1}}} \right)^{p-1} \left(\frac{1}{1 - 2^{-n+p-(n-1)\frac{\log 2}{\log \lambda} + rp}} \right) \|\nabla u\|_{L^p(\Omega_\lambda)}^p, \end{aligned}$$

where $\tilde{\mathcal{W}}_1 = \{\tilde{Q} \in \tilde{\mathcal{W}} : \#\{i : \tilde{Q} \in C_{i,0}\} > 0\}$ and $\tilde{Q}_{next} \in C_{i,0}$ for some $i \in \mathbb{N}$ such that $\tilde{Q} \in C_{i,0}$.

Choosing $r = \frac{p-1}{p^2}(n - p - \dim_{\mathcal{H}}(C_\lambda))$, we conclude that

$$\|E\| \lesssim \frac{1}{1 - 2^{\frac{1}{p}(-n+p+\dim_{\mathcal{H}}(C_\lambda))}},$$

which yields

$$\dim_{\mathcal{H}}(C_\lambda) \geq n - p - \frac{C(n, p)}{\|E\|}.$$

Let us now prove the Claim 4.2.

Proof (Proof of Claim 4.2) To prove (i) we need to estimate $|a_i - a_j|^p$ in the expression (4.4). First note that from (4.4) one gets

$$\begin{aligned} |a_i - a_j|^p &\leq \left(\sum_{\tilde{Q} \in C_{i,j}} \left| \frac{1}{\mathbf{m}_n(\tilde{Q})} \int_{\tilde{Q}} u(x) dx - \frac{1}{\mathbf{m}_n(\tilde{Q}_{next})} \int_{\tilde{Q}_{next}} u(x) dx \right| \right)^p \\ &\lesssim \left(\sum_{\tilde{Q} \in C_{i,j}} \ell(\tilde{Q})^{1-n} \int_{\tilde{Q} \cup \tilde{Q}_{next}} |\nabla u(x)| dx \right)^p, \end{aligned} \tag{4.6}$$

where we are using the Poincaré inequality in the last line (see [6, Lemma 2.2] and also [1]). Observe that if $Q_j, Q_i \in \mathcal{V}$ then $R(i) = R(j) = 0$ and $|a_j - a_i| = 0$. We now consider two cases.

- (1) Suppose i, j are so that $Q_i \cap Q_j \neq \emptyset$ and $Q_i, Q_j \notin \mathcal{V}$. Then using (4.6), that $\#C(\tilde{Q}_{R(i)}, \tilde{Q}_{R(j)}) \leq C_0(n)$, that the sides of the cubes of the chain are comparable to that of $\tilde{Q}_{R(i)}$, and hence that of Q_i , and applying Hölder inequality

$$\begin{aligned} |a_i - a_j|^p &\lesssim \sum_{\tilde{Q} \in C_{i,j}} \ell(\tilde{Q})^{(1-n)p} \left(\int_{\tilde{Q} \cup \tilde{Q}_{next}} |\nabla u(x)| dx \right)^p \\ &\lesssim \ell(Q_i)^{p-n} \sum_{\tilde{Q} \in C_{i,j}} \int_{\tilde{Q} \cup \tilde{Q}_{next}} |\nabla u(x)|^p dx. \end{aligned} \tag{4.7}$$

(2) Suppose i, j are such that $Q_i \cap Q_j \neq \emptyset, Q_j \in \mathcal{V}$ (then $R(j) = 0$) and $Q_i \notin \mathcal{V}$ then we fix $r > 0$ to be determined later and apply Hölder inequality to (4.6) to get

$$\begin{aligned}
 |a_i - a_j|^p &\lesssim \left(\sum_{\tilde{Q} \in C_{i,0}} \ell(\tilde{Q})^{-r} \ell(\tilde{Q})^{1-n+r} \int_{\tilde{Q} \cup \tilde{Q}_{\text{next}}} |\nabla u(x)| dx \right)^p \\
 &\leq \left(\sum_{\tilde{Q} \in C_{i,0}} \ell(\tilde{Q})^{-r \frac{p}{p-1}} \right)^{p-1} \left(\sum_{\tilde{Q} \in C_{i,0}} \ell(\tilde{Q})^{(1-n+r)p} \left(\int_{\tilde{Q} \cup \tilde{Q}_{\text{next}}} |\nabla u(x)| dx \right)^p \right) \\
 &\lesssim \left(\sum_{k=0}^{\infty} (2^k \ell(\tilde{Q}_{R(i)}))^{-r \frac{p}{p-1}} \right)^{p-1} \left(\sum_{\tilde{Q} \in C_{i,0}} \ell(\tilde{Q})^{p-n+rp} \int_{\tilde{Q} \cup \tilde{Q}_{\text{next}}} |\nabla u(x)|^p dx \right) \\
 &\lesssim D(r, p) \ell(\tilde{Q}_{R(i)})^{-rp} \left(\sum_{\tilde{Q} \in C_{i,0}} \ell(\tilde{Q})^{p-n+rp} \int_{\tilde{Q} \cup \tilde{Q}_{\text{next}}} |\nabla u(x)|^p dx \right).
 \end{aligned}
 \tag{4.8}$$

Going back to equation (4.4) for any $Q_i \notin \mathcal{V}$, using (4.7) and (4.8) we have

$$\begin{aligned}
 \|\nabla Eu\|_{L^p(Q_i)}^p &\lesssim \ell(Q_i)^{n-p} \left(\sum_{Q_j \cap Q_i \neq \emptyset, Q_j \notin \mathcal{V}} |a_i - a_j|^p + \sum_{Q_j \cap Q_i \neq \emptyset, Q_j \in \mathcal{V}} |a_i - a_j|^p \right) \\
 &\lesssim \ell(Q_i)^{n-p} \left(\sum_{Q_j \cap Q_i \neq \emptyset, Q_j \notin \mathcal{V}} \ell(Q_i)^{p-n} \sum_{\tilde{Q} \in C_{i,j}} \int_{\tilde{Q} \cup \tilde{Q}_{\text{next}}} |\nabla u(x)|^p dx \right. \\
 &\quad \left. + \sum_{Q_j \cap Q_i \neq \emptyset, Q_j \in \mathcal{V}} D(r, p) \ell(Q_i)^{-rp} \sum_{\tilde{Q} \in C_{i,0}} \ell(\tilde{Q})^{p-n+rp} \int_{\tilde{Q} \cup \tilde{Q}_{\text{next}}} |\nabla u(x)|^p dx \right) \\
 &\lesssim \sum_{\{\tilde{Q} \in \mathcal{I} : \#C(\tilde{Q}_{R(i)}, \tilde{Q}) \leq C_0(n)\}} \int_{\tilde{Q}} |\nabla u(x)|^p dx \\
 &\quad + \ell(Q_i)^{n-p-rp} D(r, p) \sum_{\tilde{Q} \in C_{i,0}} \ell(\tilde{Q})^{p-n+rp} \int_{\tilde{Q} \cup \tilde{Q}_{\text{next}}} |\nabla u(x)|^p dx,
 \end{aligned}$$

and if $Q_i \in \mathcal{V}$, using only (4.8)

$$\begin{aligned}
 \|\nabla Eu\|_{L^p(Q_i)}^p &\lesssim \ell(Q_i)^{n-p} \sum_{Q_j \cap Q_i \neq \emptyset, Q_j \notin \mathcal{V}} |a_i - a_j|^p \\
 &\lesssim \ell(Q_i)^{n-p-rp} \sum_{Q_j \cap Q_i \neq \emptyset, Q_j \notin \mathcal{V}} D(r, p) \sum_{\tilde{Q} \in C_{j,0}} \ell(\tilde{Q})^{p-n+rp} \\
 &\quad \times \int_{\tilde{Q} \cup \tilde{Q}_{\text{next}}} |\nabla u(x)|^p dx.
 \end{aligned}$$

which proves (i).

Let us prove (ii). Let us write the Cantor set $C_\lambda \subset [0, 1]^{n-1}$ as

$$C_\lambda = \bigcap_{i=0}^{\infty} C_\lambda^i = \bigcap_{i=0}^{\infty} \bigcup_{1 \leq j \leq 2^{(n-1)i}} I_{i,j},$$

where $I_{i,j}$ is a translated copy of $[0, \lambda^i]^{n-1}$ for all $i = 0, 1, 2, \dots$ and $j = 1, 2, \dots, 2^{(n-1)i}$. It is clear that for $i < i'$, any cube $I_{i,j}$ contains $2^{(n-1)(i'-i)}$ cubes of side length $\lambda^{i'}$.

Fix $\tilde{Q} \in \tilde{\mathcal{W}}$ and $k \in \mathbb{N}$. Let $t \in \mathbb{N}$ such that $\ell(\tilde{Q}) = 2^{-t}$. We count the cardinality of

$$A = \left\{ Q_i \in \mathcal{W} \setminus \mathcal{V} : Q_i \text{ has a neighbouring cube in } \mathcal{V}, \ell(\tilde{Q}) = 2^k \ell(Q_i), \tilde{Q} \in C_{i,0} \right\}.$$

Define $B = \{P_n(Q_i)\}_{Q_i \in A}$, where $P_n: \mathbb{R}^n \rightarrow \mathbb{R}^{n-1}: (x_1, \dots, x_n) \mapsto (x_1, \dots, x_{n-1})$.

Let m be the least positive integer such that $\lambda^m < 2^{-t}$ and let l be the least positive integer so that $\lambda^l \leq 2^{-t-k}$. By the properties of the Whitney decomposition, the construction of the Cantor set and the minimality of m it is enough to count $\#\{Q \in B : \text{dist}(Q, I_{m,j}) \leq C(n)\ell(Q)\}$ for a fixed $I_{m,j}$. Moreover, by the selection of l none of the cubes $I_{l,j}$ contains any $Q \in B$.

Because $\lambda^l \leq \ell(Q)$, we have

$$\#\{Q \in B : \text{dist}(Q, I_{l,j'}) \leq C(n)\ell(Q)\} \leq c(n)$$

for all $I_{l,j'} \subset I_{m,j}$. Finally since $I_{m,j} \cap C_\lambda^l$ is a disjoint union of $2^{(n-1)(l-m)}$ cubes $I_{l,j'}$ of side length λ^l we conclude that

$$\#A \lesssim \#B \leq c(n)2^{(n-1)(l-m)} \lesssim 2^{-k(n-1) \frac{\log 2}{\log \lambda}}.$$

Acknowledgements The authors thank Panu Lahti for valuable comments on the earlier version of this paper.

Author Contributions All the authors of this manuscript have participated in conducting the research and writing the manuscript.

Funding Open Access funding provided by University of Jyväskylä (JYU). The authors of this manuscript have received financial support from the Academy of Finland, grant no. 314789.

Declarations

Competing interests The authors of this manuscript have no relevant financial or non-financial interests to disclose.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Bojarski, B (1988) Remarks on Sobolev imbedding inequalities. Complex analysis, Joensuu 1987, pp 52–68, Lecture Notes in Math., 1351, Springer, Berlin

2. Evans, LC and Gariepy, RF (2015) Measure theory and fine properties of functions. Revised edition. Textbooks in Mathematics. CRC Press, Boca Raton, FL, xiv+299 pp
3. García-Bravo, M., Rajala, T.: Strong BV -extension and $W^{1,1}$ -extension domains. *J. Funct. Anal.* **283**(10), 109665 (2022)
4. Hajlasz, P., Koskela, P., Tuominen, H.: Sobolev embeddings, extensions and measure density condition. *J. Funct. Anal.* **254**(5), 1217–1234 (2008)
5. Hakobyan, H., Herron, D.: Euclidean quasiconvexity. *Ann. Acad. Sci. Fenn. Math.* **33**(1), 205–230 (2008)
6. Jones, P.W.: Quasiconformal mappings and extendability of Sobolev functions. *Acta Math.* **47**, 71–88 (1981)
7. Koskela, P (1990) Capacity extension domains, *Ann. Acad. Sci. Fenn. Ser. A I Math. Dissertationes No. 73*, p 42
8. Koskela, P, Miranda Jr., M and Shanmugalingam, N (2010) Geometric properties of planar BV extension domains. In: Around the Research of Prof. Maz'ya I, in: International Mathematical Series, pp 255–272, Function Spaces; Topics (Springer collection)
9. Koskela, P, Rajala, T and Zhang, Y (2015) A geometric characterization of planar Sobolev extension domains, preprint
10. Koskela, P, Rajala, T and Zhang, Y (2017) Planar $W^{1,1}$ -extension domains, preprint
11. Koskela, P., Yang, D., Zhou, Y.: A Jordan Sobolev extension domain. *Ann. Acad. Sci. Fenn. Math.* **35**(1), 309–320 (2010)
12. Koskela, P., Zhu, Z.: Product of extension domains is still an extension domain, *Indiana Univ. Math. J.* **69**, 137–150 (2020)
13. Lučić, D., Rajala, T., Takanen, J.: Dimension estimates for the boundary of planar Sobolev extension domains. *Adv. Calc. Var.* **16**(2), 517–528 (2023)
14. Maz'ya, V.: Extension of functions from Sobolev spaces. *Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI)* **113**, 231–236 (1981)
15. Romanov, AS (1993) On the extension of functions that belong to Sobolev spaces, *Sibirsk. Mat. Zh.* **34**:149–152; English transl. in *Siberian Math. J.* **34** (1993), 723–726
16. Stein, E.M.: *Singular Integrals and Differentiability Properties of Functions*, Princeton Mathematical Series, vol. 30. Princeton University Press, Princeton, N.J. (1970)
17. Takanen, J (2021) Dimension estimate for two-sided points of planar Sobolev extension domains, *Proc. Amer. Math. Soc.* (to appear)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.