

JYX



This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Mohanta, Kaushik; Tyagi, Jagmohan

Title: Improved hardy inequalities on Riemannian manifolds

Year: 2023

Version: Published version


Copyright: © 2023 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis

Rights: CC BY 4.0

Rights url: <https://creativecommons.org/licenses/by/4.0/>

Please cite the original version:

Mohanta, K., & Tyagi, J. (2023). Improved hardy inequalities on Riemannian manifolds. *Complex Variables and Elliptic Equations*, Early online. <https://doi.org/10.1080/17476933.2023.2247998>



Complex Variables and Elliptic Equations

An International Journal

ISSN: (Print) (Online) Journal homepage: <https://www.tandfonline.com/loi/gcov20>

Improved hardy inequalities on Riemannian manifolds

Kaushik Mohanta & Jagmohan Tyagi

To cite this article: Kaushik Mohanta & Jagmohan Tyagi (2023): Improved hardy inequalities on Riemannian manifolds, *Complex Variables and Elliptic Equations*, DOI: 10.1080/17476933.2023.2247998

To link to this article: <https://doi.org/10.1080/17476933.2023.2247998>



© 2023 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



Published online: 21 Aug 2023.



Submit your article to this journal [↗](#)



View related articles [↗](#)



View Crossmark data [↗](#)

Improved hardy inequalities on Riemannian manifolds

Kaushik Mohanta^a and Jagmohan Tyagi^b

^aDepartment of Mathematics and Statistics, University of Jyväskylä, Jyväskylä, Finland; ^bDiscipline of Mathematics, Indian Institute of Technology Gandhinagar, Gandhinagar, Gujarat, India

ABSTRACT

We study the following version of Hardy-type inequality on a domain Ω in a Riemannian manifold (M, g) :

$$\int_{\Omega} |\nabla u|_g^p \rho^\alpha dV_g \geq \left(\frac{|p-1+\beta|}{p} \right)^p \int_{\Omega} \frac{|u|^p |\nabla \rho|_g^p}{|\rho|^p} \rho^\alpha dV_g + \int_{\Omega} V |u|^p \rho^\alpha dV_g, \quad \forall u \in C_c^\infty(\Omega).$$

We provide sufficient conditions on p, α, β, ρ and V for which the above inequality holds. This generalizes earlier well-known works on Hardy inequalities on Riemannian manifolds. The functional setup covers a wide variety of particular cases, which are discussed briefly: for example, \mathbb{R}^N with $p < N$, $\mathbb{R}^N \setminus \{0\}$ with $p \geq N$, \mathbb{H}^N , etc.

ARTICLE HISTORY

Received 2 May 2023
Accepted 11 August 2023

COMMUNICATED BY

H. Boas

KEYWORDS

Hardy inequality; manifold; reminder term; critical case

AMS SUBJECT

CLASSIFICATIONS

58J05; 35A23; 46E35


1. Introduction

The Hardy inequality plays an important role in analysis and in the theory of partial differential equations. In \mathbb{R}^N , it reads as follows:

$$\int_{\mathbb{R}^N} \frac{|u|^p}{|x|^p} \leq c \int_{\mathbb{R}^N} |\nabla u|^p, \quad \forall u \in C_c^\infty(\mathbb{R}^N), \quad (1)$$

where the constant c is independent of u . When we restrict u to be in the space $W_0^{1,p}(\mathbb{R}^N)$, with $p < N$, or in $W_0^{1,p}(\mathbb{R}^N \setminus \{0\})$, with $p \geq N$, (1) holds; in that case the smallest possible choice for c becomes $\left(\frac{p}{N-p}\right)^p$, and this constant is never achieved (see for example [1,2]). Hence there is a scope to get an improvement in the above inequality. Brezis-Vázquez [2] have shown for the case $p=2$, that we can add an L^2 -term in the left hand side of (1) even with the best constant. This was further generalized in many directions, for example, see [1,3–13].

There are many results in literature on this subject in the context of a complete Riemannian manifolds (M, g) . An important result in this direction is due to Kombe-Ozaydin [14]. They proved that for a nonnegative function ρ with $|\nabla \rho|_g = 1$ and $\Delta \rho \geq \frac{C}{\rho}$, the following

CONTACT Kaushik Mohanta  kaushik.k.mohanta@jyu.fi
Kaushik Mohanta and Jagmohan Tyagi have contributed equally in this article.

holds

$$\left(\frac{C+1+\alpha-p}{p}\right)^p \int_M \frac{|u|^p \rho^\alpha}{\rho^p} \leq \int_M |\nabla u|_g^p \rho^\alpha \quad \text{for any } u \in C_c^\infty(M).$$

Moreover, for a bounded domain Ω with smooth boundary and in the case $p=2$, they also proved that the above inequality still holds for any u in $C_c^\infty(\Omega)$, if we add a reminder term of the form $C_1(\int_\Omega |\nabla u|^q \rho^{q\alpha/2})^{2/q}$ in the left hand side of the equation, where $1 < q < 2$ and $C = C(N, q, \Omega)$. D’Ambrosio-Dipierro [15] proved another version of Hardy inequality, where the restriction $|\nabla \rho| = 1$ is not there. They showed that if Ω is an open set in M and there is a $\rho : \Omega \rightarrow [0, \infty)$ such that $\rho \in W_{loc}^{1,p}(\Omega)$ with $\Delta_p \rho \leq 0$ weakly, then $\frac{|\nabla \rho|}{\rho} \in L_{loc}^p$, and for any $u \in C_c^\infty(\Omega)$, the following holds:

$$\left(\frac{p-1}{p}\right)^p \int_\Omega \frac{|u|^p |\nabla \rho|_g^p}{\rho^p} \leq \int_\Omega |\nabla u|_g^p.$$

They discussed, in details, the advantage and applications of this kind of Hardy inequality. For other related results, the reader may refer to [16–21] and the references therein. As of now, there is no known result regarding Hardy inequality with a reminder term as prescribed by D’Ambrosio-Dipierro [15]. We wish to address this problem in this paper by proving a slightly more general version of the result.

An interesting feature of our method is that the results of [14,15] follow immediately. Also the proof becomes much simpler (see Theorem 1.1) provided no reminder term is expected. The formulation allows greater control over the constant, and, at least in the Euclidean case, we can get the inequality with the best constants. Here we mainly focus on the improved Hardy inequality in all its generality. We refrain ourselves from studying the special cases exclusively, although that is very much possible to use our results to achieve improvement of the inequalities obtained in [15].

Throughout the article, (M, g) stands for a fixed oriented Riemannian manifold. We shall often use the notation $\langle X, Y \rangle$ to denote $g(X, Y)$ for any two vector fields X and Y . We use the symbol dV_g to denote the volume form, however, the symbol will be often dropped when there is no scope for confusion.

Now, we state the main results of this paper, which we shall prove in Section 2. The first one is the following Hardy inequality, which, in the particular case $\alpha = -\beta$, is proven in [15]. However, our proof is much more simpler.

Theorem 1.1: *Let $1 < p < \infty$, $\alpha, \beta \in \mathbb{R}$, M be a complete Riemannian manifold, Ω be a domain in M with boundary $\partial\Omega$ (possibly empty). Let there exist a function $\rho : \Omega \rightarrow (0, \infty)$ with $|\nabla \rho|_g \rho^{\frac{-p+\alpha}{p}} \in L_{loc}^p(\Omega)$ such that*

$$\frac{1}{p-1+\beta} \int_\Omega \langle \nabla \xi, |\nabla \rho|^{p-2} \nabla \rho \rangle \rho^{-p+1+\alpha} \geq \int_\Omega \frac{|\nabla \rho|_g^p}{\rho^{p-\alpha}} \xi, \quad \forall \xi \in C_c^1(\Omega) \quad \text{with } \xi \geq 0. \tag{2}$$

Then, for any $u \in W_0^{1,p}(\Omega)$, we have the following inequality:

$$\int_\Omega |\nabla u|_g^p \rho^\alpha dV_g \geq \left(\frac{p-1+\beta}{p}\right)^p \int_\Omega \frac{|u|^p |\nabla \rho|_g^p}{\rho^p} \rho^\alpha dV_g.$$

Theorem 1.2: Let $p, \alpha, \beta, M, \Omega$ and ρ be as in Theorem 1.1. Let the constant $C = C(p)$ be as in Lemma 2.2 for $p \geq 2$ and in the case $p < 2$, $C(p) := 2^{p-3}p(p-1)$. Further, in the case $p < 2$, assume that for any $\xi \in C_c^1(\Omega)$ with $\xi \geq 0$

$$\begin{cases} \rho^{\frac{-p+1-\beta}{p}} \in W_{loc}^{1,p}(\Omega), \\ (p-1+\beta) \int_{\Omega} \langle \nabla \xi, |\nabla \rho|^{p-2} \nabla \rho \rangle \rho^{\alpha+\beta} \geq \int_{\Omega} \frac{|\nabla \rho|^p}{\rho} \rho^{\alpha+\beta} \xi. \end{cases} \quad (3)$$

If there exist functions $V : \Omega \rightarrow [0, \infty)$ and $\varphi \in W_{loc}^{1,p}(\Omega)$ such that for any $\xi \in C_c^1(\Omega)$,

$$\begin{aligned} & \int_{\Omega} (|\nabla \varphi|^{p-2} \nabla \varphi, \nabla \xi) \varphi^{-p+1} \rho^{\alpha} - (p-1) \int_{\Omega} |\nabla \varphi|^p \xi \rho^{\alpha} \varphi^{-p} \\ & - (p-1+\beta) \int_{\Omega} (|\nabla \varphi|^{p-2} \nabla \varphi, \nabla \rho) \xi \varphi^{-p+1} \rho^{-1+\alpha} \\ & \geq \int_{\Omega} \frac{V}{C(p)} \xi \rho^{\alpha}, \end{aligned} \quad (4)$$

then for any $u \in W_0^{1,p}(\Omega)$, we have the following inequality:

$$\int_{\Omega} |\nabla u|_g^p \rho^{\alpha} dV_g \geq \left(\frac{|p-1+\beta|}{p} \right)^p \int_{\Omega} \frac{|u|^p |\nabla \rho|_g^p}{|\rho|^p} \rho^{\alpha} dV_g + \int_{\Omega} V |u|^p \rho^{\alpha} dV_g. \quad (5)$$

Before moving further, let us observe that the hypotheses (2), (3), (4) can be thought of as weak formulations of certain problems; we explain this in the following using the Green's theorem (see Lemma 2.1). Consider the following condition on ρ :

$$\frac{-1}{p-1+\beta} \Delta_p \rho \geq \frac{\alpha+\beta}{p-1+\beta} \frac{|\nabla \rho|^p}{\rho}. \quad (6)$$

In the special case $\alpha = -\beta$, this is just the p -superharmonicity condition. We rewrite this as

$$\frac{-1}{p-1+\beta} \rho^{-p+1+\alpha} \Delta_p \rho + \frac{p-1-\alpha}{p-1+\beta} \rho^{-p+\alpha} |\nabla \rho|^p \geq \frac{|\nabla \rho|^p}{\rho^{p-\alpha}},$$

and then multiply both sides by a test function $\xi \in C_c^1(\Omega)$, integrate over Ω , and then apply the product rule of divergence operator to get

$$\frac{-1}{p-1+\beta} \int_{\Omega} \operatorname{div} (\rho^{-p+1+\alpha} |\nabla \rho|^{p-2} \nabla \rho) \xi \geq \int_{\Omega} \frac{|\nabla \rho|^p}{\rho^{p-\alpha}} \xi.$$

An application of Green's theorem then gives (2). Thus (2) is an weak formulation of (6). Similarly, the second condition of (3) can be interpreted as the weak formulation of

$$\frac{-1}{p-1+\beta} \Delta_p \rho \geq \frac{p-1+\alpha+2\beta}{p-1+\beta} \frac{|\nabla \rho|^p}{\rho},$$

and (4) can be regarded as a weak formulation of

$$\Delta_p \varphi + (p-1+\alpha+\beta) \left\langle |\nabla \varphi|^{p-2} \nabla \varphi, \frac{\nabla \rho}{\rho} \right\rangle + \frac{V}{C(p)} \varphi^{p-1} \leq 0.$$

Now, we discuss some immediate consequences of Theorem 1.2. The following result says that if $\left(\frac{|p-1+\beta|}{p}\right)^p$ is the best constant in (5), then (4) has no solution when $V = \frac{|\nabla\rho|^p}{\rho^p}$.

Corollary 1.3: *Let p, α, β, ρ be as in Theorem 1.2, and $V = C\frac{|\nabla\rho|}{\rho}$ be such that there exists some φ for which (4) is satisfied. Then the constant $\left(\frac{|p-1+\beta|}{p}\right)^p$ in (5) is not sharp.*

Remark 1.4: The above result says nothing about when the constant $\left(\frac{|p-1+\beta|}{p}\right)^p$ in (5) is sharp. However it is easy to see that the constant is sharp if and only if we can never have $V = C\frac{|\nabla\rho|}{\rho}$ in (5) for any $C > 0$. So the question of whether the constant is sharp is related to necessity of (4) in (5); whereas result Theorem 1.2 concerns with the sufficiency part. In some very particular case the condition may also be necessary as can be seen from [22, Theorem 1].

We discuss some special cases of Theorem 1.2. In the Euclidean setup, we get the following results as a corollary. Let us take $M = \mathbb{R}^N$, $\rho = |x|^d$. Then $\nabla\rho = d|x|^{d-2}x$ and we get the following corollary:

Corollary 1.5: *Assume $1 < p < \infty$, $d \neq 0$, $\alpha, \beta \in \mathbb{R}$ be such that $p - 1 + \beta \neq 0$,*

$$\frac{\alpha + \beta}{p - 1 + \beta} \leq 0 \quad \text{and} \quad \frac{(N + (d - 1)(p - 1))}{d(p - 1 + \beta)} \leq 0,$$

and

$$\frac{(N + (d - 1)(p - 1))}{d(p - 1 + \beta)} \leq -1 \quad \text{when } p < 2.$$

Assume further that there exists a function $V : \mathbb{R}^N \rightarrow \mathbb{R}$ such that the following problem admits a weak solution $\varphi \in W_{loc}^{1,p}(\Omega)$:

$$\Delta_p\varphi + \frac{d(p - 1 + \alpha + \beta)}{|x|^2}|\nabla\varphi|^{p-2}\nabla\varphi \cdot x + \frac{V}{C(p)}\varphi^{p-1} \leq 0. \tag{7}$$

Consider the two cases:

(i) $\Omega \subseteq \mathbb{R}^N$, $p < N + \alpha d$ with

$$(p - 1 + \beta)d < N - p \quad \text{when } p < 2,$$

and

(ii) $\Omega \subseteq \mathbb{R}^N \setminus \{0\}$, $p \geq N + \alpha d$.

Then, in both cases, we have the following inequality:

$$\begin{aligned} & \int_{\Omega} |\nabla u|_g^p |x|^{\alpha d} \, dx \\ & \geq \left(\frac{|d(p - 1 + \beta)|}{p}\right)^p \int_{\Omega} \frac{|u|^p}{|x|^{p-\alpha d}} \, dx + \int_{\Omega} V|u|^p |x|^{\alpha d}, \quad \forall u \in W_0^{1,p}(\Omega). \end{aligned}$$

Remark that in the case $\alpha = \beta = 0, d = -1$ and $\Omega = \mathbb{R}^N$, [23, Proposition 1.2] implies that (7) has no solution in \mathbb{R}^N for any $1 < p < N$.

Now, let us consider the so-called critical case: $p = N \geq 2$. Set $\Omega := B_1(0) \subset \mathbb{R}^N, \rho = -\log|x|$. We have

Corollary 1.6: *Let $N \geq 2, B_1(0)$ denote the unit ball in $\mathbb{R}^N, \alpha, \beta \in \mathbb{R}$ be such that $p - 1 + \beta \neq 0$ and*

$$\frac{\alpha + \beta}{N - 1 + \beta} \leq 0.$$

Assume further that there exists a function $V : B_1(0) \subset \mathbb{R}^N \rightarrow \mathbb{R}$ such that the following problem admits a weak solution $\varphi \in W_{loc}^{1,p}(B_1(0))$:

$$\Delta_N \varphi - \frac{(N - 1 + \alpha + \beta)|\nabla \varphi|^{N-2}}{|x|^2 \log x} \nabla \varphi \cdot x + \frac{V}{C(N)} \leq 0.$$

Then for any $u \in W_0^{1,N}(B_1(0))$, we have

$$\begin{aligned} \int_{B_1(0)} |\nabla u|^N |\log|x||^\alpha dx &\geq \left(\frac{|N - 1 + \beta|}{N} \right)^N \int_{B_1(0)} \frac{|u|^N}{|x|^N |\log|x||^{N-\alpha}} dx \\ &+ \int_{B_1(0)} V |u|^N |\log|x||^\alpha dx. \end{aligned}$$

Consider the case $M = \mathbb{H}^N := \mathbb{R}^{N-1} \times (0, \infty)$ and set $\rho(x) := x_N$. This gives $|\nabla \rho| = 1$ and $\Delta_p \rho = 0$. This implies

Corollary 1.7: *Let $1 < p < \infty, \alpha, \beta \in \mathbb{R}$. Let the constant $C = C(p)$ be as in Theorem 1.2. Assume*

$$(\alpha + \beta)(p - 1 + \beta) \leq 0.$$

Further, in the case $p < 2$, assume that

$$(p - 1 + \alpha + 2\beta)(p - 1 + \beta) \leq 0.$$

If there exists a positive p -harmonic function $\varphi \in W_{loc}^{1,p}(\mathbb{H}^N)$ such that $(p - 1 + \alpha + \beta) \frac{\partial \varphi}{\partial x_N} \leq 0$, then for any $u \in W_0^{1,p}(\mathbb{H}^N)$, we have the following inequality:

$$\begin{aligned} \int_{\mathbb{H}^N} |\nabla u|_g^p x_N^\alpha dx &\geq \left(\frac{|p - 1 + \beta|}{p} \right)^p \int_{\mathbb{H}^N} \frac{|u|^p}{x_N^{p-\alpha}} dx \\ &- C(p)(p - 1 + \alpha + \beta) \int_{\mathbb{H}^N} \frac{\partial \varphi}{\partial x_N} \frac{|\nabla \varphi|^{p-2}}{\varphi^{p-1}} x_N^{\alpha+1} |u|^p dx. \end{aligned}$$

Remark 1.8: The conditions (2) in Theorem 1.1 and (4) in Theorem 1.2 may seem artificially imposed at first glance. However, in the most commonly used form of Hardy inequality, that is in the setup of Corollary 1.5 with $\alpha = \beta = 0$, (2) holds automatically with

proper choice of d . In case of Riemannian manifolds validity of Equation (2) is not immediate, it is related to p -hyperbolicity of the underlying manifold. A discussion regarding this can be found in [15].

In the particular case $p=2$, $\Omega = \mathbb{R}^N$, and when V is radial, (4) is actually a necessary condition too for Equation (5) to hold. This can be seen from some symmetrization argument (to reduce the condition to its one-dimensional analogue) and [22, Theorem 1].

In Section 2, we shall prove some preliminary results, Theorem 1.1 followed by the proof of Theorem 1.2.

2. Proof of the theorems

The proof of the following lemma can be found in [24, Theorem III.7.6.] for the case $p=2$. The proof of this version can also be done similarly as the essence of the proof lies in the Stokes theorem and in the product rule of divergence operator:

$$\operatorname{div}(fX) = f \operatorname{div}(X) + \langle \nabla f, X \rangle.$$

Lemma 2.1 (Green's Formula): *Let $p > 1$, M be complete, oriented Riemannian manifold, Ω a domain in M with smooth boundary. Let $f \in C^2(\Omega)$, $\xi \in C_c^1(\Omega)$. Then, we have*

$$\int_{\Omega} \langle |\nabla f|^{p-2} \nabla f, \nabla \xi \rangle = - \int_{\Omega} \xi \Delta_p f.$$

The following result plays a key role in the proof of the theorem.

Lemma 2.2: *Let $x \in M$ and $X_x, Y_x \in T_x M$ be two tangent vectors. Then, for $p \geq 2$, there is a constant $C = C(p) > 0$ such that*

$$|X_x + Y_x|_g^p - |X_x|_g^p \geq C(p) |Y_x|_g^p + p |X_x|_g^{p-2} \langle X_x, Y_x \rangle,$$

and for $1 \leq p \leq 2$,

$$|X_x + Y_x|_g^p - |X_x|_g^p \geq \frac{p(p-1)}{2} \frac{|Y_x|^2}{\left(|X_x| + |Y_x|\right)^{2-p}} + p |X_x|^{p-2} \langle X_x, Y_x \rangle.$$

In general, for $1 < p < \infty$, we have

$$|X_x + Y_x|_g^p - |X_x|_g^p \geq p |X_x|^{p-2} \langle X_x, Y_x \rangle.$$

Proof: For the case $p \geq 2$, refer to [25, Chapter 12] and recall that any N -dimensional Hilbert space is isomorphic to \mathbb{R}^N . We give a proof of the case $1 \leq p \leq 2$, which is adapted from [26, Lemma 1].

In the following calculation, we omit the point x for better readability. Consider the function

$$f(t) := |X + tY|^p.$$

It can be easily verified that, for $t \in (0, 1)$, $f''(t) \geq p(p-1)|Y|^2|X + tY|^{p-2}$. Using this, Taylor's theorem, the fact that $p-2 \leq 0$, we get

$$|X + Y|^p = f(1)$$

$$\begin{aligned}
 &= f(0) + f'(0) + \int_0^1 (1-t)f''(t)dt \\
 &\geq |X|^p + p|X|^{p-2} \langle X, Y \rangle \\
 &\quad + p(p-1) \int_0^1 (1-t)|Y|^2|X| + t|Y|^{p-2} dt \\
 &\geq |X|^p + p|X|^{p-2} \langle X, Y \rangle + \frac{p(p-1)}{2}|Y|^2(|X| + |Y|)^{p-2}.
 \end{aligned}$$

This proves the lemma. ■

We now present the proof of our first main result.

Proof of Theorem 1.1: By a density argument, we can always assume that $u \in C_c^1(\Omega)$. Set $w(x) = u(x)\rho^{\frac{-p+1-\beta}{p}}(x)$ in Ω . Then

$$\nabla w = \rho^{\frac{p-1+\beta}{p}} \nabla w + \frac{p-1+\beta}{p} \rho^{\frac{-1+\beta}{p}} w \nabla \rho.$$

Note that if $\int_{\Omega} |\nabla u(x)|_g^p \rho^\alpha dV_g = \infty$, then we have nothing to prove, so we assume it to be finite. In the following calculation, we need the term $\frac{|u|^p |\nabla \rho|_g^p}{\rho^{p-\alpha}}$ to be integrable over Ω ; this is indeed true, as $u \in C_c^1(\Omega)$ and $\frac{|\nabla \rho|_g^p}{\rho^{p-\alpha}} \in L_{loc}^1(\Omega)$ according to hypothesis. Using Lemma 2.2 and (2), we have

$$\begin{aligned}
 &\int_{\Omega} |\nabla u(x)|_g^p \rho^\alpha dV_g - \left(\frac{|p-1+\beta|}{p} \right)^p \int_{\Omega} \frac{|u(x)|^p |\nabla \rho(x)|_g^p}{|\rho(x)|^p} \rho^\alpha dV_g \\
 &= \int_{\Omega} \left(\left| \rho^{\frac{p-1+\beta}{p}} \nabla w + \frac{p-1+\beta}{p} \frac{w \nabla \rho}{\rho^{\frac{1-\beta}{p}}} \right|_g^p - \left(\frac{|p-1+\beta|}{p} \right)^p \left| \frac{w \nabla \rho}{\rho^{\frac{1-\beta}{p}}} \right|_g^p \right) \rho^\alpha \\
 &\geq p \left(\frac{|p-1+\beta|}{p} \right)^{p-2} \left(\frac{p-1+\beta}{p} \right) \int_{\Omega} \left| \frac{w \nabla \rho}{\rho^{\frac{1-\beta}{p}}} \right|^{p-2} \left\langle \rho^{\frac{p-1+\beta}{p}} \nabla w, \frac{w \nabla \rho}{\rho^{\frac{1-\beta}{p}}} \right\rangle \rho^\alpha \\
 &= \left(\frac{|p-1+\beta|}{p} \right)^{p-2} \left(\frac{p-1+\beta}{p} \right) \int_{\Omega} \langle \nabla |w|^p, |\nabla \rho|^{p-2} \nabla \rho \rangle \rho^{\alpha+\beta} \\
 &= \left(\frac{|p-1+\beta|}{p} \right)^{p-2} \left(\frac{p-1+\beta}{p} \right) \int_{\Omega} \langle \nabla (\rho^{-p+1-\beta} |u|^p), |\nabla \rho|^{p-2} \nabla \rho \rangle \rho^{\alpha+\beta} \\
 &= \left(\frac{|p-1+\beta|}{p} \right)^{p-2} \left(\frac{p-1+\beta}{p} \right) \int_{\Omega} \langle \nabla |u|^p, |\nabla \rho|^{p-2} \nabla \rho \rangle \rho^{-p+1+\alpha} \\
 &\quad - \left(\frac{|p-1+\beta|}{p} \right)^p \int_{\Omega} \frac{|\nabla \rho|_g^p}{\rho^{p-\alpha}} |u|^p \\
 &\geq 0.
 \end{aligned}$$

In the last line of the above calculation, we have used the hypothesis by using the fact that $|u|^p \in C_c^1(\Omega)$ as $p > 1$. ■

Next, we proceed with the proof of our second main result.

Proof of Theorem 1.2: As in the proof of Theorem 1.1, we shall assume $u \in C_c^1(\Omega)$ and we set $u(x) = w(x)\rho^{\frac{p-1+\beta}{p}}(x)$, so that we have $\nabla u = \rho^{\frac{p-1+\beta}{p}}\nabla w + \frac{p-1+\beta}{p}\rho^{\frac{-1+\beta}{p}}w\nabla\rho$. We need to estimate the term

$$\begin{aligned} I &:= \int_{\Omega} |\nabla u(x)|_g^p \rho^\alpha dV_g \\ &\quad - \left(\frac{|p-1+\beta|}{p}\right)^p \int_{\Omega} \frac{|u(x)|^p |\nabla\rho(x)|_g^p}{|\rho(x)|^p} \rho^\alpha dV_g \\ &= \int_{\Omega} \left(\left| \rho^{\frac{p-1+\beta}{p}}\nabla w + \left(\frac{p-1+\beta}{p}\right) \frac{w\nabla\rho}{\rho^{\frac{1-\beta}{p}}} \right|_g^p \right. \\ &\quad \left. - \left(\frac{|p-1+\beta|}{p}\right)^p \left| \frac{w\nabla\rho}{\rho^{\frac{1-\beta}{p}}} \right|_g^p \right) \rho^\alpha. \end{aligned}$$

Let $p \geq 2$. Using Lemma 2.2 and (2), we get

$$\begin{aligned} I &\geq C(p) \int_{\Omega} \left| \rho^{\frac{p-1+\beta}{p}}\nabla w \right|^p \rho^\alpha + p \left(\frac{|p-1+\beta|}{p}\right)^{p-2} \left(\frac{p-1+\beta}{p}\right) \\ &\quad \times \int_{\Omega} \left| \frac{w\nabla\rho}{\rho^{\frac{1-\beta}{p}}} \right|^{p-2} \left\langle \rho^{\frac{p-1+\beta}{p}}\nabla w, \frac{w\nabla\rho}{\rho^{\frac{1-\beta}{p}}} \right\rangle \rho^\alpha \\ &= C(p) \int_{\Omega} |\nabla w|^p \rho^{p-1+\alpha+\beta} + \left(\frac{|p-1+\beta|}{p}\right)^{p-2} \left(\frac{p-1+\beta}{p}\right) \\ &\quad \times \int_{\Omega} \langle \nabla|w|^p, |\nabla\rho|^{p-2}\nabla\rho \rangle \rho^{\alpha+\beta} \\ &\geq C(p) \int_{\Omega} |\nabla w|^p \rho^{p-1+\alpha+\beta}. \end{aligned} \tag{8}$$

The last line in the above calculation follows from (2). Now, set $\psi = \frac{w}{\varphi}$. Then $\nabla w = \varphi\nabla\psi + \psi\nabla\varphi$. We have, from Lemma 2.2 and (4),

$$\begin{aligned} &\int_{\Omega} \rho^{p-1+\alpha+\beta} |\nabla w|_g^p \\ &= \int_{\Omega} \rho^{p-1+\alpha+\beta} |\varphi\nabla\psi + \psi\nabla\varphi|_g^p \\ &\geq C(p) \int_{\Omega} \rho^{p-1+\alpha+\beta} |\varphi\nabla\psi|_g^p \end{aligned}$$

$$\begin{aligned}
 & + \int_{\Omega} p \rho^{p-1+\alpha+\beta} |\psi \nabla \varphi|^{p-2} < \psi \nabla \varphi, \varphi \nabla \psi > + \int_{\Omega} \rho^{p-1+\alpha+\beta} |\psi \nabla \varphi|_g^p \\
 \geq & \int_{\Omega} \langle \rho^{p-1+\alpha+\beta} \varphi |\nabla \varphi|^{p-2} \nabla \varphi, \nabla |\psi|^p \rangle + \int_{\Omega} \rho^{p-1+\alpha+\beta} |\psi \nabla \varphi|_g^p \\
 = & \int_{\Omega} \langle |\nabla \varphi|^{p-2} \nabla \varphi, \nabla (\varphi |\psi|^p) \rangle \rho^{p-1+\alpha+\beta} \\
 = & \int_{\Omega} \langle |\nabla \varphi|^{p-2} \nabla \varphi, \nabla (|u|^p \varphi^{-p+1} \rho^{-p+1-\beta}) \rangle \rho^{p-1+\alpha+\beta} \\
 = & \int_{\Omega} \langle |\nabla \varphi|^{p-2} \nabla \varphi, \nabla (|u|^p) \rangle \varphi^{-p+1} \rho^{\alpha} - (p-1) \int_{\Omega} |\nabla \varphi|^p |u|^p \rho^{\alpha} \varphi^{-p} \\
 & - (p-1+\beta) \int_{\Omega} \langle |\nabla \varphi|^{p-2} \nabla \varphi, \nabla \rho \rangle |u|^p \varphi^{-p+1} \rho^{-1+\alpha} \\
 \geq & \int_{\Omega} \frac{V}{C(p)} |u|^p \rho^{\alpha}. \tag{9}
 \end{aligned}$$

Combining (8) and (9), we get the desired result.

Now let us consider the case $p < 2$. Using Lemma 2.2 and (2), we get

$$\begin{aligned}
 I \geq & \int_{\Omega} \frac{p(p-1)}{2} \frac{\left| \rho^{\frac{p-1+\beta}{p}} \nabla w \right|^2 \rho^{\alpha}}{\left| \left| \frac{p-1+\beta}{p} \frac{w \nabla \rho}{\rho^{\frac{1-\beta}{p}}} \right| + \left| \rho^{\frac{p-1+\beta}{p}} \nabla w \right| \right|^{2-p}} \\
 & + p \left| \frac{p-1+\beta}{p} \frac{w \nabla \rho}{\rho^{\frac{1-\beta}{p}}} \right|^{p-2} \left\langle \frac{p-1+\beta}{p} \frac{w \nabla \rho}{\rho^{\frac{1-\beta}{p}}}, \rho^{\frac{p-1+\beta}{p}} \nabla w \right\rangle \rho^{\alpha} \\
 = & \int_{\Omega} \frac{p(p-1)}{2} \frac{\rho^{1+\alpha+\beta} |\nabla w|^2}{\left| \left| \frac{p-1+\beta}{p} w \nabla \rho \right| + |\rho \nabla w| \right|^{2-p}} \\
 & + \left(\frac{|p-1+\beta|}{p} \right)^{p-2} \left(\frac{p-1+\beta}{p} \right) \int_{\Omega} \langle |\nabla \rho|^{p-2} \nabla \rho, \nabla |w|^p \rangle \rho^{\alpha+\beta} \\
 \geq & \frac{p^{3-p}(p-1)}{2} \int_{\Omega} \frac{\rho^{p-1+\alpha+\beta} |\nabla w|^2}{\left| |(p-1+\beta)w\rho^{-1}\nabla\rho| + p|\nabla w| \right|^{2-p}}. \tag{10}
 \end{aligned}$$

Now, we estimate the last term using Hölder's inequality, convexity of $t \mapsto t^p$ in the following:

$$\begin{aligned}
 & \int_{\Omega} \rho^{p-1+\alpha+\beta} |\nabla w|^p \\
 = & \int_{\Omega} \left(\frac{\rho^{p-1+\alpha+\beta} |\nabla w|^2}{\left| |(p-1+\beta)w\rho^{-1}\nabla\rho| + p|\nabla w| \right|^{2-p}} \right)^{\frac{p}{2}}
 \end{aligned}$$

$$\begin{aligned}
& \times \left(\rho^{\frac{(p-1+\alpha+\beta)(2-p)}{2}} \left| |(p-1+\beta)w\rho^{-1}\nabla\rho| + p|\nabla w| \right|^{\frac{2p-p^2}{2}} \right) \\
& \leq \left(\int_{\Omega} \left(\frac{\rho^{p-1+\alpha+\beta}|\nabla w|^2}{\left| |(p-1+\beta)w\rho^{-1}\nabla\rho| + p|\nabla w| \right|^{2-p}} \right) \right)^{\frac{p}{2}} \\
& \times \left(\int_{\Omega} \left(\rho^{\frac{(p-1+\alpha+\beta)(2-p)}{2}} \left| |(p-1+\beta)w\rho^{-1}\nabla\rho| + p|\nabla w| \right|^{\frac{2p-p^2}{2}} \right)^{\frac{2}{2-p}} \right)^{\frac{2-p}{2}} \\
& = \left(\int_{\Omega} \frac{\rho^{p-1+\alpha+\beta}|\nabla w|^2}{\left| |(p-1+\beta)w\rho^{-1}\nabla\rho| + p|\nabla w| \right|^{2-p}} \right)^{\frac{p}{2}} \\
& \times \left(\int_{\Omega} \rho^{p-1+\alpha+\beta} \left| |(p-1+\beta)w\rho^{-1}\nabla\rho| + p|\nabla w| \right|^p \right)^{\frac{2-p}{2}} \\
& \leq \left(\int_{\Omega} \frac{\rho^{p-1+\alpha+\beta}|\nabla w|^2}{\left| |(p-1+\beta)w\rho^{-1}\nabla\rho| + p|\nabla w| \right|^{2-p}} \right)^{\frac{p}{2}} \\
& \times \left(2^{p-1}|p-1+\beta|^p \int_{\Omega} \rho^{p-1+\alpha+\beta} |w\rho^{-1}\nabla\rho|^p \right. \\
& \left. + p^p 2^{p-1} \int_{\Omega} \rho^{p-1+\alpha+\beta} |\nabla w|^p \right)^{\frac{2-p}{2}}. \tag{11}
\end{aligned}$$

Observing that since we have assumed (3), we can apply Theorem 1.1 to get

$$\begin{aligned}
\int_{\Omega} \rho^{p-1+\alpha+\beta} |\nabla w|^p & \leq \left(\int_{\Omega} \frac{\rho^{p-1+\alpha+\beta} |\nabla w|^2}{\left| |(p-1+\beta)w\rho^{-1}\nabla\rho| + p|\nabla w| \right|^{2-p}} \right)^{\frac{p}{2}} \\
& \times \left(p^p 2^p \int_{\Omega} \rho^{p-1+\alpha+\beta} |\nabla w|^p \right)^{\frac{2-p}{2}}.
\end{aligned}$$

After setting $\psi = \frac{w}{\varphi}$, so that $\nabla w = \varphi \nabla \psi + \psi \nabla \varphi$ and then applying Lemma 2.2 and (4), we get

$$\begin{aligned}
& \frac{p^3-p(p-1)}{2} \int_{\Omega} \frac{\rho^{p-1+\alpha+\beta} |\nabla w|^2}{\left| |(p-1+\beta)w\rho^{-1}\nabla\rho| + p|\nabla w| \right|^{2-p}} \\
& \geq 2^{p-3} p(p-1) \int_{\Omega} \rho^{p-1+\alpha+\beta} |\nabla w|^p \\
& = 2^{p-3} p(p-1) \int_{\Omega} \rho^{p-1+\alpha+\beta} |\varphi \nabla \psi + \psi \nabla \varphi|^p \\
& \geq 2^{p-3} p^2(p-1) \int_{\Omega} \rho^{p-1+\alpha+\beta} |\psi \nabla \varphi|^{p-2} \langle \psi \nabla \varphi, \varphi \nabla \psi \rangle
\end{aligned}$$

$$\begin{aligned}
 &+ 2^{p-3}p(p-1) \int_{\Omega} \rho^{p-1+\alpha+\beta} |\psi \nabla \varphi|^p \\
 &= 2^{p-3}p(p-1) \int_{\Omega} \langle \rho^{p-1+\alpha+\beta} \varphi |\nabla \varphi|^{p-2} \nabla \varphi, \nabla |\psi|^p \rangle \\
 &\quad + 2^{p-3}p(p-1) \int_{\Omega} \rho^{p-1+\alpha+\beta} |\psi \nabla \varphi|^p \\
 &= 2^{p-3}p(p-1) \int_{\Omega} \langle |\nabla \varphi|^{p-2} \nabla \varphi, \nabla (\varphi |\psi|^p) \rangle \rho^{p-1+\alpha+\beta} \\
 &\geq \int_{\Omega} V |u|^p \rho^{\alpha}.
 \end{aligned}$$

The last line follows following the same equalities as in (9). This with (10) proves the theorem. ■

Acknowledgments

Authors thank the anonymous referee for the valuable suggestions and corrections.

Disclosure statement

No potential conflict of interest was reported by the author(s).

Funding

Authors thank IIT Gandhinagar for the financial support under the grant MIS/IITGN/R&D/MA/JT/202122/069.

References

- [1] Adimurthi, Chaudhuri N, Ramaswamy M. An improved hardy-Sobolev inequality and its application. *Proc Amer Math Soc.* 2002;130(2):489–505. doi: [10.1090/proc/2002-130-02](https://doi.org/10.1090/proc/2002-130-02)
- [2] Brezis H, Vázquez J. Blow-up solutions of some nonlinear elliptic problems. *Rev Mat Univ Complut Madrid.* 1997;10(2):443–469.
- [3] Abdellaoui B, Colorado E, Peral I. Some improved Caffarelli-Kohn-Nirenberg inequalities. *Calc Var Partial Differ Equ.* 2005;23(3):327–345. doi: [10.1007/s00526-004-0303-8](https://doi.org/10.1007/s00526-004-0303-8)
- [4] Adimurthi, Esteban M. An improved Hardy-Sobolev inequality in $W^{1,p}$ and its application to Schrödinger operators. *NoDEA Nonlinear Differ Equ Appl.* 2005;12(2):243–263. doi: [10.1007/s00030-005-0009-4](https://doi.org/10.1007/s00030-005-0009-4)
- [5] Adimurthi, Filippas S, Tertikas A. On the best constant of Hardy-Sobolev inequalities. *Nonlinear Anal.* 2009;70(8):2826–2833. doi: [10.1016/j.na.2008.12.019](https://doi.org/10.1016/j.na.2008.12.019)
- [6] Adimurthi, Sekar A. Role of the fundamental solution in Hardy-Sobolev-type inequalities. *Proc Roy Soc Edinb Sect A.* 2006;136(6):1111–1130. doi: [10.1017/S030821050000490X](https://doi.org/10.1017/S030821050000490X)
- [7] Barbatis G, Filippas S, Tertikas A. A unified approach to improved L^p Hardy inequalities with best constants. *Trans Amer Math Soc.* 2004;356(6):2169–2196. doi: [10.1090/tran/2004-356-06](https://doi.org/10.1090/tran/2004-356-06)
- [8] Chaudhuri N. Bounds for the best constant in an improved Hardy-Sobolev inequality. *Z Anal Anwend.* 2003;22(4):757–765. doi: [10.4171/ZAA](https://doi.org/10.4171/ZAA)
- [9] Cuomo S, Perrotta A. On best constants in Hardy inequalities with a remainder term. *Nonlinear Anal.* 2011;74(16):5784–5792. doi: [10.1016/j.na.2011.05.069](https://doi.org/10.1016/j.na.2011.05.069)
- [10] Duy N, Lam N, Phi L. Improved Hardy inequalities and weighted Hardy type inequalities with spherical derivatives. *Rev Mat Complut.* 2022;35(1):1–23. doi: [10.1007/s13163-020-00379-3](https://doi.org/10.1007/s13163-020-00379-3)

- [11] Duy N, Lam N, Triet N, et al. Improved Hardy inequalities with exact remainder terms. *Math Inequal Appl.* **2020**;23(4):1205–1226.
- [12] Nguyen V. Improved critical Hardy inequality and Leray-Trudinger type inequalities in Carnot groups. *Ann Fenn Math.* **2022**;47(1):121–138. doi: [10.54330/afm.112567](https://doi.org/10.54330/afm.112567)
- [13] Tertikas A, Zographopoulos NB. Best constants in the Hardy-Rellich inequalities and related improvements. *Adv Math.* **2007**;209(2):407–459. doi: [10.1016/j.aim.2006.05.011](https://doi.org/10.1016/j.aim.2006.05.011)
- [14] Kombe I, Özaydin M. Improved Hardy and Rellich inequalities on Riemannian manifolds. *Trans Amer Math Soc.* **2009**;361(12):6191–6203. doi: [10.1090/S0002-9947-09-04642-X](https://doi.org/10.1090/S0002-9947-09-04642-X)
- [15] D’Ambrosio L, Dipierro S. Hardy inequalities on Riemannian manifolds and applications. *Ann Inst H Poincaré C Anal Non Linéaire.* **2014**;31(3):449–475. doi: [10.4171/aihpc](https://doi.org/10.4171/aihpc)
- [16] Ahmetolan S, Kombe I. Improved Hardy and Rellich type inequalities with two weight functions. *Math Inequal Appl.* **2018**;21(3):885–896.
- [17] Devyver B, Fraas M, Pinchover Y. Optimal Hardy weight for second-order elliptic operator: an answer to a problem of agmon. *J Funct Anal.* **2014**;266(7):4422–4489. doi: [10.1016/j.jfa.2014.01.017](https://doi.org/10.1016/j.jfa.2014.01.017)
- [18] Kombe I, Yener A. Weighted Hardy and Rellich type inequalities on Riemannian manifolds. *Math Nachr.* **2016**;289(8-9):994–1004. doi: [10.1002/mana.201500237](https://doi.org/10.1002/mana.201500237)
- [19] Kristály A. Sharp uncertainty principles on Riemannian manifolds: the influence of curvature. *J Math Pures Appl (9).* **2018**;119:326–346. doi: [10.1016/j.matpur.2017.09.002](https://doi.org/10.1016/j.matpur.2017.09.002)
- [20] Meng C, Wang H, Zhao W. Hardy type inequalities on closed manifolds via ricci curvature. *Proc Roy Soc Edinb Sect A.* **2021**;151(3):993–1020. doi: [10.1017/prm.2020.47](https://doi.org/10.1017/prm.2020.47)
- [21] Thiam E. Weighted Hardy inequality on Riemannian manifolds. *Commun Contemp Math.* **2016**;18(06):1550072. doi: [10.1142/S021919971550072825](https://doi.org/10.1142/S021919971550072825).
- [22] Ghoussoub N, Moradifam A. On the best possible remaining term in the Hardy inequality. *Proc Natl Acad Sci USA.* **2008**;105(37):13746–13751. doi: [10.1073/pnas.0803703105](https://doi.org/10.1073/pnas.0803703105)
- [23] Cazacu C, Krejcirik D, Laptev A. Hardy inequalities for magnetic p -Laplacians, *arXiv preprint arXiv:2201.02482*, 2022.
- [24] Chavel I. *Riemannian geometry*. 2nd ed., Cambridge: Cambridge University Press; **2006**. (Cambridge Studies in Advanced Mathematics; vol. 98). A modern introduction.
- [25] Lindqvist P. Notes on the p -Laplace equation. Jyväskylä: University of Jyväskylä; **2006**. (Report. University of Jyväskylä Department of Mathematics and Statistics; vol. 102).
- [26] Gazzola F, Grunau H, Mitidieri E. Hardy inequalities with optimal constants and remainder terms. *Trans Amer Math Soc.* **2004**;356(6):2149–2168. doi: [10.1090/tran/2004-356-06](https://doi.org/10.1090/tran/2004-356-06)