

This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Rissanen, Antti J.; Jilbert, Tom; Simojoki, Asko; Mangayil, Rahul; Aalto, Sanni L.; Khanongnuch, Ramita; Peura, Sari; Jäntti, Helena

Title: Organic matter lability modifies the vertical structure of methane-related microbial communities in lake sediments

Year: 2023

Version: Published version

Copyright: © 2023 Rissanen et al.

Rights: CC BY 4.0

Rights url: <https://creativecommons.org/licenses/by/4.0/>

Please cite the original version:

Rissanen, A. J., Jilbert, T., Simojoki, A., Mangayil, R., Aalto, S. L., Khanongnuch, R., Peura, S., & Jäntti, H. (2023). Organic matter lability modifies the vertical structure of methane-related microbial communities in lake sediments. *Microbiology Spectrum*, 11(5), Article e01955-23. <https://doi.org/10.1128/spectrum.01955-23>

Organic matter lability modifies the vertical structure of methane-related microbial communities in lake sediments

Antti J. Rissanen,^{1,2} Tom Jilbert,³ Asko Simojoki,⁴ Rahul Mangayil,¹ Sanni L. Aalto,^{5,6} Ramita Khanongnuch,¹ Sari Peura,⁷ Helena Jantti⁵

AUTHOR AFFILIATIONS See affiliation list on p. 12.

ABSTRACT Eutrophication increases the input of labile, algae-derived, organic matter (OM) into lake sediments. This potentially increases methane (CH₄) emissions from sediment to water through increased methane production rates and decreased methane oxidation efficiency in sediments. However, the effect of OM lability on the structure of methane oxidizing (methanotrophic) and methane producing (methanogenic) microbial communities in lake sediments is still understudied. We studied the vertical profiles of the sediment and porewater geochemistry and the microbial communities (16S rRNA gene amplicon sequencing) at five profundal stations of an oligo-mesotrophic, boreal lake (Lake Pääjärvi, Finland), varying in surface sediment OM sources (assessed via sediment C:N ratio). Porewater profiles of methane, dissolved inorganic carbon (DIC), acetate, iron, and sulfur suggested that sites with more autochthonous OM showed higher overall OM lability, which increased remineralization rates, leading to increased electron acceptor (EA) consumption and methane emissions from sediment to water. When OM lability increased, the abundance of anaerobic nitrite-reducing methanotrophs (*Candidatus Methyloirabilis*) relative to aerobic methanotrophs (*Methylococcales*) in the methane oxidation layer of sediment surface decreased, suggesting that *Methylococcales* were more competitive than *Ca. Methyloirabilis* under decreasing redox conditions and increasing methane availability due to their more diverse metabolism (fermentation and anaerobic respiration) and lower affinity for methane. Furthermore, when OM lability increased, the abundance of methanotrophic community in the sediment surface layer, especially *Ca. Methyloirabilis*, relative to the methanogenic community decreased. We conclude that increasing input of labile OM, subsequently affecting the redox zonation of sediments, significantly modifies the methane producing and consuming microbial community of lake sediments.

IMPORTANCE Lakes are important natural emitters of the greenhouse gas methane (CH₄). It has been shown that eutrophication, via increasing the input of labile organic matter (OM) into lake sediments and subsequently affecting the redox conditions, increases methane emissions from lake sediments through increased sediment methane production rates and decreased methane oxidation efficiency. However, the effect of organic matter lability on the structure of the methane-related microbial communities of lake sediments is not known. In this study, we show that, besides the activity, also the structure of lake sediment methane producing and consuming microbial community is significantly affected by changes in the sediment organic matter lability.

KEYWORDS greenhouse gas, freshwater, methanotroph, methanogen, 16S rRNA gene, eutrophication

The concentration of atmospheric methane (CH₄), a critical greenhouse gas, has increased substantially since industrialization, with current total emissions of 550–

Editor Sandi Orlic, Institut Ruder Boskovic, Zagreb, Croatia

Ad Hoc Peer Reviewer Lisa Y. Stein, University of Alberta, Edmonton, Alberta, Canada

Address correspondence to Antti J. Rissanen, antti.rissanen@tuni.fi.

The authors declare no conflict of interest.

See the funding table on p. 12.

Received 12 May 2023

Accepted 17 July 2023

Published 12 September 2023

Copyright © 2023 Rissanen et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

600 Tg/year (top-down estimates) of which approximately 40% stem from natural sources (1). Lakes are important natural emitters of CH₄. The numerous lakes and ponds of the northern boreal zone, with annual emissions of ~16.5 Tg, are especially important contributors to the global CH₄ budget (2, 3). Thus, knowledge of the factors affecting CH₄ emissions from northern lakes is essential for accurate estimates and modeling of the global CH₄ budget and its changes due to global change (e.g., eutrophication and climate warming).

The CH₄ emissions from lake sediments are controlled by the balance between methanogenesis and CH₄ oxidation. Methanogenesis is the final step in anoxic organic matter (OM) degradation by methanogenic archaea that produce CH₄ from acetate or from oxidation of H₂ using CO₂ as an electron acceptor (EA) (4). Some methanogens can also produce CH₄ from other methyl compounds (e.g., methanol) (4, 5). In oxic surface sediments, CH₄ is consumed through aerobic CH₄ oxidation by methanotrophic bacteria (MOB) using O₂ as an EA (6). In anoxic conditions, CH₄ is consumed through anaerobic methane oxidation (AOM) by anaerobic methanotrophic archaea (ANME) utilizing various inorganic or organic compounds (7–9) or by bacteria within genus *Candidatus* *Methylomirabilis* utilizing NO₂⁻ as EAs (10). Furthermore, MOB belonging to *Methylococcales* have been shown to be capable of metabolizing CH₄ in hypoxic and anoxic conditions via fermentation and anaerobic respiration of various EAs, including NO₃⁻, NO₂⁻, Fe³⁺, and organic EAs (11–19).

Eutrophication was recently shown to increase CH₄ emissions from lake sediments through lowered CH₄ oxidation efficiency at increasing sediment methanogenesis rate (20). However, whether this phenomenon is due to change in the activity of methanogens and methanotrophs or also due to change in their community structure and abundance is not known. In support of the latter hypothesis, the results from the recent studies comparing lakes with different trophic states suggest that the density of sediment methanogens is higher while that of methanotrophs is lower, in eutrophic lakes that contain more labile algae-based sediment OM and where CH₄ fluxes from the sediment are higher than in oligotrophic and mesotrophic lakes (21–23). However, no previous study has simultaneously studied the change in both methanotrophic and methanogenic communities within a gradient of changing OM lability. The close spatial proximity of MOB and AOM-driving *Ca. Methylomirabilis* bacteria in lake sediments (24) also suggests that competition for CH₄ exists between these groups. Indeed, a recent modeling study suggested that an increase in OM quantity decreases the abundance of *Ca. Methylomirabilis* while increasing the abundance of MOB (25). Furthermore, in the comparison [using the 16S rRNA gene dataset by Han et al. (21)] of sediment methanotroph communities between lakes with varying trophic status by van Grinsven et al. (22), *Ca. Methylomirabilis* sp. was as abundant as MOB in the sediments of an oligotrophic lake but had negligible abundance in the sediments of meso- and eutrophic lakes, where MOB dominated the methanotrophic community. These differences suggest that MOB are more competitive in conditions of higher OM lability, when increased OM mineralization decreases redox potential and increases CH₄ availability. This is possibly due to MOB, especially *Methylococcales*, having a lower affinity for CH₄ and more diverse metabolism in hypoxic and anoxic conditions (incl. capability for fermentation and anaerobic respiration of various EAs as explained above) than *Ca. Methylomirabilis* (12, 15, 17, 18, 26–28). Hence, it could be expected that changes in the OM quality of lake sediments affect differently the abundances of MOB and *Ca. Methylomirabilis* sp. bacteria.

In this study, we investigated how spatial variability in OM lability in a single lake affects the community structure of CH₄ producing and consuming sedimentary microbes. Using 16S rRNA gene sequencing, we studied the variation in the methanogenic and methanotrophic community in the uppermost sediment layers at five sites with naturally varying OM quality of surface sediment within an oligo-mesotrophic boreal Lake Pääjärvi, Southern Finland. To determine OM lability, we used bulk sediment C:N ratios, which have been shown to broadly reflect relative contributions

of (autochthonous) phytoplankton and (allochthonous) terrestrial sources to bulk sedimentary OM (29, 30). Higher C:N ratios indicate a greater proportion of terrestrial OM, which is considered less labile (less available for microbial degradation) due to a combination of primary chemical composition, degradation during transport and protection by aggregation with mineral material [(31) and references therein], whereas lower C:N indicates more labile, phytoplankton-derived material [(32) and references therein]. We hypothesized that under increasing OM lability, (i) the relative contribution of methanotrophs within the CH₄ cycling community decreases while that of methanogens increases and (ii) the relative contribution of *Ca. Methyloirabilis* sp. bacteria within the CH₄ oxidizing community decreases while that of MOB increases.

MATERIALS AND METHODS

Study lake

Lake Pääjärvi is a NO₃⁻-rich, oligo-mesotrophic lake in Southern Finland (61.04N, 25.08E; A = 13.4 km², max. depth = 87 m, mean residence time = 3.3 yr). The water column circulates twice per year and is always well oxygenated (33). Field measurements of dissolved oxygen during this study (determined with a handheld YSI ODO probe) confirmed the presence of oxygen throughout the water column at all sediment sampling locations (Fig. S1). The hypolimnetic NO₃⁻ concentration, measured 2–5 cm above the sediment surface, is typically 46–75 μmol L⁻¹ (34, 35). The large catchment area of Lake Pääjärvi (244 km²) is dominated by forests and agriculture. The nutrient concentrations of the water have increased since the 1970s (36).

Porewater and sediment sampling

Vertical profiles of porewater and sediment samples were collected from five stations in Lake Pääjärvi on 9 August 2017 using a handheld HTH/Kajak corer with plexiglass tubes (Table 1). This study uses data from the top-most 10 cm layer (i.e., 0–10 cm from the sediment-water interface surface). The study stations followed a water depth gradient, Station 1 being the shallowest and Station 5 the deepest (Table 1). The core tubes were pre-drilled with two vertical series of 4 mm holes (each at 2 cm resolution), and then taped, in preparation for porewater sampling with Rhizons (Rhizosphere research products, Wageningen, Netherlands). Rhizon sampling automatically filters the porewaters at 0.15 μm into attached 10 mL syringes under vacuum. One vertical series of samples was taken for analysis of dissolved CH₄ and dissolved inorganic carbon (DIC). The syringes were pre-filled with 1 mL of 0.1M HNO₃ to immediately convert all DIC to CO₂. A known volume of N₂ gas headspace was injected into the syringes after sampling, and the samples were shaken to equilibrate the dissolved gases with the headspace. The subsamples of the headspace were then extracted into 3 mL Exetainers (Labco Limited, Lampeter, UK) and stored at room temperature (RT) until analysis [for full details see (37)]. The second vertical series was taken for S and Fe (by inductively coupled plasma atomic emission spectroscopy, ICP-OES) and short chain organic acid analysis. Short chain organic acid subsamples were stored frozen at –20°C until analysis. The subsamples for ICP-OES were acidified with 1 M HNO₃ and stored at RT. After porewater sampling, the sediment cores were sliced into plastic bags at a resolution of 1 cm. The subsamples of 400–500 μL wet sediment were collected from each slice and stored frozen at –20°C for DNA-based molecular microbiological analyses. The remaining wet sediment samples were stored frozen at –20°C under N₂ until further processing.

Porewater and sediment bulk geochemical analysis

The S and Fe concentrations in the porewater samples were determined by ICP-OES (Thermo iCAP 6000, Thermo Fisher Scientific, Waltham, MA, USA). In this system, S is expected to be dominated by sulfate (SO₄²⁻) and Fe by Fe²⁺, although in each case, other forms are possible. The porewater CH₄ and DIC concentrations were determined

TABLE 1 C:N ratios of the surface sediment [0–1 cm layer and 0–2 cm layer (i.e., average of 0–1 cm and 0–2 cm layers)], sediment porewater acetate and dissolved inorganic carbon (DIC) concentrations, and Shannon diversity index of prokaryotic diversity in the sediment (average \pm SD within the 0–10 cm layer) as well as the estimated CH₄ emissions from sediment to water (based on porewater CH₄ profiles) at the study stations^b

Station	Depth (m)	C:N (0–1)	C:N (0–2)	Acetate ($\mu\text{mol L}^{-1}$)	SD	DIC ($\mu\text{mol L}^{-1}$)	SD	CH ₄ flux ($\mu\text{mol m}^{-2} \text{d}^{-1}$)	Shannon	SD
3	52	14.74	14.73	9.3	8.5	553.4	269.3	280.6	6.71	0.16
2	22	13.40	13.78	5.3	7.3	589.7	322.6	99.5	6.80	0.21
4	60	12.90	13.38	15.4	1.3	675.8	140.0	1,148.2	6.93	0.11
1	14	12.78	13.25	13.7	2.5	900.8	93.0	1,262.6 ^a	7.01	0.10
5	80	12.54	12.50	14.4	0.8	965.4	144.8	4,324.8 ^a	6.99	0.13

^aCH₄ flux estimates for stations 1 and 5 should be considered as maximum estimates because the sampling resolution was lower at the top of the core (due to geometry of the core in the tube) and, therefore, the gradient at the sediment-water-interface may be less steep in reality (see Fig. 1).

^bStudy stations are organized in the order of increasing organic matter lability based on decreasing surface sediment C:N ratios. See full vertical profiles (for the 0–10 cm layer) of C:N ratio, DIC, acetate, and Shannon diversity index in Fig. S2A through D, respectively.

by gas chromatography (GC) as described in Jilbert et al. (37). Briefly, the sample vials (Exetainers) were pressurized with helium (He) to 2.0 bar before loading into the GC (Agilent technologies 7890B GC system, Agilent Technologies, Santa Clara, Ca, USA). CH₄ was determined by a flame-ionization detector (FID) and CO₂ by a thermal conductivity detector (TCD). The instrument simultaneously measures N₂ and O₂ + Ar (in TCD), from which a 100% sum can be calculated for the estimation of CH₄ and CO₂ concentrations in the original sample, in ppm by volume. For a full description of the calculations, see Jilbert et al. (37). Diffusive fluxes of methane across the sediment-water-interface were estimated using Fick's Law:

$$F = \frac{\varphi \cdot D}{\theta^2} \frac{\partial C}{\partial z}$$

where F = flux in $\mu\text{mol m}^{-2} \text{d}^{-1}$, D = diffusion coefficient of CH₄ in freshwater, based on a value of $1.67 \times 10^{-9} \text{ m}^2 \text{ s}^{-1}$ at 25°C and adjusted downward according to bottom water temperature, using Eq. 4.57 in Boudreau (38); φ = volume fraction of total porosity and θ = tortuosity, as related by $\theta^2 = 1 - \ln(\varphi^2)$; and $\partial C/\partial z$ is the partial differential gradient estimated from the finite difference gradient $\Delta C/\Delta z$, the concentration gradient of methane between the uppermost porewater sample and the overlying water in the sediment core tube.

Sediment samples were freeze-dried, grounded in an agate mortar, and weighed into tin cups for carbon (C) and nitrogen (N) content determinations, which were determined using an elemental analyzer (LECO TruSpec Micro, LECO Corp., St. Joseph, MI, USA). In accordance with extensive previous studies on Finnish lake sediments (39, 40), acidification was not applied prior to the determinations. High levels of organic acidity from Finnish river catchments (41) maintain low annual mean pH values in most lakes and, therefore, there is a negligible occurrence of carbonates in lake sediments. Hence, our total C data is considered equivalent to organic C (C_{org}), and total N is considered equivalent to organic N (N_{org}).

Porewater short chain organic acids were analyzed using high performance liquid chromatography (HPLC) equipped with Shodex SUGAR column (300 mm \times 8 mm, Phenomenex, Torrance, CA, USA), autosampler (SIL-20AC HT, Shimadzu, Kyoto, Japan), refractive index detector (RID-10A, Shimadzu), and 0.01 M H₂SO₄ as the mobile phase. The HPLC samples were prepared as described in Salmela et al. (42). The identification and quantification of the liquid metabolites were conducted using external standards.

Molecular microbiological analyses

DNA was extracted from the frozen sediment samples using DNeasy PowerSoil Kit (Qiagen, Hilden, Germany). DNA concentration was measured using a Qubit 2.0 Fluorometer and a dsDNA HS Assay Kit (Thermo Fisher Scientific, Waltham, MA, USA).

PCR and 16S rRNA gene amplicon sequencing was performed commercially by the Foundation for the Promotion of Health and Biomedical Research of Valencia Region

(FISABIO, Valencia, Spain). In the PCR reactions, the V4 region of the bacterial and archaeal 16S rRNA genes was simultaneously targeted using primer pair 515FB (5'-GTGYCAGCMGCCGCGGTAA-3')/806FB (5'-GGACTACNVGGGTWTCTAAT-3') (43, 44). PCR, library preparation, and paired-end sequencing (Illumina MiSeq, Illumina, San Diego, CA, USA) were performed as previously described (45), except that, in PCR reactions, approximately 15 ng of DNA were used.

The quality assessment of the raw sequence reads, merging of paired-end reads, alignment, chimera removal, preclustering, taxonomic classification (using Silva database 132), and removal of chloroplast, mitochondria, and eukaryote sequences, and division of 16S rRNA gene sequences into operational taxonomic units (OTUs) at 97% similarity level was conducted as described in Rissanen et al. (11) (the detailed description is available also in Supplementary Methods). Singleton OTUs (OTUs with only one sequence) were removed, and the data were then normalized by subsampling to the same size, 77,340 sequences. One sample, representing the layer 8–9 cm depth at Station 3, was discarded from the analyses since it had only ~10,000 joined sequence reads. Good coverage was 0.95–0.97 in each library confirming that sequence variation was adequately covered. Prokaryotic diversity was assessed via calculation of the Shannon diversity index for each library. To test the hypotheses, this study focused specifically on the relative abundance (% of prokaryotic 16S rRNA genes) of known aerobic (46) and anaerobic methanotrophic bacteria (10), anaerobic methanotrophic archaea (47) as well as methanogenic archaea (5, 48).

Statistical analyses

The relationship between microbial variables and the OM quality of sediment was analyzed using the Spearman's rank correlation test. Data showing significant correlations ($P < 0.05$) were further studied using linear and non-linear regression analyses. The microbial variables included the Shannon diversity index and the relative abundances (percentage of 16S rRNA genes) of methanotrophs and methanogens (both total and different taxonomic groups), while C:N ratio of the surface sediment (0–1 and 0–2 cm layer) was used as a proxy for the OM quality of sedimenting OM (i.e., decreasing C:N ratio indicates increasing lability of sedimenting OM). Furthermore, sediment porewater DIC concentration was used as a further indicator of sediment OM lability in the correlation analyses (i.e., increasing DIC indicates increasing OM mineralization due to increased OM lability). We acknowledge that the relative abundances of microbes do not predict their absolute abundances. However, ratios of the relative abundances of different organisms to each other are robust against variations in their absolute abundances. Therefore, besides relative abundances, we also analyzed abundance ratios of microbes, such as the ratio of *Ca. Methyloirabilis*/*Methylococcales*, Methanotrophs/Methanogens, and *Ca. Methyloirabilis*/Methanogens. Correlation analyses were done using IBM SPSS Statistics 26 (IBM SPSS Statistics for Windows, Version 26.0, IBM Corp., Armonk, NY, USA), while the regression analyses were done using Minitab software (Minitab Statistical Software for Windows, Version 16.2.0.0, Minitab Inc., PA, USA). To determine the best-fitting models to the experimental data, along with their 95% confidence intervals (CIs), the parameter values of models were adjusted so as to minimize the squared deviation between the data values and the fitted values (S) and via checking the normality of residuals (P -value > 0.05).

RESULTS AND DISCUSSION

Sediment OM quality, vertical porewater profiles, and prokaryotic diversity

Stations were ordered according to increasing OM lability as determined by decreasing C:N ratio in surface sediment as follows: Station 3, Station 2, Station 4, Station 1, and Station 5 (Table 1; Fig. S2A). This ordering did not follow the depth gradient, and therefore likely represents the heterogeneity of sedimentation of autochthonous and allochthonous OM within the lake. Surface sediment values were used here due to

assumed minimum overprinting of sediment diagenetic processes on the C:N ratio; hence, the values should represent the C:N ratio (lability) of the OM sedimenting to the lake bottom at the stations. Using C:N ratio of either the 0–1 cm layer or the average of 0–1 cm and 0–2 cm layers had no impact on the order of stations in terms of OM lability (Table 1; Fig. S2A). Hence, the possible mixing of surface sediment by bioturbation did not affect the major pattern in the OM lability between stations. The range of observed molar C:N ratios is quite narrow (approx 12.5–14.7 for the 0–1 cm interval) indicating an overall dominance of allochthonous terrestrial OM according to the end-member values of Goñi et al. (29), but small differences in the contribution of autochthonous OM appear to strongly influence the overall OM lability. Evidence for an OM lability gradient was shown in porewater DIC and acetate data. Average concentration of DIC within the 0–10 cm layer increased when there was a decrease in surface sediment C:N (Table 1, Fig. S2B) ($\rho = -1.00$, $P < 0.0001$, for both C:N ratio of 0–1 cm layer and average of C:N ratio of 0–1 and 0–2 cm layers). Although not significantly correlated with surface sediment C:N ratio ($\rho = -0.6$, $P = 0.285$, for both C:N ratio of 0–1 cm layer and average of C:N ratio of 0–1 and 0–2 cm layers), acetate was also higher in stations with lower (i.e., Stations 4, 1, and 5) than with higher surface sediment C:N (Stations 3 and 2) (Table 1; Fig. S2C). Lactate was also detected but only within the sediments of Station 1, which was among the stations with the lowest surface sediment C:N ratio (Fig. S2C). These results suggest that OM remineralization rates (producing DIC, acetate, and lactate) were higher in stations with lower surface sediment C:N, which reflects their higher sediment OM lability.

According to visual analysis of the porewater data, the differences in the vertical profiles of CH₄, S (assumed to be mostly SO₄²⁻) and Fe (assumed to be mostly Fe²⁺ from reduction of solid-phase Fe oxides) between the stations followed broadly the differences in OM lability (Fig. 1). CH₄ concentrations increased from the sediment surface downward with depth at each station (Fig. 1A through E). However, compared to other stations, they remained at very low levels at surface 0–4 cm layer at Stations 3 and 2 with the lowest sediment OM lability, while the highest surface sediment CH₄ concentrations were observed at Station 5 with the highest OM lability (Fig. 1A through E). This pattern generally agrees with the comparison of sediment CH₄ profiles between eutrophic (labile sediment OM) and oligotrophic (less labile OM) lakes by van Grinsven et al. (22). This suggests that active zones of methanogenesis extend closer to the sediment surface, and CH₄ oxidation takes place in a thinner surface layer in stations with higher sediment OM lability than in those with lower OM lability (Fig. 1A through E). In addition, S (i.e., SO₄²⁻, see above) and Fe (Fe²⁺, see above) profiles suggested that the zone of reduction of SO₄²⁻ and Fe oxides were located deeper from the sediment surface in the stations with less labile OM, as seen in S accumulation zone extending deeper and Fe accumulation zone starting deeper from the sediment surface in stations with less labile OM (Fig. 1F through O). As the water column was well oxygenated at each station (Fig. S1), the differences in the porewater profiles of CH₄, S, and Fe between stations are not explained by differences in oxygen availability in the water overlying the sediment but are very likely driven by the differences in the sediment OM lability. Altogether, the porewater profiles suggested a higher rate of OM processing, higher consumption of O₂ and alternative EAs, lower redox potential, and subsequently higher rate of methanogenesis in the stations with higher sediment OM lability (Fig. 1). This agrees with previous results on the comparison of lakes with different trophic status (22, 23). In accordance, the modeled diffusive methane emissions from sediment to water were generally higher in the stations with higher sediment OM lability (Table 1) ($\rho = -0.90$, $P < 0.05$, for both C:N ratio of 0–1 cm layer and average of C:N ratio of 0–1 and 0–2 cm layers), which agrees with the results by van Grinsven et al. (22). The magnitudes of estimated methane emissions from sediment to water, 99.5–4324.8 $\mu\text{mol m}^{-2} \text{d}^{-1}$ (Table 1), agree well with the range of previously measured data for boreal lakes, i.e., 300–6562.5 $\mu\text{mol m}^{-2} \text{d}^{-1}$ (49).

Prokaryotic diversity, assessed via analysis of Shannon diversity index, increased when sediment OM lability increased (i.e., when C:N ratio decreased) (Table 1; Fig. S2D) ($\rho = -0.90$, $P < 0.05$, for both C:N ratio of 0–1 cm layer and average of C:N ratio of 0–1 and




FIG 1 Concentrations of CH₄ (A-E), Fe (F-J), and S (K-O) in the porewater at different depths of sediments (incl. water overlying the sediment) at the five study stations shown in the order of increasing OM lability (from left to right; St3, St2, St4, St1, and St5) based on surface sediment C:N ratio (see Table 1). Depth 0 cm (dashed line) indicates the sediment-water interface.

0–2 cm layers). This result agrees with those from soil ecosystems showing negative correlation between bacterial diversity and soil C:N ratio (50). This can be explained by high resource quality (i.e., labile OM with low C:N ratio) leading to a greater variety of resources for bacterial communities, which enhances their diversity by promoting greater niche differentiation (50, 51). Hence, variation in the prokaryotic diversity further highlights the differences in OM lability between stations.

Methanotrophic community

Aerobic MOB in the order *Methylococcales* as well as anaerobic methanotrophs in genera *Ca. Methyloirabilis* sp. (within bacterial phylum *NC10*) and *Ca. Methanoperedens* sp. (also known as ANME 2D archaea) were the most abundant groups of methanotrophs present in the studied sediments (Fig. 2A through E). This agrees with a previous study on boreal lake sediments (52). Aerobic MOB in the family *Methylacidiphilaceae* (i.e., Verrucomicrobial methanotrophs) were rare, while alphaproteobacterial MOB were not detected (Fig. 2A through E). The relative abundance of both *Methylococcales* and *Ca. Methyloirabilis* sp. generally peaked in the surface 0–3 cm layer at each station, while *Ca. Methyloirabilis* was occasionally observed also at deeper depths (Fig. 2A through E). In contrast to *Methylococcales* and *Ca. Methyloirabilis* sp., *Ca. Methanoperedens*

sp. archaea had peaks in its relative abundance only at deeper layers, clearly below the sediment surface (i.e., below 5 cm depth) (Fig. 2A through E). In the deep layers, *Ca. Methanoperedens* sp. archaea can potentially use a wide variety of EAs in AOM, i.e., SO_4^{2-} , Fe^{3+} minerals, and organic compounds (8, 47, 53–56), but as the scope of this study is focused on *Ca. Methyloirabilis* sp. and MOB, the putative role of *Ca. Methanoperedens* sp. archaea in the biogeochemical processes of the study lake is not considered further.

The CH_4 profiles in porewater show clearly depleted CH_4 concentrations near the sediment surface (Fig. 1A through E), suggesting that abundant *Methylococcales* and *Ca. Methyloirabilis* sp. act as a crucial filter in the top 0–3 cm sediment layer, reducing CH_4 fluxes from sediment to the overlying water column (Fig. 2A through E). The overlap in the vertical distribution of *Methylococcales* and *Ca. Methyloirabilis* sp. agrees with previous results from lake sediments and suggests competition between these groups (22, 24), as further indicated by a recent modeling study (25). Therefore, we tested the hypothesis that the relative contribution of *Ca. Methyloirabilis* sp. and MOB within the CH_4 oxidizing community is sensitive to OM lability. In this analysis, we specifically focused on the top 0–3 cm sediment layer, as based on depleted CH_4 concentrations and high methanotroph relative abundances at that layer, it is considered to be the key CH_4 filter layer in reducing CH_4 emissions from sediment to water (Fig. 1A through E and 2A through E). In partial support for the hypothesis, we found that within that layer, the relative abundance of *Ca. Methyloirabilis* decreased, when OM lability increased, i.e., *Ca. Methyloirabilis* correlated positively with sediment C:N ratio (Table 2; Fig. 3A), while the relative abundance of *Methylococcales* was not correlated with the C:N ratio (Table 2). Consequently, the ratio of *Ca. Methyloirabilis* to *Methylococcales* decreased alongside when OM lability increased (C:N decreased) (Table 2; Fig. 3B). As it is challenging to quantify the relative abundance of taxonomic groups in a layer (i.e., 0–3 cm) consisting of multiple sublayers (i.e., 0–1 cm, 0–2 cm, and 0–3 cm), we increased the robustness in our results by carrying out these analyses both with the average relative abundance and the maximum relative abundance of *Methylococcales* and *Ca. Methyloirabilis* within the 0–3 cm sediment layer, with both choices giving similar results for the change in *Ca. Methyloirabilis* and ratio of *Ca. Methyloirabilis* to *Methylococcales* alongside the OM




FIG 2 Relative abundance (percentage of prokaryotic 16S rRNA gene reads) of methanotrophs (A–E) and methanogens (F–J) at different depths in the sediment at the five study stations shown in the order of increasing OM lability (from left to right; St3, St2, St4, St1, and St5) based on the surface sediment C:N ratio (see Table 1).

lability gradient (Table 2). Besides surface sediment C:N ratio, correlation analyses were conducted using porewater DIC concentrations, which also indicate sediment OM lability (i.e., increasing DIC indicate increasing OM lability), with results identical to those with C:N ratios (Table 2). Hence, in accordance with He et al. (25) and van Grinsven et al. (22), our results suggest that increasing input of labile OM leads to outcompetition of *Ca. Methyloirabilis* sp. by *Methylococcales* in lake sediments, which is very likely explained by their different metabolisms. While *Ca. Methyloirabilis* sp. bacteria are restricted in using NO_2^- as an EA in CH_4 oxidation and have been reported to have a high affinity for CH_4 (10, 28), *Methylococcales* are potentially capable of coupling CH_4 oxidation with fermentation and with reduction of a variety of EAs (e.g., O_2 , NO_3^- , NO_2^- , Fe^{3+} , and organic EAs) and have a low affinity for CH_4 (11–19, 26, 27). This gives *Methylococcales* advantage over *Ca. Methyloirabilis* in conditions of increased OM lability, when redox potential and availability of EAs is decreased and availability of CH_4 is increased.

Methanogenic community

In accordance with lower redox potential and more suitable conditions for methanogenesis at stations with higher OM lability, the relative abundance of methanogens generally increased when the sediment C:N ratio decreased (Table 2; Fig. 2F through J; Fig. 3C). Methanogens were present all through the 0–10 cm layer at all stations. However, they had very low relative abundance in the surface 0–5 cm layer at all stations except for Station 5 with the highest sediment OM lability. This agrees with the CH_4 profile data described above, further suggesting that the methanogenesis zone extends closer to the sediment surface at stations with high OM lability (Fig. 1E and 2J). Our results agree with those of Yang et al. (23) showing a higher density of methanogens in sediments of an eutrophic than a mesotrophic lake. Based on the study by D'Ambrosio and Harrison (20) on the effect of eutrophication increasing the CH_4 emissions from lake sediments due to lowered CH_4 oxidation efficiency at increased methanogenesis rate, we hypothesized that increasing sediment OM lability would similarly lead to decreased abundance of methanotrophs in relation to methanogens. As above, we considered the top 0–3 cm CH_4 filter layer for methanotrophs in these analyses. For methanogens, we chose the layer 4–10 cm as it is below the CH_4 filter layer, and the relative abundance of methanogens is clearly highest below than above 4 cm depth at all stations except Station 5 (Fig. 2F through J). However, due to methanogens being abundant also above 4 cm depth at Station 5, we did the analyses for methanogens also by considering the whole 0–10 cm layer. In support of our hypothesis, the ratio of relative abundance of methanotrophs (sum of all methanotrophs) in the top 0–3 cm aerobic CH_4 filter layer to the relative abundance of methanogens in the 4–10 cm or 0–10 cm layer decreased when sediment C:N ratio decreased (Table 2). More specifically, this was due to the fact that the ratio *Ca. Methyloirabilis* to methanogens decreased strongly when the sediment C:N ratio decreased (Table 2; Fig. 3D). Similar to the hypothesis testing considering methanotrophs (see above), to increase robustness of the results, these analyses were carried out using both the average and the maximum relative abundance of methanotrophs and methanogens within the chosen layers (i.e., 0–3 cm for methanotrophs and 4–10 cm or 0–10 cm for methanogens), with similar results for the change in the ratio of *Ca. Methyloirabilis* to methanogens alongside the OM lability gradient (Table 2). As above for the hypothesis testing considering methanotrophs, the correlation analyses were also conducted using porewater DIC concentrations, with results identical to those with C:N ratios (Table 2). Hence, based on our results, it can be suggested that increasing input of labile OM changes lake sediment microbial community toward a lower genetic potential for methanotrophy relative to the genetic potential for methanogenesis (Table 2; Fig. 3). Further studies are required to assess whether the observed changes in the genetic potential have any role in controlling the sediment-to-water CH_4 emissions or whether the CH_4 emissions are solely determined by the activity of methanotrophs and methanogens.

TABLE 2 Spearman correlation analysis results^a on the relationship between microbial variables and environmental variables indicating sediment OM lability, i.e., C:N ratio^b of the surface sediments (representing C:N ratio of sedimenting OM) and concentration of porewater dissolved inorganic carbon (DIC)^c

Microbial variables	C:N ^b	C:N ^b	DIC ^c	DIC ^c
	(aver.)	(max.)	(aver.)	(max.)
Relative abundances:				
Total methanotrophs (0–3 cm)	0.7	0.9	–0.7	–0.9
<i>Methylococcales</i> (0–3 cm)	–0.7	–0.6	0.7	0.6
<i>Ca. Methyloirabilis</i> (0–3 cm)	0.9	0.9	–0.9	–0.9
<i>Ca. Methanoperedens</i> (0–3 cm)	0.1	–0.2	–0.1	0.2
Total methanogens (4–10 cm)	–0.9	–0.7	0.9	0.7
Total methanogens (0–10 cm)	–0.9	–0.7	0.9	0.7
Ratios of relative abundances:				
<i>Ca. Methyloirabilis</i> (0–3 cm)/ <i>Methylococcales</i> (0–3 cm)	1	1	–1	–1
Total methanotrophs (0–3 cm)/Total methanogens (4–10 cm)	0.9	0.6	–0.9	–0.6
Total methanotrophs (0–3 cm)/Total methanogens (0–10 cm)	0.9	0.6	–0.9	–0.6
<i>Methylococcales</i> (0–3 cm)/Total methanogens (4–10 cm)	0.1	–0.4	–0.1	0.4
<i>Methylococcales</i> (0–3 cm)/Total methanogens (0–10 cm)	0.1	–0.4	–0.1	0.4
<i>Ca. Methyloirabilis</i> (0–3 cm)/Total methanogens (4–10 cm)	1	1	–1	–1
<i>Ca. Methyloirabilis</i> (0–3 cm)/Total methanogens (0–10 cm)	1	1	–1	–1

^a $n=5$, statistically significant results ($P < 0.05$) highlighted in bold. Negative and positive correlations with C:N ratio and positive and negative correlation with DIC indicate increase and decrease, respectively, with increasing sediment OM lability. Aver. and max in brackets below the column title denote whether average or maximum relative abundance, respectively, in 0–3 cm (for methanotrophs), 4–10 cm or 0–10 cm sediment layers (for methanogens) was used in the analysis (see text).

^bUsing either C:N ratio in 0–1 cm sediment layer or C:N ratio of 0–2 cm sediment layer (i.e., average C:N ratio of 0–1 cm and 1–2 cm sediment layers) gave identical results. The C:N ratios are shown in Table 1.

^cAverage porewater DIC concentration within the 0–10 cm sediment layer. The values are shown in Table 1.

The order *Methanomassiliicoccales*, whose cultivated members use the methyl reducing methanogenic pathway (5), dominated the methanogen communities at stations with the lowest OM lability in sediment (Stations 3 and 2), while in addition to these, other methanogenic groups, such as the hydrogenotrophic (H₂ consuming) CO₂ reducing *Methanomicrobiales* (48), were relatively abundant members of methanogenic community at stations with higher OM lability (Stations 4, 1, and 5) (Fig. 2F through J). Also, acetoclastic (acetate using) *Methanosaetaeaceae* (48) was relatively abundant in Stations 4 and 5 (Fig. 2H and J).

The dominance of *Methanomassiliicoccales* is surprising because methanogenesis in freshwater sediments is typically dominated by the acetoclastic and hydrogenotrophic CO₂ reducing pathways (4). Yet, also in the eutrophic Lake Dianchi, *Methanomassiliicoccales* were dominant members of the methanogenic community, constituting at maximum over 20% of methanogens (23). However, recent metagenomic data suggested the presence of the acetoclastic pathway in genomes of uncultivated *Methanomassiliicoccales*, which could explain these findings (57). On the other hand, being only a very recently described order with limited information on its functional diversity (58), *Methanomassiliicoccales* might also contain non-methanogenic species. Therefore, the correlation analyses were made also by excluding *Methanomassiliicoccales* from the sum of total methanogens, but it did not affect the outcome of the study (Table 2; Table S2). The higher methanogenic functional diversity due to the increasing presence of hydrogenotrophic, CO₂ reducing *Methanomicrobiales*, and acetoclastic *Methanosaetaeaceae* at stations with higher sediment OM lability (i.e., Stations 4, 1, and 5) further reflects a lower redox potential and more suitable conditions for methanogenesis at these stations (Fig. 2F through J).

Conclusion

Our results on the porewater profiles of lake sediments suggest modified redox zonation caused by changing sediment OM lability. This was because increased OM lability likely enhanced OM processing and consumption of electron acceptors. We also provide evidence that changes in lake sediment OM lability affect the structure of the CH₄ cycling




FIG 3 Regression analysis results [observations in black dots, regression line in black line, and 95% confidence intervals (CIs) in red dot line] on the dependence of (A) the relative abundance of *Ca. Methylomirabilis* sp., (B) the ratio of abundances of *Ca. Methylomirabilis* to *Methylococcales*, (C) the relative abundance of total methanogens, and (D) the ratio of abundances of *Ca. Methylomirabilis* to total methanogens, with the surface sediment C:N ratios (i.e., C:N ratio of 0–1 cm layer). The average relative abundance of *Ca. Methylomirabilis* and *Methylococcales* within the 0–3 cm layer and the average relative abundance of methanogens within the 4–10 cm layer were used in the analyses. See Table S1 for detailed statistics of the regression models.

microbial community. By generating suitable conditions for methanogenesis (i.e., higher availability of methanogenic substrates and decreased availability of alternative EAs), increasing sediment OM lability increased the relative abundance of methanogens. Furthermore, increasing OM lability caused changes in the structure of the CH₄ filtering methanotrophic community by reducing the abundance of anaerobic nitrite-reducing methanotrophic *Ca. Methylomirabilis* sp. in relative to aerobic methanotrophic *Methylococcales* in lake sediments, which suggests that *Methylococcales* thrived better than *Ca. Methylomirabilis* under decreasing redox conditions and increasing methane availability due to their more diverse metabolism (fermentation and anaerobic respiration of various EAs) and lower affinity for CH₄. The results also suggest that increased OM lability decreases the abundance of methanotrophs, especially *Ca. Methylomirabilis*, relative to methanogens. Altogether, our results suggest that increasing input of labile algae-based OM changes the redox zonation of the sediment and exerts significant changes on the methanotrophic and methanogenic microbial community of lake sediments.

ACKNOWLEDGMENTS

We thank the staff at the Lammi Biological Station for their support in field and laboratory work. Reviewers are acknowledged for their valuable comments and suggestions that improved the paper.

This study was supported by Academy of Finland (Grant No. 286642, 346751, and 353750 for A.J.R., 323214 for R.M., 307331 for H.J., and 310302 for S.L.A.), joint funding by Olvi Foundation, Jenny and Antti Wihuri Foundation, and Saastamoinen Foundation (for H.J.) and Kone Foundation (Grant No. 201803224 for A.J.R.), and a Tenure Track starting package from University of Helsinki (T.J.). Open access funding was provided by Tampere University.

Author contributions: Antti J. Rissanen: Conceptualization, Methodology, Formal analysis, Investigation, Funding acquisition, Project administration, Writing—original draft, Writing—review and editing; Tom Jilbert: Conceptualization, Methodology, Formal analysis, Investigation, Funding acquisition, Writing—review and editing; Asko Simojoki: Methodology, Formal analysis, Writing—review and editing; Rahul Mangayil: Methodology, Formal analysis, Writing—review and editing; Sanni L. Aalto: Methodology, Formal analysis, Writing—review and editing; Ramita Khanongnuch: Methodology, Formal analysis, Writing—review and editing; Sari Peura: Methodology, Formal analysis, Writing—review and editing; Helena Jäntti: Conceptualization, Methodology, Formal analysis, Investigation, Funding acquisition, Project administration, Writing—review and editing.

AUTHOR AFFILIATIONS

¹Faculty of Engineering and Natural Sciences, Tampere University, Tampere, Finland

²Natural Resources Institute Finland (Luke), Helsinki, Finland

³Environmental Geochemistry Group, Department of Geosciences and Geography, Faculty of Science, Helsinki, Finland

⁴Department of Agricultural Sciences (Environmental Soil Science), Faculty of Agriculture and Forestry, University of Helsinki, Helsinki, Finland

⁵Department of Environmental and Biological Sciences, University of Eastern Finland, Kuopio, Finland

⁶Department of Biological and Environmental Sciences, University of Jyväskylä, Jyväskylä, Finland

⁷Department of Forest Mycology and Plant Pathology, Science for Life Laboratory, Swedish University of Agricultural Sciences, Uppsala, Sweden

PRESENT ADDRESS

Rahul Mangayil, Department of Bioproducts and Biosystems, School of Chemical Engineering, Aalto University, Aalto, Finland

Sanni L. Aalto, Technical University of Denmark, DTU Aqua, Section for Aquaculture, The North Sea Research Centre, Hirtshals, Denmark

AUTHOR ORCID*s*

Antti J. Rissanen  <http://orcid.org/0000-0002-5678-3361>

FUNDING

Funder	Grant(s)	Author(s)
Academy of Finland (AKA)	286642, 346751, 353750	Antti J. Rissanen
Academy of Finland (AKA)	323214	Rahul Mangayil
Academy of Finland (AKA)	307331	Helena Jäntti
Academy of Finland (AKA)	310302	Sanni L. Aalto
OLVI-Säätiö (Olvi Foundation)		Helena Jäntti

Funder	Grant(s)	Author(s)
WRI Jenny ja Antti Wihurin Rahasto (Jenny JA Antti Wihurin Rahasto sr)		Helena Jäntti
Saastamoisen säätiö (Saastamoinen Foundation)		Helena Jäntti
Koneen Säätiö (Kone Foundation)	201803224	Antti J. Rissanen
University of Helsinki		Tom Jilbert

DATA AVAILABILITY

The reads of the 16S rRNA gene amplicon data set are deposited in NCBI's Sequence Read Archive (SRA) under the accession number [PRJNA627335](#) for Stations 1–3 and [PRJNA924099](#) for Stations 4 and 5. The other data are available in the supplemental data sets (see Supplemental Material).

ADDITIONAL FILES

The following material is available [online](#).

Supplemental Material

Supplemental methods, tables, and figures (Spectrum01955-23-S0001.doc). Appendix A: Supplemental methods, Tables S1 and S2, Figures S1 and S2.

Datasets (Spectrum01955-23-S0002.xlsx). Contains all datasets except for 16S rRNA gene sequence data, which has been stored in NCBI's SRA database.

Open Peer Review

PEER REVIEW HISTORY (review-history.pdf). An accounting of the reviewer comments and feedback.

REFERENCES

- Saunio M, Stavert AR, Poulter B, Bousquet P, Canadell JG, Jackson RB, Raymond PA, Dlugokencky EJ, Houweling S, Patra PK, Ciais P, Arora VK, Bastviken D, Bergamaschi P, Blake DR, Brailsford G, Bruhwiler L, Carlson KM, Carrol M, Castaldi S, Chandra N, Crevoisier C, Crill PM, Covey K, Curry CL, Etiope G, Frankenberg C, Gedney N, Hegglin MI, Höglund-Isaksson L, Hugelius G, Ishizawa M, Ito A, Janssens-Maenhout G, Jensen KM, Joos F, Kleinen T, Krummel PB, Langenfelds RL, Laruelle GG, Liu L, Machida T, Maksyutov S, McDonald KC, McNorton J, Miller PA, Melton JR, Morino I, Müller J, Murguía-Flores F, Naik V, Niwa Y, Noce S, O'Doherty S, Parker RJ, Peng C, Peng S, Peters GP, Prigent C, Prinn R, Ramonet M, Regnier P, Riley WJ, Rosentretter JA, Segers A, Simpson IJ, Shi H, Smith SJ, Steele LP, Thornton BF, Tian H, Tohjima Y, Tubiello FN, Tsuruta A, Viovy N, Voulgarakis A, Weber TS, van Weele M, van der Werf GR, Weiss RF, Worthy D, Wunch D, Yin Y, Yoshida Y, Zhang W, Zhang Z, Zhao Y, Zheng B, Zhu Q, Zhu Q, Zhuang Q. 2020. The global methane budget 2000–2017. *Earth Syst Sci Data* 12:1561–1623. <https://doi.org/10.5194/essd-12-1561-2020>
- Bastviken D, Tranvik LJ, Downing JA, Crill PM, Enrich-Prast A. 2011. Freshwater methane emissions offset the continental carbon sink. *Science* 331:50. <https://doi.org/10.1126/science.1196808>
- Wik M, Varner RK, Anthony KW, MacIntyre S, Bastviken D. 2016. Climate-sensitive northern lakes and ponds are critical components of methane release. *Nature Geosci* 9:99–105. <https://doi.org/10.1038/ngeo2578>
- Conrad R. 1999. Contribution of hydrogen to methane production and control of hydrogen concentrations in methanogenic soils and sediments. *FEMS Microbiol Ecol* 28:193–202. <https://doi.org/10.1111/j.1574-6941.1999.tb00575.x>
- Borrel G, O'Toole PW, Harris HMB, Peyret P, Brugère J-F, Gribaldo S. 2013. Phylogenomic data support a seventh order of methylophilic methanogens and provide insights into the evolution of methanogenesis. *Genome Biol Evol* 5:1769–1780. <https://doi.org/10.1093/gbe/evt128>
- Hanson RS, Hanson TE. 1996. Methanotrophic bacteria. *Microbiol Rev* 60:439–471. <https://doi.org/10.1128/mr.60.2.439-471.1996>
- Beal EJ, House CH, Orphan VJ. 2009. Manganese- and iron-dependent marine methane oxidation. *Science* 325:184–187. <https://doi.org/10.1126/science.1169984>
- Ettwig KF, Zhu B, Speth D, Keltjens JT, Jetten MSM, Kartal B. 2016. Archaea catalyze iron-dependent anaerobic oxidation of methane. *Proc Natl Acad Sci U S A* 113:12792–12796. <https://doi.org/10.1073/pnas.1609534113>
- Scheller S, Yu H, Chadwick GL, McGlynn SE, Orphan VJ. 2016. Artificial electron acceptors decouple archaeal methane oxidation from sulfate reduction. *Science* 351:703–707. <https://doi.org/10.1126/science.aad7154>
- Ettwig Katharina F, Butler MK, Le Paslier D, Pelletier E, Mangenot S, Kuypers MMM, Schreiber F, Dutilh BE, Zedelius J, de Beer D, Gloerich J, Wessels HJCT, van Alen T, Luesken F, Wu ML, van de Pas-Schoonen KT, Op den Camp HJM, Janssen-Megens EM, Francoijs K-J, Stunnenberg H, Weissenbach J, Jetten MSM, Strous M. 2010. Nitrite-driven anaerobic methane oxidation by oxygenic bacteria. *Nature* 464:543–548. <https://doi.org/10.1038/nature08883>
- Rissanen AJ, Saarela T, Jäntti H, Buck M, Peura S, Aalto SL, Ojala A, Pumpanen J, Tirola M, Elvert M, Nykänen H. 2021. Vertical stratification patterns of methanotrophs and their genetic controllers in water columns of oxygen-stratified boreal lakes. *FEMS Microbiol Ecol* 97:fiaa252. <https://doi.org/10.1093/femsec/fiaa252>
- Kits KD, Klotz MG, Stein LY. 2015. Methane oxidation coupled to nitrate reduction under hypoxia by the Gammaproteobacterium *Methylomonas denitrificans*. *Environ Microbiol* 17:3219–3232. <https://doi.org/10.1111/1462-2920.12772>
- Rissanen A, Saareheimo J, Tirola M, Peura S, Aalto S, Karvinen A, Nykänen H. 2018. Gammaproteobacterial methanotrophs dominate

- methanotrophy in aerobic and anaerobic layers of boreal lake waters. *Aquat Microb Ecol* 81:257–276. <https://doi.org/10.3354/ame01874>
14. Martinez-Cruz K, Leewis M-C, Herriott IC, Sepulveda-Jauregui A, Anthony KW, Thalasso F, Leigh MB. 2017. Anaerobic oxidation of methane by aerobic methanotrophs in sub-Arctic lake sediments. *Sci Total Environ* 607:23–31. <https://doi.org/10.1016/j.scitotenv.2017.06.187>
 15. Oswald K, Milucka J, Brand A, Hach P, Littmann S, Wehrli B, Kuypers MMM, Schubert CJ. 2016. Aerobic gammaproteobacterial methanotrophs mitigate methane emissions from oxic and anoxic lake waters. *Limnol Oceanogr* 61:S101–S118. <https://doi.org/10.1002/lno.10312>
 16. Oswald K, Graf JS, Littmann S, Tienken D, Brand A, Wehrli B, Albertsen M, Daims H, Wagner M, Kuypers MMM, Schubert CJ, Milucka J. 2017. Crenothrix are major methane consumers in stratified lakes. *ISME J* 11:2124–2140. <https://doi.org/10.1038/ismej.2017.77>
 17. Zheng Y, Wang H, Liu Y, Zhu B, Li J, Yang Y, Qin W, Chen L, Wu X, Chistoserdova L, Zhao F. 2020. Methane-dependent mineral reduction by aerobic methanotrophs under hypoxia. *Environ Sci Technol Lett* 7:606–612. <https://doi.org/10.1021/acs.estlett.0c00436>
 18. Kalyuzhnaya MG, Yang S, Rozova ON, Smalley NE, Clubb J, Lamb A, Gowda GAN, Raftery D, Fu Y, Bringel F, Vuilleumier S, Beck DAC, Trotsenko YA, Khmelenina VN, Lidstrom ME. 2013. Highly efficient methane biocatalysis revealed in a methanotrophic bacterium. *Nat Commun* 4:2785–2785. <https://doi.org/10.1038/ncomms3785>
 19. Khanongnuch R, Mangayil R, Svenning MM, Rissanen AJ. 2022. Characterization and genome analysis of a psychrophilic methanotroph representing a ubiquitous *Methylobacter* spp. cluster in boreal Lake ecosystems. *ISME Commun* 2:85. <https://doi.org/10.1038/s43705-022-00172-x>
 20. D'Ambrosio SL, Harrison JA. 2021. Methanogenesis exceeds CH₄ consumption in eutrophic lake sediments. *Limnol Oceanogr Lett* 6:173–181. <https://doi.org/10.1002/lol2.10192>
 21. Han X, Schubert CJ, Fiskal A, Dubois N, Lever MA. 2020. Eutrophication as a driver of microbial community structure in lake sediments. *Environ Microbiol* 22:3446–3462. <https://doi.org/10.1111/1462-2920.15115>
 22. van Grinsven S, Meier DV, Michel A, Han X, Schubert CJ, Lever MA. 2022. Redox zone and trophic state as drivers of methane-oxidizing bacterial abundance and community structure in lake sediments. *Front Environ Sci* 10:857358. <https://doi.org/10.3389/fenvs.2022.857358>
 23. Yang Y, Chen J, Tong T, Xie S, Liu Y. 2020. Influences of eutrophication on methanogenesis pathways and methanogenic microbial community structures in freshwater lakes. *Environ Pollut* 260:114106. <https://doi.org/10.1016/j.envpol.2020.114106>
 24. Deutzmann JS, Stief P, Brandes J, Schink B. 2014. Anaerobic methane oxidation coupled to denitrification is the dominant methane sink in a deep lake. *Proc Natl Acad Sci U S A* 111:18273–18278. <https://doi.org/10.1073/pnas.1411617111>
 25. He Z, Xu S, Zhao Y, Pan X. 2019. Methane emissions from aqueous sediments are influenced by complex interactions among microbes and environmental factors: a modeling study. *Water Res* 166:115086. <https://doi.org/10.1016/j.watres.2019.115086>
 26. Bagnoud A, Pramateftaki P, Bogard MJ, Battin TJ, Peter H. 2020. Microbial ecology of methanotrophy in streams along a gradient of CH₄ availability. *Front Microbiol* 11:771. <https://doi.org/10.3389/fmicb.2020.00771>
 27. Webster KD, Schimmelmann A, Drobniak A, Mastalerz M, Rosales Lagarde L, Boston PJ, Lennon JT. 2022. Diversity and composition of methanotroph communities in caves. *Microbiol Spectr* 10:e0156621. <https://doi.org/10.1128/spectrum.01566-21>
 28. Guerrero-Cruz S, Stultiens K, van Kessel MAHJ, Versantvoort W, Jetten MSM, Op den Camp HJM, Kartal B. 2019. Key physiology of a nitrite-dependent methane-oxidizing enrichment culture. *Appl Environ Microbiol* 85:e00124-19. <https://doi.org/10.1128/AEM.00124-19>
 29. Goñi MA, Teixeira MJ, Perkey DW. 2003. Sources and distribution of organic matter in a river-dominated estuary (Winyah Bay, SC, USA). *Estuarine, Coastal and Shelf Science* 57:1023–1048. [https://doi.org/10.1016/S0272-7714\(03\)00008-8](https://doi.org/10.1016/S0272-7714(03)00008-8)
 30. Hedges JI, Clark WA, Come GL. 1988. Organic matter sources to the water column and surficial sediments of a marine bay. *Limnol Oceanogr* 33:1116–1136. <https://doi.org/10.4319/lno.1988.33.5.1116>
 31. Arndt S, Jørgensen BB, LaRowe DE, Middelburg JJ, Pancost RD, Regnier P. 2013. Quantifying the degradation of organic matter in marine sediments: a review and synthesis. *Earth-Science Reviews* 123:53–86. <https://doi.org/10.1016/j.earscirev.2013.02.008>
 32. Meyers PA, Teranes JL. 2001. Sediment organic matter, p 239–269. In Last WM, JP Smol (ed), *Tracking environmental change using Lake sediments: physical and geochemical methods*. Springer Netherlands, Dordrecht. <https://doi.org/10.1007/0-306-47670-3>
 33. Salonen K. 1981. The ecosystem of the oligotrophic Lake Pääjärvi 2. Bacterioplankton. *SIL Proceedings, 1922-2010* 21:448–453. <https://doi.org/10.1080/03680770.1980.11897021>
 34. Jäntti H, Jilbert T, Aalto SL, Simojoki A, Mangayil R, Peura S, Rissanen AJ. 2022. The role of organic matter and microbial community controlling nitrate reduction under elevated ferrous iron concentrations in boreal lake sediments. *Hydrobiologia* 849:2145–2160. <https://doi.org/10.1007/s10750-022-04858-0>
 35. Rissanen A, Tirola M, Hietanen S, Ojala A. 2013. Interlake variation and environmental controls of denitrification across different geographical scales. *Aquat Microb Ecol* 69:1–16. <https://doi.org/10.3354/ame01619>
 36. Arvola L, Kankaala P, Tulonen T, Ojala A. 1996. Effects of phosphorus and allochthonous humic matter enrichment on metabolic processes and community structure of plankton in a boreal lake (Lake Pääjärvi). *Can J Fish Aquat Sci* 53:1646–1662. <https://doi.org/10.1139/f96-083>
 37. Jilbert T, Jokinen S, Saarinen T, Mattus-Kumpunen U, Simojoki A, Saarni S, Salminen S, Niemistö J, Horppila J. 2020. Impacts of a deep reactive layer on sedimentary phosphorus dynamics in a boreal lake recovering from eutrophication. *Hydrobiologia* 847:4401–4423. <https://doi.org/10.1007/s10750-020-04289-9>
 38. Boudreau BP. 1997. Diagenetic models and their implementation. Springer, Berlin, Heidelberg. <https://doi.org/10.1007/978-3-642-60421-8>
 39. Pajunen H. 2000. Carbon in Finnish Lake sediments. special paper 29, geological survey of Finland. Geological Survey of Finland, Espoo, Finland.
 40. Kortelainen P, Pajunen H, Rantakari M, Saarnisto M. 2004. A large carbon pool and small sink in boreal Holocene lake sediments. *Glob Chang Biol* 10:1648–1653. <https://doi.org/10.1111/j.1365-2486.2004.00848.x>
 41. Kortelainen P, Saukkonen S. 1995. Organic vs. minerogenic acidity in headwater streams in Finland. *Water Air Soil Pollut* 85:559–564. <https://doi.org/10.1007/BF00476888>
 42. Salmela M, Lehtinen T, Efimova E, Santala S, Mangayil R. 2018. Metabolic pairing of aerobic and anaerobic production in a one-pot batch cultivation. *Biotechnol Biofuels* 11:187. <https://doi.org/10.1186/s13068-018-1186-9>
 43. Apprill A, McNally S, Parsons R, Weber L. 2015. Minor revision to V4 region SSU rRNA 806R gene primer greatly increases detection of SAR11 Bacterioplankton. *Aquat Microb Ecol* 75:129–137. <https://doi.org/10.3354/ame01753>
 44. Parada AE, Needham DM, Fuhrman JA. 2016. Every base matters: assessing small subunit rRNA primers for marine microbiomes with mock communities, time series and global field samples. *Environ Microbiol* 18:1403–1414. <https://doi.org/10.1111/1462-2920.13023>
 45. Myllykangas J-P, Rissanen AJ, Hietanen S, Jilbert T. 2020. Influence of electron acceptor availability and microbial community structure on sedimentary methane oxidation in a boreal estuary. *Biogeochemistry* 148:291–309. <https://doi.org/10.1007/s10533-020-00660-z>
 46. Chowdhury TR, Dick RP. 2013. Ecology of aerobic methanotrophs in controlling methane fluxes from wetlands. *Applied Soil Ecology* 65:8–22. <https://doi.org/10.1016/j.apsoil.2012.12.014>
 47. Knittel K, Boetius A. 2009. Anaerobic oxidation of methane: progress with an unknown process. *Annu Rev Microbiol* 63:311–334. <https://doi.org/10.1146/annurev.micro.61.080706.093130>
 48. Nazaries L, Murrell JC, Millard P, Baggs L, Singh BK. 2013. Methane, microbes and models: fundamental understanding of the soil methane cycle for future predictions. *Environ Microbiol* 15:2395–2417. <https://doi.org/10.1111/1462-2920.12149>
 49. Huttunen JT, Väisänen T, Hellsten S, Martikainen PJ. 2006. Methane fluxes at the sediment-water interface in some boreal lakes and reservoirs. *Boreal Environment Research* 11:27–34. <https://doi.org/https://www.borenv.net/BER/archive/pdfs/ber11/ber11-027.pdf>
 50. Delgado-Baquerizo M, Reich PB, Khachane AN, Campbell CD, Thomas N, Freitag TE, Abu Al-Soud W, Sørensen S, Bardgett RD, Singh BK. 2017. It is elemental: soil nutrient stoichiometry drives bacterial diversity. *Environ Microbiol* 19:1176–1188. <https://doi.org/10.1111/1462-2920.13642>

51. Hooper DU, Bignell DE, Brown VK, Brussard L, Dangerfield JM, Wall DH, Wardle DA, Coleman DC, Giller KE, Lavelle P, Van Der WH, De Ruiter PC, Rusek J, Silver WL, Tiedje JM, Wolters V. 2000. Interactions between aboveground and belowground Biodiversity in terrestrial ecosystems: patterns, mechanisms, and feedbacks: we assess the evidence for correlation between aboveground and belowground diversity and conclude that a variety of mechanisms could lead to positive, negative, or no relationship—depending on the strength and type of interactions among species. *BioScience* 50:1049–1061. [https://doi.org/10.1641/0006-3568\(2000\)050\[1049:IBAABB\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2000)050[1049:IBAABB]2.0.CO;2)
52. Rissanen AJ, Karvinen A, Nykänen H, Peura S, Tirola M, Mäki A, Kankaala P. 2017. Effects of alternative electron acceptors on the activity and community structure of methane-producing and consuming microbes in the sediments of two shallow boreal lakes. *FEMS Microbiol Ecol* 93. <https://doi.org/10.1093/femsec/fix078>
53. Bai Y-N, Wang X-N, Wu J, Lu Y-Z, Fu L, Zhang F, Lau T-C, Zeng RJ. 2019. Humic substances as electron acceptors for anaerobic oxidation of methane driven by ANME-2D. *Water Res* 164:114935. <https://doi.org/10.1016/j.watres.2019.114935>
54. Su G, Zopfi J, Yao H, Steinle L, Niemann H, Lehmann MF. 2020. Manganese/iron-supported sulfate-dependent anaerobic oxidation of methane by archaea in lake sediments. *Limnol Oceanogr* 65:863–875. <https://doi.org/10.1002/lno.11354>
55. Timmers PHA, Welte CU, Koehorst JJ, Plugge CM, Jetten MSM, Stams AJM. 2017. Reverse methanogenesis and respiration in methanotrophic archaea. *Archaea* 2017:1654237. <https://doi.org/10.1155/2017/1654237>
56. Weber HS, Habicht KS, Thamdrup B. 2017. Anaerobic methanotrophic archaea of the ANME-2D cluster are active in a low-sulfate, iron-rich freshwater sediment. *Front Microbiol* 8:619. <https://doi.org/10.3389/fmicb.2017.00619>
57. Speth DR, Orphan VJ. 2018. Metabolic marker gene mining provides insight in global *mcrA* diversity and, coupled with targeted genome reconstruction, sheds further light on metabolic potential of the *Methanomassiliicoccales* PeerJ 6:e5614. <https://doi.org/10.7717/peerj.5614>
58. Iino T, Tamaki H, Tamazawa S, Ueno Y, Ohkuma M, Suzuki K-I, Igarashi Y, Haruta S. 2013. *Candidatus* methanogranum caenicola: a novel methanogen from the anaerobic digested sludge, and proposal of *Methanomassiliicocaceae* fam. nov. and *Methanomassiliicoccales* ord. nov., for a Methanogenic lineage of the class *Thermoplasmata*. *Microbes Environ* 28:244–250. <https://doi.org/10.1264/jsme2.me12189>