

JYX



This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): ALICE Collaboration

Title: Elliptic flow of charged particles at midrapidity relative to the spectator plane in Pb–Pb and Xe–Xe collisions

Year: 2023

Version: Published version

Copyright: © 2022 The Author(s). Published by Elsevier B.V. Funded by SCOAP3.

Rights: CC BY 4.0

Rights url: <https://creativecommons.org/licenses/by/4.0/>

Please cite the original version:

ALICE Collaboration. (2023). Elliptic flow of charged particles at midrapidity relative to the spectator plane in Pb–Pb and Xe–Xe collisions. *Physics Letters B*, 846, Article 137453. <https://doi.org/10.1016/j.physletb.2022.137453>



Elliptic flow of charged particles at midrapidity relative to the spectator plane in Pb–Pb and Xe–Xe collisions

ALICE Collaboration*

ARTICLE INFO

Article history:

Received 5 May 2022

Received in revised form 5 September 2022

Accepted 10 September 2022

Available online 19 September 2022

Editor: M. Doser

Dataset link: <https://www.hepdata.net/record/ins2070420>

ABSTRACT

Measurements of the elliptic flow coefficient relative to the collision plane defined by the spectator neutrons $v_2\{\Psi_{SP}\}$ in collisions of Pb ions at center-of-mass energy per nucleon–nucleon pair $\sqrt{s_{NN}} = 2.76$ TeV and Xe ions at $\sqrt{s_{NN}} = 5.44$ TeV are reported. The results are presented for charged particles produced at midrapidity as a function of centrality and transverse momentum for the 5–70% and 0.2–6 GeV/c ranges, respectively. The ratio between $v_2\{\Psi_{SP}\}$ and the elliptic flow coefficient relative to the participant plane $v_2\{4\}$, estimated using four-particle correlations, deviates by up to 20% from unity depending on centrality. This observation differs strongly from the magnitude of the corresponding eccentricity ratios predicted by the T_RENTo and the elliptic power models of initial state fluctuations that are tuned to describe the participant plane anisotropies. The differences can be interpreted as a decorrelation of the neutron spectator plane and the reaction plane because of fragmentation of the remnants from the colliding nuclei, which points to an incompleteness of current models describing the initial state fluctuations. A significant transverse momentum dependence of the ratio $v_2\{\Psi_{SP}\}/v_2\{4\}$ is observed in all but the most central collisions, which may help to understand whether momentum anisotropies at low and intermediate transverse momentum have a common origin in initial state fluctuations. The ratios of $v_2\{\Psi_{SP}\}$ and $v_2\{4\}$ to the corresponding initial state eccentricities for Xe–Xe and Pb–Pb collisions at similar initial entropy density show a difference of $(7.0 \pm 0.9)\%$ with an additional variation of +1.8% when including RHIC data in the T_RENTo parameter extraction. These observations provide new experimental constraints for viscous effects in the hydrodynamic modeling of the expanding quark–gluon plasma produced in heavy-ion collisions at the LHC.

© 2022 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

Determining the properties of deconfined quark–gluon matter, called quark–gluon plasma (QGP), is the goal of the heavy-ion program at the Large Hadron Collider (LHC). Previous studies of data collected at the Brookhaven Relativistic Heavy-Ion Collider (RHIC) and the LHC have shown that the QGP behaves like a liquid with very small specific shear and bulk viscosities [1,2]. In heavy-ion collisions, these properties are encoded in the collective expansion of the strongly interacting QGP. Measurements of this collective behavior, in particular the anisotropic flow driven by the spatial anisotropy of the shape of the overlap region of the colliding nuclei, can be used to infer the QGP transport properties. The anisotropic flow is quantified by the coefficients v_n of a Fourier decomposition of the momentum anisotropy of emitted particles relative to the collision symmetry planes with angles Ψ_n of the harmonic n . The dominant coefficient in off-center collision of heavy ions is the elliptic flow v_2 . The shape of the initial energy density distribution in the overlap region of the nuclei (participant

zone) fluctuates from collision to collision due to the motion of nucleons and the quantum mechanical nature of the nucleus–nucleus interaction, which is typically modeled by spatial position fluctuations of the interacting nucleons. This was initially discussed in the context of the elliptic flow fluctuations [3–5] and later confirmed by observations [6–10] of significant non-zero triangular flow v_3 and higher flow harmonics. A detailed picture of fluctuations has already been inferred from the comparison of initial state models to measurements of flow fluctuations relative to the participant symmetry planes [8,11–28]. Additional information about the pattern of initial state fluctuations can be obtained from measurements using spectator nucleons, which decouple very fast from the participant zone.

Fig. 1 sketches the geometry of a non-central heavy-ion collision, divided into the regions of the participant nucleons (i.e. those encountering strong interactions) and the deflected spectator nucleons of the target (T) and projectile (P) nuclei. The geometrical reaction plane, denoted by its angle Ψ_{RP} in Fig. 1a, is spanned by the impact parameter, pointing in the direction of the x_{RP} axis, and the movement direction of the colliding nuclei, which is indicated

* E-mail address: alice-publications@cern.ch.




Fig. 1. A sketch of the geometry of a non-central heavy-ion collision in the (a) transverse (x_{RP}, y) and (b) reaction (x_{RP}, z) planes. The x_{RP} axis points along the direction of the impact parameter given by the distance between the centers of the colliding nuclei, while the z axis is oriented along the direction of the colliding nuclei. The full (open) circles represent the protons (neutrons) from projectile (red) and target (blue) nuclei. The participant nucleons are shown in purple. In panel (a), the arrows indicate the eccentricity vectors (ϵ_n is shown as example) and corresponding reaction (Ψ_{RP}), projectile/target spectator ($\Psi_{SP}^{P,T}$), and participant (multiple $\Psi_n, n \geq 1$) plane angles. Panel (b) shows the outward deflection, indicated by curved arrows, of the spectators and fragmentation products of the projectile ($\eta > 0$) and target ($\eta < 0$) recoil nuclei, such as protons (p), neutrons (n), and nuclei (deuteron, helium, etc.).

by the z axis. The fluctuating shape of the energy density distribution of the collision can be characterized by the eccentricities ϵ_n for $n > 1$

$$\epsilon_n e^{in\Psi_n} = -\frac{\int r^n e^{in\varphi} \rho(r, \varphi) r dr d\varphi}{\int r^n \rho(r, \varphi) r dr d\varphi}, \quad (1)$$

where $\rho(r, \varphi)$ is the initial energy or entropy density near midrapidity and (r, φ) are the polar coordinates. Eq. (1) for $n = 2$ gives the standard formula for the ellipticity [29,30]. In the absence of fluctuations, the azimuthal symmetry plane of the collision is given by the reaction plane. In the presence of fluctuations, the collision symmetry is not described by a single reaction plane, but by multiple participant planes with angles Ψ_n and spectator planes with angles Ψ_{SP}^T and Ψ_{SP}^P . The spectator planes are spanned by the beam direction and the deflection of the spectators, which is determined by the sum of the momenta of the spectator nucleons and fragments. For small fluctuations of the spectator deflection, the target and projectile spectator planes share a common symmetry plane with angle Ψ_{SP} . The sketch is oriented along the reaction plane x_{RP} (i.e. $\Psi_{RP} = 0$), while in the experiment the orientation of the collision relative to the laboratory frame fluctuates from one collision to another. This causes the reaction plane angle to be different from zero.

Fig. 1b illustrates a longitudinal view of the early time of a heavy-ion collision, in particular the deflection of the spectators. The fragmentation of the nuclear recoil is a complex process [31] involving the release of protons, neutrons, and other nuclei, which causes a decorrelation of the reaction plane and spectator planes. Additionally, a plane defined only by a subset of the spectator fragments (e.g. only neutrons or protons) can be different from the overall spectator plane orientation illustrated in Fig. 1a. This difference depends on the collision centrality due to the balance between the energy carried by free nucleons and nuclear fragments.

The deflection of the spectators along the spectator planes at LHC collision energies, as illustrated by the red and blue arrows in Fig. 1a, was shown to be outward from the nuclear overlap region, based on the measured directed flow relative to the neutron spectator plane and hydrodynamic model calculations [32]. From measurements of the neutron deflection at the LHC, the transverse momentum exchange between the participants and the spectators has been estimated to be in the order of 20 MeV/c per nucleon in midcentral collisions [33]. This is about an order of magnitude smaller than the width of the transverse momentum distribution

of emitted spectators, which is proportional [34,35] to the Fermi momentum of 265 MeV/c [36] for lead nuclei.

Due to the geometry of the collision, the direction in which the spectators are deflected is believed to be strongly correlated to the reaction plane. In collisions with large impact parameters, the smaller number of sources of particle production (number of participants) results in a stronger decorrelation of the participant symmetry plane and the reaction plane. Measurements of the directed flow using the deflection of spectator neutrons showed a small but non-zero decorrelation between the direction of the neutron deflection and the reaction plane orientation, in addition to a relative decorrelation of the neutron spectator planes originating from the projectile and target nuclei, respectively [33]. A strong correlation between the participant and spectator planes is expected in midcentral collisions because of the dominance of the elliptic flow [37]. Together with measurements relative to the participant planes, a measurement relative to the spectator deflection provides unique information about spatial orientation of the participant zone relative to the geometrical orientation of the colliding nuclei [33].

Hydrodynamic model calculations with small viscosities indicate an approximately linear response to the initial eccentricities for elliptic and triangular flow [38]. In ideal hydrodynamics, the elliptic flow v_2 is proportional to the initial eccentricity [39]

$$v_2 \propto \kappa_2 \epsilon_2. \quad (2)$$

This has been demonstrated by the comparison of measurements relative to the participant plane with eccentricities from different initial state model calculations [25]. The scaling coefficient κ_2 is related to the initial entropy density. In ideal hydrodynamics, it is estimated from the final state multiplicity dN_{ch}/dy per unit of the overlap area S [40–42]

$$\frac{v_2}{\epsilon_2} \propto \frac{1}{S} \frac{dN_{ch}}{dy}. \quad (3)$$

Following Ref. [25], the area S is defined as $S = 4\pi\sigma_x\sigma_y$, where σ_x is the spatial width of the participant zone along the reaction plane direction (x_{RP}) and σ_y is along the transverse direction (y). This scaling can be tested by comparing different collision systems and their dependence on centrality or center-of-mass energy. Such a scaling is broken by viscous effects, which depend on the magnitude of the temperature-dependent specific shear and bulk

viscosities [41]. Even for an ideal fluid, the scaling can be broken during the final stages of the evolution before freeze-out [41].

One of the largest uncertainties in the description of heavy-ion collisions is the pattern of fluctuations in the initial state that results in large uncertainties of the extracted QGP transport properties [43,44]. Among the observables that quantify the effects of initial state fluctuations are the cumulants [29], which can be used to decompose both the spatial and momentum anisotropies. The cumulants of the eccentricity distribution describe the shape of the initial state fluctuations. There are different models describing this initial state and its fluctuations. One of the simplest analytical descriptions is the Bessel-Gaussian model (BGM) [45] in which many sources of entropy production result in a two-dimensional Gaussian shape of fluctuations, where all higher-order ($m \geq 4$) eccentricity cumulants, $\varepsilon_2\{m\}$, are degenerate and equal to the reaction plane eccentricity

$$\varepsilon_2\{2\} > \varepsilon_2\{4\} = \varepsilon_{RP}. \quad (4)$$

The BGM is only applicable to small fluctuations. The elliptic-power model (EPM) is a more general analytical model, which is also applicable for the description of large fluctuations [11]. In the EPM, the degeneracy of higher-order cumulants is broken and therefore

$$\varepsilon_2\{2\} > \varepsilon_2\{4\} > \varepsilon_{RP}. \quad (5)$$

There are a number of Monte Carlo models of the initial state that either implement a specific physical mechanism of pre-equilibrium entropy production, for example IP-Glasma [46], or do not assume a specific mechanism, such as TrENTo [12]. In TrENTo, the entropy production in the participant zone is parameterized by the transverse density of the participating nucleons, i.e. the participant thickness. Each of these models is characterized by specific signatures of the eccentricity cumulants and their relationships. The importance of subnucleonic degrees of freedom for modeling fluctuations in central collisions has been highlighted in Ref. [47].

One can probe the initial state eccentricities by comparing the ratios of eccentricity cumulants defined by Eq. (1) to equivalent flow observables, if the fluid response is linear as given by Eq. (2). These observables can be constructed through angular correlations of the emitted particles and the symmetry planes

$$v_2\{\Psi_{SP}\} = \langle\langle \cos 2(\varphi_1 - \Psi_{SP}) \rangle\rangle, \quad (6)$$

$$v_2\{2, |\Delta\eta|\} = \sqrt{\langle\langle \cos 2(\varphi_1 - \varphi_2) \rangle\rangle} = \langle v_2^2 \rangle^{\frac{1}{2}}, \quad (7)$$

$$\begin{aligned} v_2\{4\} &= \left[2\langle\langle \cos 2(\varphi_1 - \varphi_2) \rangle\rangle^2 \right. \\ &\quad \left. - \langle\langle \cos 2(\varphi_1 - \varphi_2 + \varphi_3 - \varphi_4) \rangle\rangle \right]^{\frac{1}{4}} \\ &= \left[2\langle v_2^2 \rangle^2 - \langle v_4^2 \rangle \right]^{\frac{1}{4}}. \end{aligned} \quad (8)$$

Here φ_i ($i = 1 \dots 4$) are the azimuthal angles of the produced particles, $\langle \dots \rangle$ is the average over all collisions, and $\langle\langle \dots \rangle\rangle$ is the average over all particles in a given transverse momentum window in all collisions. The $v_2\{\Psi_{SP}\}$ is the elliptic flow measured relative to the spectator plane with angle Ψ_{SP} , where fluctuations of the spectator symmetry planes are assumed to be small. The $v_2\{2\}$ and $v_2\{4\}$ are estimators of elliptic flow coefficients whose equations, including p_T -dependence, are derived from the two- and four-particle cumulants [48,49]. The variable $|\Delta\eta|$ in Eq. (7) refers to the minimum pseudorapidity (η) separation between the two correlated particles. The observables $v_2\{\Psi_{SP}\}$, $v_2\{4\}$, and $v_2\{2, |\Delta\eta|\}$ may deviate from $v_2\{\Psi_{RP}\}$, which is the unmeasurable momentum anisotropy relative to the reaction plane. The comparison of ratios

of the measured anisotropic flow coefficients to their respective eccentricities is instrumental for identifying the most realistic model of the initial state.

The transverse momentum dependence of the flow coefficients v_n allows one to study if there is a common origin of anisotropy at different transverse momentum scales due to initial state eccentricity fluctuations. The main source of anisotropy for soft ($p_T < 3$ GeV/c) particle production is the collective expansion of the QGP [38]. At significantly larger p_T , the dominant contribution to v_n is the path length dependent energy loss of the jet propagating through the azimuthally asymmetric medium oriented along Ψ_n [50]. A significant transverse momentum dependence of flow fluctuations relative to the participant planes is observed [50] in central collisions and within large experimental uncertainties in midcentral collisions. Given that the spectators only probe the early time evolution of the collision and provide sensitivity to a different symmetry plane, the transverse momentum dependence of the ratio $v_2\{\Psi_{SP}\}/v_2\{4\}$ can be used as an additional observable to investigate the origin of flow fluctuations.

An accurate determination of the flow coefficients relative to the spectator plane, in particular of the elliptic flow $v_2\{\Psi_{SP}\}$, is also crucial for the search of the chiral magnetic effect (CME) [51, 52]. Because the magnetic field orientation is correlated more strongly to the spectator deflection than to the participant plane orientation, the measurement relative to both planes allows the separation of the CME signal from the background correlations coupled to the elliptic flow [52–57].

In this letter, the charged particle elliptic flow relative to the neutron spectator deflection, $v_2\{\Psi_{SP}\}$, and to the participant plane, quantified by $v_2\{2, |\Delta\eta|\} > 1$ and $v_2\{4\}$, is reported as a function of centrality and transverse momentum for collisions of Pb ions at a center-of-mass energy per nucleon–nucleon pair of $\sqrt{s_{NN}} = 2.76$ TeV and Xe ions at $\sqrt{s_{NN}} = 5.44$ TeV. The comparison with different initial state models, such as BGM, EPM, and TrENTo, and the scaling properties of v_2/ε_2 are also presented.

ALICE is a dedicated heavy-ion experiment at the LHC. A detailed description of the detector and its performance can be found elsewhere [58,59]. The trajectories of the charged particles are reconstructed using the Inner Tracking System (ITS) and the Time Projection Chamber (TPC) [60,61]. The neutron spectator deflection is estimated using the neutron Zero Degree Calorimeters (ZDC) [62]. The ZDC consist of two detectors with a 2×2 transverse segmentation. During the Pb–Pb (Xe–Xe) data-taking, the ZDC were placed 114 m (112.5 m) away from the collision point on both sides of the experiment, named A and C side, covering the pseudorapidity interval $|\eta| > 8.78$ (8.77). The collision centrality is reconstructed using the energy deposition in the forward V0 scintillator arrays [63].

About 10 (1) million of minimum bias [59] Pb–Pb (Xe–Xe) collisions collected in the year 2010 (2017) are analyzed. This corresponds to collisions in the 5–70% centrality range and a reconstructed primary collision vertex located within ± 10 cm along the beam direction from the nominal interaction point. Outside of this centrality range, the deflection of the spectators cannot be reliably reconstructed because of the small number of detected neutron spectators in central collisions and the large fraction of spectator neutrons that are not detected by the ZDC in peripheral collisions because they are bound in charged fragments and thus deflected by the LHC magnets.

The charged particles are selected within the range of pseudorapidity $|\eta| < 0.8$ and transverse momentum $p_T > 0.2$ GeV/c. The tracks are reconstructed using the combined information from the ITS and the TPC detectors. A minimum of 70 TPC space points (out of a maximum of 159) and a minimum of two ITS hits, with at least one in one of the two innermost layers, are required for all tracks. A reduced χ^2 per TPC space point (ITS hit) in the range

0.1–4 (less than 36) is required. Only tracks with a distance of closest approach to the primary vertex position smaller than 3.2 cm in the beam direction and within a radius of 2.4 cm in the transverse plane are considered.

The observables in Eqs. (6) to (8), are calculated using flow vectors q_n^K of harmonic n in each collision as

$$q_n^K = x_n^K + iy_n^K = \frac{\sum_{j \in K} (w_j)^p e^{in\varphi_j}}{\sum_{j \in K} (w_j)^p}, \quad (9)$$

where K denotes either the reconstructed tracks $K = T$ or the two ZDC subdetectors $K = A, C$.

For the second harmonic flow vector of the reconstructed tracks q_2^T , the w_j and φ_j are the weight and azimuthal angle of the j -th track, respectively. The weights in the q_2^T calculation are defined by the inverse product $w = 1/(\varepsilon dN/d\varphi)$ of the p_T -dependent reconstruction efficiency ε and the measured non-uniform azimuthal angle distribution $dN/d\varphi$ of the charged particles. The $dN/d\varphi$ is calculated as a function of the pseudorapidity of the particles and the primary vertex position along the beam direction. The impact of the correction for the non-uniform azimuthal acceptance of the tracks is negligible. The reconstruction efficiency correction is derived from a Monte Carlo simulation using the HIJING [64] heavy-ion event generator to simulate collisions and GEANT3 [65] to model particle transport through the ALICE detector material. It is applied to all measurements of the p_T -integrated flow coefficients. The power p in Eq. (9) corresponds to the correction of the non-uniform azimuthal acceptance and reconstruction efficiency in the multi-particle cumulant method [66]. It only differs from unity for q_2^T . The $v_2\{2, |\Delta\eta| > 1\}$ and $v_2\{4\}$ are calculated from flow vectors defined in Eq. (9) using the multi-particle cumulant approach [66]. In the measurement of $v_2\{2, |\Delta\eta| > 1\}$, a pseudorapidity separation of $|\Delta\eta| > 1$ is applied as it largely suppresses few-particle correlations (so called non-flow), such as those originating from jets or resonance decays [67].

For the first harmonic flow vectors of the ZDC, q_1^A and q_1^C , φ_j corresponds to the angle of the center of the j -th calorimeter segment and w_j to its measured energy deposition. The indices A and C stand for the subdetectors located on either side of the interaction point. The neutron spectator deflection is estimated with these flow vectors. The beam configuration can result in an average shift of the spectator neutron distributions in the transverse plane of the ZDC that are corrected by a data-driven procedure (re-centering)

$$q_n' = q_n - \langle q_n \rangle. \quad (10)$$

To account for changing beam conditions, the re-centering correction is calculated as a function of the centrality, and the three dimensional position of the collision vertex for each data-taking period. The elliptic flow $v_2\{\Psi_{SP}\}$ is calculated from an average of three independent estimates using the scalar product method with mixed harmonics [68]

$$v_2\{\Psi_{SP}\} = \frac{2}{3} \left(\frac{\langle x_2^T x_1^A x_1^C \rangle}{\langle x_1^A x_1^C \rangle} - \frac{\langle x_2^T y_1^A y_1^C \rangle}{\langle y_1^A y_1^C \rangle} \right) + \sqrt{\frac{\langle y_2^T x_1^A y_1^C \rangle \langle y_2^T y_1^A x_1^C \rangle}{\langle x_1^A x_1^C \rangle \langle y_1^A y_1^C \rangle}}, \quad (11)$$

which corresponds to a measurement of elliptic flow relative to the common spectator plane Ψ_{SP} . The numerical coefficient (2/3) arises from (1/3) which corresponds to an average of the three independent terms in Eq. (11) and an additional factor two

from an expectation that correlators in the numerators of individual terms gives $(1/4)v_2\{\Psi_{SP}\}R_1^2\{ZDC\}$ and in the denominators $(1/2)R_1^2\{ZDC\}$, where $R_1\{ZDC\}$ is a convolution of the directed flow of spectators with the ZDC response.

The systematic uncertainties are evaluated from variations of the analysis procedure. The total systematic uncertainty is given by the square root of the sum in quadrature of the differences in the results obtained from each significant variation. A variation is considered significant, if $\Delta R/\Delta\sigma > 1$ [69], where ΔR is the difference between results from a given variation and the default analysis procedure. The variable $\Delta\sigma$ is the square root of the difference between the corresponding squared statistical uncertainties.

The description of systematic uncertainties related to measurements of $v_2\{2, |\Delta\eta| > 1\}$ and $v_2\{4\}$ is presented in Ref. [70]. The details of the systematic uncertainty evaluation for $v_2\{\Psi_{SP}\}$ are described below. The largest source of systematic uncertainty is due to spurious correlations which are not corrected by the re-centering procedure. These affect the denominators in Eq. (11), which are determined by the diagonal correlations $\langle x_1^A x_1^C \rangle$ and $\langle y_1^A y_1^C \rangle$. The corresponding relative uncertainty, labeled as ZDC scale uncertainty, is evaluated from the off-diagonal correlations $\langle x_1^A y_1^C \rangle$ and $\langle y_1^A x_1^C \rangle$, which should not contain physical signal, divided by the diagonal correlations. This uncertainty is found to be about 4% (9%) in Pb–Pb (Xe–Xe) collisions for all centrality classes and it is assigned as a fully correlated uncertainty for $v_2\{\Psi_{SP}\}$ results. The observed negative sign of the diagonal correlations indicates a deflection of the neutron spectators in, on average, opposite directions [33]. The off-diagonal correlations may be non-zero because of the specific beam conditions during the LHC operations that may affect the distribution of neutrons impinging ZDC, such as the beam divergence at the interaction point.

The second largest contribution to the relative uncertainty comes from the differences among the three independent estimates of $v_2\{\Psi_{SP}\}$ corresponding to the three terms in Eq. (11) and amounts to lower than 2%. The uncertainty due to the implementation of the flow vector re-centering procedure, defined by Eq. (10), is evaluated by comparing to results from an iterative procedure. In the latter, each of the four iterations consists of one coarse-binned four-dimensional re-centering step followed by four fine-grained one-dimensional steps as a function of the centrality and the three dimensional position of the collision vertex. The corresponding relative variation is below 1%. Introducing an additional equalization of the average ZDC signals in the flow vector calculation in Eq. (9) amounts to a relative uncertainty less than 1%.

The systematic uncertainties related to the reconstruction of charged particle tracks and to the secondary particle contamination are evaluated by varying the track selection criteria. A relative uncertainty from changing the selection of the distance to the primary vertex of the collision as a function of particle p_T , the minimum number of TPC clusters and the reduced χ^2 per TPC cluster is about 1.5%. No significant difference between results for positively and negatively charged particles is observed. A relative uncertainty of 0.7% on the p_T -dependent track reconstruction efficiency was estimated from the difference between the results obtained with an efficiency calculation using all primary tracks and those using only tracks of primary π , K , p , μ , and e . The uncertainty due to the non-uniform acceptance correction, estimated from the difference of results with and without applying the $1/(dN/d\varphi)$ weights in Eq. (9), is not significant. An uncertainty due to the centrality determination reaching up to 2% in the most peripheral collisions is estimated from the difference of the results obtained when using an alternative centrality estimator based on the multiplicity of hits in the ITS. The effects of the TPC misalignment and the beam optics at the interaction point are estimated to be smaller than 0.5% by comparing results obtained with different magnetic field polarities of the ALICE solenoid. The impact




Fig. 2. (upper panels) Elliptic flow relative to the spectator plane, $v_2\{\Psi_{SP}\}$, and to the participant plane, $v_2\{2, |\Delta\eta| > 1\}$ and $v_2\{4\}$, as a function of centrality in Pb–Pb (left) and Xe–Xe (right) collisions. (bottom panels) Ratios of the elliptic flow $v_2\{\Psi_{SP}\}$ and $v_2\{2, |\Delta\eta| > 1\}$ to $v_2\{4\}$. The dashed (solid) lines show the eccentricity ratios of ε_{RP} ($\varepsilon_2\{2\}$) to $\varepsilon_2\{4\}$ from the elliptic power model. The corresponding eccentricity ratios for T_{RENTo} are shown as solid bands. The error bars (open boxes) indicate statistical (systematic) uncertainties. The bin-to-bin uncorrelated uncertainties and the correlated ones are combined for $v_2\{\Psi_{SP}\}$ results. For the ratio $v_2\{\Psi_{SP}\}/v_2\{4\}$, the ZDC scale uncertainties are shown separately as solid boxes centered at unity on the right side of the lower panels.

on the results of the variation of charged particle acceptance in pseudorapidity within the fiducial volume of ALICE is probed by reducing the accepted range of the position of the reconstructed primary vertex of the collisions along the beam direction from the nominal ± 10 cm to ± 8 cm. It yields no significant change of $v_2\{\Psi_{SP}\}$. For the ratios $v_2\{2, |\Delta\eta| > 1\}/v_2\{4\}$ and $v_2\{\Psi_{SP}\}/v_2\{4\}$ most of the common sources of systematic uncertainties cancel out. In addition, the bin-to-bin uncorrelated uncertainties cancel in the p_T -dependence of these ratios.

The results for the elliptic flow coefficients relative to the spectator symmetry plane, $v_2\{\Psi_{SP}\}$, and the participant plane, $v_2\{2\}$ and $v_2\{4\}$, as a function of the centrality in Pb–Pb and Xe–Xe collisions are shown in the upper panels of Fig. 2. In both Pb–Pb and Xe–Xe collisions, the maximum of $v_2\{\Psi_{SP}\}$ is located in the 35–45% centrality range with a hint of being shifted towards more peripheral collisions for smaller systems. The difference between $v_2\{2, |\Delta\eta| > 1\}$ and $v_2\{4\}$ is due to fluctuations in the participant zone, which have been extensively studied [8,71–73]. In peripheral collisions, this difference may also originate from residual few-particle correlations. In midcentral collisions, $v_2\{\Psi_{SP}\}$ and $v_2\{4\}$ are similar as expected from the BGM (see Eq. (4)).

To study the differences between the flow observables in detail, the lower panels of Fig. 2 show the centrality dependence of the ratios of $v_2\{2, |\Delta\eta| > 1\}$ and $v_2\{\Psi_{SP}\}$ to $v_2\{4\}$ as well as the ratios of the respective eccentricity cumulants calculated from the EPM and the T_{RENTo} models. The width of the bands, which represent the T_{RENTo} calculations, in Fig. 2 indicate the differences between two configurations that are tuned to different experimental data. The model in Ref. [43] uses Pb–Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV and $\sqrt{s_{NN}} = 5.02$ TeV, whereas Ref. [44] uses data of Au–Au collisions at $\sqrt{s_{NN}} = 0.2$ TeV and Pb–Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV. The deformed Woods–Saxon parametrization of Xe nuclei is adopted from Ref. [74]. The considered model calculations do not take into account the subnucleonic degrees of free-

dom. It was demonstrated in Ref. [25] based on two-particle cumulant measurements that at the LHC they only impact very central (0–5%) Pb–Pb collisions, which is outside of the reported centrality range. The EPM model parameters, which relies on significant differences between higher order cumulants, are only accessible with the larger data sample of Pb–Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV [70].

In both central and peripheral Pb–Pb collisions, the ratio $v_2\{\Psi_{SP}\}/v_2\{4\}$ strongly deviates from unity, which is the value expected from the BGM for the eccentricity ratio. In midcentral collisions, the measured ratio is smaller than in central collisions, but it is significantly larger than the BGM (unity), EPM and T_{RENTo} model calculations. The latter indicates that the spectator plane becomes decorrelated from the reaction plane and simultaneously more correlated with the participant plane. Thus $v_2\{\Psi_{SP}\}$ becomes closer to $v_2\{2, |\Delta\eta| > 1\}$, which makes the ratio $v_2\{\Psi_{SP}\}/v_2\{4\}$ above unity. Similar trends are observed in Xe–Xe collisions, however due to the limited size of the data sample the deviations from unity are not significant within the uncertainties except in the most peripheral collisions. A decorrelation in peripheral collisions is expected due to the small number of sources of particle production (number of participants), as well as due to the decreasing fraction of energy carried by the neutrons measured with the ZDC relative to the unmeasured energy of protons and other charged nuclear fragments.

In central collisions, despite the largest number of particle-producing sources, the fluctuations are also large and deviations of the $v_2\{\Psi_{SP}\}/v_2\{4\}$ ratio from unity are growing, which is in contrast to BGM expectations. The deviation from unity is also large compared to the EPM calculations, which yield a value close to unity in central collisions. This can be either due to (a) a small number of spectators emitted in these collisions which increases the spectator plane fluctuations (decorrelation) around the reaction plane and consequently increases the magnitude of $v_2\{\Psi_{SP}\}$, (b) the strengthening of the correlation between Ψ_{SP} and Ψ_2 angles,




Fig. 3. (upper panels) Elliptic flow relative to the spectator plane, $v_2\{\Psi_{SP}\}$, and to the participant plane, $v_2\{4\}$, as a function of transverse momentum in different centrality classes for Pb–Pb collisions. The linear fits to the ratio $v_2\{\Psi_{SP}\}/v_2\{4\}$ are shown as red lines for the individual centrality classes. The error bars (open boxes) indicate statistical (systematic) uncertainties. The correlated uncertainties, related to the ZDC, and bin-to-bin uncorrelated ones are combined for $v_2\{\Psi_{SP}\}$ (upper panels). For the ratio $v_2\{\Psi_{SP}\}/v_2\{4\}$ (lower panels), the correlated uncertainty is shown by the grey band at unity.

or (c) specific geometry fluctuations and/or correlated local particle production in individual participating nucleon–nucleon interactions that reduce $v_2\{4\}$. In midcentral and peripheral collisions, both models predict a small deviation from unity for $\varepsilon_{RP}/\varepsilon_2\{4\}$, which is expected for small fluctuations given by the second equality of Eq. (4) and due to the small skewness observed in Ref. [70]. The ratio of $\varepsilon_2\{2\}/\varepsilon_2\{4\}$ from the T_RENTo model shows smaller fluctuations compared to the tuned-on-data EPM calculations and the measured $v_2\{2, |\Delta\eta| > 1\}/v_2\{4\}$.

The upper panels of Fig. 3 show the dependence of the elliptic flow coefficients $v_2\{\Psi_{SP}\}$ and $v_2\{4\}$ on transverse momentum for different centrality classes in Pb–Pb collisions. The lower panels of Fig. 3 show the ratio $v_2\{\Psi_{SP}\}/v_2\{4\}$ as a function of p_T for different centrality classes. To quantify the possible dependence of $v_2\{\Psi_{SP}\}/v_2\{4\}$ on p_T , the ratio was fitted with a linear function in the range 0.2–6 GeV/c. The slopes of the linear fit in different centrality classes are: 2.31 ± 2.17 (5–10%), 1.41 ± 0.48 (10–20%), 1.48 ± 0.36 (20–30%), 1.40 ± 0.42 (30–40%), 1.74 ± 0.65 (40–50%), 4.29 ± 1.34 (50–60%) in units of $100/(\text{GeV}/c)$. There are indications for a non-flat transverse momentum dependence of $v_2\{\Psi_{SP}\}/v_2\{4\}$ with a statistical significance of about $2.5\text{--}4\sigma$ in all centrality classes except for 5–10% central collisions where it is only 1σ . These observations are important for understanding to which extent the momentum anisotropy at low and intermediate transverse momentum has a common origin from the fluctuating initial state anisotropy in the coordinate space and quantifying the effects of momentum-dependent flow magnitude and angle decorrelation [22,75].

The scaling behavior of the elliptic flow with the initial state eccentricities, as suggested by Eq. (3), is shown in Fig. 4. The top and middle panels present the ratio v_2/ε_2 as a function of the charged particle multiplicity density per unit of the nuclei overlap area $(1/S)dN_{ch}/d\eta$ for the elliptic flow coefficients $v_2\{2, |\Delta\eta| > 1\}$, $v_2\{4\}$, and $v_2\{\Psi_{SP}\}$. The area S and the corresponding eccentricities are calculated from the T_RENTo model using the configuration from Ref. [43]. The experimental data points for $dN_{ch}/d\eta$ as a function of centrality are taken from Ref. [76] for Pb–Pb and from Ref. [74] for Xe–Xe collisions. The lines represent linear fits to the ratios $v_2\{\Psi_{SP}\}/\varepsilon_{RP}$, $v_2\{2, |\Delta\eta| > 1\}/\varepsilon_2\{2\}$, and $v_2\{4\}/\varepsilon_2\{4\}$.

Deviations from the scaling behavior are quantified in the bottom panel of Fig. 4, which shows v_2/ε_2 divided by the respective linear fits of the Pb–Pb measurement. The small differences between eccentricities calculated from the two T_RENTo configurations are reflected in the height of the solid grey box on the left side. The splitting between the individual observables for a given




Fig. 4. Ratios of v_2/ε_2 as a function of $(1/S)dN_{ch}/d\eta$ in Xe–Xe (top panel) and Pb–Pb (middle panel) collisions. The linear fit functions to v_2/ε_2 in the top and middle panels are shown by lines. The bands show the uncertainty of the fit. (bottom panel) The ratio of v_2/ε_2 to the linear Pb–Pb fits shown in the middle panel. The data points of $v_2\{2, |\Delta\eta| > 1\}$, $v_2\{4\}$ are shifted by -0.1 and $+0.1$ along $(1/S)dN_{ch}/d\eta$ for better visibility. The error bars (open boxes) indicate statistical (systematic) uncertainties. The ZDC scale uncertainty, due to residual correlations in the determination of the denominators in Eq. (11), and the bin-to-bin uncorrelated one are combined for $v_2\{\Psi_{SP}\}$ (top and middle panels). For the ratio of $v_2\{\Psi_{SP}\}$ to the Pb–Pb fit (bottom panel), the ZDC scale uncertainty is separated and is shown by the solid boxes on the left side of the bottom panel. The grey box at unity in the bottom panel on the left side shows the variation depending on the T_RENTo configuration. The blue horizontal band (bottom panel) represents the relative uncertainties of the individual linear fits to the Pb–Pb data from the middle panel. The red horizontal line (bottom panel) shows a combined fit of a constant function to the Xe–Xe data and its uncertainty.

collision system may be sensitive to the chosen association of eccentricities to flow coefficients. A relative deviation of $(7.0 \pm 0.9)\%$ from the scaling given by Eq. (3) between the two collision systems is extracted from a constant line fit to the Xe–Xe data and by propagating uncertainties of the Pb–Pb data. An additional variation of $+1.8\%$ is estimated from the difference between the two $T_{\text{R}}\text{ENTo}$ configurations. It is similar for the different flow coefficients and only weakly depends on the multiplicity density or collision centrality. This collision system dependence may be sensitive to the details of the initial state models [25,77] and to viscous effects in the expansion of the QGP [41]. Together with the results on the centrality and transverse momentum dependence, shown in Figs. 2 and 3, the scaling presented in Fig. 4 provides new experimental constraints to both the initial conditions and the transport coefficients of the QGP produced in heavy-ion collisions.

In summary, the elliptic flow of charged particles at midrapidity ($|\eta| < 0.8$) relative to the neutron spectator plane $v_2\{\Psi_{\text{SP}}\}$ is measured in Pb–Pb at $\sqrt{s_{\text{NN}}} = 2.76$ TeV and Xe–Xe at $\sqrt{s_{\text{NN}}} = 5.44$ TeV collisions as a function of the centrality and as a function of the transverse momentum in different centrality classes. The measurements are compared to results of elliptic flow relative to the participant plane obtained from two and four-particle correlations. Both in central and peripheral collisions, the ratio $v_2\{\Psi_{\text{SP}}\}/v_2\{4\}$ strongly deviates from the respective ratio of the eccentricities, $\varepsilon_{\text{RP}}/\varepsilon_2\{4\}$, predicted by the $T_{\text{R}}\text{ENTo}$ and EPM models of the initial state fluctuations. The observations can be interpreted as a decorrelation of the neutron spectator plane and the reaction plane due to the fragmentation of the recoil from the colliding nuclei which indicates an incomplete description of initial state fluctuations in the models. A transverse momentum dependence of the ratio $v_2\{\Psi_{\text{SP}}\}/v_2\{4\}$ is observed with $2.5\text{--}4\sigma$ significance in all but the most central collisions. This is an important experimental input for understanding whether the momentum anisotropies at low and intermediate transverse momentum have a common origin in the initial state fluctuations. The ratios of elliptic flow coefficients to the initial state eccentricities for Xe–Xe and Pb–Pb collisions reveal a relative deviation from the scaling with the initial entropy density of $(7.0 \pm 0.9)\%$ with an additional variation of $+1.8\%$ when including RHIC data in the $T_{\text{R}}\text{ENTo}$ parameter extraction. The deviation is independent of the charged particle multiplicity density, which provides new experimental constraints for viscous effects in the expansion of the quark–gluon plasma produced in heavy-ion collisions at the LHC.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

This manuscript has associated data in a HEPData repository at: <https://www.hepdata.net/record/ins2070420>.

Acknowledgements

The ALICE Collaboration would like to thank all its engineers and technicians for their invaluable contributions to the construction of the experiment and the CERN accelerator teams for the outstanding performance of the LHC complex. The ALICE Collaboration gratefully acknowledges the resources and support provided by all Grid centres and the Worldwide LHC Computing Grid (WLCG) collaboration. The ALICE Collaboration acknowledges the following funding agencies for their support in building and running the

ALICE detector: A. I. Alikhanyan National Science Laboratory (Yerevan Physics Institute) Foundation (ANSL), State Committee of Science and World Federation of Scientists (WFS), Armenia; Austrian Academy of Sciences, Austrian Science Fund (FWF): [M 2467-N36] and Nationalstiftung für Forschung, Technologie und Entwicklung, Austria; Ministry of Communications and High Technologies, National Nuclear Research Center, Azerbaijan; Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Financiadora de Estudos e Projetos (Finep), Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP) and Universidade Federal do Rio Grande do Sul (UFRGS), Brazil; Bulgarian Ministry of Education and Science, within the National Roadmap for Research Infrastructures 2020–2027 (object CERN), Bulgaria; Ministry of Education of China (MOEC), Ministry of Science & Technology of China (MSTC) and National Natural Science Foundation of China (NSFC), China; Ministry of Science and Education and Croatian Science Foundation, Croatia; Centro de Aplicaciones Tecnológicas y Desarrollo Nuclear (CEADEN), Cubaenergía, Cuba; Ministry of Education, Youth and Sports of the Czech Republic, Czech Republic; The Danish Council for Independent Research | Natural Sciences, the Villum Fonden and Danish National Research Foundation (DNRF), Denmark; Helsinki Institute of Physics (HIP), Finland; Commissariat à l'Énergie Atomique (CEA) and Institut National de Physique Nucléaire et de Physique des Particules (IN2P3) and Centre National de la Recherche Scientifique (CNRS), France; Bundesministerium für Bildung und Forschung (BMBF) and GSI Helmholtzzentrum für Schwerionenforschung GmbH, Germany; General Secretariat for Research and Technology, Ministry of Education, Research and Religions, Greece; National Research, Development and Innovation Office, Hungary; Department of Atomic Energy Government of India (DAE), Department of Science and Technology, Government of India (DST), University Grants Commission, Government of India (UGC) and Council of Scientific and Industrial Research (CSIR), India; National Research and Innovation Agency - BRIN, Indonesia; Istituto Nazionale di Fisica Nucleare (INFN), Italy; Japanese Ministry of Education, Culture, Sports, Science and Technology (MEXT) and Japan Society for the Promotion of Science (JSPS) KAKENHI, Japan; Consejo Nacional de Ciencia (CONACYT) y Tecnología, through Fondo de Cooperación Internacional en Ciencia y Tecnología (FONCICYT) and Dirección General de Asuntos del Personal Académico (DGAPA), Mexico; Nederlandse Organisatie voor Wetenschappelijk Onderzoek (NWO), Netherlands; The Research Council of Norway, Norway; Commission on Science and Technology for Sustainable Development in the South (COMSATS), Pakistan; Pontificia Universidad Católica del Perú, Peru; Ministry of Education and Science, National Science Centre and WUT ID-UB, Poland; Korea Institute of Science and Technology Information and National Research Foundation of Korea (NRF), Republic of Korea; Ministry of Education and Scientific Research, Institute of Atomic Physics, Ministry of Research and Innovation and Institute of Atomic Physics and University Politehnica of Bucharest, Romania; Ministry of Education, Science, Research and Sport of the Slovak Republic, Slovakia; National Research Foundation of South Africa, South Africa; Swedish Research Council (VR) and Knut & Alice Wallenberg Foundation (KAW), Sweden; European Organization for Nuclear Research, Switzerland; Suranaree University of Technology (SUT), National Science and Technology Development Agency (NSTDA), Thailand Science Research and Innovation (TSRI) and National Science, Research and Innovation Fund (NSRF), Thailand; Turkish Energy, Nuclear and Mineral Research Agency (TEN-MAK), Turkey; National Academy of Sciences of Ukraine, Ukraine; Science and Technology Facilities Council (STFC), United Kingdom; National Science Foundation of the United States of America (NSF) and United States Department of Energy, Office of Nuclear Physics (DOE NP), United States of America. In addition, individual groups or members have received support from: Marie Skłodowska Curie,

Strong 2020 - Horizon 2020, European Research Council (grant nos. 824093, 896850, 950692), European Union; Academy of Finland (Center of Excellence in Quark Matter) (grant nos. 346327, 346328), Finland; Programa de Apoyos para la Superación del Personal Académico, UNAM, Mexico.

References

- [1] S. Ryu, J.F. Paquet, C. Shen, G.S. Denicol, B. Schenke, S. Jeon, C. Gale, Importance of the bulk viscosity of QCD in ultrarelativistic heavy-ion collisions, *Phys. Rev. Lett.* 115 (13) (2015) 132301, arXiv:1502.01675 [nucl-th].
- [2] B. Schenke, S. Jeon, C. Gale, Elliptic and triangular flow in event-by-event (3+1) D viscous hydrodynamics, *Phys. Rev. Lett.* 106 (2011) 042301, arXiv:1009.3244 [hep-ph].
- [3] M. Miller, R. Snellings, Eccentricity fluctuations and its possible effect on elliptic flow measurements, arXiv:nucl-ex/0312008.
- [4] PHOBOS Collaboration, B. Alver, et al., The eccentricities of flow: elliptic flow fluctuations and evidence for transverse localization in the initial state of the matter in relativistic heavy ion collisions, in: International Workshop on Hadron Physics and Property of High Baryon Density Matter, 2007, arXiv:nucl-ex/0702029.
- [5] STAR Collaboration, P. Sorensen, Elliptic flow fluctuations in Au + Au collisions at $\sqrt{s_{NN}} = 200$ GeV, *J. Phys. G* 35 (2008) 104102, arXiv:0808.0356 [nucl-ex].
- [6] B. Alver, G. Roland, Collision geometry fluctuations and triangular flow in heavy-ion collisions, *Phys. Rev. C* 81 (2010) 054905, arXiv:1003.0194 [nucl-th], Erratum: *Phys. Rev. C* 82 (2010) 039903.
- [7] PHENIX Collaboration, A. Adare, et al., Measurements of higher-order flow harmonics in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV, *Phys. Rev. Lett.* 107 (2011) 252301, arXiv:1105.3928 [nucl-ex].
- [8] ALICE Collaboration, K. Aamodt, et al., Higher harmonic anisotropic flow measurements of charged particles in Pb–Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, *Phys. Rev. Lett.* 107 (2011) 032301, arXiv:1105.3865 [nucl-ex].
- [9] STAR Collaboration, L. Adamczyk, et al., Third harmonic flow of charged particles in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV, *Phys. Rev. C* 88 (1) (2013) 014904, arXiv:1301.2187 [nucl-ex].
- [10] STAR Collaboration, L. Adamczyk, et al., Beam energy dependence of the third harmonic of azimuthal correlations in Au+Au collisions at RHIC, *Phys. Rev. Lett.* 116 (11) (2016) 112302, arXiv:1601.01999 [nucl-ex].
- [11] L. Yan, J.-Y. Ollitrault, A.M. Poskanzer, Eccentricity distributions in nucleus-nucleus collisions, *Phys. Rev. C* 90 (2) (2014) 024903, arXiv:1405.6595 [nucl-th].
- [12] J.S. Moreland, J.E. Bernhard, S.A. Bass, Alternative ansatz to wounded nucleon and binary collision scaling in high-energy nuclear collisions, *Phys. Rev. C* 92 (1) (2015) 011901, arXiv:1412.4708 [nucl-th].
- [13] ALICE Collaboration, S. Acharya, et al., Characterizing the initial conditions of heavy-ion collisions at the LHC with mean transverse momentum and anisotropic flow correlations, arXiv:2111.06106 [nucl-ex].
- [14] ATLAS Collaboration, G. Aad, et al., Measurement of the azimuthal anisotropy for charged particle production in $\sqrt{s_{NN}} = 2.76$ TeV lead-lead collisions with the ATLAS detector, *Phys. Rev. C* 86 (2012) 014907, arXiv:1203.3087 [hep-ex].
- [15] CMS Collaboration, S. Chatrchyan, et al., Measurement of higher-order harmonic azimuthal anisotropy in PbPb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, *Phys. Rev. C* 89 (4) (2014) 044906, arXiv:1310.8651 [nucl-ex].
- [16] ATLAS Collaboration, G. Aad, et al., Measurement of the distributions of event-by-event flow harmonics in lead-lead collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ATLAS detector at the LHC, *J. High Energy Phys.* 11 (2013) 183, arXiv:1305.2942 [hep-ex].
- [17] ATLAS Collaboration, G. Aad, et al., Measurement of event-plane correlations in $\sqrt{s_{NN}} = 2.76$ TeV lead-lead collisions with the ATLAS detector, *Phys. Rev. C* 90 (2) (2014) 024905, arXiv:1403.0489 [hep-ex].
- [18] ATLAS Collaboration, G. Aad, et al., Measurement of the correlation between flow harmonics of different order in lead-lead collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ATLAS detector, *Phys. Rev. C* 92 (3) (2015) 034903, arXiv:1504.01289 [hep-ex].
- [19] ALICE Collaboration, J. Adam, et al., Correlated event-by-event fluctuations of flow harmonics in Pb–Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, *Phys. Rev. Lett.* 117 (2016) 182301, arXiv:1604.07663 [nucl-ex].
- [20] ALICE Collaboration, J. Adam, et al., Anisotropic flow of charged particles in Pb–Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, *Phys. Rev. Lett.* 116 (13) (2016) 132302, arXiv:1602.01119 [nucl-ex].
- [21] CMS Collaboration, A.M. Sirunyan, et al., Non-gaussian elliptic-flow fluctuations in PbPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, *Phys. Lett. B* 789 (2019) 643–665, arXiv:1711.05594 [nucl-ex].
- [22] ALICE Collaboration, S. Acharya, et al., Searches for transverse momentum dependent flow vector fluctuations in Pb–Pb and p–Pb collisions at the LHC, *J. High Energy Phys.* 09 (2017) 032, arXiv:1707.05690 [nucl-ex].
- [23] ALICE Collaboration, S. Acharya, et al., Linear and non-linear flow modes in Pb–Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, *Phys. Lett. B* 773 (2017) 68–80, arXiv:1705.04377 [nucl-ex].
- [24] ALICE Collaboration, S. Acharya, et al., Energy dependence and fluctuations of anisotropic flow in Pb–Pb collisions at $\sqrt{s_{NN}} = 5.02$ and 2.76 TeV, *J. High Energy Phys.* 07 (2018) 103, arXiv:1804.02944 [nucl-ex].
- [25] ALICE Collaboration, S. Acharya, et al., Anisotropic flow in Xe–Xe collisions at $\sqrt{s_{NN}} = 5.44$ TeV, *Phys. Lett. B* 784 (2018) 82–95, arXiv:1805.01832 [nucl-ex].
- [26] ALICE Collaboration, S. Acharya, et al., Non-linear flow modes of identified particles in Pb–Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, *J. High Energy Phys.* 06 (2020) 147, arXiv:1912.00740 [nucl-ex].
- [27] ALICE Collaboration, S. Acharya, et al., Higher harmonic non-linear flow modes of charged hadrons in Pb–Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, *J. High Energy Phys.* 05 (2020) 085, arXiv:2002.00633 [nucl-ex].
- [28] ALICE Collaboration, S. Acharya, et al., Measurements of mixed harmonic cumulants in Pb–Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, *Phys. Lett. B* 818 (2021) 136354, arXiv:2102.12180 [nucl-ex].
- [29] D. Teaney, L. Yan, Triangularity and dipole asymmetry in heavy ion collisions, *Phys. Rev. C* 83 (2011) 064904, arXiv:1010.1876 [nucl-th].
- [30] L. Yan, A flow paradigm in heavy-ion collisions, *Chin. Phys. C* 42 (4) (2018) 042001, arXiv:1712.04580 [nucl-th].
- [31] A. Schütttauf, et al., Universality of spectator fragmentation at relativistic bombarding energies, *Nucl. Phys. A* 607 (1996) 457–486, arXiv:nucl-ex/9606001.
- [32] S.A. Voloshin, T. Niida, Ultrarelativistic nuclear collisions: direction of spectator flow, *Phys. Rev. C* 94 (2) (2016) 021901, arXiv:1604.04597 [nucl-th].
- [33] ALICE Collaboration, B. Abelev, et al., Directed flow of charged particles at midrapidity relative to the spectator plane in Pb–Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, *Phys. Rev. Lett.* 111 (23) (2013) 232302, arXiv:1306.4145 [nucl-ex].
- [34] A.S. Goldhaber, Statistical models of fragmentation processes, *Phys. Lett. B* 53 (1974) 306–308.
- [35] D.J. Morrissey, Systematics of momentum distributions from reactions with relativistic ions, *Phys. Rev. C* 39 (1989) 460–470.
- [36] E.J. Moniz, I. Sick, R.R. Whitney, J.R. Ficenec, R.D. Kephart, W.P. Trower, Nuclear Fermi momenta from quasielastic electron scattering, *Phys. Rev. Lett.* 26 (1971) 445–448.
- [37] ALICE Collaboration, K. Aamodt, et al., Harmonic decomposition of two-particle angular correlations in Pb–Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, *Phys. Lett. B* 708 (2012) 249–264, arXiv:1109.2501 [nucl-ex].
- [38] Z. Qiu, U.W. Heinz, Event-by-event shape and flow fluctuations of relativistic heavy-ion collision fireballs, *Phys. Rev. C* 84 (2011) 024911, arXiv:1104.0650 [nucl-th].
- [39] F.G. Gardim, F. Grassi, M. Luzum, J.-Y. Ollitrault, Mapping the hydrodynamic response to the initial geometry in heavy-ion collisions, *Phys. Rev. C* 85 (2012) 024908, arXiv:1111.6538 [nucl-th].
- [40] S.A. Voloshin, A.M. Poskanzer, The physics of the centrality dependence of elliptic flow, *Phys. Lett. B* 474 (2000) 27–32, arXiv:nucl-th/9906075.
- [41] H. Song, U.W. Heinz, Multiplicity scaling in ideal and viscous hydrodynamics, *Phys. Rev. C* 78 (2008) 024902, arXiv:0805.1756 [nucl-th].
- [42] U. Heinz, R. Snellings, Collective flow and viscosity in relativistic heavy-ion collisions, *Annu. Rev. Nucl. Part. Sci.* 63 (2013) 123–151, arXiv:1301.2826 [nucl-th].
- [43] J.E. Bernhard, J.S. Moreland, S.A. Bass, Bayesian estimation of the specific shear and bulk viscosity of quark–gluon plasma, *Nat. Phys.* 15 (11) (2019) 1113–1117.
- [44] JETSCAPE Collaboration, D. Everett, et al., Multisystem bayesian constraints on the transport coefficients of QCD matter, *Phys. Rev. C* 103 (5) (2021) 054904, arXiv:2011.01430 [hep-ph].
- [45] S.A. Voloshin, A.M. Poskanzer, A. Tang, G. Wang, Elliptic flow in the gaussian model of eccentricity fluctuations, *Phys. Lett. B* 659 (2008) 537–541, arXiv:0708.0800 [nucl-th].
- [46] B. Schenke, P. Tribedy, R. Venugopalan, Fluctuating glasma initial conditions and flow in heavy ion collisions, *Phys. Rev. Lett.* 108 (2012) 252301, arXiv:1202.6646 [nucl-th].
- [47] H. Song, S.A. Bass, U. Heinz, T. Hirano, C. Shen, 200 A GeV Au+Au collisions serve a nearly perfect quark–gluon liquid, *Phys. Rev. Lett.* 106 (2011) 192301, arXiv:1011.2783 [nucl-th], Erratum: *Phys. Rev. Lett.* 109 (2012) 139904.
- [48] N. Borghini, P.M. Dinh, J.-Y. Ollitrault, A new method for measuring azimuthal distributions in nucleus-nucleus collisions, *Phys. Rev. C* 63 (2001) 054906, arXiv:nucl-th/0007063.
- [49] A. Bilandzic, R. Snellings, S. Voloshin, Flow analysis with cumulants: direct calculations, *Phys. Rev. C* 83 (2011) 044913, arXiv:1010.0233 [nucl-ex].
- [50] ALICE Collaboration, B. Abelev, et al., Anisotropic flow of charged hadrons, pions and (anti-)protons measured at high transverse momentum in Pb–Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, *Phys. Lett. B* 719 (2013) 18–28, arXiv:1205.5761 [nucl-ex].
- [51] STAR Collaboration, M. Abdallah, et al., Search for the chiral magnetic effect with isobar collisions at $\sqrt{s_{NN}} = 200$ GeV by the STAR collaboration at the BNL relativistic heavy ion collider, *Phys. Rev. C* 105 (1) (2022) 014901, arXiv:2109.00131 [nucl-ex].
- [52] STAR Collaboration, M. Abdallah, et al., Search for the chiral magnetic effect via charge-dependent azimuthal correlations relative to spectator and participant planes in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV, *Phys. Rev. Lett.* 128 (9) (2022) 092301, arXiv:2106.09243 [nucl-ex].
- [53] S.A. Voloshin, Parity violation in hot QCD: how to detect it, *Phys. Rev. C* 70 (2004) 057901, arXiv:hep-ph/0406311.

- [54] F. Wang, Effects of cluster particle correlations on local parity violation observables, *Phys. Rev. C* 81 (2010) 064902, arXiv:0911.1482 [nucl-ex].
- [55] A. Bzdak, V. Koch, J. Liao, Remarks on possible local parity violation in heavy ion collisions, *Phys. Rev. C* 81 (2010) 031901, arXiv:0912.5050 [nucl-th].
- [56] H.-j. Xu, J. Zhao, X. Wang, H. Li, Z.-W. Lin, C. Shen, F. Wang, Varying the chiral magnetic effect relative to flow in a single nucleus-nucleus collision, *Chin. Phys. C* 42 (8) (2018) 084103, arXiv:1710.07265 [nucl-th].
- [57] S.A. Voloshin, Estimate of the signal from the chiral magnetic effect in heavy-ion collisions from measurements relative to the participant and spectator flow planes, *Phys. Rev. C* 98 (5) (2018) 054911, arXiv:1805.05300 [nucl-ex].
- [58] ALICE Collaboration, K. Aamodt, et al., The ALICE experiment at the CERN LHC, *J. Instrum.* 3 (08) (Aug 2008) S08002, <https://doi.org/10.1088/1748-0221/3/08/S08002>.
- [59] ALICE Collaboration, B.B. Abelev, et al., Performance of the ALICE experiment at the CERN LHC, *Int. J. Mod. Phys. A* 29 (2014) 1430044, arXiv:1402.4476 [nucl-ex].
- [60] G. Dellacasa, et al., ALICE Collaboration, ALICE technical design report of the inner tracking system (ITS), CERN-LHCC-99-12, 1999.
- [61] J. Alme, et al., The ALICE TPC, a large 3-dimensional tracking device with fast readout for ultra-high multiplicity events, *Nucl. Instrum. Methods A* 622 (2010) 316–367, arXiv:1001.1950 [physics.ins-det].
- [62] G. Dellacasa, et al., ALICE Collaboration, ALICE technical design report of the zero degree calorimeter (ZDC), CERN-LHCC-99-05, 1999.
- [63] ALICE Collaboration, E. Abbas, et al., Performance of the ALICE VZERO system, *J. Instrum.* 8 (2013) P10016, arXiv:1306.3130 [nucl-ex].
- [64] M. Gyulassy, X.-N. Wang, HIJING 1.0: a Monte Carlo program for parton and particle production in high-energy hadronic and nuclear collisions, *Comput. Phys. Commun.* 83 (1994) 307, arXiv:nucl-th/9502021.
- [65] R. Brun, F. Bruyant, M. Maire, A.C. McPherson, P. Zancarini, GEANT 3: User's Guide Geant 3.10, Geant 3.11, rev. version, CERN, Geneva, 1987, <https://cds.cern.ch/record/1119728>.
- [66] A. Bilandzic, C.H. Christensen, K. Gulbrandsen, A. Hansen, Y. Zhou, Generic framework for anisotropic flow analyses with multiparticle azimuthal correlations, *Phys. Rev. C* 89 (6) (2014) 064904, arXiv:1312.3572 [nucl-ex].
- [67] ALICE Collaboration, B.B. Abelev, et al., Multiparticle azimuthal correlations in p–Pb and Pb–Pb collisions at the CERN large hadron collider, *Phys. Rev. C* 90 (5) (2014) 054901, arXiv:1406.2474 [nucl-ex].
- [68] I. Selyuzhenkov, S. Voloshin, Effects of non-uniform acceptance in anisotropic flow measurement, *Phys. Rev. C* 77 (2008) 034904, arXiv:0707.4672 [nucl-th].
- [69] R. Barlow, Systematic errors: facts and fictions, in: *Conference on Advanced Statistical Techniques in Particle Physics*, 2002, pp. 134–144. 7, arXiv:hep-ex/0207026.
- [70] ALICE Collaboration, S. Acharya, et al., Energy dependence and fluctuations of anisotropic flow in Pb–Pb collisions at $\sqrt{s_{NN}} = 5.02$ and 2.76 TeV, *J. High Energy Phys.* 07 (2018) 103, arXiv:1804.02944 [nucl-ex].
- [71] STAR Collaboration, G. Agakishiev, et al., Energy and system-size dependence of two- and four-particle v_2 measurements in heavy-ion collisions at RHIC and their implications on flow fluctuations and nonflow, *Phys. Rev. C* 86 (2012) 014904, arXiv:1111.5637 [nucl-ex].
- [72] PHOBOS Collaboration, B. Alver, et al., Non-flow correlations and elliptic flow fluctuations in gold–gold collisions at $\sqrt{s_{NN}} = 200$ GeV, *Phys. Rev. C* 81 (2010) 034915, arXiv:1002.0534 [nucl-ex].
- [73] PHOBOS Collaboration, B. Alver, et al., Event-by-event fluctuations of azimuthal particle anisotropy in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV, *Phys. Rev. Lett.* 104 (2010) 142301, arXiv:nucl-ex/0702036.
- [74] ALICE Collaboration, S. Acharya, et al., Centrality and pseudorapidity dependence of the charged-particle multiplicity density in Xe–Xe collisions at $\sqrt{s_{NN}} = 5.44$ TeV, *Phys. Lett. B* 790 (2019) 35–48, arXiv:1805.04432 [nucl-ex].
- [75] CMS Collaboration, V. Khachatryan, et al., Evidence for transverse momentum and pseudorapidity dependent event plane fluctuations in PbPb and pPb collisions, *Phys. Rev. C* 92 (3) (2015) 034911, arXiv:1503.01692 [nucl-ex].
- [76] ALICE Collaboration, K. Aamodt, et al., Centrality dependence of the charged-particle multiplicity density at mid-rapidity in Pb–Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, *Phys. Rev. Lett.* 106 (2011) 032301, arXiv:1012.1657 [nucl-ex].
- [77] STAR Collaboration, S.A. Voloshin, Energy and system size dependence of charged particle elliptic flow and v_2/ϵ scaling, *J. Phys. G* 34 (8) (2007), S883–886, arXiv:nucl-ex/0701038.

ALICE Collaboration

S. Acharya^{123,131}, D. Adamová⁸⁵, A. Adler⁶⁹, G. Aglieri Rinella³², M. Agnello²⁹, N. Agrawal⁵⁰, Z. Ahammed¹³¹, S. Ahmad¹⁵, S.U. Ahn⁷⁰, I. Ahuja³⁷, A. Akindinov¹³⁹, M. Al-Turany⁹⁷, D. Aleksandrov¹³⁹, B. Alessandro⁵⁵, H.M. Alfanda⁶, R. Alfaro Molina⁶⁶, B. Ali¹⁵, Y. Ali¹³, A. Alici²⁵, N. Alizadehvandchali¹¹², A. Alkin³², J. Alme²⁰, G. Alocco⁵¹, T. Alt⁶³, I. Altsybeev¹³⁹, M.N. Anaam⁶, C. Andrei⁴⁵, A. Andronic¹³⁴, V. Anguelov⁹⁴, F. Antinori⁵³, P. Antonioli⁵⁰, C. Anuj¹⁵, N. Apadula⁷³, L. Aphecetche¹⁰², H. Appelshäuser⁶³, S. Arcelli²⁵, R. Arnaldi⁵⁵, I.C. Arsene¹⁹, M. Arslandok¹³⁶, A. Augustinus³², R. Averbeck⁹⁷, S. Aziz¹²⁷, M.D. Azmi¹⁵, A. Badalà⁵², Y.W. Baek⁴⁰, X. Bai⁹⁷, R. Bailhache⁶³, Y. Bailung⁴⁷, R. Bala⁹⁰, A. Balbino²⁹, A. Baldisseri¹²⁶, B. Balis², D. Banerjee⁴, Z. Banoo⁹⁰, R. Barbera²⁶, L. Barioglio⁹⁵, M. Barlou⁷⁷, G.G. Barnaföldi¹³⁵, L.S. Barnby⁸⁴, V. Barret¹²³, L. Barreto¹⁰⁸, C. Bartels¹¹⁵, K. Barth³², E. Bartsch⁶³, F. Baruffaldi²⁷, N. Bastid¹²³, S. Basu⁷⁴, G. Batigne¹⁰², D. Battistini⁹⁵, B. Batyunya¹⁴⁰, D. Bauri⁴⁶, J.L. Bazo Alba¹⁰⁰, I.G. Bearden⁸², C. Beattie¹³⁶, P. Becht⁹⁷, D. Behera⁴⁷, I. Belikov¹²⁵, A.D.C. Bell Hechavarria¹³⁴, F. Bellini²⁵, R. Bellwied¹¹², S. Belokurova¹³⁹, V. Belyaev¹³⁹, G. Bencedi^{135,64}, S. Beole²⁴, A. Bercuci⁴⁵, Y. Berdnikov¹³⁹, A. Berdnikova⁹⁴, L. Bergmann⁹⁴, M.G. Besoiu⁶², L. Betev³², P.P. Bhaduri¹³¹, A. Bhasin⁹⁰, I.R. Bhat⁹⁰, M.A. Bhat⁴, B. Bhattacharjee⁴¹, L. Bianchi²⁴, N. Bianchi⁴⁸, J. Bielčik³⁵, J. Bielčíková⁸⁵, J. Biernat¹⁰⁵, A. Bilandzic⁹⁵, G. Biro¹³⁵, S. Biswas⁴, J.T. Blair¹⁰⁶, D. Blau¹³⁹, M.B. Blidaru⁹⁷, N. Bluhme³⁸, C. Blume⁶³, G. Boca^{21,54}, F. Bock⁸⁶, T. Bodova²⁰, A. Bogdanov¹³⁹, S. Boi²², J. Bok⁵⁷, L. Boldizsár¹³⁵, A. Bolozdynya¹³⁹, M. Bombara³⁷, P.M. Bond³², G. Bonomi^{130,54}, H. Borel¹²⁶, A. Borissov¹³⁹, H. Bossi¹³⁶, E. Botta²⁴, L. Bratrud⁶³, P. Braun-Munzinger⁹⁷, M. Bregant¹⁰⁸,

M. Broz ^{35, [id](#)}, G.E. Bruno ^{96,31, [id](#)}, M.D. Buckland ^{115, [id](#)}, D. Budnikov ^{139, [id](#)}, H. Buesching ^{63, [id](#)},
 S. Bufalino ^{29, [id](#)}, O. Bugnon ¹⁰², P. Buhler ^{101, [id](#)}, Z. Buthelezi ^{67,119, [id](#)}, J.B. Butt ¹³, A. Bylinkin ^{114, [id](#)},
 S.A. Bysiak ¹⁰⁵, M. Cai ^{27,6, [id](#)}, H. Caines ^{136, [id](#)}, A. Caliva ^{97, [id](#)}, E. Calvo Villar ^{100, [id](#)}, J.M.M. Camacho ^{107, [id](#)},
 P. Camerini ^{23, [id](#)}, F.D.M. Canedo ^{108, [id](#)}, M. Carabas ^{122, [id](#)}, F. Carnesecchi ^{32, [id](#)}, R. Caron ^{124,126, [id](#)}, J. Castillo
 Castellanos ^{126, [id](#)}, F. Catalano ^{29, [id](#)}, C. Ceballos Sanchez ^{140, [id](#)}, I. Chakaberia ^{73, [id](#)}, P. Chakraborty ^{46, [id](#)},
 S. Chandra ^{131, [id](#)}, S. Chapeland ^{32, [id](#)}, M. Chartier ^{115, [id](#)}, S. Chattopadhyay ^{131, [id](#)}, S. Chattopadhyay ^{98, [id](#)},
 T.G. Chavez ^{44, [id](#)}, T. Cheng ^{6, [id](#)}, C. Cheshkov ^{124, [id](#)}, B. Cheynis ^{124, [id](#)}, V. Chibante Barroso ^{32, [id](#)},
 D.D. Chinellato ^{109, [id](#)}, E.S. Chizzali ^{95, [id](#),II}, J. Cho ^{57, [id](#)}, S. Cho ^{57, [id](#)}, P. Chochula ^{32, [id](#)}, P. Christakoglou ^{83, [id](#)},
 C.H. Christensen ^{82, [id](#)}, P. Christiansen ^{74, [id](#)}, T. Chujo ^{121, [id](#)}, M. Ciacco ^{29, [id](#)}, C. Cicalo ^{51, [id](#)}, L. Cifarelli ^{25, [id](#)},
 F. Cindolo ^{50, [id](#)}, M.R. Ciupek ⁹⁷, G. Clai ^{50,III}, F. Colamaria ^{49, [id](#)}, J.S. Colburn ⁹⁹, D. Colella ^{96,31, [id](#)},
 A. Collu ⁷³, M. Colocci ^{32, [id](#)}, M. Concas ^{55, [id](#),IV}, G. Conesa Balbastre ^{72, [id](#)}, Z. Conesa del Valle ^{127, [id](#)},
 G. Contin ^{23, [id](#)}, J.G. Contreras ^{35, [id](#)}, M.L. Coquet ^{126, [id](#)}, T.M. Cormier ^{86,I}, P. Cortese ^{129,55, [id](#)},
 M.R. Cosentino ^{110, [id](#)}, F. Costa ^{32, [id](#)}, S. Costanza ^{21,54, [id](#)}, P. Crochet ^{123, [id](#)}, R. Cruz-Torres ^{73, [id](#)}, E. Cuautle ⁶⁴,
 P. Cui ^{6, [id](#)}, L. Cunqueiro ⁸⁶, A. Dainese ^{53, [id](#)}, M.C. Danisch ^{94, [id](#)}, A. Danu ^{62, [id](#)}, P. Das ^{79, [id](#)}, P. Das ^{4, [id](#)},
 S. Das ^{4, [id](#)}, S. Dash ^{46, [id](#)}, R.M.H. David ⁴⁴, A. De Caro ^{28, [id](#)}, G. de Cataldo ^{49, [id](#)}, L. De Cilladi ^{24, [id](#)}, J. de
 Cuveland ³⁸, A. De Falco ^{22, [id](#)}, D. De Gruttola ^{28, [id](#)}, N. De Marco ^{55, [id](#)}, C. De Martin ^{23, [id](#)}, S. De
 Pasquale ^{28, [id](#)}, S. Deb ^{47, [id](#)}, H.F. Degenhardt ¹⁰⁸, K.R. Deja ¹³², R. Del Grande ^{95, [id](#)}, L. Dello Stritto ^{28, [id](#)},
 W. Deng ^{6, [id](#)}, P. Dhankher ^{18, [id](#)}, D. Di Bari ^{31, [id](#)}, A. Di Mauro ^{32, [id](#)}, R.A. Diaz ^{140,7, [id](#)}, T. Dietel ^{111, [id](#)},
 Y. Ding ^{124,6, [id](#)}, R. Divià ^{32, [id](#)}, D.U. Dixit ^{18, [id](#)}, Ø. Djuvsland ²⁰, U. Dmitrieva ^{139, [id](#)}, A. Dobrin ^{62, [id](#)},
 B. Dönigus ^{63, [id](#)}, A.K. Dubey ^{131, [id](#)}, J.M. Dubinski ^{132, [id](#)}, A. Dubla ^{97, [id](#)}, S. Dudi ^{89, [id](#)}, P. Dupieux ^{123, [id](#)},
 M. Durkac ¹⁰⁴, N. Dzalaiova ¹², T.M. Eder ^{134, [id](#)}, R.J. Ehlers ^{86, [id](#)}, V.N. Eikeland ²⁰, F. Eisenhut ^{63, [id](#)},
 D. Elia ^{49, [id](#)}, B. Erazmus ^{102, [id](#)}, F. Ercolessi ^{25, [id](#)}, F. Erhardt ^{88, [id](#)}, M.R. Ersdal ²⁰, B. Espagnon ^{127, [id](#)},
 G. Eulisse ^{32, [id](#)}, D. Evans ^{99, [id](#)}, S. Evdokimov ^{139, [id](#)}, L. Fabbietti ^{95, [id](#)}, M. Faggin ^{27, [id](#)}, J. Faivre ^{72, [id](#)},
 F. Fan ^{6, [id](#)}, W. Fan ^{73, [id](#)}, A. Fantoni ^{48, [id](#)}, M. Fasel ^{86, [id](#)}, P. Fecchio ²⁹, A. Feliciello ^{55, [id](#)}, G. Feofilov ^{139, [id](#)},
 A. Fernández Téllez ^{44, [id](#)}, M.B. Ferrer ^{32, [id](#)}, A. Ferrero ^{126, [id](#)}, A. Ferretti ^{24, [id](#)}, V.J.G. Feuillard ^{94, [id](#)},
 J. Figiel ^{105, [id](#)}, V. Filova ^{35, [id](#)}, D. Finogeev ^{139, [id](#)}, F.M. Fionda ^{51, [id](#)}, G. Fiorenza ⁹⁶, F. Flor ^{112, [id](#)},
 A.N. Flores ^{106, [id](#)}, S. Foertsch ^{67, [id](#)}, I. Fokin ^{94, [id](#)}, S. Fokin ^{139, [id](#)}, E. Fragiaco ^{56, [id](#)}, E. Frajna ^{135, [id](#)},
 U. Fuchs ^{32, [id](#)}, N. Funicello ^{28, [id](#)}, C. Furget ^{72, [id](#)}, A. Furs ^{139, [id](#)}, J.J. Gaardhøje ^{82, [id](#)}, M. Gagliardi ^{24, [id](#)},
 A.M. Gago ^{100, [id](#)}, A. Gal ¹²⁵, C.D. Galvan ^{107, [id](#)}, P. Ganoti ^{77, [id](#)}, C. Garabatos ^{97, [id](#)}, J.R.A. Garcia ^{44, [id](#)},
 E. Garcia-Solis ^{9, [id](#)}, K. Garg ^{102, [id](#)}, C. Gargiulo ^{32, [id](#)}, A. Garibli ⁸⁰, K. Garner ¹³⁴, E.F. Gauger ^{106, [id](#)},
 A. Gautam ^{114, [id](#)}, M.B. Gay Ducati ^{65, [id](#)}, M. Germain ^{102, [id](#)}, S.K. Ghosh ⁴, M. Giacalone ^{25, [id](#)},
 P. Gianotti ^{48, [id](#)}, P. Giubellino ^{97,55, [id](#)}, P. Giubilato ^{27, [id](#)}, A.M.C. Glaenger ^{126, [id](#)}, P. Glässel ^{94, [id](#)},
 E. Glimos ^{118, [id](#)}, D.J.Q. Goh ⁷⁵, V. Gonzalez ^{133, [id](#)}, L.H. González-Trueba ^{66, [id](#)}, S. Gorbunov ³⁸,
 M. Gorgon ^{2, [id](#)}, L. Görlich ^{105, [id](#)}, S. Gotovac ³³, V. Grabski ^{66, [id](#)}, L.K. Graczykowski ^{132, [id](#)}, E. Grecka ^{85, [id](#)},
 L. Greiner ^{73, [id](#)}, A. Grelli ^{58, [id](#)}, C. Grigoras ^{32, [id](#)}, V. Grigoriev ^{139, [id](#)}, S. Grigoryan ^{140,1, [id](#)}, F. Grosa ^{32, [id](#)},
 J.F. Grosse-Oetringhaus ^{32, [id](#)}, R. Grosso ^{97, [id](#)}, D. Grund ^{35, [id](#)}, G.G. Guardiano ^{109, [id](#)}, R. Guernane ^{72, [id](#)},
 M. Guilbaud ^{102, [id](#)}, K. Gulbrandsen ^{82, [id](#)}, T. Gunji ^{120, [id](#)}, W. Guo ^{6, [id](#)}, A. Gupta ^{90, [id](#)}, R. Gupta ^{90, [id](#)},
 S.P. Guzman ^{44, [id](#)}, L. Gyulai ^{135, [id](#)}, M.K. Habib ⁹⁷, C. Hadjidakis ^{127, [id](#)}, H. Hamagaki ^{75, [id](#)}, M. Hamid ⁶,
 Y. Han ^{137, [id](#)}, R. Hannigan ^{106, [id](#)}, M.R. Haque ^{132, [id](#)}, A. Harlanderova ⁹⁷, J.W. Harris ^{136, [id](#)}, A. Harton ^{9, [id](#)},
 J.A. Hasenbichler ³², H. Hassan ^{86, [id](#)}, D. Hatzifotiadou ^{50, [id](#)}, P. Hauer ^{42, [id](#)}, L.B. Havener ^{136, [id](#)},
 S.T. Heckel ^{95, [id](#)}, E. Hellbär ^{97, [id](#)}, H. Helstrup ^{34, [id](#)}, T. Herman ^{35, [id](#)}, G. Herrera Corral ^{8, [id](#)}, F. Herrmann ¹³⁴,
 K.F. Hetland ^{34, [id](#)}, B. Heybeck ^{63, [id](#)}, H. Hillemanns ^{32, [id](#)}, C. Hills ^{115, [id](#)}, B. Hippolyte ^{125, [id](#)}, B. Hofman ^{58, [id](#)},

B. Hohlweger^{83, [id](#)}, J. Honermann^{134, [id](#)}, G.H. Hong^{137, [id](#)}, D. Horak^{35, [id](#)}, A. Horzyk^{2, [id](#)}, R. Hosokawa¹⁴,
 Y. Hou^{6, [id](#)}, P. Hristov^{32, [id](#)}, C. Hughes^{118, [id](#)}, P. Huhn⁶³, L.M. Huhta^{113, [id](#)}, C.V. Hulse^{127, [id](#)},
 T.J. Humanic^{87, [id](#)}, H. Hushnud⁹⁸, A. Hutson^{112, [id](#)}, D. Hutter^{38, [id](#)}, J.P. Iddon^{115, [id](#)}, R. Ilkaev¹³⁹,
 H. Ilyas^{13, [id](#)}, M. Inaba^{121, [id](#)}, G.M. Innocenti^{32, [id](#)}, M. Ippolitov^{139, [id](#)}, A. Isakov^{85, [id](#)}, T. Isidori^{114, [id](#)},
 M.S. Islam^{98, [id](#)}, M. Ivanov^{97, [id](#)}, V. Ivanov^{139, [id](#)}, V. Izucheev¹³⁹, M. Jablonski^{2, [id](#)}, B. Jacak^{73, [id](#)},
 N. Jacazio^{32, [id](#)}, P.M. Jacobs^{73, [id](#)}, S. Jadlovská¹⁰⁴, J. Jadlovsky¹⁰⁴, L. Jaffe³⁸, C. Jahnke^{109, [id](#)},
 M.A. Janik^{132, [id](#)}, T. Janson⁶⁹, M. Jercic⁸⁸, O. Jevons⁹⁹, A.A.P. Jimenez^{64, [id](#)}, F. Jonas^{86, [id](#)}, P.G. Jones⁹⁹,
 J.M. Jowett^{32, 97, [id](#)}, J. Jung^{63, [id](#)}, M. Jung^{63, [id](#)}, A. Junique^{32, [id](#)}, A. Jusko^{99, [id](#)}, M.J. Kabus^{32, 132, [id](#)},
 J. Kaewjai¹⁰³, P. Kalinak^{59, [id](#)}, A.S. Kalteyer^{97, [id](#)}, A. Kalweit^{32, [id](#)}, V. Kaplin^{139, [id](#)}, A. Karasu Uysal^{71, [id](#)},
 D. Karatovic^{88, [id](#)}, O. Karavichev^{139, [id](#)}, T. Karavicheva^{139, [id](#)}, P. Karczmarczyk^{132, [id](#)}, E. Karpechev^{139, [id](#)},
 V. Kashyap⁷⁹, A. Kazantsev¹³⁹, U. Kebschull^{69, [id](#)}, R. Keidel^{138, [id](#)}, D.L.D. Keijdener⁵⁸, M. Keil^{32, [id](#)},
 B. Ketzer^{42, [id](#)}, A.M. Khan^{6, [id](#)}, S. Khan^{15, [id](#)}, A. Khanzadeev^{139, [id](#)}, Y. Kharlov^{139, [id](#)}, A. Khatun^{15, [id](#)},
 A. Khuntia^{105, [id](#)}, B. Kileng^{34, [id](#)}, B. Kim^{16, [id](#)}, C. Kim^{16, [id](#)}, D.J. Kim^{113, [id](#)}, E.J. Kim^{68, [id](#)}, J. Kim^{137, [id](#)},
 J.S. Kim^{40, [id](#)}, J. Kim^{94, [id](#)}, J. Kim^{68, [id](#)}, M. Kim^{94, [id](#)}, S. Kim^{17, [id](#)}, T. Kim^{137, [id](#)}, S. Kirsch^{63, [id](#)}, I. Kisel^{38, [id](#)},
 S. Kiselev^{139, [id](#)}, A. Kisiel^{132, [id](#)}, J.P. Kitowski^{2, [id](#)}, J.L. Klay^{5, [id](#)}, J. Klein^{32, [id](#)}, S. Klein^{73, [id](#)},
 C. Klein-Bösing^{134, [id](#)}, M. Kleiner^{63, [id](#)}, T. Klemenz^{95, [id](#)}, A. Kluge^{32, [id](#)}, A.G. Knospe^{112, [id](#)}, C. Kobdaj^{103, [id](#)},
 T. Kollegger⁹⁷, A. Kondratyev^{140, [id](#)}, N. Kondratyeva^{139, [id](#)}, E. Kondratyuk^{139, [id](#)}, J. König^{63, [id](#)},
 S.A. Königstorfer^{95, [id](#)}, P.J. Konopka^{32, [id](#)}, G. Kornakov^{132, [id](#)}, S.D. Koryciak^{2, [id](#)}, A. Kotliarov^{85, [id](#)},
 O. Kovalenko^{78, [id](#)}, V. Kovalenko^{139, [id](#)}, M. Kowalski^{105, [id](#)}, I. Králik^{59, [id](#)}, A. Kravčáková^{37, [id](#)}, L. Kreis⁹⁷,
 M. Krivda^{99, 59, [id](#)}, F. Krizek^{85, [id](#)}, K. Krizkova Gajdosova^{35, [id](#)}, M. Kroesen^{94, [id](#)}, M. Krüger^{63, [id](#)},
 D.M. Krupova^{35, [id](#)}, E. Kryshen^{139, [id](#)}, M. Krzewicki³⁸, V. Kučera^{32, [id](#)}, C. Kuhn^{125, [id](#)}, P.G. Kuijjer^{83, [id](#)},
 T. Kumaoka¹²¹, D. Kumar¹³¹, L. Kumar^{89, [id](#)}, N. Kumar⁸⁹, S. Kundu^{32, [id](#)}, P. Kurashvili^{78, [id](#)},
 A. Kurepin^{139, [id](#)}, A.B. Kurepin^{139, [id](#)}, S. Kushpil^{85, [id](#)}, J. Kvapil^{99, [id](#)}, M.J. Kweon^{57, [id](#)}, J.Y. Kwon^{57, [id](#)},
 Y. Kwon^{137, [id](#)}, S.L. La Pointe^{38, [id](#)}, P. La Rocca^{26, [id](#)}, Y.S. Lai⁷³, A. Lakrathok¹⁰³, M. Lamanna^{32, [id](#)},
 R. Langoy^{117, [id](#)}, P. Larionov^{48, [id](#)}, E. Laudi^{32, [id](#)}, L. Lautner^{32, 95, [id](#)}, R. Lavicka^{101, [id](#)}, T. Lazareva^{139, [id](#)},
 R. Lea^{130, 54, [id](#)}, J. Lehrbach^{38, [id](#)}, R.C. Lemmon^{84, [id](#)}, I. León Monzón^{107, [id](#)}, M.M. Lesch^{95, [id](#)},
 E.D. Lesser^{18, [id](#)}, M. Lettrich⁹⁵, P. Lévai^{135, [id](#)}, X. Li¹⁰, X.L. Li⁶, J. Lien^{117, [id](#)}, R. Lietava^{99, [id](#)}, B. Lim^{16, [id](#)},
 S.H. Lim^{16, [id](#)}, V. Lindenstruth^{38, [id](#)}, A. Lindner⁴⁵, C. Lippmann^{97, [id](#)}, A. Liu^{18, [id](#)}, D.H. Liu^{6, [id](#)}, J. Liu^{115, [id](#)},
 I.M. Lofnes^{20, [id](#)}, V. Loginov¹³⁹, C. Loizides^{86, [id](#)}, P. Loncar^{33, [id](#)}, J.A. Lopez^{94, [id](#)}, X. Lopez^{123, [id](#)}, E. López
 Torres^{7, [id](#)}, P. Lu^{97, 116, [id](#)}, J.R. Luhder^{134, [id](#)}, M. Lunardon^{27, [id](#)}, G. Luparello^{56, [id](#)}, Y.G. Ma^{39, [id](#)},
 A. Maevskaya¹³⁹, M. Mager^{32, [id](#)}, T. Mahmoud⁴², A. Maire^{125, [id](#)}, M. Malaev^{139, [id](#)}, N.M. Malik^{90, [id](#)},
 Q.W. Malik¹⁹, S.K. Malik^{90, [id](#)}, L. Malinina^{140, [id](#)}, VII, D. Mal'Kevich^{139, [id](#)}, D. Mallick^{79, [id](#)}, N. Mallick^{47, [id](#)},
 G. Mandaglio^{30, 52, [id](#)}, V. Manko^{139, [id](#)}, F. Manso^{123, [id](#)}, V. Manzari^{49, [id](#)}, Y. Mao^{6, [id](#)}, G.V. Margagliotti^{23, [id](#)},
 A. Margotti^{50, [id](#)}, A. Marín^{97, [id](#)}, C. Markert^{106, [id](#)}, M. Marquard⁶³, N.A. Martin⁹⁴, P. Martinengo^{32, [id](#)},
 J.L. Martinez¹¹², M.I. Martínez^{44, [id](#)}, G. Martínez García^{102, [id](#)}, S. Masciocchi^{97, [id](#)}, M. Masera^{24, [id](#)},
 A. Masoni^{51, [id](#)}, L. Massacrier^{127, [id](#)}, A. Mastroserio^{128, 49, [id](#)}, A.M. Mathis^{95, [id](#)}, O. Matonoha^{74, [id](#)},
 P.F.T. Matuoka¹⁰⁸, A. Matyja^{105, [id](#)}, C. Mayer^{105, [id](#)}, A.L. Mazuecos^{32, [id](#)}, F. Mazzaschi^{24, [id](#)},
 M. Mazzilli^{32, [id](#)}, J.E. Mdhluhi^{119, [id](#)}, A.F. Mechler⁶³, Y. Melikyan^{139, [id](#)}, A. Menchaca-Rocha^{66, [id](#)},
 E. Meninno^{101, 28, [id](#)}, A.S. Menon^{112, [id](#)}, M. Meres^{12, [id](#)}, S. Mhlanga^{111, 67}, Y. Miake¹²¹, L. Micheletti^{55, [id](#)},
 L.C. Migliorin¹²⁴, D.L. Mihaylov^{95, [id](#)}, K. Mikhaylov^{140, 139, [id](#)}, A.N. Mishra^{135, [id](#)}, D. Miśkowiec^{97, [id](#)},
 A. Modak^{4, [id](#)}, A.P. Mohanty^{58, [id](#)}, B. Mohanty⁷⁹, M. Mohisin Khan^{15, [id](#)}, V, M.A. Molander^{43, [id](#)},
 Z. Moravcova^{82, [id](#)}, C. Mordasini^{95, [id](#)}, D.A. Moreira De Godoy^{134, [id](#)}, I. Morozov^{139, [id](#)}, A. Morsch^{32, [id](#)},

T. Mrnjavac^{32, [id](#)}, V. Muccifora^{48, [id](#)}, E. Mudnic³³, S. Muhuri^{131, [id](#)}, J.D. Mulligan^{73, [id](#)}, A. Mulliri²², M.G. Munhoz^{108, [id](#)}, R.H. Munzer^{63, [id](#)}, H. Murakami^{120, [id](#)}, S. Murray^{111, [id](#)}, L. Musa^{32, [id](#)}, J. Musinsky^{59, [id](#)}, J.W. Myrcha^{132, [id](#)}, B. Naik^{119, [id](#)}, R. Nair^{78, [id](#)}, B.K. Nandi^{46, [id](#)}, R. Nania^{50, [id](#)}, E. Nappi^{49, [id](#)}, A.F. Nassirpour^{74, [id](#)}, A. Nath^{94, [id](#)}, C. Nattrass^{118, [id](#)}, A. Neagu¹⁹, A. Negru¹²², L. Nellen^{64, [id](#)}, S.V. Nesbo³⁴, G. Neskovic^{38, [id](#)}, D. Nesterov^{139, [id](#)}, B.S. Nielsen^{82, [id](#)}, E.G. Nielsen^{82, [id](#)}, S. Nikolaev^{139, [id](#)}, S. Nikulin^{139, [id](#)}, V. Nikulin^{139, [id](#)}, F. Noferini^{50, [id](#)}, S. Noh^{11, [id](#)}, P. Nomokonov^{140, [id](#)}, J. Norman^{115, [id](#)}, N. Novitzky^{121, [id](#)}, P. Nowakowski^{132, [id](#)}, A. Nyanin^{139, [id](#)}, J. Nystrand^{20, [id](#)}, M. Ogino^{75, [id](#)}, A. Ohlson^{74, [id](#)}, V.A. Okorokov^{139, [id](#)}, J. Oleniacz^{132, [id](#)}, A.C. Oliveira Da Silva^{118, [id](#)}, M.H. Oliver^{136, [id](#)}, A. Onnerstad^{113, [id](#)}, C. Oppedisano^{55, [id](#)}, A. Ortiz Velasquez^{64, [id](#)}, A. Oskarsson⁷⁴, J. Otwinowski^{105, [id](#)}, M. Oya⁹², K. Oyama^{75, [id](#)}, Y. Pachmayer^{94, [id](#)}, S. Padhan^{46, [id](#)}, D. Pagano^{130,54, [id](#)}, G. Paic^{64, [id](#)}, A. Palasciano^{49, [id](#)}, S. Panebianco^{126, [id](#)}, J. Park^{57, [id](#)}, J.E. Parkkila^{32,113, [id](#)}, S.P. Pathak¹¹², R.N. Patra⁹⁰, B. Paul^{22, [id](#)}, H. Pei^{6, [id](#)}, T. Peitzmann^{58, [id](#)}, X. Peng^{6, [id](#)}, L.G. Pereira^{65, [id](#)}, H. Pereira Da Costa^{126, [id](#)}, D. Peresunko^{139, [id](#)}, G.M. Perez^{7, [id](#)}, S. Perrin^{126, [id](#)}, Y. Pestov¹³⁹, V. Petráček^{35, [id](#)}, V. Petrov^{139, [id](#)}, M. Petrovici^{45, [id](#)}, R.P. Pezzi^{102,65, [id](#)}, S. Piano^{56, [id](#)}, M. Pikna^{12, [id](#)}, P. Pillot^{102, [id](#)}, O. Pinazza^{50,32, [id](#)}, L. Pinsky¹¹², C. Pinto^{95,26, [id](#)}, S. Pisano^{48, [id](#)}, M. Płoskoń^{73, [id](#)}, M. Planinic⁸⁸, F. Pliquet⁶³, M.G. Poghosyan^{86, [id](#)}, S. Politano^{29, [id](#)}, N. Poljak^{88, [id](#)}, A. Pop^{45, [id](#)}, S. Porteboeuf-Houssais^{123, [id](#)}, J. Porter^{73, [id](#)}, V. Pozdniakov^{140, [id](#)}, S.K. Prasad^{4, [id](#)}, S. Prasad^{47, [id](#)}, R. Preghenella^{50, [id](#)}, F. Prino^{55, [id](#)}, C.A. Pruneau^{133, [id](#)}, I. Pshenichnov^{139, [id](#)}, M. Puccio^{32, [id](#)}, S. Qiu^{83, [id](#)}, L. Quaglia^{24, [id](#)}, R.E. Quishpe¹¹², S. Ragoni^{99, [id](#)}, A. Rakotozafindrabe^{126, [id](#)}, L. Ramello^{129,55, [id](#)}, F. Rami^{125, [id](#)}, S.A.R. Ramirez^{44, [id](#)}, T.A. Rancien⁷², R. Raniwala^{91, [id](#)}, S. Raniwala⁹¹, S.S. Räsänen^{43, [id](#)}, R. Rath^{47, [id](#)}, I. Ravasenga^{83, [id](#)}, K.F. Read^{86,118, [id](#)}, A.R. Redelbach^{38, [id](#)}, K. Redlich^{78, [id](#)}, VI, A. Rehman²⁰, P. Reichelt⁶³, F. Reidt^{32, [id](#)}, H.A. Reme-Ness^{34, [id](#)}, Z. Rescakova³⁷, K. Reygers^{94, [id](#)}, A. Riabov^{139, [id](#)}, V. Riabov^{139, [id](#)}, R. Ricci^{28, [id](#)}, T. Richert⁷⁴, M. Richter^{19, [id](#)}, W. Riegler^{32, [id](#)}, F. Riggi^{26, [id](#)}, C. Ristea^{62, [id](#)}, M. Rodríguez Cahuantzi^{44, [id](#)}, K. Røed^{19, [id](#)}, R. Rogalev^{139, [id](#)}, E. Rogochaya^{140, [id](#)}, T.S. Rogoschinski^{63, [id](#)}, D. Rohr^{32, [id](#)}, D. Röhrich^{20, [id](#)}, P.F. Rojas⁴⁴, S. Rojas Torres^{35, [id](#)}, P.S. Rokita^{132, [id](#)}, F. Ronchetti^{48, [id](#)}, A. Rosano^{30,52, [id](#)}, E.D. Rosas⁶⁴, A. Rossi^{53, [id](#)}, A. Roy^{47, [id](#)}, P. Roy⁹⁸, S. Roy^{46, [id](#)}, N. Rubini^{25, [id](#)}, D. Ruggiano^{132, [id](#)}, R. Rui^{23, [id](#)}, B. Rumyantsev¹⁴⁰, P.G. Russek^{2, [id](#)}, R. Russo^{83, [id](#)}, A. Rustamov^{80, [id](#)}, E. Ryabinkin^{139, [id](#)}, Y. Ryabov^{139, [id](#)}, A. Rybicki^{105, [id](#)}, H. Rytönen^{113, [id](#)}, W. Rzesza^{132, [id](#)}, O.A.M. Saarimaki^{43, [id](#)}, R. Sadek^{102, [id](#)}, S. Sadovsky^{139, [id](#)}, J. Saetre^{20, [id](#)}, K. Šafařík^{35, [id](#)}, S.K. Saha^{131, [id](#)}, S. Saha^{79, [id](#)}, B. Sahoo^{46, [id](#)}, P. Sahoo⁴⁶, R. Sahoo^{47, [id](#)}, S. Sahoo⁶⁰, D. Sahu^{47, [id](#)}, P.K. Sahu^{60, [id](#)}, J. Saini^{131, [id](#)}, K. Sajdakova³⁷, S. Sakai^{121, [id](#)}, M.P. Salvan^{97, [id](#)}, S. Sambyal^{90, [id](#)}, T.B. Saramela¹⁰⁸, D. Sarkar^{133, [id](#)}, N. Sarkar¹³¹, P. Sarma^{41, [id](#)}, V. Sarritzu^{22, [id](#)}, V.M. Sarti^{95, [id](#)}, M.H.P. Sas^{136, [id](#)}, J. Schambach^{86, [id](#)}, H.S. Scheid^{63, [id](#)}, C. Schiaua^{45, [id](#)}, R. Schicker^{94, [id](#)}, A. Schmah⁹⁴, C. Schmidt^{97, [id](#)}, H.R. Schmidt⁹³, M.O. Schmidt^{32, [id](#)}, M. Schmidt⁹³, N.V. Schmidt^{86,63, [id](#)}, A.R. Schmier^{118, [id](#)}, R. Schotter^{125, [id](#)}, J. Schukraft^{32, [id](#)}, K. Schwarz⁹⁷, K. Schweda^{97, [id](#)}, G. Scioli^{25, [id](#)}, E. Scomparin^{55, [id](#)}, J.E. Seger^{14, [id](#)}, Y. Sekiguchi¹²⁰, D. Sekihata^{120, [id](#)}, I. Selyuzhenkov^{97,139, [id](#)}, S. Senyukov^{125, [id](#)}, J.J. Seo^{57, [id](#)}, D. Serebryakov^{139, [id](#)}, L. Šerkšnytė^{95, [id](#)}, A. Sevcenco^{62, [id](#)}, T.J. Shaba^{67, [id](#)}, A. Shabanov¹³⁹, A. Shabetai^{102, [id](#)}, R. Shahoyan³², W. Shaikh⁹⁸, A. Shangaraev^{139, [id](#)}, A. Sharma⁸⁹, D. Sharma^{46, [id](#)}, H. Sharma^{105, [id](#)}, M. Sharma^{90, [id](#)}, N. Sharma^{89, [id](#)}, S. Sharma^{90, [id](#)}, U. Sharma^{90, [id](#)}, A. Shatat^{127, [id](#)}, O. Sheibani¹¹², K. Shigaki^{92, [id](#)}, M. Shimomura⁷⁶, S. Shirinkin^{139, [id](#)}, Q. Shou^{39, [id](#)}, Y. Sibiriak^{139, [id](#)}, S. Siddhanta^{51, [id](#)}, T. Siemiarczuk^{78, [id](#)}, T.F. Silva^{108, [id](#)}, D. Silvermyr^{74, [id](#)}, T. Simantathammakul¹⁰³, R. Simeonov^{36, [id](#)}, G. Simonetti³², B. Singh⁹⁰, B. Singh^{95, [id](#)}, R. Singh^{79, [id](#)}, R. Singh^{90, [id](#)}, R. Singh^{47, [id](#)},

V.K. Singh ^{131, [id](#)}, V. Singhal ^{131, [id](#)}, T. Sinha ^{98, [id](#)}, B. Sitar ^{12, [id](#)}, M. Sitta ^{129,55, [id](#)}, T.B. Skaali ¹⁹,
 G. Skorodumovs ^{94, [id](#)}, M. Slupecki ^{43, [id](#)}, N. Smirnov ^{136, [id](#)}, R.J.M. Snellings ^{58, [id](#)}, E.H. Solheim ^{19, [id](#)},
 C. Soncco ¹⁰⁰, J. Song ^{112, [id](#)}, A. Songmoolnak ¹⁰³, F. Soramel ^{27, [id](#)}, S.P. Sorensen ^{118, [id](#)}, R. Soto Camacho ⁴⁴,
 R. Spijkers ^{83, [id](#)}, I. Sputowska ^{105, [id](#)}, J. Staa ^{74, [id](#)}, J. Stachel ^{94, [id](#)}, I. Stan ^{62, [id](#)}, P.J. Steffanic ^{118, [id](#)},
 S.F. Stiefelmaier ^{94, [id](#)}, D. Stocco ^{102, [id](#)}, I. Storehaug ^{19, [id](#)}, M.M. Storetvedt ^{34, [id](#)}, P. Stratmann ^{134, [id](#)},
 S. Strazzi ^{25, [id](#)}, C.P. Stylianidis ⁸³, A.A.P. Suaide ^{108, [id](#)}, C. Suire ^{127, [id](#)}, M. Sukhanov ^{139, [id](#)}, M. Suljic ^{32, [id](#)},
 V. Sumberia ^{90, [id](#)}, S. Sumowidagdo ^{81, [id](#)}, S. Swain ⁶⁰, A. Szabo ¹², I. Szarka ^{12, [id](#)}, U. Tabassam ¹³,
 S.F. Taghavi ^{95, [id](#)}, G. Taillepied ^{97,123, [id](#)}, J. Takahashi ^{109, [id](#)}, G.J. Tambave ^{20, [id](#)}, S. Tang ^{123,6, [id](#)},
 Z. Tang ^{116, [id](#)}, J.D. Tapia Takaki ^{114, [id](#)}, N. Tapus ¹²², L.A. Tarasovicova ^{134, [id](#)}, M.G. Tarzila ^{45, [id](#)},
 A. Tauro ^{32, [id](#)}, A. Telesca ^{32, [id](#)}, L. Terlizzi ^{24, [id](#)}, C. Terrevoli ^{112, [id](#)}, G. Tersimonov ³, S. Thakur ^{131, [id](#)},
 D. Thomas ^{106, [id](#)}, R. Tieulent ^{124, [id](#)}, A. Tikhonov ^{139, [id](#)}, A.R. Timmins ^{112, [id](#)}, M. Tkacik ¹⁰⁴, T. Tkacik ^{104, [id](#)},
 A. Toia ^{63, [id](#)}, N. Topilskaya ^{139, [id](#)}, M. Toppi ^{48, [id](#)}, F. Torres-Acosta ¹⁸, T. Tork ^{127, [id](#)}, A.G. Torres Ramos ^{31, [id](#)},
 A. Trifiró ^{30,52, [id](#)}, A.S. Triolo ^{30,52, [id](#)}, S. Tripathy ^{50, [id](#)}, T. Tripathy ^{46, [id](#)}, S. Trogolo ^{32, [id](#)}, V. Trubnikov ^{3, [id](#)},
 W.H. Trzaska ^{113, [id](#)}, T.P. Trzcinski ^{132, [id](#)}, R. Turrisi ^{53, [id](#)}, T.S. Tveter ^{19, [id](#)}, K. Ullaland ^{20, [id](#)}, B. Ulukutlu ^{95, [id](#)},
 A. Uras ^{124, [id](#)}, M. Urioni ^{54,130, [id](#)}, G.L. Usai ^{22, [id](#)}, M. Vala ³⁷, N. Valle ^{21, [id](#)}, S. Vallero ^{55, [id](#)}, L.V.R. van
 Doremalen ⁵⁸, M. van Leeuwen ^{83, [id](#)}, C.A. van Veen ^{94, [id](#)}, R.J.G. van Weelden ^{83, [id](#)}, P. Vande Vyvre ^{32, [id](#)},
 D. Varga ^{135, [id](#)}, Z. Varga ^{135, [id](#)}, M. Varga-Kofarago ^{135, [id](#)}, M. Vasileiou ^{77, [id](#)}, A. Vasiliev ^{139, [id](#)}, O. Vázquez
 Doce ^{95, [id](#)}, O. Vazquez Rueda ^{74, [id](#)}, V. Vechernin ^{139, [id](#)}, E. Vercellin ^{24, [id](#)}, S. Vergara Limón ⁴⁴,
 L. Vermunt ^{58, [id](#)}, R. Vértesi ^{135, [id](#)}, M. Verweij ^{58, [id](#)}, L. Vickovic ³³, Z. Vilakazi ¹¹⁹, O. Villalobos Baillie ^{99, [id](#)},
 G. Vino ^{49, [id](#)}, A. Vinogradov ^{139, [id](#)}, T. Virgili ^{28, [id](#)}, V. Vislavicius ⁸², A. Vodopyanov ^{140, [id](#)}, B. Volkel ^{32, [id](#)},
 M.A. Völkl ^{94, [id](#)}, K. Voloshin ¹³⁹, S.A. Voloshin ^{133, [id](#)}, G. Volpe ^{31, [id](#)}, B. von Haller ^{32, [id](#)}, I. Vorobyev ^{95, [id](#)},
 N. Vozniuk ^{139, [id](#)}, J. Vrláková ^{37, [id](#)}, B. Wagner ²⁰, C. Wang ^{39, [id](#)}, D. Wang ³⁹, M. Weber ^{101, [id](#)},
 A. Wegrzynek ^{32, [id](#)}, F.T. Weiglhofer ³⁸, S.C. Wenzel ^{32, [id](#)}, J.P. Wessels ^{134, [id](#)}, S.L. Weyhmler ^{136, [id](#)},
 J. Wiechula ^{63, [id](#)}, J. Wikne ^{19, [id](#)}, G. Wilk ^{78, [id](#)}, J. Wilkinson ^{97, [id](#)}, G.A. Willems ^{134, [id](#)}, B. Windelband ^{94, [id](#)},
 M. Winn ^{126, [id](#)}, J.R. Wright ^{106, [id](#)}, W. Wu ³⁹, Y. Wu ^{116, [id](#)}, R. Xu ^{6, [id](#)}, A.K. Yadav ^{131, [id](#)}, S. Yalcin ^{71, [id](#)},
 Y. Yamaguchi ^{92, [id](#)}, K. Yamakawa ⁹², S. Yang ²⁰, S. Yano ^{92, [id](#)}, Z. Yin ^{6, [id](#)}, I.-K. Yoo ^{16, [id](#)}, J.H. Yoon ^{57, [id](#)},
 S. Yuan ²⁰, A. Yuncu ^{94, [id](#)}, V. Zaccolo ^{23, [id](#)}, C. Zampolli ^{32, [id](#)}, H.J.C. Zanolli ⁵⁸, F. Zanone ^{94, [id](#)},
 N. Zardoshti ^{32,99, [id](#)}, A. Zarochentsev ^{139, [id](#)}, P. Závada ^{61, [id](#)}, N. Zaviyalov ¹³⁹, M. Zhalov ^{139, [id](#)}, B. Zhang ^{6, [id](#)},
 S. Zhang ^{39, [id](#)}, X. Zhang ^{6, [id](#)}, Y. Zhang ¹¹⁶, M. Zhao ^{10, [id](#)}, V. Zhrebchevskii ^{139, [id](#)}, Y. Zhi ¹⁰,
 N. Zhigareva ¹³⁹, D. Zhou ^{6, [id](#)}, Y. Zhou ^{82, [id](#)}, J. Zhu ^{97,6, [id](#)}, Y. Zhu ⁶, G. Zinovjev ^{3,1}, N. Zurlo ^{130,54, [id](#)}

¹ A.I. Alikhanyan National Science Laboratory (Yerevan Physics Institute) Foundation, Yerevan, Armenia

² AGH University of Krakow, Cracow, Poland

³ Bogolyubov Institute for Theoretical Physics, National Academy of Sciences of Ukraine, Kiev, Ukraine

⁴ Bose Institute, Department of Physics and Centre for Astroparticle Physics and Space Science (CAPSS), Kolkata, India

⁵ California Polytechnic State University, San Luis Obispo, CA, United States

⁶ Central China Normal University, Wuhan, China

⁷ Centro de Aplicaciones Tecnológicas y Desarrollo Nuclear (CEADEN), Havana, Cuba

⁸ Centro de Investigación y de Estudios Avanzados (CINVESTAV), Mexico City and Mérida, Mexico

⁹ Chicago State University, Chicago, IL, United States

¹⁰ China Institute of Atomic Energy, Beijing, China

¹¹ Chungbuk National University, Cheongju, Republic of Korea

¹² Comenius University Bratislava, Faculty of Mathematics, Physics and Informatics, Bratislava, Slovak Republic

¹³ COMSATS University Islamabad, Islamabad, Pakistan

¹⁴ Creighton University, Omaha, NE, United States

¹⁵ Department of Physics, Aligarh Muslim University, Aligarh, India

¹⁶ Department of Physics, Pusan National University, Pusan, Republic of Korea

¹⁷ Department of Physics, Sejong University, Seoul, Republic of Korea

¹⁸ Department of Physics, University of California, Berkeley, CA, United States

¹⁹ Department of Physics, University of Oslo, Oslo, Norway

²⁰ Department of Physics and Technology, University of Bergen, Bergen, Norway

²¹ Dipartimento di Fisica, Università di Pavia, Pavia, Italy

²² Dipartimento di Fisica dell'Università and Sezione INFN, Cagliari, Italy

- 23 Dipartimento di Fisica dell'Università and Sezione INFN, Trieste, Italy
- 24 Dipartimento di Fisica dell'Università and Sezione INFN, Turin, Italy
- 25 Dipartimento di Fisica e Astronomia dell'Università and Sezione INFN, Bologna, Italy
- 26 Dipartimento di Fisica e Astronomia dell'Università and Sezione INFN, Catania, Italy
- 27 Dipartimento di Fisica e Astronomia dell'Università and Sezione INFN, Padova, Italy
- 28 Dipartimento di Fisica 'E.R. Caianiello' dell'Università and Gruppo Collegato INFN, Salerno, Italy
- 29 Dipartimento DISAT del Politecnico and Sezione INFN, Turin, Italy
- 30 Dipartimento di Scienze MIFT, Università di Messina, Messina, Italy
- 31 Dipartimento Interateneo di Fisica 'M. Merlin' and Sezione INFN, Bari, Italy
- 32 European Organization for Nuclear Research (CERN), Geneva, Switzerland
- 33 Faculty of Electrical Engineering, Mechanical Engineering and Naval Architecture, University of Split, Split, Croatia
- 34 Faculty of Engineering and Science, Western Norway University of Applied Sciences, Bergen, Norway
- 35 Faculty of Nuclear Sciences and Physical Engineering, Czech Technical University in Prague, Prague, Czech Republic
- 36 Faculty of Physics, Sofia University, Sofia, Bulgaria
- 37 Faculty of Science, P.J. Šafárik University, Košice, Slovak Republic
- 38 Frankfurt Institute for Advanced Studies, Johann Wolfgang Goethe-Universität Frankfurt, Frankfurt, Germany
- 39 Fudan University, Shanghai, China
- 40 Gangneung-Wonju National University, Gangneung, Republic of Korea
- 41 Gauhati University, Department of Physics, Guwahati, India
- 42 Helmholtz-Institut für Strahlen- und Kernphysik, Rheinische Friedrich-Wilhelms-Universität Bonn, Bonn, Germany
- 43 Helsinki Institute of Physics (HIP), Helsinki, Finland
- 44 High Energy Physics Group, Universidad Autónoma de Puebla, Puebla, Mexico
- 45 Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest, Romania
- 46 Indian Institute of Technology Bombay (IIT), Mumbai, India
- 47 Indian Institute of Technology Indore, Indore, India
- 48 INFN, Laboratori Nazionali di Frascati, Frascati, Italy
- 49 INFN, Sezione di Bari, Bari, Italy
- 50 INFN, Sezione di Bologna, Bologna, Italy
- 51 INFN, Sezione di Cagliari, Cagliari, Italy
- 52 INFN, Sezione di Catania, Catania, Italy
- 53 INFN, Sezione di Padova, Padova, Italy
- 54 INFN, Sezione di Pavia, Pavia, Italy
- 55 INFN, Sezione di Torino, Turin, Italy
- 56 INFN, Sezione di Trieste, Trieste, Italy
- 57 Inha University, Incheon, Republic of Korea
- 58 Institute for Gravitational and Subatomic Physics (GRASP), Utrecht University/Nikhef, Utrecht, Netherlands
- 59 Institute of Experimental Physics, Slovak Academy of Sciences, Košice, Slovak Republic
- 60 Institute of Physics, Homi Bhabha National Institute, Bhubaneswar, India
- 61 Institute of Physics of the Czech Academy of Sciences, Prague, Czech Republic
- 62 Institute of Space Science (ISS), Bucharest, Romania
- 63 Institut für Kernphysik, Johann Wolfgang Goethe-Universität Frankfurt, Frankfurt, Germany
- 64 Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México, Mexico City, Mexico
- 65 Instituto de Física, Universidade Federal do Rio Grande do Sul (UFRGS), Porto Alegre, Brazil
- 66 Instituto de Física, Universidad Nacional Autónoma de México, Mexico City, Mexico
- 67 iThemba LABS, National Research Foundation, Somerset West, South Africa
- 68 Jeonbuk National University, Jeonju, Republic of Korea
- 69 Johann-Wolfgang-Goethe Universität Frankfurt Institut für Informatik, Fachbereich Informatik und Mathematik, Frankfurt, Germany
- 70 Korea Institute of Science and Technology Information, Daejeon, Republic of Korea
- 71 KTO Karatay University, Konya, Turkey
- 72 Laboratoire de Physique Subatomique et de Cosmologie, Université Grenoble-Alpes, CNRS-IN2P3, Grenoble, France
- 73 Lawrence Berkeley National Laboratory, Berkeley, CA, United States
- 74 Lund University Department of Physics, Division of Particle Physics, Lund, Sweden
- 75 Nagasaki Institute of Applied Science, Nagasaki, Japan
- 76 Nara Women's University (NWU), Nara, Japan
- 77 National and Kapodistrian University of Athens, School of Science, Department of Physics, Athens, Greece
- 78 National Centre for Nuclear Research, Warsaw, Poland
- 79 National Institute of Science Education and Research, Homi Bhabha National Institute, Jatni, India
- 80 National Nuclear Research Center, Baku, Azerbaijan
- 81 National Research and Innovation Agency - BRIN, Jakarta, Indonesia
- 82 Niels Bohr Institute, University of Copenhagen, Copenhagen, Denmark
- 83 Nikhef, National institute for subatomic physics, Amsterdam, Netherlands
- 84 Nuclear Physics Group, STFC Daresbury Laboratory, Daresbury, United Kingdom
- 85 Nuclear Physics Institute of the Czech Academy of Sciences, Husinec-Řež, Czech Republic
- 86 Oak Ridge National Laboratory, Oak Ridge, TN, United States
- 87 Ohio State University, Columbus, OH, United States
- 88 Physics department, Faculty of science, University of Zagreb, Zagreb, Croatia
- 89 Physics Department, Panjab University, Chandigarh, India
- 90 Physics Department, University of Jammu, Jammu, India
- 91 Physics Department, University of Rajasthan, Jaipur, India
- 92 Physics Program and International Institute for Sustainability with Knotted Chiral Meta Matter (SKCM2), Hiroshima University, Hiroshima, Japan
- 93 Physikalisches Institut, Eberhard-Karls-Universität Tübingen, Tübingen, Germany
- 94 Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany
- 95 Physik Department, Technische Universität München, Munich, Germany
- 96 Politecnico di Bari and Sezione INFN, Bari, Italy
- 97 Research Division and ExtreMe Matter Institute EMMI, GSI Helmholtzzentrum für Schwerionenforschung GmbH, Darmstadt, Germany
- 98 Saha Institute of Nuclear Physics, Homi Bhabha National Institute, Kolkata, India
- 99 School of Physics and Astronomy, University of Birmingham, Birmingham, United Kingdom
- 100 Sección Física, Departamento de Ciencias, Pontificia Universidad Católica del Perú, Lima, Peru
- 101 Stefan Meyer Institut für Subatomare Physik (SMI), Vienna, Austria
- 102 SUBATECH, IMT Atlantique, Nantes Université, CNRS-IN2P3, Nantes, France

- 103 Suranaree University of Technology, Nakhon Ratchasima, Thailand
 104 Technical University of Košice, Košice, Slovak Republic
 105 The Henryk Niewodniczanski Institute of Nuclear Physics, Polish Academy of Sciences, Cracow, Poland
 106 The University of Texas at Austin, Austin, TX, United States
 107 Universidad Autónoma de Sinaloa, Culiacán, Mexico
 108 Universidade de São Paulo (USP), São Paulo, Brazil
 109 Universidade Estadual de Campinas (UNICAMP), Campinas, Brazil
 110 Universidade Federal do ABC, Santo Andre, Brazil
 111 University of Cape Town, Cape Town, South Africa
 112 University of Houston, Houston, TX, United States
 113 University of Jyväskylä, Jyväskylä, Finland
 114 University of Kansas, Lawrence, KS, United States
 115 University of Liverpool, Liverpool, United Kingdom
 116 University of Science and Technology of China, Hefei, China
 117 University of South-Eastern Norway, Kongsberg, Norway
 118 University of Tennessee, Knoxville, TN, United States
 119 University of the Witwatersrand, Johannesburg, South Africa
 120 University of Tokyo, Tokyo, Japan
 121 University of Tsukuba, Tsukuba, Japan
 122 University Politehnica of Bucharest, Bucharest, Romania
 123 Université Clermont Auvergne, CNRS/IN2P3, LPC, Clermont-Ferrand, France
 124 Université de Lyon, CNRS/IN2P3, Institut de Physique des 2 Infinis de Lyon, Lyon, France
 125 Université de Strasbourg, CNRS, IPHC UMR 7178, F-67000 Strasbourg, France
 126 Université Paris-Saclay, Centre d'Etudes de Saclay (CEA), IRFU, Département de Physique Nucléaire (DPhN), Saclay, France
 127 Université Paris-Saclay, CNRS/IN2P3, IJCLab, Orsay, France
 128 Università degli Studi di Foggia, Foggia, Italy
 129 Università del Piemonte Orientale, Vercelli, Italy
 130 Università di Brescia, Brescia, Italy
 131 Variable Energy Cyclotron Centre, Homi Bhabha National Institute, Kolkata, India
 132 Warsaw University of Technology, Warsaw, Poland
 133 Wayne State University, Detroit, MI, United States
 134 Westfälische Wilhelms-Universität Münster, Institut für Kernphysik, Münster, Germany
 135 Wigner Research Centre for Physics, Budapest, Hungary
 136 Yale University, New Haven, CT, United States
 137 Yonsei University, Seoul, Republic of Korea
 138 Zentrum für Technologie und Transfer (ZTT), Worms, Germany
 139 Affiliated with an institute covered by a cooperation agreement with CERN
 140 Affiliated with an international laboratory covered by a cooperation agreement with CERN

^I Deceased.

^{II} Also at: Max-Planck-Institut für Physik, Munich, Germany.

^{III} Also at: Italian National Agency for New Technologies, Energy and Sustainable Economic Development (ENEA), Bologna, Italy.

^{IV} Also at: Dipartimento DET del Politecnico di Torino, Turin, Italy.

^V Also at: Department of Applied Physics, Aligarh Muslim University, Aligarh, India.

^{VI} Also at: Institute of Theoretical Physics, University of Wrocław, Poland.

^{VII} Also at: An institution covered by a cooperation agreement with CERN.