

JYX



This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): CUPID-Mo Collaboration

Title: Measurement of the $2\nu\beta\beta$ Decay Rate and Spectral Shape of ^{100}Mo from the CUPID-Mo Experiment

Year: 2023

Version: Published version

Copyright: © 2023 American Physical Society

Rights: In Copyright

Rights url: <http://rightsstatements.org/page/InC/1.0/?language=en>

Please cite the original version:

CUPID-Mo Collaboration. (2023). Measurement of the $2\nu\beta\beta$ Decay Rate and Spectral Shape of ^{100}Mo from the CUPID-Mo Experiment. *Physical Review Letters*, 131, Article 162501.

<https://doi.org/10.1103/PhysRevLett.131.162501>

Measurement of the $2\nu\beta\beta$ Decay Rate and Spectral Shape of ^{100}Mo from the CUPID-Mo Experiment

C. Augier,¹ A. S. Barabash,² F. Bellini,^{3,4} G. Benato,^{5,6} M. Beretta,⁷ L. Bergé,⁸ J. Billard,¹ Yu. A. Borovlev,⁹ L. Cardani,⁴ N. Casali,⁴ A. Cazes,¹ E. Celi,^{5,6} M. Chapellier,⁸ D. Chiesa,^{10,11} I. Dafinei,⁴ F. A. Danevich,^{12,13} M. De Jesus,¹ T. Dixon,^{8,14,*} L. Dumoulin,⁸ K. Eitel,¹⁵ F. Ferri,¹⁴ B. K. Fujikawa,¹⁶ J. Gascon,¹ L. Gironi,^{10,11} A. Giuliani,⁸ V. D. Grigorieva,⁹ M. Gros,¹⁴ D. L. Helis,^{14,5} H. Z. Huang,¹⁷ R. Huang,⁷ L. Imbert,⁸ J. Johnston,¹⁸ A. Juillard,¹ H. Khalife,⁸ M. Kleifges,¹⁹ V. V. Kobychyev,¹² Yu. G. Kolomensky,^{7,16} S. I. Konovalov,²⁰ J. Kotila,^{21,22,23} P. Loaiza,⁸ L. Ma,¹⁷ E. P. Makarov,⁹ P. de Marcillac,⁸ R. Mariam,⁸ L. Marini,^{7,16,5} S. Marnieros,⁸ X.-F. Navick,¹⁴ C. Nones,¹⁴ E. B. Norman,⁷ E. Olivieri,⁸ J. L. Ouellet,¹⁸ L. Pagnanini,^{6,5} L. Pattavina,^{5,24} B. Paul,¹⁴ M. Pavan,^{10,11} H. Peng,²⁵ G. Pessina,¹¹ S. Pirro,⁵ D. V. Poda,⁸ O. G. Polischuk,^{12,4} S. Pozzi,¹¹ E. Previtali,^{10,11} Th. Redon,⁸ A. Rojas,²⁶ S. Rozov,²⁷ V. Sanglard,¹ J. A. Scarpaci,⁸ B. Schmidt,¹⁴ Y. Shen,¹⁷ V. N. Shlegel,⁹ F. Šimkovic,^{28,29} V. Singh,⁷ C. Tomei,⁴ V. I. Tretyak,^{12,5} V. I. Umatov,²⁰ L. Vagneron,¹ M. Velázquez,³⁰ B. Ware,³¹ B. Welliver,⁷ L. Winslow,¹⁸ M. Xue,²⁵ E. Yakushev,²⁷ M. Zarytsky,¹² and A. S. Zolotarova⁸

(CUPID-Mo Collaboration)

¹Univ Lyon, Université Lyon 1, CNRS/IN2P3, IP2I-Lyon, F-69622 Villeurbanne, France

²National Research Centre “Kurchatov Institute,” Kurchatov Complex of Theoretical and Experimental Physics, 117218 Moscow, Russia

³Dipartimento di Fisica, Sapienza Università di Roma, Piazzale Aldo Moro 2, I-00185 Rome, Italy

⁴INFN, Sezione di Roma, P.le Aldo Moro 2, I-00185 Rome, Italy

⁵INFN, Laboratori Nazionali del Gran Sasso, I-67100 Assergi (AQ), Italy

⁶INFN, Gran Sasso Science Institute, I-67100 L'Aquila, Italy

⁷University of California, Berkeley, California 94720, USA

⁸Université Paris-Saclay, CNRS/IN2P3, IJCLab, 91405 Orsay, France

⁹Nikolaev Institute of Inorganic Chemistry, 630090 Novosibirsk, Russia

¹⁰Dipartimento di Fisica, Università di Milano-Bicocca, I-20126 Milano, Italy

¹¹INFN, Sezione di Milano-Bicocca, I-20126 Milano, Italy

¹²Institute for Nuclear Research of NASU, 03028 Kyiv, Ukraine

¹³INFN, Sezione di Roma Tor Vergata, Via della Ricerca Scientifica 1, I-00133 Rome, Italy

¹⁴IRFU, CEA, Université Paris-Saclay, F-91191 Gif-sur-Yvette, France

¹⁵Karlsruhe Institute of Technology, Institute for Astroparticle Physics, 76021 Karlsruhe, Germany

¹⁶Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA

¹⁷Key Laboratory of Nuclear Physics and Ion-beam Application (MOE), Fudan University, Shanghai 200433, People's Republic of China

¹⁸Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA

¹⁹Karlsruhe Institute of Technology, Institute for Data Processing and Electronics, 76021 Karlsruhe, Germany

²⁰National Research Centre Kurchatov Institute, Institute of Theoretical and Experimental Physics, 117218 Moscow, Russia

²¹Department of Physics, University of Jyväskylä, P.O. Box 35, FI-40014 Jyväskylä, Finland

²²Finnish Institute for Educational Research, University of Jyväskylä, P.O. Box 35, FI-40014 Jyväskylä, Finland

²³Center for Theoretical Physics, Sloane Physics Laboratory, Yale University, New Haven, Connecticut 06520-8120, USA

²⁴Physik Department, Technische Universität München, Garching D-85748, Germany

²⁵Department of Modern Physics, University of Science and Technology of China, Hefei 230027, People's Republic of China

²⁶LSM, Laboratoire Souterrain de Modane, 73500 Modane, France

²⁷Laboratory of Nuclear Problems, JINR, 141980 Dubna, Moscow region, Russia

²⁸Faculty of Mathematics, Physics and Informatics, Comenius University in Bratislava, 842 48 Bratislava, Slovakia

²⁹Institute of Experimental and Applied Physics, Czech Technical University in Prague, 128 00 Prague, Czech Republic

³⁰Université Grenoble Alpes, CNRS, Grenoble INP, SIMAP, 38420 Saint Martin d'Hères, France

³¹John de Laeter Centre for Isotope Research, GPO Box U 1987, Curtin University, Bentley, Western Australia, Australia



(Received 26 July 2023; revised 29 August 2023; accepted 21 September 2023; published 18 October 2023)

Neutrinoless double beta decay ($0\nu\beta\beta$) is a yet unobserved nuclear process that would demonstrate Lepton number violation, a clear evidence of beyond standard model physics. The process two neutrino double beta decay ($2\nu\beta\beta$) is allowed by the standard model and has been measured in numerous experiments. In this Letter, we report a measurement of $2\nu\beta\beta$ decay half-life of ^{100}Mo to the ground state of ^{100}Ru of $[7.07 \pm 0.02(\text{stat}) \pm 0.11(\text{syst})] \times 10^{18}$ yr by the CUPID-Mo experiment. With a relative precision of $\pm 1.6\%$ this is the most precise measurement to date of a $2\nu\beta\beta$ decay rate in ^{100}Mo . In addition, we constrain higher-order corrections to the spectral shape, which provides complementary nuclear structure information. We report a novel measurement of the shape factor $\xi_{3,1} = 0.45 \pm 0.03(\text{stat}) \pm 0.05(\text{syst})$ based on a constraint on the ratio of higher-order terms from theory, which can be reliably calculated. This is compared to theoretical predictions for different nuclear models. We also extract the first value for the effective axial vector coupling constant obtained from a spectral shape study of $2\nu\beta\beta$ decay.

DOI: 10.1103/PhysRevLett.131.162501

For more than 20 years it has been known that neutrinos have mass via measurements of neutrino oscillations [1,2]. This raises the question of the nature of this mass. If the neutrino is its own antiparticle, a *Majorana* particle, then a decay mode of some nuclei would become possible, neutrinoless double beta decay ($0\nu\beta\beta$) (see reviews [3–5]). This decay could be observed in nuclei for which single beta decay is energetically disallowed (or disfavored by angular momentum). Two neutrons would be transformed into two protons, with the emission of only two electrons. The observation of this decay would have profound consequences for particle physics by showing that the Lepton number is not a fundamental symmetry of nature and providing clear evidence of beyond standard model physics.

The measurement of the decay rate could also provide a method to measure the effective neutrino mass [6]. Under the light Majorana neutrino exchange mechanism the decay rate would be related to the effective Majorana mass $\langle m_{\beta\beta} \rangle$ by

$$1/T_{1/2}^{0\nu} = G_{0\nu} \cdot g_A^4 \cdot |M_{0\nu}|^2 \cdot \langle m_{\beta\beta} \rangle^2 / m_e^2, \quad (1)$$

where $G_{0\nu}$ is the phase space factor, $M_{0\nu}$ the nuclear matrix element (NME), g_A the effective axial-vector coupling constant, and m_e the electron mass. While $G_{0\nu}$ can be calculated almost exactly [7], the NME is the result of complex many-body nuclear physics calculations (see the review [8]) and is only known to a factor of a few. To interpret the results of next-generation experiments these calculations must be improved. In addition, it has been observed that nuclear models often overpredict the decay rate of β^- and $2\nu\beta\beta$. To account for this g_A can be replaced with an effective value $g_{A,\text{eff}}$ [9–11]. Therefore there is still a possibility the $0\nu\beta\beta$ decay rate could be much lower than expected for an unrenormalized value of g_A (1.27). This would have significant impact on the discovery probability of next-generation experiments [12,13]. To constrain this possibility new measurements are needed.

Two neutrino double beta decay ($2\nu\beta\beta$) conserves Lepton number and is allowed within the standard model. It has been observed in a number of nuclei [14]. The decay rate of $2\nu\beta\beta$ decay can be described to a good approximation as

$$1/T_{1/2}^{2\nu} = G_{2\nu} \cdot g_A^4 \cdot |M_{2\nu}|^2, \quad (2)$$

where $G_{2\nu}$ is the phase space factor, $M_{2\nu}$ is the NME. Since $0\nu\beta\beta$ and $2\nu\beta\beta$ share the same initial and final nuclear states, an accurate prediction of $T_{1/2}^{2\nu}$ and therefore an accurate description of the nuclear structure, is a necessary condition to obtain reliable estimates of $M_{0\nu}$. These measurements are often used to tune the parameters of the nuclear models. However, they cannot alone answer questions about the value of $g_{A,\text{eff}}$ since only the product $M_{2\nu} \times g_{A,\text{eff}}^2$ is measured.

The $2\nu\beta\beta$ decay spectrum is typically described using two approximations: the single and higher state dominance hypotheses (SSD/HSD) [15]. In these approximations, the decay is supposed to proceed via a single intermediate 1^+ state. For the HSD model this state is an average higher energy state from the region of the Gamow-Teller resonance, while for SSD it is the lowest energy 1^+ state.

The description of the $2\nu\beta\beta$ decay spectrum was improved in [16,17]. In this approach, a Taylor expansion is performed in terms of the Lepton energies. The differential decay rate relates to phase space factors and NMEs as

$$\frac{d\Gamma}{dE} = g_{A,\text{eff}}^4 |M_{GT-1}|^2 \left(\frac{dG_0}{dE} + \xi_{3,1} \frac{dG_2}{dE} + \frac{1}{3} \xi_{3,1}^2 \frac{dG_{22}}{dE} + \left(\frac{1}{3} \xi_{3,1}^2 + \xi_{5,1} \right) \frac{dG_4}{dE} \right). \quad (3)$$

Here, G_0 , G_2 , G_{22} , G_4 are the phase space factors for different terms in the Taylor expansion. $\xi_{3,1} = M_{GT-3}/M_{GT-1}$ and $\xi_{5,1} = M_{GT-5}/M_{GT-1}$ are ratios of NMEs. By fitting the energy distribution of electrons to

this model, constraints on $\xi_{3,1}, \xi_{5,1}$ can be obtained that can be compared to theoretical predictions. M_{GT-3} and M_{GT-5} are expected to be dominated by contributions from lower-lying states due to the higher power of the energy denominators so measurement of $\xi_{3,1}, \xi_{5,1}$ provide complementary nuclear structure information to the half-life. Within this model the HSD spectrum can be recovered by fixing $\xi_{3,1}, \xi_{5,1}$ to zero, and the SSD approximation can be used to predict nonzero $\xi_{3,1}, \xi_{5,1}$ as in [16]. ξ values larger than the SSD values would indicate mutual cancellation between lower and higher lying states.

As described in [16], a measurement of $\xi_{3,1}$ and the half-life can be used to extract a value for $g_{A,\text{eff}}$ of

$$g_{A,\text{eff}}^4 = \frac{T_{1/2}^{-1} \times \xi_{3,1}^2}{M_{GT-3}^2 G}, \quad (4)$$

where $G = G_0 + \xi_{3,1}G_2 + \xi_{3,1}^2 G_{22}/3 + (\xi_{3,1}^2/3 + \xi_{5,1})G_4$. M_{GT-3} can be computed reliably within the interacting shell model (ISM), which describes accurately low lying states of nuclei.

So far the analysis to extract the ξ factors has only been performed by the KamLAND-Zen experiment [18], which established an upper bound on $\xi_{3,1}$ in ^{136}Xe decays, which is still compatible with both the ISM and pn-QRPA calculations.

We compute $2\nu\beta\beta$ NMEs M_{GT-1} , M_{GT-3} , and M_{GT-5} within the proton-neutron quasiparticle random-phase approximation (pn-QRPA) [19,20] and described in more detail in the Supplemental Material [21]. These calculations are performed for a range of $g_{A,\text{eff}}$ values. We compute the phase space factors G_0, G_2, G_4, G_{22} considering Dirac wave functions with finite nuclear size and electron screening as in [7].

The experimental signature of $2\nu\beta\beta$ decay is a continuous spectrum in the summed energies of the electrons. Differentiation of the signal from background is more challenging than for $0\nu\beta\beta$ decay: the decay rate must be extracted from a fit to the full spectrum using detailed simulations of the various contributions to the experimental background (see, for example, [22–25]).

Therefore, a very low background is imperative to make a precise measurement. Scintillating cryogenic calorimeters provide a technique to reach very low background rates [26–28]. In particular, a scintillation light signal in coincidence with a heat signal in the calorimeter can be used to remove α particle backgrounds [28–31].

In this Letter, we describe a measurement of the $2\nu\beta\beta$ decay rate and spectral shape of ^{100}Mo using the CUPID-Mo experiment, a demonstrator for the next-generation $0\nu\beta\beta$ decay experiment CUPID [32]. A detailed description of the experiment can be found in [33]. It consisted of an array of 20 lithium molybdate (LMO) cryogenic calorimeters, enriched in ^{100}Mo ($96.6 \pm 0.2\%$ isotope

abundance) each of around 200 g mass. In addition, 20 germanium light detectors (LD), also operated as cryogenic calorimeters, were employed to readout the scintillation light signal used for particle identification to remove the α particle background. An individual module of CUPID-Mo consisted of an LMO crystal attached to a copper holder and a Ge LD. Both the LMO and LD signals were read out using neutron transmutation doped germanium thermistors [34]. These modules were then arranged into five towers of four LMOs each and installed in the EDELWEISS cryostat [35] at the Laboratoire Souterrain de Modane, France. It collected a total exposure of $1.48 \text{ kg} \times \text{yr}$ of ^{100}Mo between 2019 and 2020. The scintillation light signal allowed a complete rejection of α particles, while an excellent energy resolution of $7.7 \pm 0.4 \text{ keV}$ FWHM was measured at 3034 keV [36]. This performance lead to a limit on $0\nu\beta\beta$ in ^{100}Mo of $T_{1/2}^{0\nu} > 1.8 \times 10^{24} \text{ yr}$ (90% credible interval) [36].

For this analysis we use the full data collected by CUPID-Mo. A detailed description of the data processing is given in [36] and was also used for [37,38]. An optimal filter based analysis chain [39], which maximizes the signal to noise ratio, is used to select physics events and estimate pulse amplitudes. Spurious events, such as pileup or spikes induced by electronics, are removed using a principal component analysis based pulse shape cut [36,40], normalized to ensure an energy independent efficiency. Because of the relatively short range of electrons in LMO, both $0\nu\beta\beta$ and $2\nu\beta\beta$ to ground states are likely to deposit energy in just a single LMO detector. However, background events induced by γ quanta are more likely to deposit energy in multiple crystals. As such we define the “multiplicity” (\mathcal{M}) of an event as the number of LMO detectors with a pulse above the 40 keV energy threshold within a $\pm 10 \text{ ms}$ window. In addition, muon induced events are excluded using a dedicated muon veto system [41]. We select β, γ -like events using the scintillation light signal as described in detail in [36]. We also remove events with a trigger in one LD with high ^{60}Co contamination as described in [37,38]. Multiplicity one γ/β ($\mathcal{M}_{1,\gamma/\beta}$) events are used to extract the signal rates while \mathcal{M}_2 are used to constrain the γ background. We also extract the spectra at high energy without any α rejection, this dataset ($\mathcal{M}_{1,\alpha}$) is used to constrain the radioactivity of the LMO crystals and other nearby components.

The energy resolution and bias in the energy scale are measured using γ lines. The efficiency of all selection cuts has been estimated as $88.9 \pm 1.1\%$ for $\mathcal{M}_{1,\gamma/\beta}$ [36,38]. No evidence of energy dependence was found, over the range of the fit, and our cuts are normalized to have an energy-independent efficiency.

To extract the rate of $2\nu\beta\beta$ decay we construct a model of the data described in detail in [38]. We simulate using Geant4 [42] the $2\nu\beta\beta$ signal, using both the SSD and HSD models parametrized from [7] and the contributions to the




FIG. 1. Fit of the $\mathcal{M}_{1,\gamma/\beta}$ spectrum showing the main contributions to the model and the residuals defined as $(\text{data} - \text{fit})/\sqrt{\text{data}}$. The model describes well the experimental data and the spectrum above ~ 500 keV is dominated by $2\nu\beta\beta$ events.

improved $2\nu\beta\beta$ model from Eq. (3) [16]. We also simulate radioactive contaminations in the various components of the experimental setup. These simulations are then convolved with a detector response model consisting of the energy resolution of the detectors, energy threshold, coincidences, and dead times of the detectors.

We use a Bayesian analysis based on JAGS [43,44] to fit our three experimental spectra ($\mathcal{M}_{1,\gamma/\beta}$, $\mathcal{M}_{1,\alpha}$, \mathcal{M}_2) to a sum of Monte Carlo (MC) simulations. The details of the choices of the model components are given in [38]. The fit to the $\mathcal{M}_{1,\gamma/\beta}$ spectrum uses a range of 100–4000 keV. A variable binning is used so that at minimum 15 keV bins are used in the continuum region and then bins are combined so at least 15 events are in each bin. Each γ or α peak is placed in one bin to avoid the systematic effect of the peak line shape. We show the $\mathcal{M}_{1,\gamma/\beta}$ fit in Fig. 1. We call this fit our “reference fit”: we see that this model is able to describe all the features of the experimental data, and that the data are dominated by $2\nu\beta\beta$ decay events. While in [38] the SSD model of $2\nu\beta\beta$ was used by default, for this work we instead use the improved $2\nu\beta\beta$ model, and consider SSD as a cross-check. By using the improved model, which allows the spectral shape to vary during the fit, we marginalize over the theoretical uncertainty in the spectral shape.

We study the consistency between our model and data using pseudoexperiments. We generate from the best fit model a set of 1000 pseudoexperiments, and for each we perform the background model fit and extract $-\log(\mathcal{L})$. The value obtained for the $\mathcal{M}_{1,\gamma/\beta}$ data is consistent with the expected distribution. In particular, we extract a p value, or the probability of observing equal or larger fluctuations of 0.54.

From the background model fit we extract the $2\nu\beta\beta$ decay rate. We consider systematic uncertainties related to the number of reconstructed events and the efficiency and

isotopic abundance conversion factors. We have performed a series of tests varying the assumptions of our background model to assess the dependence of $T_{1/2}$ on these choices. For each test a probability distribution is assumed for the systematic uncertainty based on the change in the best fit value with respect to our reference fit. We then compute a convolution of these distributions and the posterior distribution from the fit to obtain the posterior distribution considering all systematic uncertainties. This can be considered a generalization of adding in quadrature to non-Gaussian uncertainties.

First we perform tests to check the dependence of our results on the γ radioactivity source location. We remove far sources of Th/U radioactivity leading to a slightly lower $2\nu\beta\beta$ rate (-0.83%) and then close (10 mK) sources of Th/U radioactivity leading to a higher rate ($+0.22\%$). In principle, this uncertainty is already marginalized over in our analysis. However, our fit favors far sources of radioactivity, possibly due to some other effects such as pure β decays, so we take a conservative approach considering an uncertainty of $\pm 0.83\%$ from the first test. This is assigned a Gaussian distribution to account for the possibility of even further sources than those included in our model.

Anthropogenic β^- decays could contribute to our background. In our model we include a source of $^{90}\text{Sr} + ^{90}\text{Y}$, consisting of two pure β^- decays with Q values 546, 2276 keV and ~ 60 h delay. This is one of the only anthropogenic contaminations with a large enough Q value to correlate with $2\nu\beta\beta$ decay and a relatively long half-life. In our model the activity is constrained as 179_{-32}^{+36} $\mu\text{Bq/kg}$. Since the convergence of this parameter is driven by events at low energy that could have several origins we repeat the fit without this contribution. We obtain a half-life value $+1.0\%$ higher than the reference and we assign a uniform probability distribution between the reference and this fit.

We repeat the fit removing any contributions where the smallest 68% interval contains zero activity, which we call the “minimal model.” In our analysis, all the contributions are assigned non-negative uniform priors; therefore, a large number of parameters could bias the fit leading to a smaller $2\nu\beta\beta$ rate. We find a small shift of $+0.24\%$ in the $2\nu\beta\beta$ decay rate for this fit. We assign a Gaussian distribution with 0.24% uncertainty for this systematic uncertainty. We also check that our fit is not biased using our set of pseudoexperiments. The distribution of obtained $T_{1/2}$ is consistent with the fit to data.

We perform fits varying the energy scale by ∓ 1 keV resulting in a $2\nu\beta\beta$ decay rate shifted by ${}_{-0.16}^{+0.11}\%$, which we assign an asymmetric-Gaussian distribution.

Our reference fit uses a variable binning described in [38]. We repeat the fit using fixed binning of 1, 2, 10, 20, and 30 keV. The largest effect is for a binning of 2 keV where the rate is reduced by -0.37% . We take a conservative approach considering a Gaussian distribution with $\pm 0.37\%$ standard deviation.

TABLE I. Systematic uncertainties in the determination of the $2\nu\beta\beta$ decay rate and $\xi_{3,1}$. All uncertainties are assigned either a Gaussian or asymmetric-Gaussian (for asymmetric uncertainties) posterior distribution with the exception of the $^{90}\text{Sr} + ^{90}\text{Y}$, where we assign a uniform distribution.

Systematic test	Uncertainties	
	$T_{1/2}$ [%]	$\xi_{3,1}$ [%]
Source location	0.83	0.9
$^{90}\text{Sr} + ^{90}\text{Y}$	+1.0 ^a	-4.9 ^a
Minimal model	0.24	7.7
Binning	0.37	1.4
Energy bias	+0.11 -0.16	+3.5 -3.7
Bremsstrahlung	+0.13 -0.22	+6.0 -6.8
MC statistics	0.11	1.4
Efficiency	1.2	...
Isotopic abundance	0.2	...

^aUniform distribution.

To assess the dependence on the accuracy of the MC simulations we generate simulations of $2\nu\beta\beta$ decay where we vary the Bremsstrahlung cross section by $\pm 10\%$. These lead to $^{+0.13}_{-0.22}\%$ change in the $2\nu\beta\beta$ rate, to which we assign an asymmetric-Gaussian distribution.

To account for the statistical uncertainty in the MC simulations we perform a fit adding nuisance parameters to the model as is done in [22]. This leads to a -0.11% smaller $2\nu\beta\beta$ rate, which we consider a systematic with a Gaussian distribution.

The final systematic uncertainties are on selection efficiency and ^{100}Mo abundance, which are 1.2% and 0.2%, respectively, and are assigned Gaussian distributions. These systematic uncertainties are summarized in Table I.

Computing the convolution of all systematic uncertainties in Table I and converting to the decay rate we compute the posterior distribution of $T_{1/2}^{-1}$ (see Ref. [21]) for both the statistical only uncertainty and the combined uncertainty. From the central 68% credible interval we extract a measurement of

$$T_{1/2}^{2\nu} = [7.07 \pm 0.02(\text{stat}) \pm 0.11(\text{syst})] \times 10^{18} \text{ yr}. \quad (5)$$

With a relative uncertainty of $\pm 1.6\%$ this is one of the most precise determinations of a $2\nu\beta\beta$ decay half-life. The half-life is in agreement with our previous result obtained with a much smaller exposure [24], the value from NEMO-3 [45] and one obtained using the SSD $2\nu\beta\beta$ spectral shape model.

Next, we extract the values of the shape factors from the fit. We find a clear contribution from higher-order terms with a mild preference for a contribution from $\xi_{5,1}$ instead of $\xi_{3,1}$ (see more details in Supplemental Material [21]). However, the parameters $\xi_{3,1}$ and $\xi_{5,1}$ are strongly anti-correlated with $\rho = -0.92$. Thus, the fit is not sensitive to whether this contribution originates from $\xi_{3,1}$ or $\xi_{5,1}$.




FIG. 2. Posterior distribution of $\xi_{3,1}$ both with and without convolution with the systematic uncertainties (upper figure). In the lower panel we compare to the pn-QRPA, ISM and SSD theoretical values as a function of $g_{A,\text{eff}}$ for two potentials (CD-Bonn and Argonne V18; see Supplemental Material [21] for more details).

However, within nuclear structure calculations the value of $\xi_{5,1}/\xi_{3,1}$ can be calculated reliably since M_{GT-3} and M_{GT-5} depend on contributions from low lying states. The value of $\xi_{5,1}/\xi_{3,1}$ within pn-QRPA is 0.364–0.368 depending on $g_{A,\text{eff}}$ and the nuclear potential. Within the SSD hypothesis the value is 0.367 [16] and within the ISM it is 0.349 [46,47]. To reduce the degeneracy in our model we perform a fit with a Gaussian prior on $\xi_{5,1}/\xi_{3,1}$ with a mean of the SSD prediction and a conservative 5% uncertainty.

From this fit we extract the value of $\xi_{3,1}$ to compare to theoretical predictions. We consider the same systematic uncertainties as for the half-life (also shown in Table I). The largest effects are found to be from the MC bremsstrahlung cross section and the choice of parameters of the model. The posterior distribution of this observable both before and after convolution with the systematics is shown in Fig. 2. We extract a measurement:

$$\xi_{3,1} = 0.45 \pm 0.03(\text{stat}) \pm 0.05(\text{syst}). \quad (6)$$

We compare our measurement of $\xi_{3,1}$ to pn-QRPA theoretical predictions in the lower panel of Fig. 2. Within pn-QRPA the g_{pp} parameter (the strength of the particle-particle interaction) is tuned using our measurement of the half-life for each $g_{A,\text{eff}}$. Since the calculated values of $\xi_{3,1}$ depend on $g_{A,\text{eff}}$, our measurement of $\xi_{3,1}$

provides complementary information on $g_{A,\text{eff}}$. We find the experimental value is incompatible ($\sim 8\sigma$) with the prediction of the HSD hypothesis of $\xi_{3,1} = \xi_{5,1} = 0$, somewhat incompatible with that from the ISM ($\sim 2.1\sigma$) but mostly compatible with that from the SSD hypothesis ($\sim 1.4\sigma$) and the pn-QRPA predictions if the value of $g_{A,\text{eff}}$ is moderately quenched (> 0.8) or unquenched. We encourage computation of $\xi_{3,1}$ and $\xi_{5,1}$ in additional theoretical frameworks such as the interacting boson model [48]. To extract a value for $g_{A,\text{eff}}$ within the pn-QRPA framework, we sample from the distribution of $\xi_{3,1}$ from our fit and for each sample we extract the corresponding $g_{A,\text{eff}}$ values. Assigning equal weights to the CD-Bonn and Argonne V-18 nuclear potentials we extract a posterior distribution on $g_{A,\text{eff}}$ and thus we extract a value:

$$g_{A,\text{eff}}(\text{pn-QRPA}) = 1.0 \pm 0.1(\text{stat}) \pm 0.2(\text{syst}). \quad (7)$$

As mentioned previously an analysis of $\xi_{3,1}$ and the half-life can be used to extract a measurement of $g_{A,\text{eff}}$ if M_{GT-3} is known [see Eq. (4)]. Using the value of M_{GT-3} from the ISM [46,47] and our fit we reconstruct

$$g_{A,\text{eff}}(\text{ISM}) = 1.11 \pm 0.03(\text{stat}) \pm 0.05(\text{syst}). \quad (8)$$

The statistical uncertainty is obtained by sampling from the Markov chain; therefore, combining the uncertainties on $T_{1/2}$ and $\xi_{3,1}$, the systematic uncertainty is obtained from the same tests as previously considered. This is the first measurement of $g_{A,\text{eff}}$ from a spectral shape study of $2\nu\beta\beta$ decay.

In this Letter, we have reported a measurement of the $2\nu\beta\beta$ decay half-life of ^{100}Mo . Utilizing excellent background rejection, a very clean spectrum is obtained that allowed us to obtain the most precise ever measurement of a $2\nu\beta\beta$ decay rate in this isotope. Special attention was paid to the systematic uncertainties affecting the result, particularly to the source location, model choices, and MC accuracy.

In addition, we obtained a first of its kind measurement of a novel nuclear structure observable $\xi_{3,1}$ based on an improved description of the $2\nu\beta\beta$ decay process. The value of this observable is found to be incompatible with an HSD prediction, mildly incompatible with predictions from the ISM, but compatible with pn-QRPA predictions and a moderately quenched or unquenched value of $g_{A,\text{eff}}$. Finally, we report two novel measurements of $g_{A,\text{eff}}$, the first of their kind obtained from a spectral shape study of a $2\nu\beta\beta$ decay.

This work has been performed in the framework of the CUPID-1 and P2IO LabEx programs, funded by the Agence Nationale de la Recherche (ANR, France). F. A. D., V. V. K., V. I. T., and M. M. Z. were supported in part by the National Research Foundation of Ukraine. A. S. B., S. I. K.,

I. M. M., V. N. S., and V. I. U. were supported by the Russian Science Foundation. J. K. is supported by Academy of Finland. F. D. was partly supported by the Slovak Research and Development Agency. Additionally the work is supported by the Istituto Nazionale di Fisica Nucleare (INFN), by the US Department of Energy (DOE) Office of Science. This work makes use of the *Diana* data analysis software and the background model based on JAGS, developed by the CUORICINO, CUORE, LUCIFER, and CUPID-0 collaborations.

*Corresponding author: toby.dixon@ijclab.in2p3.fr

- [1] Y. Fukuda *et al.* (Super-Kamiokande Collaboration), *Phys. Rev. Lett.* **81**, 1562 (1998).
- [2] Q. R. Ahmad *et al.*, *Phys. Rev. Lett.* **89**, 011301 (2002).
- [3] S. Dell’Oro, S. Marcocci, M. Viel, and F. Vissani, *Adv. High Energy Phys.* **2016**, 2162659 (2016).
- [4] M. Agostini, G. Benato, J. A. Detwiler, J. Menéndez, and F. Vissani, *Rev. Mod. Phys.* **95**, 025002 (2023).
- [5] M. J. Dolinski, A. W. Poon, and W. Rodejohann, *Annu. Rev. Nucl. Part. Sci.* **69**, 219 (2019).
- [6] G. Benato, *Eur. Phys. J. C* **75**, 563 (2015).
- [7] J. Kotila and F. Iachello, *Phys. Rev. C* **85**, 034316 (2012).
- [8] J. Engel and J. Menéndez, *Rep. Prog. Phys.* **80**, 046301 (2017).
- [9] J. T. Suhonen, *Front. Phys.* **5**, 55 (2017).
- [10] H. Ejiri, J. Suhonen, and K. Zuber, *Phys. Rep.* **797**, 1 (2019).
- [11] G. Martínez-Pinedo, A. Poves, E. Caurier, and A. P. Zuker, *Phys. Rev. C* **53**, R2602 (1996).
- [12] M. Agostini, G. Benato, and J. A. Detwiler, *Phys. Rev. D* **96**, 053001 (2017).
- [13] M. Ettengruber, M. Agostini, A. Caldwell, P. Eller, and O. Schulz, *Phys. Rev. D* **106**, 073004 (2022).
- [14] A. Barabash, *Universe* **6**, 159 (2020).
- [15] F. Šimkovic, P. Domin, and S. V. Semenov, *J. Phys. G* **27**, 2233 (2001).
- [16] F. Šimkovic, R. Dvornický, D. Stefánik, and A. Faessler, *Phys. Rev. C* **97**, 034315 (2018).
- [17] O. Nițescu, R. Dvornický, S. Stoica, and F. Šimkovic, *Universe* **7**, 147 (2021).
- [18] A. Gando *et al.* (KamLAND-Zen Collaboration), *Phys. Rev. Lett.* **122**, 192501 (2019).
- [19] F. Šimkovic, V. Rodin, A. Faessler, and P. Vogel, *Phys. Rev. C* **87**, 045501 (2013).
- [20] F. Šimkovic, L. Paceaescu, and A. Faessler, *Nucl. Phys.* **A733**, 321 (2004).
- [21] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.131.162501> for a brief description of the proton-neutron quasiparticle random-phase approximation calculations of nuclear matrix elements and the posterior distributions from the fit.
- [22] O. Azzolini *et al.* (CUPID-0 Collaboration), *Eur. Phys. J. C* **79**, 583 (2019).
- [23] D. Q. Adams *et al.* (CUORE Collaboration), *Phys. Rev. Lett.* **126**, 171801 (2021).
- [24] E. Armengaud *et al.*, *Eur. Phys. J. C* **80**, 674 (2020).
- [25] M. Agostini *et al.* (GERDA Collaboration), *J. High Energy Phys.* **03** (2020) 139.

- [26] S. Pirro, J. W. Beeman, S. Capelli, M. Pavan, E. Previtali, and P. Gorla, *Phys. At. Nucl.* **69**, 2109 (2006).
- [27] D. Poda, *Physics* **3**, 473 (2021).
- [28] E. Armengaud *et al.*, *Eur. Phys. J. C* **77**, 785 (2017).
- [29] L. Cardani *et al.*, *J. Instrum.* **8**, P10002 (2013).
- [30] O. Azzolini *et al.* (CUPID-0 Collaboration), *Eur. Phys. J. C* **78**, 428 (2018).
- [31] J. W. Beeman *et al.* (LUCIFER Collaboration), *Eur. Phys. J. C* **75**, 591 (2015).
- [32] W. R. Armstrong *et al.* (CUPID Collaboration), [arXiv:1907.09376](https://arxiv.org/abs/1907.09376).
- [33] E. Armengaud *et al.*, *Eur. Phys. J. C* **80**, 44 (2020).
- [34] E. E. Haller, N. P. Palaio, M. Rodder, W. L. Hansen, and E. Kreysa, in *Neutron Transmutation Doping of Semiconductor Materials*, edited by R. D. Larrabee (Springer US, Boston, MA, 1984), pp. 21–36.
- [35] E. Armengaud *et al.* (EDELWEISS Collaboration), *J. Instrum.* **12**, P08010 (2017).
- [36] C. Augier *et al.*, *Eur. Phys. J. C* **82**, 1033 (2022).
- [37] C. Augier *et al.* (CUPID-Mo Collaboration), *Phys. Rev. C* **107**, 025503 (2023).
- [38] C. Augier *et al.* (CUPID-Mo Collaboration), *Eur. Phys. J. C* **83**, 675 (2023).
- [39] E. Gatti and P. F. Manfredi, *Riv. Nuovo Cimento* **9N1**, 1 (1986).
- [40] R. Huang *et al.* (CUPID-Mo Collaboration), *J. Instrum.* **16**, P03032 (2021).
- [41] B. Schmidt *et al.* (EDELWEISS Collaboration), *Astropart. Phys.* **44**, 28 (2013).
- [42] J. Allison *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **835**, 186 (2016).
- [43] M. Plummer, in *Proceedings of the 3rd International Workshop on Distributed Statistical Computing (DSC 2003)* (DSC proceedings, Vienna, 2003), p. 124.
- [44] M. Plummer, JAGS version 3.3.0 user manual (2012).
- [45] R. Arnold *et al.* (NEMO-3 Collaboration), *Eur. Phys. J. C* **79**, 440 (2019).
- [46] L. Coraggio, N. Itaco, G. De Gregorio, A. Gargano, R. Mancino, and F. Nowacki, *Phys. Rev. C* **105**, 034312 (2022).
- [47] L. Coraggio and G. De Gregorio (private communication).
- [48] K. Nomura, *Phys. Rev. C* **105**, 044301 (2022).