

JYX



This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Mohanta, Kaushik

Title: Bourgain-Brezis-Mironescu formula for $W_{s,p,q}$ -spaces in arbitrary domains

Year: 2024

Version: Published version

Copyright: © The Author(s) 2024

Rights: CC BY 4.0

Rights url: <https://creativecommons.org/licenses/by/4.0/>

Please cite the original version:

Mohanta, K. (2024). Bourgain-Brezis-Mironescu formula for $W_{s,p,q}$ -spaces in arbitrary domains. *Calculus of Variations and Partial Differential Equations*, 63(2), Article 31.
<https://doi.org/10.1007/s00526-023-02637-w>



Bourgain-Brezis-Mironescu formula for $W_q^{s,p}$ -spaces in arbitrary domains

Kaushik Mohanta¹

Received: 19 August 2023 / Accepted: 3 December 2023
© The Author(s) 2024

Abstract

Under certain restrictions on s, p, q , the Triebel-Lizorkin spaces can be viewed as generalised fractional Sobolev spaces $W_q^{s,p}$. In this article, we show that the Bourgain-Brezis-Mironescu formula holds for $W_q^{s,p}$ -seminorms in arbitrary domain. This addresses an open question raised by Brazke-Schikorra-Yung (Calc Var Partial Differ Equ 62(2):41–33, (2023)).

Mathematics Subject Classification 46E35 · 42B35

1 Introduction

Sobolev spaces arise naturally in the study of partial differential equations. They are defined in terms of weak derivatives. For an open set $\Omega \subseteq \mathbb{R}^N$, and $1 \leq p < \infty$, the Sobolev space $W^{1,p}(\Omega)$ is defined to be $\{f \in L^p(\Omega) \mid [f]_{W^{1,p}(\Omega)} < \infty\}$, where

$$[f]_{W^{1,p}(\Omega)}^p := \int_{\Omega} |\nabla f(x)|^p dx.$$

Some closely related spaces are Triebel-Lizorkin spaces $F_{p,q}^s(\Omega)$, which are defined to be $\{f|_{\Omega} \mid f \in F_{p,q}^s(\mathbb{R}^N)\}$, and are equipped with the norm $[\cdot]_{F_{p,q}^s(\mathbb{R}^N)}$. We shall not define the norm $[\cdot]_{F_{p,q}^s(\mathbb{R}^N)}$ as it will not be necessary for the present article. Instead, we refer the reader, for definition and classical results regarding Triebel-Lizorkin spaces, to [59], or more modern references like [36, 56]. For $1 \leq p, q < \infty$, $\max\{0, \frac{N(q-p)}{pq}\} < s < 1$, we have the characterisation (see Theorem 1.2 of [55])

$$F_{p,q}^s(\mathbb{R}^N) := \left\{ f \in L^{\max\{p,q\}}(\mathbb{R}^N) \mid \|f\|_{L^p(\mathbb{R}^N)} + [f]_{W_q^{s,p}(\mathbb{R}^N)} < \infty \right\},$$

Communicated by A. Mondino.

✉ Kaushik Mohanta
kaushik.k.mohanta@jyu.fi

¹ Department of Mathematics and Statistics, University of Jyväskylä, Jyväskylä, Finland

where

$$[f]_{W_q^{s,p}(\Omega)} := \left(\int_{\Omega} \left(\int_{\Omega} \frac{|f(x) - f(y)|^q}{|x - y|^{N+sq}} dy \right)^{\frac{p}{q}} dx \right)^{\frac{1}{p}}. \tag{1}$$

In the special case of $p = q$, these spaces are related to the so-called fractional Sobolev spaces $W^{s,p}(\Omega)$, defined by

$$W^{s,p}(\Omega) := \left\{ f \in L^p(\Omega) \mid \|f\|_{L^p(\Omega)} + [f]_{W^{s,p}(\Omega)} < \infty \right\},$$

where $[f]_{W^{s,p}(\Omega)} := [f]_{W_p^{s,p}(\Omega)}$. Of course, when $\Omega = \mathbb{R}^N$, or when Ω is fractional extension domain (see [25]), we have $W^{s,p}(\Omega) = F_{p,p}^s(\Omega)$.

Bourgain-Brezis-Mironescu [8] showed that for any smooth and bounded domain Ω , $1 \leq p < \infty$, and any $f \in W^{1,p}(\Omega)$,

$$\lim_{s \rightarrow 1^-} (1 - s)[f]_{W^{s,p}(\Omega)}^p = K \|\nabla f\|_{L^p(\Omega)}^p.$$

Conversely, for any $f \in L^p(\Omega)$, if we have

$$\lim_{s \rightarrow 1^-} (1 - s)[f]_{W^{s,p}(\Omega)}^p < \infty,$$

then $f \in W^{1,p}(\Omega)$ if $p > 1$ and $f \in BV(\Omega)$ if $p = 1$. Later Dávila [23] extended this result and proved that for any $f \in BV(\Omega)$, $\lim_{s \rightarrow 1^-} (1 - s)[f]_{W^{s,1}(\Omega)}^p = K |\nabla f|(\Omega)$. This is commonly known as the Bourgain-Brezis-Mironescu formula (BBM formula for short). The subject was further developed in [9, 10, 18, 47, 48, 53]. The BBM formula has been generalised to Orlicz and generalised Orlicz setup in [1–3, 28, 31, 61], to variable exponent setup in [30, 39], to magnetic Sobolev spaces [29, 49, 50, 52, 57], to anisotropic setup [37, 45, 51], to Riemannian manifolds in [42], to metric spaces in [24, 35], to Banach function spaces in [22, 62]. Similar studies are possible in the context of Besov spaces also [41, 60]. More recent developments on this topic can be found in [14–17]. For further reading, we refer the reader to [4, 6, 7, 11, 19, 21, 26, 32–34, 38, 46, 54].

Our first interest lies in works regarding the domain. That is, how far the smooth-bounded condition, on the domain, can be relaxed. In this direction, we mention three works. The first is due to Lioni-Spector [43, 44]. They showed that for an arbitrary domain Ω , and $f \in L^p(\Omega)$,

$$\lim_{\lambda \rightarrow 0^+} \lim_{s \rightarrow 1^-} (1 - s) \int_{\Omega_\lambda} \int_{\Omega_\lambda} \frac{|f(x) - f(y)|^p}{|x - y|^{N+sp}} dy dx = K [f]_{W^{1,p}(\Omega)}^p.$$

where

$$\Omega_\lambda := \{x \in \Omega \mid \text{dist}(x, \partial\Omega) > \lambda\} \cap B(0, \lambda^{-1}). \tag{2}$$

The second result is due to the author with Bal-Roy [5], where it has been shown that the BBM-formula [8], holds if we take Ω to be a $W^{1,p}$ -extension domain. The third result is due to Drelichman-Duran [27]. They showed that for $1 < p < \infty$, and an arbitrary bounded domain Ω , and any $\tau \in (0, 1)$, we have

$$\lim_{s \rightarrow 1^-} (1 - s) \int_{\Omega} \int_{B(x, \tau \text{dist}(x, \partial\Omega))} \frac{|f(x) - f(y)|^p}{|x - y|^{N+sp}} dy dx = K [f]_{W^{1,p}(\Omega)}^p.$$

The second direction of work regarding the BBM-formula, that we are interested in, is its extension for Triebel-Lizorkin spaces. The first work in this direction was done by

Brazke-Schikorra-Yung [12]. They explained via examining thoroughly various constants of embeddings that although $F_{p,p}^s = W^{s,p}$, when $s \in (0, 1)$ and $F_{p,2}^s = W^{1,p}$ it makes sense for the scaled $W^{s,p}$ seminorm to converge to $W^{1,p}$ -seminorm, even when $p \neq 2$. They posed the open problem (see [12, Question 1.12]) about the asymptotic constant in the identification of the $\|\cdot\|_{W_q^{s,p}} \approx \|\cdot\|_{F_{p,q}^s}$.

The current article addresses this question by showing the asymptotic behaviour (as $s \rightarrow 1-$) of $W_q^{s,p}$ -seminorms. Similar studies has been done, when $1 < q < p < \infty$, in [22] for \mathbb{R}^N , and in [62, Theorem 6.1] for a special class of bounded extension domains (called (ε, ∞) -domains).

We concentrate our focus on the following seminorm (for some $\tau \in (0, 1)$)

$$[f]_{\tilde{W}_q^{s,p}(\Omega)} := \left(\int_{\Omega} \left(\int_{B(x,\tau \text{dist}(x,\partial\Omega))} \frac{|f(x) - f(y)|^q}{|x - y|^{N+sq}} dy \right)^{\frac{p}{q}} dx \right)^{\frac{1}{p}}, \tag{3}$$

as one can then extend the above question for arbitrary bounded domains, motivated by [27]. We go one step further and show that the boundedness of the domain is not necessary. Our main results are the following:

Theorem 1 *Let $\Omega \subset \mathbb{R}^N$ be an open set $\tau \in (0, 1)$. Assume one of the following conditions*

- (1) $1 \leq q \leq p < \infty$,
- (2) $1 < p < q < \infty$ with $p \leq N$ and $q < \frac{Np}{N-p}$,
- (3) $N < p < q < \infty$.

Then there is a constant $K = K(N, p, q) > 0$ such that for any $f \in W^{1,p}(\Omega)$, we have,

$$\lim_{s \rightarrow 1-} (1 - s)^{\frac{p}{q}} \int_{\Omega} \left(\int_{B(x,\tau \text{dist}(x,\partial\Omega))} \frac{|f(x) - f(y)|^q}{|x - y|^{N+sq}} dy \right)^{\frac{p}{q}} dx = K \int_{\Omega} |\nabla f(x)|^p dx. \tag{4}$$

Theorem 2 *Let $\Omega \subset \mathbb{R}^N$ be an open set, $\tau \in (0, 1)$ and $1 \leq p, q < \infty$. If $f \in L^p(\Omega) \cap L^q(\Omega)$ is such that*

$$L_{p,q}^*(f) := \lim_{s \rightarrow 1-} \int_{\Omega} \left((1 - s) \int_{B(x,\tau \text{dist}(x,\partial\Omega))} \frac{|f(x) - f(y)|^q}{|x - y|^{N+sq}} dy \right)^{\frac{p}{q}} dx < \infty,$$

then $f \in W^{1,p}(\Omega)$ when $p > 1$, and $f \in BV(\Omega)$ when $p = 1$.

Remark 3 Before proceeding further, let us discuss some difficulties that arise here, and strategies for overcoming them. The proof of the main results roughly follow the outline of [8]. However, there are certain obstacles to that path. The first obstacle arises when we want to apply dominated convergence theorem to interchange limit and integral. Similar difficulty was faced and overcome in [27], but we had to take a different route (see lemma 9) for this purpose. The introduction of the second exponent q forces us to deviate from the usual route again; The case $q \leq p$ is rather easy to handle, for the case $p < q$, a careful use of Sobolev embedding is needed. To take into account the case where the domain Ω is unbounded, we need to restrict the seminorm further and define some new fractional Sobolev spaces (see eq. 5) and prove a version of the main result theorem 1 in that context (see theorem 14), and then finally derive the proof of the main results from there.

The article is organised as follows: In section 2, we list some preliminary results, already known in literature, which shall be useful for the proof of our main results. In section 3, we

introduce a variant of fractional Sobolev spaces and prove some relevant embedding results. In section 4 we prove the main results in the context of these new spaces (see theorems 14 and 15). Finally in section 5, we prove theorems 1 and 2.

2 Preliminary results

For the sake of completeness, we first state the well-known Sobolev inequality (also known as (q, p) -Poincaré inequality):

Lemma 4 *Let $1 \leq p, q \leq \infty, \tau \in (0, 1)$, and one of the following hold*

- (1) $p < N$, and $q \leq \frac{Np}{N-p}$,
- (2) $p = N$, and $q < \infty$
- (3) $p > N$.

Then there is a constants $C = C(p, q, N) > 0$ such that the following holds for any $f \in W^{1,p}(B(0, \frac{1}{\tau}))$:

$$\begin{aligned} \frac{1}{t^N} \int_{B(0,t)} |f(y)|^q dy &\leq C(N, p, q)t^q \left(\frac{1}{t^N} \int_{B(0,t)} |\nabla f(y)|^p dy \right)^{\frac{q}{p}} \\ &\quad + C(N, p, q) \left(\frac{1}{t^N} \int_{B(0,t)} |f(y)|^p dy \right)^{\frac{q}{p}}. \end{aligned}$$

The following lemma was established in [20] for $1 < p < \infty$; the $p \geq 1$ case can be found in [58, Chapter-VI, Theorems 5 and 5’].

Lemma 5 *Let $\Omega \subseteq \mathbb{R}^N$ be an open set with Lipschitz boundary and $1 \leq p < \infty$. Then for any $f \in W^{1,p}(\Omega)$ there is some $\tilde{f} \in W^{1,p}(\mathbb{R}^N)$ such that $\tilde{f}|_{\Omega} = f$ and for some constant $C = C(N, \Omega, p)$,*

$$\|\tilde{f}\|_{W^{1,p}(\mathbb{R}^N)} \leq C \|f\|_{W^{1,p}(\Omega)}.$$

The following result can be found in Proposition 9.3 and Remark 6 of [13].

Lemma 6 *Let $\Omega \subseteq \mathbb{R}^N$ be open, $1 \leq p < \infty, \frac{1}{p} + \frac{1}{p'} = 1$, and $f \in L^p(\Omega)$. Assume that there is constant $C > 0$ such that for any $\varphi \in C_c^\infty(\Omega)$*

$$\left| \int_{\Omega} f(x) \frac{\partial \varphi(x)}{\partial x_i} dx \right| \leq C \|\varphi\|_{L^{p'}(\Omega)} \quad \text{for } i = 1, 2, \dots, N.$$

Then $f \in W^{1,p}(\Omega)$ when $1 < p$, and $f \in BV(\Omega)$ when $p = 1$.

Next we list a special case of Proposition 2/(ii) of [59, Chapter 2.3.3], combined with the fact that $W^{1,p}(\mathbb{R}^N) = F_{p,2}^1(\mathbb{R}^N)$.

Lemma 7 *Let $1 \leq p, q < \infty, s \in (0, 1)$. Then*

$$W^{1,p}(\mathbb{R}^N) \subseteq F_{p,q}^s(\mathbb{R}^N) = W_q^{s,p}(\mathbb{R}^N).$$

The following result is taken from Lemma 8 of [5].

Lemma 8 *Let $\Omega \subseteq \mathbb{R}^N$ be open and $\lambda > 0$ be sufficiently small. Then there is a bounded open set Ω_λ^* with smooth boundary such that $\Omega_\lambda \subseteq \Omega_\lambda^* \subseteq \Omega$, where Ω_λ is as in (2).*

The next result can be found in Theorem 2.1 of [40]. It will play a crucial in this article.

Lemma 9 *Let $\Omega \subseteq \mathbb{R}^N$, $\{\Omega_i\}_{i \in \mathbb{N}}$ be such that $\Omega = \cup_i \Omega_i$, $F_n, F \in L^1(\Omega)$ for $n \in \mathbb{N}$ be such that for a.e. $x \in \Omega$, $F_n(x) \rightarrow F(x)$ as $n \rightarrow \infty$. Assume that*

(1)

$$\limsup_{n \rightarrow \infty} \sup_{x \in \Omega_i} |F_n(x) - F(x)| < \infty \text{ for all } i \in \mathbb{N},$$

(2)

$$\liminf_{i \rightarrow \infty} \limsup_{n \rightarrow \infty} \int_{\Omega \setminus \Omega_i} F_n(x) dx = 0.$$

Then

$$\lim_{n \rightarrow \infty} \int_{\Omega} F_n(x) dx = \int_{\Omega} F(x).$$

3 Fractional Sobolev space with restricted internal distance

Fix $R > 0$ and $\tau \in (0, 1)$ once and for all. Denote $\delta_{x,R,\tau} = \min\{R, \tau \text{dist}(x, \partial\Omega)\}$. We shall often drop the R and τ in the above notation and write δ_x to denote $\delta_{x,R,\tau}$.

Remark 10 If the function $x \mapsto \text{dist}(x, \partial\Omega)$ is bounded in Ω , we can choose $R > 0$ large enough, so that $\delta_x = \tau \text{dist}(x, \partial\Omega)$. Then the particular case $p = q$ of theorems 14 and 15 are similar to the results proved in [27], but here Ω need not be a bounded domain; for example, it can be a cylindrical domain or any open subset of $\mathbb{R}^N \setminus \mathbb{Z}^N$.

Define, for any open set $\Omega \subseteq \mathbb{R}^N$, $1 \leq p, q < \infty$, $0 < s < 1$, $\hat{W}_q^{s,p}(\Omega) := \{f \in L^p(\Omega) \mid [f]_{\hat{W}_q^{s,p}(\Omega)} < \infty\}$ where

$$[f]_{\hat{W}_q^{s,p}(\Omega)}^p := \int_{x \in \Omega} \left(\int_{y \in B(x, \delta_x)} \frac{|f(x) - f(y)|^q}{|x - y|^{N+sq}} dy \right)^{\frac{p}{q}} dx. \tag{5}$$

We shall need some embedding results for these new fractional Sobolev spaces for our purpose. As expected, the case $q \leq p$ and $p < q$ are treated separately.

Lemma 11 *Let $\Omega \subseteq \mathbb{R}^N$ be open and $1 \leq q \leq p < \infty$. Assume either $D = \tilde{D} = \Omega$ or, for some $\frac{1}{2R} > \alpha > 0$, $D = \{x \in \Omega \mid \text{dist}(x, \partial\Omega) < \alpha\}$ with $\tilde{D} = \{x \in \Omega \mid \text{dist}(x, \partial\Omega) < 2\alpha \text{ or } |x| > \frac{1}{2\alpha}\}$. Then there is a constant $C = C(p, q, R, \Omega, N)$ such that for any f in $W^{1,p}(\Omega)$,*

$$(1 - s)^{\frac{p}{q}} \int_{x \in D} \left(\int_{y \in B(x, \delta_x)} \frac{|f(x) - f(y)|^q}{|x - y|^{N+sq}} dy \right)^{\frac{p}{q}} dx \leq C [f]_{W^{1,p}(\tilde{D})}^p. \tag{6}$$

Proof We have

$$\begin{aligned} & (1 - s)^{\frac{p}{q}} \int_{x \in D} \left(\int_{y \in B(x, \delta_x)} \frac{|f(x) - f(y)|^q}{|x - y|^{N+sq}} dy \right)^{\frac{p}{q}} dx \\ &= (1 - s)^{\frac{p}{q}} \int_{x \in D} \left(\int_{h \in B(0, \delta_x)} \frac{|f(x+h) - f(x)|^q}{|h|^{N+sq}} dh \right)^{\frac{p}{q}} dx \end{aligned}$$

$$\begin{aligned} &= (1-s)^{\frac{p}{q}} \int_{x \in D} \left(\int_{h \in B(0, \delta_x)} \frac{|f(x+h) - f(x)|^q}{|h|^p} \frac{dh}{|h|^{N+sq-q}} \right)^{\frac{p}{q}} dx \\ &\leq (1-s)^{\frac{p}{q}} \int_{x \in D} \left(\int_{h \in B(0, \delta_x)} \int_0^1 |\nabla f(x+th)|^q dt \frac{dh}{|h|^{N+sq-q}} \right)^{\frac{p}{q}} dx. \end{aligned}$$

The last inequality follows from the absolute continuity on lines of the $W^{1,p}$ -functions. We now have, after a change of variable $y = x + th$, (using that $B(x, t\delta_x) \subset B(x, \delta_x)$)

$$\begin{aligned} &(1-s)^{\frac{p}{q}} \int_{x \in D} \left(\int_{y \in B(x, \delta_x)} \frac{|f(x) - f(y)|^q}{|x-y|^{N+sq}} dy \right)^{\frac{p}{q}} dx \\ &\leq (1-s)^{\frac{p}{q}} \int_{x \in D} \left(\int_{y \in B(x, \delta_x)} \int_0^1 |\nabla f(y)|^q t^{sq-q} dt \frac{dy}{|x-y|^{N+sq-q}} \right)^{\frac{p}{q}} dx \\ &= \frac{(1-s)^{\frac{p}{q}}}{(sq-q+1)^{\frac{p}{q}}} \int_{x \in D} \left(\int_{y \in B(x, \delta_x)} |\nabla f(y)|^q \frac{dy}{|x-y|^{N+sq-q}} \right)^{\frac{p}{q}} dx. \end{aligned}$$

Note that in the above inequality, ∇f is required to be defined only inside \tilde{D} . So, we shall take a 0-extension of ∇f outside \tilde{D} . Since we have $\frac{p}{q} \geq 1$, we can use Young’s convolution inequality to get

$$\begin{aligned} &(1-s)^{\frac{p}{q}} \int_{x \in D} \left(\int_{y \in B(x, \delta_x)} \frac{|f(x) - f(y)|^q}{|x-y|^{N+sq}} dy \right)^{\frac{p}{q}} dx \\ &\leq \frac{(1-s)^{\frac{p}{q}}}{(sq-q+1)^{\frac{p}{q}}} \int_{x \in D} \left(\int_{y \in B(x, R)} |\nabla f(y)|^q \frac{dy}{|x-y|^{N+sq-q}} \right)^{\frac{p}{q}} dx \\ &\leq \frac{(1-s)^{\frac{p}{q}}}{(sq-q+1)^{\frac{p}{q}}} \int_{x \in \tilde{D}} |\nabla f(x)|^p dx \left(\int_{x \in B(0, R)} \frac{dx}{|x|^{N+sq-q}} \right)^{\frac{p}{q}} \\ &= \frac{R^{p-sp}}{q^{\frac{p}{q}}} \int_{x \in \tilde{D}} |\nabla f(x)|^p dx. \end{aligned}$$

□

Lemma 12 Let $\Omega \subseteq \mathbb{R}^N$ be open, $1 < p < q < \infty$, and one of the following hold

- (1) $p < N$, and $q < \frac{Np}{N-p}$,
- (2) $N \leq p$.

Assume either $D = \tilde{D} = \Omega$ or, for some $\frac{1}{2R} > \alpha > 0$, $D = \{x \in \Omega \mid \text{dist}(x, \partial\Omega) < \alpha\}$ and $\tilde{D} = \{x \in \Omega \mid \text{dist}(x, \partial\Omega) < 2\alpha \text{ or } |x| > \frac{1}{2\alpha}\}$. Then there is a constant $C = C(N, R, p, q) > 0$ such that for any $f \in W^{1,p}(\Omega)$, eq. (6) holds.

Proof Note that

$$\int_{t=|h|}^{\delta_x} \frac{dt}{t^{N+sq+1}} = \frac{1}{N+sq} \left(\frac{1}{|h|^{N+sq}} - \frac{1}{\delta_x^{N+sq}} \right),$$

which gives

$$\frac{1}{|h|^{N+sq}} = (N+sq) \int_{t=|h|}^{\delta_x} \frac{dt}{t^{N+sq+1}} + \delta_x^{-N-sq}.$$

So, we can write

$$\begin{aligned}
 C(p, q) &\int_{x \in D} \left((1-s) \int_{h \in B(0, \delta_x)} \frac{|f(x+h) - f(x)|^q}{|h|^{N+sq}} dh \right)^{\frac{p}{q}} dx \\
 &\leq \int_{x \in D} \left((N+sq)(1-s) \int_{h \in B(0, \delta_x)} \int_{t=|h|}^{\delta_x} \frac{|f(x+h) - f(x)|^q}{t^{N+sq+1}} dt dh \right)^{\frac{p}{q}} dx \\
 &\quad + \int_{x \in D} \left((1-s) \int_{h \in B(0, \delta_x)} \frac{|f(x+h) - f(x)|^q}{\delta_x^{N+sq}} dh \right)^{\frac{p}{q}} dx \\
 &= I_1 + I_2.
 \end{aligned}
 \tag{7}$$

According to either $p < N$ or $p \geq N$, fix $\beta \in (0, 1)$, depending on p, q, N , such that $(q, \beta p)$ -type Poincaré inequality (lemma 4) is satisfied. We use this to estimate I_1 below. First, we change the order of integration between t and h , then apply lemma 4.

$$\begin{aligned}
 I_1 &= \int_{x \in D} \left((N+sq)(1-s) \int_{t=0}^{\delta_x} \frac{1}{t^{N+sq+1}} \int_{h \in B(0,t)} |f(x+h) - f(x)|^q dh dt \right)^{\frac{p}{q}} dx \\
 &\leq \int_{x \in D} \left((N+sq)(1-s) \int_{t=0}^{\delta_x} \frac{1}{t^{sq+1-q}} \left(\frac{1}{t^N} \int_{h \in B(x,t)} |\nabla f(h)|^{\beta p} dh \right)^{\frac{q}{\beta p}} dt \right)^{\frac{p}{q}} dx \\
 &\quad + \int_{x \in D} \left((N+sq)(1-s) \int_{t=0}^{\delta_x} \frac{1}{t^{sq+1}} \left(\frac{1}{t^N} \int_{h \in B(0,t)} |f(x+h) - f(x)|^{\beta p} dh \right)^{\frac{q}{\beta p}} dt \right)^{\frac{p}{q}} dx \\
 &= I_{1,1} + I_{1,2}.
 \end{aligned}
 \tag{8}$$

As in the proof of the previous lemma, we shall take a 0-extension of ∇f outside \tilde{D} . Now using Hardy-Littlewood maximal inequality, we get

$$\begin{aligned}
 I_{1,1} &\leq \int_{x \in D} (M|\nabla f(x)|^{\beta p})^{\frac{1}{\beta}} \left((N+sq)(1-s) \int_{t=0}^{\delta_x} \frac{1}{t^{sq+1-q}} dt \right)^{\frac{p}{q}} dx \\
 &= C(N, p, q, R) \int_{x \in \mathbb{R}^N} (M|\nabla f(x)|^{\beta p})^{\frac{1}{\beta}} dx \\
 &\leq C(N, p, q, R) \int_{x \in \tilde{D}} |\nabla f(x)|^p dx.
 \end{aligned}
 \tag{9}$$

Again,

$$\begin{aligned}
 I_{1,2} &\leq \int_{x \in D} \left((N+sq)(1-s) \int_{t=0}^{\delta_x} \frac{1}{t^{sq+1-q}} \left(\frac{1}{t^N} \int_{h \in B(0,t)} \frac{|f(x+h) - f(x)|^{\beta p}}{|h|^{\beta p}} dh \right)^{\frac{q}{\beta p}} dt \right)^{\frac{p}{q}} dx \\
 &\leq \int_{x \in D} \left((N+sq)(1-s) \int_{t=0}^{\delta_x} \frac{1}{t^{sq+1-q}} \left(\int_{r=0}^1 \frac{1}{t^N} \int_{h \in B(0,t)} |\nabla f(x+rh)|^{\beta p} dh dr \right)^{\frac{q}{\beta p}} dt \right)^{\frac{p}{q}} dx \\
 &= \int_{x \in D} \left((N+sq)(1-s) \int_{t=0}^{\delta_x} \frac{1}{t^{sq+1-q}} \left(\int_{r=0}^1 \frac{1}{(rt)^N} \int_{h \in B(x,rt)} |\nabla f(h)|^{\beta p} dh dr \right)^{\frac{q}{\beta p}} dt \right)^{\frac{p}{q}} dx \\
 &\leq \int_{x \in \mathbb{R}^N} (M|\nabla f(x)|^{\beta p})^{\frac{1}{\beta}} \left((N+sq)(1-s) \int_{t=0}^R \frac{1}{t^{sq+1-q}} \left(\int_{r=0}^1 dr \right)^{\frac{q}{\beta p}} dt \right)^{\frac{p}{q}} dx \\
 &\leq C(p, q, N, R) \int_{x \in \tilde{D}} |\nabla f(x)|^p dx.
 \end{aligned}
 \tag{10}$$

Combining eqs. (9) and (10), we get

$$I_1 \leq C(p, q, N, R) \|\nabla f\|_{L^p(\tilde{D})}^p. \tag{11}$$

Again, we can estimate I_2 , in similar way as above with δ_x in place of t . We have a better estimate this time. Also, we can apply (q, p) -Poincaré inequality this time.

$$\begin{aligned} I_2 &= \int_{x \in D} \left((1-s) \int_{h \in B(0, \delta_x)} \frac{|f(x+h) - f(x)|^q}{\delta_x^{N+sq}} dh \right)^{\frac{p}{q}} dx \\ &\leq \int_{x \in D} \left(\frac{(1-s)}{\delta_x^{sq}} \left(\delta_x^{p-N} \int_{h \in B(0, \delta_x)} |\nabla f(x+h)|^p dh + \delta_x^{-N} \int_{h \in B(0, \delta_x)} |f(x+h) - f(x)|^p dh \right)^{\frac{q}{p}} \right)^{\frac{p}{q}} dx \\ &\leq \int_{x \in D} \frac{(1-s)^{\frac{p}{q}}}{\delta_x^{sp-p}} \left(\left(\delta_x^{-N} \int_{h \in B(0, \delta_x)} |\nabla f(x+h)|^p dh + \delta_x^{-N} \int_{h \in B(0, \delta_x)} \frac{|f(x+h) - f(x)|^p}{|h|^p} dh \right)^{\frac{q}{p}} \right)^{\frac{p}{q}} dx \\ &\leq \int_{x \in D} \frac{(1-s)^{\frac{p}{q}}}{\delta_x^{sp-p}} \left(\delta_x^{-N} \int_{h \in B(0, \delta_x)} |\nabla f(x+h)|^p dh + \int_{r=0}^1 \frac{1}{(r\delta_x)^N} \int_{h \in B(0, r\delta_x)} |\nabla f(x+h)|^p dh dr \right) dx \\ &\leq (1-s)^{\frac{p}{q}} \int_{x \in D} \delta_x^{p-sp} |\nabla f(x)|^p dx. \end{aligned} \tag{12}$$

Combing eqs. (7), (11) and (12),

$$\int_{x \in D} \left((1-s) \int_{h \in B(0, \delta_x)} \frac{|f(x+h) - f(x)|^q}{|h|^{N+sq}} dh \right)^{\frac{p}{q}} dx \leq C(N, p, q, R) \|\nabla f\|_{L^p(\tilde{D})}^p.$$

This proves the lemma. □

4 BBM formula for $\widehat{W}_q^{s,p}$ -seminorms

First, we state the following result whose proof can be found in the proof of Theorem of [8] as the quantity δ_x is bounded by R .

Lemma 13 *Let $\Omega \subset \mathbb{R}^N$ be any open set, $1 \leq q < \infty$, $0 < s < 1$. Then for any $f \in C^2(\Omega)$, we have for all $x \in \Omega$,*

$$\lim_{s \rightarrow 1^-} (1-s) \int_{B(x, \delta_x)} \frac{|f(x) - f(y)|^q}{|x-y|^{N+sq}} dy = K |\nabla f(x)|^q. \tag{13}$$

We now prove the following BBM-type results which are closely related with theorems 1 and 2.

Theorem 14 *Let $\Omega \subset \mathbb{R}^N$ be any open set. Assume one of the following conditions*

- (1) $1 \leq q \leq p < \infty$,
- (2) $1 < p < q < \infty$ with $p \leq N$ and $q < \frac{Np}{N-p}$,
- (3) $N < p < q < \infty$.

Then there is a constant $K = K(N, p, q) > 0$ such that for any $f \in W^{1,p}(\Omega)$, we have for all $x \in \Omega$,

$$\lim_{s \rightarrow 1^-} (1-s)^{\frac{p}{q}} \int_{\Omega} \left(\int_{B(x, \delta_x)} \frac{|f(x) - f(y)|^q}{|x-y|^{N+sq}} dy \right)^{\frac{p}{q}} dx = K \int_{\Omega} |\nabla f(x)|^p dx. \tag{14}$$

Proof of Theorem 14 Step-1: We show that it is enough to prove eq. (14) for $f \in W^{1,p}(\Omega) \cap C^2(\Omega)$.

Let $f \in W^{1,p}(\Omega)$ and $\varepsilon > 0$ be fixed. Since $C^2(\Omega) \cap W^{1,p}(\Omega)$ is dense in $W^{1,p}(\Omega)$, there exists $g \in C^2(\Omega) \cap W^{1,p}(\Omega)$ such that

$$[f - g]_{W^{1,p}(\Omega)} < \varepsilon. \tag{15}$$

Since we have assumed that eq. (14) holds for functions in $C^2(\Omega) \cap W^{1,p}(\Omega)$, for $s > \frac{1}{2}$, we have

$$\left| (1-s)^{\frac{1}{q}} [g]_{s,p,q,\Omega,R} - K^{\frac{1}{p}} [g]_{W^{1,p}(\Omega)} \right| < \varepsilon. \tag{16}$$

Using triangle inequality, and then eq. (16) followed by eq. (15) and either lemma 11 or lemma 12, we get

$$\begin{aligned} & \left| (1-s)^{\frac{1}{q}} [f]_{s,p,q,\Omega,R} - K^{\frac{1}{p}} [f]_{W^{1,p}(\Omega)} \right| \leq (1-s)^{\frac{1}{q}} |[f]_{s,p,q,\Omega,R} - [g]_{s,p,q,\Omega,R}| \\ & + \left| (1-s)^{\frac{1}{q}} [g]_{s,p,\Omega,R} - K^{\frac{1}{q}} [g]_{W^{1,p}(\Omega)} \right| + K^{\frac{1}{q}} |[g]_{W^{1,p}(\Omega)} - [f]_{W^{1,p}(\Omega)}| \\ & \leq (1-s)^{\frac{1}{q}} [f - g]_{s,p,q,\Omega,R} + \varepsilon + K^{\frac{1}{p}} [f - g]_{W^{1,p}(\Omega)} \\ & \leq [f - g]_{W^{1,p}(\Omega)} + \varepsilon + K^{\frac{1}{p}} [f - g]_{W^{1,p}(\Omega)} \\ & \leq C(K, q)\varepsilon. \end{aligned}$$

The proof of step-1 follows.

Step-2:

In view of the previous step, it is now enough to assume that $f \in C^2(\Omega) \cap W^{1,p}(\Omega)$ and prove eq. (14). Let us take an arbitrary sequence $s_n \in (0, 1)$ such that $s_n \rightarrow 1$ as $n \rightarrow \infty$. Set

$$F_n(x) := \left((1-s_n) \int_{B(0, \delta_x)} \frac{|f(x+h) - f(x)|^p}{|h|^{N+s_n p}} dh \right)^{\frac{p}{q}},$$

and

$$F(x) := K |\nabla f(x)|^p.$$

Also note that, lemma 13 implies that $F_n \rightarrow F$ pointwise a.e in Ω . To complete the proof, it is enough to show that

$$\lim_{n \rightarrow \infty} \int_{\Omega} F_n(x) dx = \int_{\Omega} F(x) dx.$$

We shall apply lemma 9 on F_n to show that the interchange of limit and integral is valid.

For any $i \in \mathbb{N}$, consider the sets $\Omega_i := \{x \in \Omega \mid \text{dist}(x, \partial\Omega) > \frac{1}{i}\} \cap B(0, i)$. We need to verify the hypotheses of lemma 9. First, note that for $x \in \Omega_i, h \in B(0, \delta_x), t \in (0, 1)$, we have

$$\text{dist}(x + th, \partial\Omega) > \text{dist}(x, \partial\Omega) - |h| \geq (1 - \tau)\text{dist}(x, \partial\Omega).$$

Thus $x + th \in \Omega_{i2}$ for $i > \frac{1}{(1-\tau)}$. Thus, we have using triangle inequality and then mean value inequality,

$$\begin{aligned} |F_n(x) - F(x)| &= \left| \left((1 - s_n) \int_{B(0, \delta_x)} \frac{|f(x+h) - f(x)|^q}{|h|^{N+s_nq}} dh \right)^{\frac{p}{q}} - K |\nabla f(x)|^p \right| \\ &\leq \sup_{y \in \Omega_{i2}} |\nabla f(y)|^p \left((1 - s_n) \int_{B(0, R)} \frac{dh}{|h|^{N+s_nq-q}} \right)^{\frac{p}{q}} + K |\nabla f(x)|^p \\ &\leq C(N, p, q, R) \sup_{y \in \Omega_{i2}} |\nabla f(y)|^p. \end{aligned}$$

Since f is continuous in the closure of the bounded open set Ω_{i2} , we have the hypothesis (1) of lemma 9 satisfied for sufficiently large $i \in \mathbb{N}$.

Note that, to show that hypothesis (2) of lemma 9 is satisfied, it is enough to show that

$$\lim_{i \rightarrow \infty} \lim_{n \rightarrow \infty} \int_{\Omega \setminus \Omega_{2i}} F_n(x) dx = 0.$$

We start with an arbitrary $x \in \Omega \setminus \Omega_{2i}, h \in B(0, \delta_x)$ and $t \in (0, 1)$. There can be two cases:

Case:1 $\delta_x = \text{dist}(x, \partial\Omega) < \frac{1}{2i}$.

We have $\text{dist}(x + th, \partial\Omega) \leq |x + th - x| + \text{dist}(x, \partial\Omega) < \frac{\tau}{2i} + \frac{1}{2i} < \frac{1}{i}$. Thus $x + th \in \Omega \setminus \Omega_i$.

Case:2 $|x| > 2i$ and $\frac{1}{2i} < \delta_x = R < \text{dist}(x, \partial\Omega)$. Moreover, we can assume $R < i$ without loss of generality.

We have $|x + th| \geq |x| - \tau R \geq 2i - \tau R \geq i$. Thus $x + th \in \Omega \setminus B(0, i) \subseteq \Omega \setminus \Omega_i$. Hence we always have

$$x + th \in \Omega \setminus \Omega_i \text{ whenever } x \in \Omega \setminus \Omega_{2i}. \tag{17}$$

From eq. (17), we get

$$\lim_{i \rightarrow \infty} \lim_{n \rightarrow \infty} \int_{\Omega \setminus \Omega_{2i}} F_n(x) dx = \lim_{i \rightarrow \infty} \lim_{n \rightarrow \infty} \int_{x \in \Omega \setminus \Omega_{2i}} \left((1 - s_n) \int_{y \in B(x, \delta_x)} \frac{|f(x) - f(y)|^q}{|x - y|^{N+s_nq}} dy \right)^{\frac{p}{q}} dx.$$

Now we apply lemma 11 or lemma 12 with $D = \Omega \setminus \Omega_{2i}$ (so that $\tilde{D} = \Omega \setminus \Omega_i$) to get

$$\begin{aligned} \lim_{i \rightarrow \infty} \lim_{n \rightarrow \infty} \int_{\Omega \setminus \Omega_{2i}} F_n(x) dx &\leq \lim_{i \rightarrow \infty} \lim_{n \rightarrow \infty} C(p, q, R, N) [f]_{W^{1,p}(\Omega \setminus \Omega_i)}^p \\ &= \lim_{i \rightarrow \infty} C(p, q, R, N) [f]_{W^{1,p}(\Omega \setminus \Omega_i)}^p = 0. \end{aligned}$$

Hence we can integrate eq. (13) and interchange the limit and the integral to get the result. □

Theorem 15 Let $\Omega \subset \mathbb{R}^N$ be an open set. If $f \in L^p(\Omega) \cap L^p(\Omega)$ is such that

$$L_{p,q}(f) := \lim_{s \rightarrow 1^-} \int_{\Omega} \left((1 - s) \int_{B(x, \delta_x)} \frac{|f(x) - f(y)|^q}{|x - y|^{N+sq}} dy \right)^{\frac{p}{q}} dx < \infty,$$

then $f \in W^{1,p}(\Omega)$ when $p > 1$, and $f \in BV(\Omega)$ when $p = 1$.

Proof of Theorem 15 We divide the proof into two parts. First, we prove it for a particular case with a bit stronger assumptions, and then give the general proof.

Step-1: Ω is bounded with $\Omega \subseteq B(0, \lambda)$, and

$$\tilde{L}_{p,q}(f) := \lim_{s \rightarrow 1^-} \int_{\Omega} \left((1-s) \int_{\Omega} \frac{|f(x) - f(y)|^q}{|x-y|^{N+sq}} dy \right)^{\frac{p}{q}} dx < \infty.$$

Extend f by 0, outside Ω . From the proof of Theorem 2 and 3 in [8] we can see that for any $i = 1, 2, \dots, N$, and $\varphi \in C_c^\infty(\Omega)$,

$$\left| \int_{\Omega} f(x) \frac{\partial \varphi(x)}{\partial x_i} dx \right| \leq C(\Omega, N, p, q)(1-s)(J_{1,s} + J_{2,s}), \tag{18}$$

where,

$$J_{1,s} = \int_{\Omega} \int_{\Omega} \frac{|f(x) - f(y)|}{|x-y|^{1+N+sq-q}} |\varphi(y)| dy dx$$

and

$$J_{2,s} = \int_{\mathbb{R}^N \setminus \Omega} \int_{\text{supp} \varphi} \frac{|f(y)| |\varphi(y)|}{|x-y|^{1+N+sq-q}} dy dx.$$

We estimate $J_{1,s}$ using Fubini's theorem to change the order of integration, then using Hölder's inequality twice, first with respect to the measure $\frac{dx}{|x-y|^{N+sq-q}}$ and then with respect to dy . We get

$$\begin{aligned} J_{1,s} &\leq \int_{\Omega} \left(\int_{\Omega} \frac{|f(x) - f(y)|^q}{|x-y|^{q+N+sq-q}} dx \right)^{\frac{1}{q}} \left(\int_{\Omega} \frac{|\varphi(y)|^{q'}}{|x-y|^{N+sq-q}} dx \right)^{\frac{1}{q'}} dy \\ &\leq \left(\int_{\Omega} \left(\int_{\Omega} \frac{|f(x) - f(y)|^q}{|x-y|^{N+sq}} dx \right)^{\frac{p}{q}} dy \right)^{\frac{1}{p}} \left(\int_{\Omega} \left(\int_{\Omega} \frac{|\varphi(y)|^{q'}}{|x-y|^{N+sq-q}} dx \right)^{\frac{p'}{q'}} dy \right)^{\frac{1}{p'}} \\ &\leq \left(\int_{\Omega} \left(\int_{\Omega} \frac{|f(x) - f(y)|^q}{|x-y|^{N+sq}} dy \right)^{\frac{p}{q}} dx \right)^{\frac{1}{p}} \left(\int_{\Omega} |\varphi(y)|^{p'} \left(\int_{B(0,\lambda)} \frac{dx}{|x-y|^{N+sq-q}} \right)^{\frac{p'}{q'}} dy \right)^{\frac{1}{p'}} \\ &= C(p, q, N, \lambda)(1-s)^{-\frac{1}{q'}} \left(\int_{\Omega} \left(\int_{\Omega} \frac{|f(x) - f(y)|^q}{|x-y|^{N+sq}} dy \right)^{\frac{p}{q}} dx \right)^{\frac{1}{p}} \|\varphi\|_{L^{p'}(\Omega)} \end{aligned}$$

Now using the hypothesis of Step-1, we have

$$(1-s)J_{1,s} \leq C(p, q, N, \lambda) \tilde{L}_{p,q}(f) \|\varphi\|_{L^{p'}(\Omega)}. \tag{19}$$

Using Hölder's inequality, we estimate $J_{2,s}$ as in [8] to get

$$(1-s)J_{2,s} \leq C(N, p, q, \lambda) \|\varphi\|_{L^{p'}(\Omega)} \|f\|_{L^p(\Omega)} \tag{20}$$

Using eqs. (18) to (20), we get

$$\left| \int_{\Omega} f(x) \frac{\partial \varphi(x)}{\partial x_i} dx \right| \leq C(\Omega, N, p, q, \lambda, f) \|\varphi\|_{L^{p'}(\Omega)},$$

Hence by lemma 6, the result follows.

Step-2: We now prove the theorem in full generality.

For $1 < p < \infty$, define $X^{1,p}(\Omega) := W^{1,p}(\Omega)$, and $X^{1,1}(\Omega) := BV(\Omega)$. Using lemma 8, choose an increasing sequence of bounded open sets $\{\Omega_n\}_n$ with smooth boundary such that $\cup_n \Omega_n = \Omega$, and $\text{dist}(x, \partial\Omega) > \frac{1}{n}$, for $x \in \Omega_n$. From the hypothesis, it follows that

$$\lim_{s \rightarrow 1^-} \int_{x \in \Omega_n} \left((1-s) \int_{y \in \Omega_n \cap B(x, \delta_x)} \frac{|f(x) - f(y)|^q}{|x-y|^{N+sq}} dy \right)^{\frac{p}{q}} dx < \infty.$$

We also have, for $s > \frac{1}{2}$, and $R > \frac{1}{n}$,

$$\begin{aligned} \int_{\Omega_n} \left(\int_{\Omega_n \setminus B(x, \delta_x)} \frac{|f(x) - f(y)|^q}{|x-y|^{N+sq}} dy \right)^{\frac{p}{q}} dx &\leq \int_{\Omega_n} \left(\int_{\Omega_n, |x-y| > \frac{1}{n}} \frac{|f(x) - f(y)|^q}{|x-y|^{N+sq}} dy \right)^{\frac{p}{q}} dx \\ &\leq n^{\frac{Np}{q} + sp} \int_{\Omega_n} \left(\int_{\Omega_n, |x-y| > \frac{1}{n}} |f(x) - f(y)|^q dy \right)^{\frac{p}{q}} dx \\ &\leq C(n, N, p, q) \left[\|f\|_{L^p(\Omega_n)}^p |\Omega_n|^{\frac{p}{q}} + \|f\|_{L^q(\Omega_n)}^p |\Omega_n| \right]. \end{aligned}$$

Since, we have $f \in L^p(\Omega) \cap L^q(\Omega)$ from the hypotheses, and Ω_n are bounded domains, we have

$$\lim_{s \rightarrow 1^-} \int_{\Omega_n} \left((1-s) \int_{\Omega_n} \frac{|f(x) - f(y)|^q}{|x-y|^{N+sq}} dy \right)^{\frac{p}{q}} dx < \infty.$$

From Step-1, we can conclude that $f \in X^{1,p}(\Omega_n)$ for all n . Further the $X^{1,p}$ -seminorms are uniformly bounded (independent of n) as can be seen from the following calculation, where we use theorem 14,

$$\begin{aligned} K[f]_{X^{1,p}(\Omega_n)}^p &= \lim_{s \rightarrow 1^-} \int_{x \in \Omega_n} \left((1-s) \int_{y \in B(x, \delta_x, \Omega_n)} \frac{|f(x) - f(y)|^p}{|x-y|^{N+sp}} dy \right)^{\frac{p}{q}} dx \\ &\leq \lim_{s \rightarrow 1^-} \int_{x \in \Omega} \left((1-s) \int_{y \in B(x, \delta_x)} \frac{|f(x) - f(y)|^p}{|x-y|^{N+sp}} dy \right)^{\frac{p}{q}} dx \\ &= L_{p,q}(f) < \infty. \end{aligned}$$

The proof follows from the observation that $K[f]_{X^{1,p}(\Omega)}^p = \sup_n K[f]_{X^{1,p}(\Omega_n)}^p$. □

5 Proofs of Theorems 1 and 2

Note that theorem 2 is a straightforward consequence of theorem 15, as $L_{p,q}(f) \leq L_{p,q}^*(f)$. theorem 1 is also a consequence of theorem 14, but it requires a bit more work. To complete the proof of theorem 1, we only need the following lemma:

Lemma 16 *Let $\Omega \subseteq \mathbb{R}^N$ be an open set, $1 \leq p < \infty$, $\tau \in (0, 1)$, $f \in W^{1,p}(\Omega)$ and $R > 0$. Assume one of the following conditions*

- (1) $1 \leq q \leq \frac{Np}{N-p}$ with $p < N$,
- (2) $1 \leq q < \infty$ with $p \geq N$.

Then eq. (14) implies eq. (4).

In order to prove this, we first prove a bit more general result.

Proposition 17 Let $\Omega \subseteq \mathbb{R}^N$ be an open set, $1 \leq p, q < \infty$, $\tau \in (0, 1)$, $f \in L^p(\Omega) \cap L^q(\Omega)$ and $R > 0$. Additionally, in the case $p < q$, assume that for some $s_0 \in (0, 1)$,

$$\int_{\Omega} \left(\int_{R \leq |h| \leq \tau \text{dist}(x, \partial\Omega)} \frac{|f(x+h) - f(x)|^q}{|h|^{N+s_0q}} dh \right)^{\frac{p}{q}} dx < \infty. \tag{21}$$

Then eq. (14) implies eq. (4).

Proof of Proposition 17 Note that, since

$$\int_{\Omega} \left(\int_{B(x, \delta_x)} \frac{|f(x) - f(y)|^q}{|x - y|^{N+s_0q}} dy \right)^{\frac{p}{q}} dx \leq \int_{\Omega} \left(\int_{B(x, \tau \text{dist}(x, \partial\Omega))} \frac{|f(x) - f(y)|^q}{|x - y|^{N+s_0q}} dy \right)^{\frac{p}{q}} dx,$$

from eq. (14) we have

$$\lim_{s \rightarrow 1-} (1 - s)^{\frac{p}{q}} \int_{\Omega} \left(\int_{B(x, \tau \text{dist}(x, \partial\Omega))} \frac{|f(x) - f(y)|^q}{|x - y|^{N+s_0q}} dy \right)^{\frac{p}{q}} dx \geq K \int_{\Omega} |\nabla f(x)|^p dx.$$

We focus on the reverse inequality. Observe that for $x, y \in \Omega$, $\delta_x < |x - y| \leq \tau \text{dist}(x, \partial\Omega)$ implies $R < |x - y| \leq \tau \text{dist}(x, \partial\Omega)$. Hence we can write, using triangle inequality for L^p -norms,

$$\begin{aligned} & \left(\int_{\Omega} \left(\int_{B(x, \tau \text{dist}(x, \partial\Omega))} \frac{|f(x) - f(y)|^q}{|x - y|^{N+s_0q}} dy \right)^{\frac{p}{q}} dx \right)^{\frac{1}{p}} \\ &= \left(\int_{\Omega} \left(\int_{B(x, \delta_x)} \frac{|f(x) - f(y)|^q}{|x - y|^{N+s_0q}} dy + \int_{\delta_x \leq |x-y| \leq \tau \text{dist}(x, \partial\Omega)} \frac{|f(x) - f(y)|^q}{|x - y|^{N+s_0q}} dy \right)^{\frac{p}{q}} dx \right)^{\frac{1}{p}} \\ &\leq \left(\int_{\Omega} \left(\int_{B(x, \delta_x)} \frac{|f(x) - f(y)|^q}{|x - y|^{N+s_0q}} dy \right)^{\frac{p}{q}} dx \right)^{\frac{1}{p}} \\ &\quad + \left(\int_{\Omega} \left(\int_{R \leq |x-y| \leq \tau \text{dist}(x, \partial\Omega)} \frac{|f(x) - f(y)|^q}{|x - y|^{N+s_0q}} dy \right)^{\frac{p}{q}} dx \right)^{\frac{1}{p}} \\ &=: I_1^{\frac{1}{p}} + I_2^{\frac{1}{p}}. \end{aligned}$$

In order to complete the proof, in view of eq. (14), we need to show that I_2 is bounded as $s \rightarrow 1-$. We estimate I_2 in two separate cases.

Case-1: $1 \leq q \leq p < \infty$.

Using Minkowsky’s integral inequality and taking the 0-extension of f outside Ω , we have

$$\begin{aligned} I_2^{\frac{q}{p}} &\leq \left(\int_{\Omega} \left(\int_{R \leq |h| \leq \tau \text{dist}(x, \partial\Omega)} \frac{|f(x+h) - f(x)|^q}{|h|^{N+s_0q}} dh \right)^{\frac{p}{q}} dx \right)^{\frac{q}{p}} \\ &\leq \int_{\mathbb{R}^N} \left(\int_{\Omega} \frac{|f(x+h) - f(x)|^p}{|h|^{\frac{Np}{q} + s_0p}} \chi_{B(0, \tau \text{dist}(x, \partial\Omega)) \setminus B(0, R)}(h) dx \right)^{\frac{q}{p}} dh \\ &\leq \int_{|h| \geq R} \frac{1}{|h|^{N+s_0q}} \left(\int_{\Omega} |f(x+h) - f(x)|^p dx \right)^{\frac{q}{p}} dh \\ &\leq C(p, q) \int_{|h| \geq R} \frac{1}{|h|^{N+s_0q}} \left(\int_{\Omega} |f(x)|^p dx \right)^{\frac{q}{p}} dh \end{aligned}$$

$$\leq C(p, q, R, N) \|f\|_{L^p(\Omega)}^q.$$

Hence the proof follows in this case.

Case-2: $1 \leq p \leq q < \infty$.

From eq. (21) we get that there is some $\lambda_f > 0$ such that for $s \in (s_0, 1)$,

$$\begin{aligned} I_2 &\leq \int_{\Omega} \left(\int_{R \leq |h| \leq \tau \text{dist}(x, \partial\Omega)} \frac{|f(x+h) - f(x)|^q}{|h|^{N+s_0q}} dh \right)^{\frac{p}{q}} dx \\ &\leq 2 \int_{\Omega \cap B(0, \lambda_f)} \left(\int_{R \leq |h| \leq \tau \text{dist}(x, \partial\Omega)} \frac{|f(x+h) - f(x)|^q}{|h|^{N+s_0q}} dh \right)^{\frac{p}{q}} dx. \end{aligned}$$

Using Hölder’s inequality, we get

$$I_2 \leq 2\lambda_f^{N(1-\frac{p}{q})} \left(\int_{\Omega \cap B(0, \lambda_f)} \int_{R \leq |h| \leq \tau \text{dist}(x, \partial\Omega)} \frac{|f(x+h) - f(x)|^q}{|h|^{N+s_0q}} dh dx \right)^{\frac{p}{q}}.$$

Now we can proceed as in Case-1 to show that

$$I_2 \leq C(N, p, q, R, f) \|f\|_{L^q(\Omega)}^p.$$

This completes the proof. □

Now we can prove lemma 16 and thereby complete the proof of theorem 1.

Proof of lemma 16 By the standard embedding theorems, we already know that $f \in L^q(\Omega)$. In order to prove the statement, we need to show that when $p < q$, eq. (21) holds. Let Ω_1 be a smooth domain such that

$$\{x \in \Omega \mid \text{dist}(x, \partial\Omega) > R\} \subseteq \Omega_1 \subseteq \Omega.$$

Clearly Ω_1 is a $W^{1,p}$ -extension domain (by lemma 5). Let $\tilde{f} \in W^{1,p}(\mathbb{R}^N)$ be an extension of $f|_{\Omega_1}$, that is

$$\|\tilde{f}\|_{W^{1,p}(\mathbb{R}^N)} \leq C(p, q, N, \Omega) \|f\|_{W^{1,p}(\Omega_1)} \leq C(p, q, N, \Omega) \|f\|_{W^{1,p}(\Omega)} < \infty.$$

This, along with lemma 7

$$\begin{aligned} &\int_{\Omega} \left(\int_{R \leq |h| \leq \tau \text{dist}(x, \partial\Omega)} \frac{|f(x+h) - f(x)|^q}{|h|^{N+sq}} dh \right)^{\frac{p}{q}} dx \\ &= \int_{\Omega_1} \left(\int_{R \leq |h| \leq \tau \text{dist}(x, \partial\Omega)} \frac{|f(x+h) - f(x)|^q}{|h|^{N+sq}} dh \right)^{\frac{p}{q}} dx \\ &\leq \int_{\mathbb{R}^N} \left(\int_{\mathbb{R}^N} \frac{|\tilde{f}(x+h) - \tilde{f}(x)|^q}{|h|^{N+sq}} dh \right)^{\frac{p}{q}} dx \\ &< \infty. \end{aligned}$$

□

Acknowledgements I extend my heartfelt gratitude to Antti Vähäkangas for his invaluable discussions, meticulous review of the manuscript, and unwavering support throughout this project. I want to thank Pekka Koskela for his insightful contributions during the manuscript preparation. Additionally, my appreciation goes to Emiel Lorist for pointing out an error in the initial version of the article, and to the anonymous reviewer for his/her valuable comments and suggestions.

Funding Open Access funding provided by University of Jyväskylä (JYU). The research is funded by Academy of Finland grant: Geometrinen Analyysi (21000046081).

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Alberico, Angela, Cianchi, Andrea, Pick, Luboš, Slavíková, Lenka: On the limit as $s \rightarrow 1^-$ of possibly non-separable fractional Orlicz-Sobolev spaces. *Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl.* **31**(4), 879–899 (2020)
2. Alberico, Angela, Cianchi, Andrea, Pick, Luboš, Slavíková, Lenka: Fractional Orlicz-Sobolev embeddings. *J. Math. Pures Appl.* **9**(149), 216–253 (2021)
3. Alberico, Angela, Cianchi, Andrea, Pick, Luboš, Slavíková, Lenka: On fractional Orlicz-Sobolev spaces. *Anal. Math. Phys.* **11**(2), 84, 21, (2021)
4. Arroyo-Rabasa, Adolfo, Bonicatto, Paolo: A Bourgain-Brezis-Mironescu representation for functions with bounded deformation. *Calc. Var. Partial Differ. Equ.* **62**(1), 33, 22, (2023)
5. Bal, Kaushik, Mohanta, Kaushik, Roy, Prosenjit: Bourgain-Brezis-Mironescu domains. *Nonlinear Anal.* **199**, 111928, 10, (2020)
6. Barza, Sorina, Lind, Martin: A new variational characterization of Sobolev spaces. *J. Geom. Anal.* **25**(4), 2185–2195 (2015)
7. Bojarski, B.: Remarks on the Bourgain-Brezis-Mironescu approach to Sobolev spaces. *Bull. Pol. Acad. Sci. Math.* **59**(1), 65–75 (2011)
8. Bourgain, Jean, Brezis, Haim, Mironescu, Petru. Another look at Sobolev spaces. In *Optimal control and partial differential equations*, 439–455. IOS, Amsterdam. (2001)
9. Bourgain, Jean, Brezis, Haim, Mironescu, Petru, Limiting embedding theorems for $W^{s,p}$ when $s \uparrow 1$ and applications. **87**, 77–101. (2002). Dedicated to the memory of Thomas H. Wolff
10. Bourgain, Jean, Nguyen, Hoai-Minh.: A new characterization of Sobolev spaces. *C. R. Math. Acad. Sci. Paris* **343**(2), 75–80 (2006)
11. Brasco, Lorenzo, Gómez-Castro, David, Vázquez, Juan Luis: Characterisation of homogeneous fractional Sobolev spaces. *Calc. Var. Partial Differ. Equ.* **60**(2), 60, 40 (2021)
12. Brazke, Denis, Schikorra, Armin, Yung, Po-Lam.: Bourgain-Brezis-Mironescu convergence via Triebel-Lizorkin spaces. *Calc. Var. Partial Differ. Equ.* **62**(2), 41, 33, (2023)
13. Brezis, Haim: *Functional analysis. Sobolev spaces and partial differential equations*. Universitext. Springer, New York (2011)
14. Brezis, Haïm., Nguyen, Hoai-Minh.: The BBM formula revisited. *Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl.* **27**(4), 515–533 (2016)
15. Brezis, Haïm., Seeger, Andreas, Van Schaftingen, Jean, Yung, Po-Lam.: Sobolev spaces revisited. *Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl.* **33**(2), 413–437 (2022)
16. Brezis, Haïm., Van Schaftingen, Jean, Yung, Po-Lam.: Going to Lorentz when fractional Sobolev, Gagliardo and Nirenberg estimates fail. *Calc. Var. Partial Differ. Equ.* **60**(4), 129, 12 (2021)
17. Brezis, Haïm., Van Schaftingen, Jean, Yung, Po-Lam.: A surprising formula for Sobolev norms. *Proc. Natl. Acad. Sci. USA* **118**(8), e2025254118, 6, (2021)
18. Brezis, Kh.: How to recognize constant functions A. connection with Sobolev spaces. *Uspekhi Mat. Nauk* **57**(4(346)), 59–74 (2002)

19. Buseghin, Federico, Garofalo, Nicola, Tralli, Giulio: On the limiting behaviour of some nonlocal semi-norms: a new phenomenon (5). *Ann. Sc. Norm. Super. Pisa Cl. Sci.* **23**(2), 837–875 (2022)
20. Calderón, A.-P.: Lebesgue spaces of differentiable functions and distributions. In *Proc. Sympos. Pure Math., Vol. IV*, pages 33–49. American Mathematical Society, Providence, R.I., (1961)
21. Cui, Xiaoyue, Lam, Nguyen, Guozhen, Lu.: Characterizations of second order Sobolev spaces. *Nonlinear Anal.* **121**, 241–261 (2015)
22. Dai, Feng, Grafakos, Loukas, Pan, Zhulei, Yang, Dachun, Yuan, Wen, Zhang, Yangyang: The Bourgain-Brezis-Mironescu formula on ball Banach function spaces. *Math. Ann.* (2023)
23. Dávila, J.: On an open question about functions of bounded variation. *Calc. Var. Partial Differ. Equ.* **15**(4), 519–527 (2002)
24. Di Marino, Simone, Squassina, Marco: New characterizations of Sobolev metric spaces. *J. Funct. Anal.* **276**(6), 1853–1874 (2019)
25. Di Nezza, Eleonora, Palatucci, Giampiero, Valdinoci, Enrico: Hitchhiker’s guide to the fractional Sobolev spaces. *Bull. Sci. Math.* **136**(5), 521–573 (2012)
26. Domínguez, Oscar, Milman, Mario: Bourgain-Brezis-Mironescu-Maz’ya-Shaposhnikova limit formulae for fractional Sobolev spaces via interpolation and extrapolation. *Calc. Var. Partial Differ. Equ.* **62**(2), 43, 37, (2023)
27. Drelichman, Irene, Durán, Ricardo G.: The Bourgain-Brezis-Mironescu formula in arbitrary bounded domains. *Proc. Amer. Math. Soc.* **150**(2), 701–708 (2022)
28. Bonder, Julián Fernández., Salort, Ariel M.: Fractional order Orlicz-Sobolev spaces. *J. Funct. Anal.* **277**(2), 333–367 (2019)
29. Bonder, Julián Fernández., Salort, Ariel M.: Magnetic fractional order Orlicz-Sobolev spaces. *Studia Math.* **259**(1), 1–24 (2021)
30. Ferrari, Gianluca, Squassina, Marco: Nonlocal characterizations of variable exponent Sobolev spaces. *Asymptot. Anal.* **127**(1–2), 121–142 (2022)
31. Ferreira, Rita, Hästö, Peter, Ribeiro, Ana Margarida: Characterization of generalized Orlicz spaces. *Commun. Contemp. Math.* **22**(2), 1850079, 25, (2020)
32. Foghem, Guy, L2-theory for nonlocal operators on domains. *Publikationen an der Universität Bielefeld*, (2020)
33. Gounoue, Guy Fabrice Foghem.: A remake of Bourgain-Brezis-Mironescu characterization of Sobolev spaces. *Partial Differ. Equ. Appl.* **4**(2), 16, 36, (2023)
34. Gounoue, Guy Fabrice Foghem., Kassmann, Moritz, Voigt, Paul: Mosco convergence of nonlocal to local quadratic forms. *Nonlinear Anal.* **193**, 111504, 22, (2020)
35. Górný, Wojciech: Bourgain-Brezis-Mironescu approach in metric spaces with Euclidean tangents. *J. Geom. Anal.* **32**(4), 128, 22, (2022)
36. Grafakos, Loukas: Classical and modern Fourier analysis. Pearson Education Inc, Upper Saddle River, NJ (2004)
37. Qingsong, Gu., Huang, Qingzhong: Anisotropic versions of the Brezis-Van Schaftingen-Yung approach at $s = 1$ and $s = 0$. *J. Math. Anal. Appl.* **525**(2), 127156, 15, (2023)
38. Hajl asz, Piotr, A new characterization of the Sobolev space. **159**, pages 263–275. (2003). Dedicated to Professor Aleksander Pełczyński on the occasion of his 70th birthday (Polish)
39. Hästö, Peter: Ribeiro, Ana Margarida, Characterization of the variable exponent Sobolev norm without derivatives. *Commun. Contemp. Math.* **19**(3), 1650022 (2017)
40. Kamihigashi, Takashi: Interchanging a limit and an integral: necessary and sufficient conditions. *J. Inequal. Appl.* **243**, 9 (2020)
41. Kolyada, V.I., Lerner, A.K.: On limiting embeddings of Besov spaces. *Studia Math.* **171**(1), 1–13 (2005)
42. Kreuml, Andreas, Mordhorst, Olaf: Fractional Sobolev norms and BV functions on manifolds. *Nonlinear Anal.* **187**, 450–466 (2019)
43. Leoni, Giovanni, Spector, Daniel: Characterization of Sobolev and BV spaces. *J. Funct. Anal.* **261**(10), 2926–2958 (2011)
44. Leoni, Giovanni, Spector, Daniel: Corrigendum to Characterization of Sobolev and BV spaces. [*J. Funct. Anal.* **261** (10) (2011) 2926–2958]. *J. Funct. Anal.* **266**(2), 1106–1114 (2014)
45. Ludwig, Monika: Anisotropic fractional Sobolev norms. *Adv. Math.* **252**, 150–157 (2014)
46. Milman, Mario: Notes on limits of Sobolev spaces and the continuity of interpolation scales. *Trans. Amer. Math. Soc.* **357**(9), 3425–3442 (2005)
47. Nguyen, Hoai-Minh.: Some new characterizations of Sobolev spaces. *J. Funct. Anal.* **237**(2), 689–720 (2006)
48. Nguyen, Hoai-Minh.: Further characterizations of Sobolev spaces. *J. Eur. Math. Soc. (JEMS)* **10**(1), 191–229 (2008)

49. Nguyen, Hoai-Minh., Pinamonti, Andrea, Squassina, Marco, Vecchi, Eugenio: New characterizations of magnetic Sobolev spaces. *Adv. Nonlinear Anal.* **7**(2), 227–245 (2018)
50. Nguyen, Hoai-Minh., Pinamonti, Andrea, Squassina, Marco, Vecchi, Eugenio: Some characterizations of magnetic Sobolev spaces. *Complex Var. Elliptic Equ.* **65**(7), 1104–1114 (2020)
51. Nguyen, Hoai-Minh., Squassina, Marco: On anisotropic Sobolev spaces. *Commun. Contemp. Math.* **21**(1), 1850017, 13, (2019)
52. Pinamonti, Andrea, Squassina, Marco, Vecchi, Eugenio: Magnetic BV-functions and the Bourgain-Brezis-Mironescu formula. *Adv. Calc. Var.* **12**(3), 225–252 (2019)
53. Ponce, Augusto C.: A new approach to Sobolev spaces and connections to Γ -convergence. *Calc. Var. Partial Differential Equations* **19**(3), 229–255 (2004)
54. Ponce, Augusto C., Spector, Daniel: On formulae decoupling the total variation of BV functions. *Nonlinear Anal.* **154**, 241–257 (2017)
55. Prats, Martí, Saksman, Eero: A $T(1)$ theorem for fractional Sobolev spaces on domains. *J. Geom. Anal.* **27**(3), 2490–2538 (2017)
56. Runst, Thomas, Sickel, Winfried: Sobolev spaces of fractional order, Nemytskij operators, and nonlinear partial differential equations. *De Gruyter Series in Nonlinear Analysis and Applications*, vol. 3. Walter de Gruyter & Co., Berlin (1996)
57. Squassina, Marco, Volzone, Bruno: Bourgain-Brézis-Mironescu formula for magnetic operators. *C. R. Math. Acad. Sci. Paris* **354**(8), 825–831 (2016)
58. Stein, Elias M.: *Singular integrals and differentiability properties of functions*. Princeton Mathematical Series, No. 30. Princeton University Press, Princeton, N.J., (1970)
59. Triebel, Hans: *Theory of function spaces*. Monographs in Mathematics, vol. 78. Birkhäuser Verlag, Basel (1983)
60. Triebel, Hans: Limits of Besov norms. *Arch. Math. (Basel)* **96**(2), 169–175 (2011)
61. Yang, Sibe, Yang, Dachun, Yuan, Wen: New characterizations of Musielak-Orlicz-Sobolev spaces via sharp ball averaging functions. *Front. Math. China* **14**(1), 177–201 (2019)
62. Zhu, Chenfeng, Yang, Dachun, Yuan, Wen, Extension theorem and bourgain-brezis-mironescu-type characterization of ball banach sobolev spaces on domains. *ArXiv preprint arXiv: 2307.11392*, (2023)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.