

This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Iso-Markku, Paula; Aaltonen, Sari; Kujala, Urho M.; Halme, Hanna-Leena; Phipps, Daniel; Knittle, Keegan; Vuoksimaa, Eero; Waller, Katja

Title: Physical Activity and Cognitive Decline Among Older Adults : A Systematic Review and Meta-Analysis

Year: 2024

Version: Published version

Copyright: © 2024 the Authors

Rights: CC BY 4.0

Rights url: <https://creativecommons.org/licenses/by/4.0/>

Please cite the original version:

Iso-Markku, P., Aaltonen, S., Kujala, U. M., Halme, H.-L., Phipps, D., Knittle, K., Vuoksimaa, E., & Waller, K. (2024). Physical Activity and Cognitive Decline Among Older Adults : A Systematic Review and Meta-Analysis. *JAMA Network Open*, 7(2), Article e2354285.
<https://doi.org/10.1001/jamanetworkopen.2023.54285>



Original Investigation | Neurology

Physical Activity and Cognitive Decline Among Older Adults A Systematic Review and Meta-Analysis

Paula Iso-Markku, MD, PhD; Sari Aaltonen, PhD; Urho M. Kujala, MD, PhD; Hanna-Leena Halme, PhD; Daniel Phipps, PhD; Keegan Knittle, PhD; Eero Vuoksimaa, PhD; Katja Waller, PhD

Abstract

IMPORTANCE Physical activity is associated with the risk for cognitive decline, but much of the evidence in this domain comes from studies with short follow-ups, which is prone to reverse causation bias.

OBJECTIVE To examine how length of follow-up, baseline age, physical activity amount, and study quality modify the longitudinal associations of physical activity with cognition.

DATA SOURCES Observational studies of adults with a prospective follow-up of at least 1 year, a valid baseline cognitive measure or midlife cohort, and an estimate of the association of baseline physical activity and follow-up cognition were sought from PsycInfo, Scopus, CINAHL, Web of Science, SPORTDiscus, and PubMed, with the final search conducted on November 2, 2022.

STUDY SELECTION Two independent researchers screened titles with abstracts and full-text reports.

DATA EXTRACTION AND SYNTHESIS Two reviewers independently assessed study quality and extracted data. Pooled estimates of association were calculated with random-effects meta-analyses. An extensive set of moderators, funnel plots, and scatter plots of physical activity amount were examined. This study is reported following the Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) reporting guideline.

MAIN OUTCOMES AND MEASURES Pooled estimates of the associations between physical activity and global cognition, as well as specific cognitive domains, were examined.

RESULTS A total of 104 studies with 341 471 participants were assessed. Analysis of binary outcomes included 45 studies with 102 452 individuals, analysis of follow-up global cognition included 14 studies with 41 045 individuals, and analysis of change in global cognition included 25 studies with 67 463 individuals. Physical activity was associated with a decreased incidence of cognitive impairment or decline after correction for funnel plot asymmetry (pooled risk ratio, 0.97; 95% CI, 0.97-0.99), but there was no significant association in follow-ups longer than 10 years. Physical activity was associated with follow-up global cognition (standardized regression coefficient, 0.03; 95% CI, 0.02-0.03) and change in global cognition (standardized regression coefficient, 0.01; 95% CI, 0.01 to 0.02) from trim-and-fill analyses, with no clear dose-response or moderation by follow-up length, baseline age, study quality or adjustment for baseline cognition. The specific cognitive domains associated with physical activity were episodic memory (standardized regression coefficient, 0.03; 95% CI, 0.02-0.04) and verbal fluency (standardized regression coefficient, 0.05; 95% CI, 0.03-0.08).

(continued)

Key Points

Question Is physical activity associated with cognitive decline?

Findings This systematic review and meta-analysis including a total of 104 studies with 341 471 participants found a weak association between baseline physical activity and follow-up global cognition that was evident also in episodic memory and verbal fluency domains. Neither study quality, follow-up length, baseline age, nor adjustment for preceding level of cognition moderated the association, and there was no clear dose-response association between the amount of physical activity and global cognition.

Meaning These findings suggest that physical activity might postpone cognitive decline at a population health level but only to a very small extent.

+ Supplemental content

Author affiliations and article information are listed at the end of this article.

Open Access. This is an open access article distributed under the terms of the CC-BY License.

Abstract (continued)

CONCLUSIONS AND RELEVANCE In this meta-analysis of the association of physical activity with cognitive decline, physical activity was associated with better late-life cognition, but the association was weak. However, even a weak association is important from a population health perspective.

JAMA Network Open. 2024;7(2):e2354285. doi:10.1001/jamanetworkopen.2023.54285

Introduction

Extensive research links physical activity with better cognitive outcomes across the lifespan¹ and a decreased risk of dementia.^{2,3} Many different specific cognitive domains have been suggested to be associated with physical activity, but the evidence is inconsistent.^{4,5} Despite the optimism surrounding physical activity as a means to preserve or improve cognition, many recent high-quality interventional studies urge caution in claims linking cognitive benefits to physical activity,⁶⁻⁸ and most existing evidence comes from observational studies with short follow-ups and no information on preceding levels of cognition.

Observational prospective cohort studies can provide information on the association between risk factors and outcomes over very long follow-up periods, despite their inherent weaknesses. Earlier meta-analyses on physical activity and cognition have found a clear association between higher levels of physical activity and a decreased risk of subsequent cognitive decline⁹⁻¹²; however, these meta-analyses combined only studies with binary outcomes: cognitive impairment or decline or no cognitive impairment or decline⁹⁻¹² and have examined only a few moderators. While binary outcomes are clinically relevant, modeling cognition as a continuous variable would improve statistical power.

This meta-analysis investigates whether physical activity is associated with global and domain-specific cognitive decline, examining cognition both categorically and continuously. Furthermore, we explore a possible dose-response association of physical activity with cognition and a broad range of possible moderators of this association. Taken together, this meta-analysis accounts for known limitations of research in this area to improve understanding of the association between physical activity participation and cognitive decline.

Methods

This systematic review and meta-analysis was preregistered in PROSPERO (CRD42018083236) and follows the Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) reporting guideline.¹³ The original registered plan was adapted according to data found. These changes are described in the eMethods in Supplement 1.

Eligibility Criteria

Types of Studies and Participants

We included prospective cohort studies or case-control studies presenting an estimate of association between baseline physical activity and cognition after at least 1 year of follow-up. Participants were adults aged at least 20 years. We excluded cohorts with a specific disease, established dementia, or cognitive impairment at baseline. To be included, studies with baseline in later life (defined as mean or median age ≥ 55 years and maximum age ≥ 65 years or mean age within 1 SD of ≥ 60 years) were required to report a valid measure of cognition at baseline to reduce possible reverse causality.

Types of Exposure

Studies measuring contemporaneous physical activity with interviews, questionnaires, or devices were included. We excluded studies measuring retrospective physical activity level, cardiorespiratory

or other fitness levels, single bouts of physical activity, physical activity extending over the follow-up, or statistical reallocations of physical activity.

Types of Outcomes

The association of baseline physical activity and cognition or specific cognitive domain (ie, executive function, episodic memory, processing speed, verbal fluency and naming, verbal ability, working memory, and visuospatial ability) at follow-up. We excluded studies reporting subjective estimates of cognition and studies in which cognition was based on registers of disability level.

Types of Reports

We included reports with full texts in English. We previously reported specifications to the exclusion and inclusion criteria in the screening and full-text review phases.³ Additional specifications in cases of disagreements were made during the update in 2022 to 2023. Changes are described in the eMethods in [Supplement 1](#).

Search Strategy

We conducted a systematic literature search in 6 databases (PubMed, PsycInfo, CINAHL, Scopus, SPORTDiscus, Web of Science). The last search was conducted November 2, 2022. Keywords included were *physical activity, sport, athletics, walking, physical training, cognition, cognitive, executive function, TELE* (telephone assessment for dementia), *TICS* (Telephone Interview for Cognitive Status), *MMSE* (Mini-Mental State Examination), *3-MS* (the Modified Mini-Mental State Examination), *memory, processing speed, verbal fluency, semantic fluency, reasoning, delayed recall, prospective, longitudinal, follow-up, observational, and cohort*. Individual additional articles known to the authors were also added. Example searches and further details of the search have been described in eTable 1 in [Supplement 1](#).

Study Selection

Two reviewers (P.I.-M. and S.A., D.P., or K.W.) independently assessed the eligibility of each study in 2 phases: title and abstract screening and full-text review. Disagreements were discussed, and a third reviewer (U.M.K.) made the final decision of inclusion if consensus was not reached. In cases of multiple studies reporting results for the same outcome from the same cohort from an overlapping time period, the study with a higher study quality, longer follow-up, larger sample, or which adjusted for baseline cognition was chosen.

Quality Assessment

We developed a quality assessment tool with 3 tiered ranking (high, moderate, low) specifically for this meta-analysis.³ This tool provides high transparency of the quality assessment and accounts for the special characteristics of measuring physical activity and its association with an outcome with a long preclinical period extending over decades (cognitive impairment or dementia).¹⁴ One modification to this tool from our previous meta-analysis³ is described in the eMethods in [Supplement 1](#). Two reviewers (P.I.-M. and D.P. or K.W.) conducted the quality review at the outcome level independently. Any disagreements were resolved with discussion.

Data Extraction

Two reviewers (P.I.-M. and S.A., H.-L.H., D.P., or K.W.) extracted the following data: estimate of the association between physical activity and cognition or specific cognitive domain, measure of cognition, proportion of the sample who became cognitively impaired (or SD for continuous cognition outcomes), physical activity levels, sample size, country of origin, publication year, length of follow-up, age at baseline, work-related or leisure-time physical activity, confounders (cognition at baseline, chronic diseases, education, vascular risk factors, APOE ϵ 4 allele), the number of confounders, device-based measure or self-report of physical activity, cohort, and follow-up rate.

The extracted data were compared, and disagreements were solved by discussion. Follow-up rate was extracted by a single reviewer (P.I.-M.). Additional details of the data extraction are provided in eMethods and eTables 2-4 in [Supplement 1](#) and eTables 1-12 in [Supplement 2](#).

Statistical Analysis

The results extracted from the original studies were risk ratios (RRs), hazard ratios, odds ratios, regression coefficients for cognition, number of participants with and without cognitive impairment at follow-up, regression coefficients for change in cognition, regression coefficients for rate of change in cognition, mean changes with 95% CIs, pre- and post-follow-up means and SDs, adjusted *P* values from analysis of covariance, means from repeated measures analysis of variance, and post-follow-up means and SDs with *P* values for a difference from a generalized linear regression model. We performed 3 separate sets of analyses for 3 different outcome types: a first set of analyses with risk of cognitive impairment or decline as the binary outcome, a second analysis with follow-up global cognition as a continuous outcome, and a third analysis with change in global cognition as a continuous outcome. Full details of transformations, standardizations, and the use of studies with cognition trajectories are described in eMethods in [Supplement 1](#) and eTable 11 and eTable 12 in [Supplement 2](#).

Binary Outcomes

We transformed hazard and odds ratios and number of participants with and without event at follow-up into risk ratios and pooled these risk ratios. In the main analyses, we compared all other physical activity levels with the lowest physical activity level.³

Continuous Outcomes

Follow-Up Cognition | We standardized unstandardized regression coefficients by dividing them by the SD of the outcome measure of cognition and multiplying them by the SD of physical activity.¹⁵ After standardization, we pooled the standardized regression coefficients for follow-up cognition with standardized mean differences in post-follow-up scores to yield us a pooled magnitude of association for physical activity and cognition. The basis for pooling standardized regression coefficients for cognition with standardized mean differences of the outcome post follow-up is that, although apparently different, the comparison of means of 2 independent groups measured at a single time with multiple regression analysis is mathematically equivalent with Cohen *d* from a *t* test for independent groups or between-participants analysis of variance with 2 groups.^{15,16} Thus, for 2 physical activity groups, the difference in cognition can be calculated with either of the 2 following equations:

$$d = SD(X) \frac{M_T - M_C}{SD(Y)}$$

$$d = SD(X) \frac{b}{SD(Y)}$$

Where *d* indicates Cohen *d*; *M_T*, the mean of the treatment group; *M_C*, the mean of the control group; *SD(Y)*, the pooled within-group SD of the outcome measure; *SD(X)*, the SD of the independent variable; and *b*, the unstandardized regression coefficient for the association of the group.

Change in Cognition | In the set of analyses with change in cognition during the follow-up, we standardized the regression coefficients in a similar manner as for follow-up global cognition, but instead used the SD of cognition at baseline (between-participant variability at baseline).^{16,17} We pooled all standardized regression coefficients for change with standardized mean differences of change¹⁶ to yield 1 estimate for the association of physical activity with change in cognition.

Pooling the Studies

We used random-effects models with inverse variance as the weighting method for our meta-analysis. Heterogeneity was estimated with DerSimonian and Laird method. The possibility of publication bias was estimated with contour-enhanced funnel plots. We used the trim-and-fill method with run estimator to correct any funnel plot asymmetry. We present the results as pooled RRs with 95% CIs for studies with binary outcomes and as pooled standardized regression coefficients with 95% CIs for studies with continuous outcomes.

The main analyses were risk of cognitive impairment or decline (ie, analysis of binary outcomes), standardized regression coefficient for follow-up global cognition, and standardized regression coefficient for change in global cognition during the follow-up. In addition, we performed separate analyses for the following specific cognitive domains: executive function, working memory, processing speed, episodic memory, verbal fluency and naming, verbal ability, and visuospatial ability. In the sensitivity analyses, we examined categorical moderators with subgroup analyses and continuous moderators with meta-regressions. The following moderators were examined: follow-up length, baseline and follow-up age, study quality, type of physical activity, measurement of physical activity, validity of physical activity measurement, sample size, number of confounders, measurement of cognition (at least 1 neuropsychological test vs dementia screening tool), follow-up rate, and adjustment for preceding level of cognition, education, chronic diseases, APOE ϵ 4 allele, and other vascular risk factors.

In addition, we examined a possible dose-response association between physical activity and global cognition with scatter plots and fitted lines. The details of this analysis are described in the eMethods in [Supplement 1](#). *P* values were 2-sided, and statistical significance was set at *P* = .05. All analyses were performed with Stata statistical software version 18.0 (StataCorp). Data were analyzed from January to August 2023, with a final analysis in December 2023.

Results


The searches yielded 18 669 articles, of which 17 861 were excluded in title and abstract review phase and 703 were excluded in full-text review (**Figure 1**). This resulted in 104 included studies addressing physical activity and cognition, assessing a total of 341 471 participants. Of these, 45 had a binary outcome,¹⁸⁻⁶² 14 addressed follow-up global cognition,⁶³⁻⁷⁶ 25 addressed change or rate of change in global cognition,^{74,76-99} and 37 addressed physical activity and a specific cognitive domain of cognition.^{20,25,47,57,64,66,71,74,75,78,85,89,91,92,95,98-122} The quality assessments of all included studies are presented in eFigure 1 in [Supplement 1](#).

Physical Activity and Cognition With Binary Outcomes

Among 45 studies¹⁸⁻⁶² with a binary outcome, 1 study²⁹ was of high quality, 13 of moderate quality,^{18,22,24,28,32,37,38,41,43,46,47,55,59} and 31 of low quality.^{19-21,23,25-27,30,31,33-36,39,40,42,44,45,48-54,56-58,60-62} Physical activity was associated with a decreased risk of cognitive impairment or decline (RR, 0.89; 95% CI, 0.86-0.92; I^2 = 69.6%) (**Table 1** and **Table 2**; eFigure 2 in [Supplement 1](#)). Contour-enhanced funnel plot showed asymmetry (eFigure 3 in [Supplement 1](#)) due to reasons other than publication bias, like study quality. Trim-and-fill analysis correcting for this asymmetry showed a weaker association between physical activity and risk of cognitive impairment (RR, 0.97; 95% CI, 0.97-0.99) (Table 1). From the extensive set of moderators, only quality of physical activity measurement, sample size, follow-up rate, and length of follow-up significantly moderated the association between physical activity and cognition (Table 1; eTable 4 and eFigures 4-6 in [Supplement 1](#)). There was no significant association in follow-ups longer than 10 years (Table 1), while higher quality physical activity measurements and higher follow-up rate were associated with better cognition (eTable 4 in [Supplement 1](#)). The association was weaker in larger studies, but this result seemed largely driven by a few studies with very large sample sizes (>10 000 persons) (eTable 4 and eFigure 4 in [Supplement 1](#)).

Contour-enhanced funnel plot showed asymmetry (eFigure 3 in Supplement 1) due to reasons other than publication bias, like study quality. Trim-and-fill analysis correcting for this asymmetry showed a weaker association between physical activity and risk of cognitive impairment (RR, 0.97; 95% CI, 0.97-0.99) (Table 1). The amount of physical activity had a larger inverse association with cognitive impairment or decline until 5000 metabolic equivalent of task-minutes per week (ie, 16 hours of moderate to vigorous physical activity per week) (Figure 2A).

Figure 1. Study Selection Flowchart



Physical Activity and Follow-Up Global Cognition

Among 14 studies that assessed follow-up global cognition, there were no high-quality studies, 4 moderate-quality studies,^{63,65,67,71} and 10 low-quality studies.^{64,66,68-70,72-76} Pooled analysis showed a significant positive association between physical activity and follow-up global cognition (pooled $\beta = 0.025$; 95% CI, 0.004-0.047) (Table 1 and **Figure 3A**). The heterogeneity between the results from different studies was large ($I^2 = 75.8%$) (Table 1). None of the examined moderators significantly moderated the association between physical activity and cognition (Table 1; eTable 5 in [Supplement 1](#)). The funnel plot showed minimal asymmetry (eFigure 7 in [Supplement 1](#)), and the trim-and-fill analysis correcting for this asymmetry showed a very similar pooled standardized regression coefficient for physical activity and follow-up global cognition with narrower CIs ($\beta = 0.025$; 95% CI, 0.017-0.034) (Table 1). No dose-response association was seen (Figure 2B).

Table 1. Main Analysis of the Association of Physical Activity With Cognition

Analysis	Estimate (95% CI)	I^2 , %	No. of studies combined	Regression coefficient of the meta-regression	P value
Physical activity: binary outcomes					
Overall	0.89 (0.86 to 0.92) ^a	69.6	45	NA	NA
Meta trim-and-fill analysis	0.97 (0.97 to 0.99) ^a	NA	NA	NA	NA
Baseline age (continuous)	NA	NA	NA	0.0002	.93
Follow-up length					
Continuous	NA	NA	NA	0.008	.007
Categorical, y					
<5	0.81 (0.76 to 0.86) ^a	74.0	22	NA	NA
5-9	0.90 (0.86 to 0.95) ^a	58.7	13	NA	NA
10-14	1.06 (0.95 to 1.17) ^a	16.2	4	NA	NA
≥15	0.93 (0.82 to 1.05) ^a	77.2	6	NA	NA
Study quality (continuous) ^b	NA	NA	NA	0.03	.41
Adjusting for preceding level of cognition					
No	0.89 (0.85 to 0.93) ^a	69.2	31	NA	.23
Yes	0.85 (0.79 to 0.91) ^a	71.6	14	NA	
Physical activity and follow-up global cognition					
Overall	0.025 (0.004 to 0.047) ^c	75.8	14	NA	NA
Meta trim- and fill analysis	0.025 (0.017 to 0.034) ^c	NA	NA	NA	NA
Age at baseline (continuous)	NA	NA	NA	-0.0006	.94
Follow-up length (continuous)	NA	NA	NA	0.0008	.69
Study quality (continuous) ^b	NA	NA	NA	-0.03	.35
Adjustment for preceding level of cognition					
No	0.005 (-0.033 to 0.043) ^c	76.6	8	NA	.14
Yes	0.041 (0.012 to 0.071) ^c	77.2	6	NA	
Physical activity and change in global cognition					
Overall	0.016 (0.002 to 0.030) ^c	67.4	25	NA	NA
Meta trim- and fill analysis	0.013 (0.008 to 0.018) ^c	NA	NA	NA	NA
Age at baseline (continuous)	NA	NA	NA	-0.0005	.75
Follow-up length (continuous)	NA	NA	NA	-0.0009	.69
Study quality (continuous) ^b	NA	NA	NA	0.003	.90
Adjustment for preceding level of cognition					
No	0.023 (-0.010 to 0.057) ^c	79.0	14	NA	.64
Yes	0.015 (0.010 to 0.020) ^c	0	11	NA	

Abbreviation: NA, not applicable.

^a Expressed as pooled risk ratios.

^b Study quality was assessed with a quality assessment tool we developed (range, 1-3; higher score denotes worse quality). Further details are provided in the eMethods in [Supplement 1](#).

^c Expressed as pooled standardized regression coefficients.

Table 2. Studies Assessing Physical Activity and Cognition as Binary Outcomes


Study	Risk ratio 95% (CI)	Weight, %
Beauchet et al, ⁵² 2020	1.21 (0.65-2.19)	0.29
Brunner et al, ¹⁸ 2017		
Moderately active	0.99 (0.89-1.11)	3.14
Sufficiently active	0.89 (0.79-1.01)	2.88
Chen et al, ¹⁹ 2016	1.08 (0.93-1.27)	2.33
Clark et al, ²⁰ 2016		
Second quintile	0.83 (0.55-1.24)	0.60
Third quintile	1.27 (0.90-1.77)	0.81
Fourth quintile	0.95 (0.65-1.35)	0.72
Fifth quintile	0.77 (0.50-1.15)	0.58
de Frias et al, ²¹ 2014	1.08 (0.93-1.26)	2.42
de Looze et al, ⁵¹ 2022	0.93 (0.85-1.02)	3.50
Dupré et al, ⁵⁵ 2020		
Second category	1.01 (0.81-1.29)	1.43
Third category	0.75 (0.50-1.15)	0.57
Elwood et al, ²² 2013	0.79 (0.64-0.96)	1.73
Etgen et al, ²³ 2010		
High activity	0.54 (0.32-0.87)	0.42
Moderate activity	0.59 (0.36-0.95)	0.44
Fassier et al, ⁵⁷ 2022		
Tertile 2	0.94 (0.75-1.19)	1.42
Tertile 3	0.81 (0.63-1.02)	1.37
Gao et al, ²⁴ 2017	1.19 (1.07-1.31)	3.26
He et al, ⁵⁹ 2021	0.78 (0.64-0.94)	1.89
Hildreth et al, ²⁵ 2014	0.75 (0.55-1.00)	1.00
Ho et al, ²⁶ 2001	0.69 (0.55-0.88)	1.42
Hughes et al, ²⁷ 2015	0.42 (0.25-0.72)	0.37
Infurna et al, ²⁸ 2016	1.06 (1.02-1.11)	4.28
Iso-Markku et al, ²⁹ 2016	0.78 (0.66-0.94)	2.00
Iwasa et al, ³⁰ 2012	0.97 (0.76-1.24)	1.31
Kim et al, ³¹ 2011	0.81 (0.70-0.95)	2.38
Krell-Roesch et al, ⁶² 2021	0.86 (0.74-1.00)	2.41
Laurin et al, ³² 2001		
Men with high activity	0.70 (0.41-1.18)	0.37
Men with low activity	0.67 (0.32-1.33)	0.21
Men with moderate activity	0.85 (0.55-1.30)	0.54
Women with high activity	0.49 (0.27-0.91)	0.29
Women with low activity	0.71 (0.43-1.14)	0.44
Women with moderate activity	0.57 (0.38-0.83)	0.64
Lee et al, ³³ 2013		
Quartile 2	0.76 (0.53-1.10)	0.72
Quartile 3	0.73 (0.50-1.06)	0.69
Quartile 4	0.62 (0.42-0.91)	0.66
Leung et al, ³⁴ 2011	0.97 (0.95-0.99)	4.52
Lipnicki et al, ³⁵ 2017	0.96 (0.89-1.04)	3.66
Lytle et al, ³⁶ 2004		
High exercise	0.62 (0.44-0.88)	0.76
Low exercise	0.83 (0.66-1.05)	1.42
McGarrigle et al, ⁶⁰ 2022	1.00 (0.98-1.02)	4.55
Middleton et al, ³⁷ 2011		
Highest tertile of device-measured activity	0.10 (0.01-0.80)	0.02
Lowest tertile of device-measured activity	0.30 (0.07-1.21)	0.05

(continued)

Table 2. Studies Assessing Physical Activity and Cognition as Binary Outcomes (continued)

Study	Risk ratio 95% (CI)	Weight, %
Min et al, ⁵⁶ 2018	0.67 (0.47-0.96)	0.73
Newman et al, ³⁸ 2009		
>1890 kcal	1.42 (0.56-3.18)	0.14
1-270 kcal	1.07 (0.39-2.63)	0.12
271-810 kcal	1.28 (0.49-2.93)	0.14
811-890 kcal	0.70 (0.26-1.80)	0.12
Niti et al, ³⁹ 2008	0.88 (0.78-1.01)	2.69
Pignatti et al, ⁴⁰ 2002	0.27 (0.09-0.83)	0.09
Pitrou et al, ⁶¹ 2022	0.94 (0.59-1.51)	0.46
Ramoo et al, ⁵³ 2022		
Quartile 2	0.69 (0.47-1.02)	0.65
Quartile 3	0.99 (0.68-1.50)	0.63
Quartile 4	0.53 (0.38-0.76)	0.78
Shih et al, ⁴¹ 2017	0.71 (0.50-0.99)	0.81
Stewart et al, ⁴² 2003	0.84 (0.58-1.19)	0.74
Strozza et al, ⁵⁴ 2020		
Light physical activity	0.87 (0.61-1.19)	0.84
Heavy physical activity	0.69 (0.42-1.11)	0.44
Sumic et al, ⁴³ 2007		
Men	0.94 (0.34-1.64)	0.17
Women	0.17 (0.04-0.52)	0.07
Thompson et al, ⁵⁸ 2022	1.37 (0.70-2.71)	0.23
Verdelho et al, ⁴⁴ 2012	0.72 (0.57-0.92)	1.36
Verghese et al, ⁴⁸ 2006	0.97 (0.94-1.01)	4.37
Verghese et al, ⁴⁹ 2009	0.99 (0.96-1.03)	4.44
Wang et al, ⁵⁰ 2006	0.96 (0.84-1.09)	2.68
Woodard et al, ⁴⁵ 2012	0.99 (0.38-2.55)	0.12
Yaffe et al, ⁴⁶ 2001		
Highest quartile	0.86 (0.78-0.95)	3.26
Second quartile	0.95 (0.86-1.04)	3.35
Third quartile	0.88 (0.80-0.98)	3.26
Zhu et al, ⁴⁷ 2017		
Quartile 2	0.65 (0.49-0.85)	1.16
Quartile 3	0.54 (0.39-0.75)	0.86
Quartile 4	0.58 (0.40-0.82)	0.74
Overall ($I^2 = 69.6\%$; $P < .001$)	0.89 (0.86-0.92)	100

Figure 2. Assessment of Dose-Response Association of Physical Activity With Cognition





MET indicates metabolic equivalent task; circles, individual studies; size of circles, weight of studies; blue line, estimate; outline, 95% CI.

Physical Activity and Change in Global Cognition

Among 25 studies that assessed change in global cognition, there were no high-quality studies, 5 moderate-quality studies,^{83,87-89,99} and 20 low-quality studies.^{74,76-82,84-86,90-98} Pooled analysis of all studies addressing physical activity and change in global cognition showed a significant positive association (pooled $\beta = 0.016$; 95% CI, 0.002-0.030) (Table 1 and Figure 3B). The heterogeneity among the results from different studies was large ($I^2 = 67.4%$) (Table 1). The funnel plot showed at

Figure 3. Assessment of Association of Physical Activity With Cognition Using Continuous Outcomes



most minimal asymmetry (eFigure 8 in Supplement 1). Trim-and-fill analysis showed a similar pooled standardized regression coefficient as the main analysis with a narrower CI ($\beta = 0.013$; 95% CI, 0.008-0.018) (Table 1). None of the moderators examined significantly moderated the association between physical activity and change in cognition (Table 1; eTable 6 in Supplement 1). There was no dose-response association (Figure 2C).

Physical Activity and Specific Cognitive Domains

There were no significant associations between physical activity and any of the specific cognitive domains in studies with binary outcomes (eTable 7 in Supplement 1). In the studies of physical activity and follow-up cognition, physical activity was associated with minimally better executive function (standardized regression coefficient, 0.05, 95% CI 0.01 to 0.09), episodic memory (standardized regression coefficient, 0.03; 95% CI, 0.02-0.04), and verbal fluency (standardized regression coefficient, 0.05; 95% CI, 0.03-0.08). There was only 1 study on physical activity and verbal ability.¹¹⁷ In the studies of physical activity and change in cognition, physical activity was significantly and minimally associated with episodic memory and verbal fluency. Adjustment for the preceding level of cognition did not significantly modify the associations between physical activity and specific cognitive domains. The only exception was verbal fluency, for which the association was significant only when adjusting for preceding level of cognition in studies of change in cognition and significant, in binary studies, only when not adjusting for preceding level of cognition (eTable 8 in Supplement 1).

Discussion

This systematic review and meta-analysis found only minimal associations between physical activity and cognition. These very small estimates are more in line with a recent umbrella review of randomized clinical trials showing very small effects sizes between physical activity and cognition⁶ than with earlier meta-analyses of observational studies on physical activity and cognition, which showed moderate associations.⁹⁻¹² For comparison, a 2018 study found that each 1-year increase in age was associated with a 0.037-SD decrease in global cognition.¹²³ The identified weak association between physical activity and cognition was persistent, regardless of the preceding level of cognition or cohort age, which is in line with our previous meta-analysis of physical activity and dementia.³ Although the pooled standardized magnitudes of association were very small,^{15,124} they are significant in a population health perspective for the potential to postpone the multifactorial diseases causing dementia.

Our results indicate a dose-response association between physical activity and cognition among studies with binary cognition outcomes. This association was moderated by follow-up length, follow-up rate, physical activity measurement type, and physical activity measurement quality, but funnel plots detected possible bias in this set of studies. On the contrary, while not revealing possible bias, our meta-analysis of studies with continuous outcomes found neither a dose-response association nor any significant moderation. This contradicts our earlier meta-analysis of physical activity and dementia that found a dose-response association³ and also other meta-analyses of physical activity and dementia.^{125,126}

Beyond this, study quality did not significantly moderate any of the associations. For a study to be rated as high quality in our assessment, it needed to include a follow-up of more than 10 years,⁸⁵ a measurement of baseline cognition, and very high participation and follow-up rates—all factors that are necessary to accurately examine longitudinal associations between physical activity and cognition. Further high-quality research is needed.

The analysis of physical activity and specific cognitive domains revealed similar weak associations for episodic memory and verbal fluency as the main analysis (pooled standardized regression coefficients between 0.02 and 0.05). The results for executive function were mixed between analyses of follow-up and change. The CIs for the associations were wider for verbal ability,

working memory, processing speed, and visuospatial ability, but the data for these analyses were scarcer and explain at least partly the wider CIs.

Our meta-analysis has many strengths. To our knowledge, no other meta-analysis of observational studies has examined continuous outcomes or specific cognitive domains before. In addition, we combined the data of 6 times more individuals (>300 000) than any previous meta-analysis on the topic⁹⁻¹² and examined more possible moderators.

Limitations

This meta-analysis has limitations. High-quality studies were rare and data examining midlife physical activity and midlife cognition were scarce. Thus, our meta-analysis mainly provides evidence on how physical activity was associated with cognitive aging, not cognition in midlife. In retrospect, excluding studies without valid measures of cognition at baseline perhaps limited power to detect adjustment for baseline cognition as a moderator. We also did not assess whether studies accounted for practice effects when measuring cognition. Low study quality and imprecise study-level measures of physical activity and cognition limit the robustness of our dose-response analyses.

Conclusions

This systematic review and meta-analysis found that the association between physical activity and cognitive decline was very small, with no evident dose-response association. With that said, even weak associations can be clinically significant from a population health perspective when physical activity continues over decades. It should also be noted that very few high-quality studies were included. Further high-quality cohort studies with follow-ups longer than 10 to 20 years, fine-grained measures of physical activity and cognition at baseline, and high participation and follow-up rates are needed to solidify the evidence base in this area.

ARTICLE INFORMATION

Accepted for Publication: December 11, 2023.

Published: February 1, 2024. doi:10.1001/jamanetworkopen.2023.54285

Open Access: This is an open access article distributed under the terms of the [CC-BY License](#). © 2024 Iso-Markku P et al. *JAMA Network Open*.

Corresponding Author: Paula Iso-Markku, MD, PhD, Institute for Molecular Medicine Finland, HiLIFE, University of Helsinki, PO Box 20 (Tukholmankatu 8), 00014 University of Helsinki, Finland (paula.iso-markku@helsinki.fi).

Author Affiliations: Institute for Molecular Medicine Finland, HiLIFE, University of Helsinki, Helsinki, Finland (Iso-Markku, Aaltonen, Vuoksima); Helsinki University Hospital Diagnostic Center, Clinical Physiology and Nuclear Medicine, University of Helsinki and Helsinki University Hospital, Helsinki, Finland (Iso-Markku, Halme); Faculty of Sport and Health Sciences, University of Jyväskylä, Jyväskylä, Finland (Kujala, Phipps, Knittle, Waller).

Author Contributions: Dr Iso-Markku had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Concept and design: Iso-Markku, Kujala, Vuoksima, Waller.

Acquisition, analysis, or interpretation of data: All authors.

Drafting of the manuscript: Iso-Markku, Knittle.

Critical review of the manuscript for important intellectual content: All authors.

Statistical analysis: Iso-Markku, Kujala, Knittle.

Obtained funding: Iso-Markku.

Administrative, technical, or material support: Kujala, Phipps, Knittle, Vuoksima, Waller.

Supervision: Vuoksima, Waller.

Conflict of Interest Disclosures: None reported.

Funding/Support: This work was funded by Biomedicum Helsinki Foundation (Dr Iso-Marku), Orion Research Foundation (Dr Iso-Marku), Juho Vainio Foundation (Dr Iso-Marku), HUS Diagnostic Center research funding (Dr Iso-Marku), and the Academy of Finland (grant No. 314639 and 320109; Dr Vuoksimaa). Open access was funded by Helsinki University Library.

Role of the Funder/Sponsor: The funders had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

Data Sharing Statement: See Supplement 3.

REFERENCES

1. Erickson KI, Hillman C, Stillman CM, et al; for 2018 Physical Activity Guidelines Advisory Committee. Physical activity, cognition, and brain outcomes: a review of the 2018 physical activity guidelines. *Med Sci Sports Exerc*. 2019;51(6):1242-1251. doi:10.1249/MSS.0000000000001936
2. Livingston G, Huntley J, Sommerlad A, et al. Dementia prevention, intervention, and care: 2020 report of the Lancet Commission. *Lancet*. 2020;396(10248):413-446. doi:10.1016/S0140-6736(20)30367-6
3. Iso-Markku P, Kujala UM, Knittle K, Polet J, Vuoksimaa E, Waller K. Physical activity as a protective factor for dementia and Alzheimer's disease: systematic review, meta-analysis and quality assessment of cohort and case-control studies. *Br J Sports Med*. 2022;56(12):701-709. doi:10.1136/bjsports-2021-104981
4. Turner DT, Hu MX, Generaal E, et al. Physical exercise interventions targeting cognitive functioning and the cognitive domains in nondementia samples: a systematic review of meta-analyses. *J Geriatr Psychiatry Neurol*. 2021;34(2):91-101. doi:10.1177/0891988720915523
5. Engeroff T, Ingmann T, Banzer W. Physical activity throughout the adult life span and domain-specific cognitive function in old age: a systematic review of cross-sectional and longitudinal data. *Sports Med*. 2018;48(6):1405-1436. doi:10.1007/s40279-018-0920-6
6. Ciria LF, Román-Caballero R, Vadillo MA, et al. An umbrella review of randomized control trials on the effects of physical exercise on cognition. *Nat Hum Behav*. 2023;7(6):928-941. doi:10.1038/s41562-023-01554-4
7. Sink KM, Espeland MA, Castro CM, et al; LIFE Study Investigators. Effect of a 24-month physical activity intervention vs health education on cognitive outcomes in sedentary older adults: the LIFE randomized trial. *JAMA*. 2015;314(8):781-790. doi:10.1001/jama.2015.9617
8. Komulainen P, Tuomilehto J, Savonen K, et al. Exercise, diet, and cognition in a 4-year randomized controlled trial: Dose-Responses to Exercise Training (DR's EXTRA). *Am J Clin Nutr*. 2021;113(6):1428-1439. doi:10.1093/ajcn/nqab018
9. Sofi F, Valecchi D, Bacci D, et al. Physical activity and risk of cognitive decline: a meta-analysis of prospective studies. *J Intern Med*. 2011;269(1):107-117. doi:10.1111/j.1365-2796.2010.02281.x
10. Morgan GS, Gallacher J, Bayer A, Fish M, Ebrahim S, Ben-Shlomo Y. Physical activity in middle-age and dementia in later life: findings from a prospective cohort of men in Caerphilly, South Wales and a meta-analysis. *J Alzheimers Dis*. 2012;31(3):569-580. doi:10.3233/JAD-2012-112171
11. Blondell SJ, Hammersley-Mather R, Veerman JL. Does physical activity prevent cognitive decline and dementia: a systematic review and meta-analysis of longitudinal studies. *BMC Public Health*. 2014;14:510. doi:10.1186/1471-2458-14-510
12. Guure CB, Ibrahim NA, Adam MB, Said SM. Impact of physical activity on cognitive decline, dementia, and its subtypes: meta-analysis of prospective studies. *Biomed Res Int*. 2017;2017:9016924. doi:10.1155/2017/9016924
13. Moher D, Liberati A, Tetzlaff J, Altman DG; PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med*. 2009;6(7):e1000097. doi:10.1371/journal.pmed.1000097
14. Jack CR Jr, Knopman DS, Jagust WJ, et al. Tracking pathophysiological processes in Alzheimer's disease: an updated hypothetical model of dynamic biomarkers. *Lancet Neurol*. 2013;12(2):207-216. doi:10.1016/S1474-4422(12)70291-0
15. Nieminen P. Application of standardized regression coefficient in meta-analysis. *BioMedInformatics*. 2022;2(3):434-458. doi:10.3390/biomedinformatics2030028
16. Feingold A. A regression framework for effect size assessments in longitudinal modeling of group differences. *Rev Gen Psychol*. 2013;17(1):111-121. doi:10.1037/a0030048
17. Lipsey MWD. *Practical Meta-Analysis*. Sage Publications; 2001.
18. Brunner EJ, Welch CA, Shipley MJ, Ahmadi-Abhari S, Singh-Manoux A, Kivimäki M. Midlife risk factors for impaired physical and cognitive functioning at older ages: a cohort study. *J Gerontol A Biol Sci Med Sci*. 2017;72(2):237-242. doi:10.1093/gerona/glw092

19. Chen TY, Chang HY. Developmental patterns of cognitive function and associated factors among the elderly in Taiwan. *Sci Rep*. 2016;6:33486. doi:10.1038/srep33486
20. Clark S, Parisi J, Kuo J, Carlson MC. Physical activity is associated with reduced risk of executive function impairment in older women. *J Aging Health*. 2016;28(4):726-739. doi:10.1177/0898264315609908
21. de Frias CM, Dixon RA. Lifestyle engagement affects cognitive status differences and trajectories on executive functions in older adults. *Arch Clin Neuropsychol*. 2014;29(1):16-25. doi:10.1093/arclin/act089
22. Elwood P, Galante J, Pickering J, et al. Healthy lifestyles reduce the incidence of chronic diseases and dementia: evidence from the Caerphilly cohort study. *PLoS One*. 2013;8(12):e81877. doi:10.1371/journal.pone.0081877
23. Etgen T, Sander D, Huntgeburth U, Poppert H, Förstl H, Bickel H. Physical activity and incident cognitive impairment in elderly persons: the INVADE study. *Arch Intern Med*. 2010;170(2):186-193. doi:10.1001/archinternmed.2009.498
24. Gao M, Kuang W, Qiu P, Wang H, Lv X, Yang M. The time trends of cognitive impairment incidence among older Chinese people in the community: based on the CLHLS cohorts from 1998 to 2014. *Age Ageing*. 2017;46(5):787-793. doi:10.1093/ageing/afx038
25. Hildreth KL, Grigsby J, Bryant LL, Wolfe P, Baxter J. Cognitive decline and cardiometabolic risk among Hispanic and non-Hispanic White adults in the San Luis Valley Health and Aging Study. *J Behav Med*. 2014;37(2):332-342. doi:10.1007/s10865-013-9491-z
26. Ho SC, Woo J, Sham A, Chan SG, Yu AL. A 3-year follow-up study of social, lifestyle and health predictors of cognitive impairment in a Chinese older cohort. *Int J Epidemiol*. 2001;30(6):1389-1396. doi:10.1093/ije/30.6.1389
27. Hughes TF, Becker JT, Lee CW, Chang CC, Ganguli M. Independent and combined effects of cognitive and physical activity on incident MCI. *Alzheimers Dement*. 2015;11(11):1377-1384. doi:10.1016/j.jalz.2014.11.007
28. Infurna FJ, Okun MA, Grimm KJ. Volunteering is associated with lower risk of cognitive impairment. *J Am Geriatr Soc*. 2016;64(11):2263-2269. doi:10.1111/jgs.14398
29. Iso-Markku P, Waller K, Vuoksima E, et al. Midlife physical activity and cognition later in life: a prospective twin study. *J Alzheimers Dis*. 2016;54(4):1303-1317. doi:10.3233/JAD-160377
30. Iwasa H, Yoshida Y, Kai I, Suzuki T, Kim H, Yoshida H. Leisure activities and cognitive function in elderly community-dwelling individuals in Japan: a 5-year prospective cohort study. *J Psychosom Res*. 2012;72(2):159-164. doi:10.1016/j.jpsychores.2011.10.002
31. Kim JM, Stewart R, Bae KY, et al. Role of BDNF val66met polymorphism on the association between physical activity and incident dementia. *Neurobiol Aging*. 2011;32(3):551.e5-551.e12. doi:10.1016/j.neurobiolaging.2010.01.018
32. Laurin D, Verreault R, Lindsay J, MacPherson K, Rockwood K. Physical activity and risk of cognitive impairment and dementia in elderly persons. *Arch Neurol*. 2001;58(3):498-504. doi:10.1001/archneur.58.3.498
33. Lee S, Yuki A, Nishida Y, et al. Research relationship between light-intensity physical activity and cognitive function in a community-dwelling elderly population-an 8-year longitudinal study. *J Am Geriatr Soc*. 2013;61(3):452-453. doi:10.1111/jgs.12119
34. Leung GT, Fung AW, Tam CW, et al. Examining the association between late-life leisure activity participation and global cognitive decline in community-dwelling elderly Chinese in Hong Kong. *Int J Geriatr Psychiatry*. 2011;26(1):39-47. doi:10.1002/gps.2478
35. Lipnicki DM, Crawford J, Kochan NA, et al; Sydney Memory and Ageing Study Team. Risk factors for mild cognitive impairment, dementia and mortality: the Sydney Memory and Ageing Study. *J Am Med Dir Assoc*. 2017;18(5):388-395. doi:10.1016/j.jamda.2016.10.014
36. Lytle ME, Vander Bilt J, Pandav RS, Dodge HH, Ganguli M. Exercise level and cognitive decline: the MOVIES project. *Alzheimer Dis Assoc Disord*. 2004;18(2):57-64. doi:10.1097/01.wad.0000126614.87955.79
37. Middleton LE, Manini TM, Simonsick EM, et al. Activity energy expenditure and incident cognitive impairment in older adults. *Arch Intern Med*. 2011;171(14):1251-1257. doi:10.1001/archinternmed.2011.277
38. Newman AB, Arnold AM, Sachs MC, et al. Long-term function in an older cohort—the cardiovascular health study all stars study. *J Am Geriatr Soc*. 2009;57(3):432-440. doi:10.1111/j.1532-5415.2008.02152.x
39. Niti M, Yap KB, Kua EH, Tan CH, Ng TP. Physical, social and productive leisure activities, cognitive decline and interaction with APOE-epsilon 4 genotype in Chinese older adults. *Int Psychogeriatr*. 2008;20(2):237-251. doi:10.1017/S1041610207006655
40. Pignatti F, Rozzini R, Trabucchi M. Physical activity and cognitive decline in elderly persons. *Arch Intern Med*. 2002;162(3):361-362. doi:10.1001/archinte.162.3.361

41. Shih IF, Paul K, Haan M, Yu Y, Ritz B. Physical activity modifies the influence of apolipoprotein E ϵ 4 allele and type 2 diabetes on dementia and cognitive impairment among older Mexican Americans. *Alzheimers Dement*. 2018;14(1):1-9. doi:10.1016/j.jalz.2017.05.005
42. Stewart R, Prince M, Mann A. Age, vascular risk, and cognitive decline in an older, British, African-Caribbean population. *J Am Geriatr Soc*. 2003;51(11):1547-1553. doi:10.1046/j.1532-5415.2003.51504.x
43. Sumic A, Michael YL, Carlson NE, Howieson DB, Kaye JA. Physical activity and the risk of dementia in oldest old. *J Aging Health*. 2007;19(2):242-259. doi:10.1177/0898264307299299
44. Verdelho A, Madureira S, Ferro JM, et al; LADIS Study. Physical activity prevents progression for cognitive impairment and vascular dementia: results from the LADIS (Leukoaraiosis and Disability) study. *Stroke*. 2012;43(12):3331-3335. doi:10.1161/STROKEAHA.112.661793
45. Woodard JL, Sugarman MA, Nielson KA, et al. Lifestyle and genetic contributions to cognitive decline and hippocampal structure and function in healthy aging. *Curr Alzheimer Res*. 2012;9(4):436-446. doi:10.2174/156720512800492477
46. Yaffe K, Barnes D, Nevitt M, Lui LY, Covinsky K. A prospective study of physical activity and cognitive decline in elderly women: women who walk. *Arch Intern Med*. 2001;161(14):1703-1708. doi:10.1001/archinte.161.14.1703
47. Zhu W, Wadley VG, Howard VJ, Hutto B, Blair SN, Hooker SP. Objectively measured physical activity and cognitive function in older adults. *Med Sci Sports Exerc*. 2017;49(1):47-53. doi:10.1249/MSS.0000000000001079
48. Verghese J, LeValley A, Derby C, et al. Leisure activities and the risk of amnesic mild cognitive impairment in the elderly. *Neurology*. 2006;66(6):821-827. doi:10.1212/01.wnl.0000202520.68987.48
49. Verghese J, Wang C, Katz MJ, Sanders A, Lipton RB. Leisure activities and risk of vascular cognitive impairment in older adults. *J Geriatr Psychiatry Neurol*. 2009;22(2):110-118. doi:10.1177/0891988709332938
50. Wang JY, Zhou DH, Li J, et al. Leisure activity and risk of cognitive impairment: the Chongqing aging study. *Neurology*. 2006;66(6):911-913. doi:10.1212/01.wnl.0000192165.99963.2a
51. De Looze C, Williamson W, Demnitz N, O'Connor D, Hernández B, Kenny RA. Physical function, an adjunct to brain health score for phenotyping cognitive function trajectories in older age: findings from the Irish Longitudinal Study on Ageing (TILDA). *J Gerontol A Biol Sci Med Sci*. 2022;77(8):1593-1602. doi:10.1093/gerona/glab024
52. Beauchet O, Sekhon H, Launay CP, Gaudreau P, Morais JA, Allali G. Relationship between motoric cognitive risk syndrome, cardiovascular risk factors and diseases, and incident cognitive impairment: results from the "NuAge" study. *Maturitas*. 2020;138:51-57. doi:10.1016/j.maturitas.2020.05.007
53. Ramoo K, Hairi NN, Yahya A, et al. Longitudinal Association between Sarcopenia and Cognitive Impairment among Older Adults in Rural Malaysia. *Int J Environ Res Public Health*. 2022;19(8):4723. doi:10.3390/ijerph19084723
54. Strozza C, Zarulli V, Egidi V. Understanding health deterioration and the dynamic relationship between physical ability and cognition among a cohort of Danish nonagenarians. *J Aging Res*. 2020;2020:4704305. doi:10.1155/2020/4704305
55. Dupré C, Hupin D, Goethals L, et al. Domestic activities associated with a decreased risk of cognitive disorders: results of the "Fréle" cohort. *Front Public Health*. 2020;8:602238. doi:10.3389/fpubh.2020.602238
56. Min JW. A longitudinal study of cognitive trajectories and its factors for Koreans aged 60 and over: a latent growth mixture model. *Int J Geriatr Psychiatry*. 2018;33(5):755-762. doi:10.1002/gps.4855
57. Fassier P, Kang JH, Lee IM, Grodstein F, Vercambre MN. Vigorous physical activity and cognitive trajectory later in life: prospective association and interaction by apolipoprotein E ϵ 4 in the Nurses' Health Study. *J Gerontol A Biol Sci Med Sci*. 2022;77(4):817-825. doi:10.1093/gerona/glab169
58. Thompson F, Russell SG, Harriss LR, et al. Using health check data to understand risks for dementia and cognitive impairment among Torres Strait Islander and Aboriginal peoples in northern Queensland—a data linkage study. *Front Public Health*. 2022;10:782373. doi:10.3389/fpubh.2022.782373
59. He F, Lin J, Li F, et al. Physical work and exercise reduce the risk of cognitive impairment in older adults: a population-based longitudinal study. *Curr Alzheimer Res*. 2021;18(8):638-645. doi:10.2174/156720501866621118100451
60. McGarrigle CA, Ward M, Kenny RA. Negative aging perceptions and cognitive and functional decline: are you as old as you feel? *J Am Geriatr Soc*. 2022;70(3):777-788. doi:10.1111/jgs.17561
61. Pitrou I, Vasiliadis HM, Hudon C. Body mass index and cognitive decline among community-living older adults: the modifying effect of physical activity. *Eur Rev Aging Phys Act*. 2022;19(1):3. doi:10.1186/s11556-022-00284-2

62. Krell-Roesch J, Syrjanen JA, Bezold J, et al. Lack of physical activity, neuropsychiatric symptoms and the risk of incident mild cognitive impairment in older community-dwelling individuals: a prospective cohort study. *Ger J Exerc Sport Res*. 2021;51(4):487-494. doi:10.1007/s12662-021-00732-8
63. Bowling A, Pikhartova J, Dodgeon B. Is mid-life social participation associated with cognitive function at age 50: results from the British National Child Development Study (NCDS). *BMC Psychol*. 2016;4(1):58. doi:10.1186/s40359-016-0164-x
64. Dregan A, Gulliford MC. Leisure-time physical activity over the life course and cognitive functioning in late mid-adult years: a cohort-based investigation. *Psychol Med*. 2013;43(11):2447-2458. doi:10.1017/S0033291713000305
65. Gross AL, Lu H, Meoni L, Gallo JJ, Schrack JA, Sharrett AR. Physical activity in midlife is not associated with cognitive health in later life among cognitively normal older adults. *J Alzheimers Dis*. 2017;59(4):1349-1358. doi:10.3233/JAD-170290
66. Wolinsky FD, Bentler SE, Hockenberry J, et al. A prospective cohort study of long-term cognitive changes in older Medicare beneficiaries. *BMC Public Health*. 2011;11:710. doi:10.1186/1471-2458-11-710
67. Carty CL, Noonan C, Muller C, et al. Mid-life physical activity and late-life cognitive performance among American Indians. *Neuroepidemiology*. 2022;56(2):119-126. doi:10.1159/000521791
68. Zhao X, Jin L, Sun SB. The bidirectional association between physical and cognitive function among Chinese older adults: a mediation analysis. *Int J Aging Hum Dev*. 2021;92(2):240-263. doi:10.1177/0091415020940214
69. Hu M, Liu S, Shen Y, et al. Physical activity trajectories and cognitive function: a national cohort study. *Ment Health Phys Act*. 2022;23:100482. doi:10.1016/j.mhpa.2022.100482
70. Visser M, Wijnhoven HAH, Comijs HC, Thomése FGCF, Twisk JWR, Deeg DJH. A healthy lifestyle in old age and prospective change in four domains of functioning. *J Aging Health*. 2019;31(7):1297-1314. doi:10.1177/0898264318774430
71. Gafni T, Gabriel KP, Shuval K, Yaffe K, Sidney S, Weinstein G. Physical activity trajectories, autonomic balance and cognitive function: the Coronary Artery Risk Development in Young Adults (CARDIA) study. *Prev Med*. 2022;164:107291. doi:10.1016/j.ypmed.2022.107291
72. Kim JH, Sumerlin TS, Goggins WB, et al. Does low subjective social status predict cognitive decline in Chinese older adults: a 4-year longitudinal study from Hong Kong. *Am J Geriatr Psychiatry*. 2021;29(11):1140-1151. doi:10.1016/j.jagp.2021.01.014
73. Pruzin JJ, Klein H, Rabin JS, et al. Physical activity is associated with increased resting-state functional connectivity in networks predictive of cognitive decline in clinically unimpaired older adults. *Alzheimers Dement (Amst)*. 2022;14(1):e12319. doi:10.1002/dad2.12319
74. Younan D, Wang X, Millstein J, et al. Air quality improvement and cognitive decline in community-dwelling older women in the United States: A longitudinal cohort study. *PLoS Med*. 2022;19(2):e1003893. doi:10.1371/journal.pmed.1003893
75. Wang S, Luo X, Barnes D, Sano M, Yaffe K. Physical activity and risk of cognitive impairment among oldest-old women. *Am J Geriatr Psychiatry*. 2014;22(11):1149-1157. doi:10.1016/j.jagp.2013.03.002
76. Yamamoto M, Wada-Isoe K, Yamashita F, et al. Association between exercise habits and subcortical gray matter volumes in healthy elderly people: a population-based study in Japan. *eNeurologicalSci*. 2017;7:1-6. doi:10.1016/j.ensci.2017.03.002
77. Albert MS, Jones K, Savage CR, et al. Predictors of cognitive change in older persons: MacArthur studies of successful aging. *Psychol Aging*. 1995;10(4):578-589. doi:10.1037/0882-7974.10.4.578
78. Bosma H, van Boxtel MP, Ponds RW, et al. Engaged lifestyle and cognitive function in middle and old-aged, non-demented persons: a reciprocal association? *Z Gerontol Geriatr*. 2002;35(6):575-581. doi:10.1007/s00391-002-0080-y
79. Brewster PW, Melrose RJ, Marquine MJ, et al. Life experience and demographic influences on cognitive function in older adults. *Neuropsychology*. 2014;28(6):846-858. doi:10.1037/neu0000098
80. Gow AJ, Mortensen EL, Avlund K. Activity participation and cognitive aging from age 50 to 80 in the Glostrup 1914 cohort. *J Am Geriatr Soc*. 2012;60(10):1831-1838. doi:10.1111/j.1532-5415.2012.04168.x
81. Gow AJ, Pattie A, Deary IJ. Lifecourse activity participation from early, mid, and later adulthood as determinants of cognitive aging: the Lothian Birth Cohort 1921. *J Gerontol B Psychol Sci Soc Sci*. 2017;72(1):25-37. doi:10.1093/geronb/gbw124
82. Lee Y, Kim J, Back JH. The influence of multiple lifestyle behaviors on cognitive function in older persons living in the community. *Prev Med*. 2009;48(1):86-90. doi:10.1016/j.ypmed.2008.10.021

83. Ottenbacher AJ, Snih SA, Bindawas SM, et al. Role of physical activity in reducing cognitive decline in older Mexican-American adults. *J Am Geriatr Soc*. 2014;62(9):1786-1791. doi:10.1111/jgs.12978
84. Ravona-Springer R, Moshier E, Schmeidler J, et al. Changes in glycemic control are associated with changes in cognition in non-diabetic elderly. *J Alzheimers Dis*. 2012;30(2):299-309. doi:10.3233/JAD-2012-120106
85. Sabia S, Dugravot A, Dartigues JF, et al. Physical activity, cognitive decline, and risk of dementia: 28 year follow-up of Whitehall II cohort study. *BMJ*. 2017;357:j2709. doi:10.1136/bmj.j2709
86. Steves CJ, Mehta MM, Jackson SH, Spector TD. Kicking back cognitive ageing: leg power predicts cognitive ageing after ten years in older female twins. *Gerontology*. 2016;62(2):138-149. doi:10.1159/000441029
87. Sturman MT, Morris MC, Mendes de Leon CF, Bienias JL, Wilson RS, Evans DA. Physical activity, cognitive activity, and cognitive decline in a biracial community population. *Arch Neurol*. 2005;62(11):1750-1754. doi:10.1001/archneur.62.11.1750
88. van Gelder BM, Tijhuis MA, Kalmijn S, Giampaoli S, Nissinen A, Kromhout D. Physical activity in relation to cognitive decline in elderly men: the FINE Study. *Neurology*. 2004;63(12):2316-2321. doi:10.1212/01.WNL.0000147474.29994.35
89. Weuve J, Kang JH, Manson JE, Breteler MM, Ware JH, Grodstein F. Physical activity, including walking, and cognitive function in older women. *JAMA*. 2004;292(12):1454-1461. doi:10.1001/jama.292.12.1454
90. Willey JZ, Park Moon Y, Ruder R, et al. Physical activity and cognition in the northern Manhattan study. *Neuroepidemiology*. 2014;42(2):100-106. doi:10.1159/000355975
91. Krell-Roesch J, Syrjanen JA, Bezold J, et al. Physical activity and trajectory of cognitive change in older persons: Mayo Clinic Study of Aging. *J Alzheimers Dis*. 2021;79(1):377-388. doi:10.3233/JAD-200959
92. Speh A, Wang R, Winblad B, et al. The relationship between cardiovascular health and rate of cognitive decline in young-old and old-old adults: a population-based study. *J Alzheimers Dis*. 2021;84(4):1523-1537. doi:10.3233/JAD-210280
93. Zeitzer JM, Blackwell T, Hoffman AR, Cummings S, Ancoli-Israel S, Stone K; Osteoporotic Fractures in Men (MrOS) Study Research Group. Daily patterns of accelerometer activity predict changes in sleep, cognition, and mortality in older men. *J Gerontol A Biol Sci Med Sci*. 2018;73(5):682-687. doi:10.1093/gerona/glw250
94. Fernández-Matarrubia M, Goni L, Rognoni T, et al. An active lifestyle is associated with better cognitive function over time in APOE ε4 non-carriers. *J Alzheimers Dis*. 2021;79(3):1257-1268. doi:10.3233/JAD-201090
95. Henstra M, Giltay E, van der Mast R, van der Velde N, Rhebergen D, Rius Ottenheim N. Does late-life depression counteract the beneficial effect of physical activity on cognitive decline: results from the NESDO Study. *J Geriatr Psychiatry Neurol*. 2022;35(3):450-459. doi:10.1177/08919887211002658
96. Litwin H, Shaul A. The effect of social network on the physical activity-cognitive function nexus in late life. *Int Psychogeriatr*. 2019;31(5):713-722. doi:10.1017/S1041610218001059
97. Mamalaki E, Charisis S, Anastasiou CA, et al. The Longitudinal association of lifestyle with cognitive health and dementia risk: findings from the HELIAD Study. *Nutrients*. 2022;14(14):2818. doi:10.3390/nu14142818
98. Li C, Ma Y, Hua R, Zheng F, Xie W. Long-term physical activity participation trajectories were associated with subsequent cognitive decline, risk of dementia and all-cause mortality among adults aged ≥50 years: a population-based cohort study. *Age Ageing*. 2022;51(3):afac071. doi:10.1093/ageing/afac071
99. Wang HX, Jin Y, Hendrie HC, et al. Late life leisure activities and risk of cognitive decline. *J Gerontol A Biol Sci Med Sci*. 2013;68(2):205-213. doi:10.1093/gerona/gls153
100. Chang M, Jonsson PV, Snaedal J, et al. The effect of midlife physical activity on cognitive function among older adults: AGES-Reykjavik Study. *J Gerontol A Biol Sci Med Sci*. 2010;65(12):1369-1374. doi:10.1093/gerona/glq152
101. Frederiksen KS, Verdelho A, Madureira S, et al; LADIS Study. Physical activity in the elderly is associated with improved executive function and processing speed: the LADIS Study. *Int J Geriatr Psychiatry*. 2015;30(7):744-750. doi:10.1002/gps.4220
102. Gaertner B, BATTERY AK, Finger JD, Wolfgruber S, Wagner M, Busch MA. Physical exercise and cognitive function across the life span: results of a nationwide population-based study. *J Sci Med Sport*. 2018;21(5):489-494. doi:10.1016/j.jsams.2017.08.022
103. Gardener H, Wright CB, Dong C, et al. Ideal cardiovascular health and cognitive aging in the Northern Manhattan Study. *J Am Heart Assoc*. 2016;5(3):e002731. doi:10.1161/JAHA.115.002731
104. Kooistra M, Boss HM, van der Graaf Y, Kappelle LJ, Biessels GJ, Geerlings MI; SMART-MR Study Group. Physical activity, structural brain changes and cognitive decline: the SMART-MR study. *Atherosclerosis*. 2014;234(1):47-53. doi:10.1016/j.atherosclerosis.2014.02.003

105. Singh-Manoux A, Hillsdon M, Brunner E, Marmot M. Effects of physical activity on cognitive functioning in middle age: evidence from the Whitehall II prospective cohort study. *Am J Public Health*. 2005;95(12):2252-2258. doi:10.2105/AJPH.2004.055574
106. Salthouse TA. Correlates of cognitive change. *J Exp Psychol Gen*. 2014;143(3):1026-1048. doi:10.1037/a0034847
107. Robinson SA, Lachman ME. Perceived control and cognition in adulthood: The mediating role of physical activity. *Psychol Aging*. 2018;33(5):769-781. doi:10.1037/pag0000273
108. Galle SA, Deijen JB, Milders MV, et al. The effects of a moderate physical activity intervention on physical fitness and cognition in healthy elderly with low levels of physical activity: a randomized controlled trial. *Alzheimers Res Ther*. 2023;15(1):12. doi:10.1186/s13195-022-01123-3
109. Loprinzi PD, Scott TM, Ikuta T, Addoh O, Tucker KL. Association of physical activity on changes in cognitive function: Boston Puerto Rican Health Study. *Phys Sportsmed*. 2019;47(2):227-231. doi:10.1080/00913847.2018.1547087
110. Sharifian N, Kraal AZ, Zaheed AB, Sol K, Zahodne LB. Longitudinal associations between contact frequency with friends and with family, activity engagement, and cognitive functioning. *J Int Neuropsychol Soc*. 2020;26(8):815-824. doi:10.1017/S1355617720000259
111. Taivalantti M, Barnett JH, Halt AH, et al. Depressive symptoms as predictors of visual memory deficits in middle-age. *J Affect Disord*. 2020;264:29-34. doi:10.1016/j.jad.2019.11.125
112. Dintica CS, Haaksmala ML, Olofsson JK, Bennett DA, Xu W. Joint trajectories of episodic memory and odor identification in older adults: patterns and predictors. *Aging (Albany NY)*. 2021;13(13):17080-17096. doi:10.18632/aging.203280
113. Hakala JO, Rovio SP, Pahkala K, et al. Physical activity from childhood to adulthood and cognitive performance in midlife. *Med Sci Sports Exerc*. 2019;51(5):882-890. doi:10.1249/MSS.0000000000001862
114. Stenling A, Eriksson Sörman D, Lindwall M, Machado L. Bidirectional within- and between-person relations between physical activity and cognitive function. *J Gerontol B Psychol Sci Soc Sci*. 2022;77(4):704-709. doi:10.1093/geronb/gbab234
115. Csajbók Z, Sieber S, Cullati S, Cermakova P, Cheval B. Physical activity partly mediates the association between cognitive function and depressive symptoms. *Transl Psychiatry*. 2022;12(1):414. doi:10.1038/s41398-022-02191-7
116. Huang ST, Tange C, Otsuka R, et al. Subtypes of physical frailty and their long-term outcomes: a longitudinal cohort study. *J Cachexia Sarcopenia Muscle*. 2020;11(5):1223-1231. doi:10.1002/jcsm.12577
117. Werneck AO, Stubbs B, Kandola A, et al. Prospective Associations of leisure-time physical activity with psychological distress and well-being: a 12-year cohort study. *Psychosom Med*. 2022;84(1):116-122. doi:10.1097/PSY.0000000000001023
118. Bott NT, Bettcher BM, Yokoyama JS, et al. Youthful processing speed in older adults: genetic, biological, and behavioral predictors of cognitive processing speed trajectories in aging. *Front Aging Neurosci*. 2017;9:55. doi:10.3389/fnagi.2017.00055
119. Young JC, Dowell NG, Watt PW, Tabet N, Rusted JM. Long-term high-effort endurance exercise in older adults: diminishing returns for cognitive and brain aging. *J Aging Phys Act*. 2016;24(4):659-675. doi:10.1123/japa.2015-0039
120. Jang H, Kim S, Kim B, Kim M, Jung J, Won CW. Functional constipation is associated with a decline in word recognition 2 years later in community-dwelling older adults: the Korean Frailty and Aging Cohort Study. *Ann Geriatr Med Res*. 2022;26(3):241-247. doi:10.4235/agmr.22.0092
121. Hammond NG, Stinchcombe A. Prospective associations between physical activity and memory in the Canadian Longitudinal Study on Aging: examining social determinants. *Res Aging*. 2022;44(9-10):709-723. doi:10.1177/01640275211070001
122. Bryan JW, Ward L. Smoking, alcohol use and engagement in exercise and cognitive performance among older adults. *Australas J Ageing*. 2002;21:67-73. doi:10.1111/j.1741-6612.2002.tb00420.x
123. Zaninotto P, Batty GD, Allerhand M, Deary IJ. Cognitive function trajectories and their determinants in older people: 8 years of follow-up in the English Longitudinal Study of Ageing. *J Epidemiol Community Health*. 2018;72(8):685-694. doi:10.1136/jech-2017-210116
124. Gignac GS, Szodorai ET. Effect size guidelines for individual differences researchers. *Pers Individ Dif*. 2016;102:74-78. doi:10.1016/j.paid.2016.06.069
125. Xu W, Wang HF, Wan Y, Tan CC, Yu JT, Tan L. Leisure time physical activity and dementia risk: a dose-response meta-analysis of prospective studies. *BMJ Open*. 2017;7(10):e014706. doi:10.1136/bmjopen-2016-014706

126. Wu W, Ding D, Zhao Q, et al; for Cohort Studies of Memory in an International Consortium (COSMIC). Dose-response relationship between late-life physical activity and incident dementia: a pooled analysis of 10 cohort studies of memory in an international consortium. *Alzheimers Dement*. 2023;19(1):107-122. doi:10.1002/alz.12628

SUPPLEMENT 1.

eMethods.

eTable 1. Search Strategy

eTable 2. Definitions of Cognitive Impairment and Decline

eTable 3. Tests Used to Assess Continuous Global Cognition, Verbal Fluency and Naming, Working Memory, Verbal Ability and Visuo-spatial Ability

eTable 4. Tests Used to Assess Episodic Memory, Executive Function and Processing Speed

eFigure 1. Quality of the Studies

eTable 5. Physical Activity and Cognition, Binary Outcomes, Supplementary Analyses

eFigure 2. Physical Activity and Cognitive Impairment or Decline, Forest Plot

eFigure 3. Physical Activity and Cognition, Binary Outcome, Contour-Enhanced Funnel Plot

eFigure 4. Physical Activity and Cognition, Binary Outcomes by Sample Size

eFigure 5. Physical Activity and Cognition, Binary Outcomes by Length of the Follow-Up

eFigure 6. Physical Activity and Cognition, Binary Outcomes by Follow-Up Rate (Studies Not Reporting Follow-Up Rate Are Excluded)

eTable 6. Physical Activity and Follow-Up Cognition: Supplementary Analyses

eFigure 7. PA and Follow-Up Cognition, Contour-Enhanced Funnel Plot

eTable 7. PA and Change in Cognition: Supplementary Analyses

eFigure 8. PA and Change in Cognition, Contour-Enhanced Funnel Plot

eTable 8. PA and Specific Cognitive Domains in Studies Adjusting for Preceding Level of Cognition

eReferences.

SUPPLEMENT 2.

eTable 1. Quality Review

eTable 2. Binary Outcomes

eTable 3. Continuous Outcomes

eTable 4. Executive Function

eTable 5. Episodic Memory

eTable 6. Processing Speed

eTable 7. Verbal Fluency and Naming

eTable 8. Working Memory

eTable 9. Verbal Ability

eTable 10. Visuospatial Ability

eTable 11. Dose Analysis: Binary Outcomes

eTable 12. Dose Analysis: Continuous Outcomes

SUPPLEMENT 3.

Data Sharing Statement