

**JYX**



JYVÄSKYLÄN YLIOPISTO  
UNIVERSITY OF JYVÄSKYLÄ

**This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.**

**Author(s):** Palmberg, Lotta; Löppönen, Antti; Hyvärinen, Matti; Portegijs, Erja; Rantanen, Taina; Rantalainen, Timo; Karavirta, Laura

**Title:** Physical Behavior Profiles Among Older Adults and Their Associations With Physical Capacity and Life-Space Mobility

**Year:** 2024

**Version:** Accepted version (Final draft)

**Copyright:** © 2024 Human Kinetics

**Rights:** In Copyright

**Rights url:** <http://rightsstatements.org/page/InC/1.0/?language=en>

**Please cite the original version:**

Palmberg, L., Löppönen, A., Hyvärinen, M., Portegijs, E., Rantanen, T., Rantalainen, T., & Karavirta, L. (2024). Physical Behavior Profiles Among Older Adults and Their Associations With Physical Capacity and Life-Space Mobility. *Journal of Aging and Physical Activity*, Ahead of Print. <https://doi.org/10.1123/japa.2023-0225>

1       **Physical behavior profiles among older adults and their associations with physical**  
2                                   **capacity and life-space mobility**

3  
4       Lotta Palmberg, PhD<sup>1</sup>, Antti Löppönen, MSc<sup>1,2</sup>, Matti Hyvärinen, MSc<sup>1</sup>, Erja Portegijs, PhD<sup>3</sup>,  
5                                   Taina Rantanen, PhD<sup>1</sup>, Timo Rantalainen, PhD<sup>1</sup> & Laura Karavirta, PhD<sup>1</sup>

6  
7       <sup>1</sup>Gerontology Research Center and Faculty of Sport and Health Sciences, University of  
8                                   Jyväskylä, Jyväskylä, Finland

9       <sup>2</sup>Physical Activity, Sports and Health Research Group, Department of Movement Sciences,  
10                                   KU Leuven, Leuven, Belgium

11       <sup>3</sup>University of Groningen, University Medical Center Groningen, Center for Human  
12                                   Movement Sciences, Groningen, The Netherlands.

13  
14       Corresponding author:

15       Lotta Palmberg

16       P.O. Box 35

17       40014-FI, University of Jyväskylä

18       lotta.m.palmberg@jyu.fi

19       ORCID: 0000-0001-5108-9330

20  
21       Running head: Physical behavior profiles in old age

22

## Abstract

We identified data-driven multidimensional physical activity (PA) profiles using several novel accelerometer-derived metrics.

Participants aged 75, 80, and 85 (n=441) wore tri-axial accelerometers for 3-7 days. PA profiles were formed with k-means cluster analysis based on PA minutes, intensity, fragmentation, sit-to-stand transitions, and gait bouts for men and women. Associations with physical capacity and life-space mobility were examined using age-adjusted general linear models.

Three profiles emerged: "Exercisers" and "actives" accumulated relatively high PA minutes, with actives engaging in lighter intensity PA. "Inactives" had the highest activity fragmentation and lowest PA volume, intensity, and gait bouts. Inactives showed lower scores in physical capacity and life-space mobility compared to exercisers and actives. Exercisers and actives had similar physical capacity and life-space mobility, except female exercisers had higher walking speed in the 6-minute walk test.

Our findings demonstrate the importance of assessing PA as multidimensional behavior rather than focusing on a single metric.

Keywords: physical function, physical activity, cluster analysis, aging

## Introduction

Physical activity (PA) can be defined as any bodily movement produced by skeletal muscles that results in energy expenditure (Caspersen et al., 1985). PA is a known modifiable behavior that can decrease the risk for several chronic conditions and maintain physical capacity (Paterson & Warburton, 2010; Warburton et al., 2006), defined as individual's ability to execute a task or an action in a standard environment (World Health Organization, 2001). The use of accelerometry in assessing free-living PA among older adults has increased over the past decade, but most studies assess PA and sedentary behavior using single metrics such as daily total volumes or step counts (Schrack et al., 2016; Shiroma et al., 2018). PA, however, is multidimensional and it is likely that no single metric can adequately describe individual's PA (Thompson et al., 2015). For instance, physical behavior characteristics such as hour-by-hour accumulation patterns, bouts of PA and sedentary behavior and activity fragmentation are associated with health-related outcomes (Bellettiere et al., 2017; Brady et al., 2022; Palmberg et al., 2020; Schrack et al., 2019). Building daily physical behavior profiles using a combination of several accelerometer-derived metrics could better capture the multidimensional nature of daily activity.

Data-driven, person-centered approaches such as mixture modelling and cluster analysis allow for the use of multiple variables of PA in the analyses and can provide a better understanding on how these are combined in older individuals' everyday life. These approaches for device-based physical behavior profiling have been increasingly used among different populations including children (Verswijveren et al., 2020), adult population (Farrahi et al., 2021; Gupta et al., 2020; von Rosen et al., 2020) and clinical subgroups (Geidl et al., 2019; Mesquita et al., 2017). Data-driven daily physical behavior profiling focusing on older people remains largely unexplored. A few previous studies investigated the associations of physical behavior clusters with health also among older people, but these studies focused on the total

1 volumes and patterns of sedentary time rather than PA patterns (Laudani et al., 2013; Manta et  
2 al., 2019; O'Regan et al., 2021).

3         Physical behavior assessment in old age has its unique challenges and thus, there is a  
4 need to move beyond PA metrics describing solely total volume (Shiroma et al., 2018). While  
5 there is a need to explore data-driven physical behavior profiles in old age, consideration should  
6 also be given into which metrics to use among older people. For instance, cut-points based on  
7 absolute PA intensity may not accurately describe PA among older adults with more  
8 heterogeneity in physical capacity than their younger counterparts. With declining physical  
9 capacity, the relative intensity of physical activities may become higher, despite the decline in  
10 absolute intensity (Kujala et al., 2017; Schrack et al., 2018). In the AGNES study, our aim has  
11 been to develop PA assessment methods that may be able to overcome these challenges. The  
12 methods include individually scaled cut-points for PA based on preferred walking speed that  
13 showed promise as an assessment method for relative PA (Karavirta et al., 2020). We also  
14 identified free-living sit-to-stand (STS) transitions and their intensity, which are commonly  
15 performed among older people and are essential for independence in daily life (Löppönen et  
16 al., 2022). In addition, we have used other novel promising physical behavior metrics such as  
17 active-to-sedentary transition probability, an indicator of activity fragmentation (Schrack et al.,  
18 2019). In our earlier study, we found that activity fragmentation was associated with greater  
19 physical fatigability even beyond total PA volume (Palmberg et al., 2020). In addition, we  
20 identified gait bouts and their intensity from free-living accelerometer data and developed cut-  
21 points for higher risk of walking difficulty (Skantz et al., 2021).

22         The aim of this study was to identify data-driven physical behavior profiles among  
23 community-dwelling older people including novel accelerometer-based metrics describing the  
24 volume, intensity and accumulation patterns of PA. Furthermore, we studied the utility of the  
25 profiles by examining whether older people with different physical behavior profiles differ in

1 terms of physical capacity and life-space mobility (ie. the spatial area where the person moves  
2 through in daily life), known correlates of PA.

### 3 **Methods**

4  
5 This study forms a part of the “Active Aging—resilience and external support as  
6 modifiers of the disablement outcome” (AGNES) project (Portegijs et al., 2019; Rantanen et  
7 al., 2018). Briefly, participants were community-dwelling 75-, 80-, and 85-year-old people  
8 living in the Jyväskylä area, Finland and the initial sample was recruited from the Digital and  
9 Population Data Services Agency. Eligibility criteria included willingness to participate and  
10 residing in the study area. A total of 1021 people participated in a home interview in 2017-  
11 2018, of whom 495 agreed to wear a tri-axial accelerometer (Portegijs et al., 2019). Participants  
12 with valid data on all accelerometer-derived metrics on at least three complete days were  
13 included in the present analyses (n=441). Days with complete data, including no non-wear,  
14 from midnight to midnight based on visual inspection were considered valid.

### 15 **Ethics**

16 The study protocol followed the principles laid down by the Declaration of Helsinki.  
17 The study has been approved by the the Ethical Committee of the Central Finland Health  
18 Care District. All participants signed a written informed consent.

### 19 *Accelerometer-assessed physical behavior*

20 The accelerometers (range  $\pm 16$  g, 13-bit analog-to-digital conversion, sampled at 100  
21 Hz, UKK RM42; UKK Terveyspalvelut Oy, Tampere, Finland) were attached on participants’  
22 dominant thigh using a waterproof film for 7–10 consecutive days following a home interview  
23 (Portegijs et al., 2019). The resultant accelerations were calculated for sampling instants and  
24 mean amplitude deviation (MAD, in g) calculated for non-overlapping 5-second epochs  
25 (Portegijs et al., 2019). Posture estimation was done following the approach by Vähä-Ypyä and

1 colleagues (Vähä-Ypyä et al., 2018). Posture categories (sitting/lying down or upright) for each  
2 5-second epoch were identified, and the median category for each minute of recording was  
3 used to calculate mean daily minutes in an upright posture. All epochs of a minimum duration  
4 of 20 seconds, upright posture and acceleration between 0.035 g and 1.2 g were identified as  
5 gait bouts based on laboratory experimentation, and mean number of gait bouts, mean gait bout  
6 intensity and mean duration of gait bouts were then calculated (Skantz et al., 2021).  
7 Furthermore, activity fragmentation was assessed as Active-to-Sedentary Transition  
8 Probability (ASTP) (Schrack et al., 2019). ASTP was calculated separately for mean daily  
9 minutes based on MAD values  $> 0.0167$  g classified as at least light activity and mean daily  
10 minutes spent in an upright posture by dividing the mean active daily bouts by the mean sum  
11 of active daily minutes (Palmberg et al., 2020).

12 Minutes spent in different PA intensities were categorized with the following cut points  
13 originally developed for high-pass filtered vector magnitude: any minute with a MAD value  
14 below 0.0420 g as non-movement time, from 0.0420 g to 0.2375 g as light activity,  $\geq 0.2375$  g  
15 to  $< 0.6285$  g as moderate activity and  $\geq 0.6285$  g as vigorous activity. These cut points  
16 corresponded to 1.5, 3 and 6 METs, respectively, following a linear equation by White and  
17 colleagues (White et al., 2019). Relative PA was then calculated as the number of epochs above  
18 or equal to the mean acceleration calculated during a laboratory-measured 6-minute walk test  
19 (6MWT) (Karavirta et al., 2020).

20 Daily STS transitions were detected from the accelerometer data using an open-access  
21 algorithm whose structure, code and properties are described elsewhere (Löppönen et al.,  
22 2022). The volume of the STS transitions was determined as the number of transitions per  
23 monitoring day.

24 *Descriptive and outcome measures*

1 Education was assessed by a single question asking participants to report their total  
2 years of education. Participants were asked about their living situation (alone, with spouse,  
3 with children or grandchildren, with relatives, siblings or other people), and dichotomized into  
4 alone vs. with others. Willingness and perceived opportunities for PA participation were asked  
5 using two single questions (Rantanen et al., 2018). Cognitive function was assessed using mini  
6 mental state examination (MMSE) (Folstein et al., 1975) and depressive symptoms were  
7 assessed using Centre for Epidemiologic studies Depression Scale (CES-D) (Radloff, 1977).

8 Lower-extremity physical performance was assessed during the home interview by  
9 trained a researcher using the Short Physical Performance Battery (SPPB) (Guralnik et al.,  
10 1994). Maximal knee extension force of the dominant lower leg (knee at 60 degrees) was  
11 measured in a sitting position using an adjustable dynamometer chair (Metitur LTD, Jyväskylä,  
12 Finland). The highest force of at least three attempts was selected for analysis (Rantanen et al.,  
13 1997). Self-reported habitual PA was assessed using the eight-item Yale Physical Activity  
14 Survey for older adults (YPAS, range 0-137, higher scores indicate higher PA) (Dipietro et al.,  
15 1993). Finally, life-space mobility was assessed using 15-item University of Alabama at  
16 Birmingham Study of Aging Life-Space Assessment (LSA, range 0-120, higher scores indicate  
17 higher life-space mobility) (Baker et al., 2003).

## 18 **Statistical methods**

19 Profiles with similar physical behavior characteristics were identified separately for  
20 men and women using k-means clustering algorithm (Hartigan & Wong, 1979). The k-means  
21 clustering was used instead of model-based methods due to the violation of conditional  
22 independence assumption caused by the strong direct relationships between the features  
23 (Oberski, 2016). All participants with valid accelerometer measurements and no missing values



1 (177 men, 264 women) were included in the cluster analyses. The cluster analyses were carried  
2 out in R (Team, 2013).

3 First, correlation-based principal component analysis (Hotelling, 1933) was carried out  
4 to deal with multicollinearity. For both men and women, three principal components were  
5 chosen based on the scree test (Cattell, 1966). They explained 74.9 and 73.6% of the variance,  
6 respectively. For k-means clustering, the similarity of clusters was assessed using Euclidian  
7 distance and number of clusters was determined based on the “elbow method” using the  
8 “factoextra” package (Kassambara & Mundt, 2020). The optimal number of clusters was three  
9 for both sexes. Clusters were validated using graphical inspection and studying the cluster-wise  
10 stability by assessing the bootstrap distribution of the Jaccard coefficient (Hennig, 2007). The  
11 bootstrapping with 100 resamples was conducted using the “fpc” package (*Fpc*, n.d.).

12 Descriptive characteristics according to the physical behavior clusters were studied with  
13 means or percentages. Group differences were assessed with the chi-square test for categorical  
14 variables and the Kruskal-Wallis test for continuous variables. Age-adjusted group differences  
15 were assessed using general linear models. To account for multiple testing, the p-values were  
16 corrected using the Bonferroni method. Analyses were conducted using data from the  
17 participants with no missing data in the variables of interest. The percentage of missing data  
18 varied from 0 to 2% across the variables used in the analyses. All other analyses were  
19 performed using IBM SPSS Statistics 26 (IBM Corp, Armonk, NY, USA). The level of  
20 statistical significance was set to  $p < 0.05$ .

## 21 **Results**

### 22 **Identification of physical behavior profiles**

23 The independent clustering analyses for men and women resulted in three clusters that  
24 were similar between the sexes. The bootstrap distribution of the Jaccard coefficients were  
25 0.67, 0.76, and 0.86 in men and 0.86, 0.91, and 0.91 in women indicating high cluster-wise

1 stability for most clusters (Hennig, 2008). The clusters were labelled as “the exercisers”, “the  
2 actives” and “the inactives” based on their physical behavior characteristics (Table 1 for  
3 women and Table 2 for men). Due to the similarity of clusters in men and women, the same  
4 labels were used for both sexes.

5 The exercisers included 22.7% of women and 10.7% of men. The exercisers  
6 accumulated the highest moderate PA minutes, highest relative PA minutes, the longest  
7 duration of gait bouts and highest mean daily acceleration compared to the actives and the  
8 inactives ( $p < 0.05$  for all). The difference was especially notable concerning relative PA  
9 minutes which the exercisers accumulated 227% more compared to the actives and 319% more  
10 compared to the inactives among women ( $p < 0.001$  for both), and 373% and 525% more among  
11 men ( $p < 0.001$  for both). The actives and the inactives accumulated similar relative PA in both  
12 sexes.

13 The actives included 37.9% of women and 43.5% of men. The actives formed the  
14 intermediate profile in terms of moderate PA minutes and mean acceleration but accumulated  
15 22% and 59% higher light-intensity PA minutes compared to the exercisers and the inactives  
16 among women, and 34% and 65% higher among men ( $p < 0.001$  for all), respectively. In  
17 addition, the women actives accumulated 22% and 70% higher number of gait bouts compared  
18 to the exercisers and the inactives ( $p < 0.001$  for both), and the men actives 32% ( $p = 0.002$ ) and  
19 70% ( $p < 0.001$ ) higher, respectively.

20 The inactives included 39.4% of women and 45.8% of men. While non-movement time  
21 between the exercisers and the actives was similar among both women and men, the inactives  
22 accumulated 5-6% higher non-movement time among women ( $p < 0.001$  for both) and 7%  
23 higher among men ( $p < 0.001$  for both) compared to the exercisers and the actives. Overall, the  
24 inactives formed the most inactive profile, accumulating the least PA minutes and 33% higher

1 activity fragmentation compared to the actives and 60% higher compared to the exercisers  
2 among women ( $p<0.001$ ), and 31% and 42% higher among men, respectively. Among men the  
3 actives had slightly higher vigorous PA minutes and mean gait bout duration compared to the  
4 inactives, while no difference was observed among women.

### 5 **Associations with demographic characteristics and measures of mobility and health**

6 Among men, the inactives had the fewest years of education, but among women there  
7 was no statistically significant difference. Among women, the inactives reported more  
8 depressive symptoms, but similar difference was not observed among men. There were no  
9 statistically significant differences in cognitive function or living situation. Among women,  
10 over 70% of the actives and the inactives reported that they wanted to be more physically active  
11 (vs. 39% in the exercisers). Similarly, among men over 70% of the inactives reported a  
12 willingness to increase PA (vs. 50% in actives and 35% in exercisers) (Table 3).

13 Among women, the inactives had 27.9% ( $p<0.001$ ) and 14.3% ( $p=0.013$ ) lower self-  
14 reported PA scores compared to the exercisers and the actives, respectively. In addition, women  
15 actives had 15.9% lower self-reported PA compared to the exercisers ( $p=0.004$ ). Among men,  
16 the inactives had 33.5% and 22.5% lower self-reported PA compared to the exercisers and the  
17 actives ( $p<0.001$  for both), but the difference between the exercisers and actives did not reach  
18 statistical significance. Among women, the inactives had 15.4% and 12.7% ( $p<0.001$  for both)  
19 lower scores in life-space mobility compared to the exercisers and the actives, respectively.  
20 Among men, the inactives had 10.3% lower scores in life-space mobility compared to the  
21 actives ( $p=0.004$ ), but the other differences did not reach statistical significance (Table 3). In  
22 the age-adjusted models, the findings did not materially change (Table 4).

### 23 **Associations with demographic characteristics and physical capacity**

1 Among women, the inactives had 9.2% ( $p=0.002$ ) and 6.6% ( $p=0.011$ ) lower SPPB  
2 scores, and 15.4% ( $p<0.001$ ) and 10.1% ( $p=0.019$ ) lower maximal knee extension force  
3 compared to the exercisers and the actives, respectively. The differences did not reach  
4 statistically significance between the women exercisers and the women actives. Concerning  
5 walking speed, the women inactives had 23.1% and 16.7% slower walking speed in the 6MWT,  
6 and 15.8% and 11.1 slower in the 10-m test ( $p<0.001$  for all) compared to the exercisers and  
7 the actives. The women exercisers had higher walking speed in the 6MWT compared to the  
8 women actives ( $p=0.013$ ) but did not differ in 10-m walking speed.

9 Among men, the exercisers and the actives did not differ in any of the indicators of  
10 physical capacity. The men inactives had 11.4% ( $p=0.011$ ) and 7.3% ( $p=0.013$ ) lower SPPB  
11 scores compared to the exercisers and the actives, respectively. Furthermore, the men inactives  
12 had 11.3% lower maximal knee extension force than the actives ( $p=0.009$ ) but no statistically  
13 significant difference compared to the exercisers. Concerning walking speed, the men inactives  
14 had 15.4% slower 6MWT speed compared to both the exercisers and the actives ( $p=0.001$ ). In  
15 the 10-m test, the men inactives had 14.3% slower walking speed compared to the actives  
16 ( $p<0.001$ ) but no difference compared to the exercisers. In the age-adjusted models (Table 4),  
17 the findings remained materially unchanged.

## 18 Discussion

19 In the present study, we identified three physical behavior profiles among community-  
20 dwelling older people, characterizing different phenotypes of physical activity behavior in old  
21 age. Similar profiles emerged for women and men, separately. Parallel to earlier research  
22 findings, we found that compared to the two more physically active profiles, the inactives were  
23 characterized by lower life-space mobility, self-reported PA, and physical capacity. In women,  
24 walking speed in 6MWT/10 m and self-reported PA differed statistically between the  
25 exercisers and the actives, but in men significant differences were not observed between these

1 two groups. These findings indicate that PA profiling can be a useful method for research  
2 aiming to combine multiple dimensions of physical activity in old age. The novelty of the  
3 present study is utilizing a data-driven, person-centered approach to assess the  
4 multidimensional physical activity combining a range of novel accelerometer-based metrics  
5 among older people.

6 Our findings demonstrate the relevance of assessing multiple PA dimensions among  
7 older people, rather than focusing on simple PA metrics such as activity minutes. An important  
8 observation was that although the exercisers and the actives accumulated similar total activity  
9 minutes, they did differ in several other PA characteristics such as relative PA minutes, gait  
10 bouts and mean intensity contributing towards a substantially different physical activity  
11 phenotype. These differences were also seen among women in walking speed and self-reported  
12 physical activity. The findings indicate that the use of distinctive multidimensional profiles has  
13 the potential to demonstrate differences in PA beyond what is captured by single PA metrics,  
14 and reducing accelerometer-based data into a simple physical activity metric may present a  
15 missed opportunity. Earlier studies identifying physical behavior clusters among older people  
16 found four distinct physical behavior clusters (Manta et al., 2019; O'Regan et al., 2021).  
17 Compared to these earlier studies, we used a wider range of PA metrics that address the  
18 common pitfalls of assessing PA among older adults, and capture accumulation patterns in  
19 addition to commonly used metrics of volume and intensity. Due to the differences in PA  
20 metrics, our findings are not comparable to these previous studies profiling physical activity in  
21 old age, and thus, future studies are needed to see whether similar profiles can be found among  
22 other older adult populations using a similar set of PA metrics.

23 The differences in physical capacity and life-space mobility between the inactives and  
24 the more active profiles were comparable with earlier research showing associations of higher  
25 PA levels, lower activity fragmentation and higher PA complexity with better physical function

1 (Paterson & Warburton, 2010; Rantalainen et al., 2022; Schrack et al., 2019; Simonsick et al.,  
2 2005) and higher life-space mobility (Portegijs et al., 2015; Tsai et al., 2016). We found that  
3 among both men and women, those who accumulated the least PA and had highest activity  
4 fragmentation (the inactives) had poorer physical capacity and lower life-space mobility  
5 compared to the more physically active profiles. Poorer physical capacity and health can limit  
6 the opportunities for PA among older people (Rai et al., 2020), while engaging in PA can also  
7 help in maintaining physical capacity in old age (Paterson & Warburton, 2010; Simonsick et  
8 al., 2005). Given the cross-sectional nature of our findings, future prospective studies are  
9 needed to confirm the predictive validity of the profiles in predicting changes in physical  
10 capacity and life-space mobility.

11         Interestingly, between the two more active profiles, we observed no differences in terms  
12 of physical capacity, besides the difference in walking speed among women. This finding is in  
13 line with earlier dose-response observations where the steepest risk reduction occurs at low  
14 volumes of PA, indicating that some activity is much better than none. This is important as  
15 there was a rather large difference in the amount of relative PA i.e. activity beyond the intensity  
16 of 6MWT between these profiles among both men and women. This finding is consistent with  
17 the growing research evidence that older people can benefit from a wide range of PA behaviors,  
18 including light-intensity activities, as already stated in the new World Health Organization  
19 2020 guidelines on health-enhancing PA and sedentary behaviour (Bull et al., 2020). Future  
20 prospective research is, however, needed to confirm whether older people with high activity  
21 levels can maintain their physical capacity and life-space mobility regardless of lower intensity  
22 levels.

23         The difference in the results among men and women concerning walking speed may be  
24 explained by lower robustness in the clusters among men or by overall higher walking speed  
25 among men. This would have allowed them to accumulate higher time in moderate and

1 vigorous PA. Furthermore, it may be that among highly functioning older people, the  
2 differences in higher intensity PA may rather be explained by individual preferences, than  
3 differences in physical function and health (Rai et al., 2020). The findings that the difference  
4 in relative PA was rather large between these profiles, and that the actives accumulated more  
5 light PA, can be interpreted in a way that the exercisers may be more likely to walk more for  
6 exercise, while the actives may be more likely to accumulate PA while doing household chores  
7 or running errands. This is also supported by differences in self-assessed PA (YPAS),  
8 suggesting that actives reported less walking and vigorous PA, and accumulated PA in  
9 activities that they did not perceive as PA or were not able to recall.

10         The strengths of the study include a continuous 3–7-day accelerometer recording, which  
11 allowed us to account for day-to-day variations in PA. Another strength is the availability of  
12 several novel metrics of physical behavior, that can together provide a more comprehensive  
13 description of the physical behavior of older people. There are also limitations that should be  
14 taken into account when interpreting the findings. First, the cross-sectional study setting does  
15 not allow conclusions about causality, and future longitudinal studies are needed. Second, the  
16 physical behavior clusters for older men were less robust, which may explain why we observed  
17 no differences between exercisers and actives among men. Third, we were not able to  
18 differentiate between sedentary time and sleep and thus, differences in non-movement time  
19 could also be explained by differences in sleep duration. Finally, it should be noted that the  
20 participants agreeing to participate in the PA monitoring had better health and higher PA than  
21 participants who only participated in the home interview (Portegijs et al., 2019), and hence the  
22 present sample underrepresented the less healthy and the less physically active part of the  
23 population.

## 24         **Conclusions**

1           The findings of this study demonstrate the importance of assessing multidimensional  
2 PA rather than focusing merely on single metrics. We were able to identify distinct activity  
3 phenotypes among older people which provide a more comprehensive picture of the volume,  
4 patterns and intensity of PA in which older people engage in during their everyday lives.  
5 Notably, although the two more physically active profiles had similar total activity minutes,  
6 they exhibited notable differences in various other PA characteristics, contributing to distinct  
7 PA phenotypes. This observation underscores the importance of recognizing that studying PA  
8 minutes alone is not sufficient when investigating the PA behavior of older adults. Instead,  
9 researchers should place greater emphasis on carefully selecting PA characteristics according  
10 to the context and purpose of their studies. In addition, similar profiles arose from both the  
11 women and the men independently, further supporting the utility of the profiles in  
12 characterizing PA behavior. Although these profiles showed associations with known  
13 correlates of physical activity in a cross-sectional study setting, the predictive validity of these  
14 profiles needs to be confirmed in future prospective studies.

## 15 **Acknowledgments**

16           The Gerontology Research Center is joint effort between the University of  
17 Jyväskylä, Finland and the University of Tampere, Finland.

## 18 **Funding:**

19           This work was supported by the European Research Council (693045 to Ta.R.); the  
20 Academy of Finland (339391 and 346462 to L.K., 321336 and 328818 to Ti.R. and 310526 to  
21 Ta.R.) and the University of Jyväskylä. The content of this manuscript does not reflect the  
22 official opinion of the European Union. Responsibility for the information and views  
23 expressed in the manuscript lies entirely with the authors.

## 24 **Conflict of interest:**



1            On behalf of all authors, the corresponding author states that there is no conflict of  
2 interest.

3

## 1 **References**

- 2 Baker, P. S., Bodner, E. V., & Allman, R. M. (2003). Measuring lifeâ€ space mobility in  
3 communityâ€ dwelling older adults. *Journal of the American Geriatrics Society*,  
4 *51*(11), 1610–1614.
- 5 Bellettiere, J., Winkler, E. A., Chastin, S. F., Kerr, J., Owen, N., Dunstan, D. W., & Healy, G.  
6 N. (2017). Associations of sitting accumulation patterns with cardio-metabolic risk  
7 biomarkers in Australian adults. *PloS One*, *12*(6), e0180119.
- 8 Brady, R., Brown, W. J., Hillsdon, M., & Mielke, G. I. (2022). Patterns of Accelerometer-  
9 measured Physical Activity and Health Outcomes in Adults: A Systematic Review.  
10 *Medicine and Science in Sports and Exercise*.
- 11 Bull, F. C., Al-Ansari, S. S., Biddle, S., Borodulin, K., Buman, M. P., Cardon, G., Carty, C.,  
12 Chaput, J.-P., Chastin, S., & Chou, R. (2020). World Health Organization 2020  
13 guidelines on physical activity and sedentary behaviour. *British Journal of Sports*  
14 *Medicine*, *54*(24), 1451–1462.
- 15 Caspersen, C. J., Powell, K. E., & Christenson, G. M. (1985). Physical activity, exercise, and  
16 physical fitness: Definitions and distinctions for health-related research. *Public Health*  
17 *Reports (Washington, D.C.: 1974)*, *100*(2), 126–131.
- 18 Cattell, R. B. (1966). The Scree Test For The Number Of Factors. *Multivariate Behavioral*  
19 *Research*, *1*(2), 245–276. [https://doi.org/10.1207/s15327906mbr0102\\_10](https://doi.org/10.1207/s15327906mbr0102_10)
- 20 Dipietro, L., Caspersen, C. J., Ostfeld, A. M., & Nadel, E. R. (1993). A survey for assessing  
21 physical activity among older adults. *Medicine & Science in Sports & Exercise*,  
22 *Journal Article*.
- 23 Farrahi, V., Kangas, M., Kiviniemi, A., Puukka, K., Korpelainen, R., & Jämsä, T. (2021).  
24 Accumulation patterns of sedentary time and breaks and their association with

1 cardiometabolic health markers in adults. *Scandinavian Journal of Medicine &*  
2 *Science in Sports*, 31(7), 1489–1507. <https://doi.org/10.1111/sms.13958>

3 Folstein, M. F., Folstein, S. E., & McHugh, P. R. (1975). “Mini-mental state”: A practical  
4 method for grading the cognitive state of patients for the clinician. *Journal of*  
5 *Psychiatric Research*, 12(3), 189–198.

6 *fpc: Flexible Procedures for Clustering version 2.2-9 from CRAN*. (n.d.). Retrieved June 29,  
7 2022, from <https://rdrr.io/cran/fpc/>

8 Geidl, W., Carl, J., Cassar, S., Leibert, N., Mino, E., Wittmann, M., Wagner, R., Schultz, K.,  
9 & Pfeifer, K. (2019). Physical activity and sedentary behaviour patterns in 326  
10 persons with COPD before starting a pulmonary rehabilitation: A cluster analysis.  
11 *Journal of Clinical Medicine*, 8(9), 1346.

12 Gupta, N., Hallman, D. M., Dumuid, D., Vij, A., Rasmussen, C. L., Jørgensen, M. B., &  
13 Holtermann, A. (2020). Movement behavior profiles and obesity: A latent profile  
14 analysis of 24-h time-use composition among Danish workers. *International Journal*  
15 *of Obesity*, 44(2), Article 2. <https://doi.org/10.1038/s41366-019-0419-8>

16 Guralnik, J. M., Simonsick, E. M., Ferrucci, L., Glynn, R. J., Berkman, L. F., Blazer, D. G.,  
17 Scherr, P. A., & Wallace, R. B. (1994). A short physical performance battery  
18 assessing lower extremity function: Association with self-reported disability and  
19 prediction of mortality and nursing home admission. *Journal of Gerontology*, 49(2),  
20 85.

21 Hartigan, J. A., & Wong, M. A. (1979). Algorithm AS 136: A K-Means Clustering  
22 Algorithm. *Journal of the Royal Statistical Society. Series C (Applied Statistics)*,  
23 28(1), 100–108. <https://doi.org/10.2307/2346830>

24 Hennig, C. (2007). Cluster-wise assessment of cluster stability. *Computational Statistics &*  
25 *Data Analysis*, 52(1), 258–271. <https://doi.org/10.1016/j.csda.2006.11.025>

- 1 Hennig, C. (2008). Dissolution point and isolation robustness: Robustness criteria for general  
2 cluster analysis methods. *Journal of Multivariate Analysis*, 99(6), 1154–1176.  
3 <https://doi.org/10.1016/j.jmva.2007.07.002>
- 4 Hotelling, H. (1933). Analysis of a complex of statistical variables into principal components.  
5 *Journal of Educational Psychology*, 24(6), 417–441.  
6 <https://doi.org/10.1037/h0071325>
- 7 Karavirta, L., Rantalainen, T., Skantz, H., Lisko, I., Portegijs, E., & Rantanen, T. (2020).  
8 Individual Scaling of Accelerometry to Preferred Walking Speed in the Assessment of  
9 Physical Activity in Older Adults. *The Journals of Gerontology: Series A*, 75(9),  
10 e111–e118. <https://doi.org/10.1093/gerona/glaa142>
- 11 Kassambara, A., & Mundt, F. (2020). *factoextra: Extract and Visualize the Results of*  
12 *Multivariate Data Analyses* (1.0.7). <https://CRAN.R-project.org/package=factoextra>
- 13 Kujala, U., Pietilä, J., Myllymäki, T., Mutikainen, S., Föhr, T., Korhonen, I., & Helander, E.  
14 (2017). Physical activity: Absolute intensity vs. Relative-to-fitness-level volumes.  
15 *Medicine and Science in Sports and Exercise*, 49(Journal Article).
- 16 Laudani, L., Vannozzi, G., Sawacha, Z., Della Croce, U., Cereatti, A., & Macaluso, A.  
17 (2013). Association between physical activity levels and physiological factors  
18 underlying mobility in young, middle-aged and older individuals living in a city  
19 district. *PLoS One*, 8(9), e74227.
- 20 Löppönen, A., Karavirta, L., Koivunen, K., Portegijs, E., Rantanen, T., Finni, T., Delecluse,  
21 C., Van Roie, E., & Rantalainen, T. (2022). Association Between Free-Living Sit-to-  
22 Stand Transition Characteristics, and Lower-Extremity Performance, Fear of Falling,  
23 and Stair Negotiation Difficulties Among Community-Dwelling 75 to 85-Year-Old  
24 Adults. *The Journals of Gerontology: Series A*, 77(8), 1644–1653.  
25 <https://doi.org/10.1093/gerona/glac071>

- 1 Manta, S. W., Sandreschi, P. F., Matias, T. S., Tomicki, C., & Benedetti, T. R. (2019).  
2 Clustering of physical activity and sedentary behavior associated to risk for metabolic  
3 syndrome in older adults. *Journal of Aging and Physical Activity*, 27(6), 781–786.
- 4 Mesquita, R., Spina, G., Pitta, F., Donaire-Gonzalez, D., Deering, B. M., Patel, M. S.,  
5 Mitchell, K. E., Alison, J., van Gestel, A. J., Zogg, S., Gagnon, P., Abascal-Bolado,  
6 B., Vagaggini, B., Garcia-Aymerich, J., Jenkins, S. C., Romme, E. A., Kon, S. S.,  
7 Albert, P. S., Waschki, B., ... Spruit, M. A. (2017). Physical activity patterns and  
8 clusters in 1001 patients with COPD. *Chronic Respiratory Disease*, 14(3), 256–269.  
9 <https://doi.org/10.1177/1479972316687207>
- 10 Oberski, D. (2016). Mixture Models: Latent Profile and Latent Class Analysis. In J.  
11 Robertson & M. Kaptein (Eds.), *Modern Statistical Methods for HCI* (pp. 275–287).  
12 Springer International Publishing. [https://doi.org/10.1007/978-3-319-26633-6\\_12](https://doi.org/10.1007/978-3-319-26633-6_12)
- 13 O'Regan, A., Hannigan, A., Glynn, L., Bengoechea, E. G., Donnelly, A., Hayes, G., Murphy,  
14 A. W., Clifford, A. M., Gallagher, S., & Woods, C. B. (2021). A cluster analysis of  
15 device-measured physical activity behaviours and the association with chronic  
16 conditions, multi-morbidity and healthcare utilisation in adults aged 45 years and  
17 older. *Preventive Medicine Reports*, 24, 101641.
- 18 Palmberg, L., Rantalainen, T., Rantakokko, M., Karavirta, L., Siltanen, S., Skantz, H.,  
19 Saajanaho, M., Portegijs, E., & Rantanen, T. (2020). The associations of activity  
20 fragmentation with physical and mental fatigability among community-dwelling 75-,  
21 80-, and 85-year-old people. *The Journals of Gerontology: Series A*, 75(9), e103–  
22 e110.
- 23 Paterson, D. H., & Warburton, D. E. (2010). Physical activity and functional limitations in  
24 older adults: A systematic review related to Canada's Physical Activity Guidelines.  
25 *International Journal of Behavioral Nutrition and Physical Activity*, 7(1), 38.

- 1 Portegijs, E., Karavirta, L., Saajanaho, M., Rantalainen, T., & Rantanen, T. (2019). Assessing  
2 physical performance and physical activity in large population-based aging studies:  
3 Home-based assessments or visits to the research center? *BMC Public Health*, *19*(1),  
4 1570.
- 5 Portegijs, E., Tsai, L.-T., Rantanen, T., & Rantakokko, M. (2015). Moving through life-space  
6 areas and objectively measured physical activity of older people. *PloS One*, *10*(8),  
7 e0135308.
- 8 Radloff, L. S. (1977). The CES-D scale: A self-report depression scale for research in the  
9 general population. *Applied Psychological Measurement*, *1*(3), 385–401.
- 10 Rai, R., Jongenelis, M. I., Jackson, B., Newton, R. U., & Pettigrew, S. (2020). Factors  
11 influencing physical activity participation among older people with low activity  
12 levels. *Ageing and Society*, *40*(12), 2593–2613.  
13 <https://doi.org/10.1017/S0144686X1900076X>
- 14 Rantalainen, T., Koivunen, K., Portegijs, E., Rantanen, T., Palmberg, L., Karavirta, L., &  
15 Chastin, S. (2022). Is Complexity of Daily Activity Associated with Physical  
16 Function and Life Space Mobility among Older Adults? *Medicine and Science in*  
17 *Sports and Exercise*.
- 18 Rantanen, T., Era, P., & Heikkinen, E. (1997). Physical activity and the changes in maximal  
19 isometric strength in men and women from the age of 75 to 80 years. *Journal of the*  
20 *American Geriatrics Society*, *45*(12), 1439–1445.
- 21 Rantanen, T., Saajanaho, M., Karavirta, L., Siltanen, S., Rantakokko, M., Viljanen, A.,  
22 Rantalainen, T., Pynnönen, K., Karvonen, A., & Lisko, I. (2018). Active aging–  
23 resilience and external support as modifiers of the disablement outcome: AGNES  
24 cohort study protocol. *BMC Public Health*, *18*(1), 565.

1 Schrack, J. A., Cooper, R., Koster, A., Shiroma, E. J., Murabito, J. M., Rejeski, W. J.,  
2 Ferrucci, L., & Harris, T. B. (2016). Assessing daily physical activity in older adults:  
3 Unraveling the complexity of monitors, measures, and methods. *Journals of*  
4 *Gerontology Series A: Biomedical Sciences and Medical Sciences*, 71(8), 1039–1048.

5 Schrack, J. A., Kuo, P.-L., Wanigatunga, A. A., Di, J., Simonsick, E. M., Spira, A. P.,  
6 Ferrucci, L., & Zipunnikov, V. (2019). Active-to-Sedentary Behavior Transitions,  
7 Fatigability, and Physical Functioning in Older Adults. *The Journals of Gerontology:*  
8 *Series A*, 74(4), 560–567.

9 Schrack, J. A., Leroux, A., Fleg, J. L., Zipunnikov, V., Simonsick, E. M., Studenski, S. A.,  
10 Crainiceanu, C., & Ferrucci, L. (2018). Using Heart Rate and Accelerometry to  
11 Define Quantity and Intensity of Physical Activity in Older Adults. *The Journals of*  
12 *Gerontology: Series A*, 73(5), 668–675.

13 Shiroma, E. J., Schrack, J. A., & Harris, T. B. (2018). Accelerating Accelerometer Research  
14 in Aging. *The Journals of Gerontology: Series A*, 73(5), 619–621.  
15 <https://doi.org/10.1093/gerona/gly033>

16 Simonsick, E. M., Guralnik, J. M., Volpato, S., Balfour, J., & Fried, L. P. (2005). Just get out  
17 the door! Importance of walking outside the home for maintaining mobility: Findings  
18 from the women’s health and aging study. *Journal of the American Geriatrics Society*,  
19 53(2), 198–203.

20 Skantz, H., Rantalainen, T., Karavirta, L., Rantakokko, M., Palmberg, L., Portegijs, E., &  
21 Rantanen, T. (2021). Associations Between Accelerometer-Based Free-Living  
22 Walking and Self-Reported Walking Capability Among Community-Dwelling Older  
23 People. *Journal of Aging and Physical Activity*, 1(aop), 1–8.

24 Team, R. C. (2013). R: A language and environment for statistical computing. R Foundation  
25 for Statistical Computing, Vienna, Austria. [Http://Www. R-Project. Org/](http://Www.R-Project.Org/).

- 1 Thompson, D., Peacock, O., Western, M., & Batterham, A. M. (2015). Multidimensional  
2 physical activity: An opportunity not a problem. *Exercise and Sport Sciences*  
3 *Reviews*, 43(2), 67–74. <https://doi.org/10.1249/JES.0000000000000039>
- 4 Tsai, L. T., Rantakokko, M., Rantanen, T., Viljanen, A., Kauppinen, M., & Portegijs, E.  
5 (2016). Objectively Measured Physical Activity and Changes in Life-Space Mobility  
6 Among Older People. *The Journals of Gerontology. Series A, Biological Sciences and*  
7 *Medical Sciences*, 71(11), 1466–1471.
- 8 Vähä-Ypyä, H., Husu, P., Suni, J., Vasankari, T., & Sievänen, H. (2018). Reliable recognition  
9 of lying, sitting, and standing with a hip-worn accelerometer. *Scandinavian Journal of*  
10 *Medicine & Science in Sports*, 28(3), 1092–1102.
- 11 Verswijveren, S. J., Lamb, K. E., Leech, R. M., Salmon, J. O., Timperio, A., Telford, R. M.,  
12 McNarry, M. A., Mackintosh, K. A., Daly, R. M., & Dunstan, D. W. (2020). Activity  
13 accumulation and cardiometabolic risk in youth: A latent profile approach. *Medicine*  
14 *and Science in Sports and Exercise*, 52(7), 1502–1510.
- 15 von Rosen, P., Dohrn, I.-M., & Hagströmer, M. (2020). Latent profile analysis of physical  
16 activity and sedentary behavior with mortality risk: A 15-year follow-up.  
17 *Scandinavian Journal of Medicine & Science in Sports*, 30(10), 1949–1956.
- 18 Warburton, D. E., Nicol, C. W., & Bredin, S. S. (2006). Health benefits of physical activity:  
19 The evidence. *CMAJ : Canadian Medical Association Journal = Journal de*  
20 *l'Association Médicale Canadienne*, 174(6), 801–809.
- 21 White, T., Westgate, K., Hollidge, S., Venables, M., Olivier, P., Wareham, N., & Brage, S.  
22 (2019). Estimating energy expenditure from wrist and thigh accelerometry in free-  
23 living adults: A doubly labelled water study. *International Journal of Obesity*, 43(11),  
24 Article 11. <https://doi.org/10.1038/s41366-019-0352-x>



- 1 World Health Organization (Ed.). (2001). *International classification of functioning,*
- 2 *disability and health: ICF.* World Health Organization.
- 3

Table 1. Included variables in the cluster analysis according to physical behavior clusters among women (n=264)

	Exercisers n=60	Actives n=100	Inactives n=104	P-values*		
	Mean (SD)	Mean (SD)	Mean (SD)	E vs. A	A vs. I	E vs. I
Non-movement time (min)	1255.2 (44.1)	1262.7 (32.9)	1330.7 (23.2)	1.000	<0.001	<0.001
Light PA (min/day)	122.9 (29.5)	149.5 (29.1)	93.8 (20.8)	<0.001	<0.001	<0.001
Moderate PA (min/day)	58.7 (22.6)	27.7 (13.0)	15.2 (10.7)	<0.001	<0.001	<0.001
Vigorous PA (min/day)	2.9 (5.8)	0.15 (0.35)	0.27 (0.84)	<0.001	1.000	<0.001
Relative PA (min/week)	138.3 (92.7)	42.2 (42.9)	33.0 (40.0)	<0.001	0.246	<0.001
No of sit-to-stand transitions	41.4 (13.4)	44.0 (17.3)	37.9 (15.4)	1.000	0.023	0.253
Mean gait bout duration (min)	1.17 (0.21)	0.82 (0.13)	0.78 (0.13)	<0.001	0.098	<0.001
SD of gait bout duration	2.83 (0.97)	1.11 (0.49)	1.05 (0.59)	<0.001	0.868	<0.001
Mean gait bout intensity (g)	0.13 (0.02)	0.12 (0.02)	0.11 (0.02)	0.010	0.013	<0.001
Mean no of gait bouts	114.1 (34.6)	139.2 (29.1)	82.1 (20.1)	<0.001	<0.001	<0.001
Mean MAD (g)	0.03 (0.007)	0.02 (0.004)	0.02 (0.003)	<0.001	<0.001	<0.001
Fragmentation (posture)	0.10 (0.02)	0.12 (0.04)	0.16 (0.05)	0.018	<0.001	<0.001
Fragmentation (MAD)	0.22 (0.05)	0.22 (0.04)	0.29 (0.05)	1.000	<0.001	<0.001

Note; PA=physical activity, MAD=mean amplitude deviation, E vs. A = exercisers vs. actives, A vs. I = actives vs. inactives, E vs. I = exercisers vs. inactives, \*Bonferroni-corrected p-values

Table 2. Included variables in the cluster analysis according to physical behavior clusters among men (n=177)

Men N=177	Exercisers	Actives	Inactives	P-values		
	n=19	n=77	n=81	E vs. A	A vs. I	E vs. I
	Mean (SD)	Mean (SD)	Mean (SD)			
Non-movement time (min)	1233.6 (51.1)	1228.7 (35.7)	1315.0 (31.0)	1.000	<0.001	<0.001
Light PA (min/day)	127.6 (40.4)	171.3 (35.7)	103.8 (28.2)	<0.001	<0.001	0.104
Moderate PA (min/day)	68.6 (27.4)	38.4 (15.9)	20.4 (10.5)	<0.002	<0.001	<0.001
Vigorous PA (min/day)	10.2 (9.8)	1.59 (2.95)	0.57 (1.79)	0.011	<0.001	<0.001
Relative PA (min/week)	230.1 (167.2)	48.6 (66.5)	36.8 (44.4)	<0.001	0.989	<0.001
No of sit-to-stand transitions	52.6 (20.9)	52.8 (15.7)	42.3 (14.7)	1.000	<0.001	0.054
Mean gait bout duration (min)	1.29 (0.29)	0.94 (0.15)	0.87 (0.19)	<0.001	0.027	<0.001
SD of gait bout duration	3.22 (1.21)	1.33 (0.61)	1.22 (0.79)	<0.001	0.493	<0.001
Mean gait bout intensity (g)	0.14 (0.02)	0.13 (0.02)	0.12 (0.02)	0.543	0.002	0.002
Mean no of gait bouts	117.2 (37.0)	154.2 (36.1)	91.1 (26.1)	0.002	<0.001	0.035
Mean MAD (g)	0.04 (0.007)	0.03 (0.005)	0.02 (0.004)	0.015	<0.001	<0.001
Fragmentation (posture)	0.12 (0.03)	0.13 (0.04)	0.17 (0.06)	1.000	<0.001	0.001
Fragmentation (MAD)	0.22 (0.05)	0.19 (0.03)	0.27 (0.06)	0.128	<0.001	0.001

Note; PA=physical activity, MAD=mean amplitude deviation, E vs. A = exercisers vs. actives, A vs. I = actives vs. inactives, E vs. I = exercisers vs. inactives, \*Bonferroni-corrected p-values

Table 3. Participant characteristics according to physical behavior profiles

	Women			p	Men			p
	Exercisers	Actives	Inactives		Exercisers	Actives	Inactives	
	Mean (SD)	Mean (SD)	Mean (SD)		Mean (SD)	Mean (SD)	Mean (SD)	
Age	78.0 (3.2)	77.7 (3.3)	78.8 (3.7)	0.094	77.7 (3.0.)	78.2 (3.4)	79.0 (3.5)	0.019
Years of education	11.2 (4.0)	12.5 (4.3)	11.4 (4.0)	0.092	12.2 (3.4)	12.4 (4.5)	11.0 (4.7)	0.016
MMSE score	27.7 (2.2)	27.7 (2.0)	27.6 (2.0)	0.872	26.8 (2.2)	27.3 (2.5)	27.0 (2.9)	0.758
CES-D score	6.5 (6.0)	7.7 (5.8)	9.0 (7.0)	0.028 <sup>c</sup>	7.6 (8.0)	6.5 (5.8)	7.8 (7.2)	0.581
SPPB score	10.9 (1.3)	10.6 (1.6)	9.9 (1.9)	0.002 <sup>b,c</sup>	11.4 (0.8)	10.9 (1.6)	10.1 (1.8)	<0.001 <sup>b,c</sup>
YPAS score	70.0 (22.5)	58.9 (1.1)	50.5 (18.8)	<0.001 <sup>a,b,c</sup>	79.9 (27.3)	68.5 (22.4)	53.1 (21.6)	<0.001 <sup>b,c</sup>
LSA score	75.0 (13.5)	72.7 (15.1)	63.9 (16.9)	<0.001 <sup>b,c</sup>	82.8 (15.8)	85.0 (14.5)	76.3 (17.1)	<0.001 <sup>b</sup>
6MWT walking speed (m/s)	1.3 (0.2)	1.2 (0.2)	1.0 (0.2)	<0.001 <sup>a,b,c</sup>	1.3 (0.2)	1.3 (0.2)	1.1 (0.2)	<0.001 <sup>b,c</sup>
10m walking speed (m/s)	1.9 (0.3)	1.8 (0.3)	1.6 (0.3)	<0.001 <sup>b,c</sup>	2.0 (0.3)	2.1 (0.4)	1.8 (0.4)	<0.001 <sup>b</sup>
Max. knee extension force (N)	318.5 (70.3)	299.8 (79.0)	269.5 (81.3)	<0.001 <sup>b,c</sup>	445.8 (92.8)	457.1 (91.4)	405.6 (114.4)	0.004 <sup>b</sup>
	%	%	%		%	%	%	
Living alone	56.7	40.4	51.9	0.203	15.8	19.7	12.5	0.468
Wants to be more physically active	38.6	70.7	72.6	<0.001	35.3	50.0	72.0	0.003
Good perceived opportunities for PA	88.1	69.7	50.0	<0.001	78.9	84.9	69.6	0.078

Note; SD=Standard deviation, PA=Physical activity, 6MWT=6-minute walk test, MMSE=Mini Mental State Examination, CES-D= Centre for Epidemiologic studies Depression Scale, SPPB=Short Physical Performance Battery, YPAS=Yale Physical Activity Survey, LSA= University of Alabama at Birmingham Study of Aging Life-Space Assessment (LSA) questionnaire, <sup>a</sup>Bonferroni-corrected p<0.05 between the exercisers and the actives, <sup>b</sup> Bonferroni-corrected p<0.05 between the actives and the inactives, <sup>c</sup> Bonferroni-corrected p<0.05 between the exercisers and the inactives

Table 4. Age-adjusted associations between the physical behavior profiles, physical capacity, life-space mobility and self-reported physical activity

	Exercisers	Actives	Inactives	P-values*		
	(E)	(A)	(I)	E vs. A	A vs. I	E vs. I
<b>Women n=264</b>	EMM (SE)	EMM (SE)	EMM (SE)			
10m walking speed (m/s)	1.9 (0.04)	1.8 (0.03)	1.6 (0.03)	0.071	0.001	<0.001
6MWT walking speed (m/s)	1.3 (0.02)	1.2 (0.02)	1.1 (0.02)	0.004	<0.001	<0.001
Max. knee extension force (N)	317.4 (9.70)	296.7 (7.54)	273.0 (7.44)	0.281	0.080	0.001
SPPB	10.8 (0.21)	10.6 (0.16)	10.0 (0.16)	0.877	0.041	0.004
CES-D	6.6 (0.81)	7.9 (0.64)	8.9 (0.63)	0.628	0.805	0.079
YPAS	69.9 (2.67)	58.6 (2.08)	50.9 (2.04)	0.003	0.027	<0.001
LSA	74.9 (1.97)	72.5 (1.54)	64.3 (1.50)	1.000	<0.001	<0.001
<b>Men n=177</b>						
10m walking speed (m/s)	2.0 (0.09)	2.0 (0.04)	1.8 (0.04)	1.000	0.001	0.274
6MWT walking speed (m/s)	1.3 (0.04)	1.3 (0.02)	1.1 (0.02)	1.000	<0.001	0.001
Max. knee extension force (N)	438.7 (22.84)	452.9 (11.11)	411.1 (10.79)	1.000	0.023	0.829
SPPB	11.3 (0.37)	10.9 (0.18)	10.2 (0.18)	0.856	0.027	0.021
CES-D	7.8 (1.53)	6.5 (0.77)	7.7 (0.75)	1.000	0.965	1.000
YPAS	79.7 (5.21)	68.4 (2.62)	53.2 (2.54)	0.166	<0.001	<0.001
LSA	81.9 (3.55)	84.5 (1.77)	76.9 (1.72)	1.000	0.009	0.646

Note; EMM=Age-adjusted estimated marginal means, SE=Standard error, 6MWT=6-minute walk test, CES-D= Centre for Epidemiologic studies Depression Scale, SPPB=Short Physical Performance Battery, YPAS=Yale Physical Activity Survey, LSA= University of Alabama at Birmingham Study of Aging Life-Space Assessment (LSA) questionnaire, E vs. A = exercisers vs. actives, A vs. I = actives vs. inactives, E vs. I = exercisers vs. inactives, \*Bonferroni corrected p-values

